Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
United States												
All races, both sexes	2,397,615	27,936	4,785	2,888	3,946	13,706	19,715	18,771	22,097	31,944	53,418	77,927
Male	1,181,668	15,718	2,649	1,645	2,354	9,678	14,909	13,554	14,805	20,485	33,192	48,514
Female	1,215,947	12,218	2,136	1,243	1,592	4,028	4,806	5,217	7,292	11,459	20,226	29,413
White, both sexes	2,056,643	18,231	3,363	2,043	2,857	10,450	14,563	13,421	15,854	23,686	40,354	59,181
Male	1,007,266	10,265	1,878	1,178	1,727	7,267	11,039	9,795	10,789	15,483	25,732	37,461
Female	1,049,377	7,966	1,485	865	1,130	3,183	3,524	3,626	5,065	8,203	14,622	21,720
Black, both sexes	287,315	8,494	1,164	685	899	2,680	4,369	4,616	5,309	7,098	11,376	16,423
Male	145,970	4,769	641	372	516	2,019	3,317	3,264	3,420	4,262	6,438	9,640
Female	141,345	3,725	523	313	383	661	1,052	1,352	1,889	2,836	4,938	6,783
American Indian 1/, both sexes	13,124	389	97	53	74	260	333	314	381	516	678	833
Male	7,134	237	50	28	42	171	238	222	239	338	409	508
Female	5,990	152	47	25	32	89	95	92	142	178	269	325
Asian or Pacific Islander 2/, both sexes	40,533	822	161	107	116	316	450	420	553	644	1,010	1,490
Male	21,298	447	80	67	69	221	315	273	357	402	613	905
Female	19,235	375	81	40	47	95	135	147	196	242	397	585
Alabama												
All races, both sexes	46,121	516	106	49	79	315	470	408	476	678	1,192	1,634
Male	22,739	322	65	20	50	212	348	280	307	398	736	994
Female	23,382	194	41	29	29	103	122	128	169	280	456	640
White, both sexes	34,891	275	66	28	54	220	312	237	293	441	779	1,084
Male	17,196	164	46	10	35	149	240	157	179	263	511	672
Female	17,695	111	20	18	19	71	72	80	114	178	268	412
Black, both sexes	11,095	239	37	21	25	94	152	166	178	233	409	540
Male	5,480	156	19	10	15	63	104	122	126	131	224	315
Female	5,615	83	18	11	10	31	48	44	52	102	185	225
American Indian 1/, both sexes	52	1	-	-	-	-	-	3	1	1	1	2
Male	22	1	-	-	-	-	-	1	1	1	-	-
Female	30	-	-	-	-	-	-	2	-	-	1	2
Asian or Pacific Islander 2/, both sexes	83	1	3	-	-	1	6	2	4	3	3	8
Male	41	1	-	-	-	-	4	-	1	3	1	7
Female	42	-	3	-	-	1	2	2	3	-	2	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Alaska												
All races, both sexes	3,051	69	22	5	24	64	73	58	63	102	125	211
Male	1,760	41	12	4	15	50	58	43	45	61	73	130
Female	1,291	28	10	1	9	14	15	15	18	41	52	81
White, both sexes	2,160	36	12	1	17	32	43	31	39	62	78	153
Male	1,261	18	6	1	10	25	35	26	31	40	49	98
Female	899	18	6	-	7	7	8	5	8	22	29	55
Black, both sexes	97	6	1	1	-	5	3	3	1	2	3	13
Male	60	6	1	1	-	5	3	1	1	1	-	8
Female	37	-	-	-	-	-	-	2	-	1	3	5
American Indian 1/, both sexes	700	27	7	3	6	26	25	23	21	36	38	40
Male	383	17	3	2	4	19	19	15	11	19	21	20
Female	317	10	4	1	2	7	6	8	10	17	17	20
Asian or Pacific Islander 2/, both sexes	94	-	2	-	1	1	2	1	2	2	6	5
Male	56	-	2	-	1	1	1	1	2	1	3	4
Female	38	-	-	-	-	-	1	-	-	1	3	1
Arizona												
All races, both sexes	43,198	630	111	58	88	343	483	440	447	699	1,018	1,450
Male	22,674	357	61	36	55	241	380	327	303	484	655	947
Female	20,524	273	50	22	33	102	103	113	144	215	363	503
White, both sexes	40,200	531	87	48	73	278	409	359	362	562	861	1,247
Male	21,026	299	48	32	46	196	326	267	242	385	562	805
Female	19,174	232	39	16	27	82	83	92	120	177	299	442
Black, both sexes	1,125	41	7	6	6	22	21	24	23	42	52	70
Male	615	28	4	2	4	16	17	18	17	34	30	50
Female	510	13	3	4	2	6	4	6	6	8	22	20
American Indian 1/, both sexes	1,564	48	15	4	9	41	51	49	59	85	95	119
Male	885	26	8	2	5	27	36	36	41	57	59	83
Female	679	22	7	2	4	14	15	13	18	28	36	36
Asian or Pacific Islander 2/, both sexes	309	10	2	-	-	2	2	8	3	10	10	14
Male	148	4	1	-	-	2	1	6	3	8	4	9
Female	161	6	1	-	-	-	1	2	-	2	6	5

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Arkansas												
All races, both sexes	27,528	319	78	49	51	181	242	266	299	370	632	956
Male	13,682	176	48	31	32	114	180	189	182	228	400	601
Female	13,846	143	30	18	19	67	62	77	117	142	232	355
White, both sexes	23,613	216	42	39	38	146	183	189	220	278	493	739
Male	11,693	117	24	25	24	92	134	134	139	173	316	472
Female	11,920	99	18	14	14	54	49	55	81	105	177	267
Black, both sexes	3,823	99	34	9	13	34	57	72	77	90	137	213
Male	1,938	56	22	5	8	21	44	51	41	53	82	128
Female	1,885	43	12	4	5	13	13	21	36	37	55	85
American Indian 1/, both sexes	50	3	1	-	-	-	-	4	1	1	2	1
Male	30	3	1	-	-	-	-	4	1	1	2	1
Female	20	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	42	1	1	1	-	1	2	1	1	1	-	3
Male	21	-	1	1	-	1	2	-	1	1	-	-
Female	21	1	-	-	-	-	-	1	-	-	-	3
California												
All races, both sexes	232,525	2,811	524	328	427	1,520	2,149	1,966	2,411	3,375	5,508	8,080
Male	116,978	1,583	283	183	262	1,131	1,673	1,455	1,670	2,286	3,527	5,048
Female	115,547	1,228	241	145	165	389	476	511	741	1,089	1,981	3,032
White, both sexes	195,727	2,158	396	249	330	1,140	1,682	1,492	1,799	2,641	4,399	6,267
Male	97,666	1,207	208	143	206	830	1,314	1,113	1,261	1,810	2,888	3,961
Female	98,061	951	188	106	124	310	368	379	538	831	1,511	2,306
Black, both sexes	18,685	397	67	41	58	248	289	307	378	473	696	1,129
Male	9,749	229	39	22	33	211	231	230	259	301	389	657
Female	8,936	168	28	19	25	37	58	77	119	172	307	472
American Indian 1/, both sexes	1,255	20	4	1	1	12	16	26	27	39	52	87
Male	665	13	3	-	-	7	11	19	17	29	36	57
Female	590	7	1	1	1	5	5	7	10	10	16	30
Asian or Pacific Islander 2/, both sexes	16,858	236	57	37	38	120	162	141	207	222	361	597
Male	8,898	134	33	18	23	83	117	93	133	146	214	373
Female	7,960	102	24	19	15	37	45	48	74	76	147	224

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Colorado												
All races, both sexes	28,309	434	54	42	56	241	270	294	350	456	741	1,074
Male	14,307	261	27	21	34	165	202	212	241	303	468	675
Female	14,002	173	27	21	22	76	68	82	109	153	273	399
White, both sexes	26,894	374	49	40	50	218	246	265	314	411	680	988
Male	13,536	228	25	19	29	149	183	189	221	271	431	619
Female	13,358	146	24	21	21	69	63	76	93	140	249	369
Black, both sexes	959	43	4	1	6	16	16	21	25	32	49	68
Male	529	22	2	1	5	10	14	15	14	24	30	44
Female	430	21	2	-	1	6	2	6	11	8	19	24
American Indian 1/, both sexes	141	4	1	-	-	4	5	4	6	8	5	6
Male	85	2	-	-	-	3	3	4	4	7	5	6
Female	56	2	1	-	-	1	2	-	2	1	-	-
Asian or Pacific Islander 2/, both sexes	315	13	-	1	-	3	3	4	5	5	7	12
Male	157	9	-	1	-	3	2	4	2	1	2	6
Female	158	4	-	-	-	-	1	-	3	4	5	6
Connecticut												
All races, both sexes	29,314	233	31	28	32	103	171	164	191	338	567	826
Male	13,930	129	12	19	19	72	131	118	117	196	341	543
Female	15,384	104	19	9	13	31	40	46	74	142	226	283
White, both sexes	27,092	155	23	22	23	84	136	132	161	285	458	711
Male	12,798	88	8	16	13	58	103	95	100	168	278	471
Female	14,294	67	15	6	10	26	33	37	61	117	180	240
Black, both sexes	2,019	68	7	5	9	15	30	29	27	48	103	107
Male	1,031	37	4	3	6	12	25	21	17	27	59	67
Female	988	31	3	2	3	3	5	8	10	21	44	40
American Indian 1/, both sexes	42	2	-	-	-	2	-	-	-	2	-	1
Male	24	1	-	-	-	1	-	-	-	1	-	1
Female	18	1	-	-	-	1	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	161	8	1	1	-	2	5	3	3	3	6	7
Male	77	3	-	-	-	1	3	2	-	-	4	4
Female	84	5	1	1	-	1	2	1	3	3	2	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Delaware												
All races, both sexes	7,143	98	19	16	9	41	57	49	65	106	167	239
Male	3,618	54	8	6	4	27	41	35	41	66	101	144
Female	3,525	44	11	10	5	14	16	14	24	40	66	95
White, both sexes	5,888	52	12	13	7	30	42	36	45	70	117	174
Male	2,971	27	4	6	2	20	28	26	31	42	72	101
Female	2,917	25	8	7	5	10	14	10	14	28	45	73
Black, both sexes	1,203	45	7	3	2	11	15	13	19	36	45	63
Male	622	27	4	-	2	7	13	9	9	24	26	42
Female	581	18	3	3	-	4	2	4	10	12	19	21
American Indian 1/, both sexes	23	-	-	-	-	-	-	-	1	-	3	2
Male	12	-	-	-	-	-	-	-	1	-	2	1
Female	11	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	29	1	-	-	-	-	-	-	-	-	2	-
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	16	1	-	-	-	-	-	-	-	-	1	-
Dist. of Columbia												
All races, both sexes	5,454	95	7	8	16	49	67	69	92	104	208	266
Male	2,785	47	4	6	7	43	57	44	60	64	126	164
Female	2,669	48	3	2	9	6	10	25	32	40	82	102
White, both sexes	1,145	14	1	1	2	1	11	6	16	12	29	30
Male	593	8	1	1	-	1	9	5	11	6	24	23
Female	552	6	-	-	2	-	2	1	5	6	5	7
Black, both sexes	4,258	79	6	7	14	48	55	62	76	92	179	234
Male	2,165	37	3	5	7	42	48	39	49	58	102	139
Female	2,093	42	3	2	7	6	7	23	27	34	77	95
American Indian 1/, both sexes	8	2	-	-	-	-	-	-	-	-	-	1
Male	5	2	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	43	-	-	-	-	-	1	1	-	-	-	1
Male	22	-	-	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Florida												
All races, both sexes	169,008	1,537	287	164	228	765	1,241	1,179	1,433	2,241	3,577	5,210
Male	86,544	854	158	92	148	534	923	825	935	1,431	2,169	3,243
Female	82,464	683	129	72	80	231	318	354	498	810	1,408	1,967
White, both sexes	149,959	884	188	109	147	579	898	818	1,005	1,629	2,711	3,995
Male	76,658	485	108	60	96	395	683	597	690	1,083	1,713	2,534
Female	73,301	399	80	49	51	184	215	221	315	546	998	1,461
Black, both sexes	18,089	623	94	54	76	173	327	346	411	584	842	1,167
Male	9,411	357	46	31	49	131	229	220	233	335	445	683
Female	8,678	266	48	23	27	42	98	126	178	249	397	484
American Indian 1/, both sexes	175	3	1	-	1	1	7	7	4	8	10	14
Male	87	-	1	-	1	1	4	5	3	2	3	8
Female	88	3	-	-	-	-	3	2	1	6	7	6
Asian or Pacific Islander 2/, both sexes	785	27	4	1	4	12	9	8	13	20	14	34
Male	388	12	3	1	2	7	7	3	9	11	8	18
Female	397	15	1	-	2	5	2	5	4	9	6	16
Georgia												
All races, both sexes	65,818	1,181	189	97	131	423	663	720	806	1,159	1,940	2,698
Male	32,683	658	110	52	71	288	493	494	526	747	1,179	1,625
Female	33,135	523	79	45	60	135	170	226	280	412	761	1,073
White, both sexes	47,658	545	107	53	75	277	398	404	439	670	1,162	1,661
Male	23,465	310	62	30	47	188	303	275	288	438	715	1,046
Female	24,193	235	45	23	28	89	95	129	151	232	447	615
Black, both sexes	17,702	613	75	40	52	143	256	302	355	474	766	1,012
Male	8,960	335	46	20	21	99	183	212	227	301	458	560
Female	8,742	278	29	20	31	44	73	90	128	173	308	452
American Indian 1/, both sexes	133	13	3	1	2	1	3	4	4	7	4	6
Male	73	9	2	1	1	1	2	1	4	5	2	3
Female	60	4	1	-	1	-	1	3	-	2	2	3
Asian or Pacific Islander 2/, both sexes	325	10	4	3	2	2	6	10	8	8	8	19
Male	185	4	-	1	2	-	5	6	7	3	4	16
Female	140	6	4	2	-	2	1	4	1	5	4	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Hawaii												
All races, both sexes	9,030	104	23	7	18	34	58	70	72	120	187	291
Male	4,986	66	10	2	11	24	47	48	54	77	122	199
Female	4,044	38	13	5	7	10	11	22	18	43	65	92
White, both sexes	2,529	27	8	2	6	5	26	22	20	42	60	110
Male	1,417	16	4	-	2	3	21	16	18	26	36	75
Female	1,112	11	4	2	4	2	5	6	2	16	24	35
Black, both sexes	81	9	1	1	2	1	-	1	5	4	4	5
Male	54	5	1	-	2	-	-	1	5	2	4	2
Female	27	4	-	1	-	1	-	-	-	2	-	3
American Indian 1/, both sexes	13	-	1	-	-	-	-	-	-	1	-	2
Male	6	-	-	-	-	-	-	-	-	1	-	2
Female	7	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6,407	68	13	4	10	28	32	47	47	73	123	174
Male	3,509	45	5	2	7	21	26	31	31	48	82	120
Female	2,898	23	8	2	3	7	6	16	16	25	41	54
Idaho												
All races, both sexes	10,028	139	37	15	24	74	92	87	78	134	186	321
Male	5,075	83	22	9	17	58	74	62	53	93	116	210
Female	4,953	56	15	6	7	16	18	25	25	41	70	111
White, both sexes	9,866	133	33	15	24	71	84	85	73	129	183	304
Male	4,984	79	20	9	17	56	69	61	50	89	114	197
Female	4,882	54	13	6	7	15	15	24	23	40	69	107
Black, both sexes	19	1	2	-	-	1	1	1	-	-	-	2
Male	13	1	1	-	-	-	1	-	-	-	-	2
Female	6	-	1	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	99	5	1	-	-	2	4	1	5	5	2	10
Male	53	3	-	-	-	2	2	1	3	4	1	8
Female	46	2	1	-	-	-	2	-	2	1	1	2
Asian or Pacific Islander 2/, both sexes	44	-	1	-	-	-	3	-	-	-	1	5
Male	25	-	1	-	-	-	2	-	-	-	1	3
Female	19	-	-	-	-	-	1	-	-	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Illinois												
All races, both sexes	102,670	1,349	205	124	149	558	775	745	890	1,291	2,192	3,192
Male	49,284	765	115	69	90	393	596	561	608	809	1,387	1,954
Female	53,386	584	90	55	59	165	179	184	282	482	805	1,238
White, both sexes	85,571	835	120	85	108	381	529	499	593	848	1,484	2,194
Male	40,730	483	70	46	65	268	399	384	417	558	978	1,393
Female	44,841	352	50	39	43	113	130	115	176	290	506	801
Black, both sexes	15,919	488	82	38	39	171	235	233	289	421	673	965
Male	7,949	265	44	22	24	121	190	170	186	240	388	540
Female	7,970	223	38	16	15	50	45	63	103	181	285	425
American Indian 1/, both sexes	63	7	-	1	-	-	1	-	1	2	3	4
Male	36	5	-	1	-	-	1	-	1	-	3	4
Female	27	2	-	-	-	-	-	-	-	2	-	-
Asian or Pacific Islander 2/, both sexes	1,117	19	3	-	2	6	10	13	7	20	32	29
Male	569	12	1	-	1	4	6	7	4	11	18	17
Female	548	7	2	-	1	2	4	6	3	9	14	12
Indiana												
All races, both sexes	54,211	700	127	70	96	303	446	374	472	632	1,173	1,677
Male	26,335	374	66	46	57	216	328	284	321	393	726	1,079
Female	27,876	326	61	24	39	87	118	90	151	239	447	598
White, both sexes	49,741	528	108	54	80	255	355	296	381	536	1,012	1,431
Male	23,983	281	55	35	43	177	254	223	254	332	634	923
Female	25,758	247	53	19	37	78	101	73	127	204	378	508
Black, both sexes	4,289	163	18	14	14	43	85	74	83	92	157	238
Male	2,260	87	10	9	12	35	69	57	61	59	89	152
Female	2,029	76	8	5	2	8	16	17	22	33	68	86
American Indian 1/, both sexes	32	1	1	1	-	-	1	1	-	-	1	2
Male	15	-	1	1	-	-	1	1	-	-	1	2
Female	17	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	149	8	-	1	2	5	5	3	8	4	3	6
Male	77	6	-	1	2	4	4	3	6	2	2	2
Female	72	2	-	-	-	1	1	-	2	2	1	4

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Iowa												
All races, both sexes	26,897	195	30	36	41	95	162	120	160	205	372	601
Male	12,881	106	12	18	29	66	122	82	99	132	238	381
Female	14,016	89	18	18	12	29	40	38	61	73	134	220
White, both sexes	26,368	173	25	31	40	89	149	111	151	190	351	579
Male	12,610	92	12	17	28	62	112	77	96	120	226	368
Female	13,758	81	13	14	12	27	37	34	55	70	125	211
Black, both sexes	411	14	4	3	1	4	10	8	6	11	15	19
Male	207	9	-	-	1	3	7	5	2	8	8	11
Female	204	5	4	3	-	1	3	3	4	3	7	8
American Indian 1/, both sexes	35	2	-	1	-	-	1	-	2	-	4	-
Male	17	1	-	-	-	-	1	-	1	-	3	-
Female	18	1	-	1	-	-	-	-	1	-	1	-
Asian or Pacific Islander 2/, both sexes	83	6	1	1	-	2	2	1	1	4	2	3
Male	47	4	-	1	-	1	2	-	-	4	1	2
Female	36	2	1	-	-	1	-	1	1	-	1	1
Kansas												
All races, both sexes	23,818	284	61	24	49	116	199	178	174	249	446	685
Male	11,316	163	40	13	31	83	146	118	104	166	268	427
Female	12,502	121	21	11	18	33	53	60	70	83	178	258
White, both sexes	22,297	228	51	21	41	105	165	147	144	221	388	616
Male	10,530	125	34	11	25	76	122	94	87	145	233	391
Female	11,767	103	17	10	16	29	43	53	57	76	155	225
Black, both sexes	1,280	47	10	3	6	9	33	28	25	23	46	59
Male	666	31	6	2	5	7	23	22	14	17	27	31
Female	614	16	4	1	1	2	10	6	11	6	19	28
American Indian 1/, both sexes	145	1	-	-	1	1	1	1	3	3	10	8
Male	74	1	-	-	-	-	1	-	2	2	7	5
Female	71	-	-	-	1	1	-	1	1	1	3	3
Asian or Pacific Islander 2/, both sexes	96	8	-	-	1	1	-	2	2	2	2	2
Male	46	6	-	-	1	-	-	2	1	2	1	-
Female	50	2	-	-	-	1	-	-	1	-	1	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Kentucky												
All races, both sexes	38,646	378	81	41	61	268	343	382	423	592	923	1,374
Male	19,168	225	46	24	40	177	244	268	269	377	598	907
Female	19,478	153	35	17	21	91	99	114	154	215	325	467
White, both sexes	36,001	312	68	36	57	237	298	346	386	546	835	1,217
Male	17,841	178	39	23	38	155	208	240	243	348	547	808
Female	18,160	134	29	13	19	82	90	106	143	198	288	409
Black, both sexes	2,563	62	12	5	4	29	39	35	35	44	86	153
Male	1,283	43	7	1	2	21	33	27	25	28	49	96
Female	1,280	19	5	4	2	8	6	8	10	16	37	57
American Indian 1/, both sexes	16	-	-	-	-	-	1	1	-	2	-	1
Male	10	-	-	-	-	-	-	1	-	1	-	1
Female	6	-	-	-	-	-	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	66	4	1	-	-	2	5	-	2	-	2	3
Male	34	4	-	-	-	1	3	-	1	-	2	2
Female	32	-	1	-	-	1	2	-	1	-	-	1
Louisiana												
All races, both sexes	42,215	684	125	73	107	329	574	544	551	709	1,183	1,666
Male	21,255	372	54	40	65	251	447	377	379	438	744	1,058
Female	20,960	312	71	33	42	78	127	167	172	271	439	608
White, both sexes	28,800	284	53	39	58	197	287	270	297	387	691	922
Male	14,166	159	19	16	33	141	217	179	207	247	455	599
Female	14,634	125	34	23	25	56	70	91	90	140	236	323
Black, both sexes	13,167	395	71	33	47	125	278	269	244	309	482	728
Male	6,945	209	35	24	31	105	223	194	166	182	282	447
Female	6,222	186	36	9	16	20	55	75	78	127	200	281
American Indian 1/, both sexes	72	1	-	-	-	3	4	3	1	5	2	3
Male	42	1	-	-	-	1	4	3	-	3	2	3
Female	30	-	-	-	-	2	-	-	1	2	-	-
Asian or Pacific Islander 2/, both sexes	176	4	1	1	2	4	5	2	9	8	8	13
Male	102	3	-	-	1	4	3	1	6	6	5	9
Female	74	1	1	1	1	-	2	1	3	2	3	4

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Maine												
All races, both sexes	12,443	79	22	10	14	56	88	74	62	125	213	302
Male	5,885	51	11	6	11	40	61	50	46	84	128	188
Female	6,558	28	11	4	3	16	27	24	16	41	85	114
White, both sexes	12,320	73	17	7	14	56	85	74	61	118	208	297
Male	5,816	47	9	4	11	40	58	50	45	80	125	184
Female	6,504	26	8	3	3	16	27	24	16	38	83	113
Black, both sexes	42	4	4	2	-	-	2	-	1	1	2	-
Male	27	2	2	1	-	-	2	-	1	1	1	-
Female	15	2	2	1	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	53	1	1	1	-	-	-	-	-	4	1	4
Male	30	1	-	1	-	-	-	-	-	3	-	3
Female	23	-	1	-	-	-	-	-	-	1	1	1
Asian or Pacific Islander 2/, both sexes	28	1	-	-	-	-	1	-	-	2	2	1
Male	12	1	-	-	-	-	1	-	-	-	2	1
Female	16	-	-	-	-	-	-	-	-	2	-	-
Maryland												
All races, both sexes	43,232	630	93	54	84	266	412	378	453	737	1,181	1,630
Male	21,260	352	57	33	58	192	319	296	295	491	723	995
Female	21,972	278	36	21	26	74	93	82	158	246	458	635
White, both sexes	30,793	252	43	21	42	133	202	182	201	391	586	907
Male	14,804	131	26	12	30	91	142	139	131	262	373	532
Female	15,989	121	17	9	12	42	60	43	70	129	213	375
Black, both sexes	11,740	359	48	31	42	129	203	185	243	334	574	699
Male	6,105	211	30	19	28	98	172	149	159	220	338	449
Female	5,635	148	18	12	14	31	31	36	84	114	236	250
American Indian 1/, both sexes	73	2	-	-	-	-	-	2	1	3	3	3
Male	38	2	-	-	-	-	-	1	-	3	2	2
Female	35	-	-	-	-	-	-	1	1	-	1	1
Asian or Pacific Islander 2/, both sexes	626	17	2	2	-	4	7	9	8	9	18	21
Male	313	8	1	2	-	3	5	7	5	6	10	12
Female	313	9	1	-	-	1	2	2	3	3	8	9

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Massachusetts												
All races, both sexes	54,511	380	45	36	57	195	322	286	359	620	992	1,483
Male	25,410	228	29	22	30	154	244	197	239	422	619	936
Female	29,101	152	16	14	27	41	78	89	120	198	373	547
White, both sexes	51,649	284	34	32	42	153	274	242	296	541	891	1,348
Male	23,866	175	21	18	21	116	206	166	200	370	560	851
Female	27,783	109	13	14	21	37	68	76	96	171	331	497
Black, both sexes	2,267	80	8	-	12	34	39	36	49	60	83	122
Male	1,231	44	7	-	6	30	32	28	29	41	47	75
Female	1,036	36	1	-	6	4	7	8	20	19	36	47
American Indian 1/, both sexes	55	1	-	-	-	1	-	-	-	3	4	1
Male	29	-	-	-	-	1	-	-	-	1	4	1
Female	26	1	-	-	-	-	-	-	-	2	-	-
Asian or Pacific Islander 2/, both sexes	540	15	3	4	3	7	9	8	14	16	14	12
Male	284	9	1	4	3	7	6	3	10	10	8	9
Female	256	6	2	-	-	-	3	5	4	6	6	3
Michigan												
All races, both sexes	85,169	984	136	105	138	476	594	597	770	1,101	1,779	2,877
Male	41,595	556	65	63	66	329	444	438	501	704	1,098	1,750
Female	43,574	428	71	42	72	147	150	159	269	397	681	1,127
White, both sexes	72,237	560	93	65	93	352	403	396	515	780	1,361	2,149
Male	34,914	318	51	43	49	242	307	298	326	506	863	1,345
Female	37,323	242	42	22	44	110	96	98	189	274	498	804
Black, both sexes	12,102	396	41	38	41	118	175	196	239	291	393	684
Male	6,226	219	14	18	15	84	126	136	168	178	220	383
Female	5,876	177	27	20	26	34	49	60	71	113	173	301
American Indian 1/, both sexes	406	4	-	-	2	2	8	1	6	14	12	21
Male	224	4	-	-	1	1	6	1	2	9	6	12
Female	182	-	-	-	1	1	2	-	4	5	6	9
Asian or Pacific Islander 2/, both sexes	424	24	2	2	2	4	8	4	10	16	13	23
Male	231	15	-	2	1	2	5	3	5	11	9	10
Female	193	9	2	-	1	2	3	1	5	5	4	13

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Minnesota												
All races, both sexes	37,034	332	66	42	60	194	220	218	248	357	657	950
Male	17,936	198	35	26	32	129	176	159	168	228	394	616
Female	19,098	134	31	16	28	65	44	59	80	129	263	334
White, both sexes	35,491	244	53	32	49	163	178	178	202	302	555	850
Male	17,072	144	29	19	26	105	145	134	137	198	333	553
Female	18,419	100	24	13	23	58	33	44	65	104	222	297
Black, both sexes	803	61	5	3	4	15	28	22	26	30	59	63
Male	441	39	3	1	1	12	21	13	18	16	37	41
Female	362	22	2	2	3	3	7	9	8	14	22	22
American Indian 1/, both sexes	397	16	6	3	3	9	7	12	15	16	27	26
Male	238	9	3	3	1	6	6	9	8	11	13	14
Female	159	7	3	-	2	3	1	3	7	5	14	12
Asian or Pacific Islander 2/, both sexes	343	11	2	4	4	7	7	6	5	9	16	11
Male	185	6	-	3	4	6	4	3	5	3	11	8
Female	158	5	2	1	-	1	3	3	-	6	5	3
Mississippi												
All races, both sexes	27,871	420	82	40	59	220	296	303	360	444	751	1,147
Male	14,112	247	46	26	41	160	203	218	239	272	463	712
Female	13,759	173	36	14	18	60	93	85	121	172	288	435
White, both sexes	18,790	146	32	18	28	134	150	150	194	243	380	617
Male	9,518	82	17	14	20	96	112	115	133	161	248	397
Female	9,272	64	15	4	8	38	38	35	61	82	132	220
Black, both sexes	8,948	270	47	22	30	84	143	151	165	199	361	519
Male	4,526	162	27	12	21	63	88	101	105	110	206	310
Female	4,422	108	20	10	9	21	55	50	60	89	155	209
American Indian 1/, both sexes	63	2	3	-	1	1	1	-	-	-	9	7
Male	36	2	2	-	-	-	1	-	-	-	8	2
Female	27	-	1	-	1	1	-	-	-	-	1	5
Asian or Pacific Islander 2/, both sexes	70	2	-	-	-	1	2	2	1	2	1	4
Male	32	1	-	-	-	1	2	2	1	1	1	3
Female	38	1	-	-	-	-	-	-	-	1	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Missouri												
All races, both sexes	53,950	584	111	66	97	331	437	382	455	651	1,123	1,696
Male	26,212	321	57	40	49	225	309	280	309	413	694	1,071
Female	27,738	263	54	26	48	106	128	102	146	238	429	625
White, both sexes	48,061	409	91	48	71	263	335	279	342	523	914	1,389
Male	23,219	224	49	28	36	173	229	198	234	330	578	884
Female	24,842	185	42	20	35	90	106	81	108	193	336	505
Black, both sexes	5,595	168	17	18	24	64	96	98	104	120	194	295
Male	2,847	91	6	12	12	49	76	77	71	78	108	180
Female	2,748	77	11	6	12	15	20	21	33	42	86	115
American Indian 1/, both sexes	106	-	2	-	-	2	2	4	2	5	7	5
Male	59	-	1	-	-	1	2	4	1	3	2	5
Female	47	-	1	-	-	1	-	-	1	2	5	-
Asian or Pacific Islander 2/, both sexes	188	7	1	-	2	2	4	1	7	3	8	7
Male	87	6	1	-	1	2	2	1	3	2	6	2
Female	101	1	-	-	1	-	2	-	4	1	2	5
Montana												
All races, both sexes	8,094	52	17	14	17	73	74	62	59	89	182	254
Male	4,070	29	7	9	11	55	57	45	43	55	92	139
Female	4,024	23	10	5	6	18	17	17	16	34	90	115
White, both sexes	7,627	41	14	10	11	53	59	48	42	75	156	221
Male	3,804	24	6	7	6	40	45	36	31	48	78	122
Female	3,823	17	8	3	5	13	14	12	11	27	78	99
Black, both sexes	11	1	-	-	-	1	1	-	-	-	-	3
Male	8	-	-	-	-	1	-	-	-	-	-	2
Female	3	1	-	-	-	-	1	-	-	-	-	1
American Indian 1/, both sexes	437	10	2	3	6	19	14	13	17	14	25	30
Male	250	5	-	2	5	14	12	9	12	7	13	15
Female	187	5	2	1	1	5	2	4	5	7	12	15
Asian or Pacific Islander 2/, both sexes	19	-	1	1	-	-	-	1	-	-	1	-
Male	8	-	1	-	-	-	-	-	-	-	1	-
Female	11	-	-	1	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Nebraska												
All races, both sexes	14,657	173	37	19	26	87	97	69	86	132	218	342
Male	7,132	96	17	5	16	59	70	44	57	85	125	213
Female	7,525	77	20	14	10	28	27	25	29	47	93	129
White, both sexes	14,027	140	29	15	24	76	88	64	76	115	193	305
Male	6,812	82	12	4	15	50	63	42	53	75	108	196
Female	7,215	58	17	11	9	26	25	22	23	40	85	109
Black, both sexes	493	26	6	3	2	8	6	3	9	13	18	34
Male	245	10	4	1	1	7	4	2	4	8	12	15
Female	248	16	2	2	1	1	2	1	5	5	6	19
American Indian 1/, both sexes	94	5	1	-	-	1	3	2	1	3	6	2
Male	57	2	1	-	-	1	3	-	-	1	4	2
Female	37	3	-	-	-	-	-	2	1	2	2	-
Asian or Pacific Islander 2/, both sexes	43	2	1	1	-	2	-	-	-	1	1	1
Male	18	2	-	-	-	1	-	-	-	1	1	-
Female	25	-	1	1	-	1	-	-	-	-	-	1
Nevada												
All races, both sexes	17,929	225	40	11	47	120	162	200	210	298	468	670
Male	9,752	121	26	5	27	76	127	151	144	190	318	428
Female	8,177	104	14	6	20	44	35	49	66	108	150	242
White, both sexes	16,050	153	33	8	40	95	129	162	170	245	377	562
Male	8,739	80	24	3	25	60	103	125	118	157	265	367
Female	7,311	73	9	5	15	35	26	37	52	88	112	195
Black, both sexes	1,195	55	2	2	6	18	20	33	31	44	58	72
Male	667	33	1	2	2	12	16	23	20	29	35	40
Female	528	22	1	-	4	6	4	10	11	15	23	32
American Indian 1/, both sexes	168	4	1	-	-	3	4	1	5	5	15	12
Male	98	4	1	-	-	1	2	1	2	3	7	10
Female	70	-	-	-	-	2	2	-	3	2	8	2
Asian or Pacific Islander 2/, both sexes	516	13	4	1	1	4	9	4	4	4	18	24
Male	248	4	-	-	-	3	6	2	4	1	11	11
Female	268	9	4	1	1	1	3	2	-	3	7	13

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
New Hampshire												
All races, both sexes	10,111	81	17	9	12	44	58	38	59	113	190	272
Male	4,852	47	11	7	5	29	51	31	38	72	115	168
Female	5,259	34	6	2	7	15	7	7	21	41	75	104
White, both sexes	10,043	73	16	8	12	43	56	38	59	107	190	269
Male	4,819	44	11	6	5	28	49	31	38	69	115	167
Female	5,224	29	5	2	7	15	7	7	21	38	75	102
Black, both sexes	39	7	1	1	-	-	1	-	-	3	-	1
Male	19	3	-	1	-	-	1	-	-	1	-	1
Female	20	4	1	-	-	-	-	-	-	2	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	1	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	1
Asian or Pacific Islander 2/, both sexes	25	1	-	-	-	1	1	-	-	2	-	1
Male	13	-	-	-	-	1	1	-	-	2	-	-
Female	12	1	-	-	-	-	-	-	-	-	-	1
New Jersey												
All races, both sexes	71,371	651	86	60	82	290	449	494	545	888	1,462	2,074
Male	33,724	348	48	35	47	210	334	352	367	568	891	1,250
Female	37,647	303	38	25	35	80	115	142	178	320	571	824
White, both sexes	60,889	401	50	41	50	205	289	325	346	605	990	1,459
Male	28,425	215	27	25	29	140	206	229	251	411	625	920
Female	32,464	186	23	16	21	65	83	96	95	194	365	539
Black, both sexes	9,230	211	30	19	25	78	141	148	181	269	436	570
Male	4,631	118	17	10	12	65	112	108	106	148	239	306
Female	4,599	93	13	9	13	13	29	40	75	121	197	264
American Indian 1/, both sexes	73	2	-	-	-	-	1	1	-	-	-	2
Male	30	1	-	-	-	-	1	1	-	-	-	1
Female	43	1	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1,179	37	6	-	7	7	18	20	18	14	36	43
Male	638	14	4	-	6	5	15	14	10	9	27	23
Female	541	23	2	-	1	2	3	6	8	5	9	20

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
New Mexico												
All races, both sexes	14,298	179	45	20	42	130	205	170	204	259	413	516
Male	7,550	89	23	13	19	105	160	125	154	179	274	321
Female	6,748	90	22	7	23	25	45	45	50	80	139	195
White, both sexes	12,975	138	31	14	28	104	163	130	151	202	334	454
Male	6,798	67	15	10	13	84	130	92	114	140	220	284
Female	6,177	71	16	4	15	20	33	38	37	62	114	170
Black, both sexes	247	6	2	-	1	4	3	3	11	4	10	19
Male	136	2	2	-	-	4	1	3	11	2	7	9
Female	111	4	-	-	1	-	2	-	-	2	3	10
American Indian 1/, both sexes	1,029	34	12	5	13	20	39	35	42	53	65	41
Male	597	20	6	2	6	15	29	30	29	37	45	28
Female	432	14	6	3	7	5	10	5	13	16	20	13
Asian or Pacific Islander 2/, both sexes	47	1	-	1	-	2	-	2	-	-	4	2
Male	19	-	-	1	-	2	-	-	-	-	2	-
Female	28	1	-	-	-	-	-	2	-	-	2	2
New York												
All races, both sexes	152,681	1,518	213	156	202	605	898	907	1,139	1,840	3,053	4,463
Male	71,948	860	126	84	103	423	695	642	747	1,149	1,841	2,723
Female	80,733	658	87	72	99	182	203	265	392	691	1,212	1,740
White, both sexes	127,222	891	142	95	135	419	609	587	747	1,238	2,114	3,075
Male	59,603	516	85	47	74	286	475	421	501	799	1,357	1,906
Female	67,619	375	57	48	61	133	134	166	246	439	757	1,169
Black, both sexes	21,988	540	59	52	61	163	259	287	348	546	855	1,235
Male	10,468	291	35	29	28	125	205	193	216	315	439	715
Female	11,520	249	24	23	33	38	54	94	132	231	416	520
American Indian 1/, both sexes	242	10	1	2	-	1	2	2	7	5	5	13
Male	123	8	1	1	-	1	2	2	7	3	4	7
Female	119	2	-	1	-	-	-	-	-	2	1	6
Asian or Pacific Islander 2/, both sexes	3,229	77	11	7	6	22	28	31	37	51	79	140
Male	1,754	45	5	7	1	11	13	26	23	32	41	95
Female	1,475	32	6	-	5	11	15	5	14	19	38	45

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
North Carolina												
All races, both sexes	72,384	1,053	141	92	120	441	669	623	761	1,103	1,807	2,467
Male	35,772	582	80	52	72	308	515	456	498	697	1,091	1,526
Female	36,612	471	61	40	48	133	154	167	263	406	716	941
White, both sexes	55,972	547	85	64	81	314	443	407	498	733	1,176	1,644
Male	27,559	301	50	38	50	216	348	298	338	484	739	1,041
Female	28,413	246	35	26	31	98	95	109	160	249	437	603
Black, both sexes	15,502	463	50	24	36	107	210	199	238	341	591	779
Male	7,739	256	25	12	20	79	155	148	147	192	325	460
Female	7,763	207	25	12	16	28	55	51	91	149	266	319
American Indian 1/, both sexes	649	21	3	4	1	17	12	13	14	25	26	31
Male	336	11	3	2	1	11	9	8	6	19	16	18
Female	313	10	-	2	-	6	3	5	8	6	10	13
Asian or Pacific Islander 2/, both sexes	261	22	3	-	2	3	4	4	11	4	14	13
Male	138	14	2	-	1	2	3	2	7	2	11	7
Female	123	8	1	-	1	1	1	2	4	2	3	6
North Dakota												
All races, both sexes	5,601	46	10	6	11	29	31	36	38	37	83	132
Male	2,796	36	7	5	7	17	25	28	26	25	52	87
Female	2,805	10	3	1	4	12	6	8	12	12	31	45
White, both sexes	5,362	36	7	4	6	20	28	27	25	26	74	113
Male	2,679	27	6	4	4	12	22	22	17	19	46	75
Female	2,683	9	1	-	2	8	6	5	8	7	28	38
Black, both sexes	9	2	-	-	-	-	-	2	1	1	-	-
Male	7	2	-	-	-	-	-	1	1	1	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	223	8	3	1	5	9	3	7	12	10	9	19
Male	108	7	1	1	3	5	3	5	8	5	6	12
Female	115	1	2	-	2	4	-	2	4	5	3	7
Asian or Pacific Islander 2/, both sexes	7	-	-	1	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Ohio												
All races, both sexes	106,288	1,143	180	107	137	522	703	657	794	1,123	2,084	3,327
Male	51,307	647	105	66	72	378	535	480	527	708	1,293	2,034
Female	54,981	496	75	41	65	144	168	177	267	415	791	1,293
White, both sexes	93,790	739	148	71	106	418	555	507	627	879	1,677	2,645
Male	44,965	424	88	45	54	297	422	370	425	566	1,064	1,654
Female	48,825	315	60	26	52	121	133	137	202	313	613	991
Black, both sexes	12,183	390	30	33	30	102	144	147	164	236	396	660
Male	6,188	217	15	19	18	80	111	109	99	136	226	369
Female	5,995	173	15	14	12	22	33	38	65	100	170	291
American Indian 1/, both sexes	41	3	-	-	-	-	-	1	1	1	1	6
Male	24	2	-	-	-	-	-	-	1	1	-	3
Female	17	1	-	-	-	-	-	1	-	-	1	3
Asian or Pacific Islander 2/, both sexes	274	11	2	3	1	2	4	2	2	7	10	16
Male	130	4	2	2	-	1	2	1	2	5	3	8
Female	144	7	-	1	1	1	2	1	-	2	7	8
Oklahoma												
All races, both sexes	34,483	411	84	50	43	223	314	244	347	445	843	1,126
Male	17,094	227	39	28	26	149	230	187	231	278	514	696
Female	17,389	184	45	22	17	74	84	57	116	167	329	430
White, both sexes	30,159	283	64	35	32	182	237	173	255	326	662	879
Male	14,865	155	33	19	20	123	172	131	177	201	406	553
Female	15,294	128	31	16	12	59	65	42	78	125	256	326
Black, both sexes	2,171	81	11	6	6	18	41	35	38	56	79	117
Male	1,100	43	3	4	4	11	33	29	24	32	45	65
Female	1,071	38	8	2	2	7	8	6	14	24	34	52
American Indian 1/, both sexes	2,002	44	8	8	5	21	35	33	50	57	98	123
Male	1,061	27	2	4	2	13	24	27	26	41	60	74
Female	941	17	6	4	3	8	11	6	24	16	38	49
Asian or Pacific Islander 2/, both sexes	151	3	1	1	-	2	1	3	4	6	4	7
Male	68	2	1	1	-	2	1	-	4	4	3	4
Female	83	1	-	-	-	-	-	3	-	2	1	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Oregon												
All races, both sexes	30,313	251	51	40	36	131	220	171	222	342	603	891
Male	14,949	154	29	20	26	89	155	121	151	230	353	577
Female	15,364	97	22	20	10	42	65	50	71	112	250	314
White, both sexes	29,185	220	47	34	30	118	203	144	198	319	555	826
Male	14,353	136	28	17	20	80	145	105	135	215	330	539
Female	14,832	84	19	17	10	38	58	39	63	104	225	287
Black, both sexes	413	13	2	1	2	5	7	11	10	7	18	22
Male	217	10	1	-	2	4	5	7	8	5	7	16
Female	196	3	1	1	-	1	2	4	2	2	11	6
American Indian 1/, both sexes	306	9	2	3	2	4	7	9	7	12	19	28
Male	167	4	-	1	2	2	3	4	5	8	7	15
Female	139	5	2	2	-	2	4	5	2	4	12	13
Asian or Pacific Islander 2/, both sexes	409	9	-	2	2	4	3	7	7	4	11	15
Male	212	4	-	2	2	3	2	5	3	2	9	7
Female	197	5	-	-	-	1	1	2	4	2	2	8
Pennsylvania												
All races, both sexes	127,640	1,049	164	102	146	563	795	728	849	1,332	2,291	3,388
Male	60,638	612	98	52	86	404	633	545	572	838	1,434	2,083
Female	67,002	437	66	50	60	159	162	183	277	494	857	1,305
White, both sexes	115,402	714	122	83	108	448	633	532	630	1,049	1,853	2,705
Male	54,493	403	73	42	66	317	493	404	423	669	1,191	1,696
Female	60,909	311	49	41	42	131	140	128	207	380	662	1,009
Black, both sexes	11,613	307	39	18	35	108	149	187	201	267	411	660
Male	5,819	196	24	9	19	85	132	137	136	159	225	376
Female	5,794	111	15	9	16	23	17	50	65	108	186	284
American Indian 1/, both sexes	57	1	-	-	-	1	2	-	1	2	-	1
Male	33	1	-	-	-	-	1	-	1	1	-	-
Female	24	-	-	-	-	1	1	-	-	1	-	1
Asian or Pacific Islander 2/, both sexes	568	27	3	1	3	6	11	9	17	14	27	22
Male	293	12	1	1	1	2	7	4	12	9	18	11
Female	275	15	2	-	2	4	4	5	5	5	9	11

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Rhode Island												
All races, both sexes	9,769	68	13	2	6	39	52	42	55	95	156	280
Male	4,605	35	9	1	2	27	38	32	43	63	91	170
Female	5,164	33	4	1	4	12	14	10	12	32	65	110
White, both sexes	9,345	54	11	-	5	35	41	34	45	86	142	257
Male	4,376	26	7	-	2	23	27	27	36	58	84	152
Female	4,969	28	4	-	3	12	14	7	9	28	58	105
Black, both sexes	345	13	-	1	1	3	10	7	9	7	11	20
Male	191	9	-	1	-	3	10	5	6	5	6	16
Female	154	4	-	-	1	-	-	2	3	2	5	4
American Indian 1/, both sexes	24	-	1	-	-	-	-	-	1	1	2	-
Male	9	-	1	-	-	-	-	-	1	-	1	-
Female	15	-	-	-	-	-	-	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes	55	1	1	1	-	1	1	1	-	1	1	3
Male	29	-	1	-	-	1	1	-	-	-	-	2
Female	26	1	-	1	-	-	-	1	-	1	1	1
South Carolina												
All races, both sexes	37,276	525	79	42	73	252	367	372	423	578	1,002	1,484
Male	18,889	283	48	23	40	176	274	268	290	356	630	900
Female	18,387	242	31	19	33	76	93	104	133	222	372	584
White, both sexes	26,451	243	37	23	39	167	220	201	248	339	588	843
Male	13,405	133	21	16	21	114	174	148	177	211	394	537
Female	13,046	110	16	7	18	53	46	53	71	128	194	306
Black, both sexes	10,675	277	41	18	33	81	143	167	173	236	407	630
Male	5,419	147	27	7	19	60	98	116	112	143	231	358
Female	5,256	130	14	11	14	21	45	51	61	93	176	272
American Indian 1/, both sexes	44	-	-	-	-	3	1	1	1	1	4	3
Male	20	-	-	-	-	1	1	1	-	1	2	2
Female	24	-	-	-	-	2	-	-	1	-	2	1
Asian or Pacific Islander 2/, both sexes	106	5	1	1	1	1	3	3	1	2	3	8
Male	45	3	-	-	-	1	1	3	1	1	3	3
Female	61	2	1	1	1	-	2	-	-	1	-	5

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
South Dakota												
All races, both sexes	6,833	93	20	21	16	48	56	54	52	76	114	182
Male	3,424	50	9	9	10	32	38	36	37	46	72	110
Female	3,409	43	11	12	6	16	18	18	15	30	42	72
White, both sexes	6,227	64	11	13	10	26	30	39	26	57	80	139
Male	3,088	34	5	6	7	16	21	29	20	33	52	86
Female	3,139	30	6	7	3	10	9	10	6	24	28	53
Black, both sexes	18	-	-	-	-	-	-	-	-	-	4	1
Male	13	-	-	-	-	-	-	-	-	-	4	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	572	28	9	5	6	22	26	15	26	19	30	41
Male	318	15	4	2	3	16	17	7	17	13	16	23
Female	254	13	5	3	3	6	9	8	9	6	14	18
Asian or Pacific Islander 2/, both sexes	16	1	-	3	-	-	-	-	-	-	-	1
Male	5	1	-	1	-	-	-	-	-	-	-	-
Female	11	-	-	2	-	-	-	-	-	-	-	1
Tennessee												
All races, both sexes	55,829	687	105	54	92	380	517	530	603	871	1,447	2,092
Male	27,612	398	63	31	56	265	380	384	407	551	893	1,314
Female	28,217	289	42	23	36	115	137	146	196	320	554	778
White, both sexes	47,445	402	78	39	66	302	379	368	440	681	1,122	1,597
Male	23,398	246	45	22	41	208	284	278	292	436	698	1,015
Female	24,047	156	33	17	25	94	95	90	148	245	424	582
Black, both sexes	8,192	278	26	14	26	77	135	155	157	183	321	482
Male	4,110	148	17	8	15	56	95	100	110	112	192	291
Female	4,082	130	9	6	11	21	40	55	47	71	129	191
American Indian 1/, both sexes	30	3	-	-	-	-	-	-	1	3	-	3
Male	16	1	-	-	-	-	-	-	1	1	-	2
Female	14	2	-	-	-	-	-	-	-	2	-	1
Asian or Pacific Islander 2/, both sexes	162	4	1	1	-	1	3	7	5	4	4	10
Male	88	3	1	1	-	1	1	6	4	2	3	6
Female	74	1	-	-	-	-	2	1	1	2	1	4

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Texas												
All races, both sexes	152,870	2,407	472	279	354	1,118	1,592	1,620	1,833	2,433	4,128	5,756
Male	77,412	1,304	272	165	227	786	1,182	1,182	1,233	1,599	2,650	3,646
Female	75,458	1,103	200	114	127	332	410	438	600	834	1,478	2,110
White, both sexes	132,048	1,819	369	207	279	954	1,289	1,263	1,405	1,899	3,238	4,501
Male	66,799	986	212	127	184	674	970	944	971	1,268	2,110	2,895
Female	65,249	833	157	80	95	280	319	319	434	631	1,128	1,606
Black, both sexes	19,123	539	95	58	65	149	273	336	397	488	819	1,175
Male	9,750	299	56	30	37	100	189	221	242	302	495	709
Female	9,373	240	39	28	28	49	84	115	155	186	324	466
American Indian 1/, both sexes	169	5	-	3	-	3	5	3	6	6	9	13
Male	82	3	-	1	-	3	4	2	5	6	4	7
Female	87	2	-	2	-	-	1	1	1	-	5	6
Asian or Pacific Islander 2/, both sexes	1,530	44	8	11	10	12	25	18	25	40	62	67
Male	781	16	4	7	6	9	19	15	15	23	41	35
Female	749	28	4	4	4	3	6	3	10	17	21	32
Utah												
All races, both sexes	13,331	264	68	19	37	97	166	204	195	211	298	418
Male	6,759	149	35	10	26	66	128	140	145	131	191	250
Female	6,572	115	33	9	11	31	38	64	50	80	107	168
White, both sexes	12,991	250	64	19	35	93	155	192	183	203	280	403
Male	6,569	142	34	10	25	63	120	134	137	127	177	242
Female	6,422	108	30	9	10	30	35	58	46	76	103	161
Black, both sexes	78	10	-	-	-	-	-	2	4	1	1	3
Male	44	4	-	-	-	-	-	2	3	1	1	2
Female	34	6	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	115	1	1	-	1	3	8	6	6	5	12	7
Male	68	1	-	-	1	2	6	3	3	2	10	3
Female	47	-	1	-	-	1	2	3	3	3	2	4
Asian or Pacific Islander 2/, both sexes	147	3	3	-	1	1	3	4	2	2	5	5
Male	78	2	1	-	-	1	2	1	2	1	3	3
Female	69	1	2	-	1	-	1	3	-	1	2	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Vermont												
All races, both sexes	4,995	30	2	4	6	23	23	35	33	48	76	128
Male	2,423	18	2	-	3	14	12	27	21	29	51	82
Female	2,572	12	-	4	3	9	11	8	12	19	25	46
White, both sexes	4,978	30	2	3	6	22	23	35	32	47	75	128
Male	2,415	18	2	-	3	14	12	27	21	29	50	82
Female	2,563	12	-	3	3	8	11	8	11	18	25	46
Black, both sexes	10	-	-	1	-	-	-	-	1	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	1	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Virginia												
All races, both sexes	56,550	776	103	66	88	305	467	435	523	712	1,250	1,802
Male	27,697	435	54	41	54	208	354	300	358	422	734	1,148
Female	28,853	341	49	25	34	97	113	135	165	290	516	654
White, both sexes	43,918	430	59	41	55	217	301	279	339	440	843	1,196
Male	21,376	249	31	28	32	148	231	201	239	272	514	772
Female	22,542	181	28	13	23	69	70	78	100	168	329	424
Black, both sexes	11,814	325	41	25	29	84	159	145	167	258	376	567
Male	5,938	175	22	13	20	58	118	93	111	144	208	354
Female	5,876	150	19	12	9	26	41	52	56	114	168	213
American Indian 1/, both sexes	54	2	-	-	1	-	1	-	-	1	2	2
Male	27	2	-	-	-	-	1	-	-	1	1	2
Female	27	-	-	-	1	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	764	19	3	-	3	4	6	11	17	13	29	37
Male	356	9	1	-	2	2	4	6	8	5	11	20
Female	408	10	2	-	1	2	2	5	9	8	18	17

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Washington												
All races, both sexes	44,770	451	68	51	59	255	354	285	376	568	940	1,386
Male	22,371	258	38	31	34	176	256	207	247	386	572	844
Female	22,399	193	30	20	25	79	98	78	129	182	368	542
White, both sexes	41,720	359	55	42	46	207	289	232	311	492	821	1,234
Male	20,746	204	30	25	29	139	207	170	204	333	504	753
Female	20,974	155	25	17	17	68	82	62	107	159	317	481
Black, both sexes	1,079	43	6	1	5	15	24	23	21	27	51	61
Male	612	28	2	1	3	12	19	19	14	19	29	44
Female	467	15	4	-	2	3	5	4	7	8	22	17
American Indian 1/, both sexes	592	19	2	3	3	16	18	17	13	25	33	42
Male	316	12	2	1	2	9	12	10	7	17	18	22
Female	276	7	-	2	1	7	6	7	6	8	15	20
Asian or Pacific Islander 2/, both sexes	1,379	30	5	5	5	17	23	13	31	24	35	49
Male	697	14	4	4	-	16	18	8	22	17	21	25
Female	682	16	1	1	5	1	5	5	9	7	14	24
West Virginia												
All races, both sexes	20,793	158	26	26	31	110	157	139	183	241	433	608
Male	10,181	90	11	15	21	72	109	97	124	155	290	366
Female	10,612	68	15	11	10	38	48	42	59	86	143	242
White, both sexes	20,153	149	24	24	30	106	148	136	177	228	419	591
Male	9,899	85	9	15	21	71	101	94	118	147	281	353
Female	10,254	64	15	9	9	35	47	42	59	81	138	238
Black, both sexes	623	9	2	2	1	3	8	3	6	13	14	17
Male	278	5	2	-	-	1	7	3	6	8	9	13
Female	345	4	-	2	1	2	1	-	-	5	5	4
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	1	1	-	-	-	-	-
Male	4	-	-	-	-	-	1	-	-	-	-	-
Female	10	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Wisconsin												
All races, both sexes	45,600	420	66	47	60	232	331	271	311	465	768	1,189
Male	22,324	226	35	29	38	162	264	189	206	278	491	741
Female	23,276	194	31	18	22	70	67	82	105	187	277	448
White, both sexes	43,088	274	49	37	46	202	270	224	244	390	658	1,024
Male	21,030	153	26	22	27	144	223	158	157	235	424	658
Female	22,058	121	23	15	19	58	47	66	87	155	234	366
Black, both sexes	1,955	128	12	7	8	19	46	40	56	53	88	127
Male	1,011	65	5	4	6	11	31	26	40	31	52	66
Female	944	63	7	3	2	8	15	14	16	22	36	61
American Indian 1/, both sexes	341	7	2	-	5	5	7	4	6	14	18	28
Male	174	3	2	-	4	4	5	3	5	6	11	11
Female	167	4	-	-	1	1	2	1	1	8	7	17
Asian or Pacific Islander 2/, both sexes	216	11	3	3	1	6	8	3	5	8	4	10
Male	109	5	2	3	1	3	5	2	4	6	4	6
Female	107	6	1	-	-	3	3	1	1	2	-	4
Wyoming												
All races, both sexes	3,955	60	6	4	8	29	54	34	37	50	76	144
Male	1,977	35	4	2	2	18	41	25	28	32	41	101
Female	1,978	25	2	2	6	11	13	9	9	18	35	43
White, both sexes	3,835	53	4	4	8	25	46	28	31	47	71	132
Male	1,918	30	3	2	2	16	37	21	25	30	38	94
Female	1,917	23	1	2	6	9	9	7	6	17	33	38
Black, both sexes	28	-	-	-	-	1	1	1	2	-	1	1
Male	15	-	-	-	-	-	1	1	1	-	1	-
Female	13	-	-	-	-	1	-	-	1	-	-	1
American Indian 1/, both sexes	81	7	2	-	-	3	7	5	4	3	4	9
Male	40	5	1	-	-	2	3	3	2	2	2	6
Female	41	2	1	-	-	1	4	2	2	1	2	3
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
United States												
All races, both sexes	99,770	121,163	143,534	171,984	227,682	310,746	373,484	338,140	227,661	87,446	18,526	346
Male	62,649	73,400	84,632	98,455	124,436	160,308	173,361	133,667	70,168	19,820	2,995	274
Female	37,121	47,763	58,902	73,529	103,246	150,438	200,123	204,473	157,493	67,626	15,531	72
White, both sexes	76,538	96,434	117,737	143,232	195,030	273,291	336,369	309,335	208,415	79,849	16,141	269
Male	48,805	58,771	69,804	82,606	107,517	142,168	157,174	122,938	64,196	17,919	2,536	218
Female	27,733	37,663	47,933	60,626	87,513	131,123	179,195	186,397	144,219	61,930	13,605	51
Black, both sexes	20,287	21,336	22,154	24,310	27,372	31,111	30,268	23,184	15,748	6,320	2,048	64
Male	12,116	12,646	12,706	13,384	14,171	14,830	12,865	8,253	4,520	1,412	363	46
Female	8,171	8,690	9,448	10,926	13,201	16,281	17,403	14,931	11,228	4,908	1,685	18
American Indian 1/, both sexes	906	1,006	1,067	1,177	1,152	1,204	1,171	809	453	187	54	10
Male	537	597	615	647	578	632	514	313	148	49	14	8
Female	369	409	452	530	574	572	657	496	305	138	40	2
Asian or Pacific Islander 2/, both sexes	2,039	2,387	2,576	3,265	4,128	5,140	5,676	4,812	3,045	1,090	283	3
Male	1,191	1,386	1,507	1,818	2,170	2,678	2,808	2,163	1,304	440	82	2
Female	848	1,001	1,069	1,447	1,958	2,462	2,868	2,649	1,741	650	201	1
Alabama												
All races, both sexes	2,109	2,739	3,030	3,756	4,675	6,072	6,724	5,665	3,775	1,367	286	-
Male	1,311	1,661	1,817	2,203	2,515	3,105	2,989	2,062	1,033	264	47	-
Female	798	1,078	1,213	1,553	2,160	2,967	3,735	3,603	2,742	1,103	239	-
White, both sexes	1,353	1,917	2,244	2,864	3,577	4,780	5,494	4,630	3,007	1,049	187	-
Male	854	1,142	1,362	1,702	1,987	2,508	2,478	1,698	825	188	26	-
Female	499	775	882	1,162	1,590	2,272	3,016	2,932	2,182	861	161	-
Black, both sexes	744	809	770	877	1,091	1,284	1,218	1,029	766	315	98	-
Male	451	514	444	491	526	595	509	361	207	76	21	-
Female	293	295	326	386	565	689	709	668	559	239	77	-
American Indian 1/, both sexes	3	8	8	6	2	6	7	-	1	1	1	-
Male	1	3	7	4	1	1	1	-	-	-	-	-
Female	2	5	1	2	1	5	6	-	1	1	1	-
Asian or Pacific Islander 2/, both sexes	9	5	8	9	5	2	5	6	1	2	-	-
Male	5	2	4	6	1	1	1	3	1	-	-	-
Female	4	3	4	3	4	1	4	3	-	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Alaska												
All races, both sexes	197	227	225	251	286	315	317	226	136	45	10	-
Male	130	150	140	153	170	174	157	86	51	15	2	-
Female	67	77	85	98	116	141	160	140	85	30	8	-
White, both sexes	139	165	165	181	224	236	229	173	104	32	8	-
Male	95	112	103	111	136	129	115	70	42	7	2	-
Female	44	53	62	70	88	107	114	103	62	25	6	-
Black, both sexes	5	13	6	5	3	8	8	7	3	1	-	-
Male	4	5	3	5	2	6	5	3	-	-	-	-
Female	1	8	3	-	1	2	3	4	3	1	-	-
American Indian 1/, both sexes	43	37	52	57	50	63	71	40	23	10	2	-
Male	23	24	34	34	28	34	31	13	6	6	-	-
Female	20	13	18	23	22	29	40	27	17	4	2	-
Asian or Pacific Islander 2/, both sexes	10	12	2	8	9	8	9	6	6	2	-	-
Male	8	9	-	3	4	5	6	-	3	2	-	-
Female	2	3	2	5	5	3	3	6	3	-	-	-
Arizona												
All races, both sexes	1,767	2,167	2,644	3,264	4,252	5,496	6,647	5,839	3,676	1,352	251	76
Male	1,118	1,326	1,580	1,888	2,398	2,909	3,310	2,549	1,234	402	46	68
Female	649	841	1,064	1,376	1,854	2,587	3,337	3,290	2,442	950	205	8
White, both sexes	1,575	1,940	2,392	3,015	3,990	5,265	6,402	5,643	3,556	1,300	240	65
Male	997	1,195	1,441	1,760	2,275	2,795	3,204	2,465	1,194	388	45	59
Female	578	745	951	1,255	1,715	2,470	3,198	3,178	2,362	912	195	6
Black, both sexes	66	85	103	102	115	83	98	88	43	23	3	2
Male	39	53	62	45	60	41	45	31	10	7	-	2
Female	27	32	41	57	55	42	53	57	33	16	3	-
American Indian 1/, both sexes	104	120	123	120	124	120	111	76	52	24	6	9
Male	72	71	65	69	49	61	50	39	19	3	-	7
Female	32	49	58	51	75	59	61	37	33	21	6	2
Asian or Pacific Islander 2/, both sexes	22	22	26	27	23	28	36	32	25	5	2	-
Male	10	7	12	14	14	12	11	14	11	4	1	-
Female	12	15	14	13	9	16	25	18	14	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Arkansas												
All races, both sexes	1,160	1,486	1,927	2,172	2,758	3,440	4,016	3,550	2,431	950	195	-
Male	701	916	1,153	1,249	1,496	1,788	1,835	1,358	746	231	28	-
Female	459	570	774	923	1,262	1,652	2,181	2,192	1,685	719	167	-
White, both sexes	886	1,201	1,625	1,881	2,410	3,051	3,629	3,222	2,151	821	153	-
Male	533	745	964	1,084	1,325	1,603	1,677	1,237	662	193	20	-
Female	353	456	661	797	1,085	1,448	1,952	1,985	1,489	628	133	-
Black, both sexes	269	277	292	287	338	382	379	326	271	125	42	-
Male	165	165	181	164	167	183	155	121	80	38	8	-
Female	104	112	111	123	171	199	224	205	191	87	34	-
American Indian 1/, both sexes	3	5	5	2	3	4	6	2	6	1	-	-
Male	1	4	3	-	2	1	3	-	3	-	-	-
Female	2	1	2	2	1	3	3	2	3	1	-	-
Asian or Pacific Islander 2/, both sexes	2	3	5	2	7	3	2	-	3	3	-	-
Male	2	2	5	1	2	1	-	-	1	-	-	-
Female	-	1	-	1	5	2	2	-	2	3	-	-
California												
All races, both sexes	10,328	12,141	13,318	15,770	21,152	29,029	36,697	32,561	21,889	8,566	1,918	57
Male	6,397	7,360	7,830	8,927	11,498	14,965	17,182	13,631	7,486	2,172	383	46
Female	3,931	4,781	5,488	6,843	9,654	14,064	19,515	18,930	14,403	6,394	1,535	11
White, both sexes	8,000	9,715	10,812	12,732	17,605	24,673	32,042	28,727	19,504	7,661	1,650	53
Male	5,054	5,912	6,383	7,202	9,622	12,710	15,005	12,030	6,570	1,873	321	43
Female	2,946	3,803	4,429	5,530	7,983	11,963	17,037	16,697	12,934	5,788	1,329	10
Black, both sexes	1,407	1,413	1,413	1,665	1,722	1,916	1,967	1,569	995	392	141	2
Male	816	840	806	980	915	967	842	567	299	87	27	2
Female	591	573	607	685	807	949	1,125	1,002	696	305	114	-
American Indian 1/, both sexes	108	94	110	120	125	132	134	78	47	19	3	-
Male	57	50	61	63	57	71	63	30	12	8	1	-
Female	51	44	49	57	68	61	71	48	35	11	2	-
Asian or Pacific Islander 2/, both sexes	813	919	983	1,253	1,700	2,308	2,554	2,187	1,343	494	124	2
Male	470	558	580	682	904	1,217	1,272	1,004	605	204	34	1
Female	343	361	403	571	796	1,091	1,282	1,183	738	290	90	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Colorado												
All races, both sexes	1,310	1,476	1,710	1,893	2,470	3,510	4,357	3,824	2,535	1,031	179	2
Male	806	912	1,017	1,078	1,368	1,780	2,071	1,584	814	235	31	2
Female	504	564	693	815	1,102	1,730	2,286	2,240	1,721	796	148	-
White, both sexes	1,221	1,382	1,612	1,774	2,349	3,342	4,196	3,724	2,478	1,008	171	2
Male	763	850	960	1,015	1,302	1,690	1,984	1,542	804	231	29	2
Female	458	532	652	759	1,047	1,652	2,212	2,182	1,674	777	142	-
Black, both sexes	63	68	72	80	73	102	91	63	43	17	6	-
Male	30	45	45	42	46	52	53	24	8	3	-	-
Female	33	23	27	38	27	50	38	39	35	14	6	-
American Indian 1/, both sexes	8	7	5	15	16	15	13	9	7	1	2	-
Male	5	5	3	10	8	6	6	4	1	1	2	-
Female	3	2	2	5	8	9	7	5	6	-	-	-
Asian or Pacific Islander 2/, both sexes	18	19	21	24	32	51	57	28	7	5	-	-
Male	8	12	9	11	12	32	28	14	1	-	-	-
Female	10	7	12	13	20	19	29	14	6	5	-	-
Connecticut												
All races, both sexes	949	1,213	1,518	1,755	2,473	3,671	5,033	4,902	3,420	1,394	299	3
Male	579	724	879	974	1,367	1,952	2,388	1,979	1,038	311	40	2
Female	370	489	639	781	1,106	1,719	2,645	2,923	2,382	1,083	259	1
White, both sexes	798	1,057	1,338	1,555	2,266	3,430	4,773	4,728	3,319	1,346	289	3
Male	488	629	776	867	1,256	1,833	2,277	1,917	1,015	300	40	2
Female	310	428	562	688	1,010	1,597	2,496	2,811	2,304	1,046	249	1
Black, both sexes	134	143	162	188	178	221	244	160	87	44	10	-
Male	79	86	95	100	94	111	102	55	21	10	-	-
Female	55	57	67	88	84	110	142	105	66	34	10	-
American Indian 1/, both sexes	5	2	6	2	3	6	6	2	2	1	-	-
Male	4	2	1	2	2	2	5	2	-	-	-	-
Female	1	-	5	-	1	4	1	-	2	1	-	-
Asian or Pacific Islander 2/, both sexes	12	11	12	10	26	14	10	12	12	3	-	-
Male	8	7	7	5	15	6	4	5	2	1	-	-
Female	4	4	5	5	11	8	6	7	10	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Delaware												
All races, both sexes	296	368	440	541	772	956	1,088	978	567	224	44	3
Male	182	227	262	326	437	494	514	407	177	55	7	3
Female	114	141	178	215	335	462	574	571	390	169	37	-
White, both sexes	202	276	356	446	640	817	942	876	502	197	35	1
Male	131	177	206	266	363	435	451	369	159	47	7	1
Female	71	99	150	180	277	382	491	507	343	150	28	-
Black, both sexes	90	87	82	87	129	133	138	97	63	27	9	2
Male	50	48	55	55	72	56	60	37	16	8	-	2
Female	40	39	27	32	57	77	78	60	47	19	9	-
American Indian 1/, both sexes	-	3	1	4	-	3	4	1	1	-	-	-
Male	-	1	1	2	-	2	1	-	1	-	-	-
Female	-	2	-	2	-	1	3	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	2	1	4	3	3	4	4	1	-	-	-
Male	1	1	-	3	2	1	2	1	1	-	-	-
Female	3	1	1	1	1	2	2	3	-	-	-	-
Dist. of Columbia												
All races, both sexes	348	367	355	385	501	580	688	626	414	173	34	2
Male	220	219	223	226	278	269	312	235	131	42	6	2
Female	128	148	132	159	223	311	376	391	283	131	28	-
White, both sexes	36	70	47	64	81	104	180	219	148	65	7	1
Male	23	44	32	40	49	61	97	88	49	17	3	1
Female	13	26	15	24	32	43	83	131	99	48	4	-
Black, both sexes	310	294	306	317	417	468	501	398	260	107	27	1
Male	197	172	190	182	229	204	214	142	77	25	3	1
Female	113	122	116	135	188	264	287	256	183	82	24	-
American Indian 1/, both sexes	-	1	-	-	-	-	2	2	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	2	4	3	8	5	7	6	1	-	-
Male	-	2	1	4	-	4	1	4	5	-	-	-
Female	2	-	1	-	3	4	4	3	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Florida												
All races, both sexes	6,491	7,904	9,620	11,968	16,301	22,483	27,268	24,804	16,729	6,252	1,279	47
Male	4,148	4,928	5,878	7,075	9,257	12,066	13,424	10,772	5,770	1,637	240	37
Female	2,343	2,976	3,742	4,893	7,044	10,417	13,844	14,032	10,959	4,615	1,039	10
White, both sexes	5,210	6,490	8,150	10,196	14,524	20,589	25,508	23,374	15,825	5,925	1,168	37
Male	3,419	4,109	4,999	6,065	8,314	11,113	12,647	10,238	5,498	1,563	218	31
Female	1,791	2,381	3,151	4,131	6,210	9,476	12,861	13,136	10,327	4,362	950	6
Black, both sexes	1,208	1,334	1,377	1,692	1,675	1,779	1,668	1,360	868	313	108	10
Male	688	776	834	964	886	905	739	507	258	68	21	6
Female	520	558	543	728	789	874	929	853	610	245	87	4
American Indian 1/, both sexes	17	14	12	10	16	17	16	9	8	-	-	-
Male	9	10	7	5	9	9	5	2	3	-	-	-
Female	8	4	5	5	7	8	11	7	5	-	-	-
Asian or Pacific Islander 2/, both sexes	56	66	81	70	86	98	76	61	28	14	3	-
Male	32	33	38	41	48	39	33	25	11	6	1	-
Female	24	33	43	29	38	59	43	36	17	8	2	-
Georgia												
All races, both sexes	3,546	4,188	4,711	5,245	6,480	7,923	8,955	7,563	5,052	1,756	387	5
Male	2,243	2,550	2,799	3,058	3,497	3,921	3,924	2,688	1,356	347	54	3
Female	1,303	1,638	1,912	2,187	2,983	4,002	5,031	4,875	3,696	1,409	333	2
White, both sexes	2,138	2,688	3,202	3,800	4,762	6,173	7,223	6,172	4,055	1,389	265	-
Male	1,381	1,627	1,925	2,291	2,617	3,111	3,196	2,218	1,096	270	31	-
Female	757	1,061	1,277	1,509	2,145	3,062	4,027	3,954	2,959	1,119	234	-
Black, both sexes	1,371	1,470	1,469	1,408	1,669	1,707	1,693	1,358	981	363	120	5
Male	839	900	848	747	854	792	707	458	254	74	22	3
Female	532	570	621	661	815	915	986	900	727	289	98	2
American Indian 1/, both sexes	10	6	10	12	11	11	10	9	5	1	-	-
Male	6	5	5	7	6	4	5	2	1	1	-	-
Female	4	1	5	5	5	7	5	7	4	-	-	-
Asian or Pacific Islander 2/, both sexes	27	24	30	25	38	32	29	24	11	3	2	-
Male	17	18	21	13	20	14	16	10	5	2	1	-
Female	10	6	9	12	18	18	13	14	6	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Hawaii												
All races, both sexes	410	465	490	597	815	1,132	1,412	1,339	920	376	90	-
Male	266	301	300	365	465	646	763	652	401	145	22	-
Female	144	164	190	232	350	486	649	687	519	231	68	-
White, both sexes	146	152	156	165	228	301	358	379	217	83	16	-
Male	102	105	100	103	135	171	193	189	79	22	1	-
Female	44	47	56	62	93	130	165	190	138	61	15	-
Black, both sexes	10	6	4	5	3	3	7	5	3	2	-	-
Male	6	5	4	2	3	1	5	2	3	1	-	-
Female	4	1	-	3	-	2	2	3	-	1	-	-
American Indian 1/, both sexes	1	-	-	2	2	1	1	-	2	-	-	-
Male	-	-	-	-	2	-	1	-	-	-	-	-
Female	1	-	-	2	-	1	-	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	253	307	330	425	582	827	1,046	955	698	291	74	-
Male	158	191	196	260	325	474	564	461	319	122	21	-
Female	95	116	134	165	257	353	482	494	379	169	53	-
Idaho												
All races, both sexes	399	436	570	710	919	1,269	1,536	1,525	1,020	383	74	-
Male	268	259	337	412	523	681	708	626	346	107	11	-
Female	131	177	233	298	396	588	828	899	674	276	63	-
White, both sexes	389	426	561	703	910	1,251	1,512	1,512	1,016	379	73	-
Male	263	254	331	409	519	671	694	621	345	105	11	-
Female	126	172	230	294	391	580	818	891	671	274	62	-
Black, both sexes	3	1	-	-	-	3	3	-	1	-	-	-
Male	3	1	-	-	-	2	2	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	-	-	-
American Indian 1/, both sexes	5	7	9	6	6	8	11	7	2	2	1	-
Male	2	3	6	2	3	4	4	2	1	2	-	-
Female	3	4	3	4	3	4	7	5	1	-	1	-
Asian or Pacific Islander 2/, both sexes	2	2	-	1	3	7	10	6	1	2	-	-
Male	-	1	-	1	1	4	8	3	-	-	-	-
Female	2	1	-	-	2	3	2	3	1	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Illinois												
All races, both sexes	4,132	4,887	6,074	7,204	9,546	12,961	15,858	15,160	10,429	4,047	897	5
Male	2,688	2,878	3,523	4,001	5,080	6,592	7,305	5,802	3,092	835	137	4
Female	1,444	2,009	2,551	3,203	4,466	6,369	8,553	9,358	7,337	3,212	760	1
White, both sexes	2,939	3,623	4,669	5,709	7,894	11,106	14,028	13,903	9,556	3,688	780	-
Male	1,949	2,153	2,757	3,206	4,271	5,743	6,528	5,359	2,834	753	116	-
Female	990	1,470	1,912	2,503	3,623	5,363	7,500	8,544	6,722	2,935	664	-
Black, both sexes	1,134	1,166	1,327	1,372	1,532	1,710	1,670	1,123	804	331	112	4
Male	701	668	721	714	749	779	706	387	234	76	21	3
Female	433	498	606	658	783	931	964	736	570	255	91	1
American Indian 1/, both sexes	3	7	4	6	5	6	5	3	-	4	1	-
Male	1	7	2	2	2	1	4	2	-	-	-	-
Female	2	-	2	4	3	5	1	1	-	4	1	-
Asian or Pacific Islander 2/, both sexes	56	91	74	117	115	139	155	131	69	24	4	1
Male	37	50	43	79	58	69	67	54	24	6	-	1
Female	19	41	31	38	57	70	88	77	45	18	4	-
Indiana												
All races, both sexes	2,168	2,644	3,370	3,944	5,302	7,358	8,540	7,547	4,973	1,910	384	1
Male	1,332	1,569	1,987	2,231	2,871	3,720	3,925	2,876	1,477	402	54	1
Female	836	1,075	1,383	1,713	2,431	3,638	4,615	4,671	3,496	1,508	330	-
White, both sexes	1,846	2,351	3,041	3,593	4,873	6,866	8,034	7,195	4,748	1,808	349	1
Male	1,140	1,390	1,806	2,039	2,647	3,476	3,686	2,743	1,409	384	51	1
Female	706	961	1,235	1,554	2,226	3,390	4,348	4,452	3,339	1,424	298	-
Black, both sexes	313	282	313	338	408	475	482	344	216	102	35	-
Male	187	175	176	186	214	238	228	130	65	18	3	-
Female	126	107	137	152	194	237	254	214	151	84	32	-
American Indian 1/, both sexes	2	-	2	1	6	4	8	-	1	-	-	-
Male	1	-	-	-	3	1	3	-	-	-	-	-
Female	1	-	2	1	3	3	5	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	7	11	14	12	15	13	16	8	8	-	-	-
Male	4	4	5	6	7	5	8	3	3	-	-	-
Female	3	7	9	6	8	8	8	5	5	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Iowa												
All races, both sexes	863	1,034	1,312	1,752	2,283	3,480	4,372	4,610	3,302	1,516	356	-
Male	556	644	779	1,003	1,237	1,903	2,170	1,882	1,026	342	54	-
Female	307	390	533	749	1,046	1,577	2,202	2,728	2,276	1,174	302	-
White, both sexes	812	1,004	1,275	1,691	2,244	3,424	4,319	4,578	3,272	1,505	355	-
Male	526	627	755	972	1,216	1,878	2,146	1,873	1,015	339	53	-
Female	286	377	520	719	1,028	1,546	2,173	2,705	2,257	1,166	302	-
Black, both sexes	44	24	26	46	33	44	36	26	26	10	1	-
Male	25	12	19	26	17	17	17	7	10	2	1	-
Female	19	12	7	20	16	27	19	19	16	8	-	-
American Indian 1/, both sexes	4	-	5	7	1	2	5	1	-	-	-	-
Male	3	-	2	1	1	2	1	1	-	-	-	-
Female	1	-	3	6	-	-	4	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	6	6	8	5	10	12	5	4	1	-	-
Male	2	5	3	4	3	6	6	1	1	1	-	-
Female	1	1	3	4	2	4	6	4	3	-	-	-
Kansas												
All races, both sexes	916	1,012	1,246	1,509	2,064	3,019	3,843	3,690	2,601	1,162	291	-
Male	587	604	755	830	1,149	1,571	1,772	1,405	775	267	42	-
Female	329	408	491	679	915	1,448	2,071	2,285	1,826	895	249	-
White, both sexes	794	905	1,147	1,389	1,929	2,849	3,693	3,535	2,522	1,130	277	-
Male	521	548	698	765	1,065	1,486	1,708	1,349	747	260	40	-
Female	273	357	449	624	864	1,363	1,985	2,186	1,775	870	237	-
Black, both sexes	97	87	79	97	116	144	129	129	71	28	14	-
Male	57	44	47	54	75	71	53	46	25	7	2	-
Female	40	43	32	43	41	73	76	83	46	21	12	-
American Indian 1/, both sexes	16	11	12	13	14	15	11	17	4	3	-	-
Male	7	8	6	4	7	11	7	6	-	-	-	-
Female	9	3	6	9	7	4	4	11	4	3	-	-
Asian or Pacific Islander 2/, both sexes	9	9	8	10	5	11	10	9	4	1	-	-
Male	2	4	4	7	2	3	4	4	3	-	-	-
Female	7	5	4	3	3	8	6	5	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Kentucky												
All races, both sexes	1,708	2,172	2,607	3,242	4,072	5,115	5,538	4,832	3,175	1,102	215	2
Male	1,089	1,278	1,504	1,870	2,240	2,607	2,430	1,801	906	242	24	2
Female	619	894	1,103	1,372	1,832	2,508	3,108	3,031	2,269	860	191	-
White, both sexes	1,517	2,006	2,401	3,036	3,805	4,800	5,254	4,610	3,004	1,036	192	2
Male	964	1,193	1,400	1,764	2,097	2,452	2,313	1,728	851	228	22	2
Female	553	813	1,001	1,272	1,708	2,348	2,941	2,882	2,153	808	170	-
Black, both sexes	187	160	197	200	256	305	278	219	170	65	22	-
Male	122	82	97	101	135	153	117	73	55	14	2	-
Female	65	78	100	99	121	152	161	146	115	51	20	-
American Indian 1/, both sexes	1	1	2	1	2	2	1	-	-	1	-	-
Male	1	1	2	1	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	3	5	7	5	9	8	5	3	1	-	1	-
Male	2	2	5	4	6	2	-	-	-	-	-	-
Female	1	3	2	1	3	6	5	3	1	-	1	-
Louisiana												
All races, both sexes	2,113	2,445	2,854	3,324	4,129	5,454	5,868	4,805	3,209	1,208	260	1
Male	1,327	1,451	1,692	1,895	2,251	2,736	2,633	1,783	965	251	45	1
Female	786	994	1,162	1,429	1,878	2,718	3,235	3,022	2,244	957	215	-
White, both sexes	1,168	1,438	1,762	2,176	2,880	4,003	4,527	3,787	2,494	915	164	1
Male	743	854	1,027	1,244	1,598	2,041	2,043	1,403	738	181	21	1
Female	425	584	735	932	1,282	1,962	2,484	2,384	1,756	734	143	-
Black, both sexes	930	983	1,079	1,130	1,224	1,436	1,325	1,001	696	288	94	-
Male	576	583	654	644	640	690	579	370	222	66	23	-
Female	354	400	425	486	584	746	746	631	474	222	71	-
American Indian 1/, both sexes	1	6	10	9	4	6	4	4	5	1	-	-
Male	-	5	9	2	2	3	3	1	-	-	-	-
Female	1	1	1	7	2	3	1	3	5	1	-	-
Asian or Pacific Islander 2/, both sexes	14	18	3	9	21	9	12	13	14	4	2	-
Male	8	9	2	5	11	2	8	9	5	4	1	-
Female	6	9	1	4	10	7	4	4	9	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Maine												
All races, both sexes	409	571	660	897	1,194	1,663	2,063	1,979	1,308	546	108	-
Male	260	335	398	518	626	834	977	763	368	111	19	-
Female	149	236	262	379	568	829	1,086	1,216	940	435	89	-
White, both sexes	403	560	652	885	1,183	1,652	2,046	1,972	1,304	546	107	-
Male	255	329	393	510	621	826	971	761	367	111	19	-
Female	148	231	259	375	562	826	1,075	1,211	937	435	88	-
Black, both sexes	3	5	2	1	4	4	4	1	1	-	1	-
Male	3	3	2	1	2	3	2	-	1	-	-	-
Female	-	2	-	-	2	1	2	1	-	-	1	-
American Indian 1/, both sexes	1	4	2	7	6	7	8	4	2	-	-	-
Male	-	2	2	5	2	5	4	2	-	-	-	-
Female	1	2	-	2	4	2	4	2	2	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	4	4	1	-	5	2	1	-	-	-
Male	2	1	1	2	1	-	-	-	-	-	-	-
Female	-	1	3	2	-	-	5	2	1	-	-	-
Maryland												
All races, both sexes	2,110	2,304	2,676	3,090	4,048	5,328	6,734	5,675	3,639	1,439	267	4
Male	1,266	1,364	1,579	1,715	2,173	2,690	3,073	2,152	1,082	321	31	3
Female	844	940	1,097	1,375	1,875	2,638	3,661	3,523	2,557	1,118	236	1
White, both sexes	1,185	1,360	1,717	2,022	2,878	4,085	5,424	4,720	3,027	1,195	217	3
Male	731	827	1,023	1,138	1,571	2,089	2,522	1,810	931	266	25	2
Female	454	533	694	884	1,307	1,996	2,902	2,910	2,096	929	192	1
Black, both sexes	886	892	893	1,015	1,098	1,174	1,222	863	571	231	47	1
Male	514	515	513	547	568	564	504	307	142	51	6	1
Female	372	377	380	468	530	610	718	556	429	180	41	-
American Indian 1/, both sexes	6	3	4	5	11	7	7	11	3	1	1	-
Male	4	2	3	3	8	2	3	2	1	-	-	-
Female	2	1	1	2	3	5	4	9	2	1	1	-
Asian or Pacific Islander 2/, both sexes	33	49	62	48	61	62	81	81	38	12	2	-
Male	17	20	40	27	26	35	44	33	8	4	-	-
Female	16	29	22	21	35	27	37	48	30	8	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Massachusetts												
All races, both sexes	1,830	2,321	2,742	3,250	4,891	7,030	9,328	9,099	6,180	2,549	516	-
Male	1,167	1,441	1,609	1,864	2,591	3,533	4,243	3,494	1,753	533	62	-
Female	663	880	1,133	1,386	2,300	3,497	5,085	5,605	4,427	2,016	454	-
White, both sexes	1,668	2,129	2,530	3,025	4,595	6,717	8,999	8,843	6,010	2,497	499	-
Male	1,072	1,310	1,491	1,733	2,420	3,377	4,088	3,392	1,700	521	58	-
Female	596	819	1,039	1,292	2,175	3,340	4,911	5,451	4,310	1,976	441	-
Black, both sexes	135	166	178	178	229	247	240	193	124	39	15	-
Male	79	112	97	101	137	122	113	76	41	10	4	-
Female	56	54	81	77	92	125	127	117	83	29	11	-
American Indian 1/, both sexes	-	4	4	7	8	7	4	5	2	4	-	-
Male	-	3	3	5	3	4	1	2	1	-	-	-
Female	-	1	1	2	5	3	3	3	1	4	-	-
Asian or Pacific Islander 2/, both sexes	27	22	30	40	59	59	85	58	44	9	2	-
Male	16	16	18	25	31	30	41	24	11	2	-	-
Female	11	6	12	15	28	29	44	34	33	7	2	-
Michigan												
All races, both sexes	3,678	4,310	4,999	6,018	7,982	11,438	13,365	12,244	7,978	2,947	653	-
Male	2,352	2,591	2,898	3,334	4,295	5,919	6,175	4,820	2,417	666	114	-
Female	1,326	1,719	2,101	2,684	3,687	5,519	7,190	7,424	5,561	2,281	539	-
White, both sexes	2,669	3,243	4,068	5,019	6,756	9,989	11,942	11,224	7,294	2,699	567	-
Male	1,742	1,979	2,355	2,806	3,648	5,207	5,518	4,420	2,193	600	98	-
Female	927	1,264	1,713	2,213	3,108	4,782	6,424	6,804	5,101	2,099	469	-
Black, both sexes	949	997	867	918	1,135	1,356	1,334	958	654	237	85	-
Male	571	568	501	482	600	665	614	373	212	63	16	-
Female	378	429	366	436	535	691	720	585	442	174	69	-
American Indian 1/, both sexes	29	33	38	44	52	48	48	29	9	5	1	-
Male	18	22	28	23	28	27	22	10	4	-	-	-
Female	11	11	10	21	24	21	26	19	5	5	1	-
Asian or Pacific Islander 2/, both sexes	31	37	26	37	39	45	41	33	21	6	-	-
Male	21	22	14	23	19	20	21	17	8	3	-	-
Female	10	15	12	14	20	25	20	16	13	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Minnesota												
All races, both sexes	1,264	1,558	1,876	2,311	3,102	4,549	5,868	6,198	4,609	1,917	438	-
Male	803	942	1,086	1,316	1,739	2,450	2,875	2,598	1,492	412	62	-
Female	461	616	790	995	1,363	2,099	2,993	3,600	3,117	1,505	376	-
White, both sexes	1,137	1,447	1,769	2,178	2,984	4,416	5,748	6,117	4,559	1,899	431	-
Male	727	875	1,017	1,246	1,676	2,384	2,818	2,568	1,469	408	61	-
Female	410	572	752	932	1,308	2,032	2,930	3,549	3,090	1,491	370	-
Black, both sexes	69	64	49	51	54	65	58	36	27	11	3	-
Male	36	39	31	22	27	28	27	13	12	3	1	-
Female	33	25	18	29	27	37	31	23	15	8	2	-
American Indian 1/, both sexes	42	31	31	45	32	31	18	15	7	4	1	-
Male	31	18	19	28	19	20	10	5	5	-	-	-
Female	11	13	12	17	13	11	8	10	2	4	1	-
Asian or Pacific Islander 2/, both sexes	16	16	27	37	32	37	44	30	16	3	3	-
Male	9	10	19	20	17	18	20	12	6	1	-	-
Female	7	6	8	17	15	19	24	18	10	2	3	-
Mississippi												
All races, both sexes	1,347	1,687	1,964	2,293	2,798	3,452	3,767	3,180	2,199	825	236	1
Male	841	1,066	1,195	1,313	1,563	1,745	1,730	1,151	647	189	44	1
Female	506	621	769	980	1,235	1,707	2,037	2,029	1,552	636	192	-
White, both sexes	739	1,023	1,270	1,553	1,966	2,557	2,859	2,418	1,608	578	126	1
Male	493	652	785	924	1,154	1,343	1,331	871	435	115	19	1
Female	246	371	485	629	812	1,214	1,528	1,547	1,173	463	107	-
Black, both sexes	591	651	683	730	819	882	901	757	588	247	108	-
Male	341	409	406	387	404	393	395	280	209	74	23	-
Female	250	242	277	343	415	489	506	477	379	173	85	-
American Indian 1/, both sexes	10	6	7	4	3	4	2	1	2	-	-	-
Male	5	2	4	1	2	3	2	-	2	-	-	-
Female	5	4	3	3	1	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	7	4	6	10	9	5	4	1	-	2	-
Male	2	3	-	1	3	6	2	-	1	-	2	-
Female	5	4	4	5	7	3	3	4	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Missouri												
All races, both sexes	2,014	2,598	3,327	3,898	5,286	7,079	8,260	7,505	5,481	2,118	447	4
Male	1,280	1,568	1,932	2,267	2,924	3,692	3,790	2,859	1,621	445	63	3
Female	734	1,030	1,395	1,631	2,362	3,387	4,470	4,646	3,860	1,673	384	1
White, both sexes	1,616	2,179	2,882	3,440	4,697	6,401	7,632	7,051	5,109	1,987	400	3
Male	1,029	1,327	1,680	2,005	2,638	3,383	3,527	2,682	1,517	413	53	2
Female	587	852	1,202	1,435	2,059	3,018	4,105	4,369	3,592	1,574	347	1
Black, both sexes	382	400	422	434	556	638	599	435	358	125	47	1
Male	242	231	238	251	271	294	252	172	95	30	10	1
Female	140	169	184	183	285	344	347	263	263	95	37	-
American Indian 1/, both sexes	6	7	10	8	10	13	11	5	3	4	-	-
Male	4	5	7	4	4	7	5	1	2	1	-	-
Female	2	2	3	4	6	6	6	4	1	3	-	-
Asian or Pacific Islander 2/, both sexes	10	12	13	16	23	27	18	14	11	2	-	-
Male	5	5	7	7	11	8	6	4	7	1	-	-
Female	5	7	6	9	12	19	12	10	4	1	-	-
Montana												
All races, both sexes	287	389	452	565	760	993	1,288	1,175	912	305	75	-
Male	190	242	258	320	426	550	641	480	323	84	14	-
Female	97	147	194	245	334	443	647	695	589	221	61	-
White, both sexes	248	350	410	523	724	951	1,265	1,153	898	300	75	-
Male	162	217	234	292	407	530	634	468	321	82	14	-
Female	86	133	176	231	317	421	631	685	577	218	61	-
Black, both sexes	1	-	3	-	-	-	-	1	-	-	-	-
Male	1	-	3	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	37	39	36	42	32	38	22	20	13	5	-	-
Male	26	25	20	28	18	19	7	10	1	2	-	-
Female	11	14	16	14	14	19	15	10	12	3	-	-
Asian or Pacific Islander 2/, both sexes	1	-	3	-	4	4	1	1	1	-	-	-
Male	1	-	1	-	1	1	-	1	1	-	-	-
Female	-	-	2	-	3	3	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Nebraska												
All races, both sexes	477	622	735	910	1,261	1,931	2,373	2,372	1,755	752	182	1
Male	329	377	432	538	733	1,005	1,157	985	583	173	32	1
Female	148	245	303	372	528	926	1,216	1,387	1,172	579	150	-
White, both sexes	429	573	683	862	1,200	1,863	2,304	2,328	1,732	749	178	1
Male	297	348	403	512	700	975	1,132	963	577	173	31	1
Female	132	225	280	350	500	888	1,172	1,365	1,155	576	147	-
Black, both sexes	33	37	39	38	44	59	54	33	21	3	4	-
Male	24	23	22	18	22	26	20	16	5	-	1	-
Female	9	14	17	20	22	33	34	17	16	3	3	-
American Indian 1/, both sexes	12	10	7	7	14	6	9	5	-	-	-	-
Male	7	6	5	7	9	3	4	2	-	-	-	-
Female	5	4	2	-	5	3	5	3	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	2	6	3	3	3	6	6	2	-	-	-
Male	1	-	2	1	2	1	1	4	1	-	-	-
Female	2	2	4	2	1	2	5	2	1	-	-	-
Nevada												
All races, both sexes	879	1,140	1,410	1,646	2,022	2,413	2,553	1,926	1,096	337	47	9
Male	582	684	844	988	1,172	1,301	1,272	824	377	82	5	8
Female	297	456	566	658	850	1,112	1,281	1,102	719	255	42	1
White, both sexes	758	990	1,256	1,440	1,812	2,221	2,371	1,813	1,042	319	46	8
Male	501	607	752	878	1,057	1,208	1,185	776	358	79	4	7
Female	257	383	504	562	755	1,013	1,186	1,037	684	240	42	1
Black, both sexes	79	83	105	118	123	115	121	60	36	12	1	1
Male	51	46	65	68	71	59	54	27	10	1	1	1
Female	28	37	40	50	52	56	67	33	26	11	-	-
American Indian 1/, both sexes	8	23	18	18	13	16	10	7	3	2	-	-
Male	7	11	10	13	9	7	6	3	1	-	-	-
Female	1	12	8	5	4	9	4	4	2	2	-	-
Asian or Pacific Islander 2/, both sexes	34	44	31	70	74	61	51	46	15	4	-	-
Male	23	20	17	29	35	27	27	18	8	2	-	-
Female	11	24	14	41	39	34	24	28	7	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
New Hampshire												
All races, both sexes	361	431	561	690	1,003	1,373	1,627	1,590	1,044	459	79	-
Male	221	258	335	397	547	717	756	653	292	90	12	-
Female	140	173	226	293	456	656	871	937	752	369	67	-
White, both sexes	354	430	557	685	996	1,366	1,620	1,589	1,038	458	79	-
Male	218	257	334	395	542	714	751	653	290	90	12	-
Female	136	173	223	290	454	652	869	936	748	368	67	-
Black, both sexes	5	1	2	1	3	3	5	1	3	1	-	-
Male	2	1	1	-	3	-	4	-	1	-	-	-
Female	3	-	1	1	-	3	1	1	2	1	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	1	4	3	4	2	-	3	-	-	-
Male	1	-	-	2	1	3	1	-	1	-	-	-
Female	1	-	1	2	2	1	1	-	2	-	-	-
New Jersey												
All races, both sexes	2,594	3,286	3,974	4,564	6,604	9,649	11,853	11,249	7,207	2,743	553	14
Male	1,613	1,951	2,243	2,525	3,557	4,876	5,332	4,306	2,177	593	89	12
Female	981	1,335	1,731	2,039	3,047	4,773	6,521	6,943	5,030	2,150	464	2
White, both sexes	1,915	2,542	3,067	3,629	5,529	8,482	10,745	10,383	6,766	2,565	498	7
Male	1,219	1,520	1,734	2,016	2,981	4,330	4,841	4,003	2,062	553	83	5
Female	696	1,022	1,333	1,613	2,548	4,152	5,904	6,380	4,704	2,012	415	2
Black, both sexes	629	668	802	812	931	1,002	955	742	373	155	46	7
Male	362	383	442	429	499	466	419	258	89	34	2	7
Female	267	285	360	383	432	536	536	484	284	121	44	-
American Indian 1/, both sexes	3	4	4	6	9	16	8	8	6	2	1	-
Male	2	2	2	4	4	7	4	-	1	-	-	-
Female	1	2	2	2	5	9	4	8	5	2	1	-
Asian or Pacific Islander 2/, both sexes	47	72	101	117	135	149	145	116	62	21	8	-
Male	30	46	65	76	73	73	68	45	25	6	4	-
Female	17	26	36	41	62	76	77	71	37	15	4	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
New Mexico												
All races, both sexes	678	789	903	983	1,349	1,725	2,069	1,789	1,268	468	87	7
Male	427	487	532	608	749	909	1,023	778	437	114	19	5
Female	251	302	371	375	600	816	1,046	1,011	831	354	68	2
White, both sexes	600	700	808	884	1,246	1,620	1,965	1,680	1,198	444	75	6
Male	381	435	476	547	695	854	980	722	415	108	12	4
Female	219	265	332	337	551	766	985	958	783	336	63	2
Black, both sexes	17	20	17	18	23	24	25	21	16	2	1	-
Male	10	10	11	15	15	13	10	8	2	1	-	-
Female	7	10	6	3	8	11	15	13	14	1	1	-
American Indian 1/, both sexes	59	67	73	75	75	77	74	86	51	21	11	1
Male	35	42	41	44	38	40	32	46	19	5	7	1
Female	24	25	32	31	37	37	42	40	32	16	4	-
Asian or Pacific Islander 2/, both sexes	2	2	5	6	5	4	5	2	3	1	-	-
Male	1	-	4	2	1	2	1	2	1	-	-	-
Female	1	2	1	4	4	2	4	-	2	1	-	-
New York												
All races, both sexes	6,039	7,548	8,732	10,530	14,012	19,916	24,032	22,706	16,083	6,600	1,484	5
Male	3,701	4,462	5,009	5,914	7,470	10,152	10,823	8,581	4,762	1,440	238	3
Female	2,338	3,086	3,723	4,616	6,542	9,764	13,209	14,125	11,321	5,160	1,246	2
White, both sexes	4,366	5,731	6,709	8,225	11,520	16,977	21,128	20,584	14,589	6,033	1,304	4
Male	2,717	3,404	3,936	4,686	6,205	8,785	9,641	7,887	4,357	1,309	207	2
Female	1,649	2,327	2,773	3,539	5,315	8,192	11,487	12,697	10,232	4,724	1,097	2
Black, both sexes	1,480	1,611	1,792	1,962	2,122	2,517	2,448	1,761	1,244	490	155	1
Male	856	929	935	1,038	1,055	1,137	947	542	311	101	25	1
Female	624	682	857	924	1,067	1,380	1,501	1,219	933	389	130	-
American Indian 1/, both sexes	17	18	15	35	23	27	32	13	9	3	2	-
Male	9	10	9	17	11	12	9	6	3	1	-	-
Female	8	8	6	18	12	15	23	7	6	2	2	-
Asian or Pacific Islander 2/, both sexes	176	188	216	308	347	395	424	348	241	74	23	-
Male	119	119	129	173	199	218	226	146	91	29	6	-
Female	57	69	87	135	148	177	198	202	150	45	17	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
North Carolina												
All races, both sexes	3,231	3,941	4,801	5,833	7,162	9,711	10,651	9,245	5,925	2,167	426	14
Male	2,043	2,419	2,860	3,355	3,958	5,042	4,752	3,346	1,635	412	63	10
Female	1,188	1,522	1,941	2,478	3,204	4,669	5,899	5,899	4,290	1,755	363	4
White, both sexes	2,113	2,737	3,572	4,425	5,626	7,855	8,809	7,792	4,943	1,783	317	8
Male	1,356	1,708	2,130	2,568	3,134	4,159	3,984	2,867	1,373	331	41	5
Female	757	1,029	1,442	1,857	2,492	3,696	4,825	4,925	3,570	1,452	276	3
Black, both sexes	1,061	1,132	1,163	1,333	1,452	1,761	1,752	1,381	950	369	104	6
Male	654	671	689	745	776	837	739	457	250	77	20	5
Female	407	461	474	588	676	924	1,013	924	700	292	84	1
American Indian 1/, both sexes	42	48	49	52	63	61	72	58	22	12	3	-
Male	28	26	30	31	37	31	23	16	7	2	1	-
Female	14	22	19	21	26	30	49	42	15	10	2	-
Asian or Pacific Islander 2/, both sexes	15	24	17	23	21	34	18	14	10	3	2	-
Male	5	14	11	11	11	15	6	6	5	2	1	-
Female	10	10	6	12	10	19	12	8	5	1	1	-
North Dakota												
All races, both sexes	149	184	260	343	509	667	890	945	774	363	58	-
Male	105	99	154	194	279	377	442	437	291	95	8	-
Female	44	85	106	149	230	290	448	508	483	268	50	-
White, both sexes	133	175	236	323	480	652	875	935	767	362	58	-
Male	94	95	146	187	267	373	437	433	290	95	8	-
Female	39	80	90	136	213	279	438	502	477	267	50	-
Black, both sexes	1	-	1	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	15	9	20	20	28	13	14	10	7	1	-	-
Male	10	4	6	7	11	4	5	4	1	-	-	-
Female	5	5	14	13	17	9	9	6	6	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	3	-	-	2	1	-	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	2	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Ohio												
All races, both sexes	4,210	5,144	6,197	7,834	10,515	14,799	17,476	15,289	9,733	3,589	725	-
Male	2,589	3,072	3,635	4,376	5,672	7,493	8,062	5,904	2,824	725	110	-
Female	1,621	2,072	2,562	3,458	4,843	7,306	9,414	9,385	6,909	2,864	615	-
White, both sexes	3,349	4,264	5,292	6,727	9,214	13,265	16,038	14,247	9,050	3,328	644	-
Male	2,091	2,547	3,110	3,766	4,971	6,757	7,424	5,497	2,622	675	96	-
Female	1,258	1,717	2,182	2,961	4,243	6,508	8,614	8,750	6,428	2,653	548	-
Black, both sexes	842	856	879	1,084	1,268	1,506	1,402	1,010	671	253	80	-
Male	487	509	506	597	687	726	626	394	195	48	14	-
Female	355	347	373	487	581	780	776	616	476	205	66	-
American Indian 1/, both sexes	2	5	4	4	4	3	4	1	1	-	-	-
Male	2	4	4	4	-	1	1	-	1	-	-	-
Female	-	1	-	-	4	2	3	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	19	22	19	29	25	32	31	11	8	1	-
Male	9	12	15	9	14	9	11	13	6	2	-	-
Female	8	7	7	10	15	16	21	18	5	6	1	-
Oklahoma												
All races, both sexes	1,525	1,812	2,339	2,655	3,558	4,255	5,156	4,428	3,156	1,209	258	2
Male	916	1,096	1,354	1,546	1,957	2,198	2,392	1,724	991	278	36	1
Female	609	716	985	1,109	1,601	2,057	2,764	2,704	2,165	931	222	1
White, both sexes	1,245	1,495	1,966	2,277	3,154	3,817	4,697	4,094	2,936	1,124	224	2
Male	761	916	1,140	1,333	1,758	1,953	2,194	1,616	916	255	32	1
Female	484	579	826	944	1,396	1,864	2,503	2,478	2,020	869	192	1
Black, both sexes	140	137	174	190	200	230	229	172	135	54	22	-
Male	82	82	91	114	101	128	90	58	46	12	3	-
Female	58	55	83	76	99	102	139	114	89	42	19	-
American Indian 1/, both sexes	133	165	184	176	183	197	211	154	77	28	12	-
Male	70	91	111	95	92	113	102	49	27	10	1	-
Female	63	74	73	81	91	84	109	105	50	18	11	-
Asian or Pacific Islander 2/, both sexes	7	15	15	12	21	11	19	8	8	3	-	-
Male	3	7	12	4	6	4	6	1	2	1	-	-
Female	4	8	3	8	15	7	13	7	6	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Oregon												
All races, both sexes	1,248	1,450	1,706	2,039	2,742	3,849	4,975	4,659	3,297	1,124	263	3
Male	776	865	985	1,127	1,455	1,992	2,332	2,042	1,104	310	54	2
Female	472	585	721	912	1,287	1,857	2,643	2,617	2,193	814	209	1
White, both sexes	1,168	1,367	1,635	1,939	2,650	3,722	4,847	4,563	3,235	1,107	255	3
Male	730	813	940	1,081	1,402	1,929	2,275	1,998	1,078	303	52	2
Female	438	554	695	858	1,248	1,793	2,572	2,565	2,157	804	203	1
Black, both sexes	37	31	32	38	31	44	36	36	22	5	3	-
Male	23	20	21	17	16	15	17	17	6	-	-	-
Female	14	11	11	21	15	29	19	19	16	5	3	-
American Indian 1/, both sexes	22	30	17	22	22	35	31	11	9	5	-	-
Male	12	20	12	9	12	20	17	6	5	3	-	-
Female	10	10	5	13	10	15	14	5	4	2	-	-
Asian or Pacific Islander 2/, both sexes	21	22	22	40	39	48	61	49	31	7	5	-
Male	11	12	12	20	25	28	23	21	15	4	2	-
Female	10	10	10	20	14	20	38	28	16	3	3	-
Pennsylvania												
All races, both sexes	4,321	5,424	6,627	8,352	12,092	17,787	22,061	20,721	13,022	4,843	948	35
Male	2,709	3,274	3,957	4,822	6,563	9,115	9,973	7,890	3,816	1,021	122	19
Female	1,612	2,150	2,670	3,530	5,529	8,672	12,088	12,831	9,206	3,822	826	16
White, both sexes	3,519	4,541	5,685	7,329	10,861	16,360	20,650	19,738	12,377	4,587	859	19
Male	2,249	2,766	3,417	4,238	5,943	8,433	9,411	7,540	3,628	974	107	10
Female	1,270	1,775	2,268	3,091	4,918	7,927	11,239	12,198	8,749	3,613	752	9
Black, both sexes	762	847	892	968	1,167	1,361	1,344	934	603	250	87	16
Male	441	488	510	547	586	642	535	333	171	46	13	9
Female	321	359	382	421	581	719	809	601	432	204	74	7
American Indian 1/, both sexes	-	6	8	4	4	13	8	5	-	-	1	-
Male	-	4	4	3	2	8	3	4	-	-	1	-
Female	-	2	4	1	2	5	5	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	40	30	42	51	60	53	59	44	42	6	1	-
Male	19	16	26	34	32	32	24	13	17	1	1	-
Female	21	14	16	17	28	21	35	31	25	5	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Rhode Island												
All races, both sexes	330	410	457	585	864	1,270	1,692	1,653	1,155	442	102	1
Male	215	253	281	363	463	652	784	643	320	104	15	1
Female	115	157	176	222	401	618	908	1,010	835	338	87	-
White, both sexes	311	385	427	541	809	1,228	1,656	1,610	1,135	433	99	1
Male	202	237	264	336	444	629	767	626	312	102	14	1
Female	109	148	163	205	365	599	889	984	823	331	85	-
Black, both sexes	16	20	22	33	47	38	30	33	13	9	2	-
Male	12	14	11	20	16	19	16	13	6	2	1	-
Female	4	6	11	13	31	19	14	20	7	7	1	-
American Indian 1/, both sexes	1	1	2	4	2	1	1	3	3	-	1	-
Male	-	-	1	2	1	1	-	1	-	-	-	-
Female	1	1	1	2	1	-	1	2	3	-	1	-
Asian or Pacific Islander 2/, both sexes	2	4	6	7	6	3	5	7	4	-	-	-
Male	1	2	5	5	2	3	1	3	2	-	-	-
Female	1	2	1	2	4	-	4	4	2	-	-	-
South Carolina												
All races, both sexes	1,880	2,389	2,647	3,023	3,721	4,708	5,379	4,457	2,642	1,007	221	5
Male	1,203	1,478	1,587	1,764	2,030	2,454	2,400	1,688	756	209	29	3
Female	677	911	1,060	1,259	1,691	2,254	2,979	2,769	1,886	798	192	2
White, both sexes	1,116	1,539	1,793	2,092	2,710	3,556	4,212	3,526	2,083	737	136	3
Male	734	936	1,086	1,219	1,507	1,897	1,957	1,357	598	151	15	2
Female	382	603	707	873	1,203	1,659	2,255	2,169	1,485	586	121	1
Black, both sexes	755	837	845	918	996	1,144	1,155	915	551	266	85	2
Male	465	539	496	537	519	555	439	322	157	57	14	1
Female	290	298	349	381	477	589	716	593	394	209	71	1
American Indian 1/, both sexes	3	3	3	2	3	4	3	4	3	2	-	-
Male	2	-	1	2	-	1	2	3	-	1	-	-
Female	1	3	2	-	3	3	1	1	3	1	-	-
Asian or Pacific Islander 2/, both sexes	6	10	6	11	12	4	9	12	5	2	-	-
Male	2	3	4	6	4	1	2	6	1	-	-	-
Female	4	7	2	5	8	3	7	6	4	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
South Dakota												
All races, both sexes	216	265	307	415	606	854	1,037	1,064	864	370	103	-
Male	138	160	184	261	360	494	525	481	262	89	21	-
Female	78	105	123	154	246	360	512	583	602	281	82	-
White, both sexes	179	222	254	354	557	813	1,005	1,033	848	366	101	-
Male	115	135	156	226	328	476	512	466	256	89	20	-
Female	64	87	98	128	229	337	493	567	592	277	81	-
Black, both sexes	1	-	1	2	3	2	1	1	1	-	1	-
Male	1	-	-	1	3	1	-	1	-	-	1	-
Female	-	-	1	1	-	1	1	-	1	-	-	-
American Indian 1/, both sexes	36	43	50	58	44	38	28	29	15	4	-	-
Male	22	25	26	34	28	17	13	14	6	-	-	-
Female	14	18	24	24	16	21	15	15	9	4	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	2	1	3	1	-	-	1	-
Male	-	-	2	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	1	1	3	1	-	-	1	-
Tennessee												
All races, both sexes	2,687	3,333	3,759	4,427	5,611	7,076	8,087	6,889	4,606	1,633	339	4
Male	1,700	2,038	2,183	2,555	3,073	3,630	3,509	2,510	1,270	351	48	3
Female	987	1,295	1,576	1,872	2,538	3,446	4,578	4,379	3,336	1,282	291	1
White, both sexes	2,093	2,714	3,176	3,747	4,824	6,151	7,265	6,181	4,125	1,417	275	3
Male	1,332	1,671	1,861	2,191	2,677	3,200	3,178	2,255	1,129	307	30	2
Female	761	1,043	1,315	1,556	2,147	2,951	4,087	3,926	2,996	1,110	245	1
Black, both sexes	582	606	565	661	772	899	805	694	476	215	62	1
Male	363	361	313	353	388	414	324	250	138	44	17	1
Female	219	245	252	308	384	485	481	444	338	171	45	-
American Indian 1/, both sexes	1	2	4	1	2	4	4	1	-	-	1	-
Male	-	1	3	1	1	3	1	1	-	-	-	-
Female	1	1	1	-	1	1	3	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	11	11	14	18	13	22	13	13	5	1	1	-
Male	5	5	6	10	7	13	6	4	3	-	1	-
Female	6	6	8	8	6	9	7	9	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Texas												
All races, both sexes	7,236	8,597	10,010	11,811	14,872	19,462	21,850	18,451	12,769	4,835	958	27
Male	4,591	5,351	5,995	6,742	8,158	9,807	10,209	7,210	3,850	1,053	177	23
Female	2,645	3,246	4,015	5,069	6,714	9,655	11,641	11,241	8,919	3,782	781	4
White, both sexes	5,713	7,041	8,443	10,047	12,894	17,216	19,884	16,833	11,603	4,330	795	26
Male	3,678	4,451	5,088	5,797	7,137	8,725	9,340	6,633	3,509	930	147	23
Female	2,035	2,590	3,355	4,250	5,757	8,491	10,544	10,200	8,094	3,400	648	3
Black, both sexes	1,425	1,418	1,445	1,601	1,793	2,048	1,792	1,475	1,099	478	154	1
Male	866	825	837	865	941	978	782	519	318	112	27	-
Female	559	593	608	736	852	1,070	1,010	956	781	366	127	1
American Indian 1/, both sexes	10	14	13	13	12	14	13	15	7	3	2	-
Male	2	9	11	6	2	7	4	4	2	-	-	-
Female	8	5	2	7	10	7	9	11	5	3	2	-
Asian or Pacific Islander 2/, both sexes	88	124	109	150	173	184	161	128	60	24	7	-
Male	45	66	59	74	78	97	83	54	21	11	3	-
Female	43	58	50	76	95	87	78	74	39	13	4	-
Utah												
All races, both sexes	517	594	781	905	1,091	1,571	2,103	1,995	1,303	422	70	2
Male	322	367	451	517	610	822	990	815	445	131	16	2
Female	195	227	330	388	481	749	1,113	1,180	858	291	54	-
White, both sexes	491	573	756	881	1,066	1,536	2,054	1,975	1,291	420	69	2
Male	305	356	436	506	593	804	965	807	438	131	15	2
Female	186	217	320	375	473	732	1,089	1,168	853	289	54	-
Black, both sexes	12	8	7	4	6	7	7	6	-	-	-	-
Male	7	2	6	3	3	4	3	3	-	-	-	-
Female	5	6	1	1	3	3	4	3	-	-	-	-
American Indian 1/, both sexes	6	4	10	10	5	6	14	3	7	-	-	-
Male	6	3	5	3	4	5	6	-	5	-	-	-
Female	-	1	5	7	1	1	8	3	2	-	-	-
Asian or Pacific Islander 2/, both sexes	8	9	8	10	14	22	28	11	5	2	1	-
Male	4	6	4	5	10	9	16	5	2	-	1	-
Female	4	3	4	5	4	13	12	6	3	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Vermont												
All races, both sexes	199	210	284	344	477	636	799	762	575	250	51	-
Male	131	124	166	208	266	336	387	322	163	54	7	-
Female	68	86	118	136	211	300	412	440	412	196	44	-
White, both sexes	198	210	281	344	475	633	799	761	573	250	51	-
Male	131	124	164	208	266	333	387	321	162	54	7	-
Female	67	86	117	136	209	300	412	440	411	196	44	-
Black, both sexes	1	-	2	-	1	2	-	-	1	-	-	-
Male	-	-	1	-	-	2	-	-	1	-	-	-
Female	1	-	1	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Virginia												
All races, both sexes	2,438	3,058	3,633	4,400	5,707	7,372	8,841	7,547	4,829	1,809	386	3
Male	1,475	1,852	2,134	2,568	3,108	3,719	4,003	2,899	1,407	365	57	2
Female	963	1,206	1,499	1,832	2,599	3,653	4,838	4,648	3,422	1,444	329	1
White, both sexes	1,653	2,193	2,678	3,308	4,430	5,886	7,267	6,350	4,096	1,548	307	2
Male	1,039	1,318	1,594	1,956	2,441	3,004	3,296	2,458	1,191	309	51	2
Female	614	875	1,084	1,352	1,989	2,882	3,971	3,892	2,905	1,239	256	-
Black, both sexes	741	819	885	1,022	1,190	1,392	1,462	1,110	687	251	78	1
Male	413	508	502	574	623	681	658	407	199	51	6	-
Female	328	311	383	448	567	711	804	703	488	200	72	1
American Indian 1/, both sexes	6	2	4	6	8	4	9	5	-	1	-	-
Male	3	2	-	4	2	4	4	1	-	-	-	-
Female	3	-	4	2	6	-	5	4	-	1	-	-
Asian or Pacific Islander 2/, both sexes	38	44	66	64	79	90	103	82	46	9	1	-
Male	20	24	38	34	42	30	45	33	17	5	-	-
Female	18	20	28	30	37	60	58	49	29	4	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Washington												
All races, both sexes	1,891	2,194	2,746	3,049	3,977	5,531	7,225	6,719	4,570	1,697	378	-
Male	1,182	1,295	1,604	1,769	2,125	2,891	3,440	2,914	1,564	468	70	-
Female	709	899	1,142	1,280	1,852	2,640	3,785	3,805	3,006	1,229	308	-
White, both sexes	1,715	1,953	2,528	2,797	3,671	5,189	6,907	6,463	4,408	1,639	362	-
Male	1,065	1,154	1,476	1,638	1,976	2,720	3,298	2,806	1,502	446	67	-
Female	650	799	1,052	1,159	1,695	2,469	3,609	3,657	2,906	1,193	295	-
Black, both sexes	78	116	83	74	90	108	105	75	52	16	5	-
Male	56	71	54	40	48	54	46	25	20	7	1	-
Female	22	45	29	34	42	54	59	50	32	9	4	-
American Indian 1/, both sexes	37	50	46	56	60	62	45	29	10	6	-	-
Male	22	35	24	29	28	31	18	13	3	1	-	-
Female	15	15	22	27	32	31	27	16	7	5	-	-
Asian or Pacific Islander 2/, both sexes	61	75	89	122	156	172	168	152	100	36	11	-
Male	39	35	50	62	73	86	78	70	39	14	2	-
Female	22	40	39	60	83	86	90	82	61	22	9	-
West Virginia												
All races, both sexes	797	1,063	1,354	1,694	2,119	3,018	3,297	2,809	1,769	643	117	1
Male	516	637	775	946	1,147	1,555	1,495	1,073	534	131	21	1
Female	281	426	579	748	972	1,463	1,802	1,736	1,235	512	96	-
White, both sexes	760	1,021	1,313	1,649	2,062	2,935	3,212	2,728	1,710	618	112	1
Male	490	619	754	926	1,127	1,517	1,470	1,044	512	126	18	1
Female	270	402	559	723	935	1,418	1,742	1,684	1,198	492	94	-
Black, both sexes	35	42	41	44	56	80	81	81	55	25	5	-
Male	26	18	21	19	20	37	25	29	21	5	3	-
Female	9	24	20	25	36	43	56	52	34	20	2	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	1	1	2	3	-	3	-	-	-
Male	-	-	-	1	-	1	-	-	1	-	-	-
Female	2	-	-	-	1	1	3	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Wisconsin												
All races, both sexes	1,533	1,958	2,347	3,042	4,116	5,870	7,425	7,292	5,308	2,083	466	-
Male	948	1,171	1,383	1,777	2,341	3,220	3,703	2,951	1,674	440	57	-
Female	585	787	964	1,265	1,775	2,650	3,722	4,341	3,634	1,643	409	-
White, both sexes	1,349	1,738	2,134	2,816	3,892	5,618	7,208	7,166	5,234	2,055	460	-
Male	839	1,051	1,269	1,655	2,231	3,091	3,619	2,907	1,650	435	56	-
Female	510	687	865	1,161	1,661	2,527	3,589	4,259	3,584	1,620	404	-
Black, both sexes	152	167	184	165	176	197	161	94	49	21	5	-
Male	91	93	99	90	85	103	63	29	18	3	-	-
Female	61	74	85	75	91	94	98	65	31	18	5	-
American Indian 1/, both sexes	23	41	17	47	30	27	30	11	13	5	1	-
Male	13	22	6	26	16	16	12	3	3	2	1	-
Female	10	19	11	21	14	11	18	8	10	3	-	-
Asian or Pacific Islander 2/, both sexes	9	12	12	14	18	28	26	21	12	2	-	-
Male	5	5	9	6	9	10	9	12	3	-	-	-
Female	4	7	3	8	9	18	17	9	9	2	-	-
Wyoming												
All races, both sexes	185	212	247	306	381	508	603	482	352	141	36	-
Male	110	124	139	184	209	259	284	196	107	29	7	-
Female	75	88	108	122	172	249	319	286	245	112	29	-
White, both sexes	177	203	235	296	378	502	593	478	349	139	36	-
Male	104	121	135	182	206	257	279	194	106	29	7	-
Female	73	82	100	114	172	245	314	284	243	110	29	-
Black, both sexes	2	3	2	1	-	3	5	3	-	2	-	-
Male	1	2	1	1	-	1	3	2	-	-	-	-
Female	1	1	1	-	-	2	2	1	-	2	-	-
American Indian 1/, both sexes	5	5	9	6	2	3	4	1	2	-	-	-
Male	4	1	3	1	2	1	1	-	1	-	-	-
Female	1	4	6	5	-	2	3	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	3	1	-	1	-	1	-	-	-
Male	1	-	-	-	1	-	1	-	-	-	-	-
Female	-	1	1	3	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
United States												
All races, both sexes	64,618	480	187	77	81	142	332	653	1,394	2,637	4,187	5,034
Male	34,303	288	94	41	40	73	196	356	869	1,774	2,856	3,510
Female	30,315	192	93	36	41	69	136	297	525	863	1,331	1,524
White, both sexes	47,439	315	135	58	47	84	152	270	597	1,384	2,261	2,982
Male	24,631	189	68	31	27	43	100	158	425	992	1,653	2,146
Female	22,808	126	67	27	20	41	52	112	172	392	608	836
Black, both sexes	15,593	144	38	14	29	52	168	364	755	1,186	1,835	1,928
Male	8,797	82	19	7	12	28	88	186	414	730	1,135	1,286
Female	6,796	62	19	7	17	24	80	178	341	456	700	642
American Indian 1/, both sexes	444	13	6	2	3	-	2	6	17	29	48	62
Male	244	12	4	-	1	-	1	4	10	23	33	35
Female	200	1	2	2	2	-	1	2	7	6	15	27
Asian or Pacific Islander 2/, both sexes	1,142	8	8	3	2	6	10	13	25	38	43	62
Male	631	5	3	3	-	2	7	8	20	29	35	43
Female	511	3	5	-	2	4	3	5	5	9	8	19
Alabama												
All races, both sexes	1,251	11	1	-	2	2	8	14	26	52	68	101
Male	585	7	1	-	1	-	4	9	16	31	50	61
Female	666	4	-	-	1	2	4	5	10	21	18	40
White, both sexes	800	5	-	-	2	1	2	3	6	19	26	48
Male	359	4	-	-	1	-	2	-	4	12	19	30
Female	441	1	-	-	1	1	-	3	2	7	7	18
Black, both sexes	446	6	1	-	-	1	6	11	20	31	42	52
Male	221	3	1	-	-	-	2	9	12	17	31	30
Female	225	3	-	-	-	1	4	2	8	14	11	22
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	1	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
Alaska												
All races, both sexes	53	4	2	-	1	-	1	1	1	6	2	6
Male	33	4	2	-	1	-	1	-	-	5	1	3
Female	20	-	-	-	-	-	-	1	1	1	1	3
White, both sexes	30	-	1	-	1	-	1	1	1	2	-	4
Male	18	-	1	-	1	-	1	-	-	2	-	1
Female	12	-	-	-	-	-	-	1	1	-	-	3
Black, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	18	3	1	-	-	-	-	-	-	3	1	2
Male	10	3	1	-	-	-	-	-	-	2	-	2
Female	8	-	-	-	-	-	-	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	1	1	-
Male	3	-	-	-	-	-	-	-	-	1	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1,056	13	9	1	1	1	5	11	20	46	58	77
Male	621	10	4	-	1	-	5	8	15	38	48	61
Female	435	3	5	1	-	1	-	3	5	8	10	16
White, both sexes	932	9	7	-	-	1	5	8	15	34	45	65
Male	534	6	3	-	-	-	5	6	11	26	39	49
Female	398	3	4	-	-	1	-	2	4	8	6	16
Black, both sexes	54	1	1	-	-	-	-	1	3	8	4	5
Male	43	1	1	-	-	-	-	1	3	8	3	5
Female	11	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	66	3	1	1	1	-	-	1	2	3	9	7
Male	40	3	-	-	1	-	-	-	1	3	6	7
Female	26	-	1	1	-	-	-	1	1	-	3	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	1	-	1	-	-
Male	4	-	-	-	-	-	-	1	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
Arkansas												
All races, both sexes	739	1	3	1	1	2	3	10	17	24	37	41
Male	374	1	2	-	-	1	3	4	7	14	30	25
Female	365	-	1	1	1	1	-	6	10	10	7	16
White, both sexes	579	1	2	1	1	1	2	5	7	13	20	34
Male	295	1	1	-	-	1	2	3	3	9	15	22
Female	284	-	1	1	1	-	-	2	4	4	5	12
Black, both sexes	158	-	1	-	-	1	1	5	10	11	17	6
Male	79	-	1	-	-	-	1	1	4	5	15	3
Female	79	-	-	-	-	1	-	4	6	6	2	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
California												
All races, both sexes	4,799	44	18	6	5	14	30	43	109	314	425	569
Male	3,063	23	7	4	4	6	21	32	85	257	342	431
Female	1,736	21	11	2	1	8	9	11	24	57	83	138
White, both sexes	3,786	35	13	6	5	7	25	26	84	230	329	443
Male	2,411	19	5	4	4	3	18	19	67	192	262	341
Female	1,375	16	8	2	1	4	7	7	17	38	67	102
Black, both sexes	612	6	1	-	-	3	3	14	20	67	71	93
Male	415	3	-	-	-	2	2	11	13	51	57	68
Female	197	3	1	-	-	1	1	3	7	16	14	25
American Indian 1/, both sexes	64	-	-	-	-	-	-	1	2	4	10	11
Male	41	-	-	-	-	-	-	1	2	3	9	6
Female	23	-	-	-	-	-	-	-	-	1	1	5
Asian or Pacific Islander 2/, both sexes	337	3	4	-	-	4	2	2	3	13	15	22
Male	196	1	2	-	-	1	1	1	3	11	14	16
Female	141	2	2	-	-	3	1	1	-	2	1	6

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
Colorado												
All races, both sexes	636	3	5	2	1	3	7	6	4	23	42	57
Male	379	2	2	1	1	-	3	4	3	17	33	45
Female	257	1	3	1	-	3	4	2	1	6	9	12
White, both sexes	593	3	4	2	1	3	6	6	3	22	37	47
Male	352	2	2	1	1	-	3	4	3	16	29	37
Female	241	1	2	1	-	3	3	2	-	6	8	10
Black, both sexes	30	-	1	-	-	-	1	-	1	1	4	9
Male	20	-	-	-	-	-	-	-	-	1	3	7
Female	10	-	1	-	-	-	1	-	1	-	1	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	920	5	1	1	1	1	2	4	11	37	61	69
Male	481	3	-	-	1	1	-	1	5	22	34	48
Female	439	2	1	1	-	-	2	3	6	15	27	21
White, both sexes	765	3	1	-	1	-	2	3	8	24	36	47
Male	391	2	-	-	1	-	-	1	4	14	21	32
Female	374	1	1	-	-	-	2	2	4	10	15	15
Black, both sexes	150	2	-	1	-	1	-	1	3	12	25	22
Male	87	1	-	-	-	1	-	-	1	7	13	16
Female	63	1	-	1	-	-	-	1	2	5	12	6
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
Delaware												
All races, both sexes	242	5	-	-	-	-	-	1	2	14	12	23
Male	131	4	-	-	-	-	-	-	-	7	7	16
Female	111	1	-	-	-	-	-	1	2	7	5	7
White, both sexes	165	3	-	-	-	-	-	-	-	6	4	11
Male	88	2	-	-	-	-	-	-	-	2	3	8
Female	77	1	-	-	-	-	-	-	-	4	1	3
Black, both sexes	75	2	-	-	-	-	-	1	2	8	8	11
Male	43	2	-	-	-	-	-	-	-	5	4	8
Female	32	-	-	-	-	-	-	1	2	3	4	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	395	3	-	1	1	-	2	11	28	29	55	49
Male	241	1	-	1	-	-	1	6	14	15	31	32
Female	154	2	-	-	1	-	1	5	14	14	24	17
White, both sexes	36	-	-	-	-	-	-	-	3	-	2	6
Male	27	-	-	-	-	-	-	-	3	-	2	6
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	357	3	-	1	1	-	2	11	25	29	53	43
Male	213	1	-	1	-	-	1	6	11	15	29	26
Female	144	2	-	-	1	-	1	5	14	14	24	17
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
Florida												
All races, both sexes	5,133	33	9	5	3	13	35	74	156	305	470	464
Male	2,893	19	6	4	1	6	17	35	88	194	290	319
Female	2,240	14	3	1	2	7	18	39	68	111	180	145
White, both sexes	3,627	22	8	4	2	7	11	18	49	132	220	240
Male	2,069	14	6	3	-	4	8	11	37	97	161	183
Female	1,558	8	2	1	2	3	3	7	12	35	59	57
Black, both sexes	1,472	10	1	1	1	6	24	55	104	171	249	221
Male	811	4	-	1	1	2	9	24	49	96	129	134
Female	661	6	1	-	-	4	15	31	55	75	120	87
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	1	-	-	2
Male	4	-	-	-	-	-	-	-	1	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	25	1	-	-	-	-	-	1	2	2	1	1
Male	9	1	-	-	-	-	-	-	1	1	-	1
Female	16	-	-	-	-	-	-	1	1	1	1	-
Georgia												
All races, both sexes	2,458	16	8	1	2	4	25	39	98	149	213	183
Male	1,297	10	3	-	1	3	17	22	63	107	139	115
Female	1,161	6	5	1	1	1	8	17	35	42	74	68
White, both sexes	1,307	10	3	-	1	2	8	5	22	56	66	86
Male	637	5	1	-	1	1	6	3	14	43	49	57
Female	670	5	2	-	-	1	2	2	8	13	17	29
Black, both sexes	1,137	6	5	1	1	2	16	34	75	92	145	97
Male	653	5	2	-	-	2	10	19	48	64	89	58
Female	484	1	3	1	1	-	6	15	27	28	56	39
American Indian 1/, both sexes	5	-	-	-	-	-	1	-	-	1	1	-
Male	2	-	-	-	-	-	1	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	1	-	1	-
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
Hawaii												
All races, both sexes	286	1	1	-	1	-	3	4	3	7	12	12
Male	156	1	-	-	-	-	3	3	3	4	8	8
Female	130	-	1	-	1	-	-	1	-	3	4	4
White, both sexes	66	-	-	-	1	-	-	-	1	1	4	4
Male	43	-	-	-	-	-	-	-	1	-	2	3
Female	23	-	-	-	1	-	-	-	-	1	2	1
Black, both sexes	5	-	-	-	-	-	-	-	1	-	1	-
Male	4	-	-	-	-	-	-	-	1	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	215	1	1	-	-	-	3	4	1	6	7	8
Male	109	1	-	-	-	-	3	3	1	4	5	5
Female	106	-	1	-	-	-	-	1	-	2	2	3
Idaho												
All races, both sexes	118	1	3	1	-	-	-	-	-	2	2	3
Male	62	1	1	-	-	-	-	-	-	1	2	2
Female	56	-	2	1	-	-	-	-	-	1	-	1
White, both sexes	117	1	3	1	-	-	-	-	-	2	2	3
Male	61	1	1	-	-	-	-	-	-	1	2	2
Female	56	-	2	1	-	-	-	-	-	1	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
Illinois												
All races, both sexes	3,084	33	11	4	4	3	13	23	56	89	150	166
Male	1,494	17	6	2	3	2	8	12	29	56	99	109
Female	1,590	16	5	2	1	1	5	11	27	33	51	57
White, both sexes	2,224	24	3	1	2	2	5	5	27	41	69	88
Male	1,060	13	2	-	1	1	2	3	17	27	46	63
Female	1,164	11	1	1	1	1	3	2	10	14	23	25
Black, both sexes	812	9	8	3	2	1	8	18	27	46	79	76
Male	410	4	4	2	2	1	6	9	11	28	51	46
Female	402	5	4	1	-	-	2	9	16	18	28	30
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	45	-	-	-	-	-	-	-	2	1	2	2
Male	23	-	-	-	-	-	-	-	1	1	2	-
Female	22	-	-	-	-	-	-	-	1	-	-	2
Indiana												
All races, both sexes	1,232	12	9	-	1	5	4	6	17	25	47	75
Male	629	7	6	-	1	2	3	4	7	16	31	52
Female	603	5	3	-	-	3	1	2	10	9	16	23
White, both sexes	1,056	9	9	-	1	3	2	5	13	19	41	46
Male	530	6	6	-	1	2	2	3	6	12	25	31
Female	526	3	3	-	-	1	-	2	7	7	16	15
Black, both sexes	171	3	-	-	-	2	2	1	4	6	6	29
Male	97	1	-	-	-	-	1	1	1	4	6	21
Female	74	2	-	-	-	2	1	-	3	2	-	8
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
Iowa												
All races, both sexes	388	1	2	1	-	-	2	2	3	4	15	26
Male	195	-	1	-	-	-	1	2	2	3	10	17
Female	193	1	1	1	-	-	1	-	1	1	5	9
White, both sexes	377	1	2	1	-	-	2	2	3	3	14	23
Male	188	-	1	-	-	-	1	2	2	2	9	14
Female	189	1	1	1	-	-	1	-	1	1	5	9
Black, both sexes	9	-	-	-	-	-	-	-	-	1	1	3
Male	6	-	-	-	-	-	-	-	-	1	1	3
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	433	5	2	-	3	-	3	4	6	7	18	28
Male	206	3	2	-	2	-	3	2	1	6	14	21
Female	227	2	-	-	1	-	-	2	5	1	4	7
White, both sexes	391	4	2	-	1	-	3	3	5	4	14	23
Male	184	2	2	-	-	-	3	1	1	3	10	20
Female	207	2	-	-	1	-	-	2	4	1	4	3
Black, both sexes	34	1	-	-	2	-	-	1	1	3	2	4
Male	18	1	-	-	2	-	-	1	-	3	2	1
Female	16	-	-	-	-	-	-	-	1	-	-	3
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	2	1
Male	4	-	-	-	-	-	-	-	-	-	2	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
Kentucky												
All races, both sexes	879	5	1	-	-	1	1	3	16	17	38	57
Male	458	4	1	-	-	1	1	2	13	10	22	43
Female	421	1	-	-	-	-	-	1	3	7	16	14
White, both sexes	796	5	1	-	-	-	-	2	11	11	28	48
Male	406	4	1	-	-	-	-	1	9	7	15	36
Female	390	1	-	-	-	-	-	1	2	4	13	12
Black, both sexes	82	-	-	-	-	1	1	1	5	6	10	9
Male	52	-	-	-	-	1	1	1	4	3	7	7
Female	30	-	-	-	-	-	-	-	1	3	3	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1,540	16	6	2	7	8	14	37	48	69	125	119
Male	823	9	3	2	2	6	7	22	28	42	96	78
Female	717	7	3	-	5	2	7	15	20	27	29	41
White, both sexes	814	8	2	2	2	4	1	6	5	18	53	44
Male	398	5	-	2	-	3	-	5	2	12	40	23
Female	416	3	2	-	2	1	1	1	3	6	13	21
Black, both sexes	716	7	4	-	5	4	13	31	42	51	72	74
Male	418	3	3	-	2	3	7	17	25	30	56	54
Female	298	4	1	-	3	1	6	14	17	21	16	20
American Indian 1/, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	1	-	-	1
Male	5	-	-	-	-	-	-	-	1	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
Maine												
All races, both sexes	299	2	-	-	1	2	2	1	1	4	6	10
Male	141	1	-	-	1	1	1	-	1	3	3	6
Female	158	1	-	-	-	1	1	1	-	1	3	4
White, both sexes	294	2	-	-	1	2	2	1	1	4	6	10
Male	137	1	-	-	1	1	1	-	1	3	3	6
Female	157	1	-	-	-	1	1	1	-	1	3	4
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1,972	5	2	2	1	7	10	24	51	91	165	199
Male	1,057	4	1	1	-	3	8	12	33	54	99	143
Female	915	1	1	1	1	4	2	12	18	37	66	56
White, both sexes	1,006	3	1	1	-	1	2	8	17	17	31	52
Male	499	3	1	1	-	-	2	5	10	12	20	38
Female	507	-	-	-	-	1	-	3	7	5	11	14
Black, both sexes	940	2	1	1	1	6	7	16	33	72	134	146
Male	545	1	-	-	-	3	6	7	22	40	79	105
Female	395	1	1	1	1	3	1	9	11	32	55	41
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	-	1
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	23	-	-	-	-	-	1	-	1	1	-	-
Male	12	-	-	-	-	-	-	-	1	1	-	-
Female	11	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
Massachusetts												
All races, both sexes	1,543	2	3	1	1	3	1	3	22	46	73	98
Male	750	1	2	-	-	2	1	3	14	28	45	66
Female	793	1	1	1	1	1	-	-	8	18	28	32
White, both sexes	1,388	1	2	1	1	2	1	3	14	38	61	79
Male	663	1	2	-	-	1	1	3	9	21	37	54
Female	725	-	-	1	1	1	-	-	5	17	24	25
Black, both sexes	139	1	1	-	-	1	-	-	7	8	10	18
Male	80	-	-	-	-	1	-	-	4	7	6	11
Female	59	1	1	-	-	-	-	-	3	1	4	7
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	1	-	2	1
Male	7	-	-	-	-	-	-	-	1	-	2	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1,569	8	4	1	1	2	3	12	29	43	73	93
Male	778	5	2	1	-	1	2	9	18	27	46	56
Female	791	3	2	-	1	1	1	3	11	16	27	37
White, both sexes	1,125	7	3	-	-	2	1	5	12	20	43	45
Male	525	5	2	-	-	1	-	3	5	10	26	30
Female	600	2	1	-	-	1	1	2	7	10	17	15
Black, both sexes	417	1	1	-	1	-	2	7	17	23	30	45
Male	238	-	-	-	-	-	2	6	13	17	20	25
Female	179	1	1	-	1	-	-	1	4	6	10	20
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	19	-	-	1	-	-	-	-	-	-	-	1
Male	13	-	-	1	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
Minnesota												
All races, both sexes	517	4	2	3	-	3	1	5	8	13	24	32
Male	281	2	-	3	-	2	-	2	6	9	17	27
Female	236	2	2	-	-	1	1	3	2	4	7	5
White, both sexes	451	2	2	3	-	2	-	4	6	9	15	22
Male	243	1	-	3	-	2	-	2	5	6	10	19
Female	208	1	2	-	-	-	-	2	1	3	5	3
Black, both sexes	45	1	-	-	-	1	-	1	-	3	8	7
Male	27	-	-	-	-	-	-	-	-	3	6	6
Female	18	1	-	-	-	1	-	1	-	-	2	1
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	2	-	-	3
Male	5	-	-	-	-	-	-	-	1	-	-	2
Female	5	-	-	-	-	-	-	-	1	-	-	1
Asian or Pacific Islander 2/, both sexes	11	1	-	-	-	-	1	-	-	1	1	-
Male	6	1	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	1	-	-	1	-	-
Mississippi												
All races, both sexes	827	9	5	1	1	5	10	10	20	42	57	59
Male	416	8	4	1	-	3	4	4	13	28	41	45
Female	411	1	1	-	1	2	6	6	7	14	16	14
White, both sexes	457	2	1	-	-	4	1	4	5	14	23	23
Male	228	1	1	-	-	2	-	1	4	13	20	16
Female	229	1	-	-	-	2	1	3	1	1	3	7
Black, both sexes	365	7	4	1	-	1	9	6	15	27	34	36
Male	188	7	3	1	-	1	4	3	9	15	21	29
Female	177	-	1	-	-	-	5	3	6	12	13	7
American Indian 1/, both sexes	2	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
Missouri												
All races, both sexes	1,288	5	3	1	2	5	5	9	32	27	65	76
Male	634	5	2	1	1	2	2	3	24	21	48	53
Female	654	-	1	-	1	3	3	6	8	6	17	23
White, both sexes	1,035	4	3	1	1	3	2	5	16	14	42	48
Male	485	4	2	1	1	1	-	2	11	10	32	31
Female	550	-	1	-	-	2	2	3	5	4	10	17
Black, both sexes	244	1	-	-	1	2	3	4	15	13	22	28
Male	145	1	-	-	-	1	2	1	12	11	16	22
Female	99	-	-	-	1	1	1	3	3	2	6	6
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	123	-	-	-	1	-	1	1	-	1	8	6
Male	58	-	-	-	-	-	-	-	-	1	2	4
Female	65	-	-	-	1	-	1	1	-	-	6	2
White, both sexes	112	-	-	-	-	-	1	1	-	1	5	4
Male	53	-	-	-	-	-	-	-	-	1	1	3
Female	59	-	-	-	-	-	1	1	-	-	4	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	11	-	-	-	1	-	-	-	-	-	3	2
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	6	-	-	-	1	-	-	-	-	-	2	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
Nebraska												
All races, both sexes	212	4	1	-	-	-	-	3	2	8	6	3
Male	93	1	1	-	-	-	-	1	1	7	4	3
Female	119	3	-	-	-	-	-	2	1	1	2	-
White, both sexes	197	3	1	-	-	-	-	3	1	7	6	3
Male	88	1	1	-	-	-	-	1	-	6	4	3
Female	109	2	-	-	-	-	-	2	1	1	2	-
Black, both sexes	13	1	-	-	-	-	-	-	1	1	-	-
Male	4	-	-	-	-	-	-	-	1	1	-	-
Female	9	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	605	9	-	-	2	-	-	4	13	20	34	40
Male	355	5	-	-	1	-	-	3	9	14	27	30
Female	250	4	-	-	1	-	-	1	4	6	7	10
White, both sexes	511	9	-	-	2	-	-	3	11	16	22	33
Male	304	5	-	-	1	-	-	2	9	11	19	26
Female	207	4	-	-	1	-	-	1	2	5	3	7
Black, both sexes	66	-	-	-	-	-	-	1	2	4	9	5
Male	34	-	-	-	-	-	-	1	-	3	6	3
Female	32	-	-	-	-	-	-	-	2	1	3	2
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	-	-	2	1
Male	14	-	-	-	-	-	-	-	-	-	2	-
Female	6	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
New Hampshire												
All races, both sexes	197	-	-	-	-	-	1	1	4	5	9	12
Male	99	-	-	-	-	-	1	-	2	-	4	9
Female	98	-	-	-	-	-	-	1	2	5	5	3
White, both sexes	193	-	-	-	-	-	1	1	4	4	9	11
Male	97	-	-	-	-	-	1	-	2	-	4	8
Female	96	-	-	-	-	-	-	1	2	4	5	3
Black, both sexes	3	-	-	-	-	-	-	-	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	3,156	19	6	3	3	5	14	25	50	119	189	199
Male	1,524	10	2	2	1	2	12	11	21	61	117	124
Female	1,632	9	4	1	2	3	2	14	29	58	72	75
White, both sexes	2,261	14	3	3	2	2	7	10	18	54	77	80
Male	1,035	8	1	2	1	-	6	4	11	32	54	51
Female	1,226	6	2	1	1	2	1	6	7	22	23	29
Black, both sexes	834	5	2	-	1	2	7	13	32	65	108	119
Male	453	2	-	-	-	1	6	6	10	29	59	73
Female	381	3	2	-	1	1	1	7	22	36	49	46
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	56	-	1	-	-	1	-	2	-	-	4	-
Male	35	-	1	-	-	1	-	1	-	-	4	-
Female	21	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
New Mexico												
All races, both sexes	278	5	3	1	-	-	1	2	3	13	24	16
Male	148	5	2	-	-	-	-	2	3	7	18	8
Female	130	-	1	1	-	-	1	-	-	6	6	8
White, both sexes	239	4	2	-	-	-	1	-	3	10	19	12
Male	127	4	1	-	-	-	-	-	3	5	15	5
Female	112	-	1	-	-	-	1	-	-	5	4	7
Black, both sexes	9	-	-	-	-	-	-	-	-	-	1	3
Male	5	-	-	-	-	-	-	-	-	-	1	2
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	29	1	1	1	-	-	-	2	-	3	4	1
Male	16	1	1	-	-	-	-	2	-	2	2	1
Female	13	-	-	1	-	-	-	-	-	1	2	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	5,090	17	8	6	10	13	21	47	104	250	384	498
Male	2,751	9	3	3	6	9	13	22	59	153	225	332
Female	2,339	8	5	3	4	4	8	25	45	97	159	166
White, both sexes	3,512	9	6	4	3	8	11	25	47	123	180	257
Male	1,874	7	2	2	2	7	8	14	33	79	123	182
Female	1,638	2	4	2	1	1	3	11	14	44	57	75
Black, both sexes	1,480	8	1	2	6	5	9	20	57	123	203	234
Male	821	2	1	1	4	2	4	6	26	71	102	145
Female	659	6	-	1	2	3	5	14	31	52	101	89
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	92	-	1	-	1	-	1	2	-	4	1	5
Male	53	-	-	-	-	-	1	2	-	3	-	4
Female	39	-	1	-	1	-	-	-	-	1	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
North Carolina												
All races, both sexes	2,063	20	4	3	1	4	12	19	65	62	132	162
Male	1,065	11	2	2	-	3	4	12	35	37	87	114
Female	998	9	2	1	1	1	8	7	30	25	45	48
White, both sexes	1,249	7	3	2	-	-	1	2	23	19	39	76
Male	610	4	1	1	-	-	1	2	16	10	28	58
Female	639	3	2	1	-	-	-	-	7	9	11	18
Black, both sexes	783	13	-	1	1	4	11	17	40	40	89	84
Male	439	7	-	1	-	3	3	10	18	24	56	55
Female	344	6	-	-	1	1	8	7	22	16	33	29
American Indian 1/, both sexes	23	-	1	-	-	-	-	-	1	3	2	2
Male	11	-	1	-	-	-	-	-	-	3	2	1
Female	12	-	-	-	-	-	-	-	1	-	-	1
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	1	-	2	-
Male	5	-	-	-	-	-	-	-	1	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
North Dakota												
All races, both sexes	67	-	-	-	-	-	-	-	2	-	1	4
Male	36	-	-	-	-	-	-	-	2	-	1	3
Female	31	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	60	-	-	-	-	-	-	-	-	-	1	4
Male	32	-	-	-	-	-	-	-	-	-	1	3
Female	28	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
Ohio												
All races, both sexes	2,491	20	5	1	6	3	6	14	23	53	93	131
Male	1,192	13	2	-	4	1	4	9	14	36	62	89
Female	1,299	7	3	1	2	2	2	5	9	17	31	42
White, both sexes	2,038	11	5	1	5	2	3	9	13	34	60	93
Male	953	7	2	-	3	-	3	6	8	26	43	68
Female	1,085	4	3	1	2	2	-	3	5	8	17	25
Black, both sexes	435	8	-	-	1	1	3	4	9	18	31	36
Male	231	5	-	-	1	1	1	3	5	9	18	20
Female	204	3	-	-	-	-	2	1	4	9	13	16
American Indian 1/, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	1	1	1	2	2
Male	6	-	-	-	-	-	-	-	1	1	1	1
Female	9	-	-	-	-	-	-	1	-	-	1	1
Oklahoma												
All races, both sexes	752	11	2	2	-	2	1	10	18	18	42	62
Male	398	7	1	-	-	-	-	8	11	14	27	38
Female	354	4	1	2	-	2	1	2	7	4	15	24
White, both sexes	624	8	-	2	-	2	-	6	10	12	24	46
Male	318	4	-	-	-	-	-	4	7	10	13	29
Female	306	4	-	2	-	2	-	2	3	2	11	17
Black, both sexes	79	1	1	-	-	-	1	3	2	5	12	6
Male	46	1	-	-	-	-	-	3	1	3	8	4
Female	33	-	1	-	-	-	1	-	1	2	4	2
American Indian 1/, both sexes	46	2	1	-	-	-	-	1	6	1	6	10
Male	33	2	1	-	-	-	-	1	3	1	6	5
Female	13	-	-	-	-	-	-	-	3	-	-	5
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
Oregon												
All races, both sexes	553	3	-	-	1	-	2	5	5	19	35	55
Male	311	2	-	-	-	-	1	2	4	17	23	41
Female	242	1	-	-	1	-	1	3	1	2	12	14
White, both sexes	520	3	-	-	1	-	2	4	5	16	33	50
Male	293	2	-	-	-	-	1	2	4	14	21	40
Female	227	1	-	-	1	-	1	2	1	2	12	10
Black, both sexes	6	-	-	-	-	-	-	-	-	-	2	-
Male	4	-	-	-	-	-	-	-	-	-	2	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	19	-	-	-	-	-	-	1	-	3	-	3
Male	10	-	-	-	-	-	-	-	-	3	-	1
Female	9	-	-	-	-	-	-	1	-	-	-	2
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	2
Pennsylvania												
All races, both sexes	3,861	15	6	6	1	6	9	15	51	94	166	225
Male	1,903	10	4	2	1	4	5	7	34	56	123	162
Female	1,958	5	2	4	-	2	4	8	17	38	43	63
White, both sexes	3,065	12	4	6	1	4	6	7	16	52	93	123
Male	1,447	7	3	2	1	2	3	4	11	32	72	85
Female	1,618	5	1	4	-	2	3	3	5	20	21	38
Black, both sexes	758	3	2	-	-	2	3	8	33	42	71	99
Male	437	3	1	-	-	2	2	3	21	24	49	76
Female	321	-	1	-	-	-	1	5	12	18	22	23
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	36	-	-	-	-	-	-	-	2	-	2	3
Male	19	-	-	-	-	-	-	-	2	-	2	1
Female	17	-	-	-	-	-	-	-	-	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
Rhode Island												
All races, both sexes	180	-	-	-	1	1	2	2	1	6	10	11
Male	93	-	-	-	-	1	2	-	1	3	6	4
Female	87	-	-	-	1	-	-	2	-	3	4	7
White, both sexes	158	-	-	-	1	1	1	1	1	4	6	9
Male	77	-	-	-	-	1	1	-	1	2	3	2
Female	81	-	-	-	1	-	-	1	-	2	3	7
Black, both sexes	17	-	-	-	-	-	-	1	-	2	3	2
Male	12	-	-	-	-	-	-	-	-	1	2	2
Female	5	-	-	-	-	-	-	1	-	1	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1,121	14	1	2	1	4	13	19	31	47	65	97
Male	590	6	-	-	-	1	7	12	20	36	50	67
Female	531	8	1	2	1	3	6	7	11	11	15	30
White, both sexes	611	7	1	-	-	1	5	4	7	10	20	35
Male	294	3	-	-	-	-	2	3	5	7	16	27
Female	317	4	1	-	-	1	3	1	2	3	4	8
Black, both sexes	500	5	-	2	1	3	8	15	24	35	45	61
Male	290	2	-	-	-	1	5	9	15	27	34	40
Female	210	3	-	2	1	2	3	6	9	8	11	21
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	1
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	8	2	-	-	-	-	-	-	-	1	-	-
Male	5	1	-	-	-	-	-	-	-	1	-	-
Female	3	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
South Dakota												
All races, both sexes	80	2	2	2	-	-	-	-	1	3	4	5
Male	43	1	-	2	-	-	-	-	1	2	2	4
Female	37	1	2	-	-	-	-	-	-	1	2	1
White, both sexes	57	-	1	2	-	-	-	-	-	1	-	4
Male	30	-	-	2	-	-	-	-	-	1	-	3
Female	27	-	1	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	21	2	1	-	-	-	-	-	1	2	3	1
Male	11	1	-	-	-	-	-	-	1	1	1	1
Female	10	1	1	-	-	-	-	-	-	1	2	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1,302	14	3	-	2	2	9	28	28	69	107	132
Male	676	5	2	-	1	-	6	12	17	52	69	87
Female	626	9	1	-	1	2	3	16	11	17	38	45
White, both sexes	916	6	3	-	1	1	2	8	8	29	59	74
Male	451	2	2	-	1	-	2	3	5	23	39	48
Female	465	4	1	-	-	1	-	5	3	6	20	26
Black, both sexes	385	8	-	-	1	1	7	20	20	40	48	57
Male	224	3	-	-	-	-	4	9	12	29	30	38
Female	161	5	-	-	1	1	3	11	8	11	18	19
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
Texas												
All races, both sexes	4,698	51	17	12	5	9	32	75	146	251	382	427
Male	2,661	34	10	8	2	6	17	38	98	191	295	316
Female	2,037	17	7	4	3	3	15	37	48	60	87	111
White, both sexes	3,629	39	15	10	3	8	15	46	69	172	256	296
Male	2,047	24	9	6	2	6	10	26	57	136	211	217
Female	1,582	15	6	4	1	2	5	20	12	36	45	79
Black, both sexes	1,007	12	1	-	2	-	16	29	77	77	125	127
Male	580	10	1	-	-	-	6	12	41	54	84	95
Female	427	2	-	-	2	-	10	17	36	23	41	32
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	58	-	1	2	-	1	1	-	-	2	-	3
Male	33	-	-	2	-	-	1	-	-	1	-	3
Female	25	-	1	-	-	1	-	-	-	1	-	-
Utah												
All races, both sexes	234	4	4	-	-	-	1	2	5	8	13	20
Male	120	2	2	-	-	-	1	1	3	6	9	11
Female	114	2	2	-	-	-	-	1	2	2	4	9
White, both sexes	224	4	4	-	-	-	1	2	3	7	13	19
Male	112	2	2	-	-	-	1	1	2	5	9	11
Female	112	2	2	-	-	-	-	1	1	2	4	8
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	1	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	1
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	1	1	-	-
Male	4	-	-	-	-	-	-	-	1	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
Vermont												
All races, both sexes	85	-	1	-	-	-	-	-	-	-	2	9
Male	36	-	1	-	-	-	-	-	-	-	2	5
Female	49	-	-	-	-	-	-	-	-	-	-	4
White, both sexes	85	-	1	-	-	-	-	-	-	-	2	9
Male	36	-	1	-	-	-	-	-	-	-	2	5
Female	49	-	-	-	-	-	-	-	-	-	-	4
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1,772	15	8	-	3	3	10	17	29	54	80	93
Male	902	9	2	-	1	1	5	9	23	30	53	74
Female	870	6	6	-	2	2	5	8	6	24	27	19
White, both sexes	1,123	7	8	-	1	3	5	4	7	17	32	46
Male	563	3	2	-	1	1	2	3	7	11	23	37
Female	560	4	6	-	-	2	3	1	-	6	9	9
Black, both sexes	620	8	-	-	1	-	5	13	21	37	48	44
Male	321	6	-	-	-	-	3	6	15	19	30	34
Female	299	2	-	-	1	-	2	7	6	18	18	10
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	26	-	-	-	1	-	-	-	1	-	-	3
Male	17	-	-	-	-	-	-	-	1	-	-	3
Female	9	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
Washington												
All races, both sexes	826	1	2	1	2	3	3	4	14	26	52	73
Male	457	1	1	-	1	2	2	3	10	21	37	57
Female	369	-	1	1	1	1	1	1	4	5	15	16
White, both sexes	725	-	1	1	2	2	2	4	7	21	48	61
Male	404	-	-	-	1	2	2	3	6	18	35	48
Female	321	-	1	1	1	-	-	1	1	3	13	13
Black, both sexes	43	1	1	-	-	1	-	-	1	4	4	6
Male	28	1	1	-	-	-	-	-	1	2	2	5
Female	15	-	-	-	-	1	-	-	-	2	2	1
American Indian 1/, both sexes	15	-	-	-	-	-	1	-	-	-	-	4
Male	5	-	-	-	-	-	-	-	-	-	-	2
Female	10	-	-	-	-	-	1	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	43	-	-	-	-	-	-	-	6	1	-	2
Male	20	-	-	-	-	-	-	-	3	1	-	2
Female	23	-	-	-	-	-	-	-	3	-	-	-
West Virginia												
All races, both sexes	461	1	2	-	-	2	1	-	4	11	12	26
Male	238	-	-	-	-	1	-	-	4	8	8	16
Female	223	1	2	-	-	1	1	-	-	3	4	10
White, both sexes	441	1	2	-	-	2	1	-	4	10	10	25
Male	229	-	-	-	-	1	-	-	4	7	6	16
Female	212	1	2	-	-	1	1	-	-	3	4	9
Black, both sexes	18	-	-	-	-	-	-	-	-	1	2	1
Male	9	-	-	-	-	-	-	-	-	1	2	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
I. Certain infectious and parasitic diseases (A00-B99)												
Wisconsin												
All races, both sexes	686	6	2	3	1	-	3	2	11	13	22	33
Male	340	5	1	1	1	-	-	2	9	5	17	27
Female	346	1	1	2	-	-	3	-	2	8	5	6
White, both sexes	608	5	2	3	1	-	3	1	6	10	15	20
Male	293	4	1	1	1	-	-	1	5	2	13	16
Female	315	1	1	2	-	-	3	-	1	8	2	4
Black, both sexes	58	1	-	-	-	-	-	1	5	2	6	9
Male	36	1	-	-	-	-	-	1	4	2	4	8
Female	22	-	-	-	-	-	-	-	1	-	2	1
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	1	1	2
Male	5	-	-	-	-	-	-	-	-	1	-	1
Female	8	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	2
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	68	2	-	-	-	1	-	1	-	2	2	3
Male	36	-	-	-	-	-	-	1	-	2	2	3
Female	32	2	-	-	-	1	-	-	-	-	-	-
White, both sexes	64	2	-	-	-	1	-	1	-	2	1	3
Male	33	-	-	-	-	-	-	1	-	2	1	3
Female	31	2	-	-	-	1	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
United States												
All races, both sexes	5,298	4,364	3,770	4,002	5,131	6,787	7,932	6,608	3,923	1,366	226	7
Male	3,622	2,863	2,149	2,157	2,610	3,213	3,469	2,576	1,167	336	39	5
Female	1,676	1,501	1,621	1,845	2,521	3,574	4,463	4,032	2,756	1,030	187	2
White, both sexes	3,371	2,960	2,662	2,975	4,088	5,678	6,809	5,790	3,451	1,188	177	5
Male	2,362	1,921	1,498	1,607	2,082	2,722	2,974	2,290	1,015	294	30	4
Female	1,009	1,039	1,164	1,368	2,006	2,956	3,835	3,500	2,436	894	147	1
Black, both sexes	1,789	1,284	979	885	885	955	970	708	418	159	46	2
Male	1,176	870	581	475	456	423	417	241	126	35	9	1
Female	613	414	398	410	429	532	553	467	292	124	37	1
American Indian 1/, both sexes	51	31	40	31	21	30	21	21	7	3	-	-
Male	24	15	20	17	8	16	10	9	2	-	-	-
Female	27	16	20	14	13	14	11	12	5	3	-	-
Asian or Pacific Islander 2/, both sexes	87	89	89	111	137	124	132	89	47	16	3	-
Male	60	57	50	58	64	52	68	36	24	7	-	-
Female	27	32	39	53	73	72	64	53	23	9	3	-
Alabama												
All races, both sexes	78	88	80	97	126	131	157	128	49	24	8	-
Male	48	61	36	39	58	64	43	39	11	5	1	-
Female	30	27	44	58	68	67	114	89	38	19	7	-
White, both sexes	45	54	54	72	91	91	118	106	32	19	6	-
Male	31	38	25	31	45	42	32	33	7	3	-	-
Female	14	16	29	41	46	49	86	73	25	16	6	-
Black, both sexes	33	33	25	25	35	40	39	22	17	5	2	-
Male	17	22	10	8	13	22	11	6	4	2	1	-
Female	16	11	15	17	22	18	28	16	13	3	1	-
American Indian 1/, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
Alaska												
All races, both sexes	8	2	3	5	-	1	4	3	3	-	-	-
Male	6	2	1	3	-	-	1	2	1	-	-	-
Female	2	-	2	2	-	1	3	1	2	-	-	-
White, both sexes	5	2	1	4	-	-	3	3	1	-	-	-
Male	5	2	-	2	-	-	1	2	-	-	-	-
Female	-	-	1	2	-	-	2	1	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	2	1	-	1	1	-	1	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	2	-	1	-	-	1	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	105	86	71	73	113	108	106	99	42	8	2	1
Male	73	62	36	40	55	44	49	48	20	3	-	1
Female	32	24	35	33	58	64	57	51	22	5	2	-
White, both sexes	93	75	61	64	106	100	102	92	39	8	2	1
Male	66	54	29	35	53	40	45	44	19	3	-	1
Female	27	21	32	29	53	60	57	48	20	5	2	-
Black, both sexes	6	7	5	3	4	2	1	2	1	-	-	-
Male	4	5	5	2	2	1	1	1	-	-	-	-
Female	2	2	-	1	2	1	-	1	1	-	-	-
American Indian 1/, both sexes	5	4	5	6	3	5	3	5	2	-	-	-
Male	2	3	2	3	-	2	3	3	1	-	-	-
Female	3	1	3	3	3	3	-	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
Arkansas												
All races, both sexes	42	40	39	47	57	98	101	81	63	25	6	-
Male	29	30	21	27	28	41	41	38	26	5	1	-
Female	13	10	18	20	29	57	60	43	37	20	5	-
White, both sexes	30	31	31	37	51	81	87	72	48	20	4	-
Male	21	22	17	23	24	32	39	34	20	5	1	-
Female	9	9	14	14	27	49	48	38	28	15	3	-
Black, both sexes	11	9	8	10	6	17	14	9	15	5	2	-
Male	8	8	4	4	4	9	2	4	6	-	-	-
Female	3	1	4	6	2	8	12	5	9	5	2	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	583	479	347	284	281	340	387	287	165	61	7	1
Male	425	350	231	178	145	160	163	129	51	18	-	1
Female	158	129	116	106	136	180	224	158	114	43	7	-
White, both sexes	446	378	271	216	227	274	324	239	146	56	5	1
Male	336	275	186	137	116	130	136	106	39	15	-	1
Female	110	103	85	79	111	144	188	133	107	41	5	-
Black, both sexes	99	66	38	29	14	26	27	25	7	2	1	-
Male	63	47	27	19	11	15	12	10	2	2	-	-
Female	36	19	11	10	3	11	15	15	5	-	1	-
American Indian 1/, both sexes	10	4	8	3	1	4	4	1	1	-	-	-
Male	5	3	3	2	1	4	2	-	-	-	-	-
Female	5	1	5	1	-	-	2	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	28	31	30	36	39	36	32	22	11	3	1	-
Male	21	25	15	20	17	11	13	13	10	1	-	-
Female	7	6	15	16	22	25	19	9	1	2	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
Colorado												
All races, both sexes	47	51	43	37	43	62	78	75	35	10	2	-
Male	39	41	22	20	25	30	37	38	8	7	1	-
Female	8	10	21	17	18	32	41	37	27	3	1	-
White, both sexes	46	50	39	33	41	62	71	70	35	10	2	-
Male	39	40	20	18	23	30	31	37	8	7	1	-
Female	7	10	19	15	18	32	40	33	27	3	1	-
Black, both sexes	-	1	2	3	-	-	3	4	-	-	-	-
Male	-	1	2	2	-	-	3	1	-	-	-	-
Female	-	-	-	1	-	-	-	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	1	2	-	4	-	-	-	-	-
Male	-	-	-	-	2	-	3	-	-	-	-	-
Female	1	-	2	1	-	-	1	-	-	-	-	-
Connecticut												
All races, both sexes	64	51	39	47	56	112	137	120	78	20	3	-
Male	47	38	22	26	32	63	54	55	21	6	2	-
Female	17	13	17	21	24	49	83	65	57	14	1	-
White, both sexes	40	37	30	40	48	101	131	116	74	20	3	-
Male	28	30	16	22	28	58	53	52	21	6	2	-
Female	12	7	14	18	20	43	78	64	53	14	1	-
Black, both sexes	24	14	8	6	7	11	5	4	4	-	-	-
Male	19	8	6	3	3	5	1	3	-	-	-	-
Female	5	6	2	3	4	6	4	1	4	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
Delaware												
All races, both sexes	26	15	13	9	21	30	34	19	13	5	-	-
Male	17	11	9	5	10	16	18	5	4	2	-	-
Female	9	4	4	4	11	14	16	14	9	3	-	-
White, both sexes	12	9	8	7	19	23	32	17	11	3	-	-
Male	9	6	5	4	9	12	18	5	3	2	-	-
Female	3	3	3	3	10	11	14	12	8	1	-	-
Black, both sexes	13	6	5	2	2	7	2	2	2	2	-	-
Male	8	5	4	1	1	4	-	-	1	-	-	-
Female	5	1	1	1	1	3	2	2	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	54	32	20	26	18	13	28	15	9	1	-	-
Male	39	28	15	20	7	3	18	8	2	-	-	-
Female	15	4	5	6	11	10	10	7	7	1	-	-
White, both sexes	-	2	5	2	2	1	8	2	3	-	-	-
Male	-	2	4	2	1	-	6	1	-	-	-	-
Female	-	-	1	-	1	1	2	1	3	-	-	-
Black, both sexes	54	29	15	24	15	12	20	13	6	1	-	-
Male	39	25	11	18	6	3	12	7	2	-	-	-
Female	15	4	4	6	9	9	8	6	4	1	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
Florida												
All races, both sexes	449	361	289	287	342	475	527	455	279	88	14	-
Male	329	241	187	170	177	245	246	194	97	24	4	-
Female	120	120	102	117	165	230	281	261	182	64	10	-
White, both sexes	287	243	196	202	292	429	496	415	261	82	11	-
Male	215	168	133	120	148	226	235	181	94	21	4	-
Female	72	75	63	82	144	203	261	234	167	61	7	-
Black, both sexes	158	114	92	83	48	44	28	36	18	5	3	-
Male	113	71	54	49	28	18	11	13	3	2	-	-
Female	45	43	38	34	20	26	17	23	15	3	3	-
American Indian 1/, both sexes	3	1	-	-	-	1	-	1	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	1	2	2	1	3	3	-	1	-	-
Male	1	1	-	1	1	-	-	-	-	1	-	-
Female	-	2	1	1	1	1	3	3	-	-	-	-
Georgia												
All races, both sexes	208	181	139	158	192	262	234	189	106	37	14	-
Male	144	112	68	84	111	113	83	65	25	11	1	-
Female	64	69	71	74	81	149	151	124	81	26	13	-
White, both sexes	76	82	78	96	109	191	172	139	70	28	7	-
Male	49	55	36	52	63	76	58	47	14	7	-	-
Female	27	27	42	44	46	115	114	92	56	21	7	-
Black, both sexes	132	97	61	60	82	69	61	49	36	9	7	-
Male	95	55	32	32	47	36	25	18	11	4	1	-
Female	37	42	29	28	35	33	36	31	25	5	6	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	1	1	2	-	1	-	-	-	-
Male	-	2	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
Hawaii												
All races, both sexes	25	18	22	21	29	29	38	30	21	8	1	-
Male	14	12	14	9	16	11	23	12	12	3	-	-
Female	11	6	8	12	13	18	15	18	9	5	1	-
White, both sexes	3	12	3	6	3	7	9	7	5	-	-	-
Male	3	10	2	2	2	4	5	5	4	-	-	-
Female	-	2	1	4	1	3	4	2	1	-	-	-
Black, both sexes	1	-	-	1	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	6	19	14	26	22	28	23	16	8	1	-
Male	10	2	12	7	14	7	17	7	8	3	-	-
Female	11	4	7	7	12	15	11	16	8	5	1	-
Idaho												
All races, both sexes	11	5	7	12	11	19	14	12	10	5	-	-
Male	11	3	3	6	4	11	7	6	2	2	-	-
Female	-	2	4	6	7	8	7	6	8	3	-	-
White, both sexes	11	5	7	12	11	18	14	12	10	5	-	-
Male	11	3	3	6	4	10	7	6	2	2	-	-
Female	-	2	4	6	7	8	7	6	8	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
Illinois												
All races, both sexes	177	165	182	215	269	350	443	393	244	82	12	-
Male	115	101	92	108	136	166	200	150	73	10	-	-
Female	62	64	90	107	133	184	243	243	171	72	12	-
White, both sexes	115	100	117	148	211	289	352	335	213	67	10	-
Male	81	58	56	74	104	140	164	134	64	10	-	-
Female	34	42	61	74	107	149	188	201	149	57	10	-
Black, both sexes	59	59	62	61	56	57	84	54	27	14	2	-
Male	32	39	33	31	32	24	31	15	9	-	-	-
Female	27	20	29	30	24	33	53	39	18	14	2	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	6	3	6	2	4	7	4	4	1	-	-
Male	1	4	3	3	-	2	5	1	-	-	-	-
Female	-	2	-	3	2	2	2	3	4	1	-	-
Indiana												
All races, both sexes	79	79	84	80	125	151	176	150	77	26	4	-
Male	63	47	45	41	64	64	87	53	27	8	1	-
Female	16	32	39	39	61	87	89	97	50	18	3	-
White, both sexes	60	68	71	70	111	131	158	140	72	23	4	-
Male	47	39	39	36	59	55	75	50	27	8	1	-
Female	13	29	32	34	52	76	83	90	45	15	3	-
Black, both sexes	19	11	13	10	11	20	16	10	5	3	-	-
Male	16	8	6	5	5	9	10	3	-	-	-	-
Female	3	3	7	5	6	11	6	7	5	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	3	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	3	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
Iowa												
All races, both sexes	18	20	23	27	30	60	60	55	26	12	1	-
Male	14	14	13	12	17	29	28	25	2	4	1	-
Female	4	6	10	15	13	31	32	30	24	8	-	-
White, both sexes	18	19	22	26	30	58	60	55	26	11	1	-
Male	14	13	13	12	17	29	28	25	2	3	1	-
Female	4	6	9	14	13	29	32	30	24	8	-	-
Black, both sexes	-	-	-	1	-	2	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	1	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	26	17	25	26	38	50	71	58	26	16	4	-
Male	17	8	18	11	21	22	31	15	6	2	1	-
Female	9	9	7	15	17	28	40	43	20	14	3	-
White, both sexes	21	13	24	22	35	48	66	57	26	16	4	-
Male	14	7	18	9	20	22	29	14	6	2	1	-
Female	7	6	6	13	15	26	37	43	20	14	3	-
Black, both sexes	2	4	1	4	2	2	5	-	-	-	-	-
Male	2	1	-	2	1	-	2	-	-	-	-	-
Female	-	3	1	2	1	2	3	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
Kentucky												
All races, both sexes	78	56	58	60	80	125	119	94	58	10	2	-
Male	57	31	32	34	39	61	54	36	15	2	-	-
Female	21	25	26	26	41	64	65	58	43	8	2	-
White, both sexes	64	53	52	55	74	122	116	89	55	9	1	-
Male	45	31	29	31	37	60	53	31	14	2	-	-
Female	19	22	23	24	37	62	63	58	41	7	1	-
Black, both sexes	14	2	6	5	6	3	3	5	3	1	1	-
Male	12	-	3	3	2	1	1	5	1	-	-	-
Female	2	2	3	2	4	2	2	-	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	123	98	90	104	110	136	175	128	91	28	6	-
Male	84	61	47	58	55	65	80	42	25	8	3	-
Female	39	37	43	46	55	71	95	86	66	20	3	-
White, both sexes	55	53	42	67	66	84	119	93	67	19	4	-
Male	35	32	16	35	33	42	55	32	17	6	3	-
Female	20	21	26	32	33	42	64	61	50	13	1	-
Black, both sexes	68	44	48	35	42	52	56	34	23	9	2	-
Male	49	29	31	21	20	23	25	10	8	2	-	-
Female	19	15	17	14	22	29	31	24	15	7	2	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	2	-	-	-	1	-	-	-
Male	-	-	-	1	2	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
Maine												
All races, both sexes	19	13	14	10	28	37	61	45	31	10	2	-
Male	12	10	7	6	14	17	32	16	9	1	-	-
Female	7	3	7	4	14	20	29	29	22	9	2	-
White, both sexes	18	13	14	10	27	35	60	45	31	10	2	-
Male	11	10	7	6	14	15	31	16	9	1	-	-
Female	7	3	7	4	13	20	29	29	22	9	2	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	2	1	-	-	-	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	169	134	113	115	151	165	222	194	104	42	6	-
Male	118	89	63	62	82	74	94	78	27	12	-	-
Female	51	45	50	53	69	91	128	116	77	30	6	-
White, both sexes	53	49	63	71	94	115	170	145	76	32	5	-
Male	37	31	33	39	49	56	71	62	21	8	-	-
Female	16	18	30	32	45	59	99	83	55	24	5	-
Black, both sexes	114	82	49	42	56	46	50	46	26	9	1	-
Male	79	57	30	22	32	16	22	15	6	3	-	-
Female	35	25	19	20	24	30	28	31	20	6	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	1	2	1	3	2	3	2	1	-	-
Male	2	1	-	1	1	2	1	1	-	1	-	-
Female	-	2	1	1	-	1	1	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
Massachusetts												
All races, both sexes	122	88	77	83	136	164	230	205	129	50	6	-
Male	86	55	44	45	75	70	93	68	35	16	1	-
Female	36	33	33	38	61	94	137	137	94	34	5	-
White, both sexes	105	77	62	72	119	153	222	198	124	47	6	-
Male	74	46	35	41	68	65	89	67	33	15	1	-
Female	31	31	27	31	51	88	133	131	91	32	5	-
Black, both sexes	16	11	14	10	15	11	6	4	3	3	-	-
Male	11	9	9	4	7	5	3	1	1	1	-	-
Female	5	2	5	6	8	6	3	3	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	2	-	2	3	2	-	-	-
Male	1	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	1	1	2	-	1	3	1	-	-	-
Michigan												
All races, both sexes	133	89	107	91	119	186	213	208	119	32	3	-
Male	91	60	65	42	60	78	91	83	31	9	1	-
Female	42	29	42	49	59	108	122	125	88	23	2	-
White, both sexes	75	50	69	65	95	148	173	178	103	28	3	-
Male	54	35	40	29	48	58	70	74	25	9	1	-
Female	21	15	29	36	47	90	103	104	78	19	2	-
Black, both sexes	56	34	34	24	21	35	37	29	16	4	-	-
Male	35	21	23	12	12	18	20	8	6	-	-	-
Female	21	13	11	12	9	17	17	21	10	4	-	-
American Indian 1/, both sexes	-	-	2	1	2	-	1	-	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	2	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	5	2	1	1	3	2	1	-	-	-	-
Male	2	4	-	1	-	2	1	1	-	-	-	-
Female	-	1	2	-	1	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
Minnesota												
All races, both sexes	44	31	21	28	50	51	73	71	39	14	-	-
Male	29	22	10	13	24	28	34	37	14	2	-	-
Female	15	9	11	15	26	23	39	34	25	12	-	-
White, both sexes	32	27	17	24	46	48	72	68	38	14	-	-
Male	23	20	8	13	22	26	33	35	13	2	-	-
Female	9	7	9	11	24	22	39	33	25	12	-	-
Black, both sexes	8	3	4	3	2	1	1	1	1	-	-	-
Male	4	2	2	-	1	1	1	-	1	-	-	-
Female	4	1	2	3	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	3	1	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	2	2	-	1	-	-	-	-
Male	1	-	-	-	1	1	-	1	-	-	-	-
Female	-	-	-	1	1	1	-	-	-	-	-	-
Mississippi												
All races, both sexes	58	52	57	56	66	78	76	78	57	25	5	-
Male	33	28	22	31	24	40	36	31	14	3	3	-
Female	25	24	35	25	42	38	40	47	43	22	2	-
White, both sexes	22	24	31	28	44	57	50	56	46	19	3	-
Male	15	10	13	18	18	29	25	25	12	3	2	-
Female	7	14	18	10	26	28	25	31	34	16	1	-
Black, both sexes	36	27	25	27	22	21	26	22	11	6	2	-
Male	18	18	9	13	6	11	11	6	2	-	1	-
Female	18	9	16	14	16	10	15	16	9	6	1	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
Missouri												
All races, both sexes	82	88	68	82	94	152	179	155	117	36	5	-
Male	55	62	33	44	40	63	76	59	29	10	1	-
Female	27	26	35	38	54	89	103	96	88	26	4	-
White, both sexes	54	60	51	71	78	137	160	141	108	31	5	-
Male	36	44	24	37	34	58	65	53	29	9	1	-
Female	18	16	27	34	44	79	95	88	79	22	4	-
Black, both sexes	27	24	17	11	15	14	19	14	9	5	-	-
Male	19	16	9	7	5	5	11	6	-	1	-	-
Female	8	8	8	4	10	9	8	8	9	4	-	-
American Indian 1/, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	-	-	-	1	-	-	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	1	-	-	-	-	-	-
Montana												
All races, both sexes	12	10	10	7	7	8	19	13	15	3	1	-
Male	8	6	3	4	4	6	11	6	3	-	-	-
Female	4	4	7	3	3	2	8	7	12	3	1	-
White, both sexes	11	8	10	6	7	8	19	13	14	3	1	-
Male	7	5	3	3	4	6	11	6	3	-	-	-
Female	4	3	7	3	3	2	8	7	11	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	2	-	1	-	-	-	-	1	-	-	-
Male	1	1	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
Nebraska												
All races, both sexes	17	10	9	7	17	26	44	33	17	4	1	-
Male	10	6	4	3	11	7	21	6	5	2	-	-
Female	7	4	5	4	6	19	23	27	12	2	1	-
White, both sexes	13	8	7	7	17	25	41	33	17	4	1	-
Male	8	5	4	3	11	7	21	6	5	2	-	-
Female	5	3	3	4	6	18	20	27	12	2	1	-
Black, both sexes	2	2	2	-	-	1	3	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	1	2	-	-	1	3	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	49	54	39	47	70	87	69	46	17	4	-	1
Male	35	32	25	29	42	40	36	22	3	1	-	1
Female	14	22	14	18	28	47	33	24	14	3	-	-
White, both sexes	40	42	29	40	57	81	61	44	16	4	-	1
Male	31	27	18	24	35	38	31	22	3	1	-	1
Female	9	15	11	16	22	43	30	22	13	3	-	-
Black, both sexes	7	9	5	6	5	4	7	1	1	-	-	-
Male	2	5	2	4	2	2	4	-	-	-	-	-
Female	5	4	3	2	3	2	3	1	1	-	-	-
American Indian 1/, both sexes	1	3	1	-	-	1	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	3	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	4	1	8	1	1	1	-	-	-	-
Male	1	-	4	1	5	-	1	-	-	-	-	-
Female	-	-	-	-	3	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
New Hampshire												
All races, both sexes	16	13	9	13	15	21	32	23	17	5	1	-
Male	9	6	4	8	7	9	19	10	10	1	-	-
Female	7	7	5	5	8	12	13	13	7	4	1	-
White, both sexes	16	13	8	13	14	21	32	23	17	5	1	-
Male	9	6	4	8	6	9	19	10	10	1	-	-
Female	7	7	4	5	8	12	13	13	7	4	1	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	219	166	155	155	234	372	441	417	253	97	13	2
Male	136	111	98	80	120	180	191	146	75	20	2	2
Female	83	55	57	75	114	192	250	271	178	77	11	-
White, both sexes	104	91	92	110	186	298	388	382	241	89	9	1
Male	69	56	53	56	95	143	172	131	71	18	-	1
Female	35	35	39	54	91	155	216	251	170	71	9	-
Black, both sexes	113	67	56	40	38	65	45	32	12	7	4	1
Male	66	50	39	21	22	31	16	13	4	2	2	1
Female	47	17	17	19	16	34	29	19	8	5	2	-
American Indian 1/, both sexes	-	1	1	-	2	-	-	-	-	1	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	2	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	2	7	6	5	8	9	8	3	-	-	-	-
Male	1	5	5	3	3	6	3	2	-	-	-	-
Female	1	2	1	2	5	3	5	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
New Mexico												
All races, both sexes	29	16	17	18	22	32	37	23	10	5	1	-
Male	18	11	8	11	9	17	17	7	3	2	-	-
Female	11	5	9	7	13	15	20	16	7	3	1	-
White, both sexes	26	16	11	15	19	32	35	19	10	4	1	-
Male	16	11	6	8	7	17	17	7	3	2	-	-
Female	10	5	5	7	12	15	18	12	7	2	1	-
Black, both sexes	2	-	1	-	1	-	1	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	-	5	3	1	-	1	4	-	1	-	-
Male	1	-	2	3	1	-	-	-	-	-	-	-
Female	-	-	3	-	-	-	1	4	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
New York												
All races, both sexes	538	380	286	292	358	457	533	468	288	108	23	1
Male	350	252	169	168	201	226	244	192	89	22	4	-
Female	188	128	117	124	157	231	289	276	199	86	19	1
White, both sexes	288	228	184	206	287	388	463	411	265	99	19	1
Male	192	144	107	127	159	199	214	168	81	21	3	-
Female	96	84	77	79	128	189	249	243	184	78	16	1
Black, both sexes	240	145	96	71	62	60	57	47	21	9	4	-
Male	150	106	58	34	37	23	22	19	6	1	1	-
Female	90	39	38	37	25	37	35	28	15	8	3	-
American Indian 1/, both sexes	2	1	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	6	6	14	9	9	13	10	2	-	-	-
Male	7	2	4	6	5	4	8	5	2	-	-	-
Female	1	4	2	8	4	5	5	5	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
North Carolina												
All races, both sexes	160	144	120	127	180	227	269	185	111	41	15	-
Male	116	89	68	68	77	104	112	75	34	9	6	-
Female	44	55	52	59	103	123	157	110	77	32	9	-
White, both sexes	77	77	75	71	136	177	197	144	87	26	10	-
Male	52	45	44	37	56	87	76	56	27	5	4	-
Female	25	32	31	34	80	90	121	88	60	21	6	-
Black, both sexes	80	67	42	54	42	48	68	38	24	15	5	-
Male	62	44	24	31	19	17	36	16	7	4	2	-
Female	18	23	18	23	23	31	32	22	17	11	3	-
American Indian 1/, both sexes	2	-	3	2	1	1	3	2	-	-	-	-
Male	1	-	-	-	1	-	-	2	-	-	-	-
Female	1	-	3	2	-	1	3	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	1	1	1	-	-	-	-
Male	1	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
North Dakota												
All races, both sexes	2	4	4	6	6	13	8	6	9	2	-	-
Male	2	4	-	2	3	9	4	1	5	-	-	-
Female	-	-	4	4	3	4	4	5	4	2	-	-
White, both sexes	2	4	2	4	6	13	7	6	9	2	-	-
Male	2	4	-	1	3	9	3	1	5	-	-	-
Female	-	-	2	3	3	4	4	5	4	2	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	2	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
Ohio												
All races, both sexes	145	131	161	186	229	328	391	317	194	50	4	-
Male	77	82	90	80	113	167	164	123	52	9	1	-
Female	68	49	71	106	116	161	227	194	142	41	3	-
White, both sexes	98	95	128	150	188	288	346	286	174	45	4	-
Male	45	57	67	67	93	149	142	114	44	8	1	-
Female	53	38	61	83	95	139	204	172	130	37	3	-
Black, both sexes	46	36	33	34	40	38	42	30	20	5	-	-
Male	31	25	23	13	20	17	21	9	8	1	-	-
Female	15	11	10	21	20	21	21	21	12	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	2	1	-	3	1	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	2	1	-	2	1	-	-	-	-
Oklahoma												
All races, both sexes	54	66	54	57	76	73	79	63	42	17	3	-
Male	35	45	33	25	44	32	40	24	11	3	-	-
Female	19	21	21	32	32	41	39	39	31	14	3	-
White, both sexes	41	58	48	49	66	63	71	59	40	16	3	-
Male	25	40	28	21	37	27	38	23	9	3	-	-
Female	16	18	20	28	29	36	33	36	31	13	3	-
Black, both sexes	7	7	3	5	8	5	7	3	2	1	-	-
Male	5	4	2	3	6	2	2	-	2	-	-	-
Female	2	3	1	2	2	3	5	3	-	1	-	-
American Indian 1/, both sexes	6	1	3	2	1	4	1	1	-	-	-	-
Male	5	1	3	1	1	2	-	1	-	-	-	-
Female	1	-	-	1	-	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
Oregon												
All races, both sexes	66	45	43	33	32	47	74	47	25	15	1	-
Male	45	26	26	15	20	20	34	23	7	5	-	-
Female	21	19	17	18	12	27	40	24	18	10	1	-
White, both sexes	64	41	39	30	32	44	69	46	25	15	1	-
Male	45	23	22	13	20	20	31	23	7	5	-	-
Female	19	18	17	17	12	24	38	23	18	10	1	-
Black, both sexes	-	1	2	-	-	1	-	-	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	2	2	2	2	-	2	2	-	-	-	-	-
Male	-	2	2	1	-	-	1	-	-	-	-	-
Female	2	-	-	1	-	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	3	1	-	-	-	-
Male	-	1	-	1	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Pennsylvania												
All races, both sexes	222	201	205	221	349	485	656	514	295	107	12	-
Male	147	119	119	118	168	227	297	189	89	21	1	-
Female	75	82	86	103	181	258	359	325	206	86	11	-
White, both sexes	139	131	139	172	293	417	595	475	269	100	11	-
Male	95	69	77	92	145	193	278	179	78	18	1	-
Female	44	62	62	80	148	224	317	296	191	82	10	-
Black, both sexes	80	67	63	44	49	64	58	37	25	7	1	-
Male	50	49	41	24	19	32	18	10	10	3	-	-
Female	30	18	22	20	30	32	40	27	15	4	1	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	3	3	5	6	3	3	2	1	-	-	-
Male	2	1	1	2	4	2	1	-	1	-	-	-
Female	1	2	2	3	2	1	2	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
Rhode Island												
All races, both sexes	11	17	3	9	11	20	17	33	19	5	1	-
Male	9	13	1	8	6	12	8	14	3	2	-	-
Female	2	4	2	1	5	8	9	19	16	3	1	-
White, both sexes	10	13	3	7	11	17	16	32	19	5	1	-
Male	8	9	1	6	6	10	8	14	3	2	-	-
Female	2	4	2	1	5	7	8	18	16	3	1	-
Black, both sexes	1	4	-	1	-	2	1	-	-	-	-	-
Male	1	4	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	1	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
South Carolina												
All races, both sexes	91	86	70	89	90	111	126	96	42	22	4	-
Male	59	54	37	45	47	46	52	38	12	1	-	-
Female	32	32	33	44	43	65	74	58	30	21	4	-
White, both sexes	44	47	36	54	62	80	81	74	29	13	1	-
Male	28	24	21	26	35	33	29	29	6	-	-	-
Female	16	23	15	28	27	47	52	45	23	13	1	-
Black, both sexes	47	39	33	35	27	29	45	21	13	9	3	-
Male	31	30	15	19	12	12	23	8	6	1	-	-
Female	16	9	18	16	15	17	22	13	7	8	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	2	-	1	-	-	-	-
Male	-	-	1	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
South Dakota												
All races, both sexes	3	7	5	7	6	8	7	7	9	1	1	-
Male	1	4	4	2	3	5	2	6	3	-	1	-
Female	2	3	1	5	3	3	5	1	6	1	-	-
White, both sexes	3	5	4	5	3	6	6	6	9	1	1	-
Male	1	3	3	2	1	4	1	5	3	-	1	-
Female	2	2	1	3	2	2	5	1	6	1	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	2	1	2	2	2	1	1	-	-	-	-
Male	-	1	1	-	1	1	1	1	-	-	-	-
Female	-	1	-	2	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	121	86	55	69	103	127	137	118	73	18	1	-
Male	82	59	29	28	50	64	52	40	18	3	-	-
Female	39	27	26	41	53	63	85	78	55	15	1	-
White, both sexes	80	58	40	60	88	109	118	95	61	15	1	-
Male	54	38	20	23	41	57	43	33	15	2	-	-
Female	26	20	20	37	47	52	75	62	46	13	1	-
Black, both sexes	41	28	15	9	15	18	19	23	12	3	-	-
Male	28	21	9	5	9	7	9	7	3	1	-	-
Female	13	7	6	4	6	11	10	16	9	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
Texas												
All races, both sexes	406	314	265	315	366	456	476	377	214	89	13	-
Male	272	193	158	192	186	211	203	136	61	34	-	-
Female	134	121	107	123	180	245	273	241	153	55	13	-
White, both sexes	302	235	204	252	290	389	422	330	189	77	10	-
Male	211	147	117	151	147	184	180	122	54	30	-	-
Female	91	88	87	101	143	205	242	208	135	47	10	-
Black, both sexes	97	76	58	57	68	59	46	42	24	11	3	-
Male	55	45	38	36	36	25	18	14	7	3	-	-
Female	42	31	20	21	32	34	28	28	17	8	3	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	3	3	5	8	8	8	4	1	1	-	-
Male	6	1	3	5	3	2	5	-	-	1	-	-
Female	1	2	-	-	5	6	3	4	1	-	-	-
Utah												
All races, both sexes	17	20	12	12	25	31	28	17	10	5	-	-
Male	13	14	5	3	11	16	9	7	5	2	-	-
Female	4	6	7	9	14	15	19	10	5	3	-	-
White, both sexes	16	19	12	12	25	29	27	17	9	5	-	-
Male	12	13	5	3	11	14	8	7	4	2	-	-
Female	4	6	7	9	14	15	19	10	5	3	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	2	-	-	1	-	-	-
Male	-	-	-	-	-	2	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
Vermont												
All races, both sexes	2	3	4	1	5	9	15	17	8	8	1	-
Male	1	2	2	1	2	6	7	3	2	2	-	-
Female	1	1	2	-	3	3	8	14	6	6	1	-
White, both sexes	2	3	4	1	5	9	15	17	8	8	1	-
Male	1	2	2	1	2	6	7	3	2	2	-	-
Female	1	1	2	-	3	3	8	14	6	6	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	108	108	79	129	172	211	261	219	129	37	6	1
Male	73	69	44	71	82	101	106	89	45	13	2	-
Female	35	39	35	58	90	110	155	130	84	24	4	1
White, both sexes	52	60	48	83	108	149	197	170	93	29	4	-
Male	38	34	27	47	51	75	77	75	36	12	1	-
Female	14	26	21	36	57	74	120	95	57	17	3	-
Black, both sexes	54	45	30	44	62	57	60	47	33	8	2	1
Male	34	32	16	22	30	23	28	12	9	1	1	-
Female	20	13	14	22	32	34	32	35	24	7	1	1
American Indian 1/, both sexes	2	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	1	2	1	5	4	2	3	-	-	-
Male	-	3	1	2	1	3	1	2	-	-	-	-
Female	-	-	-	-	-	2	3	-	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
Washington												
All races, both sexes	98	66	54	42	54	95	88	73	50	20	5	-
Male	80	42	25	20	25	44	35	30	17	4	-	-
Female	18	24	29	22	29	51	53	43	33	16	5	-
White, both sexes	90	54	44	34	48	84	82	71	47	18	4	-
Male	72	37	21	17	23	39	31	29	16	4	-	-
Female	18	17	23	17	25	45	51	42	31	14	4	-
Black, both sexes	6	7	4	2	-	3	2	-	1	-	-	-
Male	6	4	2	-	-	1	2	-	1	-	-	-
Female	-	3	2	2	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	1	3	2	-	2	-	-	1	-	1	-	-
Male	1	-	1	-	1	-	-	-	-	-	-	-
Female	-	3	1	-	1	-	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	2	4	6	4	8	4	1	2	1	1	-
Male	1	1	1	3	1	4	2	1	-	-	-	-
Female	-	1	3	3	3	4	2	-	2	1	1	-
West Virginia												
All races, both sexes	29	31	39	36	56	59	67	53	25	6	1	-
Male	21	15	24	19	26	33	31	21	9	2	-	-
Female	8	16	15	17	30	26	36	32	16	4	1	-
White, both sexes	26	29	38	35	53	57	65	52	24	6	1	-
Male	19	15	24	18	24	32	31	21	9	2	-	-
Female	7	14	14	17	29	25	34	31	15	4	1	-
Black, both sexes	2	2	1	1	3	2	2	1	-	-	-	-
Male	2	-	-	1	2	1	-	-	-	-	-	-
Female	-	2	1	-	1	1	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
Wisconsin												
All races, both sexes	52	42	37	40	58	94	102	86	53	21	5	-
Male	27	26	15	21	32	49	48	34	16	4	-	-
Female	25	16	22	19	26	45	54	52	37	17	5	-
White, both sexes	38	35	34	36	52	86	99	85	52	21	4	-
Male	22	23	15	17	29	45	46	33	15	4	-	-
Female	16	12	19	19	23	41	53	52	37	17	4	-
Black, both sexes	11	5	2	3	2	7	2	1	-	-	1	-
Male	4	3	-	3	1	3	1	1	-	-	-	-
Female	7	2	2	-	1	4	1	-	-	-	1	-
American Indian 1/, both sexes	3	2	1	1	2	-	-	-	-	-	-	-
Male	1	-	-	1	1	-	-	-	-	-	-	-
Female	2	2	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	1	-	1	-	-	-
Male	-	-	-	-	1	1	1	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	3	5	4	4	5	5	13	10	7	1	-	-
Male	1	4	2	2	2	4	7	2	3	1	-	-
Female	2	1	2	2	3	1	6	8	4	-	-	-
White, both sexes	3	4	4	3	5	5	12	10	7	1	-	-
Male	1	3	2	2	2	4	6	2	3	1	-	-
Female	2	1	2	1	3	1	6	8	4	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
United States												
All races, both sexes	4,391	14	15	1	1	1	2	9	4	11	21	39
Male	1,647	9	5	-	1	-	1	4	1	8	12	20
Female	2,744	5	10	1	-	1	1	5	3	3	9	19
White, both sexes	4,099	11	12	1	1	1	2	7	3	9	13	27
Male	1,519	7	3	-	1	-	1	3	1	6	7	13
Female	2,580	4	9	1	-	1	1	4	2	3	6	14
Black, both sexes	241	1	2	-	-	-	-	2	1	1	6	10
Male	99	1	1	-	-	-	-	1	-	1	3	6
Female	142	-	1	-	-	-	-	1	1	-	3	4
American Indian 1/, both sexes	18	1	-	-	-	-	-	-	-	1	1	1
Male	11	1	-	-	-	-	-	-	-	1	1	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	33	1	1	-	-	-	-	-	-	-	1	1
Male	18	-	1	-	-	-	-	-	-	-	1	-
Female	15	1	-	-	-	-	-	-	-	-	-	1
Alabama												
All races, both sexes	74	2	-	-	-	-	-	-	-	-	-	-
Male	27	1	-	-	-	-	-	-	-	-	-	-
Female	47	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	65	2	-	-	-	-	-	-	-	-	-	-
Male	24	1	-	-	-	-	-	-	-	-	-	-
Female	41	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
Alaska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	79	-	-	-	-	-	-	-	-	-	-	-
Male	29	-	-	-	-	-	-	-	-	-	-	-
Female	50	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	78	-	-	-	-	-	-	-	-	-	-	-
Male	29	-	-	-	-	-	-	-	-	-	-	-
Female	49	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
Arkansas												
All races, both sexes	56	-	1	-	-	-	-	-	-	-	1	1
Male	25	-	-	-	-	-	-	-	-	-	1	-
Female	31	-	1	-	-	-	-	-	-	-	-	1
White, both sexes	56	-	1	-	-	-	-	-	-	-	1	1
Male	25	-	-	-	-	-	-	-	-	-	1	-
Female	31	-	1	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	374	3	1	-	-	1	-	1	-	-	1	5
Male	145	1	1	-	-	-	-	1	-	-	1	3
Female	229	2	-	-	-	1	-	-	-	-	-	2
White, both sexes	338	2	-	-	-	1	-	1	-	-	1	5
Male	126	1	-	-	-	-	-	1	-	-	1	3
Female	212	1	-	-	-	1	-	-	-	-	-	2
Black, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	1	1	-	-	-	-	-	-	-	-	-
Male	11	-	1	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
Colorado												
All races, both sexes	32	-	-	-	1	-	-	-	-	-	-	1
Male	16	-	-	-	1	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	32	-	-	-	1	-	-	-	-	-	-	1
Male	16	-	-	-	1	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	58	-	-	-	-	-	-	-	-	-	1	1
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	41	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	58	-	-	-	-	-	-	-	-	-	1	1
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	41	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
Delaware												
All races, both sexes	16	1	-	-	-	-	-	-	-	-	-	-
Male	9	1	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	1	-	-	-	-	-	-	-	-	-	-
Male	9	1	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
Florida												
All races, both sexes	456	1	-	-	-	-	-	1	-	-	4	2
Male	192	1	-	-	-	-	-	1	-	-	2	1
Female	264	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	432	1	-	-	-	-	-	-	-	-	4	2
Male	185	1	-	-	-	-	-	-	-	-	2	1
Female	247	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	23	-	-	-	-	-	-	1	-	-	-	-
Male	7	-	-	-	-	-	-	1	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	76	1	2	-	-	-	-	-	-	1	-	3
Male	21	1	-	-	-	-	-	-	-	1	-	1
Female	55	-	2	-	-	-	-	-	-	-	-	2
White, both sexes	62	1	1	-	-	-	-	-	-	1	-	3
Male	16	1	-	-	-	-	-	-	-	1	-	1
Female	46	-	1	-	-	-	-	-	-	-	-	2
Black, both sexes	14	-	1	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
Hawaii												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
Illinois												
All races, both sexes	231	1	-	-	-	-	1	-	1	1	1	3
Male	81	-	-	-	-	-	-	-	-	1	1	1
Female	150	1	-	-	-	-	1	-	1	-	-	2
White, both sexes	213	1	-	-	-	-	1	-	1	-	-	1
Male	72	-	-	-	-	-	-	-	-	-	-	-
Female	141	1	-	-	-	-	1	-	1	-	-	1
Black, both sexes	16	-	-	-	-	-	-	-	-	1	1	2
Male	9	-	-	-	-	-	-	-	-	1	1	1
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	82	-	3	-	-	-	-	-	-	-	-	-
Male	36	-	2	-	-	-	-	-	-	-	-	-
Female	46	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	81	-	3	-	-	-	-	-	-	-	-	-
Male	35	-	2	-	-	-	-	-	-	-	-	-
Female	46	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
Iowa												
All races, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
Kentucky												
All races, both sexes	45	-	-	-	-	-	-	1	-	-	-	2
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	30	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	43	-	-	-	-	-	-	1	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	48	2	1	-	-	-	-	-	1	-	-	-
Male	15	2	1	-	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	1	1	-	-	-	-	-	1	-	-	-
Male	3	1	1	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
Maine												
All races, both sexes	88	-	-	-	-	-	-	-	-	-	-	-
Male	34	-	-	-	-	-	-	-	-	-	-	-
Female	54	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	85	-	-	-	-	-	-	-	-	-	-	-
Male	32	-	-	-	-	-	-	-	-	-	-	-
Female	53	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	98	-	-	-	-	-	-	-	-	-	1	1
Male	38	-	-	-	-	-	-	-	-	-	-	1
Female	60	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	91	-	-	-	-	-	-	-	-	-	-	1
Male	36	-	-	-	-	-	-	-	-	-	-	1
Female	55	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
Massachusetts												
All races, both sexes	107	-	-	-	-	-	-	1	-	-	1	1
Male	33	-	-	-	-	-	-	1	-	-	1	1
Female	74	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	104	-	-	-	-	-	-	1	-	-	1	1
Male	32	-	-	-	-	-	-	1	-	-	1	1
Female	72	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	112	-	-	-	-	-	-	-	1	-	1	2
Male	49	-	-	-	-	-	-	-	-	-	1	1
Female	63	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	104	-	-	-	-	-	-	-	1	-	1	1
Male	46	-	-	-	-	-	-	-	-	-	1	-
Female	58	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
Minnesota												
All races, both sexes	42	-	1	-	-	-	-	1	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	1	-	-	-	-	1	-	-	-	-
White, both sexes	41	-	1	-	-	-	-	1	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	1	-	-	-	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
Missouri												
All races, both sexes	177	-	1	-	-	-	-	-	-	-	1	2
Male	57	-	-	-	-	-	-	-	-	-	-	1
Female	120	-	1	-	-	-	-	-	-	-	1	1
White, both sexes	161	-	1	-	-	-	-	-	-	-	-	1
Male	49	-	-	-	-	-	-	-	-	-	-	-
Female	112	-	1	-	-	-	-	-	-	-	-	1
Black, both sexes	16	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
Nebraska												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
New Hampshire												
All races, both sexes	36	-	-	-	-	-	-	1	-	1	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	36	-	-	-	-	-	-	1	-	1	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	211	-	-	-	-	-	-	-	-	-	-	1
Male	68	-	-	-	-	-	-	-	-	-	-	-
Female	143	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	195	-	-	-	-	-	-	-	-	-	-	1
Male	60	-	-	-	-	-	-	-	-	-	-	-
Female	135	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
New Mexico												
All races, both sexes	13	-	1	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	1	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	222	-	1	-	-	-	-	-	1	2	1	1
Male	90	-	-	-	-	-	-	-	1	1	-	-
Female	132	-	1	-	-	-	-	-	-	1	1	1
White, both sexes	213	-	1	-	-	-	-	-	1	2	1	-
Male	86	-	-	-	-	-	-	-	1	1	-	-
Female	127	-	1	-	-	-	-	-	-	1	1	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
North Carolina												
All races, both sexes	92	-	-	-	-	-	-	-	-	-	-	-
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	66	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	78	-	-	-	-	-	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	57	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
Ohio												
All races, both sexes	379	-	-	-	-	-	-	-	-	2	-	5
Male	146	-	-	-	-	-	-	-	-	2	-	3
Female	233	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	356	-	-	-	-	-	-	-	-	2	-	4
Male	137	-	-	-	-	-	-	-	-	2	-	2
Female	219	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	53	1	-	-	-	-	-	-	-	-	1	-
Male	20	1	-	-	-	-	-	-	-	-	1	-
Female	33	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	52	1	-	-	-	-	-	-	-	-	-	-
Male	19	1	-	-	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
Oregon												
All races, both sexes	45	-	-	-	-	-	-	1	-	1	-	1
Male	18	-	-	-	-	-	-	-	-	1	-	-
Female	27	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	43	-	-	-	-	-	-	1	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Pennsylvania												
All races, both sexes	365	1	-	-	-	-	1	2	-	2	1	1
Male	125	1	-	-	-	-	1	1	-	1	1	-
Female	240	-	-	-	-	-	-	1	-	1	-	1
White, both sexes	344	1	-	-	-	-	1	1	-	2	-	-
Male	118	1	-	-	-	-	1	1	-	1	-	-
Female	226	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	18	-	-	-	-	-	-	1	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
Rhode Island												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	37	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
South Dakota												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	112	1	-	-	-	-	-	-	-	1	-	-
Male	41	-	-	-	-	-	-	-	-	1	-	-
Female	71	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	108	1	-	-	-	-	-	-	-	1	-	-
Male	39	-	-	-	-	-	-	-	-	1	-	-
Female	69	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
Texas												
All races, both sexes	147	-	2	-	-	-	-	-	-	-	2	2
Male	55	-	1	-	-	-	-	-	-	-	1	2
Female	92	-	1	-	-	-	-	-	-	-	1	-
White, both sexes	130	-	2	-	-	-	-	-	-	-	-	1
Male	49	-	1	-	-	-	-	-	-	-	-	1
Female	81	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	15	-	-	-	-	-	-	-	-	-	2	1
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	9	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
Vermont												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	77	-	-	-	-	-	-	-	-	-	1	2
Male	35	-	-	-	-	-	-	-	-	-	1	1
Female	42	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	70	-	-	-	-	-	-	-	-	-	1	1
Male	31	-	-	-	-	-	-	-	-	-	1	1
Female	39	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
Washington												
All races, both sexes	64	-	-	-	-	-	-	-	-	-	1	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	42	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	64	-	-	-	-	-	-	-	-	-	1	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	42	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	29	-	1	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	1	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infectious diseases (A00-A09)												
Wisconsin												
All races, both sexes	49	-	-	1	-	-	-	-	-	-	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	1	-	-	-	-	-	-	-	1
White, both sexes	46	-	-	1	-	-	-	-	-	-	-	1
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	1	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
United States												
All races, both sexes	58	96	160	221	396	694	975	877	590	181	25	-
Male	31	48	81	104	165	291	380	297	153	35	1	-
Female	27	48	79	117	231	403	595	580	437	146	24	-
White, both sexes	51	82	143	194	365	648	921	837	570	177	24	-
Male	26	36	74	90	152	273	361	285	145	34	1	-
Female	25	46	69	104	213	375	560	552	425	143	23	-
Black, both sexes	7	14	13	22	25	35	46	35	16	4	1	-
Male	5	12	5	12	10	11	15	10	5	1	-	-
Female	2	2	8	10	15	24	31	25	11	3	1	-
American Indian 1/, both sexes	-	-	2	3	3	4	2	-	-	-	-	-
Male	-	-	-	1	1	3	2	-	-	-	-	-
Female	-	-	2	2	2	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	2	3	7	6	5	4	-	-	-
Male	-	-	2	1	2	4	2	2	3	-	-	-
Female	-	-	-	1	1	3	4	3	1	-	-	-
Alabama												
All races, both sexes	-	3	5	5	13	11	12	17	5	-	1	-
Male	-	3	3	1	6	2	4	5	2	-	-	-
Female	-	-	2	4	7	9	8	12	3	-	1	-
White, both sexes	-	1	4	5	11	8	12	17	4	-	1	-
Male	-	1	3	1	6	2	4	5	1	-	-	-
Female	-	-	1	4	5	6	8	12	3	-	1	-
Black, both sexes	-	2	1	-	2	3	-	-	1	-	-	-
Male	-	2	-	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	2	3	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	5	5	4	10	10	15	19	8	3	-	-
Male	-	4	2	1	2	3	4	10	2	1	-	-
Female	-	1	3	3	8	7	11	9	6	2	-	-
White, both sexes	-	5	5	3	10	10	15	19	8	3	-	-
Male	-	4	2	1	2	3	4	10	2	1	-	-
Female	-	1	3	2	8	7	11	9	6	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
Arkansas												
All races, both sexes	2	-	1	2	4	12	12	10	7	3	-	-
Male	1	-	1	2	2	8	5	4	1	-	-	-
Female	1	-	-	-	2	4	7	6	6	3	-	-
White, both sexes	2	-	1	2	4	12	12	10	7	3	-	-
Male	1	-	1	2	2	8	5	4	1	-	-	-
Female	1	-	-	-	2	4	7	6	6	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	6	11	17	21	29	59	76	70	52	21	-	-
Male	1	4	9	9	12	27	27	28	14	7	-	-
Female	5	7	8	12	17	32	49	42	38	14	-	-
White, both sexes	5	9	14	17	27	52	70	64	49	21	-	-
Male	1	3	8	7	10	23	25	25	11	7	-	-
Female	4	6	6	10	17	29	45	39	38	14	-	-
Black, both sexes	1	2	1	4	-	2	3	3	1	-	-	-
Male	-	1	-	2	-	1	1	1	1	-	-	-
Female	1	1	1	2	-	1	2	2	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	2	4	3	3	2	-	-	-
Male	-	-	1	-	2	2	1	2	2	-	-	-
Female	-	-	-	-	-	2	2	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
Colorado												
All races, both sexes	-	-	1	2	2	4	11	5	3	1	1	-
Male	-	-	-	2	2	2	4	3	-	1	-	-
Female	-	-	1	-	-	2	7	2	3	-	1	-
White, both sexes	-	-	1	2	2	4	11	5	3	1	1	-
Male	-	-	-	2	2	2	4	3	-	1	-	-
Female	-	-	1	-	-	2	7	2	3	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	1	3	2	10	14	14	11	-	1	-
Male	-	-	1	1	-	4	3	4	3	-	-	-
Female	-	-	-	2	2	6	11	10	8	-	1	-
White, both sexes	-	-	1	3	2	10	14	14	11	-	1	-
Male	-	-	1	1	-	4	3	4	3	-	-	-
Female	-	-	-	2	2	6	11	10	8	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
Delaware												
All races, both sexes	-	-	-	1	1	3	5	-	3	2	-	-
Male	-	-	-	1	1	1	3	-	1	1	-	-
Female	-	-	-	-	-	2	2	-	2	1	-	-
White, both sexes	-	-	-	1	1	3	5	-	3	2	-	-
Male	-	-	-	1	1	1	3	-	1	1	-	-
Female	-	-	-	-	-	2	2	-	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	1	1	-	-	1	1	1	-	-	-	-
Male	-	1	-	-	-	-	1	1	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	1	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
Florida												
All races, both sexes	3	7	14	20	37	78	104	118	53	13	1	-
Male	2	2	7	12	21	37	43	45	16	2	-	-
Female	1	5	7	8	16	41	61	73	37	11	1	-
White, both sexes	3	7	11	17	35	75	101	111	51	13	1	-
Male	2	2	7	10	21	35	42	44	16	2	-	-
Female	1	5	4	7	14	40	59	67	35	11	1	-
Black, both sexes	-	-	3	3	2	3	3	6	2	-	-	-
Male	-	-	-	2	-	2	1	1	-	-	-	-
Female	-	-	3	1	2	1	2	5	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Georgia												
All races, both sexes	1	2	3	5	11	11	17	13	4	2	-	-
Male	1	1	-	-	7	4	5	-	-	-	-	-
Female	-	1	3	5	4	7	12	13	4	2	-	-
White, both sexes	1	2	3	3	7	10	12	12	4	2	-	-
Male	1	1	-	-	3	4	4	-	-	-	-	-
Female	-	1	3	3	4	6	8	12	4	2	-	-
Black, both sexes	-	-	-	2	4	1	5	1	-	-	-	-
Male	-	-	-	-	4	-	1	-	-	-	-	-
Female	-	-	-	2	-	1	4	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
Hawaii												
All races, both sexes	-	-	1	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	1	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Idaho												
All races, both sexes	-	1	1	1	-	2	2	2	-	1	-	-
Male	-	-	1	1	-	1	1	1	-	-	-	-
Female	-	1	-	-	-	1	1	1	-	1	-	-
White, both sexes	-	1	1	1	-	2	2	2	-	1	-	-
Male	-	-	1	1	-	1	1	1	-	-	-	-
Female	-	1	-	-	-	1	1	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
Illinois												
All races, both sexes	2	9	7	9	23	32	35	49	43	13	1	-
Male	1	5	4	4	8	11	16	14	13	2	-	-
Female	1	4	3	5	15	21	19	35	30	11	1	-
White, both sexes	2	7	7	7	23	30	32	46	41	13	1	-
Male	1	3	4	3	8	11	15	13	12	2	-	-
Female	1	4	3	4	15	19	17	33	29	11	1	-
Black, both sexes	-	2	-	2	-	2	3	2	1	-	-	-
Male	-	2	-	1	-	-	1	1	1	-	-	-
Female	-	-	-	1	-	2	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Indiana												
All races, both sexes	2	1	2	4	8	20	13	14	12	2	1	-
Male	1	-	2	3	3	10	7	3	4	1	-	-
Female	1	1	-	1	5	10	6	11	8	1	1	-
White, both sexes	2	1	2	4	8	20	12	14	12	2	1	-
Male	1	-	2	3	3	10	6	3	4	1	-	-
Female	1	1	-	1	5	10	6	11	8	1	1	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
Iowa												
All races, both sexes	-	-	1	2	1	5	12	7	5	1	-	-
Male	-	-	1	1	1	1	5	2	1	1	-	-
Female	-	-	-	1	-	4	7	5	4	-	-	-
White, both sexes	-	-	1	2	1	5	12	7	5	1	-	-
Male	-	-	1	1	1	1	5	2	1	1	-	-
Female	-	-	-	1	-	4	7	5	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	-	-	-	2	4	6	6	3	1	1	-
Male	1	-	-	-	1	-	1	2	1	-	-	-
Female	-	-	-	-	1	4	5	4	2	1	1	-
White, both sexes	1	-	-	-	2	4	6	6	3	1	1	-
Male	1	-	-	-	1	-	1	2	1	-	-	-
Female	-	-	-	-	1	4	5	4	2	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
Kentucky												
All races, both sexes	-	1	-	2	3	9	11	9	6	-	1	-
Male	-	-	-	-	2	4	4	4	-	-	-	-
Female	-	1	-	2	1	5	7	5	6	-	1	-
White, both sexes	-	1	-	2	3	9	11	9	5	-	1	-
Male	-	-	-	-	2	4	4	4	-	-	-	-
Female	-	1	-	2	1	5	7	5	5	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	2	3	3	3	4	14	5	5	3	1	-
Male	-	2	1	1	-	2	4	-	1	1	-	-
Female	1	-	2	2	3	2	10	5	4	2	1	-
White, both sexes	1	2	2	3	2	4	12	5	4	3	-	-
Male	-	2	-	1	-	2	4	-	1	1	-	-
Female	1	-	2	2	2	2	8	5	3	2	-	-
Black, both sexes	-	-	1	-	1	-	2	-	1	-	1	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
Maine												
All races, both sexes	-	-	1	1	7	14	28	18	12	6	1	-
Male	-	-	-	1	3	7	13	6	4	-	-	-
Female	-	-	1	-	4	7	15	12	8	6	1	-
White, both sexes	-	-	1	1	6	13	27	18	12	6	1	-
Male	-	-	-	1	3	6	12	6	4	-	-	-
Female	-	-	1	-	3	7	15	12	8	6	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	2	5	7	9	9	33	14	12	4	-	-
Male	1	1	3	5	4	3	10	6	4	-	-	-
Female	-	1	2	2	5	6	23	8	8	4	-	-
White, both sexes	1	2	5	6	8	9	32	11	12	4	-	-
Male	1	1	3	4	4	3	10	5	4	-	-	-
Female	-	1	2	2	4	6	22	6	8	4	-	-
Black, both sexes	-	-	-	-	1	-	1	3	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
Massachusetts												
All races, both sexes	2	1	2	3	4	24	24	21	17	6	-	-
Male	1	1	2	2	1	5	7	5	4	2	-	-
Female	1	-	-	1	3	19	17	16	13	4	-	-
White, both sexes	2	-	2	3	4	22	24	21	17	6	-	-
Male	1	-	2	2	1	5	7	5	4	2	-	-
Female	1	-	-	1	3	17	17	16	13	4	-	-
Black, both sexes	-	1	-	-	-	2	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	3	4	4	6	13	18	25	25	9	1	-	-
Male	3	2	3	4	8	5	11	8	3	-	-	-
Female	-	2	1	2	5	13	14	17	6	1	-	-
White, both sexes	2	4	4	5	12	17	25	23	8	1	-	-
Male	2	2	3	4	8	5	11	8	2	-	-	-
Female	-	2	1	1	4	12	14	15	6	1	-	-
Black, both sexes	1	-	-	1	-	1	-	2	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	1	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
Minnesota												
All races, both sexes	-	1	1	3	5	5	7	7	8	3	-	-
Male	-	-	-	2	-	3	2	4	4	1	-	-
Female	-	1	1	1	5	2	5	3	4	2	-	-
White, both sexes	-	1	1	3	5	5	6	7	8	3	-	-
Male	-	-	-	2	-	3	1	4	4	1	-	-
Female	-	1	1	1	5	2	5	3	4	2	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	1	1	2	-	6	5	1	2	-	-
Male	-	-	1	-	1	-	3	2	-	-	-	-
Female	-	-	-	1	1	-	3	3	1	2	-	-
White, both sexes	-	-	1	1	2	-	6	4	1	2	-	-
Male	-	-	1	-	1	-	3	1	-	-	-	-
Female	-	-	-	1	1	-	3	3	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
Missouri												
All races, both sexes	4	3	8	6	9	39	28	38	26	11	1	-
Male	2	1	4	4	4	13	11	11	4	2	-	-
Female	2	2	4	2	5	26	17	27	22	9	1	-
White, both sexes	2	3	7	4	7	36	27	36	26	10	1	-
Male	-	1	4	2	4	12	10	10	4	2	-	-
Female	2	2	3	2	3	24	17	26	22	8	1	-
Black, both sexes	2	-	1	2	2	3	1	2	-	1	-	-
Male	2	-	-	2	-	1	1	1	-	-	-	-
Female	-	-	1	-	2	2	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	1	-	-	1	3	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	1	-	-	-	2	-	-	-	-
White, both sexes	-	-	-	1	-	-	1	3	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	1	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
Nebraska												
All races, both sexes	-	1	1	1	1	1	5	2	4	-	-	-
Male	-	-	1	1	-	-	3	-	-	-	-	-
Female	-	1	-	-	1	1	2	2	4	-	-	-
White, both sexes	-	1	1	1	1	1	5	2	4	-	-	-
Male	-	-	1	1	-	-	3	-	-	-	-	-
Female	-	1	-	-	1	1	2	2	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	2	1	-	2	6	4	1	1	-	-	-
Male	1	1	-	-	-	3	-	-	1	-	-	-
Female	-	1	1	-	2	3	4	1	-	-	-	-
White, both sexes	1	2	1	-	2	6	4	1	1	-	-	-
Male	1	1	-	-	-	3	-	-	1	-	-	-
Female	-	1	1	-	2	3	4	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
New Hampshire												
All races, both sexes	-	1	1	1	7	5	6	7	4	1	1	-
Male	-	-	1	1	2	1	3	5	-	-	-	-
Female	-	1	-	-	5	4	3	2	4	1	1	-
White, both sexes	-	1	1	1	7	5	6	7	4	1	1	-
Male	-	-	1	1	2	1	3	5	-	-	-	-
Female	-	1	-	-	5	4	3	2	4	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	5	3	6	8	17	37	35	41	46	10	2	-
Male	3	1	4	5	4	11	16	10	14	-	-	-
Female	2	2	2	3	13	26	19	31	32	10	2	-
White, both sexes	4	2	3	7	16	33	32	40	45	10	2	-
Male	3	-	2	4	4	10	15	9	13	-	-	-
Female	1	2	1	3	12	23	17	31	32	10	2	-
Black, both sexes	1	1	2	1	1	3	3	1	1	-	-	-
Male	-	1	1	1	-	-	1	1	1	-	-	-
Female	1	-	1	-	1	3	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
New Mexico												
All races, both sexes	1	-	1	1	1	3	1	2	1	1	-	-
Male	1	-	-	1	1	1	1	2	-	1	-	-
Female	-	-	1	-	-	2	-	-	1	-	-	-
White, both sexes	1	-	1	-	1	3	1	2	1	1	-	-
Male	1	-	-	-	1	1	1	2	-	1	-	-
Female	-	-	1	-	-	2	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	3	1	10	11	17	23	53	50	35	11	2	-
Male	1	1	4	7	9	12	26	16	9	3	-	-
Female	2	-	6	4	8	11	27	34	26	8	2	-
White, both sexes	3	1	10	10	16	21	51	49	34	11	2	-
Male	1	1	4	7	8	11	24	16	9	3	-	-
Female	2	-	6	3	8	10	27	33	25	8	2	-
Black, both sexes	-	-	-	1	1	1	2	1	1	-	-	-
Male	-	-	-	-	1	-	2	-	-	-	-	-
Female	-	-	-	1	-	1	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
North Carolina												
All races, both sexes	1	3	2	6	9	19	24	15	10	2	1	-
Male	-	2	1	2	3	4	6	6	2	-	-	-
Female	1	1	1	4	6	15	18	9	8	2	1	-
White, both sexes	1	2	1	5	8	15	20	14	9	2	1	-
Male	-	1	1	1	2	2	6	6	2	-	-	-
Female	1	1	-	4	6	13	14	8	7	2	1	-
Black, both sexes	-	1	1	1	1	4	4	1	1	-	-	-
Male	-	1	-	1	1	2	-	-	-	-	-	-
Female	-	-	1	-	-	2	4	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	1	1	1	-	1	1	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	1	1	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	1	1	-	1	1	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
Ohio												
All races, both sexes	5	8	17	23	39	59	92	77	42	9	1	-
Male	2	4	8	7	13	30	36	28	12	1	-	-
Female	3	4	9	16	26	29	56	49	30	8	1	-
White, both sexes	5	7	14	23	36	57	86	73	39	9	1	-
Male	2	3	5	7	12	28	35	28	12	1	-	-
Female	3	4	9	16	24	29	51	45	27	8	1	-
Black, both sexes	-	1	3	-	3	1	6	4	3	-	-	-
Male	-	1	3	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	2	-	5	4	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	1	2	2	9	6	13	6	9	1	-	-
Male	1	-	1	1	3	3	5	2	2	-	-	-
Female	1	1	1	1	6	3	8	4	7	1	-	-
White, both sexes	2	1	2	2	9	6	13	6	9	1	-	-
Male	1	-	1	1	3	3	5	2	2	-	-	-
Female	1	1	1	1	6	3	8	4	7	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
Oregon												
All races, both sexes	1	-	1	2	1	1	18	11	3	4	-	-
Male	-	-	1	-	1	1	9	5	-	-	-	-
Female	1	-	-	2	-	-	9	6	3	4	-	-
White, both sexes	1	-	1	2	1	1	18	11	3	4	-	-
Male	-	-	1	-	1	1	9	5	-	-	-	-
Female	1	-	-	2	-	-	9	6	3	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	2	12	8	15	33	55	87	76	52	14	3	-
Male	-	6	3	4	17	22	37	16	11	3	1	-
Female	2	6	5	11	16	33	50	60	41	11	2	-
White, both sexes	2	10	8	13	32	52	80	73	52	14	3	-
Male	-	5	3	4	17	21	34	15	11	3	1	-
Female	2	5	5	9	15	31	46	58	41	11	2	-
Black, both sexes	-	2	-	2	1	2	5	3	-	-	-	-
Male	-	1	-	-	-	1	2	1	-	-	-	-
Female	-	1	-	2	1	1	3	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
Rhode Island												
All races, both sexes	1	1	-	-	1	5	3	7	3	2	-	-
Male	1	1	-	-	1	2	-	4	1	1	-	-
Female	-	-	-	-	-	3	3	3	2	1	-	-
White, both sexes	1	1	-	-	1	5	3	7	3	2	-	-
Male	1	1	-	-	1	2	-	4	1	1	-	-
Female	-	-	-	-	-	3	3	3	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	2	-	5	-	9	5	5	7	3	-	-
Male	1	1	-	3	-	3	3	3	1	-	-	-
Female	-	1	-	2	-	6	2	2	6	3	-	-
White, both sexes	1	2	-	3	-	9	4	4	6	1	-	-
Male	1	1	-	1	-	3	2	2	1	-	-	-
Female	-	1	-	2	-	6	2	2	5	1	-	-
Black, both sexes	-	-	-	2	-	-	1	1	1	2	-	-
Male	-	-	-	2	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
South Dakota												
All races, both sexes	-	-	-	1	1	-	2	1	2	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	-	1	-	-	1	-	2	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	2	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	4	4	9	14	18	23	14	19	2	1	-
Male	1	3	1	3	6	12	5	3	6	-	-	-
Female	1	1	3	6	8	6	18	11	13	2	1	-
White, both sexes	2	3	4	8	14	18	23	12	19	2	1	-
Male	1	2	1	2	6	12	5	3	6	-	-	-
Female	1	1	3	6	8	6	18	9	13	2	1	-
Black, both sexes	-	1	-	1	-	-	-	2	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
Texas												
All races, both sexes	2	-	3	10	21	30	30	23	14	7	1	-
Male	2	-	3	6	6	14	11	7	-	2	-	-
Female	-	-	-	4	15	16	19	16	14	5	1	-
White, both sexes	2	-	3	10	16	27	27	22	13	6	1	-
Male	2	-	3	6	4	13	11	7	-	1	-	-
Female	-	-	-	4	12	14	16	15	13	5	1	-
Black, both sexes	-	-	-	-	4	2	3	1	1	1	-	-
Male	-	-	-	-	2	1	-	-	-	1	-	-
Female	-	-	-	-	2	1	3	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	1	1	2	2	2	2	1	1	-	-
Male	-	-	1	-	1	2	-	1	-	-	-	-
Female	-	-	-	1	1	-	2	1	1	1	-	-
White, both sexes	-	-	1	1	2	2	2	2	1	1	-	-
Male	-	-	1	-	1	2	-	1	-	-	-	-
Female	-	-	-	1	1	-	2	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
Vermont												
All races, both sexes	-	-	2	-	1	2	5	7	-	5	-	-
Male	-	-	-	-	-	1	2	-	-	1	-	-
Female	-	-	2	-	1	1	3	7	-	4	-	-
White, both sexes	-	-	2	-	1	2	5	7	-	5	-	-
Male	-	-	-	-	-	1	2	-	-	1	-	-
Female	-	-	2	-	1	1	3	7	-	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	1	4	4	7	7	17	18	10	5	-	-
Male	1	-	2	2	4	6	6	9	2	1	-	-
Female	-	1	2	2	3	1	11	9	8	4	-	-
White, both sexes	-	1	4	4	5	5	16	18	10	5	-	-
Male	-	-	2	2	3	5	5	9	2	1	-	-
Female	-	1	2	2	2	-	11	9	8	4	-	-
Black, both sexes	1	-	-	-	2	2	1	-	-	-	-	-
Male	1	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
Washington												
All races, both sexes	1	-	3	2	6	10	17	9	12	2	1	-
Male	-	-	-	1	2	4	5	3	7	-	-	-
Female	1	-	3	1	4	6	12	6	5	2	1	-
White, both sexes	1	-	3	2	6	10	17	9	12	2	1	-
Male	-	-	-	1	2	4	5	3	7	-	-	-
Female	1	-	3	1	4	6	12	6	5	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	2	5	1	3	5	6	4	1	-	-	-
Male	1	1	3	-	-	3	4	1	-	-	-	-
Female	-	1	2	1	3	2	2	3	1	-	-	-
White, both sexes	-	2	5	1	3	5	5	4	1	-	-	-
Male	-	1	3	-	-	3	4	1	-	-	-	-
Female	-	1	2	1	3	2	1	3	1	-	-	-
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
Wisconsin												
All races, both sexes	-	-	2	3	4	7	13	7	8	2	1	-
Male	-	-	2	2	1	3	6	5	2	-	-	-
Female	-	-	-	1	3	4	7	2	6	2	1	-
White, both sexes	-	-	2	3	4	5	13	7	7	2	1	-
Male	-	-	2	2	1	2	6	5	1	-	-	-
Female	-	-	-	1	3	3	7	2	6	2	1	-
Black, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
United States												
All races, both sexes	4,280	14	14	-	1	1	2	9	3	11	21	39
Male	1,612	9	4	-	1	-	1	4	1	8	12	20
Female	2,668	5	10	-	-	1	1	5	2	3	9	19
White, both sexes	4,001	11	12	-	1	1	2	7	3	9	13	27
Male	1,491	7	3	-	1	-	1	3	1	6	7	13
Female	2,510	4	9	-	-	1	1	4	2	3	6	14
Black, both sexes	230	1	1	-	-	-	-	2	-	1	6	10
Male	93	1	-	-	-	-	-	1	-	1	3	6
Female	137	-	1	-	-	-	-	1	-	-	3	4
American Indian 1/, both sexes	16	1	-	-	-	-	-	-	-	1	1	1
Male	10	1	-	-	-	-	-	-	-	1	1	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	33	1	1	-	-	-	-	-	-	-	1	1
Male	18	-	1	-	-	-	-	-	-	-	1	-
Female	15	1	-	-	-	-	-	-	-	-	-	1
Alabama												
All races, both sexes	73	2	-	-	-	-	-	-	-	-	-	-
Male	26	1	-	-	-	-	-	-	-	-	-	-
Female	47	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	64	2	-	-	-	-	-	-	-	-	-	-
Male	23	1	-	-	-	-	-	-	-	-	-	-
Female	41	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
Alaska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	76	-	-	-	-	-	-	-	-	-	-	-
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	48	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	75	-	-	-	-	-	-	-	-	-	-	-
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	47	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
Arkansas												
All races, both sexes	56	-	1	-	-	-	-	-	-	-	1	1
Male	25	-	-	-	-	-	-	-	-	-	1	-
Female	31	-	1	-	-	-	-	-	-	-	-	1
White, both sexes	56	-	1	-	-	-	-	-	-	-	1	1
Male	25	-	-	-	-	-	-	-	-	-	1	-
Female	31	-	1	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	365	3	1	-	-	1	-	1	-	-	1	5
Male	143	1	1	-	-	-	-	1	-	-	1	3
Female	222	2	-	-	-	1	-	-	-	-	-	2
White, both sexes	329	2	-	-	-	1	-	1	-	-	1	5
Male	124	1	-	-	-	-	-	1	-	-	1	3
Female	205	1	-	-	-	1	-	-	-	-	-	2
Black, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	1	1	-	-	-	-	-	-	-	-	-
Male	11	-	1	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
Colorado												
All races, both sexes	32	-	-	-	1	-	-	-	-	-	-	1
Male	16	-	-	-	1	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	32	-	-	-	1	-	-	-	-	-	-	1
Male	16	-	-	-	1	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	57	-	-	-	-	-	-	-	-	-	1	1
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	40	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	57	-	-	-	-	-	-	-	-	-	1	1
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	40	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
Delaware												
All races, both sexes	16	1	-	-	-	-	-	-	-	-	-	-
Male	9	1	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	1	-	-	-	-	-	-	-	-	-	-
Male	9	1	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
Florida												
All races, both sexes	446	1	-	-	-	-	-	1	-	-	4	2
Male	189	1	-	-	-	-	-	1	-	-	2	1
Female	257	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	422	1	-	-	-	-	-	-	-	-	4	2
Male	182	1	-	-	-	-	-	-	-	-	2	1
Female	240	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	23	-	-	-	-	-	-	1	-	-	-	-
Male	7	-	-	-	-	-	-	1	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	74	1	2	-	-	-	-	-	-	1	-	3
Male	21	1	-	-	-	-	-	-	-	1	-	1
Female	53	-	2	-	-	-	-	-	-	-	-	2
White, both sexes	61	1	1	-	-	-	-	-	-	1	-	3
Male	16	1	-	-	-	-	-	-	-	1	-	1
Female	45	-	1	-	-	-	-	-	-	-	-	2
Black, both sexes	13	-	1	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
Hawaii												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
Illinois												
All races, both sexes	221	1	-	-	-	-	1	-	1	1	1	3
Male	77	-	-	-	-	-	-	-	-	1	1	1
Female	144	1	-	-	-	-	1	-	1	-	-	2
White, both sexes	206	1	-	-	-	-	1	-	1	-	-	1
Male	71	-	-	-	-	-	-	-	-	-	-	-
Female	135	1	-	-	-	-	1	-	1	-	-	1
Black, both sexes	13	-	-	-	-	-	-	-	-	1	1	2
Male	6	-	-	-	-	-	-	-	-	1	1	1
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	81	-	3	-	-	-	-	-	-	-	-	-
Male	35	-	2	-	-	-	-	-	-	-	-	-
Female	46	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	80	-	3	-	-	-	-	-	-	-	-	-
Male	34	-	2	-	-	-	-	-	-	-	-	-
Female	46	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
Iowa												
All races, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
Kentucky												
All races, both sexes	44	-	-	-	-	-	-	1	-	-	-	2
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	29	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	42	-	-	-	-	-	-	1	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	46	2	-	-	-	-	-	-	-	-	-	-
Male	14	2	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
Maine												
All races, both sexes	85	-	-	-	-	-	-	-	-	-	-	-
Male	33	-	-	-	-	-	-	-	-	-	-	-
Female	52	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	83	-	-	-	-	-	-	-	-	-	-	-
Male	32	-	-	-	-	-	-	-	-	-	-	-
Female	51	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	94	-	-	-	-	-	-	-	-	-	1	1
Male	36	-	-	-	-	-	-	-	-	-	-	1
Female	58	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	87	-	-	-	-	-	-	-	-	-	-	1
Male	34	-	-	-	-	-	-	-	-	-	-	1
Female	53	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
Massachusetts												
All races, both sexes	104	-	-	-	-	-	-	1	-	-	1	1
Male	31	-	-	-	-	-	-	1	-	-	1	1
Female	73	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	102	-	-	-	-	-	-	1	-	-	1	1
Male	31	-	-	-	-	-	-	1	-	-	1	1
Female	71	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	110	-	-	-	-	-	-	-	1	-	1	2
Male	49	-	-	-	-	-	-	-	-	-	1	1
Female	61	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	103	-	-	-	-	-	-	-	1	-	1	1
Male	46	-	-	-	-	-	-	-	-	-	1	-
Female	57	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
Minnesota												
All races, both sexes	41	-	1	-	-	-	-	1	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	1	-	-	-	-	1	-	-	-	-
White, both sexes	40	-	1	-	-	-	-	1	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	1	-	-	-	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
Missouri												
All races, both sexes	169	-	1	-	-	-	-	-	-	-	1	2
Male	57	-	-	-	-	-	-	-	-	-	-	1
Female	112	-	1	-	-	-	-	-	-	-	1	1
White, both sexes	154	-	1	-	-	-	-	-	-	-	-	1
Male	49	-	-	-	-	-	-	-	-	-	-	-
Female	105	-	1	-	-	-	-	-	-	-	-	1
Black, both sexes	15	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
Nebraska												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
New Hampshire												
All races, both sexes	36	-	-	-	-	-	-	1	-	1	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	36	-	-	-	-	-	-	1	-	1	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	206	-	-	-	-	-	-	-	-	-	-	1
Male	67	-	-	-	-	-	-	-	-	-	-	-
Female	139	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	190	-	-	-	-	-	-	-	-	-	-	1
Male	59	-	-	-	-	-	-	-	-	-	-	-
Female	131	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
New Mexico												
All races, both sexes	13	-	1	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	1	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	216	-	1	-	-	-	-	-	1	2	1	1
Male	88	-	-	-	-	-	-	-	1	1	-	-
Female	128	-	1	-	-	-	-	-	-	1	1	1
White, both sexes	208	-	1	-	-	-	-	-	1	2	1	-
Male	84	-	-	-	-	-	-	-	1	1	-	-
Female	124	-	1	-	-	-	-	-	-	1	1	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
North Carolina												
All races, both sexes	87	-	-	-	-	-	-	-	-	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	62	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	73	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	53	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
Ohio												
All races, both sexes	372	-	-	-	-	-	-	-	-	2	-	5
Male	142	-	-	-	-	-	-	-	-	2	-	3
Female	230	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	349	-	-	-	-	-	-	-	-	2	-	4
Male	133	-	-	-	-	-	-	-	-	2	-	2
Female	216	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	52	1	-	-	-	-	-	-	-	-	1	-
Male	20	1	-	-	-	-	-	-	-	-	1	-
Female	32	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	51	1	-	-	-	-	-	-	-	-	-	-
Male	19	1	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
Oregon												
All races, both sexes	45	-	-	-	-	-	-	1	-	1	-	1
Male	18	-	-	-	-	-	-	-	-	1	-	-
Female	27	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	43	-	-	-	-	-	-	1	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Pennsylvania												
All races, both sexes	359	1	-	-	-	-	1	2	-	2	1	1
Male	123	1	-	-	-	-	1	1	-	1	1	-
Female	236	-	-	-	-	-	-	1	-	1	-	1
White, both sexes	338	1	-	-	-	-	1	1	-	2	-	-
Male	116	1	-	-	-	-	1	1	-	1	-	-
Female	222	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	18	-	-	-	-	-	-	1	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
Rhode Island												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
South Dakota												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	111	1	-	-	-	-	-	-	-	1	-	-
Male	41	-	-	-	-	-	-	-	-	1	-	-
Female	70	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	107	1	-	-	-	-	-	-	-	1	-	-
Male	39	-	-	-	-	-	-	-	-	1	-	-
Female	68	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
Texas												
All races, both sexes	141	-	2	-	-	-	-	-	-	-	2	2
Male	52	-	1	-	-	-	-	-	-	-	1	2
Female	89	-	1	-	-	-	-	-	-	-	1	-
White, both sexes	124	-	2	-	-	-	-	-	-	-	-	1
Male	46	-	1	-	-	-	-	-	-	-	-	1
Female	78	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	15	-	-	-	-	-	-	-	-	-	2	1
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	9	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
Vermont												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	74	-	-	-	-	-	-	-	-	-	1	2
Male	34	-	-	-	-	-	-	-	-	-	1	1
Female	40	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	67	-	-	-	-	-	-	-	-	-	1	1
Male	30	-	-	-	-	-	-	-	-	-	1	1
Female	37	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
Washington												
All races, both sexes	63	-	-	-	-	-	-	-	-	-	1	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	63	-	-	-	-	-	-	-	-	-	1	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	28	-	1	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	1	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other intestinal infectious diseases (A01-A08)												
Wisconsin												
All races, both sexes	48	-	-	-	-	-	-	-	-	-	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	45	-	-	-	-	-	-	-	-	-	-	1
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
United States												
All races, both sexes	54	93	157	215	386	675	951	858	576	175	25	-
Male	29	45	80	101	163	285	371	291	151	35	1	-
Female	25	48	77	114	223	390	580	567	425	140	24	-
White, both sexes	48	81	140	188	356	632	898	819	557	172	24	-
Male	25	35	73	87	150	268	352	280	144	34	1	-
Female	23	46	67	101	206	364	546	539	413	138	23	-
Black, both sexes	6	12	13	22	25	33	45	34	15	3	1	-
Male	4	10	5	12	10	11	15	9	4	1	-	-
Female	2	2	8	10	15	22	30	25	11	2	1	-
American Indian 1/, both sexes	-	-	2	3	2	3	2	-	-	-	-	-
Male	-	-	-	1	1	2	2	-	-	-	-	-
Female	-	-	2	2	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	2	3	7	6	5	4	-	-	-
Male	-	-	2	1	2	4	2	2	3	-	-	-
Female	-	-	-	1	1	3	4	3	1	-	-	-
Alabama												
All races, both sexes	-	3	5	5	13	11	11	17	5	-	1	-
Male	-	3	3	1	6	2	3	5	2	-	-	-
Female	-	-	2	4	7	9	8	12	3	-	1	-
White, both sexes	-	1	4	5	11	8	11	17	4	-	1	-
Male	-	1	3	1	6	2	3	5	1	-	-	-
Female	-	-	1	4	5	6	8	12	3	-	1	-
Black, both sexes	-	2	1	-	2	3	-	-	1	-	-	-
Male	-	2	-	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	2	3	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	5	5	4	10	10	14	17	8	3	-	-
Male	-	4	2	1	2	3	4	9	2	1	-	-
Female	-	1	3	3	8	7	10	8	6	2	-	-
White, both sexes	-	5	5	3	10	10	14	17	8	3	-	-
Male	-	4	2	1	2	3	4	9	2	1	-	-
Female	-	1	3	2	8	7	10	8	6	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
Arkansas												
All races, both sexes	2	-	1	2	4	12	12	10	7	3	-	-
Male	1	-	1	2	2	8	5	4	1	-	-	-
Female	1	-	-	-	2	4	7	6	6	3	-	-
White, both sexes	2	-	1	2	4	12	12	10	7	3	-	-
Male	1	-	1	2	2	8	5	4	1	-	-	-
Female	1	-	-	-	2	4	7	6	6	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	5	11	15	21	29	58	75	69	50	20	-	-
Male	1	4	8	9	12	27	26	28	14	7	-	-
Female	4	7	7	12	17	31	49	41	36	13	-	-
White, both sexes	4	9	12	17	27	51	69	63	47	20	-	-
Male	1	3	7	7	10	23	24	25	11	7	-	-
Female	3	6	5	10	17	28	45	38	36	13	-	-
Black, both sexes	1	2	1	4	-	2	3	3	1	-	-	-
Male	-	1	-	2	-	1	1	1	1	-	-	-
Female	1	1	1	2	-	1	2	2	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	2	4	3	3	2	-	-	-
Male	-	-	1	-	2	2	1	2	2	-	-	-
Female	-	-	-	-	-	2	2	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
Colorado												
All races, both sexes	-	-	1	2	2	4	11	5	3	1	1	-
Male	-	-	-	2	2	2	4	3	-	1	-	-
Female	-	-	1	-	-	2	7	2	3	-	1	-
White, both sexes	-	-	1	2	2	4	11	5	3	1	1	-
Male	-	-	-	2	2	2	4	3	-	1	-	-
Female	-	-	1	-	-	2	7	2	3	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	1	3	2	9	14	14	11	-	1	-
Male	-	-	1	1	-	4	3	4	3	-	-	-
Female	-	-	-	2	2	5	11	10	8	-	1	-
White, both sexes	-	-	1	3	2	9	14	14	11	-	1	-
Male	-	-	1	1	-	4	3	4	3	-	-	-
Female	-	-	-	2	2	5	11	10	8	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
Delaware												
All races, both sexes	-	-	-	1	1	3	5	-	3	2	-	-
Male	-	-	-	1	1	1	3	-	1	1	-	-
Female	-	-	-	-	-	2	2	-	2	1	-	-
White, both sexes	-	-	-	1	1	3	5	-	3	2	-	-
Male	-	-	-	1	1	1	3	-	1	1	-	-
Female	-	-	-	-	-	2	2	-	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	1	1	-	-	1	1	1	-	-	-	-
Male	-	1	-	-	-	-	1	1	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	1	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
Florida												
All races, both sexes	3	7	14	20	35	77	100	117	51	13	1	-
Male	2	2	7	12	21	36	42	45	15	2	-	-
Female	1	5	7	8	14	41	58	72	36	11	1	-
White, both sexes	3	7	11	17	33	74	97	110	49	13	1	-
Male	2	2	7	10	21	34	41	44	15	2	-	-
Female	1	5	4	7	12	40	56	66	34	11	1	-
Black, both sexes	-	-	3	3	2	3	3	6	2	-	-	-
Male	-	-	-	2	-	2	1	1	-	-	-	-
Female	-	-	3	1	2	1	2	5	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Georgia												
All races, both sexes	1	2	3	5	11	11	15	13	4	2	-	-
Male	1	1	-	-	7	4	5	-	-	-	-	-
Female	-	1	3	5	4	7	10	13	4	2	-	-
White, both sexes	1	2	3	3	7	10	11	12	4	2	-	-
Male	1	1	-	-	3	4	4	-	-	-	-	-
Female	-	1	3	3	4	6	7	12	4	2	-	-
Black, both sexes	-	-	-	2	4	1	4	1	-	-	-	-
Male	-	-	-	-	4	-	1	-	-	-	-	-
Female	-	-	-	2	-	1	3	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
Hawaii												
All races, both sexes	-	-	1	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	1	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Idaho												
All races, both sexes	-	1	1	1	-	2	2	2	-	1	-	-
Male	-	-	1	1	-	1	1	1	-	-	-	-
Female	-	1	-	-	-	1	1	1	-	1	-	-
White, both sexes	-	1	1	1	-	2	2	2	-	1	-	-
Male	-	-	1	1	-	1	1	1	-	-	-	-
Female	-	1	-	-	-	1	1	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
Illinois												
All races, both sexes	2	7	7	9	22	29	34	48	41	13	1	-
Male	1	3	4	4	8	11	16	13	12	2	-	-
Female	1	4	3	5	14	18	18	35	29	11	1	-
White, both sexes	2	6	7	7	22	27	31	46	40	13	1	-
Male	1	2	4	3	8	11	15	13	12	2	-	-
Female	1	4	3	4	14	16	16	33	28	11	1	-
Black, both sexes	-	1	-	2	-	2	3	1	-	-	-	-
Male	-	1	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	2	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Indiana												
All races, both sexes	2	1	2	4	7	20	13	14	12	2	1	-
Male	1	-	2	3	2	10	7	3	4	1	-	-
Female	1	1	-	1	5	10	6	11	8	1	1	-
White, both sexes	2	1	2	4	7	20	12	14	12	2	1	-
Male	1	-	2	3	2	10	6	3	4	1	-	-
Female	1	1	-	1	5	10	6	11	8	1	1	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
Iowa												
All races, both sexes	-	-	1	2	1	5	12	7	5	1	-	-
Male	-	-	1	1	1	1	5	2	1	1	-	-
Female	-	-	-	1	-	4	7	5	4	-	-	-
White, both sexes	-	-	1	2	1	5	12	7	5	1	-	-
Male	-	-	1	1	1	1	5	2	1	1	-	-
Female	-	-	-	1	-	4	7	5	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	-	-	-	2	4	6	6	3	1	1	-
Male	1	-	-	-	1	-	1	2	1	-	-	-
Female	-	-	-	-	1	4	5	4	2	1	1	-
White, both sexes	1	-	-	-	2	4	6	6	3	1	1	-
Male	1	-	-	-	1	-	1	2	1	-	-	-
Female	-	-	-	-	1	4	5	4	2	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
Kentucky												
All races, both sexes	-	1	-	2	3	9	10	9	6	-	1	-
Male	-	-	-	-	2	4	4	4	-	-	-	-
Female	-	1	-	2	1	5	6	5	6	-	1	-
White, both sexes	-	1	-	2	3	9	10	9	5	-	1	-
Male	-	-	-	-	2	4	4	4	-	-	-	-
Female	-	1	-	2	1	5	6	5	5	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	2	3	3	3	4	14	5	5	3	1	-
Male	-	2	1	1	-	2	4	-	1	1	-	-
Female	1	-	2	2	3	2	10	5	4	2	1	-
White, both sexes	1	2	2	3	2	4	12	5	4	3	-	-
Male	-	2	-	1	-	2	4	-	1	1	-	-
Female	1	-	2	2	2	2	8	5	3	2	-	-
Black, both sexes	-	-	1	-	1	-	2	-	1	-	1	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
Maine												
All races, both sexes	-	-	1	1	7	13	28	17	11	6	1	-
Male	-	-	-	1	3	6	13	6	4	-	-	-
Female	-	-	1	-	4	7	15	11	7	6	1	-
White, both sexes	-	-	1	1	6	13	27	17	11	6	1	-
Male	-	-	-	1	3	6	12	6	4	-	-	-
Female	-	-	1	-	3	7	15	11	7	6	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	2	5	6	9	9	32	14	10	4	-	-
Male	1	1	3	4	4	3	9	6	4	-	-	-
Female	-	1	2	2	5	6	23	8	6	4	-	-
White, both sexes	1	2	5	5	8	9	31	11	10	4	-	-
Male	1	1	3	3	4	3	9	5	4	-	-	-
Female	-	1	2	2	4	6	22	6	6	4	-	-
Black, both sexes	-	-	-	-	1	-	1	3	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
Massachusetts												
All races, both sexes	1	-	2	3	4	24	23	21	17	6	-	-
Male	-	-	2	2	1	5	7	5	4	2	-	-
Female	1	-	-	1	3	19	16	16	13	4	-	-
White, both sexes	1	-	2	3	4	22	23	21	17	6	-	-
Male	-	-	2	2	1	5	7	5	4	2	-	-
Female	1	-	-	1	3	17	16	16	13	4	-	-
Black, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	3	4	4	6	11	18	25	25	9	1	-	-
Male	3	2	3	4	8	5	11	8	3	-	-	-
Female	-	2	1	2	3	13	14	17	6	1	-	-
White, both sexes	2	4	4	5	11	17	25	23	8	1	-	-
Male	2	2	3	4	8	5	11	8	2	-	-	-
Female	-	2	1	1	3	12	14	15	6	1	-	-
Black, both sexes	1	-	-	1	-	1	-	2	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	1	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
Minnesota												
All races, both sexes	-	1	1	3	4	5	7	7	8	3	-	-
Male	-	-	-	2	-	3	2	4	4	1	-	-
Female	-	1	1	1	4	2	5	3	4	2	-	-
White, both sexes	-	1	1	3	4	5	6	7	8	3	-	-
Male	-	-	-	2	-	3	1	4	4	1	-	-
Female	-	1	1	1	4	2	5	3	4	2	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	1	1	2	-	6	4	1	2	-	-
Male	-	-	1	-	1	-	3	2	-	-	-	-
Female	-	-	-	1	1	-	3	2	1	2	-	-
White, both sexes	-	-	1	1	2	-	6	3	1	2	-	-
Male	-	-	1	-	1	-	3	1	-	-	-	-
Female	-	-	-	1	1	-	3	2	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
Missouri												
All races, both sexes	3	3	8	6	9	37	28	36	23	11	1	-
Male	2	1	4	4	4	13	11	11	4	2	-	-
Female	1	2	4	2	5	24	17	25	19	9	1	-
White, both sexes	1	3	7	4	7	35	27	34	23	10	1	-
Male	-	1	4	2	4	12	10	10	4	2	-	-
Female	1	2	3	2	3	23	17	24	19	8	1	-
Black, both sexes	2	-	1	2	2	2	1	2	-	1	-	-
Male	2	-	-	2	-	1	1	1	-	-	-	-
Female	-	-	1	-	2	1	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	1	-	-	1	3	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	1	-	-	-	2	-	-	-	-
White, both sexes	-	-	-	1	-	-	1	3	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	1	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
Nebraska												
All races, both sexes	-	1	1	1	1	1	5	2	4	-	-	-
Male	-	-	1	1	-	-	3	-	-	-	-	-
Female	-	1	-	-	1	1	2	2	4	-	-	-
White, both sexes	-	1	1	1	1	1	5	2	4	-	-	-
Male	-	-	1	1	-	-	3	-	-	-	-	-
Female	-	1	-	-	1	1	2	2	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	2	1	-	2	6	3	-	1	-	-	-
Male	1	1	-	-	-	3	-	-	1	-	-	-
Female	-	1	1	-	2	3	3	-	-	-	-	-
White, both sexes	1	2	1	-	2	6	3	-	1	-	-	-
Male	1	1	-	-	-	3	-	-	1	-	-	-
Female	-	1	1	-	2	3	3	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
New Hampshire												
All races, both sexes	-	1	1	1	7	5	6	7	4	1	1	-
Male	-	-	1	1	2	1	3	5	-	-	-	-
Female	-	1	-	-	5	4	3	2	4	1	1	-
White, both sexes	-	1	1	1	7	5	6	7	4	1	1	-
Male	-	-	1	1	2	1	3	5	-	-	-	-
Female	-	1	-	-	5	4	3	2	4	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	5	3	6	8	17	35	35	39	45	10	2	-
Male	3	1	4	5	4	11	16	9	14	-	-	-
Female	2	2	2	3	13	24	19	30	31	10	2	-
White, both sexes	4	2	3	7	16	31	32	38	44	10	2	-
Male	3	-	2	4	4	10	15	8	13	-	-	-
Female	1	2	1	3	12	21	17	30	31	10	2	-
Black, both sexes	1	1	2	1	1	3	3	1	1	-	-	-
Male	-	1	1	1	-	-	1	1	1	-	-	-
Female	1	-	1	-	1	3	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
New Mexico												
All races, both sexes	1	-	1	1	1	3	1	2	1	1	-	-
Male	1	-	-	1	1	1	1	2	-	1	-	-
Female	-	-	1	-	-	2	-	-	1	-	-	-
White, both sexes	1	-	1	-	1	3	1	2	1	1	-	-
Male	1	-	-	-	1	1	1	2	-	1	-	-
Female	-	-	1	-	-	2	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	3	1	10	10	17	20	53	49	35	10	2	-
Male	1	1	4	7	9	10	26	16	9	3	-	-
Female	2	-	6	3	8	10	27	33	26	7	2	-
White, both sexes	3	1	10	9	16	19	51	48	34	10	2	-
Male	1	1	4	7	8	9	24	16	9	3	-	-
Female	2	-	6	2	8	10	27	32	25	7	2	-
Black, both sexes	-	-	-	1	1	-	2	1	1	-	-	-
Male	-	-	-	-	1	-	2	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
North Carolina												
All races, both sexes	1	3	2	6	8	19	20	15	10	2	1	-
Male	-	2	1	2	3	4	5	6	2	-	-	-
Female	1	1	1	4	5	15	15	9	8	2	1	-
White, both sexes	1	2	1	5	7	15	16	14	9	2	1	-
Male	-	1	1	1	2	2	5	6	2	-	-	-
Female	1	1	-	4	5	13	11	8	7	2	1	-
Black, both sexes	-	1	1	1	1	4	4	1	1	-	-	-
Male	-	1	-	1	1	2	-	-	-	-	-	-
Female	-	-	1	-	-	2	4	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	1	1	1	-	1	1	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	1	1	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	1	1	-	1	1	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
Ohio												
All races, both sexes	5	8	17	22	37	59	90	75	42	9	1	-
Male	2	4	8	7	12	30	34	27	12	1	-	-
Female	3	4	9	15	25	29	56	48	30	8	1	-
White, both sexes	5	7	14	22	34	57	84	71	39	9	1	-
Male	2	3	5	7	11	28	33	27	12	1	-	-
Female	3	4	9	15	23	29	51	44	27	8	1	-
Black, both sexes	-	1	3	-	3	1	6	4	3	-	-	-
Male	-	1	3	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	2	-	5	4	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	1	2	2	9	5	13	6	9	1	-	-
Male	1	-	1	1	3	3	5	2	2	-	-	-
Female	1	1	1	1	6	2	8	4	7	1	-	-
White, both sexes	2	1	2	2	9	5	13	6	9	1	-	-
Male	1	-	1	1	3	3	5	2	2	-	-	-
Female	1	1	1	1	6	2	8	4	7	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
Oregon												
All races, both sexes	1	-	1	2	1	1	18	11	3	4	-	-
Male	-	-	1	-	1	1	9	5	-	-	-	-
Female	1	-	-	2	-	-	9	6	3	4	-	-
White, both sexes	1	-	1	2	1	1	18	11	3	4	-	-
Male	-	-	1	-	1	1	9	5	-	-	-	-
Female	1	-	-	2	-	-	9	6	3	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	2	12	8	15	33	54	85	75	51	13	3	-
Male	-	6	3	4	17	22	36	15	11	3	1	-
Female	2	6	5	11	16	32	49	60	40	10	2	-
White, both sexes	2	10	8	13	32	51	78	72	51	13	3	-
Male	-	5	3	4	17	21	33	14	11	3	1	-
Female	2	5	5	9	15	30	45	58	40	10	2	-
Black, both sexes	-	2	-	2	1	2	5	3	-	-	-	-
Male	-	1	-	-	-	1	2	1	-	-	-	-
Female	-	1	-	2	1	1	3	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
Rhode Island												
All races, both sexes	1	1	-	-	1	5	3	7	3	2	-	-
Male	1	1	-	-	1	2	-	4	1	1	-	-
Female	-	-	-	-	-	3	3	3	2	1	-	-
White, both sexes	1	1	-	-	1	5	3	7	3	2	-	-
Male	1	1	-	-	1	2	-	4	1	1	-	-
Female	-	-	-	-	-	3	3	3	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	2	-	4	-	8	5	5	7	2	-	-
Male	1	1	-	3	-	2	3	3	1	-	-	-
Female	-	1	-	1	-	6	2	2	6	2	-	-
White, both sexes	1	2	-	2	-	8	4	4	6	1	-	-
Male	1	1	-	1	-	2	2	2	1	-	-	-
Female	-	1	-	1	-	6	2	2	5	1	-	-
Black, both sexes	-	-	-	2	-	-	1	1	1	1	-	-
Male	-	-	-	2	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
South Dakota												
All races, both sexes	-	-	-	1	1	-	2	1	2	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	-	1	-	-	1	-	2	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	2	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	4	4	9	14	18	23	13	19	2	1	-
Male	1	3	1	3	6	12	5	3	6	-	-	-
Female	1	1	3	6	8	6	18	10	13	2	1	-
White, both sexes	2	3	4	8	14	18	23	11	19	2	1	-
Male	1	2	1	2	6	12	5	3	6	-	-	-
Female	1	1	3	6	8	6	18	8	13	2	1	-
Black, both sexes	-	1	-	1	-	-	-	2	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
Texas												
All races, both sexes	2	-	3	10	21	28	28	21	14	7	1	-
Male	2	-	3	6	6	13	10	6	-	2	-	-
Female	-	-	-	4	15	15	18	15	14	5	1	-
White, both sexes	2	-	3	10	16	25	25	20	13	6	1	-
Male	2	-	3	6	4	12	10	6	-	1	-	-
Female	-	-	-	4	12	13	15	14	13	5	1	-
Black, both sexes	-	-	-	-	4	2	3	1	1	1	-	-
Male	-	-	-	-	2	1	-	-	-	1	-	-
Female	-	-	-	-	2	1	3	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	1	1	2	2	2	2	1	1	-	-
Male	-	-	1	-	1	2	-	1	-	-	-	-
Female	-	-	-	1	1	-	2	1	1	1	-	-
White, both sexes	-	-	1	1	2	2	2	2	1	1	-	-
Male	-	-	1	-	1	2	-	1	-	-	-	-
Female	-	-	-	1	1	-	2	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
Vermont												
All races, both sexes	-	-	2	-	1	2	5	7	-	5	-	-
Male	-	-	-	-	-	1	2	-	-	1	-	-
Female	-	-	2	-	1	1	3	7	-	4	-	-
White, both sexes	-	-	2	-	1	2	5	7	-	5	-	-
Male	-	-	-	-	-	1	2	-	-	1	-	-
Female	-	-	2	-	1	1	3	7	-	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	1	4	3	7	7	17	18	10	3	-	-
Male	1	-	2	1	4	6	6	9	2	1	-	-
Female	-	1	2	2	3	1	11	9	8	2	-	-
White, both sexes	-	1	4	3	5	5	16	18	10	3	-	-
Male	-	-	2	1	3	5	5	9	2	1	-	-
Female	-	1	2	2	2	-	11	9	8	2	-	-
Black, both sexes	1	-	-	-	2	2	1	-	-	-	-	-
Male	1	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
Washington												
All races, both sexes	1	-	2	2	6	10	17	9	12	2	1	-
Male	-	-	-	1	2	4	5	3	7	-	-	-
Female	1	-	2	1	4	6	12	6	5	2	1	-
White, both sexes	1	-	2	2	6	10	17	9	12	2	1	-
Male	-	-	-	1	2	4	5	3	7	-	-	-
Female	1	-	2	1	4	6	12	6	5	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	2	5	1	3	5	6	4	1	-	-	-
Male	-	1	3	-	-	3	4	1	-	-	-	-
Female	-	1	2	1	3	2	2	3	1	-	-	-
White, both sexes	-	2	5	1	3	5	5	4	1	-	-	-
Male	-	1	3	-	-	3	4	1	-	-	-	-
Female	-	1	2	1	3	2	1	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other intestinal infectious diseases (A01-A08)												
Wisconsin												
All races, both sexes	-	-	2	3	4	7	13	7	8	2	1	-
Male	-	-	2	2	1	3	6	5	2	-	-	-
Female	-	-	-	1	3	4	7	2	6	2	1	-
White, both sexes	-	-	2	3	4	5	13	7	7	2	1	-
Male	-	-	2	2	1	2	6	5	1	-	-	-
Female	-	-	-	1	3	3	7	2	6	2	1	-
Black, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
United States												
All races, both sexes	30	1	-	-	-	-	-	-	1	1	-	2
Male	22	1	-	-	-	-	-	-	-	1	-	2
Female	8	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	20	1	-	-	-	-	-	-	1	-	-	1
Male	14	1	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Arkansas												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Florida												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Hawaii												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Illinois												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Minnesota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Oregon												
All races, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Rhode Island												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Washington												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Wisconsin												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
United States												
All races, both sexes	1	2	2	4	3	4	3	6	-	-	-	-
Male	1	2	2	2	1	4	2	4	-	-	-	-
Female	-	-	-	2	2	-	1	2	-	-	-	-
White, both sexes	1	1	1	1	2	3	3	5	-	-	-	-
Male	1	1	1	1	-	3	2	3	-	-	-	-
Female	-	-	-	-	2	-	1	2	-	-	-	-
Black, both sexes	-	1	-	2	-	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	1	-	1	-	-	-	-
Male	-	-	1	-	1	1	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Arkansas												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	1	2	-	1	-	-	-	-
Male	-	-	-	-	1	2	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Florida												
All races, both sexes	-	-	-	1	-	-	1	1	-	-	-	-
Male	-	-	-	1	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Hawaii												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Oregon												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Rhode Island												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Wisconsin												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
United States												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other bacterial food-borne intoxications (A05)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
United States												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other bacterial food-borne intoxications (A05)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
United States												
All races, both sexes	3	-	-	-	1	-	1	-	-	-	-	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	3	-	-	-	1	-	1	-	-	-	-	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
Colorado												
All races, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
Illinois												
All races, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Amebiasis (A06)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
United States												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Amebiasis (A06)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
United States												
All races, both sexes	4,246	13	14	-	-	1	1	9	2	10	21	37
Male	1,587	8	4	-	-	-	1	4	1	7	12	18
Female	2,659	5	10	-	-	1	-	5	1	3	9	19
White, both sexes	3,977	10	12	-	-	1	1	7	2	9	13	26
Male	1,474	6	3	-	-	-	1	3	1	6	7	12
Female	2,503	4	9	-	-	1	-	4	1	3	6	14
Black, both sexes	226	1	1	-	-	-	-	2	-	1	6	9
Male	90	1	-	-	-	-	-	1	-	1	3	5
Female	136	-	1	-	-	-	-	1	-	-	3	4
American Indian 1/, both sexes	15	1	-	-	-	-	-	-	-	-	1	1
Male	9	1	-	-	-	-	-	-	-	-	1	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	28	1	1	-	-	-	-	-	-	-	1	1
Male	14	-	1	-	-	-	-	-	-	-	1	-
Female	14	1	-	-	-	-	-	-	-	-	-	1
Alabama												
All races, both sexes	71	2	-	-	-	-	-	-	-	-	-	-
Male	24	1	-	-	-	-	-	-	-	-	-	-
Female	47	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	63	2	-	-	-	-	-	-	-	-	-	-
Male	22	1	-	-	-	-	-	-	-	-	-	-
Female	41	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
Alaska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	74	-	-	-	-	-	-	-	-	-	-	-
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	47	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	73	-	-	-	-	-	-	-	-	-	-	-
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	46	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
Arkansas												
All races, both sexes	55	-	1	-	-	-	-	-	-	-	1	1
Male	24	-	-	-	-	-	-	-	-	-	1	-
Female	31	-	1	-	-	-	-	-	-	-	-	1
White, both sexes	55	-	1	-	-	-	-	-	-	-	1	1
Male	24	-	-	-	-	-	-	-	-	-	1	-
Female	31	-	1	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	361	3	1	-	-	1	-	1	-	-	1	5
Male	139	1	1	-	-	-	-	1	-	-	1	3
Female	222	2	-	-	-	1	-	-	-	-	-	2
White, both sexes	328	2	-	-	-	1	-	1	-	-	1	5
Male	123	1	-	-	-	-	-	1	-	-	1	3
Female	205	1	-	-	-	1	-	-	-	-	-	2
Black, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	1	1	-	-	-	-	-	-	-	-	-
Male	8	-	1	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
Colorado												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	55	-	-	-	-	-	-	-	-	-	1	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	55	-	-	-	-	-	-	-	-	-	1	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
Delaware												
All races, both sexes	16	1	-	-	-	-	-	-	-	-	-	-
Male	9	1	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	1	-	-	-	-	-	-	-	-	-	-
Male	9	1	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
Florida												
All races, both sexes	443	1	-	-	-	-	-	1	-	-	4	2
Male	186	1	-	-	-	-	-	1	-	-	2	1
Female	257	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	420	1	-	-	-	-	-	-	-	-	4	2
Male	180	1	-	-	-	-	-	-	-	-	2	1
Female	240	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	22	-	-	-	-	-	-	1	-	-	-	-
Male	6	-	-	-	-	-	-	1	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	73	1	2	-	-	-	-	-	-	1	-	3
Male	21	1	-	-	-	-	-	-	-	1	-	1
Female	52	-	2	-	-	-	-	-	-	-	-	2
White, both sexes	61	1	1	-	-	-	-	-	-	1	-	3
Male	16	1	-	-	-	-	-	-	-	1	-	1
Female	45	-	1	-	-	-	-	-	-	-	-	2
Black, both sexes	12	-	1	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
Hawaii												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
Illinois												
All races, both sexes	219	1	-	-	-	-	-	-	1	1	1	2
Male	76	-	-	-	-	-	-	-	-	1	1	-
Female	143	1	-	-	-	-	-	-	1	-	-	2
White, both sexes	205	1	-	-	-	-	-	-	1	-	-	1
Male	71	-	-	-	-	-	-	-	-	-	-	-
Female	134	1	-	-	-	-	-	-	1	-	-	1
Black, both sexes	12	-	-	-	-	-	-	-	-	1	1	1
Male	5	-	-	-	-	-	-	-	-	1	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	81	-	3	-	-	-	-	-	-	-	-	-
Male	35	-	2	-	-	-	-	-	-	-	-	-
Female	46	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	80	-	3	-	-	-	-	-	-	-	-	-
Male	34	-	2	-	-	-	-	-	-	-	-	-
Female	46	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
Iowa												
All races, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
Kentucky												
All races, both sexes	44	-	-	-	-	-	-	1	-	-	-	2
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	29	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	42	-	-	-	-	-	-	1	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	46	2	-	-	-	-	-	-	-	-	-	-
Male	14	2	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
Maine												
All races, both sexes	85	-	-	-	-	-	-	-	-	-	-	-
Male	33	-	-	-	-	-	-	-	-	-	-	-
Female	52	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	83	-	-	-	-	-	-	-	-	-	-	-
Male	32	-	-	-	-	-	-	-	-	-	-	-
Female	51	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	94	-	-	-	-	-	-	-	-	-	1	1
Male	36	-	-	-	-	-	-	-	-	-	-	1
Female	58	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	87	-	-	-	-	-	-	-	-	-	-	1
Male	34	-	-	-	-	-	-	-	-	-	-	1
Female	53	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
Massachusetts												
All races, both sexes	104	-	-	-	-	-	-	1	-	-	1	1
Male	31	-	-	-	-	-	-	1	-	-	1	1
Female	73	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	102	-	-	-	-	-	-	1	-	-	1	1
Male	31	-	-	-	-	-	-	1	-	-	1	1
Female	71	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	109	-	-	-	-	-	-	-	-	-	1	2
Male	49	-	-	-	-	-	-	-	-	-	1	1
Female	60	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	102	-	-	-	-	-	-	-	-	-	1	1
Male	46	-	-	-	-	-	-	-	-	-	1	-
Female	56	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
Minnesota												
All races, both sexes	40	-	1	-	-	-	-	1	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	1	-	-	-	-	1	-	-	-	-
White, both sexes	39	-	1	-	-	-	-	1	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	1	-	-	-	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
Missouri												
All races, both sexes	169	-	1	-	-	-	-	-	-	-	1	2
Male	57	-	-	-	-	-	-	-	-	-	-	1
Female	112	-	1	-	-	-	-	-	-	-	1	1
White, both sexes	154	-	1	-	-	-	-	-	-	-	-	1
Male	49	-	-	-	-	-	-	-	-	-	-	-
Female	105	-	1	-	-	-	-	-	-	-	-	1
Black, both sexes	15	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
Nebraska												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
New Hampshire												
All races, both sexes	36	-	-	-	-	-	-	1	-	1	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	36	-	-	-	-	-	-	1	-	1	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	204	-	-	-	-	-	-	-	-	-	-	1
Male	65	-	-	-	-	-	-	-	-	-	-	-
Female	139	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	189	-	-	-	-	-	-	-	-	-	-	1
Male	58	-	-	-	-	-	-	-	-	-	-	-
Female	131	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
New Mexico												
All races, both sexes	13	-	1	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	1	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	216	-	1	-	-	-	-	-	1	2	1	1
Male	88	-	-	-	-	-	-	-	1	1	-	-
Female	128	-	1	-	-	-	-	-	-	1	1	1
White, both sexes	208	-	1	-	-	-	-	-	1	2	1	-
Male	84	-	-	-	-	-	-	-	1	1	-	-
Female	124	-	1	-	-	-	-	-	-	1	1	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
North Carolina												
All races, both sexes	87	-	-	-	-	-	-	-	-	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	62	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	73	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	53	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
Ohio												
All races, both sexes	372	-	-	-	-	-	-	-	-	2	-	5
Male	142	-	-	-	-	-	-	-	-	2	-	3
Female	230	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	349	-	-	-	-	-	-	-	-	2	-	4
Male	133	-	-	-	-	-	-	-	-	2	-	2
Female	216	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	51	1	-	-	-	-	-	-	-	-	1	-
Male	20	1	-	-	-	-	-	-	-	-	1	-
Female	31	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	50	1	-	-	-	-	-	-	-	-	-	-
Male	19	1	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
Oregon												
All races, both sexes	43	-	-	-	-	-	-	1	-	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	42	-	-	-	-	-	-	1	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Pennsylvania												
All races, both sexes	356	-	-	-	-	-	1	2	-	2	1	1
Male	120	-	-	-	-	-	1	1	-	1	1	-
Female	236	-	-	-	-	-	-	1	-	1	-	1
White, both sexes	335	-	-	-	-	-	1	1	-	2	-	-
Male	113	-	-	-	-	-	1	1	-	1	-	-
Female	222	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	18	-	-	-	-	-	-	1	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
Rhode Island												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	33	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
South Dakota												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	111	1	-	-	-	-	-	-	-	1	-	-
Male	41	-	-	-	-	-	-	-	-	1	-	-
Female	70	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	107	1	-	-	-	-	-	-	-	1	-	-
Male	39	-	-	-	-	-	-	-	-	1	-	-
Female	68	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
Texas												
All races, both sexes	141	-	2	-	-	-	-	-	-	-	2	2
Male	52	-	1	-	-	-	-	-	-	-	1	2
Female	89	-	1	-	-	-	-	-	-	-	1	-
White, both sexes	124	-	2	-	-	-	-	-	-	-	-	1
Male	46	-	1	-	-	-	-	-	-	-	-	1
Female	78	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	15	-	-	-	-	-	-	-	-	-	2	1
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	9	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
Vermont												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	74	-	-	-	-	-	-	-	-	-	1	2
Male	34	-	-	-	-	-	-	-	-	-	1	1
Female	40	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	67	-	-	-	-	-	-	-	-	-	1	1
Male	30	-	-	-	-	-	-	-	-	-	1	1
Female	37	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
Washington												
All races, both sexes	62	-	-	-	-	-	-	-	-	-	1	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	62	-	-	-	-	-	-	-	-	-	1	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	28	-	1	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	1	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intestinal infections due to other specified organisms (A04,A07-A08)												
Wisconsin												
All races, both sexes	47	-	-	-	-	-	-	-	-	-	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	44	-	-	-	-	-	-	-	-	-	-	1
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
United States												
All races, both sexes	53	91	155	210	382	671	948	852	576	175	25	-
Male	28	43	78	98	161	281	369	287	151	35	1	-
Female	25	48	77	112	221	390	579	565	425	140	24	-
White, both sexes	47	80	139	186	353	629	895	814	557	172	24	-
Male	24	34	72	85	149	265	350	277	144	34	1	-
Female	23	46	67	101	204	364	545	537	413	138	23	-
Black, both sexes	6	11	13	20	25	33	45	34	15	3	1	-
Male	4	9	5	11	10	11	15	9	4	1	-	-
Female	2	2	8	9	15	22	30	25	11	2	1	-
American Indian 1/, both sexes	-	-	2	3	2	3	2	-	-	-	-	-
Male	-	-	-	1	1	2	2	-	-	-	-	-
Female	-	-	2	2	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	2	6	6	4	4	-	-	-
Male	-	-	1	1	1	3	2	1	3	-	-	-
Female	-	-	-	-	1	3	4	3	1	-	-	-
Alabama												
All races, both sexes	-	2	5	5	12	11	11	17	5	-	1	-
Male	-	2	3	1	5	2	3	5	2	-	-	-
Female	-	-	2	4	7	9	8	12	3	-	1	-
White, both sexes	-	1	4	5	10	8	11	17	4	-	1	-
Male	-	1	3	1	5	2	3	5	1	-	-	-
Female	-	-	1	4	5	6	8	12	3	-	1	-
Black, both sexes	-	1	1	-	2	3	-	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	2	3	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	5	5	3	10	10	14	16	8	3	-	-
Male	-	4	2	-	2	3	4	9	2	1	-	-
Female	-	1	3	3	8	7	10	7	6	2	-	-
White, both sexes	-	5	5	2	10	10	14	16	8	3	-	-
Male	-	4	2	-	2	3	4	9	2	1	-	-
Female	-	1	3	2	8	7	10	7	6	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
Arkansas												
All races, both sexes	2	-	-	2	4	12	12	10	7	3	-	-
Male	1	-	-	2	2	8	5	4	1	-	-	-
Female	1	-	-	-	2	4	7	6	6	3	-	-
White, both sexes	2	-	-	2	4	12	12	10	7	3	-	-
Male	1	-	-	2	2	8	5	4	1	-	-	-
Female	1	-	-	-	2	4	7	6	6	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	5	11	15	21	28	56	75	68	50	20	-	-
Male	1	4	8	9	11	25	26	27	14	7	-	-
Female	4	7	7	12	17	31	49	41	36	13	-	-
White, both sexes	4	9	12	17	27	50	69	63	47	20	-	-
Male	1	3	7	7	10	22	24	25	11	7	-	-
Female	3	6	5	10	17	28	45	38	36	13	-	-
Black, both sexes	1	2	1	4	-	2	3	3	1	-	-	-
Male	-	1	-	2	-	1	1	1	1	-	-	-
Female	1	1	1	2	-	1	2	2	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	3	3	2	2	-	-	-
Male	-	-	1	-	1	1	1	1	2	-	-	-
Female	-	-	-	-	-	2	2	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
Colorado												
All races, both sexes	-	-	1	2	2	4	11	5	3	1	1	-
Male	-	-	-	2	2	2	4	3	-	1	-	-
Female	-	-	1	-	-	2	7	2	3	-	1	-
White, both sexes	-	-	1	2	2	4	11	5	3	1	1	-
Male	-	-	-	2	2	2	4	3	-	1	-	-
Female	-	-	1	-	-	2	7	2	3	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	1	3	2	8	14	14	11	-	1	-
Male	-	-	1	1	-	3	3	4	3	-	-	-
Female	-	-	-	2	2	5	11	10	8	-	1	-
White, both sexes	-	-	1	3	2	8	14	14	11	-	1	-
Male	-	-	1	1	-	3	3	4	3	-	-	-
Female	-	-	-	2	2	5	11	10	8	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
Delaware												
All races, both sexes	-	-	-	1	1	3	5	-	3	2	-	-
Male	-	-	-	1	1	1	3	-	1	1	-	-
Female	-	-	-	-	-	2	2	-	2	1	-	-
White, both sexes	-	-	-	1	1	3	5	-	3	2	-	-
Male	-	-	-	1	1	1	3	-	1	1	-	-
Female	-	-	-	-	-	2	2	-	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	1	1	-	-	1	1	1	-	-	-	-
Male	-	1	-	-	-	-	1	1	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	1	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
Florida												
All races, both sexes	3	7	14	19	35	77	99	116	51	13	1	-
Male	2	2	7	11	21	36	41	44	15	2	-	-
Female	1	5	7	8	14	41	58	72	36	11	1	-
White, both sexes	3	7	11	17	33	74	96	109	49	13	1	-
Male	2	2	7	10	21	34	40	43	15	2	-	-
Female	1	5	4	7	12	40	56	66	34	11	1	-
Black, both sexes	-	-	3	2	2	3	3	6	2	-	-	-
Male	-	-	-	1	-	2	1	1	-	-	-	-
Female	-	-	3	1	2	1	2	5	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Georgia												
All races, both sexes	1	2	3	4	11	11	15	13	4	2	-	-
Male	1	1	-	-	7	4	5	-	-	-	-	-
Female	-	1	3	4	4	7	10	13	4	2	-	-
White, both sexes	1	2	3	3	7	10	11	12	4	2	-	-
Male	1	1	-	-	3	4	4	-	-	-	-	-
Female	-	1	3	3	4	6	7	12	4	2	-	-
Black, both sexes	-	-	-	1	4	1	4	1	-	-	-	-
Male	-	-	-	-	4	-	1	-	-	-	-	-
Female	-	-	-	1	-	1	3	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
Hawaii												
All races, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Idaho												
All races, both sexes	-	1	1	1	-	2	2	2	-	1	-	-
Male	-	-	1	1	-	1	1	1	-	-	-	-
Female	-	1	-	-	-	1	1	1	-	1	-	-
White, both sexes	-	1	1	1	-	2	2	2	-	1	-	-
Male	-	-	1	1	-	1	1	1	-	-	-	-
Female	-	1	-	-	-	1	1	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
Illinois												
All races, both sexes	2	7	7	9	22	29	34	48	41	13	1	-
Male	1	3	4	4	8	11	16	13	12	2	-	-
Female	1	4	3	5	14	18	18	35	29	11	1	-
White, both sexes	2	6	7	7	22	27	31	46	40	13	1	-
Male	1	2	4	3	8	11	15	13	12	2	-	-
Female	1	4	3	4	14	16	16	33	28	11	1	-
Black, both sexes	-	1	-	2	-	2	3	1	-	-	-	-
Male	-	1	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	2	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Indiana												
All races, both sexes	2	1	2	4	7	20	13	14	12	2	1	-
Male	1	-	2	3	2	10	7	3	4	1	-	-
Female	1	1	-	1	5	10	6	11	8	1	1	-
White, both sexes	2	1	2	4	7	20	12	14	12	2	1	-
Male	1	-	2	3	2	10	6	3	4	1	-	-
Female	1	1	-	1	5	10	6	11	8	1	1	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
Iowa												
All races, both sexes	-	-	1	2	1	5	12	7	5	1	-	-
Male	-	-	1	1	1	1	5	2	1	1	-	-
Female	-	-	-	1	-	4	7	5	4	-	-	-
White, both sexes	-	-	1	2	1	5	12	7	5	1	-	-
Male	-	-	1	1	1	1	5	2	1	1	-	-
Female	-	-	-	1	-	4	7	5	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	-	-	-	2	4	6	5	3	1	1	-
Male	1	-	-	-	1	-	1	1	1	-	-	-
Female	-	-	-	-	1	4	5	4	2	1	1	-
White, both sexes	1	-	-	-	2	4	6	5	3	1	1	-
Male	1	-	-	-	1	-	1	1	1	-	-	-
Female	-	-	-	-	1	4	5	4	2	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
Kentucky												
All races, both sexes	-	1	-	2	3	9	10	9	6	-	1	-
Male	-	-	-	-	2	4	4	4	-	-	-	-
Female	-	1	-	2	1	5	6	5	6	-	1	-
White, both sexes	-	1	-	2	3	9	10	9	5	-	1	-
Male	-	-	-	-	2	4	4	4	-	-	-	-
Female	-	1	-	2	1	5	6	5	5	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	2	3	3	3	4	14	5	5	3	1	-
Male	-	2	1	1	-	2	4	-	1	1	-	-
Female	1	-	2	2	3	2	10	5	4	2	1	-
White, both sexes	1	2	2	3	2	4	12	5	4	3	-	-
Male	-	2	-	1	-	2	4	-	1	1	-	-
Female	1	-	2	2	2	2	8	5	3	2	-	-
Black, both sexes	-	-	1	-	1	-	2	-	1	-	1	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
Maine												
All races, both sexes	-	-	1	1	7	13	28	17	11	6	1	-
Male	-	-	-	1	3	6	13	6	4	-	-	-
Female	-	-	1	-	4	7	15	11	7	6	1	-
White, both sexes	-	-	1	1	6	13	27	17	11	6	1	-
Male	-	-	-	1	3	6	12	6	4	-	-	-
Female	-	-	1	-	3	7	15	11	7	6	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	2	5	6	9	9	32	14	10	4	-	-
Male	1	1	3	4	4	3	9	6	4	-	-	-
Female	-	1	2	2	5	6	23	8	6	4	-	-
White, both sexes	1	2	5	5	8	9	31	11	10	4	-	-
Male	1	1	3	3	4	3	9	5	4	-	-	-
Female	-	1	2	2	4	6	22	6	6	4	-	-
Black, both sexes	-	-	-	-	1	-	1	3	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
Massachusetts												
All races, both sexes	1	-	2	3	4	24	23	21	17	6	-	-
Male	-	-	2	2	1	5	7	5	4	2	-	-
Female	1	-	-	1	3	19	16	16	13	4	-	-
White, both sexes	1	-	2	3	4	22	23	21	17	6	-	-
Male	-	-	2	2	1	5	7	5	4	2	-	-
Female	1	-	-	1	3	17	16	16	13	4	-	-
Black, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	3	4	4	6	11	18	25	25	9	1	-	-
Male	3	2	3	4	8	5	11	8	3	-	-	-
Female	-	2	1	2	3	13	14	17	6	1	-	-
White, both sexes	2	4	4	5	11	17	25	23	8	1	-	-
Male	2	2	3	4	8	5	11	8	2	-	-	-
Female	-	2	1	1	3	12	14	15	6	1	-	-
Black, both sexes	1	-	-	1	-	1	-	2	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	1	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
Minnesota												
All races, both sexes	-	1	1	3	4	5	7	6	8	3	-	-
Male	-	-	-	2	-	3	2	4	4	1	-	-
Female	-	1	1	1	4	2	5	2	4	2	-	-
White, both sexes	-	1	1	3	4	5	6	6	8	3	-	-
Male	-	-	-	2	-	3	1	4	4	1	-	-
Female	-	1	1	1	4	2	5	2	4	2	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	1	1	2	-	6	4	1	2	-	-
Male	-	-	1	-	1	-	3	2	-	-	-	-
Female	-	-	-	1	1	-	3	2	1	2	-	-
White, both sexes	-	-	1	1	2	-	6	3	1	2	-	-
Male	-	-	1	-	1	-	3	1	-	-	-	-
Female	-	-	-	1	1	-	3	2	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
Missouri												
All races, both sexes	3	3	8	6	9	37	28	36	23	11	1	-
Male	2	1	4	4	4	13	11	11	4	2	-	-
Female	1	2	4	2	5	24	17	25	19	9	1	-
White, both sexes	1	3	7	4	7	35	27	34	23	10	1	-
Male	-	1	4	2	4	12	10	10	4	2	-	-
Female	1	2	3	2	3	23	17	24	19	8	1	-
Black, both sexes	2	-	1	2	2	2	1	2	-	1	-	-
Male	2	-	-	2	-	1	1	1	-	-	-	-
Female	-	-	1	-	2	1	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	1	-	-	1	3	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	1	-	-	-	2	-	-	-	-
White, both sexes	-	-	-	1	-	-	1	3	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	1	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
Nebraska												
All races, both sexes	-	1	1	1	1	1	5	2	4	-	-	-
Male	-	-	1	1	-	-	3	-	-	-	-	-
Female	-	1	-	-	1	1	2	2	4	-	-	-
White, both sexes	-	1	1	1	1	1	5	2	4	-	-	-
Male	-	-	1	1	-	-	3	-	-	-	-	-
Female	-	1	-	-	1	1	2	2	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	2	1	-	2	6	3	-	1	-	-	-
Male	1	1	-	-	-	3	-	-	1	-	-	-
Female	-	1	1	-	2	3	3	-	-	-	-	-
White, both sexes	1	2	1	-	2	6	3	-	1	-	-	-
Male	1	1	-	-	-	3	-	-	1	-	-	-
Female	-	1	1	-	2	3	3	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
New Hampshire												
All races, both sexes	-	1	1	1	7	5	6	7	4	1	1	-
Male	-	-	1	1	2	1	3	5	-	-	-	-
Female	-	1	-	-	5	4	3	2	4	1	1	-
White, both sexes	-	1	1	1	7	5	6	7	4	1	1	-
Male	-	-	1	1	2	1	3	5	-	-	-	-
Female	-	1	-	-	5	4	3	2	4	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	5	3	5	7	17	35	35	39	45	10	2	-
Male	3	1	3	4	4	11	16	9	14	-	-	-
Female	2	2	2	3	13	24	19	30	31	10	2	-
White, both sexes	4	2	3	6	16	31	32	38	44	10	2	-
Male	3	-	2	3	4	10	15	8	13	-	-	-
Female	1	2	1	3	12	21	17	30	31	10	2	-
Black, both sexes	1	1	2	1	1	3	3	1	1	-	-	-
Male	-	1	1	1	-	-	1	1	1	-	-	-
Female	1	-	1	-	1	3	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
New Mexico												
All races, both sexes	1	-	1	1	1	3	1	2	1	1	-	-
Male	1	-	-	1	1	1	1	2	-	1	-	-
Female	-	-	1	-	-	2	-	-	1	-	-	-
White, both sexes	1	-	1	-	1	3	1	2	1	1	-	-
Male	1	-	-	-	1	1	1	2	-	1	-	-
Female	-	-	1	-	-	2	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	3	1	10	10	17	20	53	49	35	10	2	-
Male	1	1	4	7	9	10	26	16	9	3	-	-
Female	2	-	6	3	8	10	27	33	26	7	2	-
White, both sexes	3	1	10	9	16	19	51	48	34	10	2	-
Male	1	1	4	7	8	9	24	16	9	3	-	-
Female	2	-	6	2	8	10	27	32	25	7	2	-
Black, both sexes	-	-	-	1	1	-	2	1	1	-	-	-
Male	-	-	-	-	1	-	2	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
North Carolina												
All races, both sexes	1	3	2	6	8	19	20	15	10	2	1	-
Male	-	2	1	2	3	4	5	6	2	-	-	-
Female	1	1	1	4	5	15	15	9	8	2	1	-
White, both sexes	1	2	1	5	7	15	16	14	9	2	1	-
Male	-	1	1	1	2	2	5	6	2	-	-	-
Female	1	1	-	4	5	13	11	8	7	2	1	-
Black, both sexes	-	1	1	1	1	4	4	1	1	-	-	-
Male	-	1	-	1	1	2	-	-	-	-	-	-
Female	-	-	1	-	-	2	4	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	1	1	1	-	1	1	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	1	1	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	1	1	-	1	1	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
Ohio												
All races, both sexes	5	8	17	22	37	59	90	75	42	9	1	-
Male	2	4	8	7	12	30	34	27	12	1	-	-
Female	3	4	9	15	25	29	56	48	30	8	1	-
White, both sexes	5	7	14	22	34	57	84	71	39	9	1	-
Male	2	3	5	7	11	28	33	27	12	1	-	-
Female	3	4	9	15	23	29	51	44	27	8	1	-
Black, both sexes	-	1	3	-	3	1	6	4	3	-	-	-
Male	-	1	3	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	2	-	5	4	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	1	2	2	8	5	13	6	9	1	-	-
Male	1	-	1	1	3	3	5	2	2	-	-	-
Female	1	1	1	1	5	2	8	4	7	1	-	-
White, both sexes	2	1	2	2	8	5	13	6	9	1	-	-
Male	1	-	1	1	3	3	5	2	2	-	-	-
Female	1	1	1	1	5	2	8	4	7	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
Oregon												
All races, both sexes	1	-	1	2	1	1	17	11	3	4	-	-
Male	-	-	1	-	1	1	9	5	-	-	-	-
Female	1	-	-	2	-	-	8	6	3	4	-	-
White, both sexes	1	-	1	2	1	1	17	11	3	4	-	-
Male	-	-	1	-	1	1	9	5	-	-	-	-
Female	1	-	-	2	-	-	8	6	3	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	2	12	8	15	33	53	84	75	51	13	3	-
Male	-	6	3	4	17	21	35	15	11	3	1	-
Female	2	6	5	11	16	32	49	60	40	10	2	-
White, both sexes	2	10	8	13	32	50	77	72	51	13	3	-
Male	-	5	3	4	17	20	32	14	11	3	1	-
Female	2	5	5	9	15	30	45	58	40	10	2	-
Black, both sexes	-	2	-	2	1	2	5	3	-	-	-	-
Male	-	1	-	-	-	1	2	1	-	-	-	-
Female	-	1	-	2	1	1	3	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
Rhode Island												
All races, both sexes	1	-	-	-	1	5	3	7	3	2	-	-
Male	1	-	-	-	1	2	-	4	1	1	-	-
Female	-	-	-	-	-	3	3	3	2	1	-	-
White, both sexes	1	-	-	-	1	5	3	7	3	2	-	-
Male	1	-	-	-	1	2	-	4	1	1	-	-
Female	-	-	-	-	-	3	3	3	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	2	-	4	-	8	5	5	7	2	-	-
Male	-	1	-	3	-	2	3	3	1	-	-	-
Female	-	1	-	1	-	6	2	2	6	2	-	-
White, both sexes	-	2	-	2	-	8	4	4	6	1	-	-
Male	-	1	-	1	-	2	2	2	1	-	-	-
Female	-	1	-	1	-	6	2	2	5	1	-	-
Black, both sexes	-	-	-	2	-	-	1	1	1	1	-	-
Male	-	-	-	2	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
South Dakota												
All races, both sexes	-	-	-	1	1	-	2	1	2	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	-	1	-	-	1	-	2	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	2	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	4	4	9	14	18	23	13	19	2	1	-
Male	1	3	1	3	6	12	5	3	6	-	-	-
Female	1	1	3	6	8	6	18	10	13	2	1	-
White, both sexes	2	3	4	8	14	18	23	11	19	2	1	-
Male	1	2	1	2	6	12	5	3	6	-	-	-
Female	1	1	3	6	8	6	18	8	13	2	1	-
Black, both sexes	-	1	-	1	-	-	-	2	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
Texas												
All races, both sexes	2	-	3	10	21	28	28	21	14	7	1	-
Male	2	-	3	6	6	13	10	6	-	2	-	-
Female	-	-	-	4	15	15	18	15	14	5	1	-
White, both sexes	2	-	3	10	16	25	25	20	13	6	1	-
Male	2	-	3	6	4	12	10	6	-	1	-	-
Female	-	-	-	4	12	13	15	14	13	5	1	-
Black, both sexes	-	-	-	-	4	2	3	1	1	1	-	-
Male	-	-	-	-	2	1	-	-	-	1	-	-
Female	-	-	-	-	2	1	3	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	1	1	2	2	2	2	1	1	-	-
Male	-	-	1	-	1	2	-	1	-	-	-	-
Female	-	-	-	1	1	-	2	1	1	1	-	-
White, both sexes	-	-	1	1	2	2	2	2	1	1	-	-
Male	-	-	1	-	1	2	-	1	-	-	-	-
Female	-	-	-	1	1	-	2	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
Vermont												
All races, both sexes	-	-	2	-	1	2	5	7	-	5	-	-
Male	-	-	-	-	-	1	2	-	-	1	-	-
Female	-	-	2	-	1	1	3	7	-	4	-	-
White, both sexes	-	-	2	-	1	2	5	7	-	5	-	-
Male	-	-	-	-	-	1	2	-	-	1	-	-
Female	-	-	2	-	1	1	3	7	-	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	1	4	3	7	7	17	18	10	3	-	-
Male	1	-	2	1	4	6	6	9	2	1	-	-
Female	-	1	2	2	3	1	11	9	8	2	-	-
White, both sexes	-	1	4	3	5	5	16	18	10	3	-	-
Male	-	-	2	1	3	5	5	9	2	1	-	-
Female	-	1	2	2	2	-	11	9	8	2	-	-
Black, both sexes	1	-	-	-	2	2	1	-	-	-	-	-
Male	1	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
Washington												
All races, both sexes	1	-	2	2	6	10	17	8	12	2	1	-
Male	-	-	-	1	2	4	5	2	7	-	-	-
Female	1	-	2	1	4	6	12	6	5	2	1	-
White, both sexes	1	-	2	2	6	10	17	8	12	2	1	-
Male	-	-	-	1	2	4	5	2	7	-	-	-
Female	1	-	2	1	4	6	12	6	5	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	2	5	1	3	5	6	4	1	-	-	-
Male	-	1	3	-	-	3	4	1	-	-	-	-
Female	-	1	2	1	3	2	2	3	1	-	-	-
White, both sexes	-	2	5	1	3	5	5	4	1	-	-	-
Male	-	1	3	-	-	3	4	1	-	-	-	-
Female	-	1	2	1	3	2	1	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infections due to other specified organisms (A04,A07-A08)												
Wisconsin												
All races, both sexes	-	-	2	3	3	7	13	7	8	2	1	-
Male	-	-	2	2	1	3	6	5	2	-	-	-
Female	-	-	-	1	2	4	7	2	6	2	1	-
White, both sexes	-	-	2	3	3	5	13	7	7	2	1	-
Male	-	-	2	2	1	2	6	5	1	-	-	-
Female	-	-	-	1	2	3	7	2	6	2	1	-
Black, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
United States												
All races, both sexes	111	-	1	1	-	-	-	-	1	-	-	-
Male	35	-	1	-	-	-	-	-	-	-	-	-
Female	76	-	-	1	-	-	-	-	1	-	-	-
White, both sexes	98	-	-	1	-	-	-	-	-	-	-	-
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	70	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	11	-	1	-	-	-	-	-	1	-	-	-
Male	6	-	1	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
Florida												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
Illinois												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
Kentucky												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2	-	1	-	-	-	-	-	1	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	1	-	-	-	-	-	1	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
Maine												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
Massachusetts												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
Minnesota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
Missouri												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
North Carolina												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
Ohio												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
Texas												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
Washington												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diarrhea and gastroenteritis of infectious origin (A09)												
Wisconsin												
All races, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
United States												
All races, both sexes	4	3	3	6	10	19	24	19	14	6	-	-
Male	2	3	1	3	2	6	9	6	2	-	-	-
Female	2	-	2	3	8	13	15	13	12	6	-	-
White, both sexes	3	1	3	6	9	16	23	18	13	5	-	-
Male	1	1	1	3	2	5	9	5	1	-	-	-
Female	2	-	2	3	7	11	14	13	12	5	-	-
Black, both sexes	1	2	-	-	-	2	1	1	1	1	-	-
Male	1	2	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	2	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1	-	2	-	-	1	1	1	2	1	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	1	-	1	-	-	1	-	1	2	1	-	-
White, both sexes	1	-	2	-	-	1	1	1	2	1	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	1	-	1	-	-	1	-	1	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
Florida												
All races, both sexes	-	-	-	-	2	1	4	1	2	-	-	-
Male	-	-	-	-	-	1	1	-	1	-	-	-
Female	-	-	-	-	2	-	3	1	1	-	-	-
White, both sexes	-	-	-	-	2	1	4	1	2	-	-	-
Male	-	-	-	-	-	1	1	-	1	-	-	-
Female	-	-	-	-	2	-	3	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
Illinois												
All races, both sexes	-	2	-	-	1	3	1	1	2	-	-	-
Male	-	2	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	1	3	1	-	1	-	-	-
White, both sexes	-	1	-	-	1	3	1	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	3	1	-	1	-	-	-
Black, both sexes	-	1	-	-	-	-	-	1	1	-	-	-
Male	-	1	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
Maine												
All races, both sexes	-	-	-	-	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	1	-	-	1	-	2	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
White, both sexes	-	-	-	1	-	-	1	-	2	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
Massachusetts												
All races, both sexes	1	1	-	-	-	-	1	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
Minnesota												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
Missouri												
All races, both sexes	1	-	-	-	-	2	-	2	3	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	2	-	2	3	-	-	-
White, both sexes	1	-	-	-	-	1	-	2	3	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	2	3	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	2	-	2	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	2	-	1	1	-	-	-
White, both sexes	-	-	-	-	-	2	-	2	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	2	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	1	-	3	-	1	-	1	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	1	-	1	-	-
White, both sexes	-	-	-	1	-	2	-	1	-	1	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
North Carolina												
All races, both sexes	-	-	-	-	1	-	4	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	3	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	4	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	3	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
Ohio												
All races, both sexes	-	-	-	1	2	-	2	2	-	-	-	-
Male	-	-	-	-	1	-	2	1	-	-	-	-
Female	-	-	-	1	1	-	-	1	-	-	-	-
White, both sexes	-	-	-	1	2	-	2	2	-	-	-	-
Male	-	-	-	-	1	-	2	1	-	-	-	-
Female	-	-	-	1	1	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	1	2	1	1	1	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	1	-	-
White, both sexes	-	-	-	-	-	1	2	1	1	1	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	1	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
Texas												
All races, both sexes	-	-	-	-	-	2	2	2	-	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	2	2	2	-	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	1	-	-	-	-	-	2	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	2	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	2	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
Washington												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diarrhea and gastroenteritis of infectious origin (A09)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
United States												
All races, both sexes	657	1	1	-	2	1	6	6	10	24	29	33
Male	406	-	1	-	1	1	3	2	10	19	21	21
Female	251	1	-	-	1	-	3	4	-	5	8	12
White, both sexes	390	1	1	-	1	-	3	5	3	12	17	18
Male	237	-	1	-	1	-	2	2	3	8	11	12
Female	153	1	-	-	-	-	1	3	-	4	6	6
Black, both sexes	189	-	-	-	1	1	2	1	5	11	10	12
Male	118	-	-	-	-	1	1	-	5	10	8	6
Female	71	-	-	-	1	-	1	1	-	1	2	6
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	65	-	-	-	-	-	1	-	2	1	2	3
Male	42	-	-	-	-	-	-	-	2	1	2	3
Female	23	-	-	-	-	-	1	-	-	-	-	-
Alabama												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	3
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	3
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
Alaska												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	12	-	-	-	-	-	-	-	-	1	1	-
Male	7	-	-	-	-	-	-	-	-	1	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
Arkansas												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	117	-	-	-	-	-	1	1	1	5	6	6
Male	77	-	-	-	-	-	-	-	1	5	4	5
Female	40	-	-	-	-	-	1	1	-	-	2	1
White, both sexes	68	-	-	-	-	-	1	1	1	3	4	4
Male	42	-	-	-	-	-	-	-	1	3	2	3
Female	26	-	-	-	-	-	1	1	-	-	2	1
Black, both sexes	19	-	-	-	-	-	-	-	-	2	1	1
Male	15	-	-	-	-	-	-	-	-	2	1	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	30	-	-	-	-	-	-	-	-	-	1	1
Male	20	-	-	-	-	-	-	-	-	-	1	1
Female	10	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
Colorado												
All races, both sexes	8	-	-	-	-	-	-	-	-	2	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	2	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	2	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
Delaware												
All races, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	6	-	-	-	-	-	-	-	2	-	-	-
Male	5	-	-	-	-	-	-	-	2	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	2	-	-	-
Male	5	-	-	-	-	-	-	-	2	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
Florida												
All races, both sexes	39	-	-	-	-	-	-	-	1	2	2	2
Male	31	-	-	-	-	-	-	-	1	2	2	2
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	1	2	-	-
Male	16	-	-	-	-	-	-	-	1	2	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	15	-	-	-	-	-	-	-	-	-	2	2
Male	13	-	-	-	-	-	-	-	-	-	2	2
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	31	-	-	-	-	-	-	-	1	3	4	2
Male	19	-	-	-	-	-	-	-	1	2	2	-
Female	12	-	-	-	-	-	-	-	-	1	2	2
White, both sexes	13	-	-	-	-	-	-	-	-	2	2	1
Male	8	-	-	-	-	-	-	-	-	1	1	-
Female	5	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	18	-	-	-	-	-	-	-	1	1	2	1
Male	11	-	-	-	-	-	-	-	1	1	1	-
Female	7	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
Hawaii												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
Illinois												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	2	-
Male	14	-	-	-	-	-	-	-	-	-	2	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	9	-	-	-	-	-	-	1	-	-	1	1
Male	5	-	-	-	-	-	-	1	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	6	-	-	-	-	-	-	1	-	-	1	1
Male	4	-	-	-	-	-	-	1	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
Iowa												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
Kentucky												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	9	-	-	-	1	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	1	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	16	-	-	-	-	-	1	-	-	1	1	2
Male	9	-	-	-	-	-	1	-	-	1	1	1
Female	7	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	6	-	-	-	-	-	1	-	-	1	-	1
Male	5	-	-	-	-	-	1	-	-	1	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
Massachusetts												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	21	-	-	-	-	-	-	-	1	-	-	-
Male	14	-	-	-	-	-	-	-	1	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
Minnesota												
All races, both sexes	3	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Mississippi												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
Missouri												
All races, both sexes	10	-	-	-	-	-	-	-	1	-	-	1
Male	4	-	-	-	-	-	-	-	1	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
Nebraska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	23	-	-	-	-	-	-	-	-	1	1	2
Male	16	-	-	-	-	-	-	-	-	-	1	2
Female	7	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	1	1	1
Male	11	-	-	-	-	-	-	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
New Mexico												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	46	-	1	-	-	-	-	3	2	2	1	-
Male	24	-	1	-	-	-	-	1	2	2	1	-
Female	22	-	-	-	-	-	-	2	-	-	-	-
White, both sexes	22	-	1	-	-	-	-	2	1	-	-	-
Male	11	-	1	-	-	-	-	1	1	-	-	-
Female	11	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	18	-	-	-	-	-	-	1	1	1	1	-
Male	9	-	-	-	-	-	-	-	1	1	1	-
Female	9	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
North Carolina												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	2	1
Male	14	-	-	-	-	-	-	-	-	-	1	1
Female	9	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	12	-	-	-	-	-	-	-	-	-	2	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
Ohio												
All races, both sexes	23	-	-	-	-	-	1	-	-	-	1	2
Male	10	-	-	-	-	-	1	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	13	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	8	-	-	-	-	-	1	-	-	-	-	1
Male	4	-	-	-	-	-	1	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	7	-	-	-	-	-	1	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
Oregon												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	16	-	-	-	-	1	-	-	1	1	-	-
Male	9	-	-	-	-	1	-	-	1	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	1	-	-	-	1	-	-
Male	3	-	-	-	-	1	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	12	-	-	-	-	-	-	-	-	1	-	2
Male	7	-	-	-	-	-	-	-	-	1	-	2
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	1	-	2
Male	5	-	-	-	-	-	-	-	-	1	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
South Dakota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	15	-	-	-	-	-	1	1	-	-	-	-
Male	9	-	-	-	-	-	1	-	-	-	-	-
Female	6	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	10	-	-	-	-	-	1	1	-	-	-	-
Male	6	-	-	-	-	-	1	-	-	-	-	-
Female	4	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
Texas												
All races, both sexes	58	1	-	-	1	-	-	-	-	2	3	6
Male	39	-	-	-	1	-	-	-	-	2	2	3
Female	19	1	-	-	-	-	-	-	-	-	1	3
White, both sexes	48	1	-	-	1	-	-	-	-	-	2	5
Male	31	-	-	-	1	-	-	-	-	-	1	2
Female	17	1	-	-	-	-	-	-	-	-	1	3
Black, both sexes	7	-	-	-	-	-	-	-	-	2	1	-
Male	6	-	-	-	-	-	-	-	-	2	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	24	-	-	-	-	-	-	-	-	1	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	12	-	-	-	-	-	-	-	-	1	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
Washington												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tuberculosis (A16-A19)												
Wisconsin												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
United States												
All races, both sexes	47	35	44	55	82	81	84	68	37	11	-	-
Male	32	26	34	38	55	37	47	38	17	3	-	-
Female	15	9	10	17	27	44	37	30	20	8	-	-
White, both sexes	20	19	23	29	49	55	58	47	24	5	-	-
Male	14	15	19	19	34	28	30	27	11	-	-	-
Female	6	4	4	10	15	27	28	20	13	5	-	-
Black, both sexes	20	11	15	15	26	14	18	16	7	4	-	-
Male	12	9	11	10	16	4	13	8	3	1	-	-
Female	8	2	4	5	10	10	5	8	4	3	-	-
American Indian 1/, both sexes	-	1	1	5	-	3	2	-	1	-	-	-
Male	-	1	1	4	-	2	1	-	-	-	-	-
Female	-	-	-	1	-	1	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	7	4	5	6	7	9	6	5	5	2	-	-
Male	6	1	3	5	5	3	3	3	3	2	-	-
Female	1	3	2	1	2	6	3	2	2	-	-	-
Alabama												
All races, both sexes	-	1	-	1	2	-	1	-	-	-	-	-
Male	-	1	-	1	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	1	-	1	-	-	1	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
Alaska												
All races, both sexes	-	-	1	-	-	1	1	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	1	1	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	1	1	2	1	3	1	-	-	1	-	-
Male	-	1	-	1	1	2	-	-	-	-	-	-
Female	-	-	1	1	-	1	1	-	-	1	-	-
White, both sexes	-	-	1	2	1	2	1	-	-	1	-	-
Male	-	-	-	1	1	1	-	-	-	-	-	-
Female	-	-	1	1	-	1	1	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
Arkansas												
All races, both sexes	1	1	-	1	-	-	1	1	-	-	-	-
Male	-	1	-	1	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	1	-	-	1	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	8	5	7	8	15	14	16	12	11	1	-	-
Male	5	4	6	6	9	4	15	6	7	-	-	-
Female	3	1	1	2	6	10	1	6	4	1	-	-
White, both sexes	4	4	4	3	7	5	12	9	5	1	-	-
Male	2	3	4	2	3	1	11	4	3	-	-	-
Female	2	1	-	1	4	4	1	5	2	1	-	-
Black, both sexes	3	1	1	1	2	3	2	-	2	-	-	-
Male	3	1	1	-	2	1	2	-	1	-	-	-
Female	-	-	-	1	-	2	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	4	6	6	2	3	4	-	-	-
Male	-	-	1	4	4	2	2	2	3	-	-	-
Female	1	-	1	-	2	4	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
Colorado												
All races, both sexes	-	1	1	-	3	-	-	-	1	-	-	-
Male	-	1	-	-	3	-	-	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	3	-	-	-	1	-	-	-
Male	-	1	-	-	3	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	2	2	2	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	1	2	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	2	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
Delaware												
All races, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	2	-	1	1	-	-	-	-	-
Male	-	-	-	2	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	2	-	1	1	-	-	-	-	-
Male	-	-	-	2	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
Florida												
All races, both sexes	5	4	5	4	4	3	3	3	-	1	-	-
Male	5	2	5	2	4	1	2	2	-	1	-	-
Female	-	2	-	2	-	2	1	1	-	-	-	-
White, both sexes	3	2	2	2	3	2	3	2	-	-	-	-
Male	3	1	2	-	3	1	2	1	-	-	-	-
Female	-	1	-	2	-	1	1	1	-	-	-	-
Black, both sexes	1	2	3	2	1	1	-	1	-	-	-	-
Male	1	1	3	2	1	-	-	1	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	3	2	1	1	4	4	2	4	-	-	-	-
Male	2	2	1	1	3	2	1	2	-	-	-	-
Female	1	-	-	-	1	2	1	2	-	-	-	-
White, both sexes	-	1	1	-	2	2	1	1	-	-	-	-
Male	-	1	1	-	2	1	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	3	1	-	1	2	2	1	3	-	-	-	-
Male	2	1	-	1	1	1	1	1	-	-	-	-
Female	1	-	-	-	1	1	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
Hawaii												
All races, both sexes	1	-	-	-	-	1	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
Illinois												
All races, both sexes	-	1	2	2	1	5	4	1	1	1	-	-
Male	-	1	1	2	1	4	2	1	-	-	-	-
Female	-	-	1	-	-	1	2	-	1	1	-	-
White, both sexes	-	-	1	-	-	5	2	1	1	-	-	-
Male	-	-	1	-	-	4	1	1	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	-	-	-
Black, both sexes	-	-	1	1	1	-	-	-	-	1	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	2	-	-	-	-	-
Male	-	1	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	2	-	1	2	-	-	1	-	-
Male	-	-	-	1	-	-	2	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	1	-	-
White, both sexes	-	-	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
Iowa												
All races, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
Kentucky												
All races, both sexes	-	-	1	1	4	2	1	1	2	1	-	-
Male	-	-	1	-	4	2	-	1	1	-	-	-
Female	-	-	-	1	-	-	1	-	1	1	-	-
White, both sexes	-	-	1	1	4	2	1	-	1	1	-	-
Male	-	-	1	-	4	2	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2	-	-	1	1	-	2	2	-	-	-	-
Male	2	-	-	1	1	-	2	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	1	-	-	1	-	-	2	1	-	-	-	-
Male	1	-	-	1	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	1	-	-	1	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2	1	-	1	3	-	2	-	1	1	-	-
Male	-	1	-	-	1	-	2	-	-	1	-	-
Female	2	-	-	1	2	-	-	-	1	-	-	-
White, both sexes	-	-	-	1	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	2	1	-	-	2	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	2	-	-	-	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
Massachusetts												
All races, both sexes	-	-	-	-	1	1	2	3	-	1	-	-
Male	-	-	-	-	-	1	1	2	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	1	-	-
White, both sexes	-	-	-	-	1	1	2	2	-	1	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	4	1	2	-	4	3	3	1	2	-	-	-
Male	3	1	2	-	3	2	-	1	1	-	-	-
Female	1	-	-	-	1	1	3	-	1	-	-	-
White, both sexes	1	1	-	-	3	1	3	-	2	-	-	-
Male	1	1	-	-	2	1	-	-	1	-	-	-
Female	-	-	-	-	1	-	3	-	1	-	-	-
Black, both sexes	1	-	2	-	1	1	-	1	-	-	-	-
Male	-	-	2	-	1	-	-	1	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	-	-	-	-	1	3	1	-	1	-	-
Male	1	-	-	-	-	-	2	1	-	-	-	-
Female	1	-	-	-	-	1	1	-	-	1	-	-
White, both sexes	1	-	-	-	-	1	2	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	1	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	1	-	-	1	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
Missouri												
All races, both sexes	-	-	1	1	-	-	2	2	2	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	2	1	2	-	-	-
White, both sexes	-	-	1	1	-	-	2	2	2	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	2	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
Nebraska												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2	-	1	-	1	-	-	1	-	-	-	-
Male	1	-	1	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	-	4	1	3	5	-	3	2	-	-	-
Male	1	-	4	1	2	2	-	3	-	-	-	-
Female	-	-	-	-	1	3	-	-	2	-	-	-
White, both sexes	1	-	3	1	1	2	-	3	2	-	-	-
Male	1	-	3	1	1	-	-	3	-	-	-	-
Female	-	-	-	-	-	2	-	-	2	-	-	-
Black, both sexes	-	-	1	-	2	3	-	-	-	-	-	-
Male	-	-	1	-	1	2	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
New Mexico												
All races, both sexes	-	-	-	2	-	1	1	1	-	-	-	-
Male	-	-	-	2	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	3	5	5	2	6	5	5	2	4	-	-	-
Male	2	2	3	1	5	1	2	-	1	-	-	-
Female	1	3	2	1	1	4	3	2	3	-	-	-
White, both sexes	2	2	-	1	4	3	2	1	3	-	-	-
Male	1	1	-	-	3	1	1	-	1	-	-	-
Female	1	1	-	1	1	2	1	1	2	-	-	-
Black, both sexes	-	2	4	1	1	1	3	1	1	-	-	-
Male	-	1	2	1	1	-	1	-	-	-	-	-
Female	-	1	2	-	-	1	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	-	1	1	-	-	-	-	-	-
Male	1	-	1	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
North Carolina												
All races, both sexes	1	2	1	1	4	1	6	3	1	-	-	-
Male	-	2	1	-	3	-	4	2	-	-	-	-
Female	1	-	-	1	1	1	2	1	1	-	-	-
White, both sexes	-	1	1	-	3	1	2	1	-	-	-	-
Male	-	1	1	-	3	-	2	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	1	1	-	1	1	-	3	2	1	-	-	-
Male	-	1	-	-	-	-	2	1	-	-	-	-
Female	1	-	-	1	1	-	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
Ohio												
All races, both sexes	1	-	-	2	3	2	3	7	-	1	-	-
Male	1	-	-	2	2	-	-	3	-	-	-	-
Female	-	-	-	-	1	2	3	4	-	1	-	-
White, both sexes	-	-	-	1	1	2	2	5	-	1	-	-
Male	-	-	-	1	1	-	-	3	-	-	-	-
Female	-	-	-	-	-	2	2	2	-	1	-	-
Black, both sexes	1	-	-	1	2	-	-	2	-	-	-	-
Male	1	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Oklahoma												
All races, both sexes	-	1	1	-	1	1	-	-	1	-	-	-
Male	-	1	1	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
White, both sexes	-	1	1	-	1	1	-	-	1	-	-	-
Male	-	1	1	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
Oregon												
All races, both sexes	-	-	-	1	3	1	3	-	-	-	-	-
Male	-	-	-	1	2	1	1	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	-	-	-	-
White, both sexes	-	-	-	-	3	1	3	-	-	-	-	-
Male	-	-	-	-	2	1	1	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	1	1	2	2	3	2	-	2	-	-	-
Male	-	-	1	2	-	2	-	-	1	-	-	-
Female	-	1	-	-	2	1	2	-	1	-	-	-
White, both sexes	-	-	1	1	1	2	2	-	2	-	-	-
Male	-	-	1	1	-	2	-	-	1	-	-	-
Female	-	-	-	-	1	-	2	-	1	-	-	-
Black, both sexes	-	-	-	1	1	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	-	-	1	3	1	1	1	-	-	-	-
Male	2	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	1	2	-	1	1	-	-	-	-
White, both sexes	1	-	-	-	1	1	1	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Black, both sexes	1	-	-	1	2	-	-	1	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
South Dakota												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	2	2	1	3	2	2	1	-	-	-
Male	-	-	1	2	1	2	-	1	1	-	-	-
Female	-	-	1	-	-	1	2	1	-	-	-	-
White, both sexes	-	-	1	2	-	2	2	1	-	-	-	-
Male	-	-	1	2	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	1	-	-	-	-
Black, both sexes	-	-	1	-	1	1	-	1	1	-	-	-
Male	-	-	-	-	1	-	-	1	1	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
Texas												
All races, both sexes	5	6	3	6	3	12	3	6	1	-	-	-
Male	4	5	2	4	1	7	2	5	1	-	-	-
Female	1	1	1	2	2	5	1	1	-	-	-	-
White, both sexes	3	4	2	5	3	12	3	6	1	-	-	-
Male	2	4	1	4	1	7	2	5	1	-	-	-
Female	1	-	1	1	2	5	1	1	-	-	-	-
Black, both sexes	1	1	1	1	-	-	-	-	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	1	3	-	6	1	6	2	2	1	-	-
Male	-	1	3	-	4	-	4	-	1	1	-	-
Female	1	-	-	-	2	1	2	2	1	-	-	-
White, both sexes	1	-	2	-	2	1	2	2	2	-	-	-
Male	-	-	2	-	2	-	-	-	1	-	-	-
Female	1	-	-	-	-	1	2	2	1	-	-	-
Black, both sexes	-	1	1	-	4	-	4	-	-	1	-	-
Male	-	1	1	-	2	-	4	-	-	1	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
Washington												
All races, both sexes	-	-	-	1	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	-	-	2	-	1	-	-	-	-	-	-
Male	1	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	1	-	-	2	-	1	-	-	-	-	-	-
Male	1	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
Wisconsin												
All races, both sexes	2	1	-	1	-	1	1	2	-	-	-	-
Male	1	-	-	1	-	1	-	1	-	-	-	-
Female	1	1	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	1	-	1	-	1	1	1	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	1	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
United States												
All races, both sexes	504	1	1	-	1	1	3	4	7	16	20	24
Male	316	-	1	-	-	1	1	1	7	12	16	14
Female	188	1	-	-	1	-	2	3	-	4	4	10
White, both sexes	300	1	1	-	-	-	1	3	1	10	10	13
Male	185	-	1	-	-	-	-	1	1	7	8	7
Female	115	1	-	-	-	-	1	2	-	3	2	6
Black, both sexes	141	-	-	-	1	1	1	1	4	6	9	8
Male	88	-	-	-	-	1	1	-	4	5	7	4
Female	53	-	-	-	1	-	-	1	-	1	2	4
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	54	-	-	-	-	-	1	-	2	-	1	3
Male	37	-	-	-	-	-	-	-	2	-	1	3
Female	17	-	-	-	-	-	1	-	-	-	-	-
Alabama												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	3
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	3
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Alaska												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Arkansas												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	92	-	-	-	-	-	1	-	-	4	2	5
Male	62	-	-	-	-	-	-	-	-	4	2	4
Female	30	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	50	-	-	-	-	-	1	-	-	3	1	3
Male	31	-	-	-	-	-	-	-	-	3	1	2
Female	19	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	14	-	-	-	-	-	-	-	-	1	1	1
Male	12	-	-	-	-	-	-	-	-	1	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	28	-	-	-	-	-	-	-	-	-	-	1
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Colorado												
All races, both sexes	7	-	-	-	-	-	-	-	-	2	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	2	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	2	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Delaware												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Florida												
All races, both sexes	27	-	-	-	-	-	-	-	1	2	1	1
Male	20	-	-	-	-	-	-	-	1	2	1	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	1	2	-	-
Male	11	-	-	-	-	-	-	-	1	2	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	25	-	-	-	-	-	-	-	1	1	4	1
Male	16	-	-	-	-	-	-	-	1	1	2	-
Female	9	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	12	-	-	-	-	-	-	-	-	1	2	1
Male	8	-	-	-	-	-	-	-	-	1	1	-
Female	4	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	13	-	-	-	-	-	-	-	1	-	2	-
Male	8	-	-	-	-	-	-	-	1	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Hawaii												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Illinois												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	2	-
Male	13	-	-	-	-	-	-	-	-	-	2	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	7	-	-	-	-	-	-	1	-	-	-	1
Male	4	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	4	-	-	-	-	-	-	1	-	-	-	1
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Iowa												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Kentucky												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	8	-	-	-	1	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	11	-	-	-	-	-	-	-	-	1	1	2
Male	6	-	-	-	-	-	-	-	-	1	1	1
Female	5	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	4	-	-	-	-	-	-	-	-	1	-	1
Male	3	-	-	-	-	-	-	-	-	1	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Massachusetts												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	19	-	-	-	-	-	-	-	1	-	-	-
Male	13	-	-	-	-	-	-	-	1	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Minnesota												
All races, both sexes	3	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Mississippi												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Missouri												
All races, both sexes	9	-	-	-	-	-	-	-	1	-	-	1
Male	4	-	-	-	-	-	-	-	1	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Nebraska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	17	-	-	-	-	-	-	-	-	1	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	1	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
New Mexico												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	33	-	1	-	-	-	-	2	1	1	1	-
Male	16	-	1	-	-	-	-	-	1	1	1	-
Female	17	-	-	-	-	-	-	2	-	-	-	-
White, both sexes	15	-	1	-	-	-	-	1	-	-	-	-
Male	6	-	1	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	14	-	-	-	-	-	-	1	1	1	1	-
Male	8	-	-	-	-	-	-	-	1	1	1	-
Female	6	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
North Carolina												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	2	-
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	2	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Ohio												
All races, both sexes	16	-	-	-	-	-	1	-	-	-	-	1
Male	8	-	-	-	-	-	1	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Oregon												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	10	-	-	-	-	1	-	-	1	-	-	-
Male	4	-	-	-	-	1	-	-	1	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	9	-	-	-	-	-	-	-	-	1	-	2
Male	6	-	-	-	-	-	-	-	-	1	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	1	-	2
Male	5	-	-	-	-	-	-	-	-	1	-	2
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
South Dakota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	10	-	-	-	-	-	-	1	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	1	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Texas												
All races, both sexes	49	1	-	-	-	-	-	-	-	2	3	5
Male	32	-	-	-	-	-	-	-	-	2	2	2
Female	17	1	-	-	-	-	-	-	-	-	1	3
White, both sexes	40	1	-	-	-	-	-	-	-	-	2	4
Male	25	-	-	-	-	-	-	-	-	-	1	1
Female	15	1	-	-	-	-	-	-	-	-	1	3
Black, both sexes	6	-	-	-	-	-	-	-	-	2	1	-
Male	5	-	-	-	-	-	-	-	-	2	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	19	-	-	-	-	-	-	-	-	1	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Washington												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Wisconsin												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
United States												
All races, both sexes	34	26	32	41	67	57	63	62	35	9	-	-
Male	26	18	25	30	44	28	36	36	17	3	-	-
Female	8	8	7	11	23	29	27	26	18	6	-	-
White, both sexes	14	14	18	24	39	37	45	42	23	4	-	-
Male	10	11	15	16	27	22	23	25	11	-	-	-
Female	4	3	3	8	12	15	22	17	12	4	-	-
Black, both sexes	14	7	10	10	21	10	13	16	6	3	-	-
Male	11	5	8	7	12	2	9	8	3	1	-	-
Female	3	2	2	3	9	8	4	8	3	2	-	-
American Indian 1/, both sexes	-	1	-	2	-	3	2	-	1	-	-	-
Male	-	1	-	2	-	2	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	6	4	4	5	7	7	3	4	5	2	-	-
Male	5	1	2	5	5	2	3	3	3	2	-	-
Female	1	3	2	-	2	5	-	1	2	-	-	-
Alabama												
All races, both sexes	-	1	-	1	2	-	1	-	-	-	-	-
Male	-	1	-	1	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	1	-	1	-	-	1	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
Alaska												
All races, both sexes	-	-	-	-	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	1	1	1	1	3	1	-	-	-	-	-
Male	-	1	-	1	1	2	-	-	-	-	-	-
Female	-	-	1	-	-	1	1	-	-	-	-	-
White, both sexes	-	-	1	1	1	2	1	-	-	-	-	-
Male	-	-	-	1	1	1	-	-	-	-	-	-
Female	-	-	1	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
Arkansas												
All races, both sexes	1	1	-	1	-	-	1	1	-	-	-	-
Male	-	1	-	1	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	1	-	-	1	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	7	4	7	7	12	9	11	11	11	1	-	-
Male	5	3	6	6	6	3	10	6	7	-	-	-
Female	2	1	1	1	6	6	1	5	4	1	-	-
White, both sexes	3	3	4	3	5	2	8	8	5	1	-	-
Male	2	2	4	2	1	-	7	4	3	-	-	-
Female	1	1	-	1	4	2	1	4	2	1	-	-
Black, both sexes	3	1	1	-	1	2	1	-	2	-	-	-
Male	3	1	1	-	1	1	1	-	1	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	4	6	5	2	3	4	-	-	-
Male	-	-	1	4	4	2	2	2	3	-	-	-
Female	1	-	1	-	2	3	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
Colorado												
All races, both sexes	-	-	1	-	3	-	-	-	1	-	-	-
Male	-	-	-	-	3	-	-	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	3	-	-	-	1	-	-	-
Male	-	-	-	-	3	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	2	2	2	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	1	2	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	2	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
Delaware												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
Florida												
All races, both sexes	4	3	3	3	2	1	2	3	-	1	-	-
Male	4	1	3	1	2	-	1	2	-	1	-	-
Female	-	2	-	2	-	1	1	1	-	-	-	-
White, both sexes	2	2	1	2	2	-	2	2	-	-	-	-
Male	2	1	1	-	2	-	1	1	-	-	-	-
Female	-	1	-	2	-	-	1	1	-	-	-	-
Black, both sexes	1	1	2	1	-	1	-	1	-	-	-	-
Male	1	-	2	1	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2	1	1	1	4	3	2	4	-	-	-	-
Male	2	1	1	1	3	1	1	2	-	-	-	-
Female	-	-	-	-	1	2	1	2	-	-	-	-
White, both sexes	-	1	1	-	2	2	1	1	-	-	-	-
Male	-	1	1	-	2	1	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	2	-	-	1	2	1	1	3	-	-	-	-
Male	2	-	-	1	1	-	1	1	-	-	-	-
Female	-	-	-	-	1	1	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
Hawaii												
All races, both sexes	1	-	-	-	-	1	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
Illinois												
All races, both sexes	-	1	1	2	1	4	2	1	1	1	-	-
Male	-	1	-	2	1	4	2	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	1	-	-
White, both sexes	-	-	-	-	-	4	1	1	1	-	-	-
Male	-	-	-	-	-	4	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	1	1	1	-	-	-	-	1	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	1	-	-	-	-	-
Male	-	1	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	2	-	-	2	-	-	1	-	-
Male	-	-	-	1	-	-	2	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
Iowa												
All races, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
Kentucky												
All races, both sexes	-	-	1	1	3	2	1	1	2	1	-	-
Male	-	-	1	-	3	2	-	1	1	-	-	-
Female	-	-	-	1	-	-	1	-	1	1	-	-
White, both sexes	-	-	1	1	3	2	1	-	1	1	-	-
Male	-	-	1	-	3	2	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	-	-	1	1	-	2	2	-	-	-	-
Male	1	-	-	1	1	-	2	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	1	-	-	1	-	-	2	1	-	-	-	-
Male	1	-	-	1	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	1	3	-	1	-	1	1	-	-
Male	-	-	-	-	1	-	1	-	-	1	-	-
Female	-	-	-	1	2	-	-	-	1	-	-	-
White, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	2	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
Massachusetts												
All races, both sexes	-	-	-	-	1	1	1	2	-	1	-	-
Male	-	-	-	-	-	1	-	2	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	1	-	-
White, both sexes	-	-	-	-	1	1	1	1	-	1	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	4	1	2	-	4	2	2	1	2	-	-	-
Male	3	1	2	-	3	1	-	1	1	-	-	-
Female	1	-	-	-	1	1	2	-	1	-	-	-
White, both sexes	1	1	-	-	3	1	2	-	2	-	-	-
Male	1	1	-	-	2	1	-	-	1	-	-	-
Female	-	-	-	-	1	-	2	-	1	-	-	-
Black, both sexes	1	-	2	-	1	1	-	1	-	-	-	-
Male	-	-	2	-	1	-	-	1	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	-	-	-	-	-	3	1	-	-	-	-
Male	1	-	-	-	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
Missouri												
All races, both sexes	-	-	-	1	-	-	2	2	2	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	2	1	2	-	-	-
White, both sexes	-	-	-	1	-	-	2	2	2	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	2	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
Nebraska												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	-	3	1	2	3	-	3	2	-	-	-
Male	1	-	3	1	2	1	-	3	-	-	-	-
Female	-	-	-	-	-	2	-	-	2	-	-	-
White, both sexes	1	-	3	1	1	1	-	3	2	-	-	-
Male	1	-	3	1	1	-	-	3	-	-	-	-
Female	-	-	-	-	-	1	-	-	2	-	-	-
Black, both sexes	-	-	-	-	1	2	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2	4	3	2	4	4	4	2	2	-	-	-
Male	1	1	2	1	3	1	2	-	1	-	-	-
Female	1	3	1	1	1	3	2	2	1	-	-	-
White, both sexes	2	1	-	1	2	2	2	1	2	-	-	-
Male	1	-	-	-	1	1	1	-	1	-	-	-
Female	1	1	-	1	1	1	1	1	1	-	-	-
Black, both sexes	-	2	2	1	1	1	2	1	-	-	-	-
Male	-	1	1	1	1	-	1	-	-	-	-	-
Female	-	1	1	-	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	1	1	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
North Carolina												
All races, both sexes	-	2	-	1	4	1	4	3	1	-	-	-
Male	-	2	-	-	3	-	3	2	-	-	-	-
Female	-	-	-	1	1	1	1	1	1	-	-	-
White, both sexes	-	1	-	-	3	1	2	1	-	-	-	-
Male	-	1	-	-	3	-	2	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	1	-	1	1	-	2	2	1	-	-	-
Male	-	1	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	1	1	-	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
Ohio												
All races, both sexes	1	-	-	2	2	1	1	6	-	1	-	-
Male	1	-	-	2	1	-	-	2	-	-	-	-
Female	-	-	-	-	1	1	1	4	-	1	-	-
White, both sexes	-	-	-	1	1	1	1	4	-	1	-	-
Male	-	-	-	1	1	-	-	2	-	-	-	-
Female	-	-	-	-	-	1	1	2	-	1	-	-
Black, both sexes	1	-	-	1	1	-	-	2	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	1	-	1	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	1	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
Oregon												
All races, both sexes	-	-	-	1	3	-	2	-	-	-	-	-
Male	-	-	-	1	2	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	3	-	2	-	-	-	-	-
Male	-	-	-	-	2	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	1	-	1	2	1	1	-	2	-	-	-
Male	-	-	-	1	-	-	-	-	1	-	-	-
Female	-	1	-	-	2	1	1	-	1	-	-	-
White, both sexes	-	-	-	1	1	-	1	-	2	-	-	-
Male	-	-	-	1	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	-	-	-	2	1	1	1	-	-	-	-
Male	1	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	1	-	-	-	2	-	-	1	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
South Dakota												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	1	1	1	2	1	2	1	-	-	-
Male	-	-	1	1	1	2	-	1	1	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	1	1	-	2	1	1	-	-	-	-
Male	-	-	1	1	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	1	1	-	-	-
Male	-	-	-	-	1	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
Texas												
All races, both sexes	4	5	3	5	2	10	3	5	1	-	-	-
Male	3	4	2	3	1	6	2	4	1	-	-	-
Female	1	1	1	2	1	4	1	1	-	-	-	-
White, both sexes	2	4	2	4	2	10	3	5	1	-	-	-
Male	1	4	1	3	1	6	2	4	1	-	-	-
Female	1	-	1	1	1	4	1	1	-	-	-	-
Black, both sexes	1	-	1	1	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	1	3	-	4	1	4	1	2	1	-	-
Male	-	1	3	-	2	-	2	-	1	1	-	-
Female	1	-	-	-	2	1	2	1	1	-	-	-
White, both sexes	1	-	2	-	1	1	2	1	2	-	-	-
Male	-	-	2	-	1	-	-	-	1	-	-	-
Female	1	-	-	-	-	1	2	1	1	-	-	-
Black, both sexes	-	1	1	-	3	-	2	-	-	1	-	-
Male	-	1	1	-	1	-	2	-	-	1	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
Washington												
All races, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	-	-	2	-	1	-	-	-	-	-	-
Male	1	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	1	-	-	2	-	1	-	-	-	-	-	-
Male	1	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)												
Wisconsin												
All races, both sexes	1	-	-	-	-	1	1	2	-	-	-	-
Male	1	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
United States												
All races, both sexes	153	-	-	-	1	-	3	2	3	8	9	9
Male	90	-	-	-	1	-	2	1	3	7	5	7
Female	63	-	-	-	-	-	1	1	-	1	4	2
White, both sexes	90	-	-	-	1	-	2	2	2	2	7	5
Male	52	-	-	-	1	-	2	1	2	1	3	5
Female	38	-	-	-	-	-	-	1	-	1	4	-
Black, both sexes	48	-	-	-	-	-	1	-	1	5	1	4
Male	30	-	-	-	-	-	-	-	1	5	1	2
Female	18	-	-	-	-	-	1	-	-	-	-	2
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	1	1	-
Male	5	-	-	-	-	-	-	-	-	1	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Alaska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	25	-	-	-	-	-	-	1	1	1	4	1
Male	15	-	-	-	-	-	-	-	1	1	2	1
Female	10	-	-	-	-	-	-	1	-	-	2	-
White, both sexes	18	-	-	-	-	-	-	1	1	-	3	1
Male	11	-	-	-	-	-	-	-	1	-	1	1
Female	7	-	-	-	-	-	-	1	-	-	2	-
Black, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Colorado												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Delaware												
All races, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Florida												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	1	1
Male	11	-	-	-	-	-	-	-	-	-	1	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	6	-	-	-	-	-	-	-	-	2	-	1
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	1	-	1
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Illinois												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Kentucky												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	5	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Massachusetts												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Missouri												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
New Mexico												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	13	-	-	-	-	-	-	1	1	1	-	-
Male	8	-	-	-	-	-	-	1	1	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	1	1	-	-	-
Male	5	-	-	-	-	-	-	1	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
North Carolina												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Ohio												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	4	-	-	-	-	-	1	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Oregon												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	5	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Texas												
All races, both sexes	9	-	-	-	1	-	-	-	-	-	-	1
Male	7	-	-	-	1	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	1	-	-	-	-	-	-	1
Male	6	-	-	-	1	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Washington												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Wisconsin												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
United States												
All races, both sexes	13	9	12	14	15	24	21	6	2	2	-	-
Male	6	8	9	8	11	9	11	2	-	-	-	-
Female	7	1	3	6	4	15	10	4	2	2	-	-
White, both sexes	6	5	5	5	10	18	13	5	1	1	-	-
Male	4	4	4	3	7	6	7	2	-	-	-	-
Female	2	1	1	2	3	12	6	3	1	1	-	-
Black, both sexes	6	4	5	5	5	4	5	-	1	1	-	-
Male	1	4	3	3	4	2	4	-	-	-	-	-
Female	5	-	2	2	1	2	1	-	1	1	-	-
American Indian 1/, both sexes	-	-	1	3	-	-	-	-	-	-	-	-
Male	-	-	1	2	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	-	2	3	1	-	-	-	-
Male	1	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	1	3	1	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
Alaska												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	1	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1	1	-	1	3	5	5	1	-	-	-	-
Male	-	1	-	-	3	1	5	-	-	-	-	-
Female	1	-	-	1	-	4	-	1	-	-	-	-
White, both sexes	1	1	-	-	2	3	4	1	-	-	-	-
Male	-	1	-	-	2	1	4	-	-	-	-	-
Female	1	-	-	-	-	2	-	1	-	-	-	-
Black, both sexes	-	-	-	1	1	1	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
Colorado												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
Delaware												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
Florida												
All races, both sexes	1	1	2	1	2	2	1	-	-	-	-	-
Male	1	1	2	1	2	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	1	-	1	-	1	2	1	-	-	-	-	-
Male	1	-	1	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	1	1	1	1	-	-	-	-	-	-	-
Male	-	1	1	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
Illinois												
All races, both sexes	-	-	1	-	-	1	2	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	2	-	-	-	-	-
White, both sexes	-	-	1	-	-	1	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
Kentucky												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	1	-	-
White, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
Missouri												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	-	1	-	-	-	-	1	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	1	-	1	2	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	1	-	1	1	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
New Mexico												
All races, both sexes	-	-	-	2	-	1	-	-	-	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1	1	2	-	2	1	1	-	2	-	-	-
Male	1	1	1	-	2	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	1	-	2	-	-	-
White, both sexes	-	1	-	-	2	1	-	-	1	-	-	-
Male	-	1	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Black, both sexes	-	-	2	-	-	-	1	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
North Carolina												
All races, both sexes	1	-	1	-	-	-	2	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
Ohio												
All races, both sexes	-	-	-	-	1	1	2	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	2	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Oklahoma												
All races, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
Oregon												
All races, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	1	1	-	2	1	-	-	-	-	-
Male	-	-	1	1	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	1	-	-	2	1	-	-	-	-	-
Male	-	-	1	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	-	-	1	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	1	1	-	1	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	1	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
Texas												
All races, both sexes	1	1	-	1	1	2	-	1	-	-	-	-
Male	1	1	-	1	-	1	-	1	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
White, both sexes	1	-	-	1	1	2	-	1	-	-	-	-
Male	1	-	-	1	-	1	-	1	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	2	-	2	1	-	-	-	-
Male	-	-	-	-	2	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	1	-	2	-	-	-	-	-
Male	-	-	-	-	1	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
Washington												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)												
Wisconsin												
All races, both sexes	1	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
United States												
All races, both sexes	35,152	307	107	48	38	65	109	132	233	332	579	964
Male	15,895	186	55	24	18	36	59	63	126	175	292	524
Female	19,257	121	52	24	20	29	50	69	107	157	287	440
White, both sexes	28,139	195	78	39	27	46	64	88	142	238	363	630
Male	12,702	119	40	21	14	26	35	42	76	131	186	343
Female	15,437	76	38	18	13	20	29	46	66	107	177	287
Black, both sexes	6,269	100	24	6	8	17	37	40	75	78	197	297
Male	2,825	57	12	2	3	9	19	19	41	34	99	159
Female	3,444	43	12	4	5	8	18	21	34	44	98	138
American Indian 1/, both sexes	203	6	2	2	3	-	2	3	8	8	11	18
Male	95	6	2	-	1	-	1	1	3	6	4	9
Female	108	-	-	2	2	-	1	2	5	2	7	9
Asian or Pacific Islander 2/, both sexes	541	6	3	1	-	2	6	1	8	8	8	19
Male	273	4	1	1	-	1	4	1	6	4	3	13
Female	268	2	2	-	-	1	2	-	2	4	5	6
Alabama												
All races, both sexes	798	6	-	-	2	1	3	3	10	12	10	31
Male	311	3	-	-	1	-	2	1	5	4	6	13
Female	487	3	-	-	1	1	1	2	5	8	4	18
White, both sexes	549	2	-	-	2	-	1	1	3	7	6	17
Male	218	2	-	-	1	-	1	-	1	3	3	8
Female	331	-	-	-	1	-	-	1	2	4	3	9
Black, both sexes	246	4	-	-	-	1	2	2	7	4	4	13
Male	90	1	-	-	-	-	1	1	4	-	3	4
Female	156	3	-	-	-	1	1	1	3	4	1	9
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	1	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
Alaska												
All races, both sexes	25	1	2	-	1	-	1	1	1	1	1	2
Male	13	1	2	-	1	-	1	-	-	1	-	1
Female	12	-	-	-	-	-	-	1	1	-	1	1
White, both sexes	17	-	1	-	1	-	1	1	1	1	-	1
Male	8	-	1	-	1	-	1	-	-	1	-	-
Female	9	-	-	-	-	-	-	1	1	-	-	1
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	1	-	-	-	-	-	-	-	1	1
Male	3	-	1	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	500	9	3	1	1	-	1	3	-	7	11	14
Male	254	7	2	-	1	-	1	1	-	5	6	7
Female	246	2	1	1	-	-	-	2	-	2	5	7
White, both sexes	441	6	2	-	-	-	1	2	-	6	7	11
Male	219	4	1	-	-	-	1	1	-	4	4	4
Female	222	2	1	-	-	-	-	1	-	2	3	7
Black, both sexes	18	-	1	-	-	-	-	-	-	-	1	1
Male	12	-	1	-	-	-	-	-	-	-	1	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	39	3	-	1	1	-	-	1	-	1	3	2
Male	21	3	-	-	1	-	-	-	-	1	1	2
Female	18	-	-	1	-	-	-	1	-	-	2	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
Arkansas												
All races, both sexes	488	-	1	1	-	1	1	3	5	6	8	7
Male	228	-	1	-	-	1	1	2	3	2	6	2
Female	260	-	-	1	-	-	-	1	2	4	2	5
White, both sexes	390	-	-	1	-	1	1	2	3	3	6	6
Male	185	-	-	-	-	1	1	2	2	1	4	2
Female	205	-	-	1	-	-	-	-	1	2	2	4
Black, both sexes	97	-	1	-	-	-	-	1	2	3	2	-
Male	43	-	1	-	-	-	-	-	1	1	2	-
Female	54	-	-	-	-	-	-	1	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
California												
All races, both sexes	1,079	21	11	1	1	4	11	2	10	12	23	43
Male	486	10	4	1	1	1	7	1	7	8	11	23
Female	593	11	7	-	-	3	4	1	3	4	12	20
White, both sexes	886	19	10	1	1	3	8	1	8	9	16	37
Male	392	10	4	1	1	1	6	1	5	6	7	20
Female	494	9	6	-	-	2	2	-	3	3	9	17
Black, both sexes	105	1	-	-	-	-	2	1	1	2	5	3
Male	52	-	-	-	-	-	1	-	1	2	2	-
Female	53	1	-	-	-	-	1	1	-	-	3	3
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	79	1	1	-	-	1	1	-	1	1	2	3
Male	37	-	-	-	-	-	-	-	1	-	2	3
Female	42	1	1	-	-	1	1	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
Colorado												
All races, both sexes	325	3	3	-	-	1	1	3	1	3	9	13
Male	159	2	2	-	-	-	1	1	-	2	4	8
Female	166	1	1	-	-	1	-	2	1	1	5	5
White, both sexes	306	3	2	-	-	1	1	3	-	3	8	12
Male	148	2	2	-	-	-	1	1	-	2	4	7
Female	158	1	-	-	-	1	-	2	-	1	4	5
Black, both sexes	13	-	1	-	-	-	-	-	1	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	1	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	543	3	1	1	1	1	2	2	1	6	9	13
Male	265	2	-	-	1	1	-	1	-	4	5	9
Female	278	1	1	1	-	-	2	1	1	2	4	4
White, both sexes	492	1	1	-	1	-	2	2	1	4	7	13
Male	240	1	-	-	1	-	-	1	-	3	4	9
Female	252	-	1	-	-	-	2	1	1	1	3	4
Black, both sexes	49	2	-	1	-	1	-	-	-	2	2	-
Male	24	1	-	-	-	1	-	-	-	1	1	-
Female	25	1	-	1	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
Delaware												
All races, both sexes	127	3	-	-	-	-	-	-	-	4	1	5
Male	61	3	-	-	-	-	-	-	-	1	1	3
Female	66	-	-	-	-	-	-	-	-	3	-	2
White, both sexes	107	1	-	-	-	-	-	-	-	2	1	4
Male	53	1	-	-	-	-	-	-	-	-	1	3
Female	54	-	-	-	-	-	-	-	-	2	-	1
Black, both sexes	19	2	-	-	-	-	-	-	-	2	-	-
Male	8	2	-	-	-	-	-	-	-	1	-	-
Female	11	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	109	3	-	-	-	-	-	-	-	-	4	3
Male	53	1	-	-	-	-	-	-	-	-	1	1
Female	56	2	-	-	-	-	-	-	-	-	3	2
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	95	3	-	-	-	-	-	-	-	-	4	3
Male	47	1	-	-	-	-	-	-	-	-	1	1
Female	48	2	-	-	-	-	-	-	-	-	3	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
Florida												
All races, both sexes	2,108	25	7	2	2	5	5	11	11	27	42	44
Male	1,022	13	6	1	-	3	2	6	6	11	15	24
Female	1,086	12	1	1	2	2	3	5	5	16	27	20
White, both sexes	1,816	15	7	2	2	3	3	4	5	23	28	27
Male	896	8	6	1	-	2	1	1	3	11	10	16
Female	920	7	1	1	2	1	2	3	2	12	18	11
Black, both sexes	278	9	-	-	-	2	2	7	6	3	13	16
Male	123	4	-	-	-	1	1	5	3	-	5	8
Female	155	5	-	-	-	1	1	2	3	3	8	8
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	10	1	-	-	-	-	-	-	-	1	1	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	1	1	-
Georgia												
All races, both sexes	1,351	9	4	1	1	2	7	5	15	20	28	43
Male	581	4	3	-	1	1	5	2	8	12	17	19
Female	770	5	1	1	-	1	2	3	7	8	11	24
White, both sexes	874	6	2	-	1	1	3	2	8	14	14	27
Male	360	2	1	-	1	-	2	1	4	9	8	14
Female	514	4	1	-	-	1	1	1	4	5	6	13
Black, both sexes	466	3	2	1	-	1	3	3	7	5	13	16
Male	217	2	2	-	-	1	2	1	4	3	9	5
Female	249	1	-	1	-	-	1	2	3	2	4	11
American Indian 1/, both sexes	4	-	-	-	-	-	1	-	-	1	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
Hawaii												
All races, both sexes	204	1	-	-	1	-	3	-	1	5	4	6
Male	101	1	-	-	-	-	3	-	1	2	-	3
Female	103	-	-	-	1	-	-	-	-	3	4	3
White, both sexes	41	-	-	-	1	-	-	-	-	1	2	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	1	-	-	-	-	1	2	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	161	1	-	-	-	-	3	-	1	4	2	6
Male	78	1	-	-	-	-	3	-	1	2	-	3
Female	83	-	-	-	-	-	-	-	-	2	2	3
Idaho												
All races, both sexes	66	1	2	1	-	-	-	-	-	2	-	1
Male	29	1	1	-	-	-	-	-	-	1	-	1
Female	37	-	1	1	-	-	-	-	-	1	-	-
White, both sexes	65	1	2	1	-	-	-	-	-	2	-	1
Male	28	1	1	-	-	-	-	-	-	1	-	1
Female	37	-	1	1	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
Illinois												
All races, both sexes	2,082	13	8	2	1	1	5	4	16	22	30	51
Male	919	8	3	1	1	1	2	2	5	14	12	27
Female	1,163	5	5	1	-	-	3	2	11	8	18	24
White, both sexes	1,581	8	3	1	1	1	3	2	10	17	19	30
Male	709	6	2	-	1	1	1	1	3	10	8	21
Female	872	2	1	1	-	-	2	1	7	7	11	9
Black, both sexes	474	5	5	1	-	-	2	2	5	5	11	20
Male	198	2	1	1	-	-	1	1	2	4	4	6
Female	276	3	4	-	-	-	1	1	3	1	7	14
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	27	-	-	-	-	-	-	-	1	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	1	-	-	1
Indiana												
All races, both sexes	858	9	4	-	1	5	2	2	8	5	11	22
Male	386	6	2	-	1	2	1	-	1	2	5	8
Female	472	3	2	-	-	3	1	2	7	3	6	14
White, both sexes	750	7	4	-	1	3	1	2	6	4	9	14
Male	340	5	2	-	1	2	1	-	1	2	3	5
Female	410	2	2	-	-	1	-	2	5	2	6	9
Black, both sexes	106	2	-	-	-	2	1	-	2	1	2	8
Male	46	1	-	-	-	-	-	-	-	-	2	3
Female	60	1	-	-	-	2	1	-	2	1	-	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
Iowa												
All races, both sexes	243	-	-	1	-	-	2	-	1	1	2	7
Male	109	-	-	-	-	-	1	-	-	1	1	3
Female	134	-	-	1	-	-	1	-	1	-	1	4
White, both sexes	238	-	-	1	-	-	2	-	1	1	2	7
Male	108	-	-	-	-	-	1	-	-	1	1	3
Female	130	-	-	1	-	-	1	-	1	-	1	4
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	293	4	1	-	2	-	2	2	3	-	4	7
Male	119	2	1	-	1	-	2	1	1	-	3	3
Female	174	2	-	-	1	-	-	1	2	-	1	4
White, both sexes	267	3	1	-	1	-	2	1	3	-	4	3
Male	109	1	1	-	-	-	2	-	1	-	3	3
Female	158	2	-	-	1	-	-	1	2	-	1	-
Black, both sexes	23	1	-	-	1	-	-	1	-	-	-	3
Male	9	1	-	-	1	-	-	1	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
Kentucky												
All races, both sexes	602	2	-	-	-	1	1	1	8	5	9	16
Male	287	2	-	-	-	1	1	1	7	1	3	11
Female	315	-	-	-	-	-	-	-	1	4	6	5
White, both sexes	561	2	-	-	-	-	-	-	7	3	7	14
Male	270	2	-	-	-	-	-	-	6	1	3	11
Female	291	-	-	-	-	-	-	-	1	2	4	3
Black, both sexes	41	-	-	-	-	1	1	1	1	2	2	2
Male	17	-	-	-	-	1	1	1	1	-	-	-
Female	24	-	-	-	-	-	-	-	-	2	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	877	7	3	1	3	6	1	8	7	7	17	20
Male	388	3	2	1	-	5	-	6	3	3	10	8
Female	489	4	1	-	3	1	1	2	4	4	7	12
White, both sexes	553	4	1	1	2	4	1	3	3	3	11	12
Male	236	2	-	1	-	3	-	3	-	1	7	3
Female	317	2	1	-	2	1	1	-	3	2	4	9
Black, both sexes	318	3	2	-	1	2	-	5	3	4	6	8
Male	148	1	2	-	-	2	-	3	2	2	3	5
Female	170	2	-	-	1	-	-	2	1	2	3	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
Maine												
All races, both sexes	146	2	-	-	1	-	2	-	-	1	1	1
Male	62	1	-	-	1	-	1	-	-	-	-	-
Female	84	1	-	-	-	-	1	-	-	1	1	1
White, both sexes	145	2	-	-	1	-	2	-	-	1	1	1
Male	61	1	-	-	1	-	1	-	-	-	-	-
Female	84	1	-	-	-	-	1	-	-	1	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1,095	3	2	2	1	1	3	4	7	10	21	36
Male	506	2	1	1	-	-	2	4	6	5	12	26
Female	589	1	1	1	1	1	1	-	1	5	9	10
White, both sexes	701	1	1	1	-	-	1	4	3	4	7	15
Male	325	1	1	1	-	-	1	4	2	3	4	9
Female	376	-	-	-	-	-	-	-	1	1	3	6
Black, both sexes	376	2	1	1	1	1	1	-	4	5	14	20
Male	174	1	-	-	-	-	1	-	4	1	8	17
Female	202	1	1	1	1	1	-	-	-	4	6	3
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	-	1
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	1	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
Massachusetts												
All races, both sexes	1,003	1	2	1	1	-	-	-	12	6	12	19
Male	437	-	1	-	-	-	-	-	8	1	5	11
Female	566	1	1	1	1	-	-	-	4	5	7	8
White, both sexes	930	-	1	1	1	-	-	-	8	6	10	16
Male	406	-	1	-	-	-	-	-	5	1	4	9
Female	524	-	-	1	1	-	-	-	3	5	6	7
Black, both sexes	64	1	1	-	-	-	-	-	3	-	2	3
Male	28	-	-	-	-	-	-	-	2	-	1	2
Female	36	1	1	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	903	2	2	-	1	1	1	2	5	5	17	19
Male	379	1	-	-	-	-	-	1	2	1	7	14
Female	524	1	2	-	1	1	1	1	3	4	10	5
White, both sexes	695	2	1	-	-	1	1	2	4	4	11	9
Male	281	1	-	-	-	-	-	1	1	1	3	8
Female	414	1	1	-	-	1	1	1	3	3	8	1
Black, both sexes	199	-	1	-	1	-	-	-	1	1	6	10
Male	94	-	-	-	-	-	-	-	1	-	4	6
Female	105	-	1	-	1	-	-	-	-	1	2	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
Minnesota												
All races, both sexes	302	4	1	1	-	1	-	2	2	3	4	8
Male	145	2	-	1	-	1	-	1	-	2	3	5
Female	157	2	1	-	-	-	-	1	2	1	1	3
White, both sexes	281	2	1	1	-	1	-	2	1	3	4	3
Male	136	1	-	1	-	1	-	1	-	2	3	1
Female	145	1	1	-	-	-	-	1	1	1	1	2
Black, both sexes	11	1	-	-	-	-	-	-	-	-	-	3
Male	3	-	-	-	-	-	-	-	-	-	-	2
Female	8	1	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	1	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	2
Female	3	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	525	6	3	-	1	4	4	4	1	8	9	19
Male	223	6	3	-	-	2	1	1	1	5	3	10
Female	302	-	-	-	1	2	3	3	-	3	6	9
White, both sexes	327	1	1	-	-	4	1	3	-	6	5	6
Male	143	1	1	-	-	2	-	1	-	5	3	1
Female	184	-	-	-	-	2	1	2	-	1	2	5
Black, both sexes	196	5	2	-	-	-	3	1	1	2	4	13
Male	80	5	2	-	-	-	1	-	1	-	-	9
Female	116	-	-	-	-	-	2	1	-	2	4	4
American Indian 1/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
Missouri												
All races, both sexes	713	3	1	1	2	1	1	2	7	1	7	19
Male	308	3	1	1	1	1	-	-	3	-	4	7
Female	405	-	-	-	1	-	1	2	4	1	3	12
White, both sexes	592	3	1	1	1	-	1	1	6	1	4	12
Male	253	3	1	1	1	-	-	-	3	-	3	4
Female	339	-	-	-	-	-	1	1	3	1	1	8
Black, both sexes	115	-	-	-	1	1	-	1	1	-	3	7
Male	52	-	-	-	-	1	-	-	-	-	1	3
Female	63	-	-	-	1	-	-	1	1	-	2	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	73	-	-	-	1	-	1	-	-	1	2	1
Male	32	-	-	-	-	-	-	-	-	1	1	1
Female	41	-	-	-	1	-	1	-	-	-	1	-
White, both sexes	68	-	-	-	-	-	1	-	-	1	1	1
Male	29	-	-	-	-	-	-	-	-	1	-	1
Female	39	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	1	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
Nebraska												
All races, both sexes	140	1	1	-	-	-	-	-	1	4	1	1
Male	61	-	1	-	-	-	-	-	-	3	1	1
Female	79	1	-	-	-	-	-	-	1	1	-	-
White, both sexes	131	1	1	-	-	-	-	-	1	4	1	1
Male	59	-	1	-	-	-	-	-	-	3	1	1
Female	72	1	-	-	-	-	-	-	1	1	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	416	7	-	-	2	-	-	1	4	7	6	13
Male	217	3	-	-	1	-	-	-	2	5	2	8
Female	199	4	-	-	1	-	-	1	2	2	4	5
White, both sexes	365	7	-	-	2	-	-	1	4	7	4	11
Male	194	3	-	-	1	-	-	-	2	5	2	7
Female	171	4	-	-	1	-	-	1	2	2	2	4
Black, both sexes	33	-	-	-	-	-	-	-	-	-	1	2
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
New Hampshire												
All races, both sexes	103	-	-	-	-	-	-	-	1	1	1	3
Male	56	-	-	-	-	-	-	-	1	-	-	2
Female	47	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	102	-	-	-	-	-	-	-	1	1	1	2
Male	55	-	-	-	-	-	-	-	1	-	-	1
Female	47	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1,947	17	3	3	1	2	6	4	4	12	27	31
Male	851	10	1	2	1	1	5	2	1	6	13	19
Female	1,096	7	2	1	-	1	1	2	3	6	14	12
White, both sexes	1,605	12	-	3	1	1	2	3	3	9	11	22
Male	694	8	-	2	1	-	2	1	-	5	7	13
Female	911	4	-	1	-	1	-	2	3	4	4	9
Black, both sexes	304	5	2	-	-	-	4	-	1	3	16	9
Male	137	2	-	-	-	-	3	-	1	1	6	6
Female	167	3	2	-	-	-	1	-	-	2	10	3
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	35	-	1	-	-	1	-	1	-	-	-	-
Male	20	-	1	-	-	1	-	1	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
New Mexico												
All races, both sexes	158	4	1	1	-	-	1	1	-	6	5	6
Male	75	4	1	-	-	-	-	1	-	4	3	3
Female	83	-	-	1	-	-	1	-	-	2	2	3
White, both sexes	136	3	1	-	-	-	1	-	-	4	3	5
Male	64	3	1	-	-	-	-	-	-	2	2	2
Female	72	-	-	-	-	-	1	-	-	2	1	3
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	18	1	-	1	-	-	-	1	-	2	2	-
Male	9	1	-	-	-	-	-	1	-	2	1	-
Female	9	-	-	1	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2,381	13	4	3	2	5	7	15	11	17	22	45
Male	1,100	7	2	2	2	5	3	10	6	9	14	24
Female	1,281	6	2	1	-	-	4	5	5	8	8	21
White, both sexes	2,001	8	2	3	1	5	6	12	6	11	13	32
Male	938	6	1	2	1	5	3	7	5	6	9	17
Female	1,063	2	1	1	-	-	3	5	1	5	4	15
Black, both sexes	341	5	1	-	1	-	1	3	5	6	9	13
Male	141	1	1	-	1	-	-	3	1	3	5	7
Female	200	4	-	-	-	-	1	-	4	3	4	6
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	37	-	1	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
North Carolina												
All races, both sexes	1,256	18	2	3	-	2	4	3	13	10	26	33
Male	562	10	1	2	-	1	3	1	7	6	13	15
Female	694	8	1	1	-	1	1	2	6	4	13	18
White, both sexes	870	7	2	2	-	-	1	1	7	7	8	16
Male	376	4	1	1	-	-	1	1	4	4	5	7
Female	494	3	1	1	-	-	-	-	3	3	3	9
Black, both sexes	368	11	-	1	-	2	3	2	5	2	17	16
Male	178	6	-	1	-	1	2	-	3	1	8	7
Female	190	5	-	-	-	1	1	2	2	1	9	9
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	1	1	-	1
Male	6	-	-	-	-	-	-	-	-	1	-	1
Female	9	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
North Dakota												
All races, both sexes	35	-	-	-	-	-	-	-	1	-	-	2
Male	16	-	-	-	-	-	-	-	1	-	-	1
Female	19	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	33	-	-	-	-	-	-	-	-	-	-	2
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
Ohio												
All races, both sexes	1,573	15	4	-	4	2	2	6	8	17	32	47
Male	699	10	1	-	2	1	1	6	4	9	16	23
Female	874	5	3	-	2	1	1	-	4	8	16	24
White, both sexes	1,298	8	4	-	3	1	1	5	7	11	25	34
Male	568	5	1	-	1	-	1	5	3	7	14	17
Female	730	3	3	-	2	1	-	-	4	4	11	17
Black, both sexes	268	6	-	-	1	1	1	1	1	6	7	13
Male	129	4	-	-	1	1	-	1	1	2	2	6
Female	139	2	-	-	-	-	1	-	-	4	5	7
American Indian 1/, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	435	6	2	1	-	1	-	2	7	5	10	17
Male	201	2	1	-	-	-	-	1	4	3	4	7
Female	234	4	1	1	-	1	-	1	3	2	6	10
White, both sexes	368	5	-	1	-	1	-	2	3	3	4	13
Male	165	1	-	-	-	-	-	1	2	1	1	6
Female	203	4	-	1	-	1	-	1	1	2	3	7
Black, both sexes	44	-	1	-	-	-	-	-	-	2	5	1
Male	21	-	-	-	-	-	-	-	-	2	2	-
Female	23	-	1	-	-	-	-	-	-	-	3	1
American Indian 1/, both sexes	22	1	1	-	-	-	-	-	4	-	1	3
Male	14	1	1	-	-	-	-	-	2	-	1	1
Female	8	-	-	-	-	-	-	-	2	-	-	2
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
Oregon												
All races, both sexes	226	-	-	-	1	-	2	3	-	4	4	10
Male	107	-	-	-	-	-	1	1	-	2	2	6
Female	119	-	-	-	1	-	1	2	-	2	2	4
White, both sexes	213	-	-	-	1	-	2	2	-	4	3	9
Male	101	-	-	-	-	-	1	1	-	2	1	6
Female	112	-	-	-	1	-	1	1	-	2	2	3
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	1	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	1	-	-	-	1
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	2,593	14	2	4	1	4	3	3	9	13	34	71
Male	1,162	9	1	1	1	2	2	-	4	4	19	39
Female	1,431	5	1	3	-	2	1	3	5	9	15	32
White, both sexes	2,186	11	1	4	1	4	2	2	3	9	23	45
Male	969	6	1	1	1	2	1	-	3	4	13	21
Female	1,217	5	-	3	-	2	1	2	-	5	10	24
Black, both sexes	384	3	1	-	-	-	1	1	6	4	10	25
Male	181	3	-	-	-	-	1	-	1	-	5	18
Female	203	-	1	-	-	-	-	1	5	4	5	7
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	-	-	1	1
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
Rhode Island												
All races, both sexes	95	-	-	-	1	1	1	-	-	-	1	1
Male	43	-	-	-	-	1	1	-	-	-	-	-
Female	52	-	-	-	1	-	-	-	-	-	1	1
White, both sexes	88	-	-	-	1	1	-	-	-	-	1	1
Male	39	-	-	-	-	1	-	-	-	-	-	-
Female	49	-	-	-	1	-	-	-	-	-	1	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	654	7	1	1	-	2	5	2	-	4	6	23
Male	276	4	-	-	-	-	1	1	-	2	4	11
Female	378	3	1	1	-	2	4	1	-	2	2	12
White, both sexes	414	2	1	-	-	1	3	2	-	2	2	5
Male	170	2	-	-	-	-	-	1	-	1	-	4
Female	244	-	1	-	-	1	3	1	-	1	2	1
Black, both sexes	233	3	-	1	-	1	2	-	-	2	4	17
Male	103	1	-	-	-	-	1	-	-	1	4	7
Female	130	2	-	1	-	1	1	-	-	1	-	10
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	6	2	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	3	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
South Dakota												
All races, both sexes	44	-	1	2	-	-	-	-	-	1	1	3
Male	21	-	-	2	-	-	-	-	-	-	-	3
Female	23	-	1	-	-	-	-	-	-	1	1	-
White, both sexes	37	-	1	2	-	-	-	-	-	-	-	2
Male	18	-	-	2	-	-	-	-	-	-	-	2
Female	19	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	1	1	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	588	8	3	-	-	1	2	4	2	7	13	28
Male	251	4	2	-	-	-	1	1	2	7	6	13
Female	337	4	1	-	-	1	1	3	-	-	7	15
White, both sexes	462	3	3	-	-	-	1	2	2	4	8	17
Male	195	1	2	-	-	-	1	1	2	4	3	7
Female	267	2	1	-	-	-	-	1	-	-	5	10
Black, both sexes	125	5	-	-	-	1	1	2	-	3	5	10
Male	55	3	-	-	-	-	-	-	-	3	3	5
Female	70	2	-	-	-	1	1	2	-	-	2	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
Texas												
All races, both sexes	2,516	39	8	10	1	4	7	16	24	30	56	89
Male	1,167	28	4	7	-	3	3	3	14	18	31	57
Female	1,349	11	4	3	1	1	4	13	10	12	25	32
White, both sexes	2,066	30	7	9	-	4	3	14	16	23	39	65
Male	957	20	3	6	-	3	2	2	9	13	23	40
Female	1,109	10	4	3	-	1	1	12	7	10	16	25
Black, both sexes	422	9	1	-	1	-	4	2	8	6	16	23
Male	196	8	1	-	-	-	1	1	5	4	8	16
Female	226	1	-	-	1	-	3	1	3	2	8	7
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	25	-	-	1	-	-	-	-	-	1	-	1
Male	14	-	-	1	-	-	-	-	-	1	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	120	-	3	-	-	-	-	2	1	1	-	6
Male	50	-	1	-	-	-	-	1	-	-	-	2
Female	70	-	2	-	-	-	-	1	1	1	-	4
White, both sexes	114	-	3	-	-	-	-	2	-	1	-	5
Male	46	-	1	-	-	-	-	1	-	-	-	2
Female	68	-	2	-	-	-	-	1	-	1	-	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	1	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
Vermont												
All races, both sexes	38	-	1	-	-	-	-	-	-	-	-	1
Male	17	-	1	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	38	-	1	-	-	-	-	-	-	-	-	1
Male	17	-	1	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1,234	10	7	-	-	1	6	5	4	4	23	29
Male	569	7	1	-	-	-	4	2	3	2	14	21
Female	665	3	6	-	-	1	2	3	1	2	9	8
White, both sexes	848	4	7	-	-	1	3	1	2	2	14	15
Male	397	3	1	-	-	-	2	1	2	1	8	10
Female	451	1	6	-	-	1	1	-	-	1	6	5
Black, both sexes	367	6	-	-	-	-	3	4	2	2	9	12
Male	162	4	-	-	-	-	2	1	1	1	6	9
Female	205	2	-	-	-	-	1	3	1	1	3	3
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	-	2
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
Washington												
All races, both sexes	340	1	2	1	-	2	1	1	5	2	7	11
Male	147	1	1	-	-	2	-	1	4	1	3	6
Female	193	-	1	1	-	-	1	-	1	1	4	5
White, both sexes	303	-	1	1	-	2	-	1	1	2	7	9
Male	126	-	-	-	-	2	-	1	1	1	3	4
Female	177	-	1	1	-	-	-	-	-	1	4	5
Black, both sexes	10	1	1	-	-	-	-	-	1	-	-	1
Male	9	1	1	-	-	-	-	-	1	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	3	-	-	1
Male	11	-	-	-	-	-	-	-	2	-	-	1
Female	9	-	-	-	-	-	-	-	1	-	-	-
West Virginia												
All races, both sexes	334	1	-	-	-	1	1	-	2	4	4	11
Male	160	-	-	-	-	-	-	-	2	3	2	4
Female	174	1	-	-	-	1	1	-	-	1	2	7
White, both sexes	324	1	-	-	-	1	1	-	2	4	3	10
Male	156	-	-	-	-	-	-	-	2	3	1	4
Female	168	1	-	-	-	1	1	-	-	1	2	6
Black, both sexes	10	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
Wisconsin												
All races, both sexes	443	4	1	2	-	-	1	-	4	3	4	12
Male	202	3	1	1	-	-	-	-	4	2	4	10
Female	241	1	-	1	-	-	1	-	-	1	-	2
White, both sexes	407	3	1	2	-	-	1	-	3	1	3	8
Male	182	2	1	1	-	-	-	-	3	-	3	7
Female	225	1	-	1	-	-	1	-	-	1	-	1
Black, both sexes	25	1	-	-	-	-	-	-	1	1	1	2
Male	13	1	-	-	-	-	-	-	1	1	1	1
Female	12	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	1	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	45	1	-	-	-	1	-	-	-	-	-	1
Male	22	-	-	-	-	-	-	-	-	-	-	1
Female	23	1	-	-	-	1	-	-	-	-	-	-
White, both sexes	44	1	-	-	-	1	-	-	-	-	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	1
Female	23	1	-	-	-	1	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
United States												
All races, both sexes	1,426	1,813	2,163	2,593	3,668	5,062	6,045	5,140	3,058	1,090	177	3
Male	761	975	1,081	1,320	1,861	2,414	2,687	2,009	913	280	35	1
Female	665	838	1,082	1,273	1,807	2,648	3,358	3,131	2,145	810	142	2
White, both sexes	981	1,328	1,584	1,959	2,915	4,183	5,124	4,443	2,644	935	131	2
Male	540	716	792	1,011	1,486	2,023	2,277	1,767	786	244	26	1
Female	441	612	792	948	1,429	2,160	2,847	2,676	1,858	691	105	1
Black, both sexes	405	447	509	576	674	793	819	608	375	139	44	1
Male	204	241	250	280	334	353	354	207	108	31	9	-
Female	201	206	259	296	340	440	465	401	267	108	35	1
American Indian 1/, both sexes	17	13	22	14	14	20	14	18	5	3	-	-
Male	8	3	12	7	6	10	6	8	2	-	-	-
Female	9	10	10	7	8	10	8	10	3	3	-	-
Asian or Pacific Islander 2/, both sexes	23	25	48	44	65	66	88	71	34	13	2	-
Male	9	15	27	22	35	28	50	27	17	5	-	-
Female	14	10	21	22	30	38	38	44	17	8	2	-
Alabama												
All races, both sexes	37	49	53	71	94	108	131	105	44	22	6	-
Male	20	27	20	31	39	58	35	31	9	5	1	-
Female	17	22	33	40	55	50	96	74	35	17	5	-
White, both sexes	20	36	35	49	68	74	95	83	28	17	5	-
Male	12	22	14	23	31	37	26	25	6	3	-	-
Female	8	14	21	26	37	37	69	58	22	14	5	-
Black, both sexes	17	13	17	22	26	34	36	22	16	5	1	-
Male	8	5	5	8	8	21	9	6	3	2	1	-
Female	9	8	12	14	18	13	27	16	13	3	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
Alaska												
All races, both sexes	2	1	2	3	-	-	3	1	2	-	-	-
Male	1	1	-	2	-	-	1	-	1	-	-	-
Female	1	-	2	1	-	-	2	1	1	-	-	-
White, both sexes	1	1	1	2	-	-	3	1	1	-	-	-
Male	1	1	-	1	-	-	1	-	-	-	-	-
Female	-	-	1	1	-	-	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	29	31	39	37	74	75	68	61	30	4	2	-
Male	18	18	19	19	39	31	34	28	16	2	-	-
Female	11	13	20	18	35	44	34	33	14	2	2	-
White, both sexes	22	27	34	32	71	68	64	55	27	4	2	-
Male	14	15	16	16	39	28	30	25	15	2	-	-
Female	8	12	18	16	32	40	34	30	12	2	2	-
Black, both sexes	3	2	1	2	1	2	1	2	1	-	-	-
Male	2	1	1	2	-	1	1	1	-	-	-	-
Female	1	1	-	-	1	1	-	1	1	-	-	-
American Indian 1/, both sexes	3	2	4	3	2	4	3	4	2	-	-	-
Male	1	2	2	1	-	1	3	2	1	-	-	-
Female	2	-	2	2	2	3	-	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
Arkansas												
All races, both sexes	21	22	32	31	46	75	82	66	53	21	6	-
Male	15	16	17	17	23	26	35	30	25	5	1	-
Female	6	6	15	14	23	49	47	36	28	16	5	-
White, both sexes	14	17	24	25	40	62	69	57	39	16	4	-
Male	9	12	13	15	19	20	33	26	19	5	1	-
Female	5	5	11	10	21	42	36	31	20	11	3	-
Black, both sexes	7	5	8	6	6	13	13	9	14	5	2	-
Male	6	4	4	2	4	6	2	4	6	-	-	-
Female	1	1	4	4	2	7	11	5	8	5	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	48	62	78	80	84	123	192	149	86	34	4	-
Male	24	30	39	43	41	59	75	69	22	10	-	-
Female	24	32	39	37	43	64	117	80	64	24	4	-
White, both sexes	36	50	60	68	75	98	161	117	77	29	2	-
Male	17	22	33	37	36	45	61	54	18	7	-	-
Female	19	28	27	31	39	53	100	63	59	22	2	-
Black, both sexes	9	8	8	5	4	13	16	20	4	2	1	-
Male	5	5	3	3	3	8	8	9	-	2	-	-
Female	4	3	5	2	1	5	8	11	4	-	1	-
American Indian 1/, both sexes	1	1	1	-	-	2	3	-	1	-	-	-
Male	1	1	-	-	-	2	1	-	-	-	-	-
Female	-	-	1	-	-	-	2	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	9	7	5	10	12	12	4	3	1	-
Male	1	2	3	3	2	4	5	6	4	1	-	-
Female	1	1	6	4	3	6	7	6	-	2	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
Colorado												
All races, both sexes	8	18	20	20	26	41	58	62	25	9	1	-
Male	5	11	10	10	13	19	28	30	6	6	1	-
Female	3	7	10	10	13	22	30	32	19	3	-	-
White, both sexes	7	17	19	18	26	41	52	58	25	9	1	-
Male	5	10	9	8	13	19	23	29	6	6	1	-
Female	2	7	10	10	13	22	29	29	19	3	-	-
Black, both sexes	-	1	1	2	-	-	2	3	-	-	-	-
Male	-	1	1	2	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	4	-	-	-	-	-
Male	-	-	-	-	-	-	3	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Connecticut												
All races, both sexes	18	18	15	36	44	82	111	98	61	18	2	-
Male	11	12	5	22	27	49	46	47	16	5	2	-
Female	7	6	10	14	17	33	65	51	45	13	-	-
White, both sexes	13	14	13	30	39	72	107	94	58	18	2	-
Male	8	10	5	18	24	44	45	44	16	5	2	-
Female	5	4	8	12	15	28	62	50	42	13	-	-
Black, both sexes	5	4	1	5	5	10	4	4	3	-	-	-
Male	3	2	-	3	3	5	1	3	-	-	-	-
Female	2	2	1	2	2	5	3	1	3	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
Delaware												
All races, both sexes	4	7	4	7	15	21	25	19	10	2	-	-
Male	2	3	3	4	8	11	13	5	3	1	-	-
Female	2	4	1	3	7	10	12	14	7	1	-	-
White, both sexes	3	6	4	6	14	17	23	17	8	1	-	-
Male	1	3	3	3	8	9	13	5	2	1	-	-
Female	2	3	1	3	6	8	10	12	6	-	-	-
Black, both sexes	1	1	-	1	1	4	2	2	2	1	-	-
Male	1	-	-	1	-	2	-	-	1	-	-	-
Female	-	1	-	-	1	2	2	2	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	7	4	9	11	13	11	22	13	8	1	-	-
Male	5	3	6	7	4	3	14	6	2	-	-	-
Female	2	1	3	4	9	8	8	7	6	1	-	-
White, both sexes	-	-	2	-	-	1	6	1	3	-	-	-
Male	-	-	2	-	-	-	4	-	-	-	-	-
Female	-	-	-	-	-	1	2	1	3	-	-	-
Black, both sexes	7	4	7	11	12	10	16	12	5	1	-	-
Male	5	3	4	7	4	3	10	6	2	-	-	-
Female	2	1	3	4	8	7	6	6	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
Florida												
All races, both sexes	69	96	126	155	214	324	354	304	203	70	12	-
Male	38	55	73	88	107	173	176	131	70	20	4	-
Female	31	41	53	67	107	151	178	173	133	50	8	-
White, both sexes	49	78	100	122	186	293	331	275	188	66	9	-
Male	27	48	59	73	90	162	168	121	67	18	4	-
Female	22	30	41	49	96	131	163	154	121	48	5	-
Black, both sexes	19	18	26	32	27	29	21	26	15	4	3	-
Male	11	7	14	15	16	10	8	10	3	2	-	-
Female	8	11	12	17	11	19	13	16	12	2	3	-
American Indian 1/, both sexes	1	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	2	2	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	2	2	-	-	-	-
Georgia												
All races, both sexes	68	78	96	120	145	209	192	161	98	35	14	-
Male	33	40	42	63	80	90	68	58	23	11	1	-
Female	35	38	54	57	65	119	124	103	75	24	13	-
White, both sexes	30	48	52	77	85	149	141	118	63	26	7	-
Male	14	26	21	41	48	59	48	42	12	7	-	-
Female	16	22	31	36	37	90	93	76	51	19	7	-
Black, both sexes	38	29	44	41	59	58	50	42	35	9	7	-
Male	19	13	21	22	31	30	20	16	11	4	1	-
Female	19	16	23	19	28	28	30	26	24	5	6	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	1	2	-	1	-	-	-	-
Male	-	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
Hawaii												
All races, both sexes	12	8	14	13	26	24	32	26	21	7	-	-
Male	4	4	8	6	15	9	20	10	12	3	-	-
Female	8	4	6	7	11	15	12	16	9	4	-	-
White, both sexes	1	3	1	5	3	6	8	5	5	-	-	-
Male	1	2	1	1	2	3	4	4	4	-	-	-
Female	-	1	-	4	1	3	4	1	1	-	-	-
Black, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	5	13	7	23	18	23	21	16	7	-	-
Male	3	2	7	5	13	6	15	6	8	3	-	-
Female	8	3	6	2	10	12	8	15	8	4	-	-
Idaho												
All races, both sexes	2	1	3	6	8	12	9	6	8	4	-	-
Male	2	1	-	3	2	7	5	2	1	2	-	-
Female	-	-	3	3	6	5	4	4	7	2	-	-
White, both sexes	2	1	3	6	8	11	9	6	8	4	-	-
Male	2	1	-	3	2	6	5	2	1	2	-	-
Female	-	-	3	3	6	5	4	4	7	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
Illinois												
All races, both sexes	80	96	129	164	212	287	371	324	191	65	10	-
Male	46	54	61	75	107	138	170	128	57	7	-	-
Female	34	42	68	89	105	149	201	196	134	58	10	-
White, both sexes	52	63	81	111	160	232	293	272	162	52	8	-
Male	32	36	37	51	79	113	138	113	49	7	-	-
Female	20	27	44	60	81	119	155	159	113	45	8	-
Black, both sexes	27	31	45	49	50	52	75	49	26	12	2	-
Male	13	17	21	22	28	24	29	14	8	-	-	-
Female	14	14	24	27	22	28	46	35	18	12	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	3	4	2	3	3	3	3	1	-	-
Male	1	1	3	2	-	1	3	1	-	-	-	-
Female	-	1	-	2	2	2	-	2	3	1	-	-
Indiana												
All races, both sexes	35	54	64	62	103	117	145	126	59	21	3	-
Male	25	31	31	30	56	47	67	42	22	6	1	-
Female	10	23	33	32	47	70	78	84	37	15	2	-
White, both sexes	29	47	56	54	91	98	131	117	54	19	3	-
Male	20	27	27	26	51	39	59	40	22	6	1	-
Female	9	20	29	28	40	59	72	77	32	13	2	-
Black, both sexes	6	7	8	8	10	19	14	9	5	2	-	-
Male	5	4	4	4	5	8	8	2	-	-	-	-
Female	1	3	4	4	5	11	6	7	5	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
Iowa												
All races, both sexes	9	10	13	18	26	47	40	38	18	9	1	-
Male	6	6	5	6	14	24	21	17	-	3	1	-
Female	3	4	8	12	12	23	19	21	18	6	-	-
White, both sexes	9	10	12	17	26	45	40	38	18	8	1	-
Male	6	6	5	6	14	24	21	17	-	2	1	-
Female	3	4	7	11	12	21	19	21	18	6	-	-
Black, both sexes	-	-	-	1	-	2	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	1	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	10	10	20	23	28	39	54	47	22	12	3	-
Male	4	5	14	10	15	18	22	10	4	2	1	-
Female	6	5	6	13	13	21	32	37	18	10	2	-
White, both sexes	9	8	19	19	25	37	49	46	22	12	3	-
Male	3	5	14	8	14	18	20	9	4	2	1	-
Female	6	3	5	11	11	19	29	37	18	10	2	-
Black, both sexes	1	2	1	4	2	2	5	-	-	-	-	-
Male	1	-	-	2	1	-	2	-	-	-	-	-
Female	-	2	1	2	1	2	3	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
Kentucky												
All races, both sexes	33	36	44	46	66	101	98	79	46	9	1	-
Male	23	18	22	24	30	52	46	30	13	2	-	-
Female	10	18	22	22	36	49	52	49	33	7	1	-
White, both sexes	31	34	40	42	60	98	95	75	45	8	-	-
Male	21	18	21	22	28	51	45	26	13	2	-	-
Female	10	16	19	20	32	47	50	49	32	6	-	-
Black, both sexes	2	2	4	4	6	3	3	4	1	1	1	-
Male	2	-	1	2	2	1	1	4	-	-	-	-
Female	-	2	3	2	4	2	2	-	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	40	51	62	75	87	111	148	112	81	25	5	-
Male	16	25	30	42	42	52	69	39	22	7	3	-
Female	24	26	32	33	45	59	79	73	59	18	2	-
White, both sexes	24	28	30	47	52	69	96	82	60	16	4	-
Male	11	13	10	25	26	33	45	30	15	5	3	-
Female	13	15	20	22	26	36	51	52	45	11	1	-
Black, both sexes	16	22	32	26	34	42	52	29	21	9	1	-
Male	5	12	20	15	15	19	24	9	7	2	-	-
Female	11	10	12	11	19	23	28	20	14	7	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
Maine												
All races, both sexes	8	7	6	8	18	18	28	21	19	4	1	-
Male	2	4	4	4	9	8	16	6	5	1	-	-
Female	6	3	2	4	9	10	12	15	14	3	1	-
White, both sexes	8	7	6	8	18	17	28	21	19	4	1	-
Male	2	4	4	4	9	7	16	6	5	1	-	-
Female	6	3	2	4	9	10	12	15	14	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	53	60	72	89	125	141	169	165	89	36	6	-
Male	30	33	32	45	67	66	74	67	23	10	-	-
Female	23	27	40	44	58	75	95	98	66	26	6	-
White, both sexes	24	27	45	55	79	94	122	121	64	28	5	-
Male	14	15	21	30	40	50	52	52	17	8	-	-
Female	10	12	24	25	39	44	70	69	47	20	5	-
Black, both sexes	28	31	26	33	45	43	45	41	25	8	1	-
Male	15	18	11	15	26	14	21	14	6	2	-	-
Female	13	13	15	18	19	29	24	27	19	6	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	1	1	3	2	3	-	-	-	-
Male	1	-	-	-	1	2	1	1	-	-	-	-
Female	-	2	1	1	-	1	1	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
Massachusetts												
All races, both sexes	34	51	57	65	112	122	189	168	105	41	5	-
Male	20	27	28	35	64	56	79	58	30	13	1	-
Female	14	24	29	30	48	66	110	110	75	28	4	-
White, both sexes	33	48	48	57	97	115	182	163	100	39	5	-
Male	20	26	24	33	57	51	75	58	28	13	1	-
Female	13	22	24	24	40	64	107	105	72	26	4	-
Black, both sexes	1	3	9	8	13	7	5	3	3	2	-	-
Male	-	1	4	2	7	5	3	-	1	-	-	-
Female	1	2	5	6	6	2	2	3	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	2	2	2	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	2	-	1	2	1	-	-	-
Michigan												
All races, both sexes	38	39	71	46	77	134	158	162	93	28	2	-
Male	16	22	40	21	37	58	68	62	21	7	1	-
Female	22	17	31	25	40	76	90	100	72	21	1	-
White, both sexes	26	22	45	34	59	106	124	138	80	24	2	-
Male	11	12	24	17	28	42	51	56	17	7	1	-
Female	15	10	21	17	31	64	73	82	63	17	1	-
Black, both sexes	12	16	23	12	17	27	32	23	13	4	-	-
Male	5	9	15	4	9	16	16	5	4	-	-	-
Female	7	7	8	8	8	11	16	18	9	4	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	-	1	1	1	1	-	-	-	-
Male	-	1	-	-	-	-	1	1	-	-	-	-
Female	-	-	2	-	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
Minnesota												
All races, both sexes	16	15	10	21	34	35	55	58	23	9	-	-
Male	8	8	5	8	16	21	28	30	6	-	-	-
Female	8	7	5	13	18	14	27	28	17	9	-	-
White, both sexes	13	13	10	18	33	34	55	55	23	9	-	-
Male	8	8	5	8	15	20	28	28	6	-	-	-
Female	5	5	5	10	18	14	27	27	17	9	-	-
Black, both sexes	2	1	-	2	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	2	1	-	2	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	29	33	38	41	53	70	63	64	51	21	3	-
Male	13	15	15	25	20	34	29	24	11	3	2	-
Female	16	18	23	16	33	36	34	40	40	18	1	-
White, both sexes	13	17	23	18	37	52	38	44	41	16	1	-
Male	8	5	10	13	16	26	19	19	9	3	1	-
Female	5	12	13	5	21	26	19	25	32	13	-	-
Black, both sexes	16	16	15	22	16	18	25	20	10	5	2	-
Male	5	10	5	12	4	8	10	5	2	-	1	-
Female	11	6	10	10	12	10	15	15	8	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
Missouri												
All races, both sexes	32	47	36	53	60	94	133	101	86	24	2	-
Male	16	30	15	29	25	40	57	42	25	8	-	-
Female	16	17	21	24	35	54	76	59	61	16	2	-
White, both sexes	21	33	27	46	47	83	116	89	77	20	2	-
Male	10	20	11	25	19	36	47	37	25	7	-	-
Female	11	13	16	21	28	47	69	52	52	13	2	-
Black, both sexes	10	11	9	7	12	10	17	12	9	4	-	-
Male	6	8	4	4	5	4	10	5	-	1	-	-
Female	4	3	5	3	7	6	7	7	9	3	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	-	-	-	1	-	-	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	1	-	-	-	-	-	-
Montana												
All races, both sexes	2	3	6	4	4	6	15	9	14	3	1	-
Male	1	1	-	3	2	6	9	5	2	-	-	-
Female	1	2	6	1	2	-	6	4	12	3	1	-
White, both sexes	1	3	6	3	4	6	15	9	13	3	1	-
Male	-	1	-	2	2	6	9	5	2	-	-	-
Female	1	2	6	1	2	-	6	4	11	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	1	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
Nebraska												
All races, both sexes	6	5	6	5	13	21	35	26	9	4	1	-
Male	4	3	3	2	10	6	16	5	4	2	-	-
Female	2	2	3	3	3	15	19	21	5	2	1	-
White, both sexes	4	3	5	5	13	20	32	26	9	4	1	-
Male	3	2	3	2	10	6	16	5	4	2	-	-
Female	1	1	2	3	3	14	16	21	5	2	1	-
Black, both sexes	1	2	1	-	-	1	3	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	1	1	-	-	1	3	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	18	37	29	40	62	70	60	39	16	4	-	1
Male	10	19	18	25	38	30	33	19	2	1	-	1
Female	8	18	11	15	24	40	27	20	14	3	-	-
White, both sexes	17	30	21	35	51	64	53	38	15	4	-	1
Male	10	19	12	21	32	28	29	19	2	1	-	1
Female	7	11	9	14	19	36	24	19	13	3	-	-
Black, both sexes	1	4	4	4	5	4	6	1	1	-	-	-
Male	-	-	2	3	2	2	3	-	-	-	-	-
Female	1	4	2	1	3	2	3	1	1	-	-	-
American Indian 1/, both sexes	-	3	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	3	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	3	1	6	1	1	-	-	-	-	-
Male	-	-	3	1	4	-	1	-	-	-	-	-
Female	-	-	-	-	2	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
New Hampshire												
All races, both sexes	4	7	3	10	6	15	20	16	12	4	-	-
Male	2	2	2	6	3	8	14	5	10	1	-	-
Female	2	5	1	4	3	7	6	11	2	3	-	-
White, both sexes	4	7	3	10	6	15	20	16	12	4	-	-
Male	2	2	2	6	3	8	14	5	10	1	-	-
Female	2	5	1	4	3	7	6	11	2	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	58	73	92	114	178	305	368	355	198	85	11	-
Male	34	40	44	58	94	158	157	125	58	20	2	-
Female	24	33	48	56	84	147	211	230	140	65	9	-
White, both sexes	40	46	65	85	140	242	326	323	187	77	7	-
Male	24	24	30	44	75	127	145	113	55	18	-	-
Female	16	22	35	41	65	115	181	210	132	59	7	-
Black, both sexes	17	22	24	27	31	56	36	29	11	7	4	-
Male	10	11	11	14	17	27	11	10	3	2	2	-
Female	7	11	13	13	14	29	25	19	8	5	2	-
American Indian 1/, both sexes	-	-	-	-	2	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	5	3	2	5	7	6	3	-	-	-	-
Male	-	5	3	-	2	4	1	2	-	-	-	-
Female	1	-	-	2	3	3	5	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
New Mexico												
All races, both sexes	12	5	11	9	18	18	27	19	9	4	1	-
Male	7	3	6	6	7	10	12	4	3	1	-	-
Female	5	2	5	3	11	8	15	15	6	3	1	-
White, both sexes	10	5	7	9	16	18	25	16	9	3	1	-
Male	6	3	4	6	5	10	12	4	3	1	-	-
Female	4	2	3	3	11	8	13	12	6	2	1	-
Black, both sexes	1	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	-	4	-	1	-	1	3	-	1	-	-
Male	1	-	2	-	1	-	-	-	-	-	-	-
Female	-	-	2	-	-	-	1	3	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	73	118	118	174	244	363	424	386	234	85	17	1
Male	41	56	59	90	132	182	195	165	74	18	4	-
Female	32	62	59	84	112	181	229	221	160	67	13	1
White, both sexes	55	89	94	135	204	309	368	340	215	79	13	1
Male	33	43	49	75	111	161	171	146	67	17	3	-
Female	22	46	45	60	93	148	197	194	148	62	10	1
Black, both sexes	17	27	21	33	38	51	45	38	17	6	4	-
Male	7	13	8	12	19	20	18	15	5	1	1	-
Female	10	14	13	21	19	31	27	23	12	5	3	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	3	5	2	3	11	8	2	-	-	-
Male	1	-	2	2	2	1	6	4	2	-	-	-
Female	-	1	1	3	-	2	5	4	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
North Carolina												
All races, both sexes	53	64	92	95	143	180	213	159	93	36	14	-
Male	25	34	51	49	58	85	92	64	30	9	6	-
Female	28	30	41	46	85	95	121	95	63	27	8	-
White, both sexes	31	38	58	57	108	145	153	124	72	24	9	-
Male	12	21	32	29	43	73	59	46	24	5	4	-
Female	19	17	26	28	65	72	94	78	48	19	5	-
Black, both sexes	20	26	32	36	33	34	58	32	21	12	5	-
Male	12	13	19	20	13	12	33	15	6	4	2	-
Female	8	13	13	16	20	22	25	17	15	8	3	-
American Indian 1/, both sexes	2	-	2	2	1	1	2	2	-	-	-	-
Male	1	-	-	-	1	-	-	2	-	-	-	-
Female	1	-	2	2	-	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	2	1	2	4	10	5	2	4	2	-	-
Male	-	2	-	-	1	7	2	-	2	-	-	-
Female	-	-	1	2	3	3	3	2	2	2	-	-
White, both sexes	-	2	1	2	4	10	4	2	4	2	-	-
Male	-	2	-	-	1	7	1	-	2	-	-	-
Female	-	-	1	2	3	3	3	2	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
Ohio												
All races, both sexes	60	81	108	129	159	239	265	215	145	33	2	-
Male	23	47	59	52	84	118	113	83	38	8	1	-
Female	37	34	49	77	75	121	152	132	107	25	1	-
White, both sexes	40	58	85	101	128	204	234	191	128	28	2	-
Male	11	34	44	44	68	105	95	75	30	7	1	-
Female	29	24	41	57	60	99	139	116	98	21	1	-
Black, both sexes	20	23	23	27	30	34	29	23	17	5	-	-
Male	12	13	15	8	16	13	17	8	8	1	-	-
Female	8	10	8	19	14	21	12	15	9	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	1	-	1	1	-	-	-	-
Oklahoma												
All races, both sexes	9	27	42	38	55	55	59	52	30	14	3	-
Male	5	17	24	15	35	23	31	19	7	3	-	-
Female	4	10	18	23	20	32	28	33	23	11	3	-
White, both sexes	6	24	37	33	46	47	51	48	28	13	3	-
Male	3	16	20	13	28	18	29	18	5	3	-	-
Female	3	8	17	20	18	29	22	30	23	10	3	-
Black, both sexes	1	3	2	4	8	4	7	3	2	1	-	-
Male	1	1	1	2	6	2	2	-	2	-	-	-
Female	-	2	1	2	2	2	5	3	-	1	-	-
American Indian 1/, both sexes	2	-	3	1	1	3	1	1	-	-	-	-
Male	1	-	3	-	1	2	-	1	-	-	-	-
Female	1	-	-	1	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
Oregon												
All races, both sexes	13	9	21	18	15	23	41	33	19	10	-	-
Male	8	3	12	7	9	8	20	17	7	4	-	-
Female	5	6	9	11	6	15	21	16	12	6	-	-
White, both sexes	12	9	20	17	15	21	37	32	19	10	-	-
Male	8	3	11	6	9	8	17	17	7	4	-	-
Female	4	6	9	11	6	13	20	15	12	6	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	-	1	1	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	3	1	-	-	-	-
Male	-	-	-	1	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Pennsylvania												
All races, both sexes	83	113	137	164	281	398	516	413	231	90	9	-
Male	51	60	72	85	133	185	238	166	73	17	-	-
Female	32	53	65	79	148	213	278	247	158	73	9	-
White, both sexes	57	82	101	126	234	337	469	377	207	83	8	-
Male	35	38	53	65	112	154	224	157	64	14	-	-
Female	22	44	48	61	122	183	245	220	143	69	8	-
Black, both sexes	24	30	33	33	43	58	47	34	23	7	1	-
Male	15	21	18	18	18	29	14	9	8	3	-	-
Female	9	9	15	15	25	29	33	25	15	4	1	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	3	5	3	3	-	2	1	-	-	-
Male	1	1	1	2	3	2	-	-	1	-	-	-
Female	1	-	2	3	-	1	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
Rhode Island												
All races, both sexes	1	8	1	6	9	13	11	23	15	3	-	-
Male	-	5	-	5	4	8	6	10	2	1	-	-
Female	1	3	1	1	5	5	5	13	13	2	-	-
White, both sexes	1	7	1	5	9	11	10	22	15	3	-	-
Male	-	4	-	4	4	7	6	10	2	1	-	-
Female	1	3	1	1	5	4	4	12	13	2	-	-
Black, both sexes	-	1	-	-	-	1	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	1	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
South Carolina												
All races, both sexes	33	47	52	66	71	90	108	81	33	18	4	-
Male	15	23	25	35	37	36	43	29	9	1	-	-
Female	18	24	27	31	34	54	65	52	24	17	4	-
White, both sexes	21	29	29	41	50	60	68	63	23	11	1	-
Male	11	12	15	22	29	24	23	21	5	-	-	-
Female	10	17	14	19	21	36	45	42	18	11	1	-
Black, both sexes	12	18	22	25	20	29	40	17	10	7	3	-
Male	4	11	9	13	8	12	20	7	4	1	-	-
Female	8	7	13	12	12	17	20	10	6	6	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	1	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
South Dakota												
All races, both sexes	2	3	3	4	3	5	4	6	5	1	-	-
Male	1	1	3	1	1	2	-	5	2	-	-	-
Female	1	2	-	3	2	3	4	1	3	1	-	-
White, both sexes	2	3	2	4	2	4	4	5	5	1	-	-
Male	1	1	2	1	1	2	-	4	2	-	-	-
Female	1	2	-	3	1	2	4	1	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	1	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	31	27	30	39	63	87	96	83	50	14	-	-
Male	19	16	16	13	29	38	41	29	11	3	-	-
Female	12	11	14	26	34	49	55	54	39	11	-	-
White, both sexes	23	24	24	34	50	72	79	66	39	11	-	-
Male	15	13	13	11	22	32	33	24	9	2	-	-
Female	8	11	11	23	28	40	46	42	30	9	-	-
Black, both sexes	8	3	6	5	13	15	17	17	11	3	-	-
Male	4	3	3	2	7	6	8	5	2	1	-	-
Female	4	-	3	3	6	9	9	12	9	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
Texas												
All races, both sexes	122	148	159	207	270	342	393	321	184	75	11	-
Male	52	76	81	122	143	155	169	113	59	29	-	-
Female	70	72	78	85	127	187	224	208	125	46	11	-
White, both sexes	94	112	122	164	215	288	347	279	162	65	8	-
Male	45	60	62	97	114	133	147	100	52	26	-	-
Female	49	52	60	67	101	155	200	179	110	39	8	-
Black, both sexes	27	36	35	40	54	50	39	37	22	9	3	-
Male	7	16	17	23	28	22	17	13	7	2	-	-
Female	20	20	18	17	26	28	22	24	15	7	3	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	2	1	4	7	4	-	1	-	-
Male	-	-	2	2	1	-	5	-	-	1	-	-
Female	1	-	-	-	-	4	2	4	-	-	-	-
Utah												
All races, both sexes	1	9	6	9	15	24	21	11	9	2	-	-
Male	-	4	1	2	7	12	8	5	5	2	-	-
Female	1	5	5	7	8	12	13	6	4	-	-	-
White, both sexes	1	9	6	9	15	22	20	11	8	2	-	-
Male	-	4	1	2	7	10	7	5	4	2	-	-
Female	1	5	5	7	8	12	13	6	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	2	-	-	1	-	-	-
Male	-	-	-	-	-	2	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
Vermont												
All races, both sexes	1	-	1	1	4	6	7	6	7	3	-	-
Male	-	-	1	1	2	4	4	2	1	1	-	-
Female	1	-	-	-	2	2	3	4	6	2	-	-
White, both sexes	1	-	1	1	4	6	7	6	7	3	-	-
Male	-	-	1	1	2	4	4	2	1	1	-	-
Female	1	-	-	-	2	2	3	4	6	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	49	62	52	96	135	188	222	191	113	30	6	1
Male	28	36	26	50	64	87	93	78	40	11	2	-
Female	21	26	26	46	71	101	129	113	73	19	4	1
White, both sexes	22	40	33	60	87	137	168	146	79	23	4	-
Male	13	21	17	33	39	67	70	65	32	11	1	-
Female	9	19	16	27	48	70	98	81	47	12	3	-
Black, both sexes	25	22	18	35	47	48	50	43	31	7	2	1
Male	14	15	8	16	25	18	22	11	8	-	1	-
Female	11	7	10	19	22	30	28	32	23	7	1	1
American Indian 1/, both sexes	2	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	3	4	2	3	-	-	-
Male	-	-	1	1	-	2	1	2	-	-	-	-
Female	-	-	-	-	-	1	3	-	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
Washington												
All races, both sexes	13	18	26	14	32	59	55	40	33	14	3	-
Male	9	7	9	5	18	30	23	16	8	3	-	-
Female	4	11	17	9	14	29	32	24	25	11	3	-
White, both sexes	13	14	22	12	29	56	51	38	30	12	2	-
Male	9	6	7	4	16	27	20	15	7	3	-	-
Female	4	8	15	8	13	29	31	23	23	9	2	-
Black, both sexes	-	1	1	1	-	1	1	-	1	-	-	-
Male	-	1	1	-	-	1	1	-	1	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	3	-	-	1	-	-	1	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	3	-	-	-	-	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	3	1	2	2	3	1	2	1	1	-
Male	-	-	1	1	1	2	2	1	-	-	-	-
Female	-	-	2	-	1	-	1	-	2	1	1	-
West Virginia												
All races, both sexes	16	17	27	22	49	52	57	43	21	5	1	-
Male	12	5	17	8	25	30	27	16	8	1	-	-
Female	4	12	10	14	24	22	30	27	13	4	1	-
White, both sexes	16	17	26	22	46	50	56	42	21	5	1	-
Male	12	5	17	8	23	29	27	16	8	1	-	-
Female	4	12	9	14	23	21	29	26	13	4	1	-
Black, both sexes	-	-	1	-	3	2	1	1	-	-	-	-
Male	-	-	-	-	2	1	-	-	-	-	-	-
Female	-	-	1	-	1	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
Wisconsin												
All races, both sexes	23	23	24	24	43	68	81	67	36	19	4	-
Male	10	14	6	10	24	34	39	25	11	4	-	-
Female	13	9	18	14	19	34	42	42	25	15	4	-
White, both sexes	19	19	22	21	38	63	78	67	36	19	3	-
Male	9	13	6	7	21	32	37	25	11	4	-	-
Female	10	6	16	14	17	31	41	42	25	15	3	-
Black, both sexes	4	2	1	2	2	5	2	-	-	-	1	-
Male	1	1	-	2	1	2	1	-	-	-	-	-
Female	3	1	1	-	1	3	1	-	-	-	1	-
American Indian 1/, both sexes	-	2	1	1	2	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	2	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	3	3	2	2	4	12	8	6	1	-	-
Male	1	2	2	1	1	3	6	2	2	1	-	-
Female	-	1	1	1	1	1	6	6	4	-	-	-
White, both sexes	1	3	3	2	2	4	11	8	6	1	-	-
Male	1	2	2	1	1	3	5	2	2	1	-	-
Female	-	1	1	1	1	1	6	6	4	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
United States												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
Colorado												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Plague (A20)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
United States												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
Colorado												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Plague (A20)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
United States												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
Florida												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
Kentucky												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
Texas												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Tetanus (A33-A35)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
United States												
All races, both sexes	-	-	-	-	1	1	-	1	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
White, both sexes	-	-	-	-	1	1	-	1	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
Texas												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tetanus (A33-A35)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
United States												
All races, both sexes	16	15	-	-	1	-	-	-	-	-	-	-
Male	10	9	-	-	1	-	-	-	-	-	-	-
Female	6	6	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	12	-	-	1	-	-	-	-	-	-	-
Male	8	7	-	-	1	-	-	-	-	-	-	-
Female	5	5	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	4	4	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	3	3	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	4	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	3	3	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
Illinois												
All races, both sexes	2	1	-	-	1	-	-	-	-	-	-	-
Male	2	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	1	-	-	1	-	-	-	-	-	-	-
Male	2	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
Kentucky												
All races, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
Minnesota												
All races, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
New Mexico												
All races, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	2	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
Ohio												
All races, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Whooping cough (A37)												
Wisconsin												
All races, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
United States												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
United States												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
Illinois												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
Rhode Island												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
Vermont												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Scarlet fever and erysipelas (A38,A46)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
United States												
All races, both sexes	-	-	-	1	-	-	-	1	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	-	-	-	1	-	-	-	1	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
Rhode Island												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
United States												
All races, both sexes	138	8	18	8	5	16	15	9	5	6	9	7
Male	79	2	11	5	4	11	8	7	3	3	5	5
Female	59	6	7	3	1	5	7	2	2	3	4	2
White, both sexes	110	7	15	7	5	12	12	8	2	5	5	6
Male	62	2	10	4	4	7	6	6	2	3	3	5
Female	48	5	5	3	1	5	6	2	-	2	2	1
Black, both sexes	22	-	3	1	-	3	2	1	2	1	3	1
Male	13	-	1	1	-	3	1	1	1	-	1	-
Female	9	-	2	-	-	-	1	-	1	1	2	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	1	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	-	-	-	1	1	-	-	-	-	-
Male	2	-	-	-	-	1	1	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
Alaska												
All races, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
Arkansas												
All races, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	20	2	4	-	1	2	2	1	-	-	3	1
Male	6	-	1	-	1	-	-	1	-	-	1	1
Female	14	2	3	-	-	2	2	-	-	-	2	-
White, both sexes	16	1	4	-	1	2	1	1	-	-	1	1
Male	6	-	1	-	1	-	-	1	-	-	1	1
Female	10	1	3	-	-	2	1	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	1	-	-	-	2	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	1	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
Colorado												
All races, both sexes	2	1	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	1	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
Florida												
All races, both sexes	11	1	3	-	-	2	-	1	-	1	-	1
Male	9	-	3	-	-	2	-	1	-	-	-	1
Female	2	1	-	-	-	-	-	-	-	1	-	-
White, both sexes	8	1	3	-	-	1	-	-	-	1	-	1
Male	6	-	3	-	-	1	-	-	-	-	-	1
Female	2	1	-	-	-	-	-	-	-	1	-	-
Black, both sexes	3	-	-	-	-	1	-	1	-	-	-	-
Male	3	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	4	1	-	-	1	-	-	-	1	1	-	-
Male	2	-	-	-	1	-	-	-	-	1	-	-
Female	2	1	-	-	-	-	-	-	1	-	-	-
White, both sexes	3	1	-	-	1	-	-	-	-	1	-	-
Male	2	-	-	-	1	-	-	-	-	1	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
Hawaii												
All races, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
Illinois												
All races, both sexes	8	-	2	1	-	-	-	-	-	1	1	2
Male	4	-	-	1	-	-	-	-	-	-	1	1
Female	4	-	2	-	-	-	-	-	-	1	-	1
White, both sexes	4	-	1	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	1	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	1	1	-	-	-	-	-	1	-	1
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	3	-	1	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
Iowa												
All races, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	3	1	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	1	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	6	-	-	-	-	3	1	-	-	-	1	-
Male	3	-	-	-	-	3	-	-	-	-	-	-
Female	3	-	-	-	-	-	1	-	-	-	1	-
White, both sexes	4	-	-	-	-	2	1	-	-	-	1	-
Male	2	-	-	-	-	2	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
Maine												
All races, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
Massachusetts												
All races, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	5	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	1	-	-	-	1	-	-
White, both sexes	5	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	1	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
Minnesota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	4	-	2	-	-	1	-	-	-	-	-	-
Male	4	-	2	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	1	-	-	1	-	-	-	-	-	-
Male	2	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
Missouri												
All races, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
Nebraska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	9	-	1	2	1	1	1	1	-	1	-	1
Male	6	-	-	1	1	1	1	-	-	1	-	1
Female	3	-	1	1	-	-	-	1	-	-	-	-
White, both sexes	7	-	-	2	1	-	1	1	-	1	-	1
Male	5	-	-	1	1	-	1	-	-	1	-	1
Female	2	-	-	1	-	-	-	1	-	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
New Mexico												
All races, both sexes	4	1	-	-	-	-	-	-	-	-	1	1
Male	4	1	-	-	-	-	-	-	-	-	1	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	1	-	-	-	-	-	-	-	-	-	1
Male	2	1	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	10	-	-	1	-	-	1	2	2	-	-	1
Male	8	-	-	1	-	-	1	1	2	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	8	-	-	1	-	-	1	2	1	-	-	1
Male	6	-	-	1	-	-	1	1	1	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
North Carolina												
All races, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
Ohio												
All races, both sexes	7	-	1	-	1	1	-	1	-	1	1	-
Male	6	-	1	-	1	1	-	1	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	5	-	1	-	1	-	-	1	-	1	1	-
Male	4	-	1	-	1	-	-	1	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	-	-	-	-	-	-	1	1	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
Oregon												
All races, both sexes	4	-	-	-	1	-	1	-	-	-	1	-
Male	2	-	-	-	-	-	1	-	-	-	1	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	1	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
Rhode Island												
All races, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
Texas												
All races, both sexes	3	-	-	1	-	1	-	-	-	-	-	-
Male	3	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	1	-	1	-	-	-	-	-	-
Male	3	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	2	-	1	-	-	-	-	1	-	-	-	-
Male	2	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	1	-	-	-	-	1	-	-	-	-
Male	2	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
Vermont												
All races, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
Washington												
All races, both sexes	3	-	-	-	-	1	-	1	-	-	-	-
Male	3	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	1	-	1	-	-	-	-
Male	3	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Meningococcal infection (A39)												
Wisconsin												
All races, both sexes	3	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	1	-	-	1	-	-	-	-	-
White, both sexes	3	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	1	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
United States												
All races, both sexes	8	4	2	3	4	1	2	3	2	2	1	-
Male	7	2	2	2	1	-	1	-	-	-	-	-
Female	1	2	-	1	3	1	1	3	2	2	1	-
White, both sexes	5	3	2	3	3	1	2	2	2	2	1	-
Male	4	1	2	2	-	-	1	-	-	-	-	-
Female	1	2	-	1	3	1	1	2	2	2	1	-
Black, both sexes	2	1	-	-	1	-	-	1	-	-	-	-
Male	2	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	2	1	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	1	-	-	-
White, both sexes	2	1	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
Florida												
All races, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
Illinois												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
Iowa												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	1	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	1	-	-
White, both sexes	-	1	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
Missouri												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
New Mexico												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2	-	-	1	-	-	-	-	-	-	-	-
Male	2	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
Ohio												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
Texas												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
Washington												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)												
Wisconsin												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
United States												
All races, both sexes	33,373	271	84	38	28	48	87	115	213	306	540	898
Male	15,011	167	41	18	13	25	46	51	114	162	271	487
Female	18,362	104	43	20	15	23	41	64	99	144	269	411
White, both sexes	26,678	165	60	31	19	34	48	75	131	216	343	584
Male	11,986	103	29	17	9	19	27	33	68	120	174	315
Female	14,692	62	31	14	10	15	21	42	63	96	169	269
Black, both sexes	6,010	96	19	5	7	13	33	36	68	75	182	279
Male	2,690	55	9	1	3	6	16	16	37	33	93	151
Female	3,320	41	10	4	4	7	17	20	31	42	89	128
American Indian 1/, both sexes	183	5	2	2	2	-	2	3	6	7	9	16
Male	85	5	2	-	1	-	1	1	3	5	3	8
Female	98	-	-	2	1	-	1	2	3	2	6	8
Asian or Pacific Islander 2/, both sexes	502	5	3	-	-	1	4	1	8	8	6	19
Male	250	4	1	-	-	-	2	1	6	4	1	13
Female	252	1	2	-	-	1	2	-	2	4	5	6
Alabama												
All races, both sexes	756	6	-	-	1	1	3	3	9	12	10	28
Male	293	3	-	-	1	-	2	1	5	4	6	12
Female	463	3	-	-	-	1	1	2	4	8	4	16
White, both sexes	522	2	-	-	1	-	1	1	3	7	6	15
Male	206	2	-	-	1	-	1	-	1	3	3	7
Female	316	-	-	-	-	-	-	1	2	4	3	8
Black, both sexes	231	4	-	-	-	1	2	2	6	4	4	12
Male	84	1	-	-	-	-	1	1	4	-	3	4
Female	147	3	-	-	-	1	1	1	2	4	1	8
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	1	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Alaska												
All races, both sexes	19	1	2	-	1	-	-	1	1	-	-	1
Male	9	1	2	-	1	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	1	1	-	-	1
White, both sexes	13	-	1	-	1	-	-	1	1	-	-	1
Male	5	-	1	-	1	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	1	1	-	-	1
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	456	9	3	1	1	-	1	3	-	6	11	13
Male	234	7	2	-	1	-	1	1	-	5	6	7
Female	222	2	1	1	-	-	-	2	-	1	5	6
White, both sexes	401	6	2	-	-	-	1	2	-	5	7	10
Male	201	4	1	-	-	-	1	1	-	4	4	4
Female	200	2	1	-	-	-	-	1	-	1	3	6
Black, both sexes	17	-	1	-	-	-	-	-	-	-	1	1
Male	12	-	1	-	-	-	-	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	36	3	-	1	1	-	-	1	-	1	3	2
Male	19	3	-	-	1	-	-	-	-	1	1	2
Female	17	-	-	1	-	-	-	1	-	-	2	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Arkansas												
All races, both sexes	470	-	1	-	-	1	1	2	4	6	8	7
Male	220	-	1	-	-	1	1	1	2	2	6	2
Female	250	-	-	-	-	-	-	1	2	4	2	5
White, both sexes	372	-	-	-	-	1	1	1	2	3	6	6
Male	177	-	-	-	-	1	1	1	1	1	4	2
Female	195	-	-	-	-	-	-	-	1	2	2	4
Black, both sexes	97	-	1	-	-	-	-	1	2	3	2	-
Male	43	-	1	-	-	-	-	-	1	1	2	-
Female	54	-	-	-	-	-	-	1	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
California												
All races, both sexes	931	15	7	1	-	2	8	1	9	12	16	34
Male	418	9	3	1	-	1	6	-	6	8	8	18
Female	513	6	4	-	-	1	2	1	3	4	8	16
White, both sexes	768	14	6	1	-	1	6	-	7	9	14	29
Male	340	9	3	1	-	1	5	-	4	6	6	15
Female	428	5	3	-	-	-	1	-	3	3	8	14
Black, both sexes	88	1	-	-	-	-	1	1	1	2	2	2
Male	46	-	-	-	-	-	1	-	1	2	2	-
Female	42	1	-	-	-	-	-	1	-	-	-	2
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	67	-	1	-	-	1	1	-	1	1	-	3
Male	28	-	-	-	-	-	-	-	1	-	-	3
Female	39	-	1	-	-	1	1	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Colorado												
All races, both sexes	301	2	3	-	-	1	-	2	1	3	8	13
Male	144	2	2	-	-	-	-	-	-	2	3	8
Female	157	-	1	-	-	1	-	2	1	1	5	5
White, both sexes	284	2	2	-	-	1	-	2	-	3	7	12
Male	134	2	2	-	-	-	-	-	-	2	3	7
Female	150	-	-	-	-	1	-	2	-	1	4	5
Black, both sexes	13	-	1	-	-	-	-	-	1	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	1	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	516	3	1	1	1	1	2	2	1	5	9	10
Male	249	2	-	-	1	1	-	1	-	3	5	6
Female	267	1	1	1	-	-	2	1	1	2	4	4
White, both sexes	467	1	1	-	1	-	2	2	1	3	7	10
Male	225	1	-	-	1	-	-	1	-	2	4	6
Female	242	-	1	-	-	-	2	1	1	1	3	4
Black, both sexes	47	2	-	1	-	1	-	-	-	2	2	-
Male	23	1	-	-	-	1	-	-	-	1	1	-
Female	24	1	-	1	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Delaware												
All races, both sexes	121	2	-	-	-	-	-	-	-	4	1	5
Male	59	2	-	-	-	-	-	-	-	1	1	3
Female	62	-	-	-	-	-	-	-	-	3	-	2
White, both sexes	102	1	-	-	-	-	-	-	-	2	1	4
Male	52	1	-	-	-	-	-	-	-	-	1	3
Female	50	-	-	-	-	-	-	-	-	2	-	1
Black, both sexes	18	1	-	-	-	-	-	-	-	2	-	-
Male	7	1	-	-	-	-	-	-	-	1	-	-
Female	11	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	106	3	-	-	-	-	-	-	-	-	4	3
Male	51	1	-	-	-	-	-	-	-	-	1	1
Female	55	2	-	-	-	-	-	-	-	-	3	2
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	93	3	-	-	-	-	-	-	-	-	4	3
Male	46	1	-	-	-	-	-	-	-	-	1	1
Female	47	2	-	-	-	-	-	-	-	-	3	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Florida												
All races, both sexes	1,981	23	4	2	2	2	5	9	10	23	40	38
Male	947	13	3	1	-	1	2	4	5	10	14	20
Female	1,034	10	1	1	2	1	3	5	5	13	26	18
White, both sexes	1,715	14	4	2	2	2	3	4	5	19	28	24
Male	838	8	3	1	-	1	1	1	3	10	10	14
Female	877	6	1	1	2	1	2	3	2	9	18	10
Black, both sexes	254	8	-	-	-	-	2	5	5	3	11	14
Male	107	4	-	-	-	-	1	3	2	-	4	6
Female	147	4	-	-	-	-	1	2	3	3	7	8
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	1	-	-	-	-	-	-	-	1	1	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	1	1	-
Georgia												
All races, both sexes	1,305	8	4	1	-	2	7	5	14	18	28	39
Male	559	4	3	-	-	1	5	2	8	10	17	16
Female	746	4	1	1	-	1	2	3	6	8	11	23
White, both sexes	840	5	2	-	-	1	3	2	8	12	14	25
Male	342	2	1	-	-	-	2	1	4	7	8	12
Female	498	3	1	-	-	1	1	1	4	5	6	13
Black, both sexes	454	3	2	1	-	1	3	3	6	5	13	14
Male	213	2	2	-	-	1	2	1	4	3	9	4
Female	241	1	-	1	-	-	1	2	2	2	4	10
American Indian 1/, both sexes	4	-	-	-	-	-	1	-	-	1	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Hawaii												
All races, both sexes	188	1	-	-	1	-	1	-	1	5	3	6
Male	93	1	-	-	-	-	1	-	1	2	-	3
Female	95	-	-	-	1	-	-	-	-	3	3	3
White, both sexes	37	-	-	-	1	-	-	-	-	1	1	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	1	-	-	-	-	1	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	149	1	-	-	-	-	1	-	1	4	2	6
Male	72	1	-	-	-	-	1	-	1	2	-	3
Female	77	-	-	-	-	-	-	-	-	2	2	3
Idaho												
All races, both sexes	55	-	2	-	-	-	-	-	-	2	-	1
Male	23	-	1	-	-	-	-	-	-	1	-	1
Female	32	-	1	-	-	-	-	-	-	1	-	-
White, both sexes	54	-	2	-	-	-	-	-	-	2	-	1
Male	22	-	1	-	-	-	-	-	-	1	-	1
Female	32	-	1	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Illinois												
All races, both sexes	1,997	12	5	1	-	1	4	3	16	19	26	46
Male	874	7	2	-	-	1	2	2	5	13	10	25
Female	1,123	5	3	1	-	-	2	1	11	6	16	21
White, both sexes	1,513	7	2	1	-	1	2	1	10	15	16	27
Male	673	5	2	-	-	1	1	1	3	9	6	19
Female	840	2	-	1	-	-	1	-	7	6	10	8
Black, both sexes	459	5	3	-	-	-	2	2	5	4	10	18
Male	191	2	-	-	-	-	1	1	2	4	4	6
Female	268	3	3	-	-	-	1	1	3	-	6	12
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	25	-	-	-	-	-	-	-	1	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	1	-	-	1
Indiana												
All races, both sexes	820	7	4	-	1	5	2	2	8	5	11	21
Male	359	4	2	-	1	2	1	-	1	2	5	7
Female	461	3	2	-	-	3	1	2	7	3	6	14
White, both sexes	716	5	4	-	1	3	1	2	6	4	9	13
Male	316	3	2	-	1	2	1	-	1	2	3	4
Female	400	2	2	-	-	1	-	2	5	2	6	9
Black, both sexes	103	2	-	-	-	2	1	-	2	1	2	8
Male	43	1	-	-	-	-	-	-	-	-	2	3
Female	60	1	-	-	-	2	1	-	2	1	-	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Iowa												
All races, both sexes	228	-	-	1	-	-	1	-	1	1	2	7
Male	103	-	-	-	-	-	1	-	-	1	1	3
Female	125	-	-	1	-	-	-	-	1	-	1	4
White, both sexes	224	-	-	1	-	-	1	-	1	1	2	7
Male	102	-	-	-	-	-	1	-	-	1	1	3
Female	122	-	-	1	-	-	-	-	1	-	1	4
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	280	3	1	-	2	-	1	1	3	-	4	6
Male	114	2	1	-	1	-	1	-	1	-	3	3
Female	166	1	-	-	1	-	-	1	2	-	1	3
White, both sexes	257	2	1	-	1	-	1	1	3	-	4	3
Male	105	1	1	-	-	-	1	-	1	-	3	3
Female	152	1	-	-	1	-	-	1	2	-	1	-
Black, both sexes	20	1	-	-	1	-	-	-	-	-	-	2
Male	8	1	-	-	1	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Kentucky												
All races, both sexes	573	1	-	-	-	1	1	1	8	5	9	16
Male	275	1	-	-	-	1	1	1	7	1	3	11
Female	298	-	-	-	-	-	-	-	1	4	6	5
White, both sexes	534	1	-	-	-	-	-	-	7	3	7	14
Male	259	1	-	-	-	-	-	-	6	1	3	11
Female	275	-	-	-	-	-	-	-	1	2	4	3
Black, both sexes	39	-	-	-	-	1	1	1	1	2	2	2
Male	16	-	-	-	-	1	1	1	1	-	-	-
Female	23	-	-	-	-	-	-	-	-	2	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	837	7	3	1	3	3	-	8	7	6	14	17
Male	367	3	2	1	-	2	-	6	3	2	8	8
Female	470	4	1	-	3	1	-	2	4	4	6	9
White, both sexes	527	4	1	1	2	2	-	3	3	3	9	10
Male	223	2	-	1	-	1	-	3	-	1	6	3
Female	304	2	1	-	2	1	-	-	3	2	3	7
Black, both sexes	304	3	2	-	1	1	-	5	3	3	5	7
Male	140	1	2	-	-	1	-	3	2	1	2	5
Female	164	2	-	-	1	-	-	2	1	2	3	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Maine												
All races, both sexes	139	2	-	-	1	-	1	-	-	1	1	1
Male	58	1	-	-	1	-	1	-	-	-	-	-
Female	81	1	-	-	-	-	-	-	-	1	1	1
White, both sexes	138	2	-	-	1	-	1	-	-	1	1	1
Male	57	1	-	-	1	-	1	-	-	-	-	-
Female	81	1	-	-	-	-	-	-	-	1	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1,063	3	2	1	-	1	3	4	7	8	21	33
Male	487	2	1	-	-	-	2	4	6	4	12	23
Female	576	1	1	1	-	1	1	-	1	4	9	10
White, both sexes	685	1	1	-	-	-	1	4	3	2	7	14
Male	316	1	1	-	-	-	1	4	2	2	4	8
Female	369	-	-	-	-	-	-	-	1	-	3	6
Black, both sexes	361	2	1	1	-	1	1	-	4	5	14	18
Male	165	1	-	-	-	-	1	-	4	1	8	15
Female	196	1	1	1	-	1	-	-	-	4	6	3
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	-	1
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	1	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Massachusetts												
All races, both sexes	963	1	1	1	1	-	-	-	11	4	11	19
Male	415	-	-	-	-	-	-	-	7	1	4	11
Female	548	1	1	1	1	-	-	-	4	3	7	8
White, both sexes	892	-	-	1	1	-	-	-	7	4	9	16
Male	386	-	-	-	-	-	-	-	4	1	3	9
Female	506	-	-	1	1	-	-	-	3	3	6	7
Black, both sexes	63	1	1	-	-	-	-	-	3	-	2	3
Male	27	-	-	-	-	-	-	-	2	-	1	2
Female	36	1	1	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	852	2	2	-	1	-	-	1	4	4	15	18
Male	362	1	-	-	-	-	-	-	2	1	6	14
Female	490	1	2	-	1	-	-	1	2	3	9	4
White, both sexes	651	2	1	-	-	-	-	1	3	3	9	8
Male	267	1	-	-	-	-	-	-	1	1	2	8
Female	384	1	1	-	-	-	-	1	2	2	7	-
Black, both sexes	193	-	1	-	1	-	-	-	1	1	6	10
Male	91	-	-	-	-	-	-	-	1	-	4	6
Female	102	-	1	-	1	-	-	-	-	1	2	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Minnesota												
All races, both sexes	269	1	-	1	-	1	-	2	1	3	3	8
Male	132	1	-	1	-	1	-	1	-	2	2	5
Female	137	-	-	-	-	-	-	1	1	1	1	3
White, both sexes	250	-	-	1	-	1	-	2	-	3	3	3
Male	123	-	-	1	-	1	-	1	-	2	2	1
Female	127	-	-	-	-	-	-	1	-	1	1	2
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	3
Male	3	-	-	-	-	-	-	-	-	-	-	2
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	1	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	2
Female	3	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	500	6	-	-	1	3	4	4	1	8	9	18
Male	208	6	-	-	-	1	1	1	1	5	3	10
Female	292	-	-	-	1	2	3	3	-	3	6	8
White, both sexes	311	1	-	-	-	3	1	3	-	6	5	5
Male	135	1	-	-	-	1	-	1	-	5	3	1
Female	176	-	-	-	-	2	1	2	-	1	2	4
Black, both sexes	187	5	-	-	-	-	3	1	1	2	4	13
Male	73	5	-	-	-	-	1	-	1	-	-	9
Female	114	-	-	-	-	-	2	1	-	2	4	4
American Indian 1/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Missouri												
All races, both sexes	669	3	1	1	2	1	-	-	6	1	5	16
Male	287	3	1	1	1	1	-	-	2	-	2	6
Female	382	-	-	-	1	-	-	-	4	1	3	10
White, both sexes	555	3	1	1	1	-	-	-	5	1	2	10
Male	234	3	1	1	1	-	-	-	2	-	1	3
Female	321	-	-	-	-	-	-	-	3	1	1	7
Black, both sexes	108	-	-	-	1	1	-	-	1	-	3	6
Male	50	-	-	-	-	1	-	-	-	-	1	3
Female	58	-	-	-	1	-	-	-	1	-	2	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	70	-	-	-	-	-	1	-	-	1	2	1
Male	31	-	-	-	-	-	-	-	-	1	1	1
Female	39	-	-	-	-	-	1	-	-	-	1	-
White, both sexes	66	-	-	-	-	-	1	-	-	1	1	1
Male	28	-	-	-	-	-	-	-	-	1	-	1
Female	38	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Nebraska												
All races, both sexes	133	1	1	-	-	-	-	-	1	4	1	1
Male	58	-	1	-	-	-	-	-	-	3	1	1
Female	75	1	-	-	-	-	-	-	1	1	-	-
White, both sexes	124	1	1	-	-	-	-	-	1	4	1	1
Male	56	-	1	-	-	-	-	-	-	3	1	1
Female	68	1	-	-	-	-	-	-	1	1	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	395	6	-	-	1	-	-	1	3	7	6	12
Male	207	3	-	-	1	-	-	-	1	5	2	8
Female	188	3	-	-	-	-	-	1	2	2	4	4
White, both sexes	347	6	-	-	1	-	-	1	3	7	4	11
Male	185	3	-	-	1	-	-	-	1	5	2	7
Female	162	3	-	-	-	-	-	1	2	2	2	4
Black, both sexes	30	-	-	-	-	-	-	-	-	-	1	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
New Hampshire												
All races, both sexes	91	-	-	-	-	-	-	-	1	1	1	3
Male	50	-	-	-	-	-	-	-	1	-	-	2
Female	41	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	90	-	-	-	-	-	-	-	1	1	1	2
Male	49	-	-	-	-	-	-	-	1	-	-	1
Female	41	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1,903	16	2	1	-	1	5	3	4	10	27	29
Male	834	9	1	1	-	-	4	2	1	5	13	17
Female	1,069	7	1	-	-	1	1	1	3	5	14	12
White, both sexes	1,566	11	-	1	-	1	1	2	3	7	11	20
Male	679	7	-	1	-	-	1	1	-	4	7	11
Female	887	4	-	-	-	1	-	1	3	3	4	9
Black, both sexes	301	5	1	-	-	-	4	-	1	3	16	9
Male	136	2	-	-	-	-	3	-	1	1	6	6
Female	165	3	1	-	-	-	1	-	-	2	10	3
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	33	-	1	-	-	-	-	1	-	-	-	-
Male	19	-	1	-	-	-	-	1	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
New Mexico												
All races, both sexes	144	1	1	1	-	-	1	1	-	5	4	5
Male	66	1	1	-	-	-	-	1	-	3	2	2
Female	78	-	-	1	-	-	1	-	-	2	2	3
White, both sexes	125	-	1	-	-	-	1	-	-	4	3	4
Male	58	-	1	-	-	-	-	-	-	2	2	1
Female	67	-	-	-	-	-	1	-	-	2	1	3
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	15	1	-	1	-	-	-	1	-	1	1	-
Male	6	1	-	-	-	-	-	1	-	1	-	-
Female	9	-	-	1	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2,278	13	4	2	2	5	6	13	8	17	21	40
Male	1,048	7	2	1	2	5	2	9	4	9	14	20
Female	1,230	6	2	1	-	-	4	4	4	8	7	20
White, both sexes	1,917	8	2	2	1	5	5	10	5	11	12	28
Male	893	6	1	1	1	5	2	6	4	6	9	14
Female	1,024	2	1	1	-	-	3	4	1	5	3	14
Black, both sexes	324	5	1	-	1	-	1	3	3	6	9	12
Male	134	1	1	-	1	-	-	3	-	3	5	6
Female	190	4	-	-	-	-	1	-	3	3	4	6
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	35	-	1	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
North Carolina												
All races, both sexes	1,200	17	2	3	-	2	2	3	11	10	24	33
Male	534	9	1	2	-	1	1	1	6	6	13	15
Female	666	8	1	1	-	1	1	2	5	4	11	18
White, both sexes	826	6	2	2	-	-	1	1	5	7	8	16
Male	354	3	1	1	-	-	1	1	3	4	5	7
Female	472	3	1	1	-	-	-	-	2	3	3	9
Black, both sexes	357	11	-	1	-	2	1	2	5	2	15	16
Male	172	6	-	1	-	1	-	-	3	1	8	7
Female	185	5	-	-	-	1	1	2	2	1	7	9
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	1	1	-	1
Male	6	-	-	-	-	-	-	-	-	1	-	1
Female	8	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
North Dakota												
All races, both sexes	33	-	-	-	-	-	-	-	1	-	-	2
Male	16	-	-	-	-	-	-	-	1	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	2
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Ohio												
All races, both sexes	1,488	11	3	-	3	1	2	5	7	16	29	46
Male	651	9	-	-	1	-	1	5	3	8	16	22
Female	837	2	3	-	2	1	1	-	4	8	13	24
White, both sexes	1,232	4	3	-	2	1	1	4	7	10	24	33
Male	533	4	-	-	-	-	1	4	3	6	14	16
Female	699	-	3	-	2	1	-	-	4	4	10	17
Black, both sexes	249	6	-	-	1	-	1	1	-	6	5	13
Male	116	4	-	-	1	-	-	1	-	2	2	6
Female	133	2	-	-	-	-	1	-	-	4	3	7
American Indian 1/, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	414	5	2	1	-	1	-	1	5	5	10	17
Male	191	1	1	-	-	-	-	-	4	3	4	7
Female	223	4	1	1	-	1	-	1	1	2	6	10
White, both sexes	351	5	-	1	-	1	-	1	3	3	4	13
Male	157	1	-	-	-	-	-	-	2	1	1	6
Female	194	4	-	1	-	1	-	1	1	2	3	7
Black, both sexes	43	-	1	-	-	-	-	-	-	2	5	1
Male	20	-	-	-	-	-	-	-	-	2	2	-
Female	23	-	1	-	-	-	-	-	-	-	3	1
American Indian 1/, both sexes	19	-	1	-	-	-	-	-	2	-	1	3
Male	13	-	1	-	-	-	-	-	2	-	1	1
Female	6	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Oregon												
All races, both sexes	198	-	-	-	-	-	1	3	-	4	3	9
Male	94	-	-	-	-	-	-	1	-	2	1	6
Female	104	-	-	-	-	-	1	2	-	2	2	3
White, both sexes	187	-	-	-	-	-	1	2	-	4	3	8
Male	90	-	-	-	-	-	-	1	-	2	1	6
Female	97	-	-	-	-	-	1	1	-	2	2	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	1	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	1	-	-	-	1
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	2,491	13	2	4	1	4	3	3	8	13	34	69
Male	1,115	8	1	1	1	2	2	-	3	4	19	37
Female	1,376	5	1	3	-	2	1	3	5	9	15	32
White, both sexes	2,095	10	1	4	1	4	2	2	2	9	23	44
Male	928	5	1	1	1	2	1	-	2	4	13	20
Female	1,167	5	-	3	-	2	1	2	-	5	10	24
Black, both sexes	373	3	1	-	-	-	1	1	6	4	10	24
Male	175	3	-	-	-	-	1	-	1	-	5	17
Female	198	-	1	-	-	-	-	1	5	4	5	7
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	-	-	1	1
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Rhode Island												
All races, both sexes	83	-	-	-	1	-	1	-	-	-	1	1
Male	37	-	-	-	-	-	1	-	-	-	-	-
Female	46	-	-	-	1	-	-	-	-	-	1	1
White, both sexes	76	-	-	-	1	-	-	-	-	-	1	1
Male	33	-	-	-	-	-	-	-	-	-	-	-
Female	43	-	-	-	1	-	-	-	-	-	1	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	625	7	1	1	-	1	4	2	-	4	5	23
Male	265	4	-	-	-	-	-	1	-	2	4	11
Female	360	3	1	1	-	1	4	1	-	2	1	12
White, both sexes	395	2	1	-	-	-	3	2	-	2	1	5
Male	164	2	-	-	-	-	-	1	-	1	-	4
Female	231	-	1	-	-	-	3	1	-	1	1	1
Black, both sexes	223	3	-	1	-	1	1	-	-	2	4	17
Male	98	1	-	-	-	-	-	-	-	1	4	7
Female	125	2	-	1	-	1	1	-	-	1	-	10
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	6	2	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	3	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
South Dakota												
All races, both sexes	42	-	-	2	-	-	-	-	-	1	1	3
Male	20	-	-	2	-	-	-	-	-	-	-	3
Female	22	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	35	-	-	2	-	-	-	-	-	-	-	2
Male	17	-	-	2	-	-	-	-	-	-	-	2
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	1	1	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	538	7	2	-	-	1	2	4	2	6	12	26
Male	227	3	1	-	-	-	1	1	2	6	5	13
Female	311	4	1	-	-	1	1	3	-	-	7	13
White, both sexes	419	2	2	-	-	-	1	2	2	3	8	15
Male	176	-	1	-	-	-	1	1	2	3	3	7
Female	243	2	1	-	-	-	-	1	-	-	5	8
Black, both sexes	118	5	-	-	-	1	1	2	-	3	4	10
Male	50	3	-	-	-	-	-	-	-	3	2	5
Female	68	2	-	-	-	1	1	2	-	-	2	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Texas												
All races, both sexes	2,424	39	7	8	1	3	6	16	24	28	53	88
Male	1,115	28	3	5	-	2	2	3	14	17	28	56
Female	1,309	11	4	3	1	1	4	13	10	11	25	32
White, both sexes	1,987	30	6	8	-	3	2	14	16	22	37	65
Male	910	20	2	5	-	2	1	2	9	12	21	40
Female	1,077	10	4	3	-	1	1	12	7	10	16	25
Black, both sexes	412	9	1	-	1	-	4	2	8	5	15	22
Male	192	8	1	-	-	-	1	1	5	4	7	15
Female	220	1	-	-	1	-	3	1	3	1	8	7
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	23	-	-	-	-	-	-	-	-	1	-	1
Male	13	-	-	-	-	-	-	-	-	1	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	112	-	2	-	-	-	-	1	1	1	-	5
Male	46	-	-	-	-	-	-	-	-	-	-	2
Female	66	-	2	-	-	-	-	1	1	1	-	3
White, both sexes	106	-	2	-	-	-	-	1	-	1	-	4
Male	42	-	-	-	-	-	-	-	-	-	-	2
Female	64	-	2	-	-	-	-	1	-	1	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	1	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Vermont												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1,195	9	6	-	-	1	6	5	3	4	22	27
Male	550	6	1	-	-	-	4	2	2	2	13	20
Female	645	3	5	-	-	1	2	3	1	2	9	7
White, both sexes	823	4	6	-	-	1	3	1	2	2	14	14
Male	386	3	1	-	-	-	2	1	2	1	8	9
Female	437	1	5	-	-	1	1	-	-	1	6	5
Black, both sexes	353	5	-	-	-	-	3	4	1	2	8	11
Male	154	3	-	-	-	-	2	1	-	1	5	9
Female	199	2	-	-	-	-	1	3	1	1	3	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	-	2
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Washington												
All races, both sexes	311	1	2	1	-	1	1	-	5	2	7	10
Male	139	1	1	-	-	1	-	-	4	1	3	6
Female	172	-	1	1	-	-	1	-	1	1	4	4
White, both sexes	276	-	1	1	-	1	-	-	1	2	7	8
Male	119	-	-	-	-	1	-	-	1	1	3	4
Female	157	-	1	1	-	-	-	-	-	1	4	4
Black, both sexes	10	1	1	-	-	-	-	-	1	-	-	1
Male	9	1	1	-	-	-	-	-	1	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	3	-	-	1
Male	11	-	-	-	-	-	-	-	2	-	-	1
Female	9	-	-	-	-	-	-	-	1	-	-	-
West Virginia												
All races, both sexes	323	-	-	-	-	1	1	-	2	4	4	10
Male	154	-	-	-	-	-	-	-	2	3	2	3
Female	169	-	-	-	-	1	1	-	-	1	2	7
White, both sexes	313	-	-	-	-	1	1	-	2	4	3	9
Male	150	-	-	-	-	-	-	-	2	3	1	3
Female	163	-	-	-	-	1	1	-	-	1	2	6
Black, both sexes	10	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Wisconsin												
All races, both sexes	416	3	1	1	-	-	-	-	4	2	4	12
Male	189	2	1	1	-	-	-	-	4	2	4	10
Female	227	1	-	-	-	-	-	-	-	-	-	2
White, both sexes	381	2	1	1	-	-	-	-	3	-	3	8
Male	169	1	1	1	-	-	-	-	3	-	3	7
Female	212	1	-	-	-	-	-	-	-	-	-	1
Black, both sexes	24	1	-	-	-	-	-	-	1	1	1	2
Male	13	1	-	-	-	-	-	-	1	1	1	1
Female	11	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	1	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	40	1	-	-	-	-	-	-	-	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	1
Female	20	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	39	1	-	-	-	-	-	-	-	-	-	1
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	20	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
United States												
All races, both sexes	1,353	1,700	2,045	2,481	3,502	4,828	5,758	4,914	2,943	1,045	173	3
Male	720	910	1,024	1,247	1,770	2,308	2,562	1,904	869	267	34	1
Female	633	790	1,021	1,234	1,732	2,520	3,196	3,010	2,074	778	139	2
White, both sexes	931	1,237	1,490	1,868	2,781	3,990	4,869	4,240	2,540	897	127	2
Male	510	664	747	951	1,419	1,936	2,169	1,672	746	232	25	1
Female	421	573	743	917	1,362	2,054	2,700	2,568	1,794	665	102	1
Black, both sexes	385	428	488	561	650	756	792	593	366	133	44	1
Male	194	230	238	269	316	338	341	200	105	30	9	-
Female	191	198	250	292	334	418	451	393	261	103	35	1
American Indian 1/, both sexes	15	11	22	12	13	19	12	17	5	3	-	-
Male	7	2	12	7	5	9	4	8	2	-	-	-
Female	8	9	10	5	8	10	8	9	3	3	-	-
Asian or Pacific Islander 2/, both sexes	22	24	45	40	58	63	85	64	32	12	2	-
Male	9	14	27	20	30	25	48	24	16	5	-	-
Female	13	10	18	20	28	38	37	40	16	7	2	-
Alabama												
All races, both sexes	33	46	49	67	91	103	125	100	42	21	6	-
Male	16	25	19	28	38	56	34	29	8	5	1	-
Female	17	21	30	39	53	47	91	71	34	16	5	-
White, both sexes	18	34	32	46	66	70	91	80	27	17	5	-
Male	10	20	13	21	31	35	26	24	5	3	-	-
Female	8	14	19	25	35	35	65	56	22	14	5	-
Black, both sexes	15	12	16	21	25	33	34	20	15	4	1	-
Male	6	5	5	7	7	21	8	5	3	2	1	-
Female	9	7	11	14	18	12	26	15	12	2	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Alaska												
All races, both sexes	2	1	2	2	-	-	3	-	2	-	-	-
Male	1	1	-	1	-	-	1	-	1	-	-	-
Female	1	-	2	1	-	-	2	-	1	-	-	-
White, both sexes	1	1	1	1	-	-	3	-	1	-	-	-
Male	1	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	1	-	-	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	27	28	37	35	68	70	56	53	29	4	1	-
Male	17	17	18	18	34	31	27	25	15	2	-	-
Female	10	11	19	17	34	39	29	28	14	2	1	-
White, both sexes	21	24	32	30	65	64	54	47	26	4	1	-
Male	13	14	15	15	34	28	25	22	14	2	-	-
Female	8	10	17	15	31	36	29	25	12	2	1	-
Black, both sexes	3	2	1	2	1	1	1	2	1	-	-	-
Male	2	1	1	2	-	1	1	1	-	-	-	-
Female	1	1	-	-	1	-	-	1	1	-	-	-
American Indian 1/, both sexes	2	2	4	3	2	4	1	4	2	-	-	-
Male	1	2	2	1	-	1	1	2	1	-	-	-
Female	1	-	2	2	2	3	-	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Arkansas												
All races, both sexes	21	21	31	29	43	72	80	66	52	19	6	-
Male	15	16	16	15	22	25	35	30	25	4	1	-
Female	6	5	15	14	21	47	45	36	27	15	5	-
White, both sexes	14	16	23	23	37	59	67	57	38	14	4	-
Male	9	12	12	13	18	19	33	26	19	4	1	-
Female	5	4	11	10	19	40	34	31	19	10	3	-
Black, both sexes	7	5	8	6	6	13	13	9	14	5	2	-
Male	6	4	4	2	4	6	2	4	6	-	-	-
Female	1	1	4	4	2	7	11	5	8	5	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	43	53	69	73	76	108	166	131	73	30	4	-
Male	23	26	36	40	34	50	64	58	18	9	-	-
Female	20	27	33	33	42	58	102	73	55	21	4	-
White, both sexes	32	44	52	63	69	86	138	105	65	25	2	-
Male	16	20	30	35	31	39	52	47	14	6	-	-
Female	16	24	22	28	38	47	86	58	51	19	2	-
Black, both sexes	8	6	8	4	4	11	13	18	3	2	1	-
Male	5	4	3	3	3	6	6	8	-	2	-	-
Female	3	2	5	1	1	5	7	10	3	-	1	-
American Indian 1/, both sexes	1	-	1	-	-	2	3	-	1	-	-	-
Male	1	-	-	-	-	2	1	-	-	-	-	-
Female	-	-	1	-	-	-	2	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	8	6	3	9	12	8	4	3	1	-
Male	1	2	3	2	-	3	5	3	4	1	-	-
Female	1	1	5	4	3	6	7	5	-	2	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Colorado												
All races, both sexes	7	16	19	17	24	38	56	59	24	7	1	-
Male	4	10	10	7	12	18	26	28	6	5	1	-
Female	3	6	9	10	12	20	30	31	18	2	-	-
White, both sexes	6	15	18	15	24	38	51	56	24	7	1	-
Male	4	9	9	5	12	18	22	27	6	5	1	-
Female	2	6	9	10	12	20	29	29	18	2	-	-
Black, both sexes	-	1	1	2	-	-	2	3	-	-	-	-
Male	-	1	1	2	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	3	-	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Connecticut												
All races, both sexes	18	17	13	33	42	79	107	95	57	18	1	-
Male	11	11	5	21	25	47	45	45	14	5	1	-
Female	7	6	8	12	17	32	62	50	43	13	-	-
White, both sexes	13	13	11	27	37	70	103	91	55	18	1	-
Male	8	9	5	17	22	43	44	42	14	5	1	-
Female	5	4	6	10	15	27	59	49	41	13	-	-
Black, both sexes	5	4	1	5	5	9	4	4	2	-	-	-
Male	3	2	-	3	3	4	1	3	-	-	-	-
Female	2	2	1	2	2	5	3	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Delaware												
All races, both sexes	4	7	4	7	15	20	24	17	9	2	-	-
Male	2	3	3	4	8	10	13	5	3	1	-	-
Female	2	4	1	3	7	10	11	12	6	1	-	-
White, both sexes	3	6	4	6	14	16	22	15	7	1	-	-
Male	1	3	3	3	8	8	13	5	2	1	-	-
Female	2	3	1	3	6	8	9	10	5	-	-	-
Black, both sexes	1	1	-	1	1	4	2	2	2	1	-	-
Male	1	-	-	1	-	2	-	-	1	-	-	-
Female	-	1	-	-	1	2	2	2	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	7	4	8	11	13	9	22	13	8	1	-	-
Male	5	3	5	7	4	2	14	6	2	-	-	-
Female	2	1	3	4	9	7	8	7	6	1	-	-
White, both sexes	-	-	1	-	-	1	6	1	3	-	-	-
Male	-	-	1	-	-	-	4	-	-	-	-	-
Female	-	-	-	-	-	1	2	1	3	-	-	-
Black, both sexes	7	4	7	11	12	8	16	12	5	1	-	-
Male	5	3	4	7	4	2	10	6	2	-	-	-
Female	2	1	3	4	8	6	6	6	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Florida												
All races, both sexes	63	87	120	144	199	303	339	295	197	64	12	-
Male	34	49	69	78	98	162	169	126	68	17	4	-
Female	29	38	51	66	101	141	170	169	129	47	8	-
White, both sexes	46	71	94	112	175	275	316	267	182	61	9	-
Male	26	43	55	64	84	152	161	116	65	16	4	-
Female	20	28	39	48	91	123	155	151	117	45	5	-
Black, both sexes	16	16	26	31	23	27	21	25	15	3	3	-
Male	8	6	14	14	13	10	8	10	3	1	-	-
Female	8	10	12	17	10	17	13	15	12	2	3	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	2	2	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	2	2	-	-	-	-
Georgia												
All races, both sexes	63	76	92	116	142	203	184	158	96	35	14	-
Male	31	39	39	60	79	88	66	56	23	11	1	-
Female	32	37	53	56	63	115	118	102	73	24	13	-
White, both sexes	27	47	50	74	82	143	135	116	61	26	7	-
Male	12	25	19	39	47	57	46	41	12	7	-	-
Female	15	22	31	35	35	86	89	75	49	19	7	-
Black, both sexes	36	28	42	40	59	58	48	41	35	9	7	-
Male	19	13	20	21	31	30	20	15	11	4	1	-
Female	17	15	22	19	28	28	28	26	24	5	6	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	1	2	-	1	-	-	-	-
Male	-	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Hawaii												
All races, both sexes	12	8	14	12	22	22	31	23	20	6	-	-
Male	4	4	8	6	12	8	19	9	12	3	-	-
Female	8	4	6	6	10	14	12	14	8	3	-	-
White, both sexes	1	3	1	5	3	5	7	4	5	-	-	-
Male	1	2	1	1	2	3	3	3	4	-	-	-
Female	-	1	-	4	1	2	4	1	1	-	-	-
Black, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	5	13	6	19	17	23	19	15	6	-	-
Male	3	2	7	5	10	5	15	6	8	3	-	-
Female	8	3	6	1	9	12	8	13	7	3	-	-
Idaho												
All races, both sexes	2	-	3	4	7	11	6	6	7	4	-	-
Male	2	-	-	2	1	7	3	2	1	2	-	-
Female	-	-	3	2	6	4	3	4	6	2	-	-
White, both sexes	2	-	3	4	7	10	6	6	7	4	-	-
Male	2	-	-	2	1	6	3	2	1	2	-	-
Female	-	-	3	2	6	4	3	4	6	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Illinois												
All races, both sexes	79	93	123	160	208	277	357	311	182	64	10	-
Male	45	52	56	71	105	133	164	120	54	7	-	-
Female	34	41	67	89	103	144	193	191	128	57	10	-
White, both sexes	51	62	76	110	156	222	281	260	154	51	8	-
Male	31	36	33	50	77	108	132	106	46	7	-	-
Female	20	26	43	60	79	114	149	154	108	44	8	-
Black, both sexes	27	30	44	47	50	52	73	48	25	12	2	-
Male	13	16	20	20	28	24	29	13	8	-	-	-
Female	14	14	24	27	22	28	44	35	17	12	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	3	3	2	3	3	3	3	1	-	-
Male	1	-	3	1	-	1	3	1	-	-	-	-
Female	-	1	-	2	2	2	-	2	3	1	-	-
Indiana												
All races, both sexes	35	51	62	57	96	113	138	123	57	19	3	-
Male	25	29	30	25	50	45	62	42	20	5	1	-
Female	10	22	32	32	46	68	76	81	37	14	2	-
White, both sexes	29	45	54	49	86	94	125	114	52	17	3	-
Male	20	26	26	21	46	37	55	40	20	5	1	-
Female	9	19	28	28	40	57	70	74	32	12	2	-
Black, both sexes	6	6	8	8	9	19	13	9	5	2	-	-
Male	5	3	4	4	4	8	7	2	-	-	-	-
Female	1	3	4	4	5	11	6	7	5	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Iowa												
All races, both sexes	9	7	13	16	25	44	38	36	17	9	1	-
Male	6	4	5	5	14	23	20	16	-	3	1	-
Female	3	3	8	11	11	21	18	20	17	6	-	-
White, both sexes	9	7	12	15	25	43	38	36	17	8	1	-
Male	6	4	5	5	14	23	20	16	-	2	1	-
Female	3	3	7	10	11	20	18	20	17	6	-	-
Black, both sexes	-	-	-	1	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	10	9	20	21	27	38	51	47	22	11	3	-
Male	4	4	14	10	15	18	20	10	4	2	1	-
Female	6	5	6	11	12	20	31	37	18	9	2	-
White, both sexes	9	7	19	17	24	37	46	46	22	11	3	-
Male	3	4	14	8	14	18	18	9	4	2	1	-
Female	6	3	5	9	10	19	28	37	18	9	2	-
Black, both sexes	1	2	1	4	2	1	5	-	-	-	-	-
Male	1	-	-	2	1	-	2	-	-	-	-	-
Female	-	2	1	2	1	1	3	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Kentucky												
All races, both sexes	33	33	40	44	65	94	94	74	45	8	1	-
Male	23	15	21	24	30	49	44	28	13	2	-	-
Female	10	18	19	20	35	45	50	46	32	6	1	-
White, both sexes	31	31	38	40	59	91	91	70	44	7	-	-
Male	21	15	21	22	28	48	43	24	13	2	-	-
Female	10	16	17	18	31	43	48	46	31	5	-	-
Black, both sexes	2	2	2	4	6	3	3	4	1	1	1	-
Male	2	-	-	2	2	1	1	4	-	-	-	-
Female	-	2	2	2	4	2	2	-	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	38	47	58	73	84	106	144	111	79	24	4	-
Male	16	23	26	41	40	50	66	39	21	7	3	-
Female	22	24	32	32	44	56	78	72	58	17	1	-
White, both sexes	23	27	29	46	49	65	92	82	58	15	3	-
Male	11	13	9	24	24	31	42	30	14	5	3	-
Female	12	14	20	22	25	34	50	52	44	10	-	-
Black, both sexes	15	19	29	25	34	41	52	28	21	9	1	-
Male	5	10	17	15	15	19	24	9	7	2	-	-
Female	10	9	12	10	19	22	28	19	14	7	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Maine												
All races, both sexes	8	7	5	8	15	18	28	20	18	4	1	-
Male	2	4	3	4	7	8	16	5	5	1	-	-
Female	6	3	2	4	8	10	12	15	13	3	1	-
White, both sexes	8	7	5	8	15	17	28	20	18	4	1	-
Male	2	4	3	4	7	7	16	5	5	1	-	-
Female	6	3	2	4	8	10	12	15	13	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	49	60	71	87	122	136	167	161	87	34	6	-
Male	28	33	32	43	64	63	73	65	22	10	-	-
Female	21	27	39	44	58	73	94	96	65	24	6	-
White, both sexes	22	27	44	54	78	92	121	117	64	28	5	-
Male	13	15	21	29	39	49	52	50	17	8	-	-
Female	9	12	23	25	39	43	69	67	47	20	5	-
Black, both sexes	26	31	26	32	43	41	44	41	23	6	1	-
Male	14	18	11	14	24	13	20	14	5	2	-	-
Female	12	13	15	18	19	28	24	27	18	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	1	1	2	2	3	-	-	-	-
Male	1	-	-	-	1	1	1	1	-	-	-	-
Female	-	2	1	1	-	1	1	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Massachusetts												
All races, both sexes	34	50	53	63	105	116	184	164	100	40	5	-
Male	20	26	26	34	61	51	75	57	28	13	1	-
Female	14	24	27	29	44	65	109	107	72	27	4	-
White, both sexes	33	47	45	55	90	109	177	159	96	38	5	-
Male	20	25	23	32	54	46	71	57	27	13	1	-
Female	13	22	22	23	36	63	106	102	69	25	4	-
Black, both sexes	1	3	8	8	13	7	5	3	3	2	-	-
Male	-	1	3	2	7	5	3	-	1	-	-	-
Female	1	2	5	6	6	2	2	3	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	2	2	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	2	-	1	2	1	-	-	-
Michigan												
All races, both sexes	36	36	65	45	75	125	149	155	91	26	2	-
Male	15	21	39	20	37	54	64	60	20	7	1	-
Female	21	15	26	25	38	71	85	95	71	19	1	-
White, both sexes	24	19	42	33	57	99	117	131	78	22	2	-
Male	10	11	24	16	28	39	48	54	16	7	1	-
Female	14	8	18	17	29	60	69	77	62	15	1	-
Black, both sexes	12	16	21	12	17	25	30	23	13	4	-	-
Male	5	9	14	4	9	15	15	5	4	-	-	-
Female	7	7	7	8	8	10	15	18	9	4	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	1	1	1	1	-	-	-	-
Male	-	1	-	-	-	-	1	1	-	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Minnesota												
All races, both sexes	14	12	10	20	30	32	48	51	23	9	-	-
Male	6	8	5	8	15	19	24	28	6	-	-	-
Female	8	4	5	12	15	13	24	23	17	9	-	-
White, both sexes	11	10	10	18	29	31	48	48	23	9	-	-
Male	6	8	5	8	14	18	24	26	6	-	-	-
Female	5	2	5	10	15	13	24	22	17	9	-	-
Black, both sexes	2	1	-	2	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	2	1	-	2	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	29	31	36	38	49	65	62	63	49	21	3	-
Male	13	13	14	22	19	32	29	23	10	3	2	-
Female	16	18	22	16	30	33	33	40	39	18	1	-
White, both sexes	13	17	21	17	34	48	37	43	40	16	1	-
Male	8	5	9	12	15	25	19	18	8	3	1	-
Female	5	12	12	5	19	23	18	25	32	13	-	-
Black, both sexes	16	14	15	20	15	17	25	20	9	5	2	-
Male	5	8	5	10	4	7	10	5	2	-	1	-
Female	11	6	10	10	11	10	15	15	7	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Missouri												
All races, both sexes	31	44	34	49	56	88	128	94	84	23	2	-
Male	15	27	15	29	23	39	55	37	23	7	-	-
Female	16	17	19	20	33	49	73	57	61	16	2	-
White, both sexes	20	30	26	42	44	78	111	84	75	19	2	-
Male	9	17	11	25	18	35	45	33	23	6	-	-
Female	11	13	15	17	26	43	66	51	52	13	2	-
Black, both sexes	10	11	8	7	11	9	17	10	9	4	-	-
Male	6	8	4	4	4	4	10	4	-	1	-	-
Female	4	3	4	3	7	5	7	6	9	3	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	-	-	-	1	-	-	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	1	-	-	-	-	-	-
Montana												
All races, both sexes	2	3	6	4	4	6	15	8	13	3	1	-
Male	1	1	-	3	2	6	9	4	2	-	-	-
Female	1	2	6	1	2	-	6	4	11	3	1	-
White, both sexes	1	3	6	3	4	6	15	8	12	3	1	-
Male	-	1	-	2	2	6	9	4	2	-	-	-
Female	1	2	6	1	2	-	6	4	10	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	1	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Nebraska												
All races, both sexes	6	5	6	5	13	18	34	25	8	4	-	-
Male	4	3	3	2	10	6	15	4	3	2	-	-
Female	2	2	3	3	3	12	19	21	5	2	-	-
White, both sexes	4	3	5	5	13	17	31	25	8	4	-	-
Male	3	2	3	2	10	6	15	4	3	2	-	-
Female	1	1	2	3	3	11	16	21	5	2	-	-
Black, both sexes	1	2	1	-	-	1	3	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	1	1	-	-	1	3	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	17	34	28	39	58	69	55	38	16	4	-	1
Male	9	17	18	24	36	30	31	18	2	1	-	1
Female	8	17	10	15	22	39	24	20	14	3	-	-
White, both sexes	16	27	21	34	47	63	49	37	15	4	-	1
Male	9	17	12	20	30	28	28	18	2	1	-	1
Female	7	10	9	14	17	35	21	19	13	3	-	-
Black, both sexes	1	4	3	4	5	4	5	1	1	-	-	-
Male	-	-	2	3	2	2	2	-	-	-	-	-
Female	1	4	1	1	3	2	3	1	1	-	-	-
American Indian 1/, both sexes	-	3	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	3	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	3	1	6	1	1	-	-	-	-	-
Male	-	-	3	1	4	-	1	-	-	-	-	-
Female	-	-	-	-	2	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
New Hampshire												
All races, both sexes	3	7	3	9	4	14	18	14	9	4	-	-
Male	2	2	2	6	2	8	13	4	7	1	-	-
Female	1	5	1	3	2	6	5	10	2	3	-	-
White, both sexes	3	7	3	9	4	14	18	14	9	4	-	-
Male	2	2	2	6	2	8	13	4	7	1	-	-
Female	1	5	1	3	2	6	5	10	2	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	57	71	90	112	174	299	362	352	192	85	11	-
Male	34	40	44	57	92	157	155	125	55	20	2	-
Female	23	31	46	55	82	142	207	227	137	65	9	-
White, both sexes	40	44	63	83	136	237	320	320	182	77	7	-
Male	24	24	30	43	73	126	143	113	53	18	-	-
Female	16	20	33	40	63	111	177	207	129	59	7	-
Black, both sexes	17	22	24	27	31	55	36	29	10	7	4	-
Male	10	11	11	14	17	27	11	10	2	2	2	-
Female	7	11	13	13	14	28	25	19	8	5	2	-
American Indian 1/, both sexes	-	-	-	-	2	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	5	3	2	5	7	6	3	-	-	-	-
Male	-	5	3	-	2	4	1	2	-	-	-	-
Female	-	-	-	2	3	3	5	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
New Mexico												
All races, both sexes	9	5	11	9	18	16	26	17	9	4	1	-
Male	5	3	6	6	7	9	12	4	3	1	-	-
Female	4	2	5	3	11	7	14	13	6	3	1	-
White, both sexes	8	5	7	9	16	16	24	14	9	3	1	-
Male	5	3	4	6	5	9	12	4	3	1	-	-
Female	3	2	3	3	11	7	12	10	6	2	1	-
Black, both sexes	1	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	4	-	1	-	1	3	-	1	-	-
Male	-	-	2	-	1	-	-	-	-	-	-	-
Female	-	-	2	-	-	-	1	3	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	65	113	110	169	236	348	400	377	228	83	17	1
Male	37	53	55	87	129	176	187	159	70	16	4	-
Female	28	60	55	82	107	172	213	218	158	67	13	1
White, both sexes	51	85	88	131	198	297	346	332	209	77	13	1
Male	30	40	46	72	109	156	163	141	63	15	3	-
Female	21	45	42	59	89	141	183	191	146	62	10	1
Black, both sexes	13	26	20	32	36	48	44	37	17	6	4	-
Male	6	13	7	12	18	19	18	14	5	1	1	-
Female	7	13	13	20	18	29	26	23	12	5	3	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	5	2	3	10	8	2	-	-	-
Male	1	-	2	2	2	1	6	4	2	-	-	-
Female	-	1	-	3	-	2	4	4	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
North Carolina												
All races, both sexes	53	58	87	87	141	172	203	152	90	36	14	-
Male	25	30	48	46	57	80	87	62	29	9	6	-
Female	28	28	39	41	84	92	116	90	61	27	8	-
White, both sexes	31	33	54	51	106	140	144	117	69	24	9	-
Male	12	18	30	26	42	70	54	44	23	5	4	-
Female	19	15	24	25	64	70	90	73	46	19	5	-
Black, both sexes	20	25	31	35	33	31	57	32	21	12	5	-
Male	12	12	18	20	13	10	33	15	6	4	2	-
Female	8	13	13	15	20	21	24	17	15	8	3	-
American Indian 1/, both sexes	2	-	2	1	1	1	2	2	-	-	-	-
Male	1	-	-	-	1	-	-	2	-	-	-	-
Female	1	-	2	1	-	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	2	1	2	4	10	3	2	4	2	-	-
Male	-	2	-	-	1	7	2	-	2	-	-	-
Female	-	-	1	2	3	3	1	2	2	2	-	-
White, both sexes	-	2	1	2	4	10	2	2	4	2	-	-
Male	-	2	-	-	1	7	1	-	2	-	-	-
Female	-	-	1	2	3	3	1	2	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Ohio												
All races, both sexes	59	76	104	125	150	223	253	203	140	30	2	-
Male	22	43	58	49	77	111	106	74	38	7	1	-
Female	37	33	46	76	73	112	147	129	102	23	1	-
White, both sexes	39	54	82	98	124	192	225	179	123	25	2	-
Male	10	30	43	42	65	101	91	66	30	6	1	-
Female	29	24	39	56	59	91	134	113	93	19	1	-
Black, both sexes	20	22	22	26	25	30	26	23	17	5	-	-
Male	12	13	15	7	12	10	14	8	8	1	-	-
Female	8	9	7	19	13	20	12	15	9	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	1	-	1	1	-	-	-	-
Oklahoma												
All races, both sexes	9	27	39	38	54	54	56	46	29	12	3	-
Male	5	17	23	15	34	23	30	15	6	3	-	-
Female	4	10	16	23	20	31	26	31	23	9	3	-
White, both sexes	6	24	34	33	45	46	48	42	28	11	3	-
Male	3	16	19	13	27	18	28	14	5	3	-	-
Female	3	8	15	20	18	28	20	28	23	8	3	-
Black, both sexes	1	3	2	4	8	4	7	3	1	1	-	-
Male	1	1	1	2	6	2	2	-	1	-	-	-
Female	-	2	1	2	2	2	5	3	-	1	-	-
American Indian 1/, both sexes	2	-	3	1	1	3	1	1	-	-	-	-
Male	1	-	3	-	1	2	-	1	-	-	-	-
Female	1	-	-	1	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Oregon												
All races, both sexes	12	8	19	17	14	20	36	28	16	8	-	-
Male	7	3	10	7	8	6	17	16	6	4	-	-
Female	5	5	9	10	6	14	19	12	10	4	-	-
White, both sexes	11	8	18	16	14	18	33	27	16	8	-	-
Male	7	3	9	6	8	6	15	16	6	4	-	-
Female	4	5	9	10	6	12	18	11	10	4	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	-	1	1	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	2	1	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Pennsylvania												
All races, both sexes	81	109	128	162	264	382	497	394	222	89	9	-
Male	50	57	68	84	125	178	229	159	71	16	-	-
Female	31	52	60	78	139	204	268	235	151	73	9	-
White, both sexes	56	79	92	124	221	323	451	359	198	82	8	-
Male	34	36	49	64	107	148	215	150	62	13	-	-
Female	22	43	43	60	114	175	236	209	136	69	8	-
Black, both sexes	23	29	33	33	39	56	46	33	23	7	1	-
Male	15	20	18	18	15	28	14	9	8	3	-	-
Female	8	9	15	15	24	28	32	24	15	4	1	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	3	5	3	3	-	2	1	-	-	-
Male	1	1	1	2	3	2	-	-	1	-	-	-
Female	1	-	2	3	-	1	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Rhode Island												
All races, both sexes	-	5	1	4	9	12	11	22	12	3	-	-
Male	-	4	-	3	4	7	6	9	2	1	-	-
Female	-	1	1	1	5	5	5	13	10	2	-	-
White, both sexes	-	4	1	3	9	10	10	21	12	3	-	-
Male	-	3	-	2	4	6	6	9	2	1	-	-
Female	-	1	1	1	5	4	4	12	10	2	-	-
Black, both sexes	-	1	-	-	-	1	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	1	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
South Carolina												
All races, both sexes	31	46	49	62	69	86	105	75	32	18	4	-
Male	15	22	24	33	36	34	42	28	8	1	-	-
Female	16	24	25	29	33	52	63	47	24	17	4	-
White, both sexes	19	28	28	38	49	57	67	59	22	11	1	-
Male	11	11	15	21	29	22	23	20	4	-	-	-
Female	8	17	13	17	20	35	44	39	18	11	1	-
Black, both sexes	12	18	20	24	19	28	38	15	10	7	3	-
Male	4	11	8	12	7	12	19	7	4	1	-	-
Female	8	7	12	12	12	16	19	8	6	6	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	1	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
South Dakota												
All races, both sexes	2	3	3	4	3	5	4	5	5	1	-	-
Male	1	1	3	1	1	2	-	4	2	-	-	-
Female	1	2	-	3	2	3	4	1	3	1	-	-
White, both sexes	2	3	2	4	2	4	4	4	5	1	-	-
Male	1	1	2	1	1	2	-	3	2	-	-	-
Female	1	2	-	3	1	2	4	1	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	1	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	28	23	28	37	56	80	90	73	48	13	-	-
Male	17	14	15	11	26	34	39	25	11	3	-	-
Female	11	9	13	26	30	46	51	48	37	10	-	-
White, both sexes	21	21	22	32	44	66	74	56	37	11	-	-
Male	14	12	12	9	20	29	31	20	9	2	-	-
Female	7	9	10	23	24	37	43	36	28	9	-	-
Black, both sexes	7	2	6	5	12	14	16	17	11	2	-	-
Male	3	2	3	2	6	5	8	5	2	1	-	-
Female	4	-	3	3	6	9	8	12	9	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Texas												
All races, both sexes	115	141	152	199	261	331	376	310	181	74	11	-
Male	49	73	75	115	138	151	163	107	57	29	-	-
Female	66	68	77	84	123	180	213	203	124	45	11	-
White, both sexes	88	105	116	158	206	280	330	269	159	65	8	-
Male	42	57	57	91	109	129	141	94	50	26	-	-
Female	46	48	59	67	97	151	189	175	109	39	8	-
Black, both sexes	26	36	34	39	54	47	39	37	22	8	3	-
Male	7	16	16	22	28	22	17	13	7	2	-	-
Female	19	20	18	17	26	25	22	24	15	6	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	2	1	4	7	3	-	1	-	-
Male	-	-	2	2	1	-	5	-	-	1	-	-
Female	1	-	-	-	-	4	2	3	-	-	-	-
Utah												
All races, both sexes	1	8	6	9	14	23	20	11	8	2	-	-
Male	-	4	1	2	7	11	8	5	4	2	-	-
Female	1	4	5	7	7	12	12	6	4	-	-	-
White, both sexes	1	8	6	9	14	21	19	11	7	2	-	-
Male	-	4	1	2	7	9	7	5	3	2	-	-
Female	1	4	5	7	7	12	12	6	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	2	-	-	1	-	-	-
Male	-	-	-	-	-	2	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Vermont												
All races, both sexes	1	-	-	-	4	5	7	6	6	1	-	-
Male	-	-	-	-	2	4	4	2	1	1	-	-
Female	1	-	-	-	2	1	3	4	5	-	-	-
White, both sexes	1	-	-	-	4	5	7	6	6	1	-	-
Male	-	-	-	-	2	4	4	2	1	1	-	-
Female	1	-	-	-	2	1	3	4	5	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	46	61	49	96	133	185	209	185	112	29	6	1
Male	25	35	25	50	64	85	88	76	40	10	2	-
Female	21	26	24	46	69	100	121	109	72	19	4	1
White, both sexes	21	39	31	60	86	135	159	141	78	22	4	-
Male	12	20	16	33	39	65	67	64	32	10	1	-
Female	9	19	15	27	47	70	92	77	46	12	3	-
Black, both sexes	23	22	17	35	46	47	46	42	31	7	2	1
Male	12	15	8	16	25	18	20	10	8	-	1	-
Female	11	7	9	19	21	29	26	32	23	7	1	1
American Indian 1/, both sexes	2	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	3	4	2	3	-	-	-
Male	-	-	1	1	-	2	1	2	-	-	-	-
Female	-	-	-	-	-	1	3	-	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Washington												
All races, both sexes	11	15	22	13	29	59	50	35	31	13	3	-
Male	8	7	9	4	16	30	21	16	8	3	-	-
Female	3	8	13	9	13	29	29	19	23	10	3	-
White, both sexes	11	12	18	11	27	56	46	33	28	11	2	-
Male	8	6	7	3	15	27	18	15	7	3	-	-
Female	3	6	11	8	12	29	28	18	21	8	2	-
Black, both sexes	-	1	1	1	-	1	1	-	1	-	-	-
Male	-	1	1	-	-	1	1	-	1	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	2	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	2	-	-	-	-	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	3	1	2	2	3	1	2	1	1	-
Male	-	-	1	1	1	2	2	1	-	-	-	-
Female	-	-	2	-	1	-	1	-	2	1	1	-
West Virginia												
All races, both sexes	15	16	26	22	48	50	56	42	21	4	1	-
Male	11	4	16	8	25	29	27	15	8	1	-	-
Female	4	12	10	14	23	21	29	27	13	3	1	-
White, both sexes	15	16	25	22	45	48	55	41	21	4	1	-
Male	11	4	16	8	23	28	27	15	8	1	-	-
Female	4	12	9	14	22	20	28	26	13	3	1	-
Black, both sexes	-	-	1	-	3	2	1	1	-	-	-	-
Male	-	-	-	-	2	1	-	-	-	-	-	-
Female	-	-	1	-	1	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Wisconsin												
All races, both sexes	22	17	23	24	41	67	76	61	35	19	4	-
Male	9	10	5	10	23	33	38	23	10	4	-	-
Female	13	7	18	14	18	34	38	38	25	15	4	-
White, both sexes	18	14	21	21	36	62	73	61	35	19	3	-
Male	8	9	5	7	20	31	36	23	10	4	-	-
Female	10	5	16	14	16	31	37	38	25	15	3	-
Black, both sexes	4	1	1	2	2	5	2	-	-	-	1	-
Male	1	1	-	2	1	2	1	-	-	-	-	-
Female	3	-	1	-	1	3	1	-	-	-	1	-
American Indian 1/, both sexes	-	2	1	1	2	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	2	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	3	3	2	2	4	9	7	6	1	-	-
Male	1	2	2	1	1	3	5	1	2	1	-	-
Female	-	1	1	1	1	1	4	6	4	-	-	-
White, both sexes	1	3	3	2	2	4	8	7	6	1	-	-
Male	1	2	2	1	1	3	4	1	2	1	-	-
Female	-	1	1	1	1	1	4	6	4	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
United States												
All races, both sexes	1,617	13	5	2	4	1	7	8	15	20	30	59
Male	792	8	3	1	-	-	5	5	9	10	16	32
Female	825	5	2	1	4	1	2	3	6	10	14	27
White, both sexes	1,330	11	3	1	2	-	4	5	9	17	15	40
Male	643	7	1	-	-	-	2	3	6	8	9	23
Female	687	4	2	1	2	-	2	2	3	9	6	17
Black, both sexes	235	2	2	-	1	1	2	3	5	2	12	17
Male	121	1	2	-	-	-	2	2	3	1	5	8
Female	114	1	-	-	1	1	-	1	2	1	7	9
American Indian 1/, both sexes	16	-	-	-	1	-	-	-	1	1	1	2
Male	7	-	-	-	-	-	-	-	-	1	-	1
Female	9	-	-	-	1	-	-	-	1	-	1	1
Asian or Pacific Islander 2/, both sexes	36	-	-	1	-	-	1	-	-	-	2	-
Male	21	-	-	1	-	-	1	-	-	-	2	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	41	-	-	-	1	-	-	-	1	-	-	3
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	23	-	-	-	1	-	-	-	1	-	-	2
White, both sexes	26	-	-	-	1	-	-	-	-	-	-	2
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	1	-	-	-	-	-	-	1
Black, both sexes	15	-	-	-	-	-	-	-	1	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Alaska												
All races, both sexes	5	-	-	-	-	-	-	-	-	1	1	1
Male	3	-	-	-	-	-	-	-	-	1	-	1
Female	2	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	44	-	-	-	-	-	-	-	-	1	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	40	-	-	-	-	-	-	-	-	1	-	1
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Arkansas												
All races, both sexes	17	-	-	-	-	-	-	1	1	-	-	-
Male	8	-	-	-	-	-	-	1	1	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	1	1	-	-	-
Male	8	-	-	-	-	-	-	1	1	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	124	-	-	-	-	-	1	-	1	-	4	8
Male	61	-	-	-	-	-	1	-	1	-	2	4
Female	63	-	-	-	-	-	-	-	-	-	2	4
White, both sexes	98	-	-	-	-	-	1	-	1	-	1	7
Male	45	-	-	-	-	-	1	-	1	-	-	4
Female	53	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	14	-	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	2	-
Male	9	-	-	-	-	-	-	-	-	-	2	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Colorado												
All races, both sexes	21	-	-	-	-	-	-	1	-	-	1	-
Male	13	-	-	-	-	-	-	1	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	1	-	-	1	-
Male	12	-	-	-	-	-	-	1	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	26	-	-	-	-	-	-	-	-	1	-	3
Male	16	-	-	-	-	-	-	-	-	1	-	3
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	1	-	3
Male	15	-	-	-	-	-	-	-	-	1	-	3
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Delaware												
All races, both sexes	6	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Florida												
All races, both sexes	115	1	-	-	-	1	-	1	1	3	2	5
Male	66	-	-	-	-	-	-	1	1	1	1	3
Female	49	1	-	-	-	1	-	-	-	2	1	2
White, both sexes	92	-	-	-	-	-	-	-	-	3	-	2
Male	52	-	-	-	-	-	-	-	-	1	-	1
Female	40	-	-	-	-	-	-	-	-	2	-	1
Black, both sexes	21	1	-	-	-	1	-	1	1	-	2	2
Male	13	-	-	-	-	-	-	1	1	-	1	2
Female	8	1	-	-	-	1	-	-	-	-	1	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	42	-	-	-	-	-	-	-	-	1	-	4
Male	20	-	-	-	-	-	-	-	-	1	-	3
Female	22	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	31	-	-	-	-	-	-	-	-	1	-	2
Male	16	-	-	-	-	-	-	-	-	1	-	2
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Hawaii												
All races, both sexes	15	-	-	-	-	-	1	-	-	-	1	-
Male	7	-	-	-	-	-	1	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	1	-	-	-	-	-
Male	5	-	-	-	-	-	1	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	11	1	-	1	-	-	-	-	-	-	-	-
Male	6	1	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	11	1	-	1	-	-	-	-	-	-	-	-
Male	6	1	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Illinois												
All races, both sexes	74	-	1	-	-	-	1	1	-	2	3	3
Male	39	-	1	-	-	-	-	-	-	1	1	1
Female	35	-	-	-	-	-	1	1	-	1	2	2
White, both sexes	61	-	-	-	-	-	1	1	-	2	2	2
Male	31	-	-	-	-	-	-	-	-	1	1	1
Female	30	-	-	-	-	-	1	1	-	1	1	1
Black, both sexes	11	-	1	-	-	-	-	-	-	-	1	1
Male	6	-	1	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	37	1	-	-	-	-	-	-	-	-	-	1
Male	26	1	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	33	1	-	-	-	-	-	-	-	-	-	1
Male	23	1	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Iowa												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	10	-	-	-	-	-	-	1	-	-	-	1
Male	4	-	-	-	-	-	-	1	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	1	-	-	-	1
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Kentucky												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	34	-	-	-	-	-	-	-	-	1	2	3
Male	18	-	-	-	-	-	-	-	-	1	2	-
Female	16	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	22	-	-	-	-	-	-	-	-	-	1	2
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	12	-	-	-	-	-	-	-	-	1	1	1
Male	7	-	-	-	-	-	-	-	-	1	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Maine												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	31	-	-	-	1	-	-	-	-	2	-	3
Male	18	-	-	-	-	-	-	-	-	1	-	3
Female	13	-	-	-	1	-	-	-	-	1	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	2	-	1
Male	8	-	-	-	-	-	-	-	-	1	-	1
Female	7	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	15	-	-	-	1	-	-	-	-	-	-	2
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	6	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Massachusetts												
All races, both sexes	39	-	-	-	-	-	-	-	1	2	1	-
Male	21	-	-	-	-	-	-	-	1	-	1	-
Female	18	-	-	-	-	-	-	-	-	2	-	-
White, both sexes	37	-	-	-	-	-	-	-	1	2	1	-
Male	19	-	-	-	-	-	-	-	1	-	1	-
Female	18	-	-	-	-	-	-	-	-	2	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	46	-	-	-	-	-	1	1	1	-	2	1
Male	17	-	-	-	-	-	-	1	-	-	1	-
Female	29	-	-	-	-	-	1	-	1	-	1	1
White, both sexes	39	-	-	-	-	-	1	1	1	-	2	1
Male	14	-	-	-	-	-	-	1	-	-	1	-
Female	25	-	-	-	-	-	1	-	1	-	1	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Minnesota												
All races, both sexes	30	2	1	-	-	-	-	-	1	-	-	-
Male	11	1	-	-	-	-	-	-	-	-	-	-
Female	19	1	1	-	-	-	-	-	1	-	-	-
White, both sexes	29	2	1	-	-	-	-	-	1	-	-	-
Male	11	1	-	-	-	-	-	-	-	-	-	-
Female	18	1	1	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	21	-	1	-	-	-	-	-	-	-	-	1
Male	11	-	1	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	7	-	1	-	-	-	-	-	-	-	-	-
Male	5	-	1	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Missouri												
All races, both sexes	42	-	-	-	-	-	-	2	1	-	2	3
Male	21	-	-	-	-	-	-	-	1	-	2	1
Female	21	-	-	-	-	-	-	2	-	-	-	2
White, both sexes	35	-	-	-	-	-	-	1	1	-	2	2
Male	19	-	-	-	-	-	-	-	1	-	2	1
Female	16	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	7	-	-	-	-	-	-	1	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	3	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Nebraska												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	21	1	-	-	1	-	-	-	1	-	-	1
Male	10	-	-	-	-	-	-	-	1	-	-	-
Female	11	1	-	-	1	-	-	-	-	-	-	1
White, both sexes	18	1	-	-	1	-	-	-	1	-	-	-
Male	9	-	-	-	-	-	-	-	1	-	-	-
Female	9	1	-	-	1	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
New Hampshire												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	35	1	-	-	-	-	-	-	-	1	-	1
Male	11	1	-	-	-	-	-	-	-	-	-	1
Female	24	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	32	1	-	-	-	-	-	-	-	1	-	1
Male	10	1	-	-	-	-	-	-	-	-	-	1
Female	22	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
New Mexico												
All races, both sexes	8	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	92	-	-	-	-	-	-	-	1	-	1	4
Male	44	-	-	-	-	-	-	-	-	-	-	4
Female	48	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	75	-	-	-	-	-	-	-	-	-	1	3
Male	39	-	-	-	-	-	-	-	-	-	-	3
Female	36	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	15	-	-	-	-	-	-	-	1	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
North Carolina												
All races, both sexes	55	1	-	-	-	-	1	-	2	-	2	-
Male	27	1	-	-	-	-	1	-	1	-	-	-
Female	28	-	-	-	-	-	-	-	1	-	2	-
White, both sexes	44	1	-	-	-	-	-	-	2	-	-	-
Male	22	1	-	-	-	-	-	-	1	-	-	-
Female	22	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	10	-	-	-	-	-	1	-	-	-	2	-
Male	5	-	-	-	-	-	1	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Ohio												
All races, both sexes	77	3	-	-	-	-	-	-	1	-	2	1
Male	42	1	-	-	-	-	-	-	1	-	-	1
Female	35	2	-	-	-	-	-	-	-	-	2	-
White, both sexes	60	3	-	-	-	-	-	-	-	-	-	1
Male	31	1	-	-	-	-	-	-	-	-	-	1
Female	29	2	-	-	-	-	-	-	-	-	-	-
Black, both sexes	17	-	-	-	-	-	-	-	1	-	2	-
Male	11	-	-	-	-	-	-	-	1	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	18	-	-	-	-	-	-	-	1	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Oregon												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	100	-	-	-	-	-	-	-	-	-	-	2
Male	45	-	-	-	-	-	-	-	-	-	-	2
Female	55	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	89	-	-	-	-	-	-	-	-	-	-	1
Male	39	-	-	-	-	-	-	-	-	-	-	1
Female	50	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Rhode Island												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	27	-	-	-	-	-	1	-	-	-	1	-
Male	11	-	-	-	-	-	1	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	10	-	-	-	-	-	1	-	-	-	-	-
Male	5	-	-	-	-	-	1	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
South Dakota												
All races, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	48	1	-	-	-	-	-	-	-	1	1	2
Male	23	1	-	-	-	-	-	-	-	1	1	-
Female	25	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	41	1	-	-	-	-	-	-	-	1	-	2
Male	18	1	-	-	-	-	-	-	-	1	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Texas												
All races, both sexes	88	-	1	1	-	-	1	-	-	2	3	1
Male	48	-	1	1	-	-	1	-	-	1	3	1
Female	40	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	75	-	1	-	-	-	1	-	-	1	2	-
Male	43	-	1	-	-	-	1	-	-	1	2	-
Female	32	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	1	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	1
Female	6	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Vermont												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	36	-	-	-	-	-	-	-	1	-	1	2
Male	17	-	-	-	-	-	-	-	1	-	1	1
Female	19	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	13	-	-	-	-	-	-	-	1	-	1	1
Male	7	-	-	-	-	-	-	-	1	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Washington												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Wisconsin												
All races, both sexes	23	-	-	-	-	-	-	-	-	1	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	1	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
United States												
All races, both sexes	65	109	116	107	161	232	285	221	112	42	3	-
Male	34	63	55	69	89	106	124	105	44	13	1	-
Female	31	46	61	38	72	126	161	116	68	29	2	-
White, both sexes	45	88	92	86	130	191	253	199	101	35	3	-
Male	26	51	43	56	66	87	107	95	40	12	1	-
Female	19	37	49	30	64	104	146	104	61	23	2	-
Black, both sexes	18	18	21	15	23	37	27	14	9	6	-	-
Male	8	10	12	11	17	15	13	7	3	1	-	-
Female	10	8	9	4	6	22	14	7	6	5	-	-
American Indian 1/, both sexes	1	2	-	2	1	1	2	1	-	-	-	-
Male	-	1	-	-	1	1	2	-	-	-	-	-
Female	1	1	-	2	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	3	4	7	3	3	7	2	1	-	-
Male	-	1	-	2	5	3	2	3	1	-	-	-
Female	1	-	3	2	2	-	1	4	1	1	-	-
Alabama												
All races, both sexes	4	3	4	4	3	4	6	5	2	1	-	-
Male	4	2	1	3	1	2	1	2	1	-	-	-
Female	-	1	3	1	2	2	5	3	1	1	-	-
White, both sexes	2	2	3	3	2	3	4	3	1	-	-	-
Male	2	2	1	2	-	2	-	1	1	-	-	-
Female	-	-	2	1	2	1	4	2	-	-	-	-
Black, both sexes	2	1	1	1	1	1	2	2	1	1	-	-
Male	2	-	-	1	1	-	1	1	-	-	-	-
Female	-	1	1	-	-	1	1	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Alaska												
All races, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	3	2	2	6	5	12	8	1	-	1	-
Male	1	1	1	1	5	-	7	3	1	-	-	-
Female	1	2	1	1	1	5	5	5	-	-	1	-
White, both sexes	1	3	2	2	6	4	10	8	1	-	1	-
Male	1	1	1	1	5	-	5	3	1	-	-	-
Female	-	2	1	1	1	4	5	5	-	-	1	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Arkansas												
All races, both sexes	-	1	1	2	3	3	2	-	1	2	-	-
Male	-	-	1	2	1	1	-	-	-	1	-	-
Female	-	1	-	-	2	2	2	-	1	1	-	-
White, both sexes	-	1	1	2	3	3	2	-	1	2	-	-
Male	-	-	1	2	1	1	-	-	-	1	-	-
Female	-	1	-	-	2	2	2	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	3	8	9	7	8	15	26	18	12	4	-	-
Male	-	4	3	3	7	9	11	11	4	1	-	-
Female	3	4	6	4	1	6	15	7	8	3	-	-
White, both sexes	2	5	8	5	6	12	23	12	11	4	-	-
Male	-	2	3	2	5	6	9	7	4	1	-	-
Female	2	3	5	3	1	6	14	5	7	3	-	-
Black, both sexes	1	2	-	1	-	2	3	2	1	-	-	-
Male	-	1	-	-	-	2	2	1	-	-	-	-
Female	1	1	-	1	-	-	1	1	1	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	2	1	-	4	-	-	-	-
Male	-	-	-	1	2	1	-	3	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Colorado												
All races, both sexes	1	2	1	2	2	3	2	3	1	2	-	-
Male	1	1	-	2	1	1	2	2	-	1	-	-
Female	-	1	1	-	1	2	-	1	1	1	-	-
White, both sexes	1	2	1	2	2	3	1	2	1	2	-	-
Male	1	1	-	2	1	1	1	2	-	1	-	-
Female	-	1	1	-	1	2	-	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	1	2	3	2	3	3	3	4	-	1	-
Male	-	1	-	1	2	2	1	2	2	-	1	-
Female	-	-	2	2	-	1	2	1	2	-	-	-
White, both sexes	-	1	2	3	2	2	3	3	3	-	1	-
Male	-	1	-	1	2	1	1	2	2	-	1	-
Female	-	-	2	2	-	1	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Delaware												
All races, both sexes	-	-	-	-	-	1	1	2	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	1	-	-	-
White, both sexes	-	-	-	-	-	1	1	2	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	1	-	-	2	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Florida												
All races, both sexes	5	9	5	11	15	21	15	9	5	6	-	-
Male	3	6	3	10	9	11	7	5	2	3	-	-
Female	2	3	2	1	6	10	8	4	3	3	-	-
White, both sexes	3	7	5	10	11	18	15	8	5	5	-	-
Male	1	5	3	9	6	10	7	5	2	2	-	-
Female	2	2	2	1	5	8	8	3	3	3	-	-
Black, both sexes	2	2	-	1	4	2	-	1	-	1	-	-
Male	2	1	-	1	3	-	-	-	-	1	-	-
Female	-	1	-	-	1	2	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	5	2	4	4	3	6	8	3	2	-	-	-
Male	2	1	3	3	1	2	2	2	-	-	-	-
Female	3	1	1	1	2	4	6	1	2	-	-	-
White, both sexes	3	1	2	3	3	6	6	2	2	-	-	-
Male	2	1	2	2	1	2	2	1	-	-	-	-
Female	1	-	-	1	2	4	4	1	2	-	-	-
Black, both sexes	2	1	2	1	-	-	2	1	-	-	-	-
Male	-	-	1	1	-	-	-	1	-	-	-	-
Female	2	1	1	-	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Hawaii												
All races, both sexes	-	-	-	1	4	2	1	3	1	1	-	-
Male	-	-	-	-	3	1	1	1	-	-	-	-
Female	-	-	-	1	1	1	-	2	1	1	-	-
White, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	4	1	-	2	1	1	-	-
Male	-	-	-	-	3	1	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	2	1	1	-	-
Idaho												
All races, both sexes	-	1	-	2	1	1	3	-	1	-	-	-
Male	-	1	-	1	1	-	2	-	-	-	-	-
Female	-	-	-	1	-	1	1	-	1	-	-	-
White, both sexes	-	1	-	2	1	1	3	-	1	-	-	-
Male	-	1	-	1	1	-	2	-	-	-	-	-
Female	-	-	-	1	-	1	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Illinois												
All races, both sexes	1	3	5	4	4	10	14	12	9	1	-	-
Male	1	2	4	4	2	5	6	8	3	-	-	-
Female	-	1	1	-	2	5	8	4	6	1	-	-
White, both sexes	1	1	4	1	4	10	12	11	8	1	-	-
Male	1	-	3	1	2	5	6	7	3	-	-	-
Female	-	1	1	-	2	5	6	4	5	1	-	-
Black, both sexes	-	1	1	2	-	-	2	1	1	-	-	-
Male	-	1	1	2	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	3	2	5	7	4	7	3	2	2	-	-
Male	-	2	1	5	6	2	5	-	2	1	-	-
Female	-	1	1	-	1	2	2	3	-	1	-	-
White, both sexes	-	2	2	5	5	4	6	3	2	2	-	-
Male	-	1	1	5	5	2	4	-	2	1	-	-
Female	-	1	1	-	-	2	2	3	-	1	-	-
Black, both sexes	-	1	-	-	1	-	1	-	-	-	-	-
Male	-	1	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Iowa												
All races, both sexes	-	2	-	2	1	3	2	2	1	-	-	-
Male	-	1	-	1	-	1	1	1	-	-	-	-
Female	-	1	-	1	1	2	1	1	1	-	-	-
White, both sexes	-	2	-	2	1	2	2	2	1	-	-	-
Male	-	1	-	1	-	1	1	1	-	-	-	-
Female	-	1	-	1	1	1	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	1	-	1	1	1	3	-	-	1	-	-
Male	-	1	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	1	1	1	1	-	-	1	-	-
White, both sexes	-	1	-	1	1	-	3	-	-	1	-	-
Male	-	1	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	1	1	-	1	-	-	1	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Kentucky												
All races, both sexes	-	3	4	2	1	7	4	4	1	1	-	-
Male	-	3	1	-	-	3	2	2	-	-	-	-
Female	-	-	3	2	1	4	2	2	1	1	-	-
White, both sexes	-	3	2	2	1	7	4	4	1	1	-	-
Male	-	3	-	-	-	3	2	2	-	-	-	-
Female	-	-	2	2	1	4	2	2	1	1	-	-
Black, both sexes	-	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2	4	4	2	3	5	4	-	2	1	1	-
Male	-	2	4	1	2	2	3	-	1	-	-	-
Female	2	2	-	1	1	3	1	-	1	1	1	-
White, both sexes	1	1	1	1	3	4	4	-	2	1	1	-
Male	-	-	1	1	2	2	3	-	1	-	-	-
Female	1	1	-	-	1	2	1	-	1	1	1	-
Black, both sexes	1	3	3	1	-	1	-	-	-	-	-	-
Male	-	2	3	-	-	-	-	-	-	-	-	-
Female	1	1	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Maine												
All races, both sexes	-	-	1	-	3	-	-	1	1	-	-	-
Male	-	-	1	-	2	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
White, both sexes	-	-	1	-	3	-	-	1	1	-	-	-
Male	-	-	1	-	2	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	4	-	1	2	3	5	2	4	2	2	-	-
Male	2	-	-	2	3	3	1	2	1	-	-	-
Female	2	-	1	-	-	2	1	2	1	2	-	-
White, both sexes	2	-	1	1	1	2	1	4	-	-	-	-
Male	1	-	-	1	1	1	-	2	-	-	-	-
Female	1	-	1	-	-	1	1	2	-	-	-	-
Black, both sexes	2	-	-	1	2	2	1	-	2	2	-	-
Male	1	-	-	1	2	1	1	-	1	-	-	-
Female	1	-	-	-	-	1	-	-	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Massachusetts												
All races, both sexes	-	1	4	2	7	6	5	4	5	1	-	-
Male	-	1	2	1	3	5	4	1	2	-	-	-
Female	-	-	2	1	4	1	1	3	3	1	-	-
White, both sexes	-	1	3	2	7	6	5	4	4	1	-	-
Male	-	1	1	1	3	5	4	1	1	-	-	-
Female	-	-	2	1	4	1	1	3	3	1	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2	2	6	1	2	9	9	6	2	1	-	-
Male	1	1	1	1	-	4	4	2	1	-	-	-
Female	1	1	5	-	2	5	5	4	1	1	-	-
White, both sexes	2	2	3	1	2	7	7	6	2	1	-	-
Male	1	1	-	1	-	3	3	2	1	-	-	-
Female	1	1	3	-	2	4	4	4	1	1	-	-
Black, both sexes	-	-	2	-	-	2	2	-	-	-	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	-	-	1	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Minnesota												
All races, both sexes	2	3	-	1	4	3	6	7	-	-	-	-
Male	2	-	-	-	1	2	3	2	-	-	-	-
Female	-	3	-	1	3	1	3	5	-	-	-	-
White, both sexes	2	3	-	-	4	3	6	7	-	-	-	-
Male	2	-	-	-	1	2	3	2	-	-	-	-
Female	-	3	-	-	3	1	3	5	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	1	2	3	4	5	1	1	2	-	-	-
Male	-	1	1	3	1	2	-	1	1	-	-	-
Female	-	-	1	-	3	3	1	-	1	-	-	-
White, both sexes	-	-	2	1	3	4	1	1	1	-	-	-
Male	-	-	1	1	1	1	-	1	1	-	-	-
Female	-	-	1	-	2	3	1	-	-	-	-	-
Black, both sexes	-	1	-	2	1	1	-	-	1	-	-	-
Male	-	1	-	2	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Missouri												
All races, both sexes	1	3	2	4	3	6	5	7	2	1	-	-
Male	1	3	-	-	2	1	2	5	2	1	-	-
Female	-	-	2	4	1	5	3	2	-	-	-	-
White, both sexes	1	3	1	4	2	5	5	5	2	1	-	-
Male	1	3	-	-	1	1	2	4	2	1	-	-
Female	-	-	1	4	1	4	3	1	-	-	-	-
Black, both sexes	-	-	1	-	1	1	-	2	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	1	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Nebraska												
All races, both sexes	-	-	-	-	-	3	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	1	1	-	-	-
Female	-	-	-	-	-	3	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	3	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	1	1	-	-	-
Female	-	-	-	-	-	3	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	3	1	1	4	1	5	1	-	-	-	-
Male	1	2	-	1	2	-	2	1	-	-	-	-
Female	-	1	1	-	2	1	3	-	-	-	-	-
White, both sexes	1	3	-	1	4	1	4	1	-	-	-	-
Male	1	2	-	1	2	-	1	1	-	-	-	-
Female	-	1	-	-	2	1	3	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
New Hampshire												
All races, both sexes	1	-	-	1	2	1	2	2	3	-	-	-
Male	-	-	-	-	1	-	1	1	3	-	-	-
Female	1	-	-	1	1	1	1	1	-	-	-	-
White, both sexes	1	-	-	1	2	1	2	2	3	-	-	-
Male	-	-	-	-	1	-	1	1	3	-	-	-
Female	1	-	-	1	1	1	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	2	2	2	4	6	6	3	6	-	-	-
Male	-	-	-	1	2	1	2	-	3	-	-	-
Female	1	2	2	1	2	5	4	3	3	-	-	-
White, both sexes	-	2	2	2	4	5	6	3	5	-	-	-
Male	-	-	-	1	2	1	2	-	2	-	-	-
Female	-	2	2	1	2	4	4	3	3	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
New Mexico												
All races, both sexes	2	-	-	-	-	2	1	2	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	1	1	2	-	-	-	-
White, both sexes	2	-	-	-	-	2	1	2	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	1	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	6	5	8	4	8	14	24	9	6	2	-	-
Male	2	3	4	2	3	6	8	6	4	2	-	-
Female	4	2	4	2	5	8	16	3	2	-	-	-
White, both sexes	3	4	6	3	6	11	22	8	6	2	-	-
Male	2	3	3	2	2	5	8	5	4	2	-	-
Female	1	1	3	1	4	6	14	3	2	-	-	-
Black, both sexes	3	1	1	1	2	3	1	1	-	-	-	-
Male	-	-	1	-	1	1	-	1	-	-	-	-
Female	3	1	-	1	1	2	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
North Carolina												
All races, both sexes	-	6	5	8	2	8	10	7	3	-	-	-
Male	-	4	3	3	1	5	5	2	1	-	-	-
Female	-	2	2	5	1	3	5	5	2	-	-	-
White, both sexes	-	5	4	6	2	5	9	7	3	-	-	-
Male	-	3	2	3	1	3	5	2	1	-	-	-
Female	-	2	2	3	1	2	4	5	2	-	-	-
Black, both sexes	-	1	1	1	-	3	1	-	-	-	-	-
Male	-	1	1	-	-	2	-	-	-	-	-	-
Female	-	-	-	1	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Ohio												
All races, both sexes	1	5	4	4	8	16	12	12	5	3	-	-
Male	1	4	1	3	6	7	7	9	-	1	-	-
Female	-	1	3	1	2	9	5	3	5	2	-	-
White, both sexes	1	4	3	3	4	12	9	12	5	3	-	-
Male	1	4	1	2	3	4	4	9	-	1	-	-
Female	-	-	2	1	1	8	5	3	5	2	-	-
Black, both sexes	-	1	1	1	4	4	3	-	-	-	-	-
Male	-	-	-	1	3	3	3	-	-	-	-	-
Female	-	1	1	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	3	-	1	1	3	6	1	2	-	-
Male	-	-	1	-	1	-	1	4	1	-	-	-
Female	-	-	2	-	-	1	2	2	-	2	-	-
White, both sexes	-	-	3	-	1	1	3	6	-	2	-	-
Male	-	-	1	-	1	-	1	4	-	-	-	-
Female	-	-	2	-	-	1	2	2	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Oregon												
All races, both sexes	1	1	2	1	1	3	5	5	2	2	-	-
Male	1	-	2	-	1	2	3	1	1	-	-	-
Female	-	1	-	1	-	1	2	4	1	2	-	-
White, both sexes	1	1	2	1	1	3	4	5	2	2	-	-
Male	1	-	2	-	1	2	2	1	1	-	-	-
Female	-	1	-	1	-	1	2	4	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	2	4	9	2	17	16	19	19	9	1	-	-
Male	1	3	4	1	8	7	9	7	2	1	-	-
Female	1	1	5	1	9	9	10	12	7	-	-	-
White, both sexes	1	3	9	2	13	14	18	18	9	1	-	-
Male	1	2	4	1	5	6	9	7	2	1	-	-
Female	-	1	5	1	8	8	9	11	7	-	-	-
Black, both sexes	1	1	-	-	4	2	1	1	-	-	-	-
Male	-	1	-	-	3	1	-	-	-	-	-	-
Female	1	-	-	-	1	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Rhode Island												
All races, both sexes	1	3	-	1	-	1	-	1	3	-	-	-
Male	-	1	-	1	-	1	-	1	-	-	-	-
Female	1	2	-	-	-	-	-	-	3	-	-	-
White, both sexes	1	3	-	1	-	1	-	1	3	-	-	-
Male	-	1	-	1	-	1	-	1	-	-	-	-
Female	1	2	-	-	-	-	-	-	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	1	3	4	2	4	3	5	1	-	-	-
Male	-	1	1	2	1	2	1	1	1	-	-	-
Female	2	-	2	2	1	2	2	4	-	-	-	-
White, both sexes	2	1	1	3	1	3	1	3	1	-	-	-
Male	-	1	-	1	-	2	-	1	1	-	-	-
Female	2	-	1	2	1	1	1	2	-	-	-	-
Black, both sexes	-	-	2	1	1	1	2	2	-	-	-	-
Male	-	-	1	1	1	-	1	-	-	-	-	-
Female	-	-	1	-	-	1	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3	4	2	2	6	7	6	10	2	1	-	-
Male	2	2	1	2	3	4	2	4	-	-	-	-
Female	1	2	1	-	3	3	4	6	2	1	-	-
White, both sexes	2	3	2	2	5	6	5	10	2	-	-	-
Male	1	1	1	2	2	3	2	4	-	-	-	-
Female	1	2	1	-	3	3	3	6	2	-	-	-
Black, both sexes	1	1	-	-	1	1	1	-	-	1	-	-
Male	1	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Texas												
All races, both sexes	7	7	7	7	8	11	17	11	3	1	-	-
Male	3	3	6	6	4	4	6	6	2	-	-	-
Female	4	4	1	1	4	7	11	5	1	1	-	-
White, both sexes	6	7	6	5	8	8	17	10	3	-	-	-
Male	3	3	5	5	4	4	6	6	2	-	-	-
Female	3	4	1	-	4	4	11	4	1	-	-	-
Black, both sexes	1	-	1	1	-	3	-	-	-	1	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	3	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Utah												
All races, both sexes	-	1	-	-	1	1	1	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	1	-	-	1	-	1	-	-	-	-	-
White, both sexes	-	1	-	-	1	1	1	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	1	-	-	1	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Vermont												
All races, both sexes	-	-	1	1	-	1	-	-	1	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
White, both sexes	-	-	1	1	-	1	-	-	1	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	1	3	-	2	3	13	6	1	1	-	-
Male	2	1	1	-	-	2	5	2	-	1	-	-
Female	-	-	2	-	2	1	8	4	1	-	-	-
White, both sexes	-	1	2	-	1	2	9	5	1	1	-	-
Male	-	1	1	-	-	2	3	1	-	1	-	-
Female	-	-	1	-	1	-	6	4	1	-	-	-
Black, both sexes	2	-	1	-	1	1	4	1	-	-	-	-
Male	2	-	-	-	-	-	2	1	-	-	-	-
Female	-	-	1	-	1	1	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Washington												
All races, both sexes	1	3	4	1	3	-	5	5	2	1	-	-
Male	-	-	-	1	2	-	2	-	-	-	-	-
Female	1	3	4	-	1	-	3	5	2	1	-	-
White, both sexes	1	2	4	1	2	-	5	5	2	1	-	-
Male	-	-	-	1	1	-	2	-	-	-	-	-
Female	1	2	4	-	1	-	3	5	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	1	1	-	1	2	1	1	-	1	-	-
Male	1	1	1	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	1	-	-
White, both sexes	1	1	1	-	1	2	1	1	-	1	-	-
Male	1	1	1	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Wisconsin												
All races, both sexes	1	6	1	-	1	1	5	6	1	-	-	-
Male	1	4	1	-	1	1	1	2	1	-	-	-
Female	-	2	-	-	-	-	4	4	-	-	-	-
White, both sexes	1	5	1	-	1	1	5	6	1	-	-	-
Male	1	4	1	-	1	1	1	2	1	-	-	-
Female	-	1	-	-	-	-	4	4	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	3	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	3	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
United States												
All races, both sexes	52	-	-	-	-	-	-	-	3	1	2	2
Male	26	-	-	-	-	-	-	-	2	1	1	1
Female	26	-	-	-	-	-	-	-	1	-	1	1
White, both sexes	26	-	-	-	-	-	-	-	-	1	2	2
Male	12	-	-	-	-	-	-	-	-	1	1	1
Female	14	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	23	-	-	-	-	-	-	-	3	-	-	-
Male	13	-	-	-	-	-	-	-	2	-	-	-
Female	10	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
Arkansas												
All races, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
Florida												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
Illinois												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
Massachusetts												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
Nebraska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
New Mexico												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
North Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
Ohio												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	2
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	2
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
Texas												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	-	-	-	-	-	-	-	1	-	1	-
Male	2	-	-	-	-	-	-	-	1	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections with a predominately sexual mode of transmission (A50-A64)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
United States												
All races, both sexes	3	3	3	3	4	8	4	10	5	1	-	-
Male	1	2	2	3	2	4	2	2	3	-	-	-
Female	2	1	1	-	2	4	2	8	2	1	-	-
White, both sexes	1	1	2	-	4	5	1	4	3	-	-	-
Male	-	1	2	-	2	2	-	-	2	-	-	-
Female	1	-	-	-	2	3	1	4	1	-	-	-
Black, both sexes	1	2	1	3	-	3	2	5	2	1	-	-
Male	1	1	-	3	-	2	1	2	1	-	-	-
Female	-	1	1	-	-	1	1	3	1	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	1	-	-	1	1	-	1	-	-	-	-
Male	-	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	1	-	-	1	1	-	1	-	-	-	-
Male	-	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
Florida												
All races, both sexes	-	-	1	1	1	2	-	2	-	-	-	-
Male	-	-	-	1	-	1	-	1	-	-	-	-
Female	-	-	1	-	1	1	-	1	-	-	-	-
White, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Black, both sexes	-	-	1	1	-	1	-	2	-	-	-	-
Male	-	-	-	1	-	1	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	1	-	-	-	-	1	1	-	-	-
Male	-	-	1	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
Massachusetts												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	1	-	-	-	1	-	1	1	-	-
Male	-	-	1	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
North Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
Texas												
All races, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections with a predominately sexual mode of transmission (A50-A64)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
United States												
All races, both sexes	43	-	-	-	-	-	-	-	3	1	2	-
Male	21	-	-	-	-	-	-	-	2	1	1	-
Female	22	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	17	-	-	-	-	-	-	-	-	1	2	-
Male	7	-	-	-	-	-	-	-	-	1	1	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	23	-	-	-	-	-	-	-	3	-	-	-
Male	13	-	-	-	-	-	-	-	2	-	-	-
Female	10	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
Arkansas												
All races, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
Florida												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
Illinois												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
Massachusetts												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
New Mexico												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
North Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
Ohio												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
Texas												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	-	-	-	-	-	-	-	1	-	1	-
Male	2	-	-	-	-	-	-	-	1	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Syphilis (A50-A53)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
United States												
All races, both sexes	3	2	2	3	3	8	4	8	3	1	-	-
Male	1	1	1	3	1	4	2	2	2	-	-	-
Female	2	1	1	-	2	4	2	6	1	1	-	-
White, both sexes	1	-	1	-	3	5	1	2	1	-	-	-
Male	-	-	1	-	1	2	-	-	1	-	-	-
Female	1	-	-	-	2	3	1	2	-	-	-	-
Black, both sexes	1	2	1	3	-	3	2	5	2	1	-	-
Male	1	1	-	3	-	2	1	2	1	-	-	-
Female	-	1	1	-	-	1	1	3	1	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
Florida												
All races, both sexes	-	-	1	1	1	2	-	2	-	-	-	-
Male	-	-	-	1	-	1	-	1	-	-	-	-
Female	-	-	1	-	1	1	-	1	-	-	-	-
White, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Black, both sexes	-	-	1	1	-	1	-	2	-	-	-	-
Male	-	-	-	1	-	1	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	1	-	-	-	-	1	1	-	-	-
Male	-	-	1	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
Massachusetts												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	1	-	1	1	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
North Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
Texas												
All races, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
United States												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiovascular syphilis (A52.0)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
United States												
All races, both sexes	-	-	1	-	-	2	1	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
White, both sexes	-	-	1	-	-	1	1	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiovascular syphilis (A52.0)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
United States												
All races, both sexes	30	-	-	-	-	-	-	-	3	1	1	-
Male	16	-	-	-	-	-	-	-	2	1	1	-
Female	14	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	1	1	-
Male	5	-	-	-	-	-	-	-	-	1	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	18	-	-	-	-	-	-	-	3	-	-	-
Male	11	-	-	-	-	-	-	-	2	-	-	-
Female	7	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
Arkansas												
All races, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
Florida												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
Illinois												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
Massachusetts												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
North Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
Texas												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	-	-	-	-	-	-	-	1	-	1	-
Male	2	-	-	-	-	-	-	-	1	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Neurosyphilis (A52.1-A52.3)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
United States												
All races, both sexes	3	2	1	3	3	5	1	4	2	1	-	-
Male	1	1	-	3	1	3	-	1	2	-	-	-
Female	2	1	1	-	2	2	1	3	-	1	-	-
White, both sexes	1	-	-	-	3	3	-	1	1	-	-	-
Male	-	-	-	-	1	1	-	-	1	-	-	-
Female	1	-	-	-	2	2	-	1	-	-	-	-
Black, both sexes	1	2	1	3	-	2	1	3	1	1	-	-
Male	1	1	-	3	-	2	-	1	1	-	-	-
Female	-	1	1	-	-	-	1	2	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
Florida												
All races, both sexes	-	-	1	1	1	1	-	1	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	1	1	-	1	-	1	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
Massachusetts												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
North Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
Texas												
All races, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Neurosyphilis (A52.1-A52.3)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
United States												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Florida												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
New Mexico												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Ohio												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
United States												
All races, both sexes	-	-	-	-	-	1	2	4	1	-	-	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	1	-	3	1	-	-	-
White, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	2	1	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Florida												
All races, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
United States												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
Nebraska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Gonococcal infection (A54)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
United States												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Gonococcal infection (A54)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
United States												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections with a predominately sexual mode of transmission (A55-A64)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
United States												
All races, both sexes	-	1	1	-	1	-	-	2	1	-	-	-
Male	-	1	1	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
White, both sexes	-	1	1	-	1	-	-	2	1	-	-	-
Male	-	1	1	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	1	-	-	1	-	-	1	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	1	-	-	1	-	-	1	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections with a predominately sexual mode of transmission (A55-A64)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
United States												
All races, both sexes	17	-	-	1	-	-	-	-	-	-	3	3
Male	9	-	-	-	-	-	-	-	-	-	1	3
Female	8	-	-	1	-	-	-	-	-	-	2	-
White, both sexes	15	-	-	1	-	-	-	-	-	-	3	3
Male	9	-	-	-	-	-	-	-	-	-	1	3
Female	6	-	-	1	-	-	-	-	-	-	2	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Illinois												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Kentucky												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Missouri												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
North Carolina												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Texas												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
United States												
All races, both sexes	1	3	-	-	1	2	1	2	-	-	-	-
Male	1	2	-	-	-	1	-	1	-	-	-	-
Female	-	1	-	-	1	1	1	1	-	-	-	-
White, both sexes	1	3	-	-	-	2	1	1	-	-	-	-
Male	1	2	-	-	-	1	-	1	-	-	-	-
Female	-	1	-	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Illinois												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Kentucky												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
North Carolina												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Texas												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
United States												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	2
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	2
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
Illinois												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Lyme disease (A69.2)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
United States												
All races, both sexes	1	1	-	-	-	1	1	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
White, both sexes	1	1	-	-	-	1	1	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
Illinois												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Lyme disease (A69.2)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
United States												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
North Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Typhus fever (A75)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
United States												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
North Carolina												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Typhus fever (A75)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
United States												
All races, both sexes	10	-	-	1	-	-	-	-	-	-	3	1
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	5	-	-	1	-	-	-	-	-	-	2	-
White, both sexes	9	-	-	1	-	-	-	-	-	-	3	1
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	4	-	-	1	-	-	-	-	-	-	2	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Kentucky												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Missouri												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
North Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Texas												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
United States												
All races, both sexes	-	2	-	-	-	1	-	2	-	-	-	-
Male	-	1	-	-	-	1	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	2	-	-	-	1	-	1	-	-	-	-
Male	-	1	-	-	-	1	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Illinois												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Kentucky												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Texas												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each State, 2004

[Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming reported multiple-race data in 2004. The multiple-race data for these states were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes,see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2004"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-