

Table 9. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by selected first-listed diagnostic categories: United States, 2001

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

| Category of first-listed diagnosis and ICD-9-CM code | Discharges | | Days of care | | Average length of stay in days |
|--|---|---|---------------------|---|--------------------------------|
| | Number in thousands | Rate per 10,000 population ¹ | Number in thousands | Rate per 10,000 population ¹ | |
| All conditions ² | 32,653 | 1,151.3 | 159,365 | 5,618.9 | 4.9 |
| Heart disease | 391–392,0,393–398,402,404,410–416,420–429 | 4,323 | 152.4 | 20,015 | 705.7 |
| Acute myocardial infarction | 410 | 795 | 28.0 | 4,595 | 162.0 |
| Coronary atherosclerosis | 414.0 | 1,061 | 37.4 | 3,803 | 134.1 |
| Other ischemic heart disease | 411–413,414.1–414.9 | 234 | 8.2 | 583 | 20.6 |
| Cardiac dysrhythmias | 427 | 723 | 25.5 | 2,623 | 92.5 |
| Congestive heart failure | 428.0 | 995 | 35.1 | 5,416 | 191.0 |
| Females with deliveries | V27 | 3,839 | 135.4 | 9,702 | 342.1 |
| Psychoses | 290–299 | 1,633 | 57.6 | 13,223 | 466.2 |
| Pneumonia | 480–486 | 1,300 | 45.8 | 7,411 | 261.3 |
| Malignant neoplasms | 140–208,230–234 | 1,212 | 42.7 | 8,514 | 300.2 |
| Malignant neoplasm of large intestine and rectum | 153–154,197.5 | 157 | 5.6 | 1,356 | 47.8 |
| Malignant neoplasm of trachea, bronchus, and lung | 162,176.4,197.0,197.3 | 153 | 5.4 | 1,160 | 40.9 |
| Malignant neoplasm of breast | 174–175,198.81 | 107 | 3.8 | 309 | 10.9 |
| Fractures, all sites ³ | 800–829 | 999 | 35.2 | 5,369 | 189.3 |
| Fracture of neck of femur ³ | 820 | 324 | 11.4 | 2,118 | 74.7 |
| Cerebrovascular disease | 430–438 | 931 | 32.8 | 4,821 | 170.0 |
| Diabetes mellitus | 250 | 562 | 19.8 | 2,761 | 97.4 |
| Volume depletion | 276.5 | 534 | 18.8 | 2,025 | 71.4 |
| Chronic bronchitis | 491 | 505 | 17.8 | 2,641 | 93.1 |
| Osteoarthritis and allied disorders | 715 | 496 | 17.5 | 2,197 | 77.5 |
| Asthma | 493 | 454 | 16.0 | 1,469 | 51.8 |
| Cellulitis and abscess | 681–682 | 404 | 14.2 | 1,933 | 68.1 |
| Benign neoplasms | 210–229 | 387 | 13.6 | 1,240 | 43.7 |
| Cholelithiasis | 574 | 367 | 12.9 | 1,429 | 50.4 |
| Intervertebral disc disorder | 722 | 325 | 11.5 | 955 | 33.7 |
| Septicemia | 038 | 315 | 11.1 | 2,798 | 98.7 |
| Noninfectious enteritis and colitis | 555–558 | 301 | 10.6 | 1,305 | 46.0 |

¹Rates were calculated using U.S. Census Bureau estimates of the civilian population based on the 2000 census.

²Includes data for diagnostic conditions not shown in table.

³Excludes fractures coded as 733.1, pathologic fracture.