Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
All causes														
All origins[3]														
Both sexes														
2001	848.5	683.4	33.3	15.3	19.2	66.9	95.0	96.2	113.5	165.9	240.5	354.8	512.4	771.8
2000[4]	854.0	736.7	32.4	15.8	20.3	67.1	93.6	91.2	111.0	159.2	239.1	352.5	509.0	792.6
1999[4]	857.0	736.0	34.2	16.9	20.4	68.6	90.8	90.4	113.4	160.8	236.3	346.6	501.4	801.6
Male														
2001	846.4	749.8	37.0	16.7	22.8	93.7	141.1	135.6	151.2	213.7	304.8	455.4	646.8	958.4
2000[4]	853.0	806.5	35.9	17.6	24.2	93.3	138.0	128.8	147.9	205.4	305.9	456.3	642.2	989.8
1999[4]	859.2	805.0	37.9	18.6	24.4	94.7	133.1	127.0	152.0	207.6	304.0	448.0	632.4	1,003.8
Female														
2001	850.4	613.9	29.5	13.9	15.3	38.5	46.8	55.8	75.2	118.2	177.2	257.2	383.7	596.9
2000[4]	855.0	663.4	28.7	14.0	16.1	39.3	47.1	52.8	73.6	113.2	173.4	252.0	381.3	608.1
1999[4]	854.9	663.6	30.3	15.0	16.2	41.0	46.6	53.2	74.4	114.4	169.8	248.5	375.9	612.6
Hispanic														
Both sexes														
2001	306.8	572.7	30.6	13.0	16.6	63.1	86.9	80.6	88.7	129.5	185.3	280.9	403.9	580.6
2000[4]	303.8	596.3	29.6	14.1	17.5	61.6	83.3	76.7	93.2	129.4	192.2	281.1	394.3	612.1
1999[4]	305.7	583.8	30.9	13.3	18.3	61.0	81.1	77.5	94.1	132.9	189.4	283.6	398.1	602.7
Male														
2001	332.9	624.4	33.8	14.9	18.6	92.0	128.2	115.2	121.1	175.5	249.0	379.7	529.6	751.2
2000[4]	331.3	637.1	31.5	15.0	21.3	90.5	123.1	110.3	131.1	172.5	259.1	378.6	519.5	801.5
1999[4]	332.6	623.4	32.9	14.9	21.2	87.8	119.1	111.1	130.8	183.0	255.7	383.9	525.5	788.0
Female														
2001	279.0	518.9	27.2	11.0	14.5	31.1	36.1	39.4	51.3	79.0	118.3	181.7	283.8	426.0
2000[4]	274.6	553.6	27.5	13.2	13.6	28.7	34.6	37.5	49.8	82.4	122.2	183.8	274.9	440.6
1999[4]	277.2	542.3	28.7	11.6	15.2	30.5	35.3	38.0	52.1	78.3	120.4	184.2	277.2	435.3
Non-Hispanic[5]														
Both sexes														
2001	926.2	703.5	33.8	15.8	19.6	67.3	96.2	99.2	117.5	170.6	245.8	360.4	519.3	783.7
2000[4]	929.6	760.4	32.9	16.1	20.7	67.7	95.3	93.8	113.5	162.4	242.8	357.3	515.8	802.9
1999[4]	929.9	760.7	34.8	17.5	20.7	69.7	92.4	92.5	116.1	163.7	239.9	350.4	507.2	812.7
Male														
2001	923.6	773.2	37.6	17.0	23.6	93.6	143.3	139.8	156.2	218.5	309.6	460.5	653.3	969.6
2000[4]	928.1	836.8	36.8	18.1	24.7	93.4	140.9	132.2	150.3	208.8	309.0	460.9	648.3	998.9
1999[4]	932.2	834.9	38.8	19.3	24.9	95.8	135.7	129.6	154.9	209.7	307.1	450.9	637.0	1,014.4
Female														
2001	928.6	630.4	29.9	14.4	15.4	39.7	48.7	58.8	79.1	123.4	183.5	263.6	391.0	608.9
2000[4]	931.0	680.1	28.8	14.0	16.4	41.0	49.3	55.6	77.1	116.8	178.2	257.3	388.8	619.1
1999[4]	927.8	682.9	30.6	15.7	16.3	42.5	48.6	55.7	77.7	118.5	174.2	253.4	382.7	623.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All causes						
All origins[3]						
Both sexes						
2001	1,210.7	1,869.7	2,878.3	4,494.0	7,151.9	15,112.8
2000[4]	1,241.0	1,906.3	2,929.4	4,553.8	7,335.0	15,524.4
1999[4]	1,254.0	1,956.7	2,995.3	4,608.0	7,378.5	15,554.6
Male						
2001	1,496.8	2,315.6	3,587.4	5,603.2	8,781.9	16,744.8
2000[4]	1,535.9	2,362.0	3,676.1	5,693.2	9,095.4	17,501.4
1999[4]	1,560.8	2,444.1	3,785.1	5,806.4	9,216.8	17,597.2
Female						
2001	951.3	1,485.9	2,314.8	3,717.6	6,175.1	14,429.9
2000[4]	973.7	1,515.7	2,341.2	3,760.2	6,296.6	14,719.2
1999[4]	976.4	1,539.6	2,378.2	3,774.8	6,302.8	14,731.3
Hispanic						
Both sexes						
2001	910.1	1,412.8	2,229.5	3,556.3	5,501.5	11,821.8
2000[4]	913.8	1,425.9	2,278.4	3,543.6	5,777.6	11,822.9
1999[4]	940.8	1,527.8	2,264.3	3,577.0	5,774.4	12,177.9
Male						
2001	1,158.6	1,817.5	2,791.5	4,456.8	6,783.5	12,888.3
2000[4]	1,181.3	1,804.0	2,918.9	4,480.2	7,220.4	13,086.2
1999[4]	1,215.5	1,942.0	2,931.5	4,571.8	7,005.8	13,485.9
Female						
2001	695.1	1,085.0	1,800.8	2,917.0	4,707.4	11,300.5
2000[4]	683.1	1,119.7	1,793.2	2,880.3	4,899.7	11,202.8
1999[4]	704.0	1,192.9	1,762.5	2,874.8	5,039.8	11,547.3
Non-Hispanic[5]						
Both sexes						
2001	1,228.7	1,896.2	2,908.7	4,528.0	7,201.4	15,199.0
2000[4]	1,259.9	1,931.9	2,956.6	4,586.6	7,375.0	15,619.8
1999[4]	1,271.3	1,977.5	3,025.1	4,639.4	7,416.1	15,631.0
Male						
2001	1,514.1	2,341.5	3,621.9	5,644.0	8,842.8	16,866.3
2000[4]	1,554.1	2,388.4	3,704.3	5,730.8	9,144.1	17,639.8
1999[4]	1,577.2	2,466.2	3,815.8	5,842.6	9,272.9	17,710.4
Female						
2001	968.9	1,511.2	2,340.5	3,747.4	6,219.1	14,505.6
2000[4]	992.4	1,539.1	2,366.3	3,790.4	6,332.6	14,802.9
1999[4]	993.6	1,557.9	2,406.0	3,803.4	6,330.3	14,798.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
All causes														
Non-Hispanic White														
Both sexes														
2001	991.1	555.4	30.1	14.5	18.0	63.4	85.4	87.2	104.4	153.1	220.9	324.7	474.0	730.8
2000[4]	993.2	596.5	28.5	14.7	19.4	64.0	84.1	80.6	98.7	144.5	216.4	318.7	470.1	749.9
1999[4]	990.7	593.1	30.1	15.6	19.1	65.3	80.2	80.3	101.3	144.5	211.0	310.1	461.9	754.0
Male														
2001	975.6	611.6	33.7	15.2	21.7	86.1	125.6	122.4	139.3	197.9	282.4	417.3	596.5	901.6
2000[4]	978.5	658.7	32.4	16.6	23.3	86.1	122.9	112.7	132.2	188.0	278.5	415.7	588.9	928.4
1999[4]	979.6	658.1	33.4	17.0	22.7	87.1	116.1	112.1	135.7	187.1	272.8	399.6	578.8	939.4
Female														
2001	1,006.1	496.4	26.3	13.7	14.1	39.6	44.3	51.4	69.0	108.2	159.7	233.1	353.9	567.1
2000[4]	1,007.3	530.9	24.4	12.6	15.2	41.0	44.5	48.0	64.7	100.8	154.5	222.8	353.6	579.3
1999[4]	1,001.3	524.6	26.7	14.1	15.2	42.4	43.3	48.2	66.5	101.7	149.2	221.7	347.5	576.9
Non-Hispanic Black														
Both sexes														
2001	798.1	1,354.4	49.5	21.4	26.6	88.7	156.7	174.9	207.1	294.6	438.1	656.8	946.1	1,351.9
2000[4]	805.5	1,477.8	51.7	22.6	27.5	89.9	160.5	175.2	213.6	290.5	443.1	679.7	956.6	1,368.9
1999[4]	812.1	1,510.2	54.6	26.2	29.0	94.8	160.6	169.8	212.0	301.4	457.4	681.6	945.4	1,422.8
Male														
2001	849.7	1,484.2	55.1	24.6	32.5	133.9	244.7	255.7	279.6	375.1	538.9	846.0	1,222.7	1,746.2
2000[4]	859.5	1,623.2	56.1	25.4	33.0	134.0	249.2	256.9	283.2	371.1	558.6	869.4	1,248.2	1,802.9
1999[4]	872.8	1,650.8	61.5	30.1	36.3	142.3	249.7	247.5	288.0	382.5	583.2	902.1	1,238.5	1,851.6
Female														
2001	751.2	1,217.8	43.6	18.1	20.5	42.0	72.2	101.4	142.2	223.2	349.0	493.1	710.9	1,028.9
2000[4]	756.7	1,327.7	47.1	19.7	21.7	44.9	76.5	101.3	151.1	219.2	341.0	515.5	708.3	1,013.7
1999[4]	757.3	1,365.4	47.4	22.1	21.5	46.3	76.1	99.5	144.0	229.5	346.8	491.7	697.1	1,073.0
Salmonella infections (A01-A02)														
All origins[3]														
Both sexes														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All causes						
Non-Hispanic White						
Both sexes						
2001	1,169.7	1,827.4	2,849.2	4,476.8	7,196.1	15,371.2
2000[4]	1,199.4	1,865.4	2,889.5	4,525.0	7,369.2	15,799.7
1999[4]	1,206.2	1,911.7	2,949.5	4,570.3	7,406.6	15,795.3
Male						
2001	1,439.8	2,256.8	3,543.1	5,580.2	8,828.0	17,055.3
2000[4]	1,477.3	2,305.7	3,615.5	5,653.4	9,137.4	17,853.2
1999[4]	1,494.4	2,384.2	3,724.5	5,759.4	9,266.7	17,871.2
Female						
2001	919.2	1,448.1	2,285.2	3,695.5	6,211.0	14,670.6
2000[4]	941.7	1,478.3	2,305.2	3,731.9	6,318.4	14,971.7
1999[4]	939.4	1,497.5	2,331.8	3,735.9	6,309.1	14,967.5
Non-Hispanic Black						
Both sexes						
2001	1,920.5	2,777.9	3,951.1	5,712.5	8,098.0	14,660.8
2000[4]	1,959.9	2,809.4	4,065.0	5,837.2	8,234.9	14,837.5
1999[4]	1,998.3	2,851.2	4,191.4	5,927.6	8,223.1	14,839.2
Male						
2001	2,456.8	3,547.9	5,105.6	7,313.8	10,346.7	16,576.2
2000[4]	2,498.8	3,559.1	5,258.8	7,462.1	10,491.3	16,944.8
1999[4]	2,583.0	3,644.4	5,379.6	7,591.6	10,490.8	17,385.3
Female						
2001	1,498.8	2,209.4	3,182.0	4,756.8	6,952.7	13,951.4
2000[4]	1,534.7	2,256.3	3,272.3	4,857.7	7,099.0	14,054.9
1999[4]	1,539.8	2,263.4	3,402.8	4,925.0	7,099.0	13,887.7
Salmonella infections (A01-A02)						
All origins[3]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Salmonella infections (A01-A02)														
All origins[3]														
Female														
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Salmonella infections (A01-A02)						
All origins[3]						
Female						
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic White						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Salmonella infections (A01-A02)														
Non-Hispanic White														
Male														
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Shigellosis and amebiasis (A03,A06)														
All origins[3]														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Salmonella infections (A01-A02)						
Non-Hispanic White						
Male						
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Shigellosis and amebiasis (A03,A06)						
All origins[3]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Shigellosis and amebiasis (A03,A06)														
Hispanic														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Shigellosis and amebiasis (A03,A06)						
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic White						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Shigellosis and amebiasis (A03,A06)														
Non-Hispanic White														
Female														
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Certain other intestinal infections (A04,A07-A09)														
All origins[3]														
Both sexes														
2001	0.5	*	*	*	*	*	*	*	*	*	*	*	0.2	0.2
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	0.2
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	0.3
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.7	*	*	*	*	*	*	*	*	*	*	*	0.2	*
2000[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Shigellosis and amebiasis (A03,A06)						
Non-Hispanic White						
Female						
1999[4]	*	*	*	*	*	*
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Certain other intestinal infections (A04,A07-A09)						
All origins[3]						
Both sexes						
2001	0.5	0.8	1.8	3.2	6.4	12.8
2000[4]	0.3	0.9	1.5	2.7	5.3	11.8
1999[4]	0.3	0.7	1.2	2.3	4.5	10.3
Male						
2001	0.4	0.7	2.0	3.5	6.2	12.1
2000[4]	*	1.0	1.6	2.9	5.8	11.4
1999[4]	*	0.6	1.2	2.7	4.9	9.5
Female						
2001	0.5	0.8	1.8	3.0	6.5	13.1
2000[4]	*	0.8	1.5	2.6	4.9	11.9
1999[4]	*	0.8	1.3	2.0	4.3	10.7
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Certain other intestinal infections (A04,A07-A09)														
Hispanic														
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.6	*	*	*	*	*	*	*	*	*	*	*	0.2	0.2
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	0.2
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.7	*	*	*	*	*	*	*	*	*	*	*	0.2	*
2000[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	0.7	*	*	*	*	*	*	*	*	*	*	*	0.2	0.2
2000[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	*	0.2
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.9	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.7	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Certain other intestinal infections (A04,A07-A09)						
Hispanic						
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	0.5	0.8	1.9	3.2	6.4	12.8
2000[4]	0.3	0.9	1.6	2.8	5.3	11.9
1999[4]	0.3	0.7	1.3	2.3	4.7	10.4
Male						
2001	0.4	0.8	2.0	3.5	6.4	12.0
2000[4]	*	1.0	1.5	2.9	5.9	11.5
1999[4]	*	0.6	1.2	2.7	5.1	9.6
Female						
2001	0.6	0.8	1.8	3.0	6.4	13.2
2000[4]	*	0.8	1.6	2.7	5.0	12.1
1999[4]	*	0.8	1.3	2.1	4.4	10.8
Non-Hispanic White						
Both sexes						
2001	0.5	0.7	2.0	3.4	6.6	13.6
2000[4]	0.3	0.9	1.6	2.9	5.7	12.3
1999[4]	0.3	0.7	1.3	2.4	4.9	11.0
Male						
2001	*	0.7	2.1	3.6	6.4	12.6
2000[4]	*	1.0	1.6	3.1	6.2	12.0
1999[4]	*	0.7	1.3	2.8	5.4	10.0
Female						
2001	0.5	0.7	1.9	3.2	6.7	14.0
2000[4]	*	0.8	1.7	2.8	5.4	12.5
1999[4]	*	0.8	1.3	2.1	4.6	11.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Certain other intestinal infections (A04,A07-A09)														
Non-Hispanic Black														
Both sexes														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Tuberculosis (A16-A19)														
All origins[3]														
Both sexes														
2001	0.3	*	*	*	*	*	*	*	*	*	0.1	0.2	0.3	0.4
2000[4]	0.3	*	*	*	*	*	*	*	*	*	0.1	0.2	0.3	0.2
1999[4]	0.3	*	*	*	*	*	*	*	*	0.1	0.1	0.3	0.3	0.5
Male														
2001	0.3	*	*	*	*	*	*	*	*	*	*	0.3	0.4	0.7
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	0.4	0.5	0.4
1999[4]	0.4	*	*	*	*	*	*	*	*	0.2	0.2	0.4	0.5	0.8
Female														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Certain other intestinal infections (A04,A07-A09)						
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	8.3
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Tuberculosis (A16-A19)						
All origins[3]						
Both sexes						
2001	0.5	0.6	1.0	1.4	2.0	2.8
2000[4]	0.5	0.7	1.0	1.8	2.4	2.7
1999[4]	0.6	0.7	1.2	1.8	2.4	4.2
Male						
2001	0.7	0.9	1.5	1.7	2.4	4.4
2000[4]	0.7	1.0	1.1	2.3	3.8	4.6
1999[4]	0.9	1.1	1.7	2.4	3.7	7.1
Female						
2001	*	0.4	0.7	1.1	1.8	2.2
2000[4]	*	0.5	0.9	1.4	1.6	2.0
1999[4]	*	0.4	0.7	1.5	1.7	3.0
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Tuberculosis (A16-A19)														
Hispanic														
Female														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.3	*	*	*	*	*	*	*	*	*	*	0.2	0.2	0.4
2000[4]	0.3	*	*	*	*	*	*	*	*	*	0.1	0.2	0.3	0.2
1999[4]	0.3	*	*	*	*	*	*	*	*	*	0.1	0.2	0.3	0.5
Male														
2001	0.3	*	*	*	*	*	*	*	*	*	*	0.3	0.3	0.6
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	0.3	0.5	0.4
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	0.4	0.4	0.7
Female														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	0.2
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	0.2	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	0.3
Male														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	0.5
Female														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	0.6	*	*	*	*	*	*	*	*	*	*	*	1.1	1.7
2000[4]	0.7	*	*	*	*	*	*	*	*	*	*	0.9	1.2	*
1999[4]	0.7	*	*	*	*	*	*	*	*	*	*	0.9	1.2	1.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Tuberculosis (A16-A19)						
Hispanic						
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	0.4	0.6	1.0	1.3	1.9	2.7
2000[4]	0.4	0.7	0.9	1.6	2.3	2.5
1999[4]	0.5	0.7	1.1	1.6	2.2	3.9
Male						
2001	0.7	0.8	1.4	1.7	2.2	4.3
2000[4]	0.6	1.0	1.0	2.0	3.6	4.3
1999[4]	0.8	1.0	1.6	2.1	3.5	6.8
Female						
2001	*	*	0.7	1.0	1.7	2.1
2000[4]	*	0.5	0.9	1.4	1.6	1.8
1999[4]	*	*	0.7	1.3	1.5	2.8
Non-Hispanic White						
Both sexes						
2001	0.3	0.4	0.6	1.0	1.3	2.0
2000[4]	*	0.4	0.5	1.0	1.6	2.0
1999[4]	0.3	0.5	0.6	1.1	1.7	3.0
Male						
2001	*	*	0.6	1.3	1.5	2.9
2000[4]	*	0.6	0.7	1.2	2.0	3.6
1999[4]	*	0.7	1.0	1.3	2.8	4.9
Female						
2001	*	*	0.5	0.8	1.1	1.7
2000[4]	*	*	*	0.9	1.3	1.4
1999[4]	*	*	*	0.9	1.1	2.2
Non-Hispanic Black						
Both sexes						
2001	*	*	3.4	*	*	6.9
2000[4]	*	2.8	3.7	4.9	8.9	*
1999[4]	*	*	3.7	5.6	*	8.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Tuberculosis (A16-A19)														
Non-Hispanic Black														
Male														
2001	0.7	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.9	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.9	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Respiratory tuberculosis (A16)														
All origins[3]														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	0.1	0.2	0.4
2000[4]	0.2	*	*	*	*	*	*	*	*	*	0.1	0.1	0.2	0.2
1999[4]	0.3	*	*	*	*	*	*	*	*	*	0.1	0.2	0.2	0.4
Male														
2001	0.3	*	*	*	*	*	*	*	*	*	*	0.2	0.3	0.6
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	0.3	0.4	0.3
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	0.3	0.4	0.6
Female														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Tuberculosis (A16-A19)						
Non-Hispanic Black						
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	19.7	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Respiratory tuberculosis (A16)						
All origins[3]						
Both sexes						
2001	0.4	0.5	0.8	1.1	1.6	2.4
2000[4]	0.3	0.5	0.8	1.4	2.1	2.3
1999[4]	0.5	0.6	0.9	1.5	2.1	3.7
Male						
2001	0.7	0.7	1.2	1.5	2.0	3.8
2000[4]	0.6	0.9	0.9	1.9	3.2	4.1
1999[4]	0.9	0.9	1.4	2.0	3.3	6.4
Female						
2001	*	*	0.5	0.8	1.4	1.8
2000[4]	*	*	0.6	1.1	1.4	1.6
1999[4]	*	*	0.5	1.2	1.4	2.6
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Respiratory tuberculosis (A16)														
Non-Hispanic[5]														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	0.1	0.2	0.3
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	0.2	0.2	0.2
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	0.2	0.2	0.4
Male														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	0.3	0.5
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	0.3	0.4	0.3
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	0.3	0.3	0.6
Female														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	0.2
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	0.2	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	0.2
Male														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	*	1.6
Male														
2001	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.8	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.7	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Respiratory tuberculosis (A16)						
Non-Hispanic[5]						
Both sexes						
2001	0.4	0.4	0.7	1.1	1.5	2.4
2000[4]	0.3	0.5	0.7	1.3	2.0	2.2
1999[4]	0.5	0.6	0.8	1.4	2.0	3.5
Male						
2001	0.7	0.6	1.0	1.4	1.7	3.8
2000[4]	0.5	0.8	0.8	1.7	3.0	3.8
1999[4]	0.8	0.9	1.3	1.8	3.2	6.1
Female						
2001	*	*	0.5	0.8	1.4	1.8
2000[4]	*	*	0.6	1.1	1.4	1.5
1999[4]	*	*	0.5	1.1	1.3	2.4
Non-Hispanic White						
Both sexes						
2001	0.2	0.3	0.4	0.8	1.0	1.7
2000[4]	*	0.3	0.4	0.9	1.3	1.8
1999[4]	0.3	0.4	0.5	0.9	1.5	2.8
Male						
2001	*	*	*	1.1	*	2.4
2000[4]	*	*	*	1.1	1.6	3.2
1999[4]	*	0.6	0.8	1.1	2.5	4.8
Female						
2001	*	*	*	0.6	0.9	1.4
2000[4]	*	*	*	0.7	1.2	1.2
1999[4]	*	*	*	0.7	1.0	1.9
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	6.2
2000[4]	*	2.4	3.0	4.0	6.9	*
1999[4]	*	*	*	4.7	*	7.1
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Respiratory tuberculosis (A16)														
Non-Hispanic Black														
Female														
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Other tuberculosis (A17-A19)														
All origins[3]														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Respiratory tuberculosis (A16)						
Non-Hispanic Black						
Female						
1999[4]	*	*	*	*	*	*
Other tuberculosis (A17-A19)						
All origins[3]						
Both sexes						
2001	*	*	0.3	0.3	0.4	*
2000[4]	*	*	*	0.3	*	*
1999[4]	*	*	0.2	0.3	*	0.5
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	0.3	*	0.4	*
2000[4]	*	*	*	0.3	*	*
1999[4]	*	*	0.2	0.3	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other tuberculosis (A17-A19)														
Non-Hispanic[5]														
Male														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other tuberculosis (A17-A19)						
Non-Hispanic[5]						
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic White						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Whooping cough (A37)														
All origins[3]														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Whooping cough (A37)						
All origins[3]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Whooping cough (A37)														
Non-Hispanic[5]														
Female														
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Scarlet fever and erysipelas (A38,A46)														
All origins[3]														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Whooping cough (A37)						
Non-Hispanic[5]						
Female						
1999[4]	*	*	*	*	*	*
Non-Hispanic White						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Scarlet fever and erysipelas (A38,A46)						
All origins[3]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Scarlet fever and erysipelas (A38,A46)														
All origins[3]														
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Scarlet fever and erysipelas (A38,A46)						
All origins[3]						
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Scarlet fever and erysipelas (A38,A46)														
Non-Hispanic White														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Meningococcal infection (A39)														
All origins[3]														
Both sexes														
2001	0.1	*	*	*	*	0.2	*	*	*	*	*	*	*	*
2000[4]	0.1	0.6	*	*	*	0.2	0.1	*	*	*	*	*	*	*
1999[4]	0.1	0.8	0.2	*	*	0.1	*	*	*	*	*	*	*	*
Male														
2001	0.1	*	*	*	*	0.2	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	0.2	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Scarlet fever and erysipelas (A38,A46)						
Non-Hispanic White						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Meningococcal infection (A39)						
All origins[3]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Meningococcal infection (A39)														
All origins[3]														
Female														
2001	0.1	*	*	*	*	0.2	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.1	*	*	*	*	0.2	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	0.2	*	*	*	*	*	*	*	*
1999[4]	0.1	0.8	0.2	*	*	0.2	*	*	*	*	*	*	*	*
Male														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	0.3	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	0.3	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	0.1	*	*	*	*	0.2	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	0.2	*	*	*	*	*	*	*	*
1999[4]	0.1	1.0	0.3	*	*	0.2	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Meningococcal infection (A39)						
All origins[3]						
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic White						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Meningococcal infection (A39)														
Non-Hispanic White														
Male														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Septicemia (A40-A41)														
All origins[3]														
Both sexes														
2001	11.3	7.7	0.7	0.1	0.2	0.3	0.4	0.5	0.9	1.4	2.3	4.0	6.2	9.8
2000[4]	11.1	7.2	0.6	0.2	0.1	0.2	0.3	0.5	0.9	1.4	2.5	3.9	6.1	9.4
1999[4]	11.0	7.5	0.6	0.2	0.1	0.2	0.5	0.5	0.8	1.4	2.2	3.7	5.6	9.0
Male														
2001	10.2	8.4	0.8	*	0.2	0.3	0.4	0.5	0.9	1.5	2.5	4.3	6.7	10.6
2000[4]	9.8	7.1	0.7	0.2	*	0.2	0.4	0.6	0.9	1.4	2.6	4.4	7.0	10.1
1999[4]	9.8	9.0	0.6	0.2	*	*	0.5	0.5	0.9	1.5	2.6	4.2	6.2	9.7
Female														
2001	12.4	7.0	0.6	*	*	0.3	0.3	0.5	0.9	1.2	2.1	3.8	5.7	9.1
2000[4]	12.3	7.3	0.6	*	*	*	0.2	0.4	0.8	1.4	2.3	3.4	5.2	8.7
1999[4]	12.2	5.8	0.5	0.2	*	*	0.4	0.5	0.7	1.3	1.9	3.3	5.1	8.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Meningococcal infection (A39)						
Non-Hispanic White						
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Septicemia (A40-A41)						
All origins[3]						
Both sexes						
2001	15.5	25.3	40.9	67.3	103.9	205.9
2000[4]	15.2	24.2	38.4	64.0	105.0	215.7
1999[4]	14.4	24.7	38.3	63.8	102.9	220.7
Male						
2001	16.6	27.8	46.7	78.6	122.3	224.3
2000[4]	16.7	26.9	43.2	73.9	121.7	223.0
1999[4]	15.7	27.7	43.7	74.8	120.6	233.5
Female						
2001	14.4	23.1	36.2	59.5	92.9	198.3
2000[4]	13.8	21.8	34.6	57.1	95.2	212.7
1999[4]	13.1	22.1	34.1	56.2	92.5	215.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Septicemia (A40-A41)														
Hispanic														
Both sexes														
2001	3.7	6.1	*	*	*	*	*	*	*	0.9	1.8	3.6	4.3	5.4
2000[4]	3.5	8.0	*	*	*	*	*	*	*	1.0	2.2	2.9	5.2	6.4
1999[4]	3.6	5.7	*	*	*	*	*	*	*	1.1	2.0	2.1	4.4	6.9
Male														
2001	3.6	7.0	*	*	*	*	*	*	*	1.3	1.8	3.8	6.0	6.4
2000[4]	3.4	8.4	*	*	*	*	*	*	*	*	2.5	4.1	6.3	8.1
1999[4]	3.5	7.0	*	*	*	*	*	*	*	1.4	2.8	2.6	5.4	8.6
Female														
2001	3.9	5.2	*	*	*	*	*	*	*	*	1.7	3.4	2.6	4.5
2000[4]	3.5	7.7	*	*	*	*	*	*	*	*	1.9	*	4.2	4.8
1999[4]	3.7	*	*	*	*	*	*	*	*	*	*	*	3.5	5.4
Non-Hispanic[5]														
Both sexes														
2001	12.4	8.1	0.7	0.1	0.2	0.3	0.4	0.6	1.0	1.5	2.3	4.1	6.3	10.1
2000[4]	12.1	6.9	0.7	0.2	0.1	0.2	0.3	0.5	0.9	1.5	2.5	4.0	6.1	9.5
1999[4]	12.0	7.9	0.6	0.2	0.2	0.2	0.6	0.6	0.9	1.4	2.3	3.9	5.7	9.1
Male														
2001	11.2	8.8	0.9	*	*	0.3	0.5	0.6	0.9	1.6	2.6	4.3	6.7	10.9
2000[4]	10.7	6.7	0.7	0.2	*	0.2	0.4	0.7	1.0	1.5	2.6	4.5	7.0	10.2
1999[4]	10.7	9.4	0.6	*	*	*	0.6	0.6	1.0	1.5	2.6	4.3	6.2	9.7
Female														
2001	13.5	7.4	0.6	*	*	0.3	0.4	0.5	1.1	1.3	2.1	3.8	6.0	9.3
2000[4]	13.5	7.0	0.6	*	*	*	*	0.4	0.9	1.5	2.4	3.5	5.2	9.0
1999[4]	13.2	6.2	0.5	0.3	*	*	0.5	0.6	0.8	1.3	1.9	3.4	5.2	8.5
Non-Hispanic White														
Both sexes														
2001	12.3	5.1	0.7	*	*	0.2	0.3	0.4	0.7	1.1	1.8	3.1	5.0	8.2
2000[4]	11.9	4.8	0.5	0.2	*	0.2	0.2	0.4	0.7	1.1	2.0	3.0	4.8	7.9
1999[4]	11.7	5.3	0.5	0.2	*	0.2	0.4	0.5	0.7	1.0	1.7	2.8	4.5	7.2
Male														
2001	11.0	5.8	0.9	*	*	*	*	0.5	0.6	1.2	2.1	3.3	5.2	8.6
2000[4]	10.4	4.7	0.5	*	*	*	0.3	0.4	0.7	1.2	2.1	3.4	5.3	8.4
1999[4]	10.2	6.7	0.5	*	*	*	0.6	0.5	0.8	1.1	1.9	3.2	4.9	7.5
Female														
2001	13.6	4.4	0.5	*	*	*	*	0.4	0.8	0.9	1.5	2.9	4.7	7.9
2000[4]	13.4	4.8	0.6	*	*	*	*	*	0.6	1.0	1.9	2.5	4.2	7.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Septicemia (A40-A41)						
Hispanic						
Both sexes						
2001	11.8	20.5	31.5	54.1	77.0	171.3
2000[4]	11.2	16.2	28.9	48.1	79.6	144.7
1999[4]	9.4	20.6	31.9	52.3	78.7	180.3
Male						
2001	14.7	24.5	36.6	68.9	89.6	160.0
2000[4]	12.4	22.4	32.1	58.3	94.2	157.2
1999[4]	12.2	25.9	34.1	70.2	84.9	181.2
Female						
2001	9.4	17.2	27.6	43.6	69.2	176.9
2000[4]	10.2	11.2	26.5	40.8	70.8	138.5
1999[4]	6.9	16.3	30.3	39.6	75.0	179.9
Non-Hispanic[5]						
Both sexes						
2001	15.6	25.6	41.3	67.8	104.7	206.8
2000[4]	15.4	24.6	38.8	64.4	105.6	217.5
1999[4]	14.7	24.8	38.5	64.1	103.4	221.2
Male						
2001	16.6	28.0	47.1	78.8	123.1	226.8
2000[4]	16.9	26.9	43.6	74.3	122.0	224.8
1999[4]	16.0	27.7	44.1	74.5	121.5	234.5
Female						
2001	14.8	23.5	36.6	60.2	93.6	198.5
2000[4]	14.0	22.5	35.0	57.6	95.9	214.5
1999[4]	13.5	22.3	34.1	56.8	92.9	215.9
Non-Hispanic White						
Both sexes						
2001	13.2	22.3	36.9	62.4	99.1	195.6
2000[4]	12.7	20.9	34.8	58.2	97.7	204.3
1999[4]	12.3	21.2	33.5	57.6	95.7	206.0
Male						
2001	14.2	24.2	41.6	72.4	116.6	213.6
2000[4]	14.1	22.9	39.7	66.7	114.0	211.2
1999[4]	12.8	23.4	38.3	67.8	112.1	218.1
Female						
2001	12.3	20.6	33.1	55.2	88.6	188.1
2000[4]	11.4	19.1	30.8	52.1	88.1	201.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Septicemia (A40-A41)														
Non-Hispanic White														
Female														
1999[4]	13.1	3.8	0.5	*	*	*	*	0.4	0.6	0.9	1.5	2.5	4.2	6.9
Non-Hispanic Black														
Both sexes														
2001	16.4	20.3	0.9	*	*	*	0.8	1.1	2.7	3.9	5.7	10.9	17.1	27.6
2000[4]	16.7	16.6	1.1	*	*	*	0.8	1.5	2.6	3.8	5.9	11.3	17.3	24.7
1999[4]	16.6	18.6	*	*	*	*	1.0	1.4	1.8	3.8	6.4	11.3	15.5	25.8
Male														
2001	15.4	21.4	*	*	*	*	*	*	2.6	3.9	5.9	12.0	19.1	34.0
2000[4]	15.3	16.4	*	*	*	*	*	2.2	2.6	3.6	6.0	12.8	21.9	27.9
1999[4]	15.7	20.8	*	*	*	*	*	*	2.2	4.1	7.9	13.2	18.1	30.1
Female														
2001	17.2	19.1	*	*	*	*	*	*	2.7	3.9	5.6	10.0	15.5	22.4
2000[4]	17.9	16.9	*	*	*	*	*	*	2.6	4.1	5.8	10.0	13.4	22.0
1999[4]	17.4	16.4	*	*	*	*	*	1.7	1.5	3.6	5.0	9.6	13.3	22.2
Syphilis (A50-A53)														
All origins[3]														
Both sexes														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Septicemia (A40-A41)						
Non-Hispanic White						
Female						
1999[4]	11.8	19.4	29.7	50.4	86.1	201.2
Non-Hispanic Black						
Both sexes						
2001	38.9	58.2	90.9	141.2	194.2	368.6
2000[4]	40.5	60.8	86.7	148.0	218.1	395.9
1999[4]	37.0	58.8	95.6	143.9	211.6	414.2
Male						
2001	41.8	67.5	112.6	173.0	237.9	430.8
2000[4]	45.9	69.8	96.7	185.0	251.5	432.1
1999[4]	47.7	72.8	116.0	165.8	270.1	451.6
Female						
2001	36.6	51.4	76.4	122.2	172.0	345.6
2000[4]	36.2	54.1	80.1	125.7	201.3	382.5
1999[4]	28.7	48.4	82.0	130.8	182.6	400.2
Syphilis (A50-A53)						
All origins[3]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Syphilis (A50-A53)														
Hispanic														
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Syphilis (A50-A53)						
Hispanic						
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic White						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Syphilis (A50-A53)														
Non-Hispanic Black														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Acute poliomyelitis (A80)														
All origins[3]														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Syphilis (A50-A53)						
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Acute poliomyelitis (A80)						
All origins[3]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute poliomyelitis (A80)														
Hispanic														
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute poliomyelitis (A80)						
Hispanic						
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic White						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute poliomyelitis (A80)														
Non-Hispanic Black														
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Arthropod-borne viral encephalitis (A83-A84,A85.2)														
All origins[3]														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute poliomyelitis (A80)						
Non-Hispanic Black						
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Arthropod-borne viral encephalitis (A83-A84,A85.2)						
All origins[3]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Arthropod-borne viral encephalitis (A83-A84,A85.2)														
Non-Hispanic[5]														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Arthropod-borne viral encephalitis (A83-A84,A85.2)						
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic White						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Arthropod-borne viral encephalitis (A83-A84,A85.2)														
Non-Hispanic Black														
Female														
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Measles (B05)														
All origins[3]														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Arthropod-borne viral encephalitis (A83-A84,A85.2)						
Non-Hispanic Black						
Female						
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Measles (B05)						
All origins[3]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Measles (B05)														
Non-Hispanic[5]														
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Measles (B05)						
Non-Hispanic[5]						
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic White						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Viral hepatitis (B15-B19)														
All origins[3]														
Both sexes														
2001	2.0	*	*	*	*	*	*	*	0.3	0.9	2.6	5.3	5.2	4.1
2000[4]	1.9	*	*	*	*	*	*	0.1	0.3	1.0	2.5	5.2	4.8	3.7
1999[4]	1.7	*	*	*	*	*	*	*	0.3	0.9	2.3	4.4	3.8	3.3
Male														
2001	2.5	*	*	*	*	*	*	*	0.3	1.2	3.7	8.1	8.2	5.9
2000[4]	2.4	*	*	*	*	*	*	*	0.3	1.3	3.5	8.0	7.4	5.2
1999[4]	2.2	*	*	*	*	*	*	*	0.4	1.2	3.5	6.7	5.9	4.7
Female														
2001	1.4	*	*	*	*	*	*	*	0.2	0.6	1.5	2.5	2.4	2.5
2000[4]	1.4	*	*	*	*	*	*	*	0.2	0.6	1.5	2.5	2.4	2.2
1999[4]	1.3	*	*	*	*	*	*	*	0.2	0.6	1.2	2.2	1.7	2.1
Hispanic														
Both sexes														
2001	2.1	*	*	*	*	*	*	*	*	1.2	3.7	8.1	7.5	11.1
2000[4]	2.0	*	*	*	*	*	*	*	*	1.4	3.8	6.7	7.3	7.0
1999[4]	1.9	*	*	*	*	*	*	*	*	1.4	3.0	5.9	8.1	7.6
Male														
2001	2.6	*	*	*	*	*	*	*	*	1.5	5.2	12.4	12.1	17.0
2000[4]	2.5	*	*	*	*	*	*	*	*	2.0	5.9	10.3	11.1	9.4
1999[4]	2.3	*	*	*	*	*	*	*	*	2.1	4.8	9.1	12.3	9.9
Female														
2001	1.7	*	*	*	*	*	*	*	*	*	2.1	3.8	3.2	5.8
2000[4]	1.5	*	*	*	*	*	*	*	*	*	*	3.2	3.6	4.8
1999[4]	1.5	*	*	*	*	*	*	*	*	*	*	2.7	4.1	5.4
Non-Hispanic[5]														
Both sexes														
2001	1.9	*	*	*	*	*	*	*	0.3	0.9	2.5	5.0	5.0	3.5
2000[4]	1.9	*	*	*	*	*	*	0.1	0.2	0.9	2.3	5.0	4.6	3.4
1999[4]	1.7	*	*	*	*	*	*	*	0.3	0.8	2.2	4.3	3.4	3.0
Male														
2001	2.5	*	*	*	*	*	*	*	0.4	1.1	3.5	7.7	7.8	5.0
2000[4]	2.4	*	*	*	*	*	*	*	0.3	1.2	3.2	7.7	7.0	4.8
1999[4]	2.2	*	*	*	*	*	*	*	0.4	1.1	3.3	6.5	5.3	4.3
Female														
2001	1.4	*	*	*	*	*	*	*	*	0.6	1.5	2.4	2.3	2.2
2000[4]	1.4	*	*	*	*	*	*	*	*	0.6	1.5	2.4	2.3	2.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Viral hepatitis (B15-B19)						
All origins[3]						
Both sexes						
2001	3.8	4.3	4.9	5.1	4.6	3.8
2000[4]	3.8	4.3	4.9	5.4	5.2	4.0
1999[4]	3.3	4.5	5.1	5.3	5.4	4.7
Male						
2001	4.5	4.7	4.9	5.1	5.5	4.4
2000[4]	4.7	4.6	5.0	6.5	6.2	5.7
1999[4]	3.8	4.9	5.4	6.1	7.3	6.9
Female						
2001	3.3	3.9	4.9	5.0	4.1	3.5
2000[4]	3.1	4.1	4.9	4.6	4.7	3.3
1999[4]	2.8	4.1	4.9	4.7	4.2	3.9
Hispanic						
Both sexes						
2001	9.5	8.2	12.1	12.2	*	*
2000[4]	10.7	10.3	12.2	12.9	13.4	*
1999[4]	8.0	11.7	13.3	13.6	13.5	*
Male						
2001	10.3	*	*	*	*	*
2000[4]	14.7	10.4	11.7	*	*	*
1999[4]	9.5	12.4	13.7	*	*	*
Female						
2001	8.7	10.0	15.0	13.6	*	*
2000[4]	7.2	10.3	12.5	13.1	*	*
1999[4]	6.7	11.3	13.0	15.4	*	*
Non-Hispanic[5]						
Both sexes						
2001	3.4	4.0	4.4	4.7	4.4	3.6
2000[4]	3.3	3.9	4.5	5.1	4.9	3.8
1999[4]	2.9	4.0	4.7	4.9	5.1	4.7
Male						
2001	4.0	4.6	4.7	4.8	5.3	4.3
2000[4]	3.9	4.2	4.6	6.2	5.9	5.4
1999[4]	3.4	4.4	4.9	5.9	7.4	6.8
Female						
2001	2.8	3.4	4.3	4.6	3.9	3.3
2000[4]	2.7	3.6	4.4	4.3	4.3	3.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Viral hepatitis (B15-B19)														
Non-Hispanic[5]														
Female														
1999[4]	1.2	*	*	*	*	*	*	*	0.3	0.6	1.2	2.1	1.5	1.8
Non-Hispanic White														
Both sexes														
2001	1.8	*	*	*	*	*	*	*	0.2	0.9	2.4	4.8	4.4	3.0
2000[4]	1.8	*	*	*	*	*	*	*	0.2	0.9	2.2	4.7	3.9	2.7
1999[4]	1.6	*	*	*	*	*	*	*	0.3	0.8	2.0	3.9	2.9	2.4
Male														
2001	2.5	*	*	*	*	*	*	*	0.4	1.2	3.4	7.4	7.0	4.3
2000[4]	2.3	*	*	*	*	*	*	*	*	1.2	3.0	7.3	6.1	3.9
1999[4]	2.1	*	*	*	*	*	*	*	0.4	1.1	3.0	5.7	4.5	3.6
Female														
2001	1.2	*	*	*	*	*	*	*	*	0.6	1.4	2.2	1.8	1.7
2000[4]	1.3	*	*	*	*	*	*	*	*	0.6	1.4	2.1	1.8	1.5
1999[4]	1.1	*	*	*	*	*	*	*	*	0.5	1.1	2.1	1.3	1.4
Non-Hispanic Black														
Both sexes														
2001	2.3	*	*	*	*	*	*	*	*	1.0	3.4	6.8	9.6	7.3
2000[4]	2.2	*	*	*	*	*	*	*	*	1.0	3.2	7.4	8.8	6.9
1999[4]	2.1	*	*	*	*	*	*	*	*	0.9	3.5	7.2	6.8	5.7
Male														
2001	2.8	*	*	*	*	*	*	*	*	*	4.9	10.1	14.3	10.1
2000[4]	2.9	*	*	*	*	*	*	*	*	1.6	4.4	11.2	13.8	10.3
1999[4]	2.6	*	*	*	*	*	*	*	*	*	5.1	12.2	11.4	7.7
Female														
2001	1.8	*	*	*	*	*	*	*	*	*	2.0	3.9	5.6	4.9
2000[4]	1.6	*	*	*	*	*	*	*	*	*	2.2	4.2	4.5	4.0
1999[4]	1.6	*	*	*	*	*	*	*	*	*	2.1	2.9	2.9	4.1
Human immunodeficiency virus (HIV) disease (B20-B24)														
All origins[3]														
Both sexes														
2001	5.0	*	*	*	0.1	0.2	0.9	3.1	7.3	12.3	13.7	12.1	8.7	5.8
2000[4]	5.1	*	*	*	0.1	0.2	0.8	3.3	8.7	12.8	13.5	12.7	9.0	5.9
1999[4]	5.3	*	0.2	0.2	0.1	0.2	0.9	3.9	9.5	13.0	14.6	12.4	8.7	5.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Viral hepatitis (B15-B19)						
Non-Hispanic[5]						
Female						
1999[4]	2.5	3.6	4.5	4.2	3.7	3.9
Non-Hispanic White						
Both sexes						
2001	2.9	3.4	3.9	4.4	4.2	3.3
2000[4]	2.7	3.3	4.1	4.9	4.6	3.7
1999[4]	2.5	3.4	4.1	4.5	5.0	4.5
Male						
2001	3.6	4.3	4.4	4.9	5.4	4.4
2000[4]	3.3	3.6	4.2	6.0	5.6	5.4
1999[4]	3.1	4.0	4.6	5.6	7.7	6.5
Female						
2001	2.3	2.7	3.4	4.1	3.5	2.9
2000[4]	2.3	3.2	4.0	4.0	4.0	3.1
1999[4]	1.9	2.9	3.7	3.8	3.4	3.7
Non-Hispanic Black						
Both sexes						
2001	6.2	5.1	6.2	6.2	*	*
2000[4]	4.5	5.3	6.3	5.8	7.2	*
1999[4]	4.3	6.7	6.9	6.5	*	*
Male						
2001	7.5	*	*	*	*	*
2000[4]	5.3	6.9	7.5	*	*	*
1999[4]	4.6	5.6	*	*	*	*
Female						
2001	5.1	5.4	6.7	7.4	*	*
2000[4]	3.9	4.1	5.4	*	*	*
1999[4]	4.1	7.5	7.6	6.7	*	*
Human immunodeficiency virus (HIV) disease (B20-B24)						
All origins[3]						
Both sexes						
2001	4.5	2.7	1.4	0.9	0.4	*
2000[4]	4.1	2.6	1.7	0.9	*	*
1999[4]	3.5	2.7	1.5	0.7	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Human immunodeficiency virus (HIV) disease (B20-B24)														
All origins[3]														
Male														
2001	7.6	*	*	*	*	0.2	0.8	3.7	10.2	18.4	20.5	19.0	14.4	9.8
2000[4]	7.9	*	*	*	*	*	0.8	3.8	11.9	19.2	20.4	20.3	15.0	10.1
1999[4]	8.2	*	*	0.2	*	*	0.8	5.1	13.6	19.5	22.6	20.0	14.5	10.2
Female														
2001	2.5	*	*	*	*	0.2	1.0	2.5	4.4	6.3	7.0	5.4	3.3	2.0
2000[4]	2.5	*	*	*	*	*	0.7	2.9	5.5	6.4	6.7	5.3	3.4	2.0
1999[4]	2.5	*	*	*	*	*	1.0	2.7	5.4	6.6	6.9	5.0	3.1	1.9
Hispanic														
Both sexes														
2001	5.0	*	*	*	*	*	0.8	2.3	6.6	11.9	16.3	16.4	14.2	10.5
2000[4]	5.4	*	*	*	*	*	*	2.9	8.5	13.5	16.5	18.1	15.4	10.0
1999[4]	5.8	*	*	*	*	*	0.9	3.5	9.6	15.5	19.7	17.1	13.1	9.8
Male														
2001	7.6	*	*	*	*	*	*	2.9	9.4	17.5	24.9	25.0	24.8	17.6
2000[4]	8.2	*	*	*	*	*	*	3.5	12.6	21.0	24.9	28.4	26.2	17.3
1999[4]	8.7	*	*	*	*	*	*	4.9	13.9	23.8	29.2	26.8	20.8	17.6
Female														
2001	2.3	*	*	*	*	*	*	1.5	3.5	5.8	7.3	7.8	4.1	4.2
2000[4]	2.4	*	*	*	*	*	*	2.2	3.9	5.4	7.6	7.9	5.0	*
1999[4]	2.6	*	*	*	*	*	*	1.9	4.6	6.4	9.7	7.5	5.8	*
Non-Hispanic[5]														
Both sexes														
2001	4.9	*	*	*	*	0.2	1.0	3.2	7.4	12.3	13.2	11.5	8.1	5.3
2000[4]	5.0	*	*	*	0.1	0.2	0.8	3.4	8.7	12.5	12.9	12.0	8.4	5.5
1999[4]	5.2	*	*	0.2	0.1	0.2	0.9	3.9	9.5	12.6	13.9	11.7	8.1	5.5
Male														
2001	7.5	*	*	*	*	*	0.8	3.8	10.3	18.4	19.6	18.0	13.2	9.1
2000[4]	7.7	*	*	*	*	*	0.8	3.8	11.7	18.6	19.5	19.2	13.8	9.3
1999[4]	8.0	*	*	*	*	*	0.8	5.1	13.4	18.7	21.4	18.9	13.6	9.5
Female														
2001	2.5	*	*	*	*	*	1.1	2.6	4.4	6.3	6.9	5.1	3.2	1.8
2000[4]	2.5	*	*	*	*	*	0.8	3.0	5.7	6.4	6.4	5.1	3.2	1.9
1999[4]	2.5	*	*	*	*	*	1.0	2.8	5.5	6.6	6.5	4.8	2.8	1.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)						
All origins[3]						
Male						
2001	7.1	4.2	2.7	1.8	*	*
2000[4]	7.0	4.5	2.9	1.8	*	*
1999[4]	5.9	4.8	2.7	1.3	*	*
Female						
2001	2.0	1.4	0.4	*	*	*
2000[4]	1.5	1.0	0.7	*	*	*
1999[4]	1.2	1.0	0.6	*	*	*
Hispanic						
Both sexes						
2001	7.3	6.0	*	*	*	*
2000[4]	8.9	4.7	*	*	*	*
1999[4]	6.6	6.0	*	*	*	*
Male						
2001	11.1	11.3	*	*	*	*
2000[4]	12.7	8.2	*	*	*	*
1999[4]	12.5	10.8	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	5.7	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	4.1	2.4	1.3	0.8	*	*
2000[4]	3.7	2.4	1.6	0.8	*	*
1999[4]	3.1	2.4	1.4	0.7	*	*
Male						
2001	6.6	3.6	2.5	1.6	*	*
2000[4]	6.5	4.1	2.7	1.6	*	*
1999[4]	5.2	4.3	2.5	1.2	*	*
Female						
2001	1.8	1.4	*	*	*	*
2000[4]	1.2	0.9	0.7	*	*	*
1999[4]	1.2	0.9	0.6	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Human immunodeficiency virus (HIV) disease (B20-B24)														
Non-Hispanic White														
Both sexes														
2001	2.2	*	*	*	*	*	0.2	0.9	3.0	6.0	6.1	5.0	3.4	2.6
2000[4]	2.3	*	*	*	*	*	0.2	1.1	3.5	6.3	6.1	5.5	3.7	2.7
1999[4]	2.4	*	*	*	*	*	0.2	1.5	4.4	6.5	6.4	5.2	3.9	2.4
Male														
2001	3.8	*	*	*	*	*	*	1.3	4.9	10.2	10.5	8.8	6.2	4.8
2000[4]	4.0	*	*	*	*	*	*	1.5	5.6	10.6	10.2	9.6	6.5	4.9
1999[4]	4.2	*	*	*	*	*	*	2.2	7.3	10.9	10.9	9.0	7.2	4.4
Female														
2001	0.6	*	*	*	*	*	*	0.5	1.2	1.7	1.7	1.2	0.6	0.6
2000[4]	0.7	*	*	*	*	*	*	0.7	1.5	2.1	1.9	1.4	0.8	0.7
1999[4]	0.7	*	*	*	*	*	*	0.8	1.5	2.0	1.8	1.3	0.8	0.5
Non-Hispanic Black														
Both sexes														
2001	21.6	*	*	*	*	1.1	4.6	15.2	32.3	50.4	59.7	57.6	45.0	29.3
2000[4]	22.0	*	*	*	*	0.7	3.9	15.8	38.4	51.0	58.2	59.5	46.9	29.6
1999[4]	22.3	*	*	0.8	*	0.7	4.2	17.2	39.1	51.3	63.5	59.6	42.8	33.0
Male														
2001	30.8	*	*	*	*	*	3.7	17.6	43.5	71.6	84.0	88.7	72.3	50.7
2000[4]	31.7	*	*	*	*	*	3.8	17.1	49.4	73.0	85.3	93.8	77.5	51.3
1999[4]	32.6	*	*	*	*	*	3.5	21.8	52.2	71.7	96.1	96.6	71.1	57.7
Female														
2001	13.3	*	*	*	*	*	5.4	13.0	22.3	31.5	38.2	30.6	21.7	11.7
2000[4]	13.1	*	*	*	*	*	4.0	14.6	28.5	31.5	34.3	29.8	20.8	11.8
1999[4]	13.0	*	*	*	*	*	4.9	13.1	27.4	33.2	34.9	27.7	18.8	12.8
Malaria (B50-B54)														
All origins[3]														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)						
Non-Hispanic White						
Both sexes						
2001	1.5	0.8	0.6	0.3	*	*
2000[4]	1.5	1.0	0.6	0.4	*	*
1999[4]	1.4	1.1	0.5	0.3	*	*
Male						
2001	2.6	1.4	1.4	*	*	*
2000[4]	2.8	2.0	0.9	0.8	*	*
1999[4]	2.6	2.1	0.9	*	*	*
Female						
2001	0.4	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic Black						
Both sexes						
2001	26.3	16.2	8.5	6.4	*	*
2000[4]	22.2	14.7	11.5	6.0	*	*
1999[4]	18.4	14.9	11.2	4.9	*	*
Male						
2001	43.7	25.8	15.5	12.8	*	*
2000[4]	40.2	25.3	21.9	11.6	*	*
1999[4]	30.5	26.3	21.1	*	*	*
Female						
2001	12.7	9.0	*	*	*	*
2000[4]	8.1	6.9	4.5	*	*	*
1999[4]	8.9	6.5	4.6	*	*	*
Malaria (B50-B54)						
All origins[3]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malaria (B50-B54)														
All origins[3]														
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malaria (B50-B54)						
All origins[3]						
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic White						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malaria (B50-B54)														
Non-Hispanic White														
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49, A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49, B55-B99)														
All origins[3]														
Both sexes														
2001	2.0	4.7	0.4	0.2	0.1	0.2	0.2	0.3	0.4	0.7	0.9	1.5	1.9	2.4
2000[4]	2.0	4.4	0.5	0.1	0.2	0.2	0.3	0.3	0.4	0.7	0.8	1.4	1.8	2.4
1999[4]	2.6	5.3	0.5	0.2	0.2	0.2	0.3	0.5	0.7	1.0	1.4	2.0	2.6	3.3
Male														
2001	2.1	4.9	0.5	*	*	0.2	0.3	0.3	0.5	0.8	1.2	2.1	2.4	3.0
2000[4]	2.1	4.8	0.5	*	*	0.2	0.3	0.3	0.4	0.9	1.0	1.7	2.3	2.8
1999[4]	2.7	6.1	0.6	0.2	0.2	0.2	0.3	0.5	0.8	1.3	1.7	2.4	3.0	3.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malaria (B50-B54)						
Non-Hispanic White						
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49, A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49, B55-B99)						
All origins[3]						
Both sexes						
2001	3.9	5.6	6.9	9.4	12.3	20.1
2000[4]	3.6	4.7	7.0	10.0	12.9	21.2
1999[4]	4.4	6.8	8.9	12.6	16.6	27.0
Male						
2001	4.5	6.2	7.5	11.4	15.9	23.6
2000[4]	4.0	5.7	8.8	12.1	16.2	27.6
1999[4]	5.2	8.0	10.5	15.3	20.4	31.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49, A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49, B55-B99)														
All origins[3]														
Female														
2001	1.9	4.6	0.4	0.2	*	*	*	0.3	0.4	0.6	0.6	0.9	1.5	1.8
2000[4]	1.9	3.9	0.4	*	*	*	0.2	0.3	0.4	0.5	0.6	1.1	1.3	2.1
1999[4]	2.5	4.5	0.4	0.2	*	0.2	0.2	0.5	0.6	0.7	1.1	1.6	2.3	3.1
Hispanic														
Both sexes														
2001	1.1	5.0	*	*	*	*	*	*	*	0.8	0.9	2.1	2.0	3.2
2000[4]	1.2	3.0	*	*	*	*	*	*	*	0.8	1.2	1.7	2.3	3.1
1999[4]	1.5	4.8	*	*	*	*	*	*	0.7	1.4	1.4	2.0	2.7	3.0
Male														
2001	1.3	5.0	*	*	*	*	*	*	*	*	*	2.7	2.9	5.0
2000[4]	1.3	*	*	*	*	*	*	*	*	1.4	*	*	*	4.6
1999[4]	1.5	5.7	*	*	*	*	*	*	*	2.0	1.8	*	3.7	*
Female														
2001	1.0	5.0	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	2.1	4.6	0.5	0.2	0.1	0.2	0.2	0.3	0.4	0.7	0.9	1.4	1.9	2.3
2000[4]	2.1	4.7	0.4	0.1	0.2	0.2	0.2	0.3	0.4	0.7	0.8	1.3	1.7	2.3
1999[4]	2.8	5.5	0.5	0.2	0.2	0.2	0.3	0.5	0.7	1.0	1.4	2.0	2.6	3.3
Male														
2001	2.2	4.8	0.5	*	*	*	0.3	0.3	0.5	0.8	1.2	2.0	2.3	2.9
2000[4]	2.2	5.1	0.5	*	*	0.3	*	0.3	0.4	0.9	1.0	1.6	2.2	2.6
1999[4]	2.8	6.2	0.6	*	0.2	0.2	0.3	0.4	0.8	1.2	1.6	2.4	2.9	3.5
Female														
2001	2.1	4.5	0.4	*	*	*	*	0.3	0.4	0.6	0.6	0.9	1.5	1.8
2000[4]	2.0	4.3	0.4	*	*	*	*	0.3	0.4	0.5	0.6	1.1	1.2	2.1
1999[4]	2.7	4.7	0.4	*	*	*	*	0.5	0.6	0.8	1.1	1.6	2.3	3.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49, A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49, B55-B99)						
All origins[3]						
Female						
2001	3.3	5.1	6.5	8.0	10.1	18.7
2000[4]	3.2	3.9	5.6	8.5	10.9	18.6
1999[4]	3.7	5.9	7.7	10.8	14.4	25.1
Hispanic						
Both sexes						
2001	2.8	4.9	7.1	10.5	9.9	15.0
2000[4]	3.2	4.5	7.3	9.5	14.5	19.2
1999[4]	4.8	8.1	7.2	13.9	14.7	26.0
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	6.0	9.3	*	16.4	*	*
Female						
2001	*	*	*	9.7	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	7.2	8.0	12.1	*	28.1
Non-Hispanic[5]						
Both sexes						
2001	4.0	5.7	6.9	9.3	12.4	20.3
2000[4]	3.6	4.7	7.0	9.9	12.8	21.2
1999[4]	4.4	6.7	9.0	12.5	16.7	27.0
Male						
2001	4.6	6.2	7.3	11.4	16.0	24.0
2000[4]	4.0	5.9	8.8	12.1	15.9	27.8
1999[4]	5.1	7.9	10.7	15.2	20.6	32.3
Female						
2001	3.4	5.2	6.5	7.9	10.2	18.8
2000[4]	3.2	3.8	5.6	8.4	10.9	18.6
1999[4]	3.7	5.7	7.6	10.7	14.4	24.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49, A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49, B55-B99)														
Non-Hispanic White														
Both sexes														
2001	2.1	3.3	0.4	0.2	*	0.2	0.2	0.2	0.3	0.6	0.8	1.3	1.7	2.0
2000[4]	2.1	3.5	0.4	*	0.2	0.2	0.2	0.2	0.3	0.5	0.7	1.2	1.5	2.2
1999[4]	2.7	4.4	0.5	0.2	*	0.2	0.2	0.3	0.5	0.7	1.0	1.5	2.1	2.9
Male														
2001	2.2	3.3	0.4	*	*	*	*	*	0.4	0.6	1.0	1.8	2.1	2.5
2000[4]	2.2	3.9	*	*	*	*	*	*	*	0.6	0.8	1.5	1.9	2.5
1999[4]	2.7	5.5	0.6	*	*	*	*	*	0.5	1.0	1.3	2.0	2.4	3.3
Female														
2001	2.1	3.4	*	*	*	*	*	*	*	0.5	0.5	0.7	1.4	1.5
2000[4]	2.0	3.1	*	*	*	*	*	*	0.3	0.4	0.5	1.0	1.0	1.9
1999[4]	2.6	3.3	*	*	*	*	*	0.4	0.4	0.5	0.7	1.1	1.9	2.5
Non-Hispanic Black														
Both sexes														
2001	2.3	10.0	*	*	*	*	*	*	1.1	1.5	1.8	2.6	3.5	4.5
2000[4]	2.3	10.4	*	*	*	*	*	*	1.3	1.7	1.7	2.7	3.5	4.0
1999[4]	3.6	10.0	1.0	*	*	*	*	1.2	2.2	2.6	3.6	5.2	6.0	6.5
Male														
2001	2.5	11.5	*	*	*	*	*	*	*	1.7	2.6	3.7	4.4	5.5
2000[4]	2.6	10.6	*	*	*	*	*	*	*	2.3	2.2	3.3	4.8	4.6
1999[4]	3.7	10.0	*	*	*	*	*	*	2.7	2.7	3.6	6.0	6.6	5.9
Female														
2001	2.1	8.4	*	*	*	*	*	*	*	*	*	1.6	2.8	3.7
2000[4]	2.1	10.2	*	*	*	*	*	*	*	*	*	2.1	2.3	3.5
1999[4]	3.5	10.0	*	*	*	*	*	*	1.7	2.5	3.6	4.5	5.5	7.0
Malignant neoplasms (C00-C97)														
All origins[3]														
Both sexes														
2001	194.4	1.6	2.7	2.4	2.5	3.6	4.9	7.3	12.6	24.6	48.7	92.4	164.8	280.7
2000[4]	196.5	2.4	2.7	2.4	2.6	3.7	5.1	7.0	12.5	24.5	48.8	93.6	166.2	289.0
1999[4]	197.0	1.8	2.7	2.5	2.5	3.7	5.3	7.3	12.5	25.2	49.4	93.7	167.0	295.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49, A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49, B55-B99)						
Non-Hispanic White						
Both sexes						
2001	3.7	5.5	6.6	9.2	12.2	20.2
2000[4]	3.2	4.6	6.8	9.8	12.6	21.0
1999[4]	3.9	6.3	8.6	11.9	16.1	26.4
Male						
2001	4.3	5.9	7.1	11.3	16.0	24.2
2000[4]	3.6	5.6	8.6	11.6	15.8	27.0
1999[4]	4.4	7.3	10.4	14.7	18.9	30.8
Female						
2001	3.1	5.1	6.2	7.7	10.0	18.6
2000[4]	2.8	3.7	5.3	8.5	10.8	18.6
1999[4]	3.4	5.4	7.2	10.0	14.4	24.7
Non-Hispanic Black						
Both sexes						
2001	6.2	7.1	9.8	9.9	13.5	20.9
2000[4]	6.0	6.2	10.1	11.6	16.1	24.2
1999[4]	8.2	10.9	13.3	19.1	23.8	34.3
Male						
2001	7.5	9.6	10.0	10.4	16.4	*
2000[4]	7.4	8.8	12.6	16.9	19.7	31.7
1999[4]	10.4	12.5	16.3	22.2	42.5	46.4
Female						
2001	5.1	5.2	9.6	9.6	12.1	20.5
2000[4]	4.9	4.3	8.4	8.4	14.3	21.4
1999[4]	6.5	9.6	11.3	17.2	14.5	29.8
Malignant neoplasms (C00-C97)						
All origins[3]						
Both sexes						
2001	453.3	673.2	943.5	1,211.7	1,465.9	1,765.6
2000[4]	463.4	690.6	951.6	1,222.5	1,505.3	1,819.4
1999[4]	471.6	700.2	963.4	1,224.0	1,493.1	1,805.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms (C00-C97)														
All origins[3]														
Male														
2001	205.3	1.5	2.9	2.4	2.6	4.3	5.7	7.2	11.2	21.0	44.0	91.7	174.2	311.6
2000[4]	207.2	2.6	3.0	2.6	2.9	4.3	5.9	7.1	11.3	21.2	44.4	93.5	173.9	321.3
1999[4]	208.9	1.9	2.9	2.6	2.7	4.3	6.1	7.6	11.6	21.6	45.5	94.7	174.9	330.6
Female														
2001	183.9	1.8	2.5	2.4	2.3	2.9	4.1	7.4	14.0	28.1	53.5	93.2	155.9	251.7
2000[4]	186.2	2.3	2.5	2.2	2.2	3.0	4.2	7.0	13.7	27.8	53.2	93.7	158.8	258.9
1999[4]	185.6	1.7	2.5	2.3	2.3	3.0	4.4	7.0	13.4	28.7	53.3	92.7	159.5	262.3
Hispanic														
Both sexes														
2001	60.5	*	2.8	2.4	2.7	4.5	4.1	5.7	8.9	18.3	34.4	61.7	110.3	172.3
2000[4]	59.9	*	2.5	2.5	3.4	3.8	4.5	5.8	9.0	18.2	33.9	64.3	105.4	180.7
1999[4]	59.6	*	2.5	2.5	3.0	3.1	5.1	5.8	9.1	17.3	33.3	65.0	110.9	179.3
Male														
2001	62.2	*	2.9	2.6	2.4	5.7	5.4	5.2	7.5	15.1	29.9	60.1	109.9	189.4
2000[4]	61.3	*	2.9	3.0	4.1	4.3	4.8	5.7	8.2	13.8	28.0	60.9	104.2	199.7
1999[4]	61.2	*	2.3	2.5	3.1	3.4	5.8	5.6	7.6	13.9	27.2	64.4	111.7	208.3
Female														
2001	58.7	*	2.7	2.1	3.0	3.2	2.7	6.4	10.4	21.8	39.2	63.3	110.7	156.7
2000[4]	58.5	*	2.2	2.0	2.6	3.4	4.1	5.8	9.9	23.0	40.0	67.6	106.5	163.5
1999[4]	58.0	*	2.7	2.5	2.9	2.7	4.2	6.1	10.7	20.9	39.6	65.7	110.1	153.2
Non-Hispanic[5]														
Both sexes														
2001	213.9	1.6	2.7	2.4	2.4	3.4	5.1	7.7	13.3	25.4	50.3	95.3	169.1	288.6
2000[4]	215.6	2.5	2.7	2.3	2.4	3.6	5.2	7.2	13.1	25.3	50.3	96.1	170.7	296.5
1999[4]	215.5	1.7	2.7	2.5	2.4	3.8	5.3	7.6	13.0	26.2	51.0	96.1	171.2	303.1
Male														
2001	227.2	1.6	2.9	2.4	2.7	4.0	5.8	7.7	11.9	21.8	45.5	94.7	179.2	320.0
2000[4]	228.7	2.6	2.9	2.5	2.6	4.3	6.2	7.3	11.8	22.3	46.2	96.4	179.1	329.4
1999[4]	229.9	1.6	3.0	2.7	2.6	4.5	6.1	8.1	12.3	22.7	47.4	97.2	179.5	338.6
Female														
2001	201.2	1.6	2.5	2.5	2.2	2.8	4.4	7.6	14.6	29.0	55.0	95.8	159.5	259.0
2000[4]	203.1	2.4	2.6	2.2	2.2	3.0	4.2	7.2	14.3	28.3	54.4	95.8	162.7	265.6
1999[4]	201.9	1.8	2.5	2.3	2.2	3.1	4.5	7.1	13.7	29.7	54.6	95.0	163.1	269.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms (C00-C97)						
All origins[3]						
Male						
2001	526.7	814.1	1,178.0	1,568.4	1,978.7	2,613.6
2000[4]	535.7	834.7	1,190.4	1,586.0	2,050.4	2,710.7
1999[4]	550.1	848.2	1,212.9	1,600.1	2,046.0	2,722.2
Female						
2001	386.7	551.9	757.2	962.0	1,158.6	1,410.7
2000[4]	398.0	567.2	763.5	969.3	1,183.8	1,456.4
1999[4]	400.6	573.7	768.4	962.5	1,169.5	1,436.5
Hispanic						
Both sexes						
2001	291.3	413.5	616.9	846.0	1,033.1	1,363.7
2000[4]	272.5	422.5	633.8	851.8	1,138.5	1,383.5
1999[4]	282.8	448.9	611.2	834.8	1,094.3	1,399.6
Male						
2001	340.6	519.1	802.4	1,171.7	1,423.2	1,967.4
2000[4]	323.3	521.3	820.6	1,140.5	1,637.2	2,049.7
1999[4]	345.2	567.6	802.5	1,120.3	1,520.0	1,945.3
Female						
2001	248.7	328.0	475.3	614.9	791.4	1,068.6
2000[4]	228.8	342.4	492.3	647.3	835.0	1,056.5
1999[4]	228.9	352.8	467.3	633.2	840.3	1,136.5
Non-Hispanic[5]						
Both sexes						
2001	464.3	690.0	960.9	1,226.9	1,480.7	1,776.6
2000[4]	476.3	706.7	967.3	1,236.6	1,515.9	1,831.1
1999[4]	484.0	714.7	980.0	1,237.8	1,503.6	1,816.1
Male						
2001	538.6	832.2	1,196.9	1,584.2	1,998.0	2,634.8
2000[4]	549.5	852.4	1,207.8	1,602.6	2,061.5	2,732.4
1999[4]	563.0	863.4	1,231.0	1,616.8	2,060.5	2,747.5
Female						
2001	396.6	567.0	773.0	976.9	1,171.2	1,419.6
2000[4]	409.7	581.3	777.5	981.8	1,194.4	1,466.6
1999[4]	412.2	587.0	783.4	974.5	1,178.0	1,443.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms (C00-C97)														
Non-Hispanic White														
Both sexes														
2001	230.6	1.8	2.9	2.5	2.3	3.4	5.1	7.4	12.8	24.5	47.7	90.2	162.2	279.7
2000[4]	232.2	2.6	2.8	2.4	2.4	3.5	5.1	7.3	12.7	24.0	46.9	90.3	163.6	288.2
1999[4]	231.5	1.6	3.0	2.4	2.3	3.7	5.1	7.6	12.6	24.9	48.1	90.4	163.5	293.3
Male														
2001	243.4	1.7	3.1	2.3	2.4	4.0	5.8	7.7	12.0	21.2	43.2	89.1	169.4	305.9
2000[4]	244.4	2.9	3.1	2.5	2.6	4.1	6.1	7.4	11.8	21.5	42.9	90.1	168.5	314.5
1999[4]	245.1	1.7	3.3	2.7	2.5	4.2	5.8	8.1	12.0	22.1	44.8	89.9	168.3	323.3
Female														
2001	218.4	1.8	2.8	2.6	2.2	2.8	4.4	7.2	13.7	27.7	52.2	91.3	155.1	254.6
2000[4]	220.6	2.4	2.4	2.4	2.2	2.9	4.1	7.1	13.5	26.5	50.9	90.5	158.8	263.2
1999[4]	218.4	*	2.7	2.1	2.1	3.0	4.3	7.1	13.1	27.7	51.5	91.0	158.8	264.6
Non-Hispanic Black														
Both sexes														
2001	172.8	*	2.0	2.4	2.9	3.7	5.3	9.6	17.0	33.7	72.2	141.4	246.5	412.8
2000[4]	174.9	*	2.8	2.2	2.4	4.5	6.1	8.3	16.3	35.4	76.1	148.6	257.2	415.1
1999[4]	176.9	*	2.0	2.5	3.1	4.6	6.9	8.8	16.4	36.9	74.9	147.7	260.2	437.3
Male														
2001	190.8	*	2.4	2.6	3.4	4.3	6.1	8.9	12.9	27.6	64.3	147.2	286.7	504.5
2000[4]	194.9	*	2.6	2.5	2.7	5.5	7.4	7.6	13.3	28.3	71.8	155.3	299.9	521.3
1999[4]	197.4	*	2.0	2.4	3.4	5.6	7.7	9.2	14.5	28.6	69.9	164.2	305.9	543.1
Female														
2001	156.5	*	*	2.2	2.3	3.0	4.5	10.3	20.8	39.2	79.2	136.3	212.4	337.8
2000[4]	156.9	*	3.0	1.8	2.0	3.6	4.8	8.9	18.9	41.8	79.8	142.8	220.9	328.1
1999[4]	158.3	*	2.0	2.7	2.7	3.5	6.1	8.4	18.1	44.3	79.3	133.4	221.4	351.1
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)														
All origins[3]														
Both sexes														
2001	2.7	*	*	*	*	*	*	*	0.2	0.4	0.8	2.1	3.5	6.1
2000[4]	2.7	*	*	*	*	*	*	*	0.2	0.4	0.9	2.0	3.8	5.7
1999[4]	2.7	*	*	*	*	*	*	*	0.1	0.4	0.9	2.3	3.6	5.8
Male														
2001	3.7	*	*	*	*	*	*	*	0.2	0.5	1.2	3.2	5.8	10.0
2000[4]	3.6	*	*	*	*	*	*	*	*	0.5	1.4	3.1	6.0	9.5
1999[4]	3.6	*	*	*	*	*	*	*	*	0.5	1.3	3.6	5.7	9.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms (C00-C97)						
Non-Hispanic White						
Both sexes						
2001	458.2	682.7	955.9	1,222.7	1,473.0	1,775.2
2000[4]	470.0	701.3	959.9	1,227.7	1,512.6	1,826.6
1999[4]	474.6	707.7	971.9	1,226.4	1,498.7	1,808.3
Male						
2001	525.6	815.0	1,178.6	1,562.9	1,968.7	2,616.8
2000[4]	535.3	837.1	1,182.9	1,574.8	2,036.1	2,714.1
1999[4]	544.9	846.1	1,209.4	1,586.9	2,037.9	2,721.3
Female						
2001	395.8	565.8	775.0	981.8	1,173.8	1,425.1
2000[4]	409.5	582.0	780.3	983.7	1,201.6	1,468.7
1999[4]	409.6	586.3	782.7	973.5	1,180.6	1,444.2
Non-Hispanic Black						
Both sexes						
2001	598.1	871.7	1,172.3	1,446.2	1,735.8	1,947.5
2000[4]	616.2	874.9	1,193.3	1,497.2	1,718.5	2,015.1
1999[4]	644.3	891.9	1,206.6	1,516.9	1,715.6	2,024.0
Male						
2001	756.4	1,154.2	1,619.6	2,100.9	2,672.1	3,238.5
2000[4]	783.3	1,154.5	1,675.4	2,183.1	2,709.5	3,290.1
1999[4]	830.7	1,174.9	1,663.5	2,224.9	2,656.8	3,361.1
Female						
2001	473.7	663.1	874.2	1,055.5	1,258.9	1,469.3
2000[4]	484.4	668.7	873.2	1,083.7	1,219.7	1,541.6
1999[4]	498.2	682.1	903.4	1,090.3	1,249.0	1,524.4
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)						
All origins[3]						
Both sexes						
2001	7.9	9.4	11.6	13.5	15.2	18.5
2000[4]	8.1	9.9	11.5	12.6	14.0	19.4
1999[4]	8.1	9.3	11.3	13.5	15.4	19.3
Male						
2001	13.0	14.3	17.9	19.6	22.1	25.8
2000[4]	12.7	15.2	16.9	18.7	20.2	27.1
1999[4]	13.1	14.3	17.4	20.1	22.3	27.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)														
All origins[3]														
Female														
2001	1.8	*	*	*	*	*	*	*	*	0.3	0.4	1.0	1.4	2.4
2000[4]	1.8	*	*	*	*	*	*	*	*	0.2	0.4	1.0	1.6	2.2
1999[4]	1.8	*	*	*	*	*	*	*	*	0.3	0.4	1.0	1.6	2.7
Hispanic														
Both sexes														
2001	0.8	*	*	*	*	*	*	*	*	*	*	1.3	2.1	2.8
2000[4]	0.8	*	*	*	*	*	*	*	*	*	*	*	2.0	4.2
1999[4]	0.8	*	*	*	*	*	*	*	*	*	*	*	1.7	3.2
Male														
2001	1.2	*	*	*	*	*	*	*	*	*	*	*	3.3	4.6
2000[4]	1.2	*	*	*	*	*	*	*	*	*	*	*	3.2	7.5
1999[4]	1.2	*	*	*	*	*	*	*	*	*	*	*	*	6.2
Female														
2001	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	3.0	*	*	*	*	*	*	*	0.2	0.5	0.9	2.1	3.6	6.3
2000[4]	2.9	*	*	*	*	*	*	*	0.2	0.4	1.0	2.1	3.9	5.8
1999[4]	2.9	*	*	*	*	*	*	*	0.2	0.4	0.9	2.4	3.8	5.9
Male														
2001	4.1	*	*	*	*	*	*	*	0.3	0.6	1.3	3.3	6.0	10.4
2000[4]	3.9	*	*	*	*	*	*	*	*	0.6	1.5	3.3	6.2	9.6
1999[4]	4.0	*	*	*	*	*	*	*	*	0.5	1.4	3.7	6.0	9.1
Female														
2001	1.9	*	*	*	*	*	*	*	*	0.3	0.5	1.0	1.4	2.5
2000[4]	2.0	*	*	*	*	*	*	*	*	0.3	0.5	1.0	1.7	2.2
1999[4]	2.0	*	*	*	*	*	*	*	*	0.3	0.4	1.1	1.7	2.9
Non-Hispanic White														
Both sexes														
2001	3.1	*	*	*	*	*	*	*	0.2	0.3	0.7	1.8	3.3	5.7
2000[4]	3.0	*	*	*	*	*	*	*	*	0.3	0.8	1.8	3.3	5.1
1999[4]	3.0	*	*	*	*	*	*	*	*	0.3	0.8	1.8	3.1	5.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)						
All origins[3]						
Female						
2001	3.3	5.1	6.6	9.3	11.1	15.4
2000[4]	3.9	5.4	7.2	8.3	10.4	16.3
1999[4]	3.6	5.1	6.6	9.0	11.3	16.1
Hispanic						
Both sexes						
2001	5.0	6.5	7.3	11.0	*	19.4
2000[4]	4.7	6.7	6.9	8.3	11.7	18.6
1999[4]	5.1	6.4	7.7	11.6	15.9	14.0
Male						
2001	8.7	11.0	12.4	17.7	*	*
2000[4]	8.1	11.9	12.6	*	*	40.3
1999[4]	8.3	12.8	14.8	21.1	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	8.1	9.5	11.8	13.6	15.4	18.4
2000[4]	8.3	10.0	11.7	12.8	14.1	19.5
1999[4]	8.2	9.5	11.5	13.6	15.4	19.4
Male						
2001	13.3	14.5	18.2	19.5	22.4	25.8
2000[4]	13.0	15.3	17.1	19.0	20.1	26.5
1999[4]	13.4	14.3	17.5	20.0	22.7	27.3
Female						
2001	3.4	5.3	6.8	9.4	11.3	15.3
2000[4]	4.0	5.5	7.5	8.4	10.5	16.6
1999[4]	3.6	5.3	6.8	9.1	11.1	16.2
Non-Hispanic White						
Both sexes						
2001	7.5	8.9	11.6	13.6	15.7	18.9
2000[4]	7.4	9.5	11.3	12.7	14.1	19.8
1999[4]	7.6	9.0	11.1	13.3	15.5	19.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)														
Non-Hispanic White														
Male														
2001	4.1	*	*	*	*	*	*	*	*	0.4	1.1	2.7	5.4	9.2
2000[4]	3.9	*	*	*	*	*	*	*	*	0.4	1.1	2.8	5.0	8.3
1999[4]	3.9	*	*	*	*	*	*	*	*	0.4	1.2	2.8	4.9	7.7
Female														
2001	2.1	*	*	*	*	*	*	*	*	*	0.4	0.8	1.2	2.3
2000[4]	2.1	*	*	*	*	*	*	*	*	0.3	0.5	0.9	1.6	2.0
1999[4]	2.1	*	*	*	*	*	*	*	*	0.3	0.4	0.9	1.3	2.6
Non-Hispanic Black														
Both sexes														
2001	3.0	*	*	*	*	*	*	*	*	0.9	1.7	4.4	6.4	12.5
2000[4]	3.1	*	*	*	*	*	*	*	*	*	1.7	4.1	8.5	12.3
1999[4]	3.3	*	*	*	*	*	*	*	*	0.9	1.6	6.0	9.0	12.7
Male														
2001	4.6	*	*	*	*	*	*	*	*	*	2.4	7.1	11.2	22.7
2000[4]	4.9	*	*	*	*	*	*	*	*	*	2.7	6.3	16.3	22.0
1999[4]	5.0	*	*	*	*	*	*	*	*	*	2.4	10.4	14.8	21.8
Female														
2001	1.5	*	*	*	*	*	*	*	*	*	*	2.0	2.3	4.1
2000[4]	1.5	*	*	*	*	*	*	*	*	*	*	2.2	*	4.3
1999[4]	1.7	*	*	*	*	*	*	*	*	*	*	2.2	4.0	5.3
Malignant neoplasm of esophagus (C15)														
All origins[3]														
Both sexes														
2001	4.4	*	*	*	*	*	*	*	0.1	0.3	1.1	2.4	5.0	7.9
2000[4]	4.3	*	*	*	*	*	*	*	0.1	0.3	1.0	2.4	4.9	8.8
1999[4]	4.3	*	*	*	*	*	*	*	0.1	0.3	1.0	2.4	4.7	8.7
Male														
2001	6.9	*	*	*	*	*	*	*	*	0.5	1.8	4.1	8.7	13.7
2000[4]	6.7	*	*	*	*	*	*	*	0.2	0.6	1.8	4.2	8.7	15.2
1999[4]	6.6	*	*	*	*	*	*	*	0.2	0.6	1.7	4.2	8.3	15.3
Female														
2001	2.0	*	*	*	*	*	*	*	*	*	0.3	0.7	1.4	2.4
2000[4]	2.1	*	*	*	*	*	*	*	*	*	0.2	0.6	1.4	2.8
1999[4]	2.0	*	*	*	*	*	*	*	*	*	0.3	0.6	1.3	2.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)						
Non-Hispanic White						
Male						
2001	11.9	13.4	17.4	19.2	22.5	26.4
2000[4]	11.4	14.1	16.1	18.3	19.7	27.4
1999[4]	12.3	13.5	16.9	19.1	22.3	27.5
Female						
2001	3.3	5.0	6.9	9.7	11.6	15.7
2000[4]	3.7	5.5	7.4	8.7	10.7	16.8
1999[4]	3.4	5.1	6.6	9.2	11.4	16.8
Non-Hispanic Black						
Both sexes						
2001	13.4	15.5	16.0	14.2	13.5	13.1
2000[4]	15.6	15.0	16.5	14.5	16.1	14.9
1999[4]	14.7	14.5	16.6	17.4	15.8	14.5
Male						
2001	25.5	25.5	29.6	24.6	21.3	*
2000[4]	28.1	27.7	29.7	28.9	29.2	*
1999[4]	25.7	24.2	27.0	33.4	29.2	25.0
Female						
2001	3.9	8.1	6.9	7.9	9.6	10.7
2000[4]	5.7	5.7	7.7	5.8	9.5	14.4
1999[4]	6.1	7.3	9.6	7.9	9.2	10.7
Malignant neoplasm of esophagus (C15)						
All origins[3]						
Both sexes						
2001	13.0	18.5	22.9	26.1	26.5	25.8
2000[4]	13.2	17.9	22.1	24.5	27.2	25.9
1999[4]	13.0	18.2	22.0	24.3	24.6	26.8
Male						
2001	22.6	32.6	39.9	46.5	48.8	48.1
2000[4]	23.0	30.7	38.0	43.9	48.8	47.8
1999[4]	21.9	31.7	39.3	44.1	44.8	48.0
Female						
2001	4.2	6.4	9.4	11.9	13.2	16.5
2000[4]	4.4	7.1	9.6	11.0	14.5	17.1
1999[4]	5.0	6.7	8.5	10.5	12.8	18.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of esophagus (C15)														
Hispanic														
Both sexes														
2001	1.1	*	*	*	*	*	*	*	*	*	*	1.1	2.8	3.2
2000[4]	1.1	*	*	*	*	*	*	*	*	*	*	*	1.7	3.9
1999[4]	1.1	*	*	*	*	*	*	*	*	*	*	1.3	1.7	3.9
Male														
2001	1.7	*	*	*	*	*	*	*	*	*	*	*	5.8	5.0
2000[4]	1.6	*	*	*	*	*	*	*	*	*	*	*	3.2	5.7
1999[4]	1.7	*	*	*	*	*	*	*	*	*	*	2.4	*	7.9
Female														
2001	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	4.9	*	*	*	*	*	*	*	0.1	0.3	1.1	2.5	5.1	8.2
2000[4]	4.8	*	*	*	*	*	*	*	0.2	0.4	1.1	2.5	5.2	9.1
1999[4]	4.7	*	*	*	*	*	*	*	0.1	0.3	1.0	2.5	4.9	9.0
Male														
2001	7.7	*	*	*	*	*	*	*	*	0.5	1.9	4.3	9.0	14.3
2000[4]	7.5	*	*	*	*	*	*	*	0.3	0.7	1.9	4.5	9.1	15.9
1999[4]	7.4	*	*	*	*	*	*	*	0.2	0.6	1.8	4.4	8.7	15.8
Female														
2001	2.2	*	*	*	*	*	*	*	*	*	0.4	0.8	1.5	2.4
2000[4]	2.3	*	*	*	*	*	*	*	*	*	0.2	0.7	1.5	2.8
1999[4]	2.2	*	*	*	*	*	*	*	*	*	0.3	0.6	1.3	2.7
Non-Hispanic White														
Both sexes														
2001	5.1	*	*	*	*	*	*	*	*	0.4	1.1	2.4	4.8	7.4
2000[4]	5.1	*	*	*	*	*	*	*	0.2	0.4	1.0	2.4	4.8	8.4
1999[4]	4.9	*	*	*	*	*	*	*	0.2	0.4	0.9	2.3	4.5	8.3
Male														
2001	8.1	*	*	*	*	*	*	*	*	0.6	1.9	4.1	8.3	13.1
2000[4]	8.0	*	*	*	*	*	*	*	0.3	0.8	1.9	4.4	8.4	14.7
1999[4]	7.7	*	*	*	*	*	*	*	*	0.7	1.6	4.1	8.0	14.6
Female														
2001	2.3	*	*	*	*	*	*	*	*	*	0.3	0.6	1.3	1.9
2000[4]	2.3	*	*	*	*	*	*	*	*	*	*	0.4	1.2	2.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of esophagus (C15)						
Hispanic						
Both sexes						
2001	7.8	5.9	11.5	13.9	25.8	20.0
2000[4]	6.5	9.3	13.4	19.0	20.1	19.9
1999[4]	6.8	9.7	8.5	17.4	19.4	26.7
Male						
2001	13.3	11.7	21.5	22.5	45.4	41.9
2000[4]	10.9	16.4	23.3	35.4	35.3	40.3
1999[4]	13.1	16.2	16.3	32.8	33.0	49.6
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	13.3	19.3	23.5	26.7	26.5	26.0
2000[4]	13.7	18.5	22.5	24.7	27.4	26.1
1999[4]	13.4	18.7	22.7	24.5	24.8	26.8
Male						
2001	23.2	33.8	40.8	47.6	48.9	48.4
2000[4]	23.8	31.5	38.6	44.2	49.3	47.8
1999[4]	22.4	32.6	40.4	44.4	45.3	48.0
Female						
2001	4.3	6.8	9.7	12.1	13.1	16.7
2000[4]	4.5	7.3	9.8	11.1	14.5	17.3
1999[4]	5.3	6.8	8.9	10.7	12.8	18.3
Non-Hispanic White						
Both sexes						
2001	12.6	18.5	22.7	26.8	26.8	26.5
2000[4]	12.9	17.9	22.2	24.2	27.4	26.7
1999[4]	12.5	17.9	22.2	23.9	25.0	27.2
Male						
2001	22.1	32.5	39.8	48.1	49.5	49.0
2000[4]	22.8	30.8	38.2	43.4	50.1	49.4
1999[4]	21.0	31.7	39.5	43.4	46.1	48.0
Female						
2001	3.7	6.2	8.8	11.8	13.2	17.1
2000[4]	3.7	6.6	9.3	10.7	13.8	17.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of esophagus (C15)														
Non-Hispanic White														
Female														
1999[4]	2.2	*	*	*	*	*	*	*	*	*	*	0.5	1.1	2.3
Non-Hispanic Black														
Both sexes														
2001	4.7	*	*	*	*	*	*	*	*	*	1.6	3.9	8.8	16.3
2000[4]	4.7	*	*	*	*	*	*	*	*	*	1.5	4.2	10.0	17.4
1999[4]	4.7	*	*	*	*	*	*	*	*	*	1.8	4.5	9.4	17.4
Male														
2001	7.2	*	*	*	*	*	*	*	*	*	2.6	6.2	15.7	28.3
2000[4]	7.0	*	*	*	*	*	*	*	*	*	2.4	6.5	17.8	31.3
1999[4]	7.2	*	*	*	*	*	*	*	*	*	3.2	7.7	16.5	30.8
Female														
2001	2.4	*	*	*	*	*	*	*	*	*	*	1.8	2.9	6.5
2000[4]	2.6	*	*	*	*	*	*	*	*	*	*	2.2	3.4	6.0
1999[4]	2.4	*	*	*	*	*	*	*	*	*	*	1.7	3.4	6.4
Malignant neoplasm of stomach (C16)														
All origins[3]														
Both sexes														
2001	4.3	*	*	*	*	*	*	0.2	0.4	0.8	1.3	2.2	3.7	5.3
2000[4]	4.5	*	*	*	*	*	*	0.2	0.4	0.7	1.4	2.3	3.6	5.6
1999[4]	4.6	*	*	*	*	*	*	0.2	0.4	0.8	1.4	2.3	3.8	6.2
Male														
2001	5.1	*	*	*	*	*	*	0.2	0.5	0.9	1.4	2.8	4.9	7.5
2000[4]	5.3	*	*	*	*	*	*	0.2	0.5	0.9	1.7	2.9	4.8	7.8
1999[4]	5.5	*	*	*	*	*	*	0.2	0.5	0.9	1.7	3.1	5.4	8.8
Female														
2001	3.5	*	*	*	*	*	*	*	0.4	0.7	1.1	1.6	2.5	3.2
2000[4]	3.7	*	*	*	*	*	*	*	0.4	0.6	1.2	1.7	2.4	3.5
1999[4]	3.7	*	*	*	*	*	*	0.3	0.3	0.8	1.2	1.5	2.4	3.7
Hispanic														
Both sexes														
2001	3.2	*	*	*	*	*	*	*	0.8	1.3	2.3	4.1	7.0	8.8
2000[4]	3.1	*	*	*	*	*	*	*	*	1.7	2.5	4.1	5.7	7.1
1999[4]	3.1	*	*	*	*	*	*	*	0.8	1.2	2.3	3.0	5.5	9.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of esophagus (C15)						
Non-Hispanic White						
Female						
1999[4]	4.6	5.9	8.3	10.3	12.6	18.9
Non-Hispanic Black						
Both sexes						
2001	22.2	30.1	36.1	31.2	28.1	24.0
2000[4]	23.7	27.5	29.9	36.4	30.5	22.6
1999[4]	23.9	29.1	31.6	36.0	24.1	23.6
Male						
2001	39.1	52.6	60.9	55.9	54.8	47.4
2000[4]	39.1	45.6	50.2	66.0	46.3	38.7
1999[4]	40.9	46.8	56.8	68.2	37.2	49.9
Female						
2001	8.9	13.5	19.5	16.4	14.6	15.4
2000[4]	11.6	14.2	16.3	18.6	22.5	16.6
1999[4]	10.6	15.9	14.9	16.6	17.6	13.8
Malignant neoplasm of stomach (C16)						
All origins[3]						
Both sexes						
2001	9.2	13.7	19.5	27.0	34.7	45.4
2000[4]	9.7	13.8	19.6	27.1	36.9	52.7
1999[4]	9.8	14.2	20.2	28.1	36.8	50.1
Male						
2001	13.7	19.5	27.5	39.1	49.4	62.6
2000[4]	14.1	19.5	28.3	39.4	51.4	75.6
1999[4]	13.7	20.3	28.6	41.5	54.9	71.3
Female						
2001	5.1	8.8	13.1	18.5	25.9	38.3
2000[4]	5.8	8.9	12.7	18.5	28.3	43.4
1999[4]	6.4	9.1	13.7	18.8	26.2	41.6
Hispanic						
Both sexes						
2001	14.6	21.4	32.7	40.2	55.1	63.8
2000[4]	13.2	23.2	35.0	41.0	66.8	65.7
1999[4]	15.0	23.2	29.9	44.2	59.9	73.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of stomach (C16)														
Hispanic														
Male														
2001	3.6	*	*	*	*	*	*	*	*	1.5	2.4	4.5	8.5	10.2
2000[4]	3.6	*	*	*	*	*	*	*	*	1.8	2.2	3.9	6.5	9.4
1999[4]	3.5	*	*	*	*	*	*	*	*	*	2.4	4.2	6.6	12.7
Female														
2001	2.8	*	*	*	*	*	*	*	*	*	2.2	3.7	5.7	7.6
2000[4]	2.7	*	*	*	*	*	*	*	*	1.6	2.8	4.3	5.0	5.0
1999[4]	2.7	*	*	*	*	*	*	*	*	1.5	2.2	*	4.6	5.8
Non-Hispanic[5]														
Both sexes														
2001	4.5	*	*	*	*	*	*	*	0.4	0.7	1.1	2.0	3.4	5.0
2000[4]	4.7	*	*	*	*	*	*	0.2	0.4	0.6	1.3	2.2	3.4	5.4
1999[4]	4.7	*	*	*	*	*	*	0.2	0.4	0.8	1.3	2.2	3.7	5.9
Male														
2001	5.4	*	*	*	*	*	*	*	0.5	0.8	1.2	2.6	4.6	7.2
2000[4]	5.5	*	*	*	*	*	*	*	0.4	0.8	1.6	2.8	4.6	7.6
1999[4]	5.7	*	*	*	*	*	*	*	0.4	0.9	1.6	3.0	5.2	8.5
Female														
2001	3.6	*	*	*	*	*	*	*	0.3	0.6	1.0	1.3	2.2	2.9
2000[4]	3.8	*	*	*	*	*	*	*	0.4	0.4	1.0	1.5	2.2	3.4
1999[4]	3.8	*	*	*	*	*	*	*	0.3	0.7	1.1	1.4	2.2	3.5
Non-Hispanic White														
Both sexes														
2001	4.2	*	*	*	*	*	*	*	0.3	0.5	0.8	1.6	2.6	4.3
2000[4]	4.4	*	*	*	*	*	*	*	0.3	0.4	0.9	1.6	2.8	4.6
1999[4]	4.4	*	*	*	*	*	*	*	0.3	0.6	0.9	1.8	2.9	4.8
Male														
2001	5.0	*	*	*	*	*	*	*	0.3	0.6	1.0	2.1	3.6	6.1
2000[4]	5.2	*	*	*	*	*	*	*	0.3	0.5	1.1	2.1	4.0	6.5
1999[4]	5.4	*	*	*	*	*	*	*	0.3	0.7	1.0	2.5	4.1	7.1
Female														
2001	3.4	*	*	*	*	*	*	*	*	0.4	0.7	1.0	1.7	2.4
2000[4]	3.5	*	*	*	*	*	*	*	0.3	0.3	0.7	1.1	1.7	2.8
1999[4]	3.5	*	*	*	*	*	*	*	*	0.5	0.7	1.1	1.6	2.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of stomach (C16)						
Hispanic						
Male						
2001	21.5	30.5	45.7	51.8	76.6	95.2
2000[4]	19.3	34.7	54.5	53.9	104.5	98.8
1999[4]	21.1	33.2	38.7	67.9	84.9	75.5
Female						
2001	8.7	14.1	22.7	32.0	41.8	48.4
2000[4]	7.9	13.9	20.3	31.9	43.9	49.5
1999[4]	9.8	15.0	23.4	27.5	45.0	72.8
Non-Hispanic[5]						
Both sexes						
2001	8.7	13.2	18.6	26.2	33.9	44.7
2000[4]	9.4	13.1	18.7	26.4	35.6	52.0
1999[4]	9.4	13.6	19.5	27.4	35.9	49.2
Male						
2001	13.1	18.7	26.3	38.3	48.2	61.0
2000[4]	13.6	18.4	26.8	38.6	49.3	74.5
1999[4]	13.1	19.4	27.8	40.3	53.7	71.0
Female						
2001	4.8	8.4	12.5	17.8	25.3	37.9
2000[4]	5.6	8.6	12.3	17.8	27.6	42.9
1999[4]	6.1	8.6	13.1	18.4	25.5	40.5
Non-Hispanic White						
Both sexes						
2001	7.3	10.8	16.3	23.0	29.8	40.0
2000[4]	8.1	11.0	16.0	22.6	31.2	46.3
1999[4]	7.7	12.0	16.8	23.8	31.8	43.5
Male						
2001	10.9	15.1	23.0	34.2	42.4	55.7
2000[4]	11.9	15.7	23.2	33.5	43.8	67.3
1999[4]	10.7	17.3	24.3	35.2	47.6	62.6
Female						
2001	3.9	7.1	10.9	15.0	22.2	33.5
2000[4]	4.5	6.9	10.2	15.0	23.8	37.9
1999[4]	4.9	7.4	10.9	15.7	22.6	35.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of stomach (C16)														
Non-Hispanic Black														
Both sexes														
2001	5.8	*	*	*	*	*	*	*	0.8	1.7	2.6	3.8	7.5	9.3
2000[4]	6.1	*	*	*	*	*	*	*	1.0	1.3	2.9	4.8	7.2	11.2
1999[4]	6.3	*	*	*	*	*	*	*	0.8	1.4	3.3	4.0	8.6	13.3
Male														
2001	6.9	*	*	*	*	*	*	*	*	1.9	2.9	5.2	11.6	13.8
2000[4]	7.2	*	*	*	*	*	*	*	*	2.0	4.0	6.7	9.9	16.6
1999[4]	7.6	*	*	*	*	*	*	*	*	1.5	4.2	5.8	12.4	19.0
Female														
2001	4.7	*	*	*	*	*	*	*	*	1.6	2.3	2.6	4.1	5.5
2000[4]	5.2	*	*	*	*	*	*	*	*	*	2.0	3.1	4.8	6.8
1999[4]	5.1	*	*	*	*	*	*	*	*	1.3	2.5	2.4	5.4	8.7
Malignant neoplasms of colon, rectum and anus (C18-C21)														
All origins[3]														
Both sexes														
2001	20.0	*	*	*	*	*	0.2	0.4	1.1	2.3	4.1	8.4	15.2	25.4
2000[4]	20.4	*	*	*	*	*	0.1	0.4	0.9	2.1	4.0	8.2	14.7	25.2
1999[4]	20.5	*	*	*	*	*	0.2	0.4	1.0	1.9	4.0	8.3	14.4	25.3
Male														
2001	20.2	*	*	*	*	*	*	0.4	1.1	2.6	4.3	9.5	17.1	30.8
2000[4]	20.6	*	*	*	*	*	*	0.4	1.1	2.4	4.4	9.0	17.0	30.6
1999[4]	20.6	*	*	*	*	*	0.2	0.6	1.0	2.0	4.1	9.1	16.6	30.6
Female														
2001	19.8	*	*	*	*	*	*	0.5	1.0	1.9	3.8	7.4	13.4	20.4
2000[4]	20.2	*	*	*	*	*	*	0.4	0.8	1.8	3.5	7.5	12.4	20.2
1999[4]	20.3	*	*	*	*	*	*	0.3	0.9	1.9	3.9	7.5	12.3	20.4
Hispanic														
Both sexes														
2001	6.1	*	*	*	*	*	*	*	*	1.5	3.4	4.6	10.6	16.7
2000[4]	6.0	*	*	*	*	*	*	*	0.8	1.7	3.1	6.3	11.2	19.2
1999[4]	6.0	*	*	*	*	*	*	*	0.8	1.2	2.4	6.7	10.3	14.5
Male														
2001	6.2	*	*	*	*	*	*	*	*	1.8	3.8	4.6	11.1	19.8
2000[4]	6.2	*	*	*	*	*	*	*	*	1.4	2.7	6.0	13.1	22.4
1999[4]	6.1	*	*	*	*	*	*	*	*	*	2.1	7.0	11.8	16.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of stomach (C16)						
Non-Hispanic Black						
Both sexes						
2001	18.0	27.8	35.0	51.7	70.6	89.3
2000[4]	17.6	27.0	38.7	57.3	75.6	102.7
1999[4]	19.4	24.0	40.7	59.2	74.6	103.1
Male						
2001	28.0	44.8	55.1	76.8	107.1	106.3
2000[4]	26.4	38.6	57.7	83.3	104.7	138.6
1999[4]	28.7	35.6	60.6	90.9	121.3	149.8
Female						
2001	10.2	15.2	21.6	36.8	52.1	83.0
2000[4]	10.7	18.5	26.1	41.5	60.9	89.4
1999[4]	12.1	15.4	27.6	40.2	51.4	85.7
Malignant neoplasms of colon, rectum and anus (C18-C21)						
All origins[3]						
Both sexes						
2001	39.3	62.9	88.8	121.1	165.8	255.7
2000[4]	42.0	62.0	92.4	126.9	174.8	270.4
1999[4]	43.6	63.9	91.9	129.0	173.2	271.7
Male						
2001	48.0	78.9	110.4	149.4	201.3	298.3
2000[4]	51.0	77.4	115.7	159.4	213.0	315.6
1999[4]	52.9	78.0	114.3	161.3	216.2	324.0
Female						
2001	31.5	49.2	71.5	101.2	144.6	237.8
2000[4]	33.9	48.8	74.1	104.3	152.2	251.9
1999[4]	35.1	51.8	74.3	106.6	148.0	250.6
Hispanic						
Both sexes						
2001	30.1	40.4	66.7	89.8	108.3	193.8
2000[4]	28.1	40.4	60.3	85.4	121.4	181.1
1999[4]	29.2	46.8	67.8	88.4	123.9	176.8
Male						
2001	39.7	53.2	83.7	113.9	122.1	257.1
2000[4]	36.0	54.8	74.9	109.3	175.2	235.8
1999[4]	33.7	58.7	91.1	114.7	172.9	211.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of colon, rectum and anus (C18-C21)														
Hispanic														
Female														
2001	5.9	*	*	*	*	*	*	*	*	*	2.9	4.6	10.2	13.9
2000[4]	5.8	*	*	*	*	*	*	*	*	2.0	3.5	6.5	9.3	16.3
1999[4]	5.9	*	*	*	*	*	*	*	*	*	2.7	6.4	8.8	12.5
Non-Hispanic[5]														
Both sexes														
2001	22.0	*	*	*	*	*	0.2	0.5	1.2	2.4	4.2	8.8	15.6	26.0
2000[4]	22.4	*	*	*	*	*	0.1	0.4	1.0	2.2	4.0	8.4	14.9	25.6
1999[4]	22.4	*	*	*	*	*	0.2	0.5	1.0	2.0	4.2	8.4	14.7	26.0
Male														
2001	22.3	*	*	*	*	*	*	0.4	1.3	2.7	4.4	9.9	17.6	31.5
2000[4]	22.8	*	*	*	*	*	*	0.5	1.1	2.5	4.6	9.3	17.3	31.1
1999[4]	22.7	*	*	*	*	*	*	0.7	1.1	2.1	4.3	9.2	16.9	31.5
Female														
2001	21.7	*	*	*	*	*	*	0.5	1.0	2.0	3.9	7.6	13.6	20.9
2000[4]	22.1	*	*	*	*	*	*	0.4	0.8	1.8	3.5	7.5	12.6	20.4
1999[4]	22.2	*	*	*	*	*	*	0.3	0.9	1.9	4.1	7.5	12.5	20.9
Non-Hispanic White														
Both sexes														
2001	23.5	*	*	*	*	*	0.2	0.5	1.1	2.2	3.8	8.0	14.5	24.4
2000[4]	24.0	*	*	*	*	*	*	0.4	0.9	1.9	3.6	7.5	13.9	23.9
1999[4]	24.0	*	*	*	*	*	0.2	0.4	0.9	1.9	3.9	7.8	13.7	24.5
Male														
2001	23.9	*	*	*	*	*	*	0.4	1.3	2.6	4.0	9.1	16.8	29.9
2000[4]	24.2	*	*	*	*	*	*	0.4	1.1	2.3	4.1	8.5	16.3	29.0
1999[4]	24.3	*	*	*	*	*	*	0.6	1.0	2.1	4.1	8.6	16.0	29.6
Female														
2001	23.1	*	*	*	*	*	*	0.5	0.9	1.9	3.7	6.9	12.3	19.2
2000[4]	23.8	*	*	*	*	*	*	0.4	0.6	1.5	3.2	6.6	11.7	19.1
1999[4]	23.7	*	*	*	*	*	*	*	0.9	1.7	3.6	6.9	11.4	19.6
Non-Hispanic Black														
Both sexes														
2001	19.2	*	*	*	*	*	*	*	1.6	3.4	6.6	14.8	25.2	44.0
2000[4]	19.4	*	*	*	*	*	*	*	1.6	3.8	6.7	15.6	25.4	40.7
1999[4]	19.3	*	*	*	*	*	*	0.8	1.3	3.2	6.7	13.6	23.9	42.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of colon, rectum and anus (C18-C21)						
Hispanic						
Female						
2001	21.9	30.1	53.7	72.6	99.7	162.9
2000[4]	21.3	28.7	49.3	68.5	88.7	154.3
1999[4]	25.4	37.2	50.2	69.9	94.7	160.1
Non-Hispanic[5]						
Both sexes						
2001	39.9	64.4	89.9	122.4	167.9	257.4
2000[4]	43.0	63.3	94.1	128.6	176.5	273.1
1999[4]	44.5	64.8	93.0	130.5	174.2	274.3
Male						
2001	48.4	80.5	111.7	151.0	204.3	299.6
2000[4]	52.0	78.7	117.9	161.4	214.0	318.6
1999[4]	54.2	79.0	115.4	163.0	216.7	327.5
Female						
2001	32.2	50.5	72.6	102.4	146.1	239.9
2000[4]	34.8	50.1	75.4	105.7	154.4	254.7
1999[4]	35.7	52.7	75.4	108.0	149.4	253.0
Non-Hispanic White						
Both sexes						
2001	38.4	62.2	87.8	119.2	165.7	256.9
2000[4]	41.4	61.5	92.3	126.6	173.7	271.9
1999[4]	42.8	62.9	91.0	128.3	171.9	273.4
Male						
2001	47.0	78.0	109.8	148.2	201.9	298.2
2000[4]	49.9	76.3	115.6	159.5	209.2	315.8
1999[4]	52.3	77.1	114.2	159.7	214.9	328.2
Female						
2001	30.4	48.2	69.9	98.6	143.8	239.8
2000[4]	33.5	48.4	73.5	103.5	152.6	254.1
1999[4]	34.1	50.4	72.5	106.3	146.5	251.5
Non-Hispanic Black						
Both sexes						
2001	59.0	92.8	122.1	174.5	213.3	289.0
2000[4]	63.9	88.5	127.9	169.6	228.8	312.9
1999[4]	67.1	91.4	126.5	174.1	227.5	308.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of colon, rectum and anus (C18-C21)														
Non-Hispanic Black														
Male														
2001	19.0	*	*	*	*	*	*	*	1.9	4.0	7.6	16.8	27.0	52.3
2000[4]	19.8	*	*	*	*	*	*	*	1.8	4.2	8.2	17.0	29.9	52.3
1999[4]	19.2	*	*	*	*	*	*	*	*	3.1	6.5	15.4	27.4	54.9
Female														
2001	19.4	*	*	*	*	*	*	*	*	3.0	5.7	13.1	23.7	37.2
2000[4]	19.1	*	*	*	*	*	*	*	1.4	3.5	5.3	14.4	21.6	31.2
1999[4]	19.5	*	*	*	*	*	*	*	*	3.2	6.8	12.0	20.9	33.0
Malignant neoplasms of liver and intrahepatic bile ducts (C22)														
All origins[3]														
Both sexes														
2001	4.7	*	*	*	*	*	*	0.2	0.2	0.6	1.2	3.8	5.8	7.5
2000[4]	4.6	*	0.1	*	*	*	*	0.1	0.3	0.5	1.4	3.7	5.4	7.7
1999[4]	4.4	*	0.2	*	*	*	0.1	0.2	0.2	0.6	1.4	3.4	4.9	7.1
Male														
2001	6.1	*	*	*	*	*	*	*	0.3	0.9	1.8	6.0	9.7	11.4
2000[4]	5.9	*	*	*	*	*	*	*	0.4	0.7	2.0	6.0	8.5	11.7
1999[4]	5.6	*	0.3	*	*	*	*	0.2	0.3	0.8	2.1	5.5	7.6	10.6
Female														
2001	3.4	*	*	*	*	*	*	*	*	0.3	0.5	1.7	2.1	3.9
2000[4]	3.3	*	*	*	*	*	*	*	*	0.3	0.8	1.4	2.4	4.0
1999[4]	3.3	*	*	*	*	*	*	*	*	0.3	0.8	1.3	2.3	3.8
Hispanic														
Both sexes														
2001	3.4	*	*	*	*	*	*	*	*	*	1.1	5.6	8.1	12.1
2000[4]	3.3	*	*	*	*	*	*	*	*	*	2.0	5.2	6.2	11.6
1999[4]	3.2	*	*	*	*	*	*	*	*	*	1.4	4.5	7.3	9.8
Male														
2001	4.3	*	*	*	*	*	*	*	*	*	1.9	10.0	12.6	19.2
2000[4]	4.2	*	*	*	*	*	*	*	*	*	3.0	8.1	10.4	18.0
1999[4]	3.9	*	*	*	*	*	*	*	*	*	1.9	6.9	13.3	16.0
Female														
2001	2.5	*	*	*	*	*	*	*	*	*	*	*	3.7	5.6
2000[4]	2.4	*	*	*	*	*	*	*	*	*	*	2.3	*	5.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of colon, rectum and anus (C18-C21)						
Non-Hispanic Black						
Male						
2001	68.3	117.0	148.7	209.5	264.9	352.3
2000[4]	78.0	111.1	162.6	211.9	302.1	381.6
1999[4]	82.1	111.6	148.9	229.6	275.4	351.8
Female						
2001	51.6	75.0	104.4	153.6	187.0	265.6
2000[4]	52.7	71.8	104.8	144.0	191.8	287.4
1999[4]	55.3	76.4	111.7	140.7	203.7	292.7
Malignant neoplasms of liver and intrahepatic bile ducts (C22)						
All origins[3]						
Both sexes						
2001	11.6	16.3	21.5	27.0	32.0	34.1
2000[4]	11.3	16.0	21.6	27.1	31.3	32.7
1999[4]	10.9	16.3	21.8	26.1	30.9	30.6
Male						
2001	17.2	23.5	30.6	39.2	43.9	50.3
2000[4]	16.5	23.6	30.9	39.3	46.7	44.7
1999[4]	15.6	23.7	31.9	36.6	42.9	40.1
Female						
2001	6.5	10.2	14.2	18.4	24.8	27.2
2000[4]	6.5	9.5	14.2	18.6	22.2	27.8
1999[4]	6.6	10.1	13.8	18.7	23.9	26.8
Hispanic						
Both sexes						
2001	16.9	24.9	33.7	50.7	54.6	52.5
2000[4]	16.4	24.2	39.8	45.9	62.4	58.4
1999[4]	17.2	29.2	33.9	43.6	61.1	53.3
Male						
2001	24.7	35.4	48.5	68.2	67.5	64.8
2000[4]	23.9	35.4	54.9	65.0	94.2	72.6
1999[4]	26.8	36.3	51.4	56.2	77.0	66.9
Female						
2001	10.1	16.4	22.3	38.2	46.6	46.5
2000[4]	9.9	15.1	28.4	32.4	43.0	51.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of liver and intrahepatic bile ducts (C22)														
Hispanic														
Female														
1999[4]	2.4	*	*	*	*	*	*	*	*	*	*	*	*	4.2
Non-Hispanic[5]														
Both sexes														
2001	4.9	*	*	*	*	*	*	0.2	0.2	0.6	1.2	3.6	5.6	7.1
2000[4]	4.8	*	0.2	*	*	*	*	0.2	0.3	0.5	1.3	3.5	5.3	7.4
1999[4]	4.6	*	0.2	*	*	*	0.1	0.2	0.2	0.6	1.4	3.3	4.7	6.8
Male														
2001	6.3	*	*	*	*	*	*	*	0.3	0.9	1.8	5.6	9.4	10.6
2000[4]	6.2	*	*	*	*	*	*	*	0.4	0.6	1.9	5.8	8.3	11.2
1999[4]	5.8	*	*	*	*	*	*	*	0.3	0.8	2.1	5.4	7.1	10.2
Female														
2001	3.5	*	*	*	*	*	*	*	*	0.3	0.6	1.7	2.0	3.7
2000[4]	3.4	*	*	*	*	*	*	*	*	0.3	0.8	1.4	2.4	3.9
1999[4]	3.4	*	*	*	*	*	*	*	*	0.3	0.8	1.2	2.3	3.7
Non-Hispanic White														
Both sexes														
2001	4.8	*	*	*	*	*	*	*	0.1	0.4	0.8	2.9	4.6	5.7
2000[4]	4.6	*	*	*	*	*	*	*	0.2	0.3	0.9	2.8	4.2	6.1
1999[4]	4.5	*	0.2	*	*	*	*	*	*	0.4	1.1	2.5	3.9	5.5
Male														
2001	6.1	*	*	*	*	*	*	*	*	0.6	1.1	4.3	7.5	8.4
2000[4]	5.9	*	*	*	*	*	*	*	*	0.3	1.3	4.4	6.4	8.9
1999[4]	5.7	*	*	*	*	*	*	*	*	0.5	1.7	4.0	5.6	8.1
Female														
2001	3.4	*	*	*	*	*	*	*	*	*	0.4	1.4	1.7	3.2
2000[4]	3.4	*	*	*	*	*	*	*	*	0.3	0.5	1.2	2.0	3.4
1999[4]	3.4	*	*	*	*	*	*	*	*	0.3	0.5	1.0	2.1	3.0
Non-Hispanic Black														
Both sexes														
2001	4.6	*	*	*	*	*	*	*	*	1.0	2.4	7.3	10.9	15.6
2000[4]	4.5	*	*	*	*	*	*	*	*	0.8	2.8	7.1	10.9	13.0
1999[4]	4.3	*	*	*	*	*	*	*	*	1.1	2.2	6.7	8.3	12.2
Male														
2001	6.1	*	*	*	*	*	*	*	*	1.7	3.7	11.8	19.1	26.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)						
Hispanic						
Female						
1999[4]	9.0	23.4	20.7	34.7	51.6	46.8
Non-Hispanic[5]						
Both sexes						
2001	11.1	15.7	20.6	25.7	31.0	33.2
2000[4]	10.8	15.4	20.5	26.2	30.0	31.7
1999[4]	10.4	15.5	21.1	25.3	29.8	29.7
Male						
2001	16.6	22.6	29.4	37.6	42.9	49.7
2000[4]	15.9	22.7	29.5	38.0	44.6	43.4
1999[4]	14.8	22.8	30.7	35.7	41.6	38.7
Female						
2001	6.1	9.7	13.7	17.4	23.9	26.4
2000[4]	6.2	9.1	13.4	18.0	21.4	27.0
1999[4]	6.4	9.1	13.5	18.0	22.9	26.1
Non-Hispanic White						
Both sexes						
2001	9.9	14.2	19.2	24.6	29.2	32.4
2000[4]	9.4	13.8	18.8	24.7	29.2	30.8
1999[4]	9.0	14.1	19.6	24.2	28.8	28.9
Male						
2001	14.9	21.0	27.6	35.9	41.3	48.9
2000[4]	13.6	20.3	27.1	36.5	43.5	41.6
1999[4]	12.8	20.8	28.8	34.1	40.8	39.0
Female						
2001	5.2	8.2	12.3	16.6	22.0	25.5
2000[4]	5.5	8.0	12.1	16.3	20.8	26.5
1999[4]	5.6	8.1	12.3	17.2	21.7	24.8
Non-Hispanic Black						
Both sexes						
2001	14.5	20.1	24.1	26.2	41.7	35.3
2000[4]	16.2	20.7	27.3	29.2	29.0	31.8
1999[4]	14.9	21.4	28.4	26.9	34.3	34.9
Male						
2001	21.7	28.1	33.1	38.4	49.9	42.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of liver and intrahepatic bile ducts (C22)														
Non-Hispanic Black														
Male														
2000[4]	6.3	*	*	*	*	*	*	*	*	*	3.9	12.6	18.1	22.7
1999[4]	5.7	*	*	*	*	*	*	*	*	1.6	3.5	12.1	14.5	19.0
Female														
2001	3.3	*	*	*	*	*	*	*	*	*	*	3.3	3.9	7.0
2000[4]	2.9	*	*	*	*	*	*	*	*	*	1.9	2.3	4.8	5.1
1999[4]	3.1	*	*	*	*	*	*	*	*	*	*	2.1	3.0	6.7
Malignant neoplasm of pancreas (C25)														
All origins[3]														
Both sexes														
2001	10.5	*	*	*	*	*	*	*	0.2	0.8	1.8	4.0	8.6	14.9
2000[4]	10.4	*	*	*	*	*	*	*	0.2	0.6	1.8	4.2	8.7	15.1
1999[4]	10.4	*	*	*	*	*	*	*	0.2	0.8	1.9	4.5	8.7	15.6
Male														
2001	10.3	*	*	*	*	*	*	*	0.2	0.9	2.0	4.9	10.7	18.5
2000[4]	10.3	*	*	*	*	*	*	*	0.2	0.8	2.1	5.4	11.1	18.5
1999[4]	10.4	*	*	*	*	*	*	*	0.3	0.9	2.4	5.9	11.0	19.1
Female														
2001	10.6	*	*	*	*	*	*	*	*	0.6	1.6	3.1	6.5	11.5
2000[4]	10.5	*	*	*	*	*	*	*	*	0.5	1.5	3.0	6.4	11.8
1999[4]	10.5	*	*	*	*	*	*	*	0.2	0.6	1.4	3.1	6.5	12.3
Hispanic														
Both sexes														
2001	3.6	*	*	*	*	*	*	*	*	*	1.2	2.4	5.4	11.6
2000[4]	3.5	*	*	*	*	*	*	*	*	*	1.0	3.0	5.5	9.6
1999[4]	3.3	*	*	*	*	*	*	*	*	*	1.2	3.7	5.8	10.4
Male														
2001	3.4	*	*	*	*	*	*	*	*	*	*	2.9	6.2	13.6
2000[4]	3.4	*	*	*	*	*	*	*	*	*	*	4.2	6.5	12.7
1999[4]	3.2	*	*	*	*	*	*	*	*	*	*	5.1	6.6	13.4
Female														
2001	3.9	*	*	*	*	*	*	*	*	*	*	*	4.6	9.8
2000[4]	3.6	*	*	*	*	*	*	*	*	*	*	*	4.6	6.7
1999[4]	3.5	*	*	*	*	*	*	*	*	*	*	2.4	5.0	7.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)						
Non-Hispanic Black						
Male						
2000[4]	25.9	33.6	42.0	37.1	46.3	48.1
1999[4]	22.0	32.4	42.6	40.6	43.4	30.9
Female						
2001	8.9	14.2	18.1	19.0	37.5	32.5
2000[4]	8.6	11.2	17.5	24.4	20.3	25.7
1999[4]	9.4	13.2	19.1	18.6	29.9	36.4
Malignant neoplasm of pancreas (C25)						
All origins[3]						
Both sexes						
2001	24.8	37.0	52.8	67.2	84.2	100.1
2000[4]	24.8	37.2	51.4	70.4	85.7	96.4
1999[4]	25.0	38.1	52.5	67.2	82.0	99.7
Male						
2001	30.3	43.6	59.0	77.2	94.9	105.1
2000[4]	30.0	44.7	57.0	80.4	94.9	102.9
1999[4]	29.9	44.9	60.9	75.7	92.0	107.5
Female						
2001	19.8	31.3	47.8	60.1	77.8	98.1
2000[4]	20.0	30.8	47.1	63.5	80.2	93.8
1999[4]	20.5	32.2	46.0	61.2	76.1	96.6
Hispanic						
Both sexes						
2001	17.8	29.2	46.5	55.5	71.5	83.8
2000[4]	18.9	29.4	46.3	56.6	71.3	82.9
1999[4]	17.0	26.8	40.0	55.5	58.7	84.2
Male						
2001	19.8	36.1	50.3	64.1	77.9	62.9
2000[4]	21.6	39.5	52.5	57.6	73.6	80.6
1999[4]	17.6	32.8	49.4	53.1	62.9	86.3
Female						
2001	16.0	23.5	43.6	49.4	67.6	94.0
2000[4]	16.6	21.1	41.6	55.9	69.9	84.1
1999[4]	16.4	21.9	32.9	57.3	56.3	83.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of pancreas (C25)														
Non-Hispanic[5]														
Both sexes														
2001	11.5	*	*	*	*	*	*	*	0.2	0.8	1.8	4.2	8.8	15.1
2000[4]	11.4	*	*	*	*	*	*	*	0.2	0.6	1.9	4.2	8.9	15.5
1999[4]	11.4	*	*	*	*	*	*	*	0.2	0.8	1.9	4.6	8.9	16.0
Male														
2001	11.4	*	*	*	*	*	*	*	0.3	0.9	2.0	5.1	11.1	18.9
2000[4]	11.3	*	*	*	*	*	*	*	0.2	0.8	2.2	5.5	11.4	18.9
1999[4]	11.4	*	*	*	*	*	*	*	0.3	1.0	2.5	6.0	11.3	19.4
Female														
2001	11.5	*	*	*	*	*	*	*	*	0.6	1.7	3.2	6.6	11.6
2000[4]	11.4	*	*	*	*	*	*	*	*	0.5	1.6	3.1	6.5	12.2
1999[4]	11.4	*	*	*	*	*	*	*	*	0.6	1.4	3.2	6.6	12.7
Non-Hispanic White														
Both sexes														
2001	12.3	*	*	*	*	*	*	*	0.2	0.8	1.8	3.9	8.6	14.6
2000[4]	12.2	*	*	*	*	*	*	*	0.2	0.6	1.8	4.1	8.6	14.9
1999[4]	12.1	*	*	*	*	*	*	*	0.2	0.7	1.9	4.5	8.4	15.1
Male														
2001	12.4	*	*	*	*	*	*	*	*	0.9	2.0	4.9	10.8	18.6
2000[4]	12.2	*	*	*	*	*	*	*	*	0.8	2.1	5.3	11.0	18.2
1999[4]	12.1	*	*	*	*	*	*	*	*	0.9	2.4	5.8	10.5	18.2
Female														
2001	12.3	*	*	*	*	*	*	*	*	0.7	1.6	3.0	6.4	10.9
2000[4]	12.2	*	*	*	*	*	*	*	*	0.5	1.5	3.0	6.3	11.7
1999[4]	12.1	*	*	*	*	*	*	*	*	0.6	1.3	3.1	6.4	12.1
Non-Hispanic Black														
Both sexes														
2001	9.4	*	*	*	*	*	*	*	*	0.9	2.3	6.5	12.5	21.1
2000[4]	9.7	*	*	*	*	*	*	*	*	0.8	2.7	5.9	13.5	23.0
1999[4]	9.7	*	*	*	*	*	*	*	*	1.3	2.8	6.4	14.4	25.6
Male														
2001	8.9	*	*	*	*	*	*	*	*	*	2.4	8.0	15.6	24.6
2000[4]	9.3	*	*	*	*	*	*	*	*	*	3.1	7.4	18.0	29.5
1999[4]	9.7	*	*	*	*	*	*	*	*	*	3.5	8.6	20.0	34.8
Female														
2001	9.9	*	*	*	*	*	*	*	*	*	2.2	5.2	9.8	18.2
2000[4]	10.0	*	*	*	*	*	*	*	*	*	2.4	4.5	9.7	17.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of pancreas (C25)						
Non-Hispanic[5]						
Both sexes						
2001	25.3	37.4	53.0	67.6	84.6	100.6
2000[4]	25.2	37.7	51.6	70.9	85.9	96.6
1999[4]	25.5	38.7	53.0	67.5	82.6	100.1
Male						
2001	31.1	43.9	59.3	77.7	95.5	106.4
2000[4]	30.6	44.9	57.2	81.4	95.5	103.5
1999[4]	30.6	45.6	61.2	76.5	92.9	108.2
Female						
2001	20.0	31.8	48.0	60.5	78.0	98.2
2000[4]	20.3	31.4	47.3	63.7	80.3	93.8
1999[4]	20.8	32.8	46.5	61.2	76.5	96.8
Non-Hispanic White						
Both sexes						
2001	25.3	36.7	51.8	66.7	83.6	99.4
2000[4]	24.6	36.6	50.5	69.6	84.7	94.7
1999[4]	24.8	37.8	51.9	66.0	81.1	98.9
Male						
2001	31.0	43.1	58.3	77.6	95.1	108.0
2000[4]	30.1	44.5	56.0	80.9	94.5	103.0
1999[4]	29.9	44.6	60.4	75.3	91.7	108.1
Female						
2001	19.9	31.0	46.6	58.9	76.7	95.8
2000[4]	19.6	29.8	46.0	61.6	78.9	91.4
1999[4]	20.1	31.8	45.1	59.5	74.9	95.2
Non-Hispanic Black						
Both sexes						
2001	31.9	48.9	71.5	85.2	106.0	122.0
2000[4]	33.9	51.1	70.6	94.9	105.2	121.2
1999[4]	32.7	51.9	70.8	88.3	105.7	119.0
Male						
2001	38.9	60.2	78.8	94.8	110.4	100.5
2000[4]	41.0	56.8	80.0	102.6	117.6	110.4
1999[4]	40.5	61.1	78.6	101.5	118.7	115.3
Female						
2001	26.5	40.6	66.5	79.5	103.7	130.0
2000[4]	28.3	47.0	64.4	90.3	98.9	125.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of pancreas (C25)														
Non-Hispanic Black														
Female														
1999[4]	9.7	*	*	*	*	*	*	*	*	*	2.3	4.4	9.6	18.1
Malignant neoplasm of larynx (C32)														
All origins[3]														
Both sexes														
2001	1.3	*	*	*	*	*	*	*	*	*	0.3	0.7	1.5	2.9
2000[4]	1.4	*	*	*	*	*	*	*	*	*	0.2	0.8	1.5	2.9
1999[4]	1.4	*	*	*	*	*	*	*	*	0.1	0.3	0.8	1.5	3.1
Male														
2001	2.2	*	*	*	*	*	*	*	*	*	0.5	1.2	2.5	4.9
2000[4]	2.2	*	*	*	*	*	*	*	*	*	0.4	1.2	2.5	4.9
1999[4]	2.2	*	*	*	*	*	*	*	*	*	0.4	1.3	2.6	5.4
Female														
2001	0.5	*	*	*	*	*	*	*	*	*	*	0.2	0.5	1.0
2000[4]	0.6	*	*	*	*	*	*	*	*	*	*	0.3	0.6	0.9
1999[4]	0.6	*	*	*	*	*	*	*	*	*	*	0.2	0.4	1.0
Hispanic														
Both sexes														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.7	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.7	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.9	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	1.5	*	*	*	*	*	*	*	*	*	0.3	0.7	1.6	3.0
2000[4]	1.5	*	*	*	*	*	*	*	*	*	0.3	0.8	1.6	3.0
1999[4]	1.5	*	*	*	*	*	*	*	*	0.1	0.3	0.8	1.5	3.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of pancreas (C25)						
Non-Hispanic Black						
Female						
1999[4]	26.6	45.1	65.7	80.4	99.2	120.4
Malignant neoplasm of larynx (C32)						
All origins[3]						
Both sexes						
2001	4.5	5.6	7.3	7.4	6.9	6.7
2000[4]	4.5	6.0	7.3	8.0	8.1	6.5
1999[4]	4.6	6.1	7.4	7.2	7.4	7.0
Male						
2001	7.5	9.5	12.9	14.5	13.9	18.3
2000[4]	7.9	10.0	12.9	14.8	17.1	17.4
1999[4]	7.8	10.4	12.6	13.2	15.8	17.8
Female						
2001	1.7	2.2	2.8	2.5	2.8	1.9
2000[4]	1.4	2.6	2.8	3.3	2.8	2.1
1999[4]	1.7	2.5	3.3	3.0	2.6	2.6
Hispanic						
Both sexes						
2001	2.5	5.5	4.9	5.7	*	*
2000[4]	*	4.5	*	*	11.1	*
1999[4]	*	5.2	5.5	7.7	*	*
Male						
2001	*	10.3	11.0	*	*	*
2000[4]	*	8.6	*	*	*	*
1999[4]	*	10.0	11.2	15.6	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	4.6	5.5	7.4	7.5	7.0	6.7
2000[4]	4.7	6.1	7.4	8.2	8.0	6.5
1999[4]	4.8	6.2	7.5	7.2	7.3	6.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of larynx (C32)														
Non-Hispanic[5]														
Male														
2001	2.4	*	*	*	*	*	*	*	*	*	0.5	1.2	2.6	5.0
2000[4]	2.4	*	*	*	*	*	*	*	*	*	0.4	1.3	2.6	5.0
1999[4]	2.4	*	*	*	*	*	*	*	*	*	0.4	1.3	2.7	5.5
Female														
2001	0.6	*	*	*	*	*	*	*	*	*	*	0.3	0.5	1.1
2000[4]	0.6	*	*	*	*	*	*	*	*	*	*	0.4	0.6	1.0
1999[4]	0.6	*	*	*	*	*	*	*	*	*	*	0.2	0.5	1.1
Non-Hispanic White														
Both sexes														
2001	1.5	*	*	*	*	*	*	*	*	*	0.3	0.5	1.3	2.6
2000[4]	1.5	*	*	*	*	*	*	*	*	*	0.2	0.7	1.3	2.5
1999[4]	1.5	*	*	*	*	*	*	*	*	*	0.2	0.6	1.3	2.8
Male														
2001	2.3	*	*	*	*	*	*	*	*	*	0.5	0.9	2.1	4.3
2000[4]	2.4	*	*	*	*	*	*	*	*	*	0.3	1.1	2.1	4.1
1999[4]	2.3	*	*	*	*	*	*	*	*	*	0.2	1.0	2.3	4.7
Female														
2001	0.6	*	*	*	*	*	*	*	*	*	*	*	0.5	0.9
2000[4]	0.6	*	*	*	*	*	*	*	*	*	*	0.3	0.6	0.9
1999[4]	0.7	*	*	*	*	*	*	*	*	*	*	*	0.3	1.0
Non-Hispanic Black														
Both sexes														
2001	1.9	*	*	*	*	*	*	*	*	*	*	2.0	3.9	7.1
2000[4]	2.0	*	*	*	*	*	*	*	*	*	*	2.1	4.1	7.7
1999[4]	1.9	*	*	*	*	*	*	*	*	*	1.0	2.2	4.0	7.7
Male														
2001	3.1	*	*	*	*	*	*	*	*	*	*	3.7	7.1	12.9
2000[4]	3.4	*	*	*	*	*	*	*	*	*	*	3.3	7.7	14.7
1999[4]	3.2	*	*	*	*	*	*	*	*	*	1.7	4.0	7.1	14.8
Female														
2001	0.7	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.8	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.7	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of larynx (C32)						
Non-Hispanic[5]						
Male						
2001	7.6	9.4	13.0	14.5	14.0	18.4
2000[4]	8.1	10.0	13.1	15.1	16.7	17.3
1999[4]	8.1	10.4	12.6	13.0	15.6	17.3
Female						
2001	1.8	2.3	3.0	2.6	2.8	1.8
2000[4]	1.5	2.7	2.9	3.4	2.8	2.1
1999[4]	1.8	2.5	3.4	3.1	2.5	2.6
Non-Hispanic White						
Both sexes						
2001	4.2	5.2	6.8	7.0	6.8	6.5
2000[4]	4.0	5.7	6.8	7.6	7.8	6.5
1999[4]	4.3	5.7	7.2	7.0	7.2	6.5
Male						
2001	7.0	8.6	11.9	13.3	13.3	18.3
2000[4]	6.9	9.2	11.9	14.1	16.3	17.4
1999[4]	7.3	9.5	11.9	12.7	15.0	16.0
Female						
2001	1.6	2.1	2.7	2.6	2.9	1.6
2000[4]	1.4	2.6	2.7	3.1	2.8	2.1
1999[4]	1.6	2.4	3.4	3.1	2.6	2.7
Non-Hispanic Black						
Both sexes						
2001	8.9	9.6	13.9	13.6	9.4	8.4
2000[4]	11.0	10.7	15.5	15.9	11.8	7.0
1999[4]	10.0	11.5	11.9	10.4	9.4	8.4
Male						
2001	15.9	17.7	26.2	31.3	22.9	*
2000[4]	21.0	20.0	30.4	29.9	27.5	*
1999[4]	18.1	21.3	23.2	20.3	26.6	25.0
Female						
2001	3.5	*	5.7	*	*	*
2000[4]	*	3.9	5.7	7.6	*	*
1999[4]	3.8	4.3	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of trachea, bronchus and lung (C33-C34)														
All origins[3]														
Both sexes														
2001	54.8	*	*	*	*	*	*	0.2	0.6	2.9	9.5	20.5	42.3	88.8
2000[4]	55.3	*	*	*	*	*	*	0.2	0.8	2.9	9.3	21.0	43.8	92.5
1999[4]	54.5	*	*	*	*	*	*	0.2	0.7	2.9	9.3	19.9	44.4	93.0
Male														
2001	64.7	*	*	*	*	*	*	*	0.6	2.8	10.5	24.3	52.2	109.6
2000[4]	65.5	*	*	*	*	*	*	0.3	0.7	3.3	10.5	25.7	53.2	115.9
1999[4]	65.4	*	*	*	*	*	*	0.2	0.7	2.8	10.5	24.3	54.4	118.1
Female														
2001	45.3	*	*	*	*	*	*	*	0.6	2.9	8.4	16.8	32.8	69.2
2000[4]	45.4	*	*	*	*	*	*	*	0.8	2.6	8.1	16.5	34.7	70.5
1999[4]	44.1	*	*	*	*	*	*	*	0.6	3.0	8.1	15.7	34.9	69.5
Hispanic														
Both sexes														
2001	10.3	*	*	*	*	*	*	*	*	0.9	3.2	6.6	16.0	30.9
2000[4]	10.3	*	*	*	*	*	*	*	*	1.0	2.7	7.0	16.0	31.6
1999[4]	10.4	*	*	*	*	*	*	*	*	1.2	3.0	6.6	15.4	33.2
Male														
2001	13.1	*	*	*	*	*	*	*	*	1.4	3.6	8.6	21.6	43.8
2000[4]	13.3	*	*	*	*	*	*	*	*	1.4	3.1	9.1	22.3	46.0
1999[4]	13.6	*	*	*	*	*	*	*	*	*	3.6	8.8	21.4	50.6
Female														
2001	7.4	*	*	*	*	*	*	*	*	*	2.7	4.6	10.7	19.2
2000[4]	7.2	*	*	*	*	*	*	*	*	*	2.3	4.8	10.0	18.5
1999[4]	7.1	*	*	*	*	*	*	*	*	*	2.3	4.4	9.6	17.5
Non-Hispanic[5]														
Both sexes														
2001	61.3	*	*	*	*	*	*	0.2	0.6	3.2	10.2	21.9	44.4	93.2
2000[4]	61.5	*	*	*	*	*	*	0.2	0.8	3.2	10.0	22.3	45.9	96.8
1999[4]	60.5	*	*	*	*	*	*	0.2	0.7	3.2	9.9	21.1	46.6	97.2
Male														
2001	72.6	*	*	*	*	*	*	*	0.6	3.1	11.3	25.9	54.6	114.3
2000[4]	73.2	*	*	*	*	*	*	0.3	0.8	3.6	11.3	27.2	55.6	120.7
1999[4]	72.7	*	*	*	*	*	*	0.3	0.8	3.1	11.2	25.7	56.8	122.7
Female														
2001	50.5	*	*	*	*	*	*	*	0.7	3.3	9.0	17.9	34.6	73.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)						
All origins[3]						
Both sexes						
2001	154.5	239.7	323.2	373.4	368.6	302.7
2000[4]	159.8	245.8	325.4	368.2	374.7	302.1
1999[4]	160.7	243.6	322.3	361.7	362.6	297.0
Male						
2001	195.7	312.9	434.7	522.3	541.0	512.4
2000[4]	202.3	322.5	440.5	516.5	558.1	521.2
1999[4]	207.7	322.4	439.2	513.4	551.3	529.2
Female						
2001	117.2	176.7	234.5	269.1	265.2	215.0
2000[4]	121.3	180.2	234.7	264.9	266.6	212.8
1999[4]	118.1	176.1	230.9	256.2	252.2	203.5
Hispanic						
Both sexes						
2001	54.6	89.4	131.8	177.2	191.2	200.7
2000[4]	49.2	90.9	135.4	182.1	228.9	213.7
1999[4]	58.0	96.1	134.2	178.5	204.4	227.3
Male						
2001	80.9	130.8	206.3	281.7	316.8	335.2
2000[4]	75.1	130.9	214.9	304.1	374.0	368.8
1999[4]	90.5	152.2	205.7	294.1	360.0	362.3
Female						
2001	31.7	56.0	75.0	103.1	113.4	135.0
2000[4]	26.8	58.6	75.1	95.7	140.7	137.5
1999[4]	30.0	50.6	80.4	96.9	111.6	162.2
Non-Hispanic[5]						
Both sexes						
2001	161.7	249.8	334.1	382.4	375.1	305.8
2000[4]	167.6	255.5	335.4	376.0	379.5	304.5
1999[4]	167.7	252.4	331.5	368.7	367.5	298.8
Male						
2001	203.5	324.7	447.4	533.1	549.4	518.4
2000[4]	210.9	334.0	451.9	525.2	564.0	526.2
1999[4]	215.3	332.1	450.2	521.5	557.1	534.6
Female						
2001	123.7	185.0	243.9	276.9	270.8	217.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of trachea, bronchus and lung (C33-C34)														
Non-Hispanic[5]														
Female														
2000[4]	50.4	*	*	*	*	*	*	*	0.9	2.8	8.7	17.5	36.6	74.4
1999[4]	48.8	*	*	*	*	*	*	*	0.6	3.2	8.7	16.7	36.8	73.3
Non-Hispanic White														
Both sexes														
2001	67.3	*	*	*	*	*	*	0.2	0.7	3.2	9.8	20.7	43.2	92.7
2000[4]	67.4	*	*	*	*	*	*	0.2	0.9	3.0	9.4	20.8	44.4	96.3
1999[4]	65.9	*	*	*	*	*	*	0.2	0.7	3.0	9.5	19.6	45.0	96.5
Male														
2001	78.4	*	*	*	*	*	*	*	0.6	3.0	10.5	23.8	51.4	110.5
2000[4]	78.9	*	*	*	*	*	*	*	0.8	3.2	10.0	25.0	51.8	116.5
1999[4]	78.1	*	*	*	*	*	*	*	0.8	2.9	10.4	23.2	52.9	118.9
Female														
2001	56.6	*	*	*	*	*	*	*	0.7	3.4	9.1	17.7	35.1	75.5
2000[4]	56.4	*	*	*	*	*	*	*	0.9	2.7	8.7	16.7	37.2	77.0
1999[4]	54.3	*	*	*	*	*	*	*	0.6	3.1	8.6	16.1	37.2	75.1
Non-Hispanic Black														
Both sexes														
2001	44.6	*	*	*	*	*	*	*	*	3.6	14.9	33.6	64.6	121.4
2000[4]	45.4	*	*	*	*	*	*	*	0.9	5.0	16.1	36.4	70.5	124.8
1999[4]	45.6	*	*	*	*	*	*	*	*	4.7	14.9	35.8	71.8	127.4
Male														
2001	59.2	*	*	*	*	*	*	*	*	4.0	18.8	46.2	93.9	177.0
2000[4]	60.3	*	*	*	*	*	*	*	*	6.0	21.9	48.8	102.3	187.8
1999[4]	60.9	*	*	*	*	*	*	*	*	4.7	19.1	49.8	104.8	189.6
Female														
2001	31.3	*	*	*	*	*	*	*	*	3.3	11.4	22.6	39.8	75.8
2000[4]	31.9	*	*	*	*	*	*	*	*	4.1	10.9	25.7	43.4	73.3
1999[4]	31.8	*	*	*	*	*	*	*	*	4.7	11.2	23.7	43.9	76.8
Malignant melanoma of skin (C43)														
All origins[3]														
Both sexes														
2001	2.6	*	*	*	*	*	0.2	0.4	0.6	1.0	1.8	2.6	3.6	4.6
2000[4]	2.6	*	*	*	*	*	0.2	0.3	0.6	1.0	1.7	2.7	3.8	4.7
1999[4]	2.6	*	*	*	*	*	0.2	0.3	0.6	1.0	1.7	2.6	3.6	4.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)						
Non-Hispanic[5]						
Female						
2000[4]	128.2	188.0	243.4	272.1	270.8	214.8
1999[4]	124.5	183.9	238.6	262.5	256.5	204.4
Non-Hispanic White						
Both sexes						
2001	163.9	253.2	339.3	386.7	377.9	308.6
2000[4]	169.9	258.9	339.0	378.9	385.7	306.8
1999[4]	168.4	254.2	334.6	370.3	370.4	300.5
Male						
2001	201.8	322.2	446.0	530.5	545.9	520.0
2000[4]	209.8	331.3	448.7	520.4	565.8	527.3
1999[4]	211.0	328.0	446.4	517.2	553.9	536.9
Female						
2001	128.8	192.3	252.7	284.9	276.5	220.6
2000[4]	133.0	195.2	250.6	279.5	278.7	218.0
1999[4]	128.9	189.4	245.5	267.2	262.1	206.2
Non-Hispanic Black						
Both sexes						
2001	185.0	277.6	353.4	401.1	375.0	281.6
2000[4]	192.2	284.2	363.5	399.7	347.2	291.0
1999[4]	202.3	286.3	356.2	406.5	365.7	293.3
Male						
2001	271.9	426.0	561.5	668.4	667.2	550.9
2000[4]	276.6	438.7	568.2	664.3	623.1	567.1
1999[4]	307.4	434.8	564.3	662.7	669.5	580.0
Female						
2001	116.7	168.1	214.7	241.6	226.1	181.8
2000[4]	125.6	170.3	227.5	240.2	208.2	188.4
1999[4]	120.0	176.2	218.0	252.2	215.1	186.1
Malignant melanoma of skin (C43)						
All origins[3]						
Both sexes						
2001	6.1	7.8	10.7	12.4	14.7	17.8
2000[4]	6.3	7.9	10.0	12.0	14.9	18.9
1999[4]	6.1	8.3	9.7	12.9	14.2	16.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant melanoma of skin (C43)														
All origins[3]														
Male														
2001	3.4	*	*	*	*	*	*	0.4	0.6	1.2	2.2	3.2	4.7	6.3
2000[4]	3.3	*	*	*	*	*	*	0.3	0.7	1.0	2.1	3.2	5.2	6.5
1999[4]	3.3	*	*	*	*	*	*	0.4	0.7	1.3	2.2	3.5	4.6	6.7
Female														
2001	1.9	*	*	*	*	*	*	0.3	0.5	0.7	1.4	1.9	2.6	3.0
2000[4]	2.0	*	*	*	*	*	*	0.3	0.5	1.0	1.4	2.1	2.4	3.0
1999[4]	1.9	*	*	*	*	*	*	0.2	0.6	0.8	1.3	1.8	2.6	2.8
Hispanic														
Both sexes														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	3.0	*	*	*	*	*	0.2	0.4	0.6	1.1	1.9	2.8	3.9	4.9
2000[4]	3.0	*	*	*	*	*	0.2	0.4	0.7	1.1	1.9	2.8	4.0	4.9
1999[4]	2.9	*	*	*	*	*	0.2	0.4	0.7	1.2	1.9	2.8	3.8	4.9
Male														
2001	3.9	*	*	*	*	*	*	0.5	0.7	1.3	2.4	3.5	5.1	6.6
2000[4]	3.7	*	*	*	*	*	*	0.4	0.9	1.1	2.3	3.4	5.5	6.7
1999[4]	3.7	*	*	*	*	*	*	0.5	0.8	1.5	2.4	3.7	4.8	7.0
Female														
2001	2.1	*	*	*	*	*	*	0.4	0.6	0.8	1.5	2.1	2.8	3.2
2000[4]	2.2	*	*	*	*	*	*	0.3	0.6	1.1	1.5	2.2	2.6	3.1
1999[4]	2.1	*	*	*	*	*	*	0.3	0.7	0.8	1.4	2.0	2.7	2.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant melanoma of skin (C43)						
All origins[3]						
Male						
2001	8.8	11.9	16.3	19.6	24.0	29.9
2000[4]	8.7	11.2	14.7	18.4	24.7	31.8
1999[4]	8.4	12.0	14.6	19.8	22.7	27.5
Female						
2001	3.7	4.3	6.4	7.3	9.2	12.8
2000[4]	4.1	5.0	6.3	7.5	9.1	13.7
1999[4]	4.0	5.2	5.8	8.1	9.2	11.9
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	6.5	8.2	11.2	12.7	15.1	18.3
2000[4]	6.6	8.3	10.3	12.3	15.3	19.3
1999[4]	6.4	8.6	10.1	13.2	14.6	16.7
Male						
2001	9.3	12.5	17.0	20.2	24.8	31.0
2000[4]	9.2	11.9	15.2	19.0	25.2	32.6
1999[4]	8.8	12.4	15.2	20.4	23.3	28.2
Female						
2001	3.9	4.6	6.6	7.5	9.2	12.9
2000[4]	4.3	5.2	6.5	7.7	9.4	13.9
1999[4]	4.2	5.4	6.1	8.1	9.5	12.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant melanoma of skin (C43)														
Non-Hispanic White														
Both sexes														
2001	3.7	*	*	*	*	*	0.2	0.6	0.8	1.3	2.4	3.4	4.6	5.7
2000[4]	3.6	*	*	*	*	*	0.3	0.4	0.9	1.3	2.2	3.4	4.7	5.6
1999[4]	3.5	*	*	*	*	*	0.2	0.5	0.9	1.4	2.3	3.4	4.4	5.7
Male														
2001	4.8	*	*	*	*	*	*	0.6	0.9	1.6	2.9	4.2	6.0	7.6
2000[4]	4.6	*	*	*	*	*	*	0.5	1.1	1.3	2.7	4.1	6.4	7.7
1999[4]	4.5	*	*	*	*	*	*	0.6	1.0	1.8	3.0	4.4	5.6	8.1
Female														
2001	2.6	*	*	*	*	*	*	0.5	0.7	1.1	1.8	2.5	3.3	3.8
2000[4]	2.7	*	*	*	*	*	*	0.4	0.7	1.4	1.8	2.7	3.0	3.6
1999[4]	2.5	*	*	*	*	*	*	0.4	0.8	1.1	1.6	2.4	3.2	3.4
Non-Hispanic Black														
Both sexes														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Malignant neoplasm of breast (C50)														
All origins[3]														
Both sexes														
2001	14.7	*	*	*	*	*	*	0.5	1.8	4.3	8.2	13.4	20.6	27.1
2000[4]	15.0	*	*	*	*	*	*	0.4	1.9	4.3	8.3	13.6	20.8	28.1
1999[4]	14.9	*	*	*	*	*	*	0.4	1.7	4.3	7.9	13.3	21.0	28.1
Male														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	0.4	0.5
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	0.3	0.7
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	0.4	0.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant melanoma of skin (C43)						
Non-Hispanic White						
Both sexes						
2001	7.5	9.3	12.5	14.0	16.3	19.9
2000[4]	7.6	9.4	11.4	13.5	16.4	20.8
1999[4]	7.3	9.8	11.2	14.4	15.7	17.9
Male						
2001	10.7	14.0	18.8	22.2	27.0	34.1
2000[4]	10.6	13.3	16.8	20.8	27.1	35.7
1999[4]	10.0	14.0	16.9	22.2	25.2	30.5
Female						
2001	4.5	5.1	7.3	8.3	9.9	13.9
2000[4]	4.9	6.0	7.1	8.4	10.0	14.7
1999[4]	4.8	6.1	6.7	8.9	10.2	12.9
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	6.7
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Malignant neoplasm of breast (C50)						
All origins[3]						
Both sexes						
2001	34.1	41.7	54.0	69.1	88.0	134.2
2000[4]	35.0	43.4	55.2	70.9	90.9	147.4
1999[4]	34.8	43.5	55.9	70.0	92.4	144.4
Male						
2001	0.8	1.2	1.7	2.3	2.4	3.5
2000[4]	0.8	1.1	1.7	2.1	2.4	4.2
1999[4]	0.7	1.1	1.4	1.9	2.3	4.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of breast (C50)														
All origins[3]														
Female														
2001	28.6	*	*	*	*	*	*	0.9	3.6	8.6	16.2	26.3	40.1	52.0
2000[4]	29.2	*	*	*	*	*	*	0.8	3.8	8.5	16.4	26.6	40.4	53.8
1999[4]	28.9	*	*	*	*	*	*	0.9	3.4	8.5	15.6	26.1	40.8	53.9
Hispanic														
Both sexes														
2001	4.8	*	*	*	*	*	*	*	1.1	3.5	5.7	9.6	15.2	20.5
2000[4]	4.8	*	*	*	*	*	*	*	1.2	3.2	5.6	10.4	14.5	21.2
1999[4]	4.6	*	*	*	*	*	*	*	1.1	2.7	5.6	9.5	15.7	18.0
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	9.7	*	*	*	*	*	*	*	2.5	7.3	11.5	19.2	29.5	39.1
2000[4]	9.7	*	*	*	*	*	*	*	2.6	6.6	11.3	20.7	28.0	39.9
1999[4]	9.3	*	*	*	*	*	*	*	2.4	5.7	11.4	18.8	30.4	33.6
Non-Hispanic[5]														
Both sexes														
2001	16.1	*	*	*	*	*	*	0.5	1.9	4.4	8.5	13.8	21.1	27.6
2000[4]	16.5	*	*	*	*	*	*	0.4	2.0	4.5	8.6	13.9	21.3	28.6
1999[4]	16.3	*	*	*	*	*	*	0.4	1.8	4.5	8.1	13.6	21.4	28.8
Male														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	0.4	0.5
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	0.3	0.7
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	0.4	0.6
Female														
2001	31.1	*	*	*	*	*	*	0.9	3.8	8.8	16.7	27.0	40.9	53.0
2000[4]	31.8	*	*	*	*	*	*	0.8	4.0	8.8	16.9	27.1	41.3	54.8
1999[4]	31.4	*	*	*	*	*	*	0.9	3.6	8.9	16.0	26.7	41.6	55.3
Non-Hispanic White														
Both sexes														
2001	17.0	*	*	*	*	*	*	0.4	1.6	3.9	7.5	12.6	19.8	26.2
2000[4]	17.4	*	*	*	*	*	*	0.4	1.7	4.0	7.7	12.5	20.1	27.2
1999[4]	17.1	*	*	*	*	*	*	0.4	1.5	3.9	7.2	12.3	20.1	27.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of breast (C50)						
All origins[3]						
Female						
2001	64.3	76.5	95.6	115.9	139.4	188.9
2000[4]	66.0	79.6	97.3	118.8	143.1	205.7
1999[4]	65.6	79.7	98.4	117.4	145.1	200.8
Hispanic						
Both sexes						
2001	24.1	25.1	31.7	38.8	40.7	68.8
2000[4]	20.7	30.5	32.3	41.3	53.5	71.7
1999[4]	21.6	25.2	31.5	37.1	55.2	86.3
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	44.7	44.2	55.5	64.9	64.3	102.4
2000[4]	38.2	54.7	55.2	69.5	84.2	105.8
1999[4]	40.3	45.0	54.0	62.8	87.2	127.9
Non-Hispanic[5]						
Both sexes						
2001	34.7	42.7	55.3	70.4	89.8	136.4
2000[4]	35.9	44.1	56.4	72.0	92.1	149.7
1999[4]	35.6	44.5	57.1	71.3	93.5	146.2
Male						
2001	0.8	1.2	1.8	2.3	2.4	3.7
2000[4]	0.8	1.1	1.7	2.2	2.4	4.2
1999[4]	0.7	1.1	1.4	1.9	2.4	4.7
Female						
2001	65.7	78.7	97.9	118.0	142.1	191.6
2000[4]	67.8	81.1	99.6	120.6	145.0	208.6
1999[4]	67.3	81.8	100.8	119.4	146.9	202.8
Non-Hispanic White						
Both sexes						
2001	33.9	41.9	54.9	70.8	89.9	137.1
2000[4]	35.3	44.0	56.8	72.3	92.9	149.6
1999[4]	34.6	43.8	56.8	71.0	94.0	147.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of breast (C50)														
Non-Hispanic White														
Male														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	0.3	0.5
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	0.3	0.6
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	0.3	0.5
Female														
2001	32.9	*	*	*	*	*	*	0.7	3.2	7.8	15.0	24.9	38.8	50.8
2000[4]	33.8	*	*	*	*	*	*	0.7	3.4	7.8	15.3	24.7	39.4	52.6
1999[4]	33.3	*	*	*	*	*	*	0.7	3.1	7.8	14.3	24.3	39.5	53.0
Non-Hispanic Black														
Both sexes														
2001	15.5	*	*	*	*	*	*	1.2	4.2	7.8	15.2	23.2	33.7	43.4
2000[4]	15.3	*	*	*	*	*	*	0.8	3.8	8.0	15.2	24.3	33.7	44.9
1999[4]	15.5	*	*	*	*	*	*	0.9	3.8	9.1	15.1	24.6	35.2	45.2
Male														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	29.2	*	*	*	*	*	*	2.3	7.8	14.7	28.3	42.9	61.5	78.3
2000[4]	28.9	*	*	*	*	*	*	*	7.2	14.8	28.4	45.0	61.9	80.5
1999[4]	29.2	*	*	*	*	*	*	1.8	7.1	16.8	28.2	45.6	64.2	81.0
Malignant neoplasm of cervix uteri (C53)														
All origins[3]														
Both sexes														
2001	1.4	*	*	*	*	*	*	0.3	0.8	1.3	1.8	2.3	2.6	2.6
2000[4]	1.5	*	*	*	*	*	*	0.4	0.7	1.3	1.8	2.2	2.7	3.1
1999[4]	1.5	*	*	*	*	*	*	0.3	0.8	1.3	1.8	2.4	2.7	3.0
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	2.8	*	*	*	*	*	*	0.5	1.5	2.6	3.7	4.5	5.1	5.1
2000[4]	2.9	*	*	*	*	*	*	0.8	1.5	2.6	3.5	4.4	5.2	6.0
1999[4]	3.0	*	*	*	*	*	*	0.6	1.5	2.5	3.6	4.6	5.2	5.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of breast (C50)						
Non-Hispanic White						
Male						
2001	0.8	1.1	1.7	2.1	2.4	3.5
2000[4]	0.8	1.1	1.7	2.2	2.2	4.0
1999[4]	0.6	1.0	1.2	1.8	2.3	4.5
Female						
2001	64.6	78.0	98.1	119.4	142.8	192.6
2000[4]	67.2	81.7	101.1	121.5	146.9	208.3
1999[4]	66.0	81.3	101.1	119.5	148.1	204.9
Non-Hispanic Black						
Both sexes						
2001	48.5	58.9	71.7	83.8	107.3	153.9
2000[4]	48.2	53.5	66.5	85.5	101.4	178.7
1999[4]	49.6	59.0	71.1	89.0	106.3	149.1
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	85.6	101.0	117.2	131.3	160.3	208.3
2000[4]	84.9	92.4	108.9	135.6	150.4	241.6
1999[4]	86.9	101.6	116.0	140.9	156.7	201.7
Malignant neoplasm of cervix uteri (C53)						
All origins[3]						
Both sexes						
2001	3.2	2.9	3.5	4.0	5.5	7.2
2000[4]	3.1	3.6	3.9	4.3	5.6	7.3
1999[4]	3.5	3.7	3.9	4.7	4.6	7.9
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	6.0	5.4	6.3	6.8	8.9	10.1
2000[4]	5.9	6.7	6.9	7.3	9.0	10.3
1999[4]	6.7	6.8	6.9	8.0	7.2	11.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of cervix uteri (C53)														
Hispanic														
Both sexes														
2001	1.1	*	*	*	*	*	*	*	0.8	1.4	2.4	3.1	2.8	3.9
2000[4]	1.2	*	*	*	*	*	*	*	0.6	1.9	2.3	2.6	2.9	4.2
1999[4]	1.1	*	*	*	*	*	*	*	0.8	1.2	2.4	3.3	3.4	3.7
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	2.3	*	*	*	*	*	*	*	1.7	3.0	5.0	6.3	5.6	7.4
2000[4]	2.5	*	*	*	*	*	*	*	1.4	4.0	4.8	5.3	5.7	7.9
1999[4]	2.3	*	*	*	*	*	*	*	1.7	2.6	4.9	6.5	6.7	7.1
Non-Hispanic[5]														
Both sexes														
2001	1.5	*	*	*	*	*	*	0.3	0.8	1.3	1.8	2.2	2.6	2.5
2000[4]	1.5	*	*	*	*	*	*	0.4	0.8	1.2	1.7	2.2	2.6	3.0
1999[4]	1.6	*	*	*	*	*	*	0.3	0.7	1.3	1.8	2.3	2.6	2.9
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	2.9	*	*	*	*	*	*	0.5	1.5	2.5	3.5	4.3	5.0	4.9
2000[4]	3.0	*	*	*	*	*	*	0.8	1.5	2.4	3.4	4.3	5.1	5.8
1999[4]	3.0	*	*	*	*	*	*	0.7	1.5	2.5	3.5	4.5	5.1	5.7
Non-Hispanic White														
Both sexes														
2001	1.4	*	*	*	*	*	*	0.2	0.7	1.2	1.6	2.0	2.3	2.2
2000[4]	1.4	*	*	*	*	*	*	0.4	0.7	1.1	1.4	1.9	2.3	2.6
1999[4]	1.4	*	*	*	*	*	*	0.4	0.7	1.2	1.6	2.1	2.4	2.5
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	2.7	*	*	*	*	*	*	0.5	1.4	2.4	3.2	4.0	4.5	4.2
2000[4]	2.7	*	*	*	*	*	*	0.8	1.4	2.1	2.9	3.8	4.5	5.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of cervix uteri (C53)						
Hispanic						
Both sexes						
2001	4.3	*	4.6	5.9	*	*
2000[4]	5.1	5.0	6.3	7.3	*	13.9
1999[4]	4.1	4.7	5.2	6.5	*	*
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	8.0	*	8.0	10.2	*	*
2000[4]	9.4	9.1	11.0	12.5	*	20.8
1999[4]	7.7	8.4	9.2	11.0	*	*
Non-Hispanic[5]						
Both sexes						
2001	3.1	3.0	3.4	3.9	5.4	7.0
2000[4]	2.9	3.5	3.7	4.1	5.5	7.1
1999[4]	3.5	3.6	3.8	4.6	4.5	7.9
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	5.9	5.5	6.1	6.7	8.6	10.0
2000[4]	5.5	6.5	6.7	7.0	8.7	9.9
1999[4]	6.6	6.7	6.8	7.8	7.1	11.0
Non-Hispanic White						
Both sexes						
2001	2.7	2.4	2.9	3.3	4.5	6.1
2000[4]	2.4	2.8	3.1	3.6	4.8	6.0
1999[4]	2.9	3.0	2.9	3.9	4.0	6.7
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	5.2	4.4	5.3	5.6	7.3	8.6
2000[4]	4.6	5.2	5.5	6.1	7.6	8.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of cervix uteri (C53)														
Non-Hispanic White														
Female														
1999[4]	2.8	*	*	*	*	*	*	0.8	1.5	2.3	3.2	4.2	4.7	5.0
Non-Hispanic Black														
Both sexes														
2001	2.2	*	*	*	*	*	*	*	1.2	1.9	3.0	3.3	4.5	4.7
2000[4]	2.5	*	*	*	*	*	*	*	1.5	2.1	3.2	3.8	5.0	6.2
1999[4]	2.4	*	*	*	*	*	*	*	1.0	2.1	3.0	3.1	4.2	6.3
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	4.2	*	*	*	*	*	*	*	2.2	3.6	5.6	6.1	8.3	8.6
2000[4]	4.7	*	*	*	*	*	*	*	2.8	4.0	6.0	7.0	9.3	11.4
1999[4]	4.5	*	*	*	*	*	*	*	1.9	4.0	5.6	5.7	7.7	11.4
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)														
All origins[3]														
Both sexes														
2001	2.4	*	*	*	*	*	*	*	0.1	0.2	0.4	0.9	2.0	3.2
2000[4]	2.3	*	*	*	*	*	*	*	0.1	0.2	0.4	0.9	1.8	3.5
1999[4]	2.3	*	*	*	*	*	*	*	*	0.2	0.5	0.9	1.7	3.3
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	4.7	*	*	*	*	*	*	*	0.3	0.4	0.8	1.8	4.0	6.2
2000[4]	4.6	*	*	*	*	*	*	*	0.2	0.4	0.9	1.9	3.5	6.9
1999[4]	4.5	*	*	*	*	*	*	*	*	0.3	0.9	1.8	3.3	6.3
Hispanic														
Both sexes														
2001	0.9	*	*	*	*	*	*	*	*	*	*	*	2.2	3.0
2000[4]	0.8	*	*	*	*	*	*	*	*	*	*	*	1.8	3.2
1999[4]	0.8	*	*	*	*	*	*	*	*	*	*	*	1.8	3.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of cervix uteri (C53)						
Non-Hispanic White						
Female						
1999[4]	5.5	5.7	5.3	6.6	6.3	9.3
Non-Hispanic Black						
Both sexes						
2001	6.1	8.0	7.8	9.7	15.5	19.0
2000[4]	6.6	9.6	10.0	10.9	13.8	19.7
1999[4]	8.3	8.7	13.3	12.4	9.7	21.7
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	10.8	13.8	13.1	15.6	23.3	26.1
2000[4]	11.7	16.7	16.6	17.4	20.7	27.0
1999[4]	14.9	15.2	22.1	19.8	14.5	29.8
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)						
All origins[3]						
Both sexes						
2001	5.8	8.2	11.4	14.5	19.2	24.7
2000[4]	6.0	8.4	11.1	14.0	18.6	24.7
1999[4]	5.5	8.8	10.9	14.4	19.4	24.0
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	11.1	15.3	20.4	24.6	30.7	35.0
2000[4]	11.4	15.6	19.8	23.8	29.5	34.8
1999[4]	10.5	16.2	19.5	24.5	30.7	33.7
Hispanic						
Both sexes						
2001	4.9	6.2	9.1	9.6	15.4	18.8
2000[4]	4.1	5.8	8.4	10.7	*	19.9
1999[4]	4.5	7.8	9.0	9.7	12.3	15.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)														
Hispanic														
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	1.8	*	*	*	*	*	*	*	*	*	*	*	4.2	5.8
2000[4]	1.6	*	*	*	*	*	*	*	*	*	*	*	3.4	6.2
1999[4]	1.6	*	*	*	*	*	*	*	*	*	*	*	3.5	5.8
Non-Hispanic[5]														
Both sexes														
2001	2.6	*	*	*	*	*	*	*	0.1	0.2	0.4	0.9	2.0	3.2
2000[4]	2.6	*	*	*	*	*	*	*	0.1	0.2	0.4	1.0	1.8	3.6
1999[4]	2.5	*	*	*	*	*	*	*	*	0.2	0.5	1.0	1.7	3.3
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	5.1	*	*	*	*	*	*	*	0.2	0.5	0.8	1.8	3.9	6.2
2000[4]	5.0	*	*	*	*	*	*	*	0.3	0.4	0.8	1.9	3.5	6.9
1999[4]	4.9	*	*	*	*	*	*	*	*	0.3	0.9	1.9	3.3	6.3
Non-Hispanic White														
Both sexes														
2001	2.7	*	*	*	*	*	*	*	*	0.2	0.4	0.9	1.9	3.1
2000[4]	2.6	*	*	*	*	*	*	*	*	0.2	0.4	1.0	1.7	3.4
1999[4]	2.6	*	*	*	*	*	*	*	*	0.2	0.4	0.9	1.6	3.0
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	5.2	*	*	*	*	*	*	*	*	0.4	0.8	1.7	3.7	6.1
2000[4]	5.1	*	*	*	*	*	*	*	*	0.4	0.7	1.9	3.4	6.7
1999[4]	5.1	*	*	*	*	*	*	*	*	0.3	0.8	1.9	3.2	5.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)						
Hispanic						
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	9.2	11.2	16.1	16.5	24.9	27.9
2000[4]	7.7	10.6	14.7	18.3	*	29.7
1999[4]	8.5	14.1	15.7	16.5	19.7	22.9
Non-Hispanic[5]						
Both sexes						
2001	5.9	8.4	11.5	14.7	19.3	24.8
2000[4]	6.1	8.6	11.2	14.2	18.8	24.9
1999[4]	5.6	8.8	11.0	14.6	19.6	24.3
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	11.2	15.6	20.6	25.0	30.9	35.1
2000[4]	11.6	15.9	20.0	24.0	29.9	34.9
1999[4]	10.7	16.4	19.7	24.8	31.1	34.1
Non-Hispanic White						
Both sexes						
2001	5.2	7.3	10.5	13.9	18.7	24.5
2000[4]	5.4	7.4	9.9	13.1	17.7	24.5
1999[4]	5.2	7.6	10.3	13.9	19.0	23.6
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	10.0	13.7	19.0	23.8	30.0	34.7
2000[4]	10.5	13.8	17.9	22.3	28.2	34.3
1999[4]	10.0	14.2	18.5	23.6	30.3	33.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)														
Non-Hispanic Black														
Both sexes														
2001	2.9	*	*	*	*	*	*	*	*	*	*	1.4	3.0	4.6
2000[4]	3.0	*	*	*	*	*	*	*	*	*	*	1.0	2.4	5.3
1999[4]	2.8	*	*	*	*	*	*	*	*	*	0.8	*	2.3	5.9
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	5.5	*	*	*	*	*	*	*	*	*	*	2.7	5.6	8.3
2000[4]	5.7	*	*	*	*	*	*	*	*	*	*	2.0	4.5	9.7
1999[4]	5.4	*	*	*	*	*	*	*	*	*	1.5	*	4.2	10.8
Malignant neoplasm of ovary (C56)														
All origins[3]														
Both sexes														
2001	5.1	*	*	*	*	*	*	0.2	0.3	0.7	1.4	2.9	5.5	8.3
2000[4]	5.0	*	*	*	*	*	*	0.2	0.3	0.6	1.5	3.1	5.5	8.3
1999[4]	4.9	*	*	*	*	*	*	0.1	0.2	0.7	1.5	3.1	5.5	8.3
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	9.9	*	*	*	*	*	*	0.4	0.6	1.4	2.8	5.8	10.8	16.0
2000[4]	9.8	*	*	*	*	*	*	0.4	0.6	1.3	2.9	6.0	10.8	16.1
1999[4]	9.6	*	*	*	*	*	*	0.3	0.5	1.4	3.0	6.2	10.8	16.0
Hispanic														
Both sexes														
2001	1.6	*	*	*	*	*	*	*	*	*	1.2	2.0	3.5	5.4
2000[4]	1.6	*	*	*	*	*	*	*	*	*	*	2.4	3.8	5.7
1999[4]	1.4	*	*	*	*	*	*	*	*	*	*	1.7	4.0	4.5
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)						
Non-Hispanic Black						
Both sexes						
2001	11.7	19.1	24.6	26.6	31.7	32.8
2000[4]	12.3	20.0	26.5	28.3	35.6	33.7
1999[4]	10.8	21.1	22.0	25.8	30.2	35.9
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	20.9	33.3	41.1	42.4	47.9	44.9
2000[4]	22.0	34.8	44.0	45.3	53.6	46.2
1999[4]	19.3	36.8	36.5	41.4	45.2	49.3
Malignant neoplasm of ovary (C56)						
All origins[3]						
Both sexes						
2001	13.0	17.6	22.6	30.2	37.4	39.6
2000[4]	12.2	17.5	21.8	30.9	36.5	40.9
1999[4]	12.6	17.0	22.4	29.2	35.5	37.6
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	24.7	32.8	40.5	51.4	59.8	56.1
2000[4]	23.3	32.5	39.0	52.5	58.0	57.6
1999[4]	24.0	31.6	39.9	49.5	56.2	52.8
Hispanic						
Both sexes						
2001	9.3	14.1	14.8	16.4	26.3	30.6
2000[4]	7.5	12.5	16.6	24.5	25.6	32.5
1999[4]	8.1	8.8	16.4	19.0	24.1	23.9
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of ovary (C56)														
Hispanic														
Female														
2001	3.4	*	*	*	*	*	*	*	*	*	2.5	4.1	6.9	10.3
2000[4]	3.4	*	*	*	*	*	*	*	*	*	*	4.7	7.5	10.9
1999[4]	2.9	*	*	*	*	*	*	*	*	*	*	3.4	7.7	8.6
Non-Hispanic[5]														
Both sexes														
2001	5.6	*	*	*	*	*	*	0.2	0.4	0.7	1.4	3.0	5.7	8.5
2000[4]	5.5	*	*	*	*	*	*	0.2	0.3	0.7	1.5	3.1	5.6	8.5
1999[4]	5.3	*	*	*	*	*	*	0.1	0.3	0.7	1.6	3.3	5.6	8.5
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	10.9	*	*	*	*	*	*	0.4	0.7	1.4	2.8	5.9	11.1	16.4
2000[4]	10.7	*	*	*	*	*	*	0.4	0.7	1.3	3.0	6.1	11.0	16.5
1999[4]	10.4	*	*	*	*	*	*	0.3	0.6	1.5	3.1	6.4	11.0	16.5
Non-Hispanic White														
Both sexes														
2001	6.2	*	*	*	*	*	*	0.2	0.3	0.7	1.5	3.2	5.9	8.7
2000[4]	6.2	*	*	*	*	*	*	0.2	0.3	0.7	1.6	3.3	5.8	8.9
1999[4]	6.0	*	*	*	*	*	*	*	0.3	0.8	1.7	3.4	5.8	8.9
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	12.2	*	*	*	*	*	*	0.4	0.7	1.4	2.9	6.4	11.8	17.1
2000[4]	12.0	*	*	*	*	*	*	0.5	0.7	1.4	3.2	6.6	11.4	17.5
1999[4]	11.7	*	*	*	*	*	*	*	0.5	1.5	3.3	6.8	11.4	17.3
Non-Hispanic Black														
Both sexes														
2001	3.2	*	*	*	*	*	*	*	*	0.9	1.4	2.0	4.5	8.1
2000[4]	3.0	*	*	*	*	*	*	*	*	*	1.6	2.2	4.8	6.6
1999[4]	3.1	*	*	*	*	*	*	*	*	*	1.2	2.4	5.2	7.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of ovary (C56)						
Hispanic						
Female						
2001	17.4	25.5	26.2	28.1	42.6	45.6
2000[4]	13.9	22.6	29.1	41.8	41.2	48.5
1999[4]	15.1	15.9	28.7	32.5	38.4	35.4
Non-Hispanic[5]						
Both sexes						
2001	13.2	17.8	23.0	30.9	37.8	39.8
2000[4]	12.6	17.8	22.1	31.2	36.8	41.2
1999[4]	12.9	17.5	22.6	29.5	35.7	38.1
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	25.2	33.3	41.4	52.4	60.4	56.4
2000[4]	24.0	33.1	39.5	52.9	58.4	57.8
1999[4]	24.5	32.6	40.3	50.0	56.6	53.3
Non-Hispanic White						
Both sexes						
2001	13.4	18.3	23.8	31.8	39.0	40.6
2000[4]	13.1	18.3	22.7	32.3	37.9	42.2
1999[4]	13.4	18.1	23.1	30.0	36.7	39.2
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	25.9	34.4	43.2	54.3	62.5	57.5
2000[4]	25.2	34.4	41.0	54.9	60.4	59.2
1999[4]	25.8	33.9	41.5	51.0	58.3	54.9
Non-Hispanic Black						
Both sexes						
2001	12.8	16.6	20.8	26.4	28.4	36.5
2000[4]	11.2	16.9	19.9	26.6	27.9	34.7
1999[4]	11.2	15.3	22.9	28.9	28.5	29.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of ovary (C56)														
Non-Hispanic Black														
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	6.1	*	*	*	*	*	*	*	*	1.7	2.7	3.7	8.3	14.8
2000[4]	5.8	*	*	*	*	*	*	*	*	*	3.0	4.1	9.0	11.9
1999[4]	5.8	*	*	*	*	*	*	*	*	*	2.2	4.5	9.6	13.4
Malignant neoplasm of prostate (C61)														
All origins[3]														
Both sexes														
2001	10.8	*	*	*	*	*	*	*	*	*	0.1	0.5	1.7	4.6
2000[4]	11.0	*	*	*	*	*	*	*	*	*	*	0.5	1.5	4.6
1999[4]	11.4	*	*	*	*	*	*	*	*	*	0.1	0.5	1.6	4.9
Male														
2001	22.0	*	*	*	*	*	*	*	*	*	0.2	1.0	3.5	9.4
2000[4]	22.5	*	*	*	*	*	*	*	*	*	*	1.1	3.1	9.6
1999[4]	23.2	*	*	*	*	*	*	*	*	*	0.3	1.0	3.3	10.1
Female														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Hispanic														
Both sexes														
2001	3.3	*	*	*	*	*	*	*	*	*	*	*	1.5	3.8
2000[4]	2.9	*	*	*	*	*	*	*	*	*	*	*	*	2.2
1999[4]	3.1	*	*	*	*	*	*	*	*	*	*	*	1.6	4.4
Male														
2001	6.4	*	*	*	*	*	*	*	*	*	*	*	3.0	8.0
2000[4]	5.7	*	*	*	*	*	*	*	*	*	*	*	*	4.6
1999[4]	6.0	*	*	*	*	*	*	*	*	*	*	*	3.2	9.2
Female														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of ovary (C56)						
Non-Hispanic Black						
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	22.8	28.9	34.6	42.2	42.9	50.0
2000[4]	20.0	29.3	33.1	42.7	41.9	47.5
1999[4]	20.0	26.6	38.1	46.3	42.6	40.0
Malignant neoplasm of prostate (C61)						
All origins[3]						
Both sexes						
2001	11.3	25.0	47.4	82.4	133.2	200.9
2000[4]	12.1	25.9	49.0	85.1	136.6	209.3
1999[4]	12.3	28.4	51.9	89.0	141.4	207.5
Male						
2001	23.8	53.9	107.1	200.1	355.4	681.2
2000[4]	25.4	56.0	111.3	207.4	368.2	723.1
1999[4]	25.8	61.6	118.3	216.9	383.0	722.4
Female						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Hispanic						
Both sexes						
2001	8.6	22.5	37.6	78.7	103.8	170.1
2000[4]	7.6	15.9	38.1	63.7	107.5	165.9
1999[4]	7.2	23.3	35.0	67.5	104.0	176.1
Male						
2001	18.5	50.3	86.9	189.6	271.4	518.0
2000[4]	16.4	35.4	88.5	153.5	284.2	503.9
1999[4]	15.5	52.2	81.5	163.1	278.2	541.3
Female						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of prostate (C61)														
Non-Hispanic[5]														
Both sexes														
2001	11.9	*	*	*	*	*	*	*	*	*	0.1	0.5	1.7	4.6
2000[4]	12.2	*	*	*	*	*	*	*	*	*	*	0.6	1.5	4.8
1999[4]	12.5	*	*	*	*	*	*	*	*	*	0.1	0.5	1.6	4.9
Male														
2001	24.3	*	*	*	*	*	*	*	*	*	0.3	1.0	3.5	9.5
2000[4]	25.0	*	*	*	*	*	*	*	*	*	*	1.1	3.1	9.9
1999[4]	25.6	*	*	*	*	*	*	*	*	*	0.3	1.1	3.3	10.1
Female														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic White														
Both sexes														
2001	12.0	*	*	*	*	*	*	*	*	*	*	0.4	1.4	3.9
2000[4]	12.3	*	*	*	*	*	*	*	*	*	*	0.4	1.3	4.0
1999[4]	12.6	*	*	*	*	*	*	*	*	*	*	0.4	1.4	4.1
Male														
2001	24.6	*	*	*	*	*	*	*	*	*	*	0.8	2.8	8.0
2000[4]	25.2	*	*	*	*	*	*	*	*	*	*	0.8	2.6	8.1
1999[4]	25.9	*	*	*	*	*	*	*	*	*	*	0.9	2.7	8.5
Female														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic Black														
Both sexes														
2001	14.6	*	*	*	*	*	*	*	*	*	*	1.2	4.9	11.8
2000[4]	15.1	*	*	*	*	*	*	*	*	*	*	1.5	3.6	13.3
1999[4]	15.3	*	*	*	*	*	*	*	*	*	*	1.4	4.3	12.3
Male														
2001	30.8	*	*	*	*	*	*	*	*	*	*	2.5	10.7	26.2
2000[4]	31.7	*	*	*	*	*	*	*	*	*	*	3.3	7.8	29.5
1999[4]	32.2	*	*	*	*	*	*	*	*	*	*	3.0	9.4	27.4
Female														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of prostate (C61)						
Non-Hispanic[5]						
Both sexes						
2001	11.5	25.1	47.9	82.3	134.0	201.6
2000[4]	12.4	26.5	49.5	85.8	137.4	210.4
1999[4]	12.6	28.7	52.6	89.7	142.5	208.2
Male						
2001	24.1	54.1	108.1	199.9	357.9	686.3
2000[4]	26.0	57.3	112.1	209.2	370.6	730.7
1999[4]	26.4	62.1	119.8	218.8	386.3	728.2
Female						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic White						
Both sexes						
2001	9.7	22.0	42.4	74.8	123.6	189.5
2000[4]	10.6	22.8	43.3	78.1	126.3	199.3
1999[4]	10.3	24.9	46.8	81.5	133.8	195.4
Male						
2001	20.2	46.9	94.5	180.5	328.4	645.0
2000[4]	22.0	48.7	97.0	189.3	338.7	693.6
1999[4]	21.5	53.3	105.6	197.6	360.7	685.4
Female						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic Black						
Both sexes						
2001	29.3	58.2	116.5	182.0	283.4	365.8
2000[4]	30.0	66.1	122.3	190.2	295.1	359.6
1999[4]	34.0	70.0	125.0	198.4	274.4	368.6
Male						
2001	66.5	137.0	291.3	486.8	839.7	1,353.6
2000[4]	68.0	155.6	306.5	505.8	881.4	1,328.0
1999[4]	77.3	164.4	313.3	527.8	828.0	1,354.9
Female						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of prostate (C61)														
Non-Hispanic Black														
Female														
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Malignant neoplasms of kidney and renal pelvis (C64-C65)														
All origins[3]														
Both sexes														
2001	4.2	*	0.1	0.1	*	*	*	*	0.2	0.3	1.2	2.4	4.4	7.1
2000[4]	4.2	*	*	0.1	*	*	*	*	0.1	0.4	1.1	2.2	4.5	7.6
1999[4]	4.0	*	*	0.1	*	*	*	0.1	0.1	0.5	1.1	2.2	4.3	7.3
Male														
2001	5.4	*	*	*	*	*	*	*	*	0.5	1.6	3.5	6.5	10.5
2000[4]	5.3	*	*	*	*	*	*	*	*	0.5	1.6	3.1	6.6	11.2
1999[4]	5.0	*	*	*	*	*	*	*	0.2	0.6	1.5	3.2	6.1	10.6
Female														
2001	3.1	*	*	0.2	*	*	*	*	*	*	0.9	1.4	2.4	3.9
2000[4]	3.1	*	*	*	*	*	*	*	*	0.3	0.7	1.4	2.4	4.3
1999[4]	3.0	*	*	*	*	*	*	*	*	0.3	0.6	1.2	2.5	4.3
Hispanic														
Both sexes														
2001	1.7	*	*	*	*	*	*	*	*	*	1.3	2.6	4.3	4.6
2000[4]	1.6	*	*	*	*	*	*	*	*	*	1.1	2.4	3.4	6.2
1999[4]	1.6	*	*	*	*	*	*	*	*	*	1.0	1.9	4.1	6.7
Male														
2001	2.2	*	*	*	*	*	*	*	*	*	2.1	3.8	6.0	6.6
2000[4]	2.0	*	*	*	*	*	*	*	*	*	1.8	3.4	4.5	10.1
1999[4]	2.0	*	*	*	*	*	*	*	*	*	*	3.3	5.6	8.8
Female														
2001	1.2	*	*	*	*	*	*	*	*	*	*	*	2.6	*
2000[4]	1.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.2	*	*	*	*	*	*	*	*	*	*	*	*	4.8
Non-Hispanic[5]														
Both sexes														
2001	4.6	*	*	0.2	*	*	*	*	0.2	0.3	1.2	2.4	4.4	7.3
2000[4]	4.5	*	*	*	*	*	*	*	0.1	0.4	1.1	2.2	4.6	7.7
1999[4]	4.3	*	*	*	*	*	*	0.1	0.1	0.5	1.1	2.2	4.3	7.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of prostate (C61)						
Non-Hispanic Black						
Female						
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Malignant neoplasms of kidney and renal pelvis (C64-C65)						
All origins[3]						
Both sexes						
2001	11.9	14.6	20.8	24.4	29.1	32.3
2000[4]	10.8	15.0	19.4	24.5	29.6	33.8
1999[4]	10.8	14.4	19.2	23.1	26.9	31.4
Male						
2001	17.2	21.0	30.0	34.4	42.2	52.9
2000[4]	15.6	21.4	27.4	34.6	44.2	55.1
1999[4]	15.8	20.3	27.3	33.7	38.9	49.4
Female						
2001	7.0	9.1	13.5	17.4	21.2	23.7
2000[4]	6.5	9.5	13.0	17.5	21.0	25.1
1999[4]	6.2	9.3	12.8	15.7	19.9	24.1
Hispanic						
Both sexes						
2001	10.0	12.2	17.0	24.4	21.4	31.9
2000[4]	10.0	10.8	14.0	21.4	30.6	25.9
1999[4]	9.4	13.3	14.6	21.0	25.3	28.8
Male						
2001	14.4	17.7	26.1	35.5	35.1	57.1
2000[4]	13.2	14.9	21.9	34.7	45.6	40.3
1999[4]	15.8	18.9	23.9	29.6	*	*
Female						
2001	6.1	7.7	10.1	16.5	*	19.5
2000[4]	7.2	7.5	8.1	12.0	21.5	*
1999[4]	*	8.8	7.7	14.9	22.5	23.9
Non-Hispanic[5]						
Both sexes						
2001	12.0	14.8	21.0	24.4	29.3	32.3
2000[4]	10.9	15.2	19.5	24.6	29.5	33.9
1999[4]	10.9	14.4	19.4	23.1	26.9	31.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of kidney and renal pelvis (C64-C65)														
Non-Hispanic[5]														
Male														
2001	5.9	*	*	*	*	*	*	*	*	0.5	1.5	3.4	6.5	10.8
2000[4]	5.7	*	*	*	*	*	*	*	*	0.5	1.5	3.1	6.8	11.2
1999[4]	5.4	*	*	*	*	*	*	*	0.2	0.6	1.5	3.2	6.1	10.8
Female														
2001	3.4	*	*	0.3	*	*	*	*	*	*	1.0	1.4	2.4	4.0
2000[4]	3.4	*	*	*	*	*	*	*	*	0.3	0.7	1.4	2.4	4.4
1999[4]	3.2	*	*	*	*	*	*	*	*	0.3	0.6	1.2	2.5	4.2
Non-Hispanic White														
Both sexes														
2001	5.1	*	*	*	*	*	*	*	*	0.3	1.2	2.4	4.4	7.5
2000[4]	5.0	*	*	*	*	*	*	*	*	0.3	1.1	2.3	4.7	7.9
1999[4]	4.7	*	*	*	*	*	*	*	*	0.4	1.1	2.1	4.3	7.5
Male														
2001	6.5	*	*	*	*	*	*	*	*	0.4	1.5	3.6	6.5	11.1
2000[4]	6.3	*	*	*	*	*	*	*	*	0.5	1.6	3.2	6.9	11.4
1999[4]	6.0	*	*	*	*	*	*	*	*	0.5	1.5	3.1	6.0	10.9
Female														
2001	3.8	*	*	*	*	*	*	*	*	*	0.8	1.3	2.4	4.0
2000[4]	3.8	*	*	*	*	*	*	*	*	*	0.7	1.4	2.5	4.6
1999[4]	3.5	*	*	*	*	*	*	*	*	0.3	0.6	1.2	2.6	4.4
Non-Hispanic Black														
Both sexes														
2001	3.1	*	*	*	*	*	*	*	*	0.7	1.8	2.6	5.1	7.5
2000[4]	2.9	*	*	*	*	*	*	*	*	0.7	1.1	2.5	4.6	7.5
1999[4]	3.0	*	*	*	*	*	*	*	*	1.0	1.3	3.0	5.1	7.7
Male														
2001	4.0	*	*	*	*	*	*	*	*	*	1.9	3.5	8.0	10.8
2000[4]	3.7	*	*	*	*	*	*	*	*	*	*	3.3	7.1	11.5
1999[4]	3.7	*	*	*	*	*	*	*	*	*	1.8	4.3	8.2	12.5
Female														
2001	2.3	*	*	*	*	*	*	*	*	*	1.8	1.9	2.7	4.9
2000[4]	2.2	*	*	*	*	*	*	*	*	*	*	1.9	2.5	4.3
1999[4]	2.2	*	*	*	*	*	*	*	*	*	*	1.9	2.4	3.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of kidney and renal pelvis (C64-C65)						
Non-Hispanic[5]						
Male						
2001	17.4	21.2	30.1	34.3	42.4	52.6
2000[4]	15.8	21.7	27.5	34.5	44.1	55.4
1999[4]	15.8	20.4	27.5	33.8	39.2	49.8
Female						
2001	7.1	9.2	13.7	17.4	21.5	23.8
2000[4]	6.4	9.6	13.3	17.7	20.9	25.2
1999[4]	6.4	9.4	13.1	15.6	19.7	24.1
Non-Hispanic White						
Both sexes						
2001	12.2	15.4	21.5	24.7	29.7	32.8
2000[4]	11.2	15.5	19.9	24.9	29.9	34.6
1999[4]	11.3	14.8	19.7	23.3	27.3	31.5
Male						
2001	17.3	22.0	30.8	34.3	42.9	54.3
2000[4]	16.2	21.7	27.8	34.9	44.5	56.9
1999[4]	16.4	20.7	27.7	34.1	40.4	50.2
Female						
2001	7.5	9.6	14.0	17.9	21.7	23.8
2000[4]	6.4	10.1	13.5	17.9	21.3	25.7
1999[4]	6.5	9.7	13.3	15.7	19.5	24.1
Non-Hispanic Black						
Both sexes						
2001	12.5	12.0	19.4	24.3	28.7	30.0
2000[4]	10.7	15.2	18.7	23.9	29.0	29.6
1999[4]	9.9	13.8	19.5	23.4	24.7	31.7
Male						
2001	21.1	17.5	29.6	39.3	43.3	46.2
2000[4]	15.7	25.6	28.4	32.8	48.1	49.3
1999[4]	14.1	20.5	28.4	34.8	28.3	49.9
Female						
2001	5.8	7.9	12.6	15.3	21.2	24.0
2000[4]	6.7	7.5	12.3	18.6	19.4	22.2
1999[4]	6.5	8.9	13.6	16.6	22.8	24.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of bladder (C67)														
All origins[3]														
Both sexes														
2001	4.3	*	*	*	*	*	*	*	*	0.2	0.4	0.8	1.7	3.4
2000[4]	4.3	*	*	*	*	*	*	*	*	0.2	0.3	0.8	1.6	3.6
1999[4]	4.3	*	*	*	*	*	*	*	*	0.1	0.3	0.7	1.9	3.8
Male														
2001	6.0	*	*	*	*	*	*	*	*	*	0.5	1.2	2.6	5.2
2000[4]	5.9	*	*	*	*	*	*	*	*	0.2	0.4	1.1	2.5	5.7
1999[4]	5.9	*	*	*	*	*	*	*	*	0.2	0.4	1.1	2.9	5.9
Female														
2001	2.7	*	*	*	*	*	*	*	*	0.2	0.3	0.4	0.9	1.7
2000[4]	2.7	*	*	*	*	*	*	*	*	*	0.3	0.5	0.8	1.7
1999[4]	2.7	*	*	*	*	*	*	*	*	*	0.2	0.3	0.9	1.7
Hispanic														
Both sexes														
2001	0.9	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.9	*	*	*	*	*	*	*	*	*	*	*	*	2.6
1999[4]	0.9	*	*	*	*	*	*	*	*	*	*	*	*	2.3
Male														
2001	1.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.2	*	*	*	*	*	*	*	*	*	*	*	*	4.6
1999[4]	1.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	4.8	*	*	*	*	*	*	*	*	0.2	0.4	0.8	1.8	3.6
2000[4]	4.7	*	*	*	*	*	*	*	*	0.2	0.3	0.8	1.7	3.7
1999[4]	4.7	*	*	*	*	*	*	*	*	0.2	0.3	0.8	1.9	3.9
Male														
2001	6.7	*	*	*	*	*	*	*	*	*	0.5	1.2	2.8	5.4
2000[4]	6.6	*	*	*	*	*	*	*	*	*	0.4	1.2	2.6	5.7
1999[4]	6.6	*	*	*	*	*	*	*	*	*	0.4	1.2	3.0	6.0
Female														
2001	3.0	*	*	*	*	*	*	*	*	*	0.4	0.4	0.9	1.8
2000[4]	3.0	*	*	*	*	*	*	*	*	*	0.3	0.5	0.8	1.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of bladder (C67)						
All origins[3]						
Both sexes						
2001	7.0	12.0	19.0	29.3	44.8	68.9
2000[4]	6.8	11.7	19.6	29.1	45.5	69.6
1999[4]	7.0	11.8	20.2	29.7	43.8	68.5
Male						
2001	11.3	19.5	31.6	50.8	81.0	134.8
2000[4]	10.6	19.0	32.2	50.6	82.4	139.4
1999[4]	11.0	19.3	32.9	52.3	79.5	134.8
Female						
2001	3.2	5.6	9.0	14.3	23.2	41.4
2000[4]	3.4	5.5	9.7	14.0	23.8	41.2
1999[4]	3.4	5.4	10.3	13.9	22.9	41.8
Hispanic						
Both sexes						
2001	3.2	6.3	9.9	14.4	26.3	46.3
2000[4]	3.3	3.5	10.5	14.1	24.0	47.1
1999[4]	4.1	6.0	11.6	13.2	22.3	38.6
Male						
2001	6.0	9.6	14.6	21.1	46.7	87.6
2000[4]	*	*	16.5	28.8	44.2	84.6
1999[4]	7.4	11.6	19.9	21.8	37.7	77.6
Female						
2001	*	*	*	9.7	*	26.1
2000[4]	*	*	*	*	*	28.7
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	7.3	12.4	19.5	30.0	45.5	69.6
2000[4]	7.1	12.3	20.1	29.7	46.2	70.3
1999[4]	7.2	12.1	20.7	30.3	44.5	69.3
Male						
2001	11.6	20.1	32.6	52.2	82.2	136.4
2000[4]	11.1	19.9	33.0	51.5	83.5	141.3
1999[4]	11.2	19.7	33.6	53.5	81.0	136.6
Female						
2001	3.3	5.7	9.1	14.5	23.6	41.9
2000[4]	3.4	5.7	9.9	14.4	24.2	41.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of bladder (C67)														
Non-Hispanic[5]														
Female														
1999[4]	3.0	*	*	*	*	*	*	*	*	*	0.2	0.3	0.9	1.8
Non-Hispanic White														
Both sexes														
2001	5.5	*	*	*	*	*	*	*	*	0.2	0.4	0.8	1.9	3.7
2000[4]	5.4	*	*	*	*	*	*	*	*	0.2	0.4	0.9	1.8	3.9
1999[4]	5.4	*	*	*	*	*	*	*	*	0.1	0.3	0.7	2.0	3.9
Male														
2001	7.8	*	*	*	*	*	*	*	*	*	0.5	1.2	2.9	5.6
2000[4]	7.7	*	*	*	*	*	*	*	*	*	0.4	1.2	2.8	6.0
1999[4]	7.6	*	*	*	*	*	*	*	*	*	0.4	1.1	3.0	6.3
Female														
2001	3.2	*	*	*	*	*	*	*	*	*	0.4	0.3	0.9	1.8
2000[4]	3.3	*	*	*	*	*	*	*	*	*	0.3	0.5	0.8	1.8
1999[4]	3.2	*	*	*	*	*	*	*	*	*	0.3	0.3	1.0	1.6
Non-Hispanic Black														
Both sexes														
2001	2.4	*	*	*	*	*	*	*	*	*	*	0.8	1.8	3.5
2000[4]	2.4	*	*	*	*	*	*	*	*	*	*	0.9	1.3	3.0
1999[4]	2.6	*	*	*	*	*	*	*	*	*	*	1.0	2.2	4.6
Male														
2001	2.6	*	*	*	*	*	*	*	*	*	*	*	2.9	4.3
2000[4]	2.8	*	*	*	*	*	*	*	*	*	*	*	*	4.7
1999[4]	3.0	*	*	*	*	*	*	*	*	*	*	*	3.8	5.6
Female														
2001	2.3	*	*	*	*	*	*	*	*	*	*	*	*	2.8
2000[4]	2.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	2.3	*	*	*	*	*	*	*	*	*	*	*	*	3.8
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)														
All origins[3]														
Both sexes														
2001	4.4	*	0.7	0.8	0.6	0.5	0.6	0.7	1.4	1.8	2.8	4.0	6.1	8.3
2000[4]	4.5	0.5	0.7	0.8	0.7	0.5	0.6	0.8	1.2	2.0	2.9	4.1	6.3	8.2
1999[4]	4.6	*	0.7	0.9	0.6	0.6	0.6	0.9	1.3	1.9	2.8	4.3	6.1	9.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of bladder (C67)						
Non-Hispanic[5]						
Female						
1999[4]	3.6	5.6	10.6	14.1	23.2	42.3
Non-Hispanic White						
Both sexes						
2001	7.6	13.0	20.4	31.0	46.4	71.5
2000[4]	7.5	12.9	20.8	30.4	47.1	72.8
1999[4]	7.5	12.5	21.3	31.2	45.8	70.7
Male						
2001	12.4	21.4	34.3	54.7	85.0	143.0
2000[4]	11.9	21.1	34.5	53.6	85.9	148.0
1999[4]	11.7	20.6	35.2	55.6	83.7	143.0
Female						
2001	3.2	5.6	9.2	14.3	23.1	41.7
2000[4]	3.4	5.8	9.8	14.0	24.1	42.5
1999[4]	3.5	5.4	10.3	14.0	23.5	41.9
Non-Hispanic Black						
Both sexes						
2001	7.1	10.3	14.3	24.8	41.7	55.6
2000[4]	5.5	9.7	18.0	28.8	44.5	49.6
1999[4]	6.8	11.9	19.2	25.4	35.8	60.8
Male						
2001	9.4	13.0	21.0	34.1	58.9	78.5
2000[4]	7.4	12.8	24.3	41.4	73.0	88.1
1999[4]	9.6	15.7	23.9	39.2	60.2	86.8
Female						
2001	5.3	8.3	9.9	19.2	32.9	47.1
2000[4]	4.0	7.5	13.8	21.2	30.2	35.3
1999[4]	4.6	9.1	16.1	17.2	23.7	51.1
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)						
All origins[3]						
Both sexes						
2001	11.6	14.2	16.7	20.3	20.6	16.4
2000[4]	11.4	14.8	18.3	19.4	21.1	16.9
1999[4]	11.7	15.7	18.3	21.0	19.4	16.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)														
All origins[3]														
Male														
2001	5.0	*	0.7	0.8	0.6	0.6	0.7	0.9	1.7	2.2	3.5	4.9	7.5	10.7
2000[4]	5.1	*	0.7	0.8	0.8	0.6	0.7	1.0	1.6	2.6	3.7	4.9	7.7	10.1
1999[4]	5.1	*	0.7	0.9	0.6	0.7	0.7	1.1	1.7	2.3	3.4	5.2	7.4	10.6
Female														
2001	3.9	*	0.6	0.9	0.6	0.4	0.6	0.5	1.1	1.4	2.1	3.1	4.8	6.0
2000[4]	3.9	*	0.6	0.8	0.6	0.4	0.5	0.7	0.8	1.4	2.0	3.3	4.9	6.5
1999[4]	4.1	*	0.8	0.9	0.7	0.5	0.5	0.8	0.9	1.5	2.3	3.4	4.9	7.9
Hispanic														
Both sexes														
2001	1.7	*	0.7	0.6	*	*	*	*	0.7	1.3	1.5	2.2	3.0	4.5
2000[4]	1.8	*	*	1.0	0.9	*	*	0.6	0.8	1.1	2.0	2.3	3.7	3.7
1999[4]	1.7	*	0.9	0.9	*	*	*	0.6	*	1.1	1.5	2.2	4.0	5.2
Male														
2001	1.8	*	*	*	*	*	*	*	*	1.3	1.6	3.0	2.9	6.2
2000[4]	1.9	*	*	1.3	*	*	*	*	*	*	1.9	2.6	*	4.4
1999[4]	1.7	*	*	*	*	*	*	*	*	1.4	2.2	*	3.8	5.5
Female														
2001	1.6	*	*	*	*	*	*	*	*	*	*	*	3.2	*
2000[4]	1.7	*	*	*	*	*	*	*	*	*	2.1	*	4.7	*
1999[4]	1.7	*	*	*	*	*	*	*	*	*	*	*	4.1	4.8
Non-Hispanic[5]														
Both sexes														
2001	4.8	*	0.6	0.9	0.7	0.5	0.7	0.8	1.5	1.9	2.9	4.2	6.3	8.6
2000[4]	4.9	*	0.7	0.8	0.7	0.5	0.7	0.9	1.3	2.1	3.0	4.2	6.5	8.6
1999[4]	5.0	*	0.7	0.9	0.6	0.7	0.6	1.0	1.5	2.0	2.9	4.5	6.3	9.5
Male														
2001	5.5	*	0.6	0.8	0.7	0.6	0.7	1.0	1.9	2.3	3.7	5.1	7.9	11.0
2000[4]	5.5	*	0.7	0.7	0.8	0.6	0.8	1.0	1.8	2.8	3.9	5.1	8.1	10.6
1999[4]	5.6	*	0.7	0.9	0.6	0.8	0.7	1.1	1.9	2.5	3.5	5.5	7.7	10.9
Female														
2001	4.2	*	0.6	1.0	0.6	0.4	0.6	0.5	1.1	1.4	2.1	3.3	4.9	6.3
2000[4]	4.2	*	0.7	0.8	0.6	0.4	0.6	0.7	0.8	1.5	2.0	3.4	4.9	6.7
1999[4]	4.4	*	0.7	0.9	0.7	0.5	0.4	0.8	1.0	1.6	2.4	3.4	5.0	8.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)						
All origins[3]						
Male						
2001	13.9	17.3	21.2	25.4	26.5	20.1
2000[4]	14.0	17.9	22.8	23.2	26.3	22.4
1999[4]	14.7	19.5	22.3	26.1	24.5	20.7
Female						
2001	9.5	11.5	13.1	16.7	17.1	14.8
2000[4]	9.0	12.1	14.8	16.7	18.0	14.7
1999[4]	8.9	12.4	15.1	17.5	16.5	14.1
Hispanic						
Both sexes						
2001	7.8	9.5	11.3	13.3	17.4	*
2000[4]	9.3	9.7	9.8	16.2	*	15.9
1999[4]	7.2	8.1	12.5	12.3	17.0	*
Male						
2001	7.6	11.3	18.3	16.4	*	*
2000[4]	12.1	11.9	13.1	19.9	*	*
1999[4]	10.7	9.7	11.7	*	*	*
Female						
2001	8.0	8.0	*	11.1	*	*
2000[4]	6.9	7.9	7.4	13.6	*	*
1999[4]	*	6.9	13.0	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	11.8	14.5	17.1	20.6	20.7	16.5
2000[4]	11.5	15.1	18.8	19.5	21.3	16.9
1999[4]	11.9	16.2	18.5	21.3	19.5	16.1
Male						
2001	14.3	17.7	21.4	25.8	26.7	20.5
2000[4]	14.1	18.2	23.3	23.3	26.5	22.3
1999[4]	14.9	20.1	22.9	26.5	24.6	21.3
Female						
2001	9.5	11.7	13.6	16.9	17.1	14.9
2000[4]	9.1	12.4	15.2	16.8	18.3	14.7
1999[4]	9.2	12.8	15.1	17.8	16.5	14.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)														
Non-Hispanic White														
Both sexes														
2001	5.6	*	0.7	1.0	0.6	0.5	0.7	0.9	1.7	2.2	3.2	4.6	7.0	9.2
2000[4]	5.6	*	0.7	0.8	0.7	0.5	0.7	1.0	1.4	2.4	3.2	4.7	7.2	9.2
1999[4]	5.7	*	0.7	0.9	0.6	0.6	0.6	1.2	1.7	2.3	3.2	5.0	6.8	10.4
Male														
2001	6.3	*	0.7	0.8	0.6	0.6	0.8	1.2	2.2	2.8	4.1	5.6	8.6	11.8
2000[4]	6.3	*	0.7	0.7	0.8	0.6	0.8	1.2	1.9	3.1	4.2	5.7	8.7	11.4
1999[4]	6.4	*	0.8	1.0	0.6	0.7	0.7	1.4	2.1	2.8	3.7	6.0	8.2	11.8
Female														
2001	4.8	*	0.6	1.1	0.6	0.5	0.6	0.6	1.3	1.6	2.4	3.6	5.4	6.7
2000[4]	4.9	*	0.7	0.8	0.7	0.4	0.7	0.8	1.0	1.6	2.2	3.7	5.6	7.2
1999[4]	5.0	*	0.7	0.8	0.7	0.6	0.5	1.0	1.2	1.8	2.7	3.9	5.5	9.0
Non-Hispanic Black														
Both sexes														
2001	2.1	*	*	*	0.8	*	*	*	0.8	*	1.6	2.7	3.5	5.5
2000[4]	2.2	*	*	0.8	0.8	*	*	*	0.8	1.1	2.1	2.2	3.7	5.5
1999[4]	2.2	*	*	1.1	*	0.8	*	*	1.0	0.9	1.9	2.7	4.0	5.5
Male														
2001	2.2	*	*	*	*	*	*	*	*	*	1.9	3.1	4.6	7.1
2000[4]	2.4	*	*	*	*	*	*	*	*	*	2.7	2.2	5.7	6.6
1999[4]	2.4	*	*	*	*	*	*	*	*	*	2.6	3.8	5.3	6.1
Female														
2001	1.9	*	*	*	*	*	*	*	*	*	1.4	2.3	2.7	4.2
2000[4]	1.9	*	*	*	*	*	*	*	*	*	1.5	2.3	2.0	4.6
1999[4]	1.9	*	*	*	*	*	*	*	*	*	*	1.7	2.9	5.0
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)														
All origins[3]														
Both sexes														
2001	19.8	0.7	1.0	0.8	1.0	1.5	2.0	2.3	2.7	3.5	4.8	8.1	13.0	22.7
2000[4]	20.1	1.0	0.9	0.8	1.0	1.6	2.1	2.1	2.5	3.4	4.8	8.3	13.4	23.4
1999[4]	20.2	0.8	0.9	0.7	1.0	1.5	2.2	2.2	2.6	3.7	4.8	8.4	13.6	25.6
Male														
2001	21.5	*	1.0	0.9	1.1	1.8	2.3	2.6	3.1	4.2	5.9	9.8	16.0	27.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)						
Non-Hispanic White						
Both sexes						
2001	12.9	15.6	18.2	22.0	21.8	17.2
2000[4]	12.5	16.4	20.3	20.6	22.1	17.5
1999[4]	12.9	17.4	20.0	22.6	20.4	16.7
Male						
2001	15.4	19.1	22.8	27.2	28.0	20.9
2000[4]	15.3	19.8	24.9	24.5	27.7	23.5
1999[4]	16.0	21.6	24.6	27.9	25.9	22.4
Female						
2001	10.6	12.4	14.5	18.3	18.0	15.7
2000[4]	9.9	13.4	16.6	17.8	18.8	15.1
1999[4]	10.1	13.8	16.4	18.8	17.1	14.4
Non-Hispanic Black						
Both sexes						
2001	6.3	9.2	9.6	9.2	11.0	11.9
2000[4]	6.2	7.6	9.0	10.5	14.7	10.5
1999[4]	6.9	8.7	7.6	12.0	11.2	10.0
Male						
2001	9.0	9.4	11.7	11.9	*	*
2000[4]	7.9	7.7	12.6	12.0	*	*
1999[4]	8.9	11.2	9.0	14.5	*	*
Female						
2001	4.3	9.0	8.3	7.6	10.4	9.8
2000[4]	4.9	7.5	6.6	9.6	14.3	10.5
1999[4]	5.3	6.9	6.7	10.5	12.3	9.3
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)						
All origins[3]						
Both sexes						
2001	38.1	58.8	91.8	129.0	168.5	199.9
2000[4]	38.2	62.2	91.6	131.4	176.0	204.6
1999[4]	39.4	62.8	93.9	130.7	174.0	199.4
Male						
2001	47.5	75.2	116.9	165.7	220.9	275.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)														
All origins[3]														
Male														
2000[4]	21.6	1.1	1.0	0.9	1.2	1.9	2.6	2.3	3.0	3.9	5.7	10.3	16.1	28.4
1999[4]	21.7	*	1.0	0.8	1.2	1.7	2.7	2.4	3.1	4.4	5.9	10.2	16.7	31.0
Female														
2001	18.1	*	1.0	0.7	0.8	1.2	1.7	2.0	2.2	2.7	3.7	6.4	10.1	18.1
2000[4]	18.6	*	0.9	0.7	0.8	1.2	1.5	1.9	1.9	3.0	3.9	6.5	10.8	18.8
1999[4]	18.7	*	0.8	0.6	0.9	1.3	1.8	1.9	2.2	3.0	3.8	6.6	10.6	20.5
Hispanic														
Both sexes														
2001	7.4	*	1.3	1.2	1.2	2.4	1.9	2.1	2.4	3.0	4.9	6.1	10.9	17.0
2000[4]	7.5	*	1.0	0.8	1.4	1.7	2.2	1.9	1.9	3.0	5.0	7.8	10.4	17.9
1999[4]	7.5	*	1.0	0.8	1.6	1.5	2.6	1.9	2.4	3.2	4.6	8.4	12.3	20.0
Male														
2001	8.3	*	1.4	1.5	1.2	3.1	2.4	2.3	3.1	3.8	6.5	7.5	12.7	21.6
2000[4]	8.0	*	*	*	2.0	2.0	2.6	2.4	2.3	2.9	6.1	10.7	11.6	23.2
1999[4]	7.6	*	*	*	1.5	1.4	2.8	1.9	2.5	4.0	4.5	10.0	14.6	23.6
Female														
2001	6.5	*	*	*	1.3	1.7	1.3	1.8	1.6	2.2	3.2	4.8	9.1	12.8
2000[4]	6.9	*	*	*	*	1.4	1.6	1.3	1.4	3.1	3.8	4.8	9.3	13.1
1999[4]	7.4	*	*	*	1.6	1.5	2.2	1.9	2.2	2.4	4.7	6.8	10.2	16.7
Non-Hispanic[5]														
Both sexes														
2001	21.6	0.6	0.9	0.7	0.9	1.4	2.0	2.3	2.7	3.5	4.8	8.2	13.1	23.1
2000[4]	21.8	1.0	0.9	0.8	0.9	1.6	2.1	2.2	2.6	3.5	4.8	8.4	13.6	23.8
1999[4]	21.9	0.7	0.9	0.7	0.9	1.5	2.2	2.2	2.6	3.7	4.9	8.4	13.7	25.9
Male														
2001	23.5	*	1.0	0.8	1.1	1.6	2.3	2.6	3.2	4.2	5.8	10.0	16.2	28.1
2000[4]	23.7	*	1.0	1.0	1.0	1.9	2.6	2.3	3.2	4.0	5.7	10.2	16.4	28.7
1999[4]	23.7	*	0.9	0.7	1.1	1.8	2.7	2.5	3.1	4.4	6.0	10.2	16.9	31.4
Female														
2001	19.7	*	0.9	0.6	0.7	1.1	1.7	2.0	2.3	2.8	3.7	6.5	10.2	18.5
2000[4]	20.1	*	0.9	0.6	0.8	1.2	1.5	2.1	2.0	2.9	3.9	6.6	10.9	19.2
1999[4]	20.1	*	0.8	0.6	0.8	1.2	1.7	1.9	2.1	3.0	3.7	6.6	10.6	20.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)						
All origins[3]						
Male						
2000[4]	45.8	78.3	117.7	169.8	230.6	281.8
1999[4]	48.6	78.1	118.9	169.4	221.8	279.0
Female						
2001	29.5	44.6	71.8	103.4	137.1	168.4
2000[4]	31.3	48.4	71.0	104.7	143.8	173.1
1999[4]	31.0	49.8	74.3	103.8	145.9	167.3
Hispanic						
Both sexes						
2001	33.1	43.0	64.5	98.0	112.7	134.4
2000[4]	34.0	45.7	75.2	97.0	114.7	143.3
1999[4]	29.5	49.6	69.7	88.1	124.5	131.9
Male						
2001	38.8	56.3	81.9	139.8	146.7	180.9
2000[4]	40.6	58.2	88.5	114.4	151.7	197.5
1999[4]	36.9	56.0	85.0	106.9	139.9	135.9
Female						
2001	28.2	32.1	51.3	68.3	91.7	111.7
2000[4]	28.3	35.6	65.2	84.7	92.3	116.7
1999[4]	23.1	44.4	58.2	74.9	115.3	130.0
Non-Hispanic[5]						
Both sexes						
2001	38.3	59.7	93.2	130.3	170.5	202.1
2000[4]	38.4	63.2	92.3	132.7	178.1	206.5
1999[4]	40.0	63.6	95.1	132.2	175.3	201.4
Male						
2001	48.0	76.3	118.7	166.4	223.6	278.8
2000[4]	46.0	79.4	119.1	172.1	233.3	285.0
1999[4]	49.4	79.3	120.5	171.8	224.3	284.3
Female						
2001	29.5	45.4	72.9	105.0	138.7	170.2
2000[4]	31.4	49.2	71.2	105.3	145.6	174.8
1999[4]	31.4	50.1	75.2	104.7	146.7	168.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)														
Non-Hispanic White														
Both sexes														
2001	23.8	*	1.1	0.7	0.9	1.4	2.0	2.1	2.6	3.4	4.5	8.0	12.9	22.6
2000[4]	24.2	1.3	1.0	0.8	1.0	1.5	1.9	2.1	2.5	3.3	4.6	8.0	13.3	23.6
1999[4]	24.1	*	1.0	0.7	0.9	1.4	2.1	2.2	2.5	3.7	4.7	8.2	13.4	25.6
Male														
2001	25.9	*	1.0	0.7	1.0	1.6	2.3	2.3	3.2	3.9	5.5	9.9	16.0	27.4
2000[4]	26.2	*	1.1	1.0	1.1	1.8	2.5	2.2	3.0	3.9	5.7	9.8	15.9	28.4
1999[4]	26.1	*	1.0	0.8	1.0	1.7	2.8	2.4	2.9	4.3	5.7	9.9	16.6	31.0
Female														
2001	21.8	*	1.1	0.6	0.8	1.1	1.7	1.8	2.1	2.9	3.5	6.2	9.8	18.0
2000[4]	22.3	*	0.9	0.7	0.9	1.1	1.4	1.9	2.0	2.8	3.5	6.2	10.8	19.0
1999[4]	22.2	*	0.9	0.6	0.7	1.2	1.5	1.9	2.1	3.0	3.7	6.5	10.2	20.5
Non-Hispanic Black														
Both sexes														
2001	14.5	*	*	0.8	1.1	1.5	2.2	3.3	3.7	4.7	6.8	10.7	16.9	31.0
2000[4]	14.3	*	*	0.6	0.7	2.0	2.7	2.9	3.2	4.5	6.4	12.6	18.0	29.7
1999[4]	14.5	*	*	0.6	1.3	1.8	3.1	2.7	3.3	4.7	6.5	10.6	18.5	33.7
Male														
2001	15.6	*	*	*	1.7	1.8	2.2	3.9	4.1	6.8	8.4	12.1	21.4	38.5
2000[4]	15.2	*	*	*	*	2.4	3.3	2.7	4.6	4.8	6.7	15.0	22.4	37.6
1999[4]	15.6	*	*	*	1.7	2.1	3.2	3.5	4.3	5.5	8.5	13.4	22.8	42.2
Female														
2001	13.5	*	*	*	*	*	2.2	2.8	3.3	2.9	5.4	9.5	13.0	24.8
2000[4]	13.5	*	*	*	*	1.5	2.1	3.0	1.9	4.3	6.2	10.5	14.3	23.3
1999[4]	13.6	*	*	*	*	1.4	2.9	2.1	2.3	4.0	4.7	8.2	14.8	26.9
Hodgkin's disease (C81)														
All origins[3]														
Both sexes														
2001	0.5	*	*	*	*	0.1	0.3	0.4	0.4	0.4	0.4	0.4	0.5	0.4
2000[4]	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.3	0.3	0.4	0.5	0.5
1999[4]	0.5	*	*	*	*	0.2	0.3	0.4	0.3	0.4	0.3	0.4	0.5	0.7
Male														
2001	0.5	*	*	*	*	*	0.3	0.4	0.5	0.5	0.5	0.5	0.6	0.6
2000[4]	0.5	*	*	*	*	*	0.3	0.3	0.5	0.4	0.4	0.5	0.6	0.5
1999[4]	0.6	*	*	*	*	0.2	0.4	0.4	0.4	0.5	0.4	0.5	0.6	0.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)						
Non-Hispanic White						
Both sexes						
2001	38.2	60.0	95.2	133.3	174.9	208.2
2000[4]	38.6	64.5	94.1	135.7	183.7	210.7
1999[4]	39.8	64.6	96.9	134.7	179.9	206.4
Male						
2001	47.9	76.7	121.2	170.6	229.3	288.7
2000[4]	46.2	81.5	120.4	176.5	240.8	293.8
1999[4]	49.4	80.6	122.7	174.6	231.2	291.8
Female						
2001	29.3	45.2	74.0	106.8	142.1	174.6
2000[4]	31.5	49.5	73.0	107.0	149.8	177.2
1999[4]	30.9	50.5	76.3	106.6	149.7	172.4
Non-Hispanic Black						
Both sexes						
2001	44.9	68.4	92.2	117.1	144.0	157.3
2000[4]	44.8	63.5	91.5	115.8	132.8	177.1
1999[4]	46.5	65.8	91.8	120.8	140.0	162.3
Male						
2001	57.1	89.4	115.7	142.2	194.6	206.7
2000[4]	55.1	74.9	127.1	149.8	170.8	226.6
1999[4]	56.2	82.1	116.3	161.0	166.5	221.1
Female						
2001	35.3	52.9	76.6	102.1	118.3	139.0
2000[4]	36.7	55.0	67.8	95.2	113.6	158.7
1999[4]	38.9	53.7	75.6	96.7	126.9	140.4
Hodgkin's disease (C81)						
All origins[3]						
Both sexes						
2001	0.7	1.1	1.2	2.0	2.5	2.2
2000[4]	0.9	1.0	1.6	1.8	2.4	2.5
1999[4]	0.9	1.3	1.8	2.0	2.3	2.4
Male						
2001	0.9	1.5	1.6	2.5	2.9	2.6
2000[4]	1.1	1.3	1.9	2.3	2.7	3.7
1999[4]	1.2	1.6	2.3	2.3	2.8	3.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Hodgkin's disease (C81)														
All origins[3]														
Female														
2001	0.4	*	*	*	*	*	0.3	0.4	0.4	0.3	0.3	0.4	0.4	0.3
2000[4]	0.4	*	*	*	*	*	0.2	0.3	0.4	0.3	0.3	0.3	0.4	0.5
1999[4]	0.4	*	*	*	*	*	0.3	0.4	0.3	0.3	0.3	0.3	0.4	0.5
Hispanic														
Both sexes														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.5	*	*	*	*	0.1	0.3	0.4	0.5	0.4	0.4	0.5	0.5	0.4
2000[4]	0.5	*	*	*	*	*	0.3	0.3	0.5	0.4	0.3	0.4	0.5	0.5
1999[4]	0.5	*	*	*	*	0.2	0.4	0.5	0.4	0.4	0.3	0.4	0.5	0.6
Male														
2001	0.5	*	*	*	*	*	0.3	0.5	0.5	0.5	0.4	0.6	0.6	0.5
2000[4]	0.5	*	*	*	*	*	0.3	0.4	0.5	0.4	0.4	0.4	0.6	0.5
1999[4]	0.6	*	*	*	*	*	0.4	0.5	0.5	0.5	0.4	0.5	0.5	0.8
Female														
2001	0.4	*	*	*	*	*	0.3	0.4	0.4	0.3	0.3	0.4	0.4	0.3
2000[4]	0.4	*	*	*	*	*	0.3	0.3	0.4	0.3	0.3	0.3	0.4	0.5
1999[4]	0.5	*	*	*	*	*	0.3	0.4	0.3	0.3	0.2	0.3	0.4	0.5
Non-Hispanic White														
Both sexes														
2001	0.5	*	*	*	*	*	0.3	0.4	0.5	0.4	0.4	0.5	0.5	0.4
2000[4]	0.5	*	*	*	*	*	0.3	0.4	0.5	0.4	0.3	0.4	0.5	0.5
1999[4]	0.6	*	*	*	*	0.2	0.4	0.5	0.4	0.5	0.4	0.4	0.5	0.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hodgkin's disease (C81)						
All origins[3]						
Female						
2001	0.6	0.7	0.9	1.7	2.2	2.0
2000[4]	0.7	0.7	1.3	1.4	2.3	2.0
1999[4]	0.5	0.9	1.5	1.8	2.0	1.8
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	0.7	1.0	1.2	2.1	2.5	2.2
2000[4]	0.8	1.0	1.6	1.7	2.3	2.4
1999[4]	0.8	1.2	1.8	2.0	2.2	2.3
Male						
2001	0.9	1.4	1.6	2.5	2.9	2.5
2000[4]	1.0	1.3	1.9	2.1	2.6	3.7
1999[4]	1.2	1.6	2.2	2.2	2.5	3.7
Female						
2001	0.6	0.7	0.9	1.8	2.2	2.0
2000[4]	0.6	0.7	1.3	1.4	2.1	2.0
1999[4]	0.5	0.9	1.5	1.8	2.0	1.7
Non-Hispanic White						
Both sexes						
2001	0.8	1.2	1.2	2.1	2.6	2.3
2000[4]	0.8	1.1	1.6	1.9	2.4	2.6
1999[4]	0.9	1.2	1.9	2.1	2.3	2.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Hodgkin's disease (C81)														
Non-Hispanic White														
Male														
2001	0.6	*	*	*	*	*	0.4	0.4	0.5	0.5	0.5	0.6	0.6	0.5
2000[4]	0.6	*	*	*	*	*	0.3	0.4	0.6	0.4	0.4	0.5	0.6	0.5
1999[4]	0.7	*	*	*	*	*	0.4	0.5	0.5	0.6	0.5	0.5	0.5	0.9
Female														
2001	0.5	*	*	*	*	*	*	0.4	0.4	0.3	0.3	0.4	0.4	*
2000[4]	0.5	*	*	*	*	*	*	0.3	0.5	0.4	0.3	0.3	0.4	0.5
1999[4]	0.5	*	*	*	*	*	*	0.4	0.3	0.4	0.2	0.4	0.4	0.5
Non-Hispanic Black														
Both sexes														
2001	0.4	*	*	*	*	*	*	*	0.8	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hodgkin's lymphoma (C82-C85)														
All origins[3]														
Both sexes														
2001	7.8	*	*	*	0.1	0.3	0.4	0.7	0.8	1.4	2.0	3.4	5.5	9.6
2000[4]	8.1	*	*	0.1	0.1	0.3	0.6	0.6	0.8	1.3	2.1	3.6	5.8	10.6
1999[4]	8.2	*	*	*	0.1	0.3	0.5	0.6	0.9	1.4	2.1	3.5	6.2	10.8
Male														
2001	8.4	*	*	*	0.2	0.4	0.5	0.8	1.0	1.8	2.7	4.4	7.4	11.9
2000[4]	8.6	*	*	*	*	0.4	0.8	0.7	1.0	1.7	2.6	4.7	7.5	13.3
1999[4]	8.6	*	*	*	0.2	0.3	0.6	0.7	1.1	1.8	2.8	4.7	7.9	13.5
Female														
2001	7.3	*	*	*	*	*	0.4	0.6	0.6	1.0	1.3	2.5	3.8	7.5
2000[4]	7.6	*	*	*	*	*	0.4	0.6	0.6	0.9	1.5	2.6	4.1	8.2
1999[4]	7.7	*	*	*	*	0.2	0.4	0.4	0.7	1.0	1.4	2.4	4.5	8.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hodgkin's disease (C81)						
Non-Hispanic White						
Male						
2001	0.9	1.5	1.7	2.6	3.0	2.6
2000[4]	1.0	1.4	1.9	2.4	2.7	3.9
1999[4]	1.3	1.6	2.4	2.4	2.7	3.8
Female						
2001	0.7	0.8	0.9	1.8	2.4	2.3
2000[4]	0.7	0.8	1.4	1.6	2.2	2.1
1999[4]	0.5	0.8	1.5	2.0	2.1	1.7
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hodgkin's lymphoma (C82-C85)						
All origins[3]						
Both sexes						
2001	15.6	23.0	36.0	50.9	69.8	79.6
2000[4]	16.1	25.0	36.3	53.7	71.8	80.2
1999[4]	16.3	25.6	38.4	54.6	72.6	77.7
Male						
2001	19.1	29.2	44.7	61.3	87.3	106.6
2000[4]	19.2	30.5	45.2	66.3	88.1	105.9
1999[4]	19.6	30.8	46.8	68.1	85.7	104.4
Female						
2001	12.4	17.7	29.1	43.6	59.3	68.3
2000[4]	13.3	20.3	29.2	45.0	62.2	69.8
1999[4]	13.3	21.1	31.8	45.2	65.0	66.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Non-Hodgkin's lymphoma (C82-C85)														
Hispanic														
Both sexes														
2001	2.7	*	*	*	*	*	*	0.6	*	1.1	1.8	2.9	4.5	7.9
2000[4]	2.9	*	*	*	*	*	*	*	*	0.9	2.5	2.9	4.6	8.1
1999[4]	2.8	*	*	*	*	*	*	*	0.9	0.9	2.0	3.8	5.2	6.8
Male														
2001	3.0	*	*	*	*	*	*	*	*	1.4	2.7	3.6	5.4	10.0
2000[4]	3.1	*	*	*	*	*	*	*	*	*	3.0	3.8	5.7	10.5
1999[4]	2.8	*	*	*	*	*	*	*	*	*	2.1	5.0	6.7	8.8
Female														
2001	2.4	*	*	*	*	*	*	*	*	*	*	2.1	3.7	6.0
2000[4]	2.7	*	*	*	*	*	*	*	*	*	2.0	*	3.4	6.0
1999[4]	2.8	*	*	*	*	*	*	*	*	*	*	2.6	3.8	5.0
Non-Hispanic[5]														
Both sexes														
2001	8.6	*	*	*	0.1	0.3	0.5	0.7	0.9	1.4	2.0	3.5	5.6	9.8
2000[4]	8.8	*	*	0.1	0.1	0.3	0.7	0.7	0.9	1.3	2.0	3.7	5.8	10.8
1999[4]	8.9	*	*	*	0.1	0.3	0.5	0.6	0.9	1.5	2.1	3.5	6.2	11.1
Male														
2001	9.2	*	*	*	*	0.4	0.5	0.8	1.1	1.8	2.7	4.5	7.5	12.1
2000[4]	9.4	*	*	*	*	0.5	0.9	0.7	1.1	1.8	2.6	4.8	7.6	13.4
1999[4]	9.4	*	*	*	0.2	0.3	0.7	0.8	1.1	1.9	2.8	4.7	8.0	13.8
Female														
2001	8.0	*	*	*	*	*	0.4	0.6	0.7	1.0	1.4	2.5	3.8	7.6
2000[4]	8.3	*	*	*	*	*	0.5	0.6	0.6	0.9	1.5	2.6	4.2	8.3
1999[4]	8.4	*	*	*	*	0.2	0.3	0.4	0.6	1.1	1.4	2.3	4.6	8.6
Non-Hispanic White														
Both sexes														
2001	9.7	*	*	*	*	0.3	0.4	0.6	0.8	1.3	2.0	3.4	5.7	9.9
2000[4]	10.1	*	*	*	0.2	0.3	0.6	0.6	0.8	1.3	2.0	3.7	6.0	11.4
1999[4]	10.1	*	*	*	*	0.2	0.5	0.6	0.8	1.4	2.0	3.5	6.5	11.5
Male														
2001	10.4	*	*	*	*	0.4	0.5	0.7	1.1	1.7	2.6	4.5	7.6	12.1
2000[4]	10.6	*	*	*	*	0.5	0.8	0.6	1.0	1.7	2.7	4.7	7.6	13.9
1999[4]	10.6	*	*	*	*	*	0.7	0.7	1.0	1.8	2.7	4.7	8.2	14.1
Female														
2001	9.2	*	*	*	*	*	0.4	0.5	0.6	1.0	1.3	2.4	3.9	7.8
2000[4]	9.5	*	*	*	*	*	0.4	0.5	0.6	0.8	1.4	2.6	4.4	9.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)						
Hispanic						
Both sexes						
2001	14.9	14.4	23.2	41.6	46.2	59.4
2000[4]	14.9	18.5	34.8	39.2	45.7	59.7
1999[4]	12.5	20.2	31.5	37.8	51.7	56.1
Male						
2001	17.7	19.5	30.7	55.9	63.6	68.6
2000[4]	18.4	22.4	40.4	44.3	55.9	82.6
1999[4]	15.8	21.6	35.1	46.0	51.9	60.4
Female						
2001	12.5	10.3	17.4	31.5	35.4	54.9
2000[4]	11.9	15.4	30.6	35.6	39.4	48.5
1999[4]	9.8	19.1	28.7	31.9	51.6	54.1
Non-Hispanic[5]						
Both sexes						
2001	15.6	23.5	36.7	51.2	70.6	80.3
2000[4]	16.1	25.4	36.3	54.3	72.7	80.9
1999[4]	16.5	25.9	38.7	55.2	73.2	78.3
Male						
2001	19.1	29.7	45.5	61.3	88.2	108.1
2000[4]	19.1	31.0	45.3	67.2	89.3	106.7
1999[4]	19.9	31.3	47.3	68.9	86.5	106.0
Female						
2001	12.4	18.1	29.8	44.1	60.1	68.7
2000[4]	13.4	20.7	29.1	45.3	62.9	70.4
1999[4]	13.5	21.2	32.0	45.6	65.3	67.2
Non-Hispanic White						
Both sexes						
2001	16.3	24.5	38.7	54.0	74.0	84.1
2000[4]	16.9	26.8	38.1	57.1	76.9	85.1
1999[4]	17.4	27.5	40.6	57.6	77.3	82.0
Male						
2001	19.8	30.9	47.5	64.3	92.2	113.1
2000[4]	19.9	32.8	46.9	70.8	94.1	113.0
1999[4]	20.6	32.9	49.5	71.1	91.8	110.0
Female						
2001	13.1	18.8	31.6	46.7	63.0	72.0
2000[4]	14.1	21.6	31.1	47.6	66.6	73.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Non-Hodgkin's lymphoma (C82-C85)														
Non-Hispanic White														
Female														
1999[4]	9.6	*	*	*	*	*	*	0.4	0.6	1.0	1.4	2.3	4.8	9.0
Non-Hispanic Black														
Both sexes														
2001	4.1	*	*	*	*	*	*	1.2	1.2	2.1	2.5	4.3	6.1	9.9
2000[4]	3.9	*	*	*	*	*	1.0	1.2	1.0	1.9	2.4	4.5	5.6	9.1
1999[4]	4.1	*	*	*	*	*	*	0.9	1.3	2.2	2.7	4.1	5.6	10.8
Male														
2001	4.7	*	*	*	*	*	*	*	*	3.2	3.5	5.1	8.8	12.4
2000[4]	4.3	*	*	*	*	*	*	*	*	2.6	2.6	6.2	8.0	13.9
1999[4]	4.7	*	*	*	*	*	*	*	1.8	2.7	4.0	5.6	7.9	14.1
Female														
2001	3.7	*	*	*	*	*	*	*	*	*	1.6	3.7	3.7	7.9
2000[4]	3.5	*	*	*	*	*	*	*	*	*	2.1	3.1	3.6	5.1
1999[4]	3.6	*	*	*	*	*	*	*	*	1.7	1.6	2.9	3.8	8.1
Leukemia (C91-C95)														
All origins[3]														
Both sexes														
2001	7.5	0.6	1.0	0.8	0.8	1.1	1.3	1.2	1.4	1.6	1.9	3.0	4.6	7.6
2000[4]	7.6	1.0	0.9	0.7	0.8	1.2	1.2	1.2	1.2	1.6	1.9	3.1	4.5	7.7
1999[4]	7.5	0.7	0.8	0.6	0.9	1.1	1.4	1.2	1.3	1.6	1.9	3.2	4.4	8.5
Male														
2001	8.5	*	1.0	0.8	0.9	1.3	1.5	1.4	1.6	1.8	2.1	3.4	5.2	9.2
2000[4]	8.5	1.1	0.9	0.8	1.0	1.4	1.5	1.3	1.5	1.6	2.1	3.6	5.1	9.7
1999[4]	8.4	*	0.8	0.7	0.9	1.3	1.7	1.3	1.4	1.8	2.1	3.6	5.2	10.1
Female														
2001	6.6	*	1.0	0.7	0.7	0.9	1.0	1.0	1.2	1.3	1.7	2.7	4.0	6.0
2000[4]	6.7	*	0.9	0.6	0.6	0.9	0.9	1.0	0.9	1.6	1.7	2.6	3.9	5.9
1999[4]	6.7	*	0.7	0.5	0.8	0.9	1.1	1.1	1.2	1.5	1.7	2.8	3.6	7.0
Hispanic														
Both sexes														
2001	3.0	*	1.3	1.2	1.1	2.0	1.5	1.3	1.6	1.5	2.3	2.6	3.5	4.5
2000[4]	2.9	*	0.9	0.8	1.4	1.5	1.5	1.1	1.1	1.8	2.0	3.3	3.8	6.4
1999[4]	2.9	*	0.9	0.8	1.4	1.2	1.9	1.5	1.2	1.9	2.1	3.0	3.9	6.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)						
Non-Hispanic White						
Female						
1999[4]	14.4	22.7	33.4	48.2	68.8	70.8
Non-Hispanic Black						
Both sexes						
2001	12.1	19.3	24.0	25.9	38.1	40.0
2000[4]	12.8	17.0	22.9	26.3	28.5	36.6
1999[4]	11.6	17.3	24.4	32.5	31.4	40.4
Male						
2001	15.5	25.0	32.4	30.8	48.2	55.4
2000[4]	15.9	19.7	32.5	31.3	32.6	43.4
1999[4]	15.7	23.9	29.1	47.9	28.3	59.4
Female						
2001	9.5	15.0	18.4	22.9	32.9	34.2
2000[4]	10.4	14.9	16.6	23.2	26.4	34.0
1999[4]	8.4	12.4	21.4	23.3	32.9	33.3
Leukemia (C91-C95)						
All origins[3]						
Both sexes						
2001	13.1	20.6	33.3	47.7	61.3	81.4
2000[4]	12.7	21.7	33.2	47.4	65.1	83.6
1999[4]	13.1	21.8	33.1	46.3	62.7	82.4
Male						
2001	17.4	27.6	44.9	66.2	86.9	117.5
2000[4]	16.0	29.2	46.2	65.1	93.4	119.8
1999[4]	16.8	29.0	45.7	64.6	87.6	119.2
Female						
2001	9.3	14.6	24.1	34.8	45.9	66.3
2000[4]	9.7	15.3	22.9	35.1	48.4	68.8
1999[4]	9.8	15.6	23.2	33.6	48.1	67.6
Hispanic						
Both sexes						
2001	9.2	14.3	21.2	32.0	35.8	42.5
2000[4]	9.6	13.5	21.2	29.1	39.0	49.1
1999[4]	7.3	14.9	20.3	27.4	37.6	44.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Leukemia (C91-C95)														
Hispanic														
Male														
2001	3.4	*	1.4	1.3	*	2.5	2.0	1.6	2.0	2.0	2.7	3.1	3.6	5.2
2000[4]	3.3	*	*	*	2.0	1.7	1.6	1.6	1.3	1.6	2.4	4.3	3.9	9.0
1999[4]	3.1	*	*	*	1.3	*	2.3	1.5	1.4	2.0	1.8	3.6	3.5	7.2
Female														
2001	2.6	*	*	*	1.3	1.3	*	*	*	*	1.9	*	3.3	3.8
2000[4]	2.6	*	*	*	*	*	1.4	*	*	2.1	*	2.3	3.7	4.0
1999[4]	2.8	*	*	*	1.5	1.5	1.6	1.5	*	1.8	2.4	2.4	4.3	6.5
Non-Hispanic[5]														
Both sexes														
2001	8.2	*	0.9	0.7	0.8	1.0	1.2	1.2	1.3	1.5	1.8	3.1	4.7	7.8
2000[4]	8.2	1.0	0.9	0.7	0.7	1.1	1.1	1.2	1.2	1.6	1.9	3.1	4.5	7.8
1999[4]	8.2	*	0.7	0.6	0.8	1.1	1.3	1.1	1.3	1.6	1.9	3.2	4.4	8.6
Male														
2001	9.3	*	0.9	0.7	0.9	1.1	1.4	1.4	1.5	1.7	2.0	3.4	5.3	9.5
2000[4]	9.3	*	0.9	0.8	0.9	1.4	1.4	1.3	1.5	1.6	2.1	3.5	5.1	9.6
1999[4]	9.2	*	0.8	0.6	0.9	1.3	1.6	1.3	1.4	1.8	2.1	3.6	5.3	10.2
Female														
2001	7.2	*	0.9	0.6	0.6	0.9	1.0	1.0	1.1	1.4	1.7	2.8	4.1	6.2
2000[4]	7.2	*	0.9	0.5	0.6	0.9	0.7	1.1	0.9	1.5	1.7	2.6	4.0	6.0
1999[4]	7.2	*	0.7	0.6	0.7	0.8	1.0	1.0	1.2	1.4	1.7	2.8	3.5	7.1
Non-Hispanic White														
Both sexes														
2001	9.1	*	1.0	0.6	0.7	1.0	1.2	1.0	1.3	1.5	1.8	3.1	4.6	7.9
2000[4]	9.2	1.3	0.9	0.7	0.8	1.0	1.1	1.1	1.1	1.5	1.8	2.9	4.5	7.8
1999[4]	9.1	*	0.8	0.6	0.7	1.0	1.2	1.2	1.2	1.6	1.9	3.2	4.3	8.8
Male														
2001	10.3	*	1.0	0.6	0.8	1.1	1.4	1.2	1.5	1.6	1.9	3.5	5.2	9.7
2000[4]	10.4	*	1.0	0.8	0.9	1.2	1.4	1.2	1.3	1.6	2.1	3.4	5.0	9.7
1999[4]	10.2	*	0.8	0.7	0.8	1.2	1.6	1.3	1.3	1.8	2.1	3.5	5.2	10.5
Female														
2001	8.0	*	1.1	0.6	0.7	0.8	1.0	0.9	1.1	1.4	1.7	2.7	3.9	6.2
2000[4]	8.1	*	0.9	0.6	0.7	0.8	0.7	1.0	0.8	1.5	1.6	2.5	4.0	6.0
1999[4]	8.0	*	0.9	0.5	0.6	0.8	0.9	1.0	1.2	1.4	1.7	2.9	3.4	7.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Leukemia (C91-C95)						
Hispanic						
Male						
2001	12.8	16.3	23.8	49.1	39.0	68.6
2000[4]	11.8	16.8	25.8	35.4	57.4	76.6
1999[4]	10.4	17.4	31.6	35.1	47.2	49.6
Female						
2001	6.1	12.6	19.2	19.8	33.8	29.8
2000[4]	7.7	10.9	17.7	24.6	27.8	35.6
1999[4]	*	12.8	11.9	22.0	31.9	41.6
Non-Hispanic[5]						
Both sexes						
2001	13.4	21.0	33.9	48.4	62.2	82.7
2000[4]	12.9	22.2	33.8	48.1	66.0	84.7
1999[4]	13.5	22.2	33.7	47.0	63.5	83.7
Male						
2001	17.7	28.3	46.0	66.9	88.7	119.3
2000[4]	16.3	30.0	47.3	66.4	94.5	121.5
1999[4]	17.2	29.6	46.3	65.8	89.0	122.1
Female						
2001	9.5	14.7	24.3	35.5	46.4	67.5
2000[4]	9.8	15.5	23.1	35.4	49.2	69.8
1999[4]	10.1	15.8	23.8	34.0	48.5	68.3
Non-Hispanic White						
Both sexes						
2001	13.7	21.4	35.2	49.8	64.8	86.1
2000[4]	13.1	23.2	35.1	49.9	68.5	87.3
1999[4]	13.7	22.6	34.9	48.8	65.0	86.8
Male						
2001	18.1	28.6	47.8	69.0	92.0	125.3
2000[4]	16.5	31.3	48.6	68.9	97.9	125.5
1999[4]	17.5	30.2	47.9	68.0	91.7	127.2
Female						
2001	9.7	15.0	25.0	36.2	48.4	69.8
2000[4]	9.9	16.0	24.2	36.6	51.1	72.0
1999[4]	10.1	15.9	24.6	35.3	49.2	70.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Leukemia (C91-C95)														
Non-Hispanic Black														
Both sexes														
2001	4.9	*	*	0.8	0.9	0.9	1.2	1.5	1.4	1.9	2.3	3.1	5.7	8.1
2000[4]	4.9	*	*	*	0.6	1.5	1.3	1.5	1.9	1.7	2.5	4.7	5.3	8.1
1999[4]	4.9	*	*	*	1.1	1.2	1.8	1.1	1.4	1.8	1.9	3.6	5.5	8.3
Male														
2001	5.4	*	*	*	1.5	*	*	1.8	1.8	2.4	2.7	2.9	6.8	10.1
2000[4]	5.5	*	*	*	*	2.1	1.8	*	2.6	1.5	2.5	5.1	6.3	10.2
1999[4]	5.4	*	*	*	1.5	1.5	1.7	*	1.8	1.9	2.3	4.6	7.1	10.6
Female														
2001	4.4	*	*	*	*	*	*	*	*	1.4	2.0	3.4	4.8	6.5
2000[4]	4.4	*	*	*	*	*	*	1.6	*	1.9	2.4	4.3	4.5	6.4
1999[4]	4.4	*	*	*	*	*	1.9	*	*	1.8	1.6	2.7	4.2	6.4
Multiple myeloma and immunoproliferative neoplasms (C88,C90)														
All origins[3]														
Both sexes														
2001	3.9	*	*	*	*	*	*	*	*	0.2	0.5	1.2	2.3	5.1
2000[4]	3.9	*	*	*	*	*	*	*	*	0.2	0.5	1.2	2.6	4.5
1999[4]	3.9	*	*	*	*	*	*	*	*	0.2	0.5	1.2	2.5	5.6
Male														
2001	4.1	*	*	*	*	*	*	*	*	0.2	0.7	1.5	2.8	6.0
2000[4]	4.0	*	*	*	*	*	*	*	*	*	0.6	1.4	2.9	4.9
1999[4]	4.1	*	*	*	*	*	*	*	*	0.3	0.6	1.4	3.0	6.5
Female														
2001	3.8	*	*	*	*	*	*	*	*	*	0.4	0.9	1.9	4.2
2000[4]	3.8	*	*	*	*	*	*	*	*	0.2	0.4	1.0	2.3	4.2
1999[4]	3.8	*	*	*	*	*	*	*	*	*	0.4	1.1	2.1	4.7
Hispanic														
Both sexes														
2001	1.4	*	*	*	*	*	*	*	*	*	*	*	2.2	3.9
2000[4]	1.4	*	*	*	*	*	*	*	*	*	*	1.2	1.5	3.2
1999[4]	1.4	*	*	*	*	*	*	*	*	*	*	1.2	2.3	5.6
Male														
2001	1.5	*	*	*	*	*	*	*	*	*	*	*	*	4.8
2000[4]	1.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.3	*	*	*	*	*	*	*	*	*	*	*	*	6.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Leukemia (C91-C95)						
Non-Hispanic Black						
Both sexes						
2001	12.5	22.4	28.4	41.8	42.5	57.1
2000[4]	14.3	18.1	29.0	35.3	44.3	66.1
1999[4]	14.0	22.6	28.3	35.6	51.4	55.9
Male						
2001	17.6	32.6	36.8	52.6	62.1	75.1
2000[4]	18.9	22.9	44.4	50.6	65.2	98.6
1999[4]	17.6	29.5	38.4	52.2	62.0	78.4
Female						
2001	8.5	14.8	22.7	35.4	32.5	50.5
2000[4]	10.6	14.5	18.8	26.1	33.7	54.1
1999[4]	11.1	17.5	21.6	25.6	46.1	47.5
Multiple myeloma and immunoproliferative neoplasms (C88,C90)						
All origins[3]						
Both sexes						
2001	8.5	14.0	21.1	28.3	34.7	36.3
2000[4]	8.5	14.4	20.4	28.3	36.4	37.8
1999[4]	9.0	14.1	20.6	27.6	36.2	36.6
Male						
2001	10.1	16.8	25.3	35.6	43.4	48.0
2000[4]	9.5	17.1	24.2	35.7	46.1	51.8
1999[4]	11.0	16.6	24.1	34.1	45.5	51.3
Female						
2001	7.1	11.5	17.7	23.2	29.4	31.3
2000[4]	7.6	12.1	17.3	23.1	30.7	32.2
1999[4]	7.3	12.0	17.8	23.0	30.7	30.6
Hispanic						
Both sexes						
2001	8.3	12.2	18.2	22.7	28.3	29.4
2000[4]	7.9	13.0	17.4	25.1	23.4	31.2
1999[4]	8.3	12.1	15.1	19.4	30.0	28.1
Male						
2001	7.6	17.4	24.2	32.7	41.6	38.1
2000[4]	8.1	17.5	19.4	28.1	33.9	*
1999[4]	9.2	14.3	15.3	21.1	31.4	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Multiple myeloma and immunoproliferative neoplasms (C88,C90)														
Hispanic														
Female														
2001	1.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.5	*	*	*	*	*	*	*	*	*	*	*	*	4.6
Non-Hispanic[5]														
Both sexes														
2001	4.3	*	*	*	*	*	*	*	*	0.2	0.5	1.2	2.3	5.2
2000[4]	4.3	*	*	*	*	*	*	*	*	0.2	0.5	1.2	2.7	4.6
1999[4]	4.3	*	*	*	*	*	*	*	*	0.2	0.5	1.2	2.5	5.5
Male														
2001	4.5	*	*	*	*	*	*	*	*	*	0.6	1.6	2.8	6.1
2000[4]	4.4	*	*	*	*	*	*	*	*	*	0.6	1.4	3.0	5.0
1999[4]	4.5	*	*	*	*	*	*	*	*	0.3	0.6	1.4	3.0	6.5
Female														
2001	4.1	*	*	*	*	*	*	*	*	*	0.4	0.9	1.9	4.3
2000[4]	4.2	*	*	*	*	*	*	*	*	*	0.4	1.0	2.3	4.2
1999[4]	4.1	*	*	*	*	*	*	*	*	*	0.4	1.1	2.1	4.7
Non-Hispanic White														
Both sexes														
2001	4.4	*	*	*	*	*	*	*	*	0.2	0.3	1.0	2.0	4.4
2000[4]	4.3	*	*	*	*	*	*	*	*	0.1	0.4	1.0	2.3	3.9
1999[4]	4.3	*	*	*	*	*	*	*	*	0.2	0.4	1.1	2.1	4.6
Male														
2001	4.6	*	*	*	*	*	*	*	*	*	0.5	1.4	2.5	5.2
2000[4]	4.6	*	*	*	*	*	*	*	*	*	0.6	1.2	2.6	4.3
1999[4]	4.6	*	*	*	*	*	*	*	*	0.2	0.4	1.3	2.6	5.5
Female														
2001	4.1	*	*	*	*	*	*	*	*	*	*	0.7	1.6	3.7
2000[4]	4.1	*	*	*	*	*	*	*	*	*	*	0.8	2.0	3.5
1999[4]	4.1	*	*	*	*	*	*	*	*	*	0.3	0.9	1.5	3.8
Non-Hispanic Black														
Both sexes														
2001	5.0	*	*	*	*	*	*	*	*	*	1.6	2.7	4.8	12.1
2000[4]	5.2	*	*	*	*	*	*	*	*	*	1.1	3.0	6.5	11.9
1999[4]	5.1	*	*	*	*	*	*	*	*	*	1.5	2.5	6.8	13.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Multiple myeloma and immunoproliferative neoplasms (C88,C90)						
Hispanic						
Female						
2001	8.9	8.0	13.6	15.5	20.1	25.1
2000[4]	7.7	9.4	15.8	23.0	*	30.7
1999[4]	7.4	10.3	14.9	18.2	29.1	31.2
Non-Hispanic[5]						
Both sexes						
2001	8.5	14.1	21.2	28.5	34.8	36.5
2000[4]	8.5	14.5	20.5	28.4	36.8	38.0
1999[4]	9.1	14.2	20.8	27.9	36.3	36.8
Male						
2001	10.2	16.8	25.3	35.6	43.4	48.4
2000[4]	9.5	17.0	24.5	36.0	46.5	52.5
1999[4]	11.1	16.7	24.6	34.6	46.0	52.2
Female						
2001	6.9	11.8	17.9	23.5	29.7	31.5
2000[4]	7.6	12.3	17.4	23.1	31.1	32.1
1999[4]	7.3	12.1	17.9	23.2	30.7	30.6
Non-Hispanic White						
Both sexes						
2001	7.4	12.9	19.9	27.2	33.1	35.1
2000[4]	7.7	13.3	19.2	26.5	35.6	35.2
1999[4]	7.8	13.2	19.4	26.0	35.1	35.0
Male						
2001	9.1	15.5	23.9	34.6	41.6	47.3
2000[4]	8.8	15.8	22.9	34.1	45.7	50.9
1999[4]	10.0	15.8	22.9	32.9	44.7	50.6
Female						
2001	5.8	10.6	16.6	22.0	28.0	30.0
2000[4]	6.7	11.1	16.2	21.2	29.7	28.9
1999[4]	5.8	10.9	16.6	21.1	29.4	28.7
Non-Hispanic Black						
Both sexes						
2001	19.3	26.2	38.8	47.7	61.8	59.3
2000[4]	16.9	27.9	38.0	53.8	58.3	72.8
1999[4]	20.5	24.6	38.1	51.6	56.9	63.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Multiple myeloma and immunoproliferative neoplasms (C88,C90)														
Non-Hispanic Black														
Male														
2001	5.0	*	*	*	*	*	*	*	*	*	1.6	3.5	5.4	15.0
2000[4]	5.0	*	*	*	*	*	*	*	*	*	*	3.1	7.4	12.7
1999[4]	5.0	*	*	*	*	*	*	*	*	*	1.8	2.6	6.7	16.4
Female														
2001	5.0	*	*	*	*	*	*	*	*	*	1.6	2.0	4.4	9.8
2000[4]	5.4	*	*	*	*	*	*	*	*	*	*	2.9	5.7	11.2
1999[4]	5.3	*	*	*	*	*	*	*	*	*	*	2.3	6.8	11.8
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)														
All origins[3]														
Both sexes														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Multiple myeloma and immunoproliferative neoplasms (C88,C90)						
Non-Hispanic Black						
Male						
2001	23.0	31.0	45.4	56.9	81.8	72.8
2000[4]	18.7	31.7	48.5	67.4	70.4	81.0
1999[4]	22.2	26.6	47.4	59.5	75.3	80.8
Female						
2001	16.4	22.7	34.4	42.2	51.6	54.3
2000[4]	15.4	25.2	31.1	45.6	52.3	69.8
1999[4]	19.1	23.2	31.9	46.9	47.9	56.9
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)						
All origins[3]						
Both sexes						
2001	*	*	*	*	*	0.5
2000[4]	*	*	*	*	*	0.5
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)														
Hispanic														
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)						
Hispanic						
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	*	*	0.5
2000[4]	*	*	*	*	*	0.5
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic White						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)														
Non-Hispanic Black														
Both sexes														
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31, C37-C41,C44-C49,C51-C52, C57-C60,C62-C63,C66,C68-C69, C73-C80,C97)														
All origins[3]														
Both sexes														
2001	22.0	*	0.8	0.6	0.8	1.2	1.4	1.7	2.0	3.2	5.9	10.5	18.0	30.1
2000[4]	22.4	0.6	0.8	0.6	0.7	1.2	1.4	1.5	2.0	3.4	5.9	10.5	18.2	30.4
1999[4]	23.7	*	0.8	0.7	0.7	1.2	1.5	1.6	2.2	3.6	6.7	11.4	19.1	32.5
Male														
2001	22.6	*	0.9	0.6	0.8	1.5	1.9	2.0	2.4	3.5	6.4	12.0	21.5	34.9
2000[4]	23.0	*	1.0	0.6	0.7	1.4	1.8	1.9	2.4	3.7	6.4	12.0	20.6	35.1
1999[4]	24.5	*	0.8	0.8	0.8	1.5	1.8	2.1	2.8	4.1	7.6	13.2	22.0	38.2
Female														
2001	21.4	*	0.6	0.5	0.7	0.9	0.8	1.4	1.6	3.0	5.3	9.1	14.7	25.7
2000[4]	21.9	*	0.7	0.6	0.6	1.0	1.0	1.1	1.6	3.0	5.3	9.1	15.8	25.9
1999[4]	23.1	*	0.7	0.7	0.6	0.9	1.1	1.1	1.6	3.2	5.8	9.7	16.3	27.1
Hispanic														
Both sexes														
2001	7.7	*	*	*	0.9	1.3	1.2	1.0	1.4	2.8	4.3	7.4	12.9	19.0
2000[4]	7.8	*	0.8	*	0.9	1.4	1.1	1.4	1.7	2.1	3.6	6.6	12.6	23.5
1999[4]	8.1	*	*	0.7	0.7	1.3	1.4	1.5	1.7	2.5	4.9	8.6	13.2	23.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)						
Non-Hispanic Black						
Both sexes						
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31, C37-C41,C44-C49,C51-C52, C57-C60,C62-C63,C66,C68-C69, C73-C80,C97)						
All origins[3]						
Both sexes						
2001	46.4	67.2	98.1	133.4	170.9	234.5
2000[4]	48.2	71.5	100.4	136.0	177.4	239.8
1999[4]	52.5	76.1	107.8	142.3	188.5	250.4
Male						
2001	55.4	79.4	120.3	162.3	211.0	295.2
2000[4]	57.2	86.2	122.5	167.5	221.4	300.7
1999[4]	62.6	90.6	132.8	174.0	233.0	318.9
Female						
2001	38.3	56.7	80.5	113.1	146.8	209.1
2000[4]	40.0	59.0	83.1	114.1	151.4	215.0
1999[4]	43.3	63.8	88.2	120.3	162.5	222.8
Hispanic						
Both sexes						
2001	35.4	46.8	78.2	98.3	133.1	193.8
2000[4]	30.5	52.9	78.2	109.3	152.6	192.4
1999[4]	35.2	56.2	75.6	108.5	147.4	202.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31, C37-C41,C44-C49,C51-C52, C57-C60,C62-C63,C66,C68-C69, C73-C80,C97)														
Hispanic														
Male														
2001	7.7	*	*	*	*	1.5	2.0	1.1	2.0	2.7	4.2	8.1	14.3	23.0
2000[4]	7.7	*	*	*	*	1.5	1.3	1.6	2.3	2.2	3.8	6.8	13.2	24.8
1999[4]	8.0	*	*	*	*	1.7	2.0	1.9	1.8	2.5	4.8	9.0	13.9	26.6
Female														
2001	7.7	*	*	*	*	*	*	*	*	2.8	4.4	6.8	11.7	15.4
2000[4]	7.9	*	*	*	*	*	*	*	*	1.9	3.4	6.5	12.1	22.4
1999[4]	8.2	*	*	*	*	*	*	*	1.5	2.5	4.9	8.2	12.6	21.1
Non-Hispanic[5]														
Both sexes														
2001	24.1	*	0.8	0.6	0.8	1.2	1.4	1.8	2.1	3.3	6.0	10.8	18.4	30.9
2000[4]	24.4	0.7	0.8	0.6	0.6	1.2	1.5	1.5	2.0	3.6	6.1	10.9	18.5	30.8
1999[4]	25.8	*	0.8	0.7	0.7	1.2	1.5	1.6	2.3	3.8	6.8	11.7	19.5	33.0
Male														
2001	24.9	*	1.0	0.7	0.8	1.4	1.9	2.2	2.5	3.6	6.6	12.4	22.0	35.7
2000[4]	25.2	*	0.9	0.6	0.7	1.3	2.0	2.0	2.4	4.0	6.7	12.5	21.2	35.8
1999[4]	26.8	*	1.0	0.8	0.8	1.5	1.7	2.1	2.9	4.3	7.9	13.5	22.7	39.0
Female														
2001	23.3	*	0.6	0.6	0.7	0.9	0.9	1.4	1.7	3.0	5.4	9.3	15.0	26.4
2000[4]	23.7	*	0.8	0.6	0.6	1.0	1.0	1.1	1.7	3.2	5.5	9.3	16.0	26.1
1999[4]	25.0	*	0.7	0.7	0.6	0.9	1.2	1.1	1.6	3.3	5.8	9.9	16.5	27.5
Non-Hispanic White														
Both sexes														
2001	26.0	*	0.9	0.6	0.7	1.2	1.5	1.9	2.0	3.2	5.8	10.1	17.5	29.7
2000[4]	26.5	*	0.8	0.7	0.6	1.1	1.6	1.7	2.0	3.4	5.6	10.2	17.5	30.1
1999[4]	27.9	*	0.9	0.7	0.6	1.2	1.4	1.7	2.2	3.6	6.4	11.0	18.6	31.6
Male														
2001	26.8	*	1.1	0.7	0.8	1.4	2.0	2.3	2.4	3.5	6.4	11.6	20.5	33.8
2000[4]	27.2	*	1.0	0.7	0.7	1.3	2.0	2.2	2.5	3.8	6.0	11.5	20.0	34.7
1999[4]	28.8	*	1.1	0.8	0.7	1.4	1.6	2.3	2.8	4.2	7.4	12.4	21.3	37.2
Female														
2001	25.3	*	0.7	0.6	0.7	0.9	1.0	1.5	1.6	2.8	5.1	8.6	14.4	25.8
2000[4]	25.8	*	0.6	0.7	0.6	0.9	1.1	1.1	1.6	2.9	5.2	8.8	14.9	25.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31, C37-C41,C44-C49,C51-C52, C57-C60,C62-C63,C66,C68-C69, C73-C80,C97)						
Hispanic						
Male						
2001	39.9	50.0	90.6	130.3	161.0	213.3
2000[4]	35.1	60.8	90.4	138.0	179.6	231.8
1999[4]	42.0	62.2	96.7	124.0	187.1	235.1
Female						
2001	31.5	44.2	68.8	75.5	115.8	184.3
2000[4]	26.6	46.5	68.9	88.9	136.2	173.1
1999[4]	29.3	51.3	59.8	97.5	123.8	186.1
Non-Hispanic[5]						
Both sexes						
2001	47.2	68.5	98.8	134.8	172.0	235.3
2000[4]	49.3	72.6	101.4	136.7	177.9	241.0
1999[4]	53.6	77.3	109.2	143.5	189.5	251.6
Male						
2001	56.3	81.2	121.3	163.7	212.5	297.9
2000[4]	58.6	87.5	123.8	168.2	222.5	303.0
1999[4]	63.9	92.2	134.2	175.7	233.9	321.9
Female						
2001	38.8	57.5	81.0	114.6	147.7	209.3
2000[4]	40.9	59.7	83.6	114.9	151.7	215.9
1999[4]	44.2	64.5	89.6	121.1	163.5	223.5
Non-Hispanic White						
Both sexes						
2001	45.9	67.7	98.1	135.5	172.5	238.9
2000[4]	48.3	72.4	100.8	136.2	179.9	244.9
1999[4]	52.3	77.5	108.5	143.3	190.3	253.6
Male						
2001	54.3	79.8	120.8	164.4	213.7	302.9
2000[4]	56.1	87.6	123.1	166.3	226.2	309.5
1999[4]	62.1	91.7	133.2	176.2	236.1	326.9
Female						
2001	38.1	57.1	79.6	115.1	147.6	212.3
2000[4]	41.0	59.1	82.9	115.1	152.4	218.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31, C37-C41,C44-C49,C51-C52, C57-C60,C62-C63,C66,C68-C69, C73-C80,C97)														
Non-Hispanic White														
Female														
1999[4]	27.0	*	0.8	0.6	0.5	1.0	1.2	1.0	1.6	3.1	5.3	9.5	16.0	26.2
Non-Hispanic Black														
Both sexes														
2001	18.9	*	*	*	0.8	1.2	1.4	2.1	2.7	4.3	7.8	17.2	28.4	45.2
2000[4]	18.9	*	0.9	*	0.7	1.6	1.4	1.3	2.0	5.4	10.1	17.1	29.5	42.1
1999[4]	20.2	*	*	0.7	0.9	1.3	1.7	1.6	2.5	5.1	10.4	18.7	29.8	49.2
Male														
2001	20.0	*	*	*	*	1.5	2.1	2.6	3.0	4.4	8.3	19.4	36.6	58.6
2000[4]	20.0	*	*	*	*	1.8	2.3	1.6	2.1	5.6	12.0	21.4	33.9	52.1
1999[4]	21.3	*	*	*	*	1.9	2.1	1.6	3.2	5.2	11.9	23.8	37.4	62.9
Female														
2001	17.8	*	*	*	*	*	*	1.6	2.4	4.3	7.4	15.2	21.5	34.2
2000[4]	17.8	*	*	*	*	1.4	*	*	1.9	5.2	8.5	13.5	25.7	33.8
1999[4]	19.2	*	*	*	*	*	*	1.5	1.8	4.9	9.0	14.4	23.3	38.1
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)														
All origins[3]														
Both sexes														
2001	4.8	1.3	0.4	0.3	0.3	0.2	0.3	0.3	0.4	0.7	0.9	1.5	2.3	3.9
2000[4]	4.8	1.6	0.3	0.3	0.2	0.2	0.3	0.4	0.5	0.7	0.9	1.5	2.3	3.8
1999[4]	4.7	1.5	0.4	0.3	0.2	0.3	0.2	0.3	0.6	0.6	0.9	1.4	2.3	3.5
Male														
2001	4.8	1.3	0.4	0.3	0.2	0.2	0.3	0.3	0.4	0.7	1.0	1.7	2.4	4.1
2000[4]	4.8	1.6	0.4	0.3	*	0.3	0.4	0.3	0.5	0.8	0.9	1.6	2.6	4.1
1999[4]	4.8	1.5	0.4	0.3	*	0.3	0.3	0.3	0.6	0.7	1.1	1.5	2.7	4.0
Female														
2001	4.8	1.3	0.4	0.3	0.3	0.3	0.2	0.3	0.5	0.7	0.8	1.2	2.1	3.7
2000[4]	4.9	1.5	0.3	0.3	0.2	*	0.2	0.4	0.4	0.6	1.0	1.3	1.9	3.5
1999[4]	4.7	1.5	0.4	0.3	*	0.3	*	0.2	0.5	0.5	0.8	1.3	1.9	3.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31, C37-C41,C44-C49,C51-C52, C57-C60,C62-C63,C66,C68-C69, C73-C80,C97)						
Non-Hispanic White						
Female						
1999[4]	43.3	65.0	88.8	120.2	163.2	224.3
Non-Hispanic Black						
Both sexes						
2001	65.3	86.5	120.7	143.0	182.4	218.5
2000[4]	65.9	86.8	118.5	156.7	174.4	211.1
1999[4]	74.4	86.6	129.5	159.2	194.6	244.8
Male						
2001	82.6	111.8	150.5	181.6	218.3	287.6
2000[4]	90.4	103.1	150.3	211.9	212.0	253.6
1999[4]	95.8	111.0	165.1	193.8	235.6	293.6
Female						
2001	51.8	67.9	100.9	120.0	164.1	192.9
2000[4]	46.6	74.7	97.3	123.4	155.5	195.4
1999[4]	57.7	68.5	105.9	138.3	174.3	226.5
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)						
All origins[3]						
Both sexes						
2001	6.3	10.5	17.9	29.5	47.8	82.7
2000[4]	6.6	10.6	17.4	28.9	50.6	84.9
1999[4]	6.3	10.8	18.6	29.3	47.5	82.9
Male						
2001	7.5	12.8	22.5	39.2	62.4	107.6
2000[4]	7.5	13.3	21.2	36.8	65.3	108.9
1999[4]	7.9	13.4	23.9	37.2	61.0	109.2
Female						
2001	5.2	8.5	14.3	22.7	39.1	72.3
2000[4]	5.7	8.2	14.4	23.4	41.9	75.2
1999[4]	4.8	8.5	14.5	23.8	39.7	72.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)														
Hispanic														
Both sexes														
2001	1.5	*	*	*	*	*	*	*	*	*	1.0	1.1	1.9	2.4
2000[4]	1.5	*	*	*	*	*	*	*	*	*	1.0	1.1	1.7	2.4
1999[4]	1.5	*	*	*	*	*	*	*	*	*	1.1	*	1.9	3.4
Male														
2001	1.5	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	1.5	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.7	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.6	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	5.3	1.4	0.4	0.2	0.3	0.3	0.3	0.3	0.5	0.7	0.9	1.5	2.3	4.0
2000[4]	5.3	1.7	0.4	0.3	0.2	0.2	0.3	0.4	0.5	0.7	0.9	1.5	2.3	3.9
1999[4]	5.2	1.6	0.4	0.3	0.2	0.3	0.3	0.3	0.6	0.7	0.9	1.5	2.3	3.5
Male														
2001	5.3	1.5	0.4	*	0.3	*	0.4	0.4	0.5	0.8	1.0	1.7	2.4	4.2
2000[4]	5.3	1.8	0.5	0.3	*	0.3	0.4	0.4	0.5	0.8	0.9	1.7	2.7	4.2
1999[4]	5.3	1.6	0.5	0.3	*	0.2	0.3	0.3	0.7	0.8	1.0	1.5	2.6	4.1
Female														
2001	5.2	1.4	0.4	0.3	0.3	0.3	0.3	0.3	0.4	0.7	0.9	1.3	2.1	3.8
2000[4]	5.3	1.7	*	0.4	*	*	*	0.4	0.5	0.6	0.9	1.3	1.9	3.6
1999[4]	5.1	1.7	0.4	0.3	*	0.3	*	0.3	0.5	0.6	0.8	1.4	2.0	3.0
Non-Hispanic White														
Both sexes														
2001	5.9	1.3	0.4	0.2	0.2	0.2	0.3	0.4	0.5	0.7	0.9	1.5	2.3	4.0
2000[4]	5.9	1.8	0.4	0.3	0.2	0.2	0.3	0.3	0.5	0.7	0.9	1.4	2.3	4.0
1999[4]	5.7	1.5	0.4	0.3	0.2	0.2	0.2	0.2	0.5	0.6	0.8	1.3	2.2	3.4
Male														
2001	6.0	*	*	*	*	*	0.3	0.4	0.5	0.8	0.9	1.8	2.5	4.3
2000[4]	5.9	1.8	0.5	0.3	*	*	0.4	0.4	0.5	0.8	0.9	1.6	2.7	4.4
1999[4]	5.9	*	*	*	*	*	*	*	0.6	0.7	1.0	1.5	2.6	3.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)						
Hispanic						
Both sexes						
2001	5.3	7.5	10.5	19.0	30.3	57.5
2000[4]	4.0	6.7	12.2	20.5	31.2	70.3
1999[4]	4.0	6.7	15.3	18.4	31.7	51.9
Male						
2001	6.8	7.4	13.3	19.8	49.3	70.5
2000[4]	*	*	14.6	26.6	*	86.7
1999[4]	*	*	17.3	18.7	36.2	49.6
Female						
2001	*	7.5	8.4	18.4	18.5	51.2
2000[4]	*	7.2	10.3	16.2	34.0	62.3
1999[4]	*	8.1	13.8	18.2	29.1	53.0
Non-Hispanic[5]						
Both sexes						
2001	6.3	10.7	18.3	30.0	48.3	83.5
2000[4]	6.8	10.8	17.6	29.2	51.2	85.2
1999[4]	6.4	11.0	18.8	29.7	48.0	83.8
Male						
2001	7.5	13.1	23.0	40.1	62.7	109.0
2000[4]	7.8	13.8	21.6	37.2	66.6	109.4
1999[4]	8.1	13.9	24.3	37.9	61.8	111.4
Female						
2001	5.3	8.6	14.6	22.9	39.7	72.9
2000[4]	5.8	8.2	14.6	23.7	42.1	75.4
1999[4]	4.9	8.5	14.5	24.0	39.9	72.8
Non-Hispanic White						
Both sexes						
2001	6.3	11.0	18.6	30.9	49.4	84.8
2000[4]	6.8	11.0	17.6	29.8	52.8	87.6
1999[4]	6.4	10.9	19.2	30.1	49.2	86.3
Male						
2001	7.4	13.5	23.6	41.3	63.9	111.4
2000[4]	7.5	13.9	21.3	38.0	69.4	113.1
1999[4]	8.2	13.9	24.9	38.6	63.8	116.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)														
Non-Hispanic White														
Female														
2001	5.8	*	*	*	*	*	*	0.3	0.4	0.7	0.8	1.3	2.1	3.8
2000[4]	5.9	1.8	*	0.3	*	*	*	*	0.5	0.6	0.9	1.2	1.8	3.6
1999[4]	5.6	*	*	0.3	*	*	*	*	0.4	0.5	0.7	1.2	1.9	3.0
Non-Hispanic Black														
Both sexes														
2001	3.2	*	*	*	*	*	*	*	*	0.9	1.3	1.5	2.8	4.6
2000[4]	3.2	*	*	*	*	*	*	*	*	0.8	1.4	2.4	3.2	4.2
1999[4]	3.3	*	*	*	*	*	*	*	0.8	1.1	1.3	2.3	3.1	5.0
Male														
2001	3.1	*	*	*	*	*	*	*	*	*	*	*	2.5	4.5
2000[4]	3.1	*	*	*	*	*	*	*	*	*	*	2.2	3.7	4.2
1999[4]	3.1	*	*	*	*	*	*	*	*	*	*	*	3.2	6.3
Female														
2001	3.3	*	*	*	*	*	*	*	*	*	*	*	3.0	4.6
2000[4]	3.3	*	*	*	*	*	*	*	*	*	*	2.5	2.8	4.2
1999[4]	3.5	*	*	*	*	*	*	*	*	*	1.4	2.6	3.1	4.0
Anemias (D50-D64)														
All origins[3]														
Both sexes														
2001	1.6	*	0.1	0.1	0.1	0.2	0.3	0.3	0.5	0.4	0.6	0.5	0.7	0.8
2000[4]	1.6	*	0.2	0.1	0.1	0.1	0.3	0.3	0.4	0.4	0.5	0.5	0.7	0.9
1999[4]	1.6	*	0.2	0.1	0.1	0.2	0.4	0.3	0.5	0.4	0.5	0.7	0.6	0.8
Male														
2001	1.3	*	*	*	*	0.2	0.3	0.4	0.5	0.4	0.4	0.6	0.6	0.8
2000[4]	1.2	*	*	*	*	*	0.3	0.3	0.4	0.3	0.4	0.5	0.7	1.0
1999[4]	1.3	*	*	*	*	0.2	0.5	0.3	0.5	0.4	0.5	0.7	0.6	0.7
Female														
2001	2.0	*	*	*	*	0.2	0.3	0.3	0.4	0.4	0.7	0.5	0.7	0.9
2000[4]	2.0	*	*	*	*	*	0.3	0.3	0.5	0.4	0.5	0.6	0.7	0.9
1999[4]	1.9	*	*	*	*	*	0.4	0.3	0.5	0.4	0.4	0.7	0.6	0.9
Hispanic														
Both sexes														
2001	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)						
Non-Hispanic White						
Female						
2001	5.3	8.7	14.5	23.5	40.6	73.7
2000[4]	6.1	8.5	14.6	24.2	43.0	77.3
1999[4]	4.7	8.3	14.7	24.2	40.6	74.1
Non-Hispanic Black						
Both sexes						
2001	7.7	11.0	18.9	25.0	42.8	72.7
2000[4]	7.3	10.9	20.3	27.0	38.2	64.2
1999[4]	8.0	14.0	18.2	29.4	39.0	62.7
Male						
2001	9.4	13.3	23.4	32.2	59.7	87.8
2000[4]	10.4	16.0	23.9	34.7	39.5	76.3
1999[4]	8.5	17.3	21.8	36.3	43.4	64.2
Female						
2001	6.4	9.2	15.8	20.7	34.1	67.2
2000[4]	4.9	7.1	17.9	22.4	37.6	59.7
1999[4]	7.7	11.6	15.9	25.3	36.9	62.2
Anemias (D50-D64)						
All origins[3]						
Both sexes						
2001	1.2	2.0	3.5	7.1	13.6	42.0
2000[4]	1.2	2.2	3.4	6.5	13.9	42.4
1999[4]	1.2	1.8	3.5	6.8	13.7	43.8
Male						
2001	1.1	2.1	3.8	7.2	14.3	40.0
2000[4]	1.2	2.2	3.6	6.5	14.6	40.1
1999[4]	1.2	2.1	3.4	7.2	14.8	43.4
Female						
2001	1.2	1.9	3.3	7.0	13.2	42.9
2000[4]	1.2	2.2	3.3	6.5	13.6	43.3
1999[4]	1.1	1.6	3.6	6.5	13.0	44.0
Hispanic						
Both sexes						
2001	*	*	*	6.8	*	28.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Anemias (D50-D64)														
Hispanic														
Both sexes														
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	1.8	*	*	0.1	0.2	0.2	0.3	0.4	0.5	0.4	0.6	0.6	0.7	0.8
2000[4]	1.8	*	0.3	0.1	0.1	0.2	0.3	0.3	0.5	0.4	0.5	0.6	0.7	1.0
1999[4]	1.8	*	0.2	0.1	0.2	0.2	0.5	0.4	0.5	0.4	0.5	0.7	0.6	0.8
Male														
2001	1.4	*	*	*	*	0.3	0.4	0.5	0.6	0.4	0.5	0.6	0.7	0.8
2000[4]	1.3	*	*	*	*	*	0.4	0.3	0.5	0.3	0.4	0.6	0.7	1.1
1999[4]	1.4	*	*	*	*	0.3	0.5	0.4	0.5	0.4	0.5	0.7	0.6	0.7
Female														
2001	2.2	*	*	*	*	*	0.3	0.3	0.4	0.4	0.8	0.5	0.7	0.9
2000[4]	2.2	*	*	*	*	*	0.3	0.3	0.5	0.5	0.6	0.6	0.7	0.9
1999[4]	2.1	*	*	*	*	*	0.4	0.4	0.5	0.4	0.5	0.7	0.6	0.9
Non-Hispanic White														
Both sexes														
2001	1.7	*	*	*	*	*	*	*	*	*	0.2	0.2	0.4	0.5
2000[4]	1.7	*	*	*	*	*	*	*	*	*	0.1	0.2	0.4	0.7
1999[4]	1.6	*	*	*	*	*	*	*	*	*	0.1	0.3	0.3	0.4
Male														
2001	1.3	*	*	*	*	*	*	*	*	*	*	*	0.4	0.5
2000[4]	1.2	*	*	*	*	*	*	*	*	*	*	*	0.4	0.8
1999[4]	1.2	*	*	*	*	*	*	*	*	*	*	0.4	0.3	*
Female														
2001	2.1	*	*	*	*	*	*	*	*	*	0.3	*	0.4	0.5
2000[4]	2.1	*	*	*	*	*	*	*	*	*	*	*	0.4	0.6
1999[4]	2.0	*	*	*	*	*	*	*	*	*	*	0.3	*	0.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Anemias (D50-D64)						
Hispanic						
Both sexes						
2000[4]	*	*	*	*	12.8	17.3
1999[4]	*	*	4.4	*	*	35.1
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	32.6
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	32.2
Non-Hispanic[5]						
Both sexes						
2001	1.2	1.9	3.6	7.1	13.7	42.5
2000[4]	1.2	2.2	3.4	6.6	13.9	43.2
1999[4]	1.2	1.8	3.4	6.8	13.7	44.0
Male						
2001	1.1	2.0	3.9	7.1	14.4	40.8
2000[4]	1.3	2.2	3.5	6.6	14.4	41.0
1999[4]	1.2	2.1	3.2	7.2	14.7	43.3
Female						
2001	1.3	1.9	3.3	7.0	13.3	43.1
2000[4]	1.2	2.3	3.4	6.6	13.6	44.1
1999[4]	1.2	1.6	3.6	6.5	13.2	44.2
Non-Hispanic White						
Both sexes						
2001	1.1	1.7	3.3	6.7	13.2	42.1
2000[4]	0.9	2.0	3.2	6.3	13.8	42.9
1999[4]	0.8	1.6	3.1	6.5	13.7	43.2
Male						
2001	1.0	1.8	3.4	6.9	14.2	41.2
2000[4]	0.9	2.1	3.3	6.3	14.1	39.9
1999[4]	0.7	1.9	2.8	7.0	14.7	43.4
Female						
2001	1.1	1.7	3.1	6.5	12.7	42.4
2000[4]	0.9	1.8	3.1	6.3	13.5	44.1
1999[4]	1.0	1.4	3.4	6.1	13.1	43.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Anemias (D50-D64)														
Non-Hispanic Black														
Both sexes														
2001	2.8	*	*	*	*	0.9	1.6	2.0	2.5	2.2	3.3	3.3	2.9	3.3
2000[4]	2.7	*	*	*	*	*	1.7	1.8	2.9	2.3	2.7	3.0	3.2	3.0
1999[4]	2.9	*	0.9	*	*	0.8	2.4	2.1	3.2	2.3	2.8	3.3	3.0	4.3
Male														
2001	2.6	*	*	*	*	*	1.7	2.6	3.4	2.3	2.8	4.0	3.2	3.4
2000[4]	2.4	*	*	*	*	*	1.7	1.6	2.8	2.0	2.3	3.3	3.5	*
1999[4]	2.7	*	*	*	*	*	2.9	2.4	3.4	2.6	3.0	3.4	3.3	4.4
Female														
2001	2.9	*	*	*	*	*	1.5	*	1.8	2.1	3.8	2.7	2.7	3.3
2000[4]	3.1	*	*	*	*	*	1.7	2.0	3.0	2.6	3.1	2.8	3.1	3.0
1999[4]	3.1	*	*	*	*	*	1.8	1.9	2.9	2.1	2.7	3.2	2.8	4.3
Diabetes mellitus (E10-E14)														
All origins[3]														
Both sexes														
2001	25.1	*	*	*	0.2	0.2	0.5	1.1	1.8	3.3	5.4	9.9	17.8	29.5
2000[4]	24.6	*	*	*	*	0.2	0.6	1.1	2.0	3.0	5.5	9.5	17.3	29.6
1999[4]	24.5	*	*	*	0.1	0.3	0.6	1.0	1.9	3.2	5.5	9.9	16.5	29.5
Male														
2001	23.5	*	*	*	0.2	0.3	0.6	1.2	2.2	3.9	6.6	12.1	21.3	34.4
2000[4]	22.9	*	*	*	*	0.3	0.7	1.2	2.5	3.5	6.8	11.4	19.7	33.4
1999[4]	22.8	*	*	*	*	0.2	0.6	1.0	2.2	3.8	6.4	12.3	19.4	33.5
Female														
2001	26.6	*	*	*	*	0.2	0.5	1.1	1.5	2.7	4.2	7.9	14.5	24.9
2000[4]	26.3	*	*	*	*	*	0.5	0.9	1.6	2.5	4.3	7.8	14.9	26.1
1999[4]	26.2	*	*	*	*	0.3	0.5	1.0	1.6	2.6	4.6	7.5	13.8	25.8
Hispanic														
Both sexes														
2001	15.3	*	*	*	*	*	*	0.6	0.9	2.3	4.0	11.9	18.8	38.4
2000[4]	15.1	*	*	*	*	*	*	0.6	1.4	1.7	4.6	9.8	19.5	45.4
1999[4]	15.3	*	*	*	*	*	*	*	1.0	2.1	4.6	10.1	19.0	40.4
Male														
2001	13.6	*	*	*	*	*	*	*	*	2.6	5.4	14.4	23.8	43.4
2000[4]	13.8	*	*	*	*	*	*	*	1.9	1.9	5.7	11.1	21.4	52.8
1999[4]	13.4	*	*	*	*	*	*	*	*	2.2	4.7	12.5	23.5	46.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Anemias (D50-D64)						
Non-Hispanic Black						
Both sexes						
2001	2.7	4.0	7.4	12.6	21.5	49.3
2000[4]	4.0	4.9	6.3	10.5	18.1	51.8
1999[4]	4.3	3.7	6.6	11.4	16.4	53.0
Male						
2001	*	*	10.7	12.3	20.4	42.7
2000[4]	*	*	*	11.1	18.9	56.4
1999[4]	6.1	*	8.0	10.6	*	44.0
Female						
2001	*	4.0	5.3	12.7	22.1	51.8
2000[4]	4.0	5.9	6.4	10.2	17.7	50.2
1999[4]	*	*	5.7	11.9	16.2	56.4
Diabetes mellitus (E10-E14)						
All origins[3]						
Both sexes						
2001	48.4	75.5	108.6	155.5	218.7	321.8
2000[4]	48.1	74.0	108.6	156.5	213.9	319.7
1999[4]	48.9	75.6	109.2	153.5	214.8	317.2
Male						
2001	54.4	84.7	121.0	171.5	242.3	343.1
2000[4]	53.9	80.8	119.7	173.7	242.5	345.1
1999[4]	53.9	82.5	119.4	172.3	241.3	345.7
Female						
2001	42.9	67.6	98.8	144.3	204.5	312.9
2000[4]	42.8	68.1	99.9	144.6	197.0	309.4
1999[4]	44.5	69.7	101.2	140.4	199.3	305.8
Hispanic						
Both sexes						
2001	72.8	121.1	166.3	254.5	322.3	489.6
2000[4]	68.5	114.1	190.3	250.1	330.3	465.1
1999[4]	77.1	124.6	181.4	252.4	344.8	493.9
Male						
2001	79.6	132.5	175.7	262.6	328.5	422.8
2000[4]	80.3	126.0	206.1	267.2	363.7	449.4
1999[4]	82.8	127.9	192.5	262.1	355.2	468.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Diabetes mellitus (E10-E14)														
Hispanic														
Female														
2001	17.1	*	*	*	*	*	*	*	*	2.0	2.4	9.4	14.0	33.8
2000[4]	16.5	*	*	*	*	*	*	*	*	1.6	3.4	8.6	17.7	38.7
1999[4]	17.3	*	*	*	*	*	*	*	*	1.9	4.4	7.6	14.7	34.6
Non-Hispanic[5]														
Both sexes														
2001	26.4	*	*	*	0.2	0.2	0.6	1.3	2.0	3.5	5.5	9.7	17.6	28.7
2000[4]	25.9	*	*	*	*	0.3	0.7	1.2	2.1	3.2	5.6	9.5	17.0	28.3
1999[4]	25.7	*	*	*	*	0.3	0.6	1.1	2.1	3.3	5.5	9.8	16.2	28.6
Male														
2001	25.0	*	*	*	*	0.3	0.6	1.4	2.4	4.1	6.7	11.7	21.0	33.6
2000[4]	24.2	*	*	*	*	0.3	0.8	1.3	2.6	3.7	6.9	11.4	19.5	31.8
1999[4]	24.1	*	*	*	*	0.3	0.6	1.0	2.4	4.1	6.6	12.3	18.9	32.5
Female														
2001	27.8	*	*	*	*	*	0.5	1.1	1.6	2.8	4.4	7.7	14.4	24.1
2000[4]	27.6	*	*	*	*	*	0.6	1.0	1.7	2.7	4.4	7.7	14.6	25.1
1999[4]	27.3	*	*	*	*	0.3	0.6	1.1	1.7	2.6	4.6	7.5	13.7	25.0
Non-Hispanic White														
Both sexes														
2001	26.0	*	*	*	*	0.2	0.4	0.9	1.7	2.9	4.8	8.1	14.7	23.9
2000[4]	25.4	*	*	*	*	0.2	0.5	0.9	1.8	2.8	5.0	7.8	14.0	23.7
1999[4]	25.0	*	*	*	*	0.3	0.5	1.0	1.8	2.9	4.7	8.1	13.2	23.0
Male														
2001	25.1	*	*	*	*	*	0.5	0.9	1.9	3.4	6.2	9.9	17.9	29.0
2000[4]	24.3	*	*	*	*	*	0.7	1.0	2.3	3.2	6.3	9.5	16.7	27.3
1999[4]	24.0	*	*	*	*	0.3	0.5	1.0	2.0	3.6	5.6	10.3	16.0	27.2
Female														
2001	26.9	*	*	*	*	*	*	0.9	1.5	2.4	3.5	6.3	11.6	18.9
2000[4]	26.5	*	*	*	*	*	0.4	0.8	1.3	2.4	3.7	6.1	11.3	20.4
1999[4]	26.0	*	*	*	*	*	0.5	0.9	1.5	2.1	3.8	5.9	10.5	18.9
Non-Hispanic Black														
Both sexes														
2001	34.1	*	*	*	*	*	1.4	3.3	4.4	7.2	11.0	21.3	41.0	70.6
2000[4]	34.0	*	*	*	*	*	1.5	2.9	4.3	6.2	10.4	22.4	41.7	67.7
1999[4]	34.1	*	*	*	*	*	1.2	1.9	4.2	6.8	11.6	23.1	41.1	74.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diabetes mellitus (E10-E14)						
Hispanic						
Female						
2001	67.0	111.9	159.1	248.9	318.5	522.2
2000[4]	58.3	104.5	178.2	237.9	310.0	472.8
1999[4]	72.1	121.9	173.1	245.6	338.5	506.4
Non-Hispanic[5]						
Both sexes						
2001	46.4	71.9	104.7	150.1	213.9	314.8
2000[4]	46.4	71.1	103.6	151.7	209.0	313.4
1999[4]	46.7	72.3	105.0	148.7	209.5	310.0
Male						
2001	52.4	81.0	117.1	166.4	238.1	338.9
2000[4]	51.9	77.6	114.4	168.6	237.2	339.7
1999[4]	51.6	79.4	115.2	167.9	236.6	339.5
Female						
2001	41.0	64.1	94.7	138.7	199.4	304.7
2000[4]	41.4	65.5	95.1	139.9	192.4	302.8
1999[4]	42.2	66.1	97.0	135.3	193.6	298.2
Non-Hispanic White						
Both sexes						
2001	39.4	62.0	92.7	136.3	199.9	297.4
2000[4]	39.7	61.2	91.7	137.5	193.0	295.5
1999[4]	38.7	62.2	92.0	135.2	194.3	290.2
Male						
2001	45.9	72.8	108.6	155.6	226.9	332.1
2000[4]	45.7	69.6	105.1	157.9	227.9	330.4
1999[4]	44.5	70.9	106.0	158.4	227.5	326.0
Female						
2001	33.3	52.5	79.7	122.6	183.7	283.0
2000[4]	34.2	53.7	80.9	123.2	172.2	281.4
1999[4]	33.3	54.5	80.8	119.0	174.7	275.9
Non-Hispanic Black						
Both sexes						
2001	106.6	159.4	229.9	307.4	385.5	533.8
2000[4]	103.7	160.7	226.7	320.5	409.2	530.1
1999[4]	113.7	162.6	232.3	297.1	399.2	530.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Diabetes mellitus (E10-E14)														
Non-Hispanic Black														
Male														
2001	29.4	*	*	*	*	*	1.5	4.1	5.8	9.0	12.0	25.8	46.4	77.8
2000[4]	28.3	*	*	*	*	*	1.6	3.3	4.7	7.4	12.0	26.1	44.4	73.5
1999[4]	28.6	*	*	*	*	*	*	1.7	5.1	8.0	13.5	28.2	44.5	79.6
Female														
2001	38.4	*	*	*	*	*	*	2.6	3.0	5.6	10.1	17.4	36.4	64.8
2000[4]	39.1	*	*	*	*	*	1.5	2.6	3.9	5.1	8.9	19.3	39.4	63.0
1999[4]	39.0	*	*	*	*	*	*	2.0	3.4	5.6	9.9	18.7	38.2	69.8
Nutritional deficiencies (E40-E64)														
All origins[3]														
Both sexes														
2001	1.3	*	*	*	*	*	*	*	*	*	0.1	0.2	0.3	0.5
2000[4]	1.5	*	*	*	*	*	*	*	*	0.1	0.1	0.2	0.3	0.5
1999[4]	1.5	*	*	*	*	*	*	*	*	0.1	0.2	0.2	0.4	0.6
Male														
2001	0.9	*	*	*	*	*	*	*	*	*	*	0.3	0.3	0.5
2000[4]	1.0	*	*	*	*	*	*	*	*	*	*	0.3	0.2	0.7
1999[4]	1.1	*	*	*	*	*	*	*	*	*	0.2	0.2	0.4	0.7
Female														
2001	1.7	*	*	*	*	*	*	*	*	*	*	0.2	0.4	0.4
2000[4]	2.0	*	*	*	*	*	*	*	*	0.2	*	0.2	0.4	0.4
1999[4]	2.0	*	*	*	*	*	*	*	*	*	0.2	0.2	0.4	0.5
Hispanic														
Both sexes														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diabetes mellitus (E10-E14)						
Non-Hispanic Black						
Male						
2001	114.5	163.9	223.4	310.5	404.7	452.7
2000[4]	110.1	155.4	227.2	314.1	384.5	456.8
1999[4]	115.2	167.3	218.8	284.2	379.0	506.3
Female						
2001	100.4	156.0	234.2	305.5	375.6	563.8
2000[4]	98.6	164.6	226.4	324.4	421.7	557.3
1999[4]	112.5	159.1	241.2	304.9	409.2	540.1
Nutritional deficiencies (E40-E64)						
All origins[3]						
Both sexes						
2001	0.8	1.4	2.8	5.9	12.8	42.9
2000[4]	1.0	2.0	3.0	6.8	15.8	50.0
1999[4]	0.9	1.6	3.2	6.4	15.9	53.9
Male						
2001	0.9	1.4	2.9	5.6	13.2	35.2
2000[4]	1.0	2.0	3.4	6.8	14.4	43.7
1999[4]	1.1	1.6	3.8	6.8	16.0	48.4
Female						
2001	0.7	1.3	2.7	6.1	12.6	46.1
2000[4]	0.9	2.0	2.7	6.8	16.6	52.6
1999[4]	0.7	1.5	2.6	6.2	15.8	56.2
Hispanic						
Both sexes						
2001	*	*	*	6.2	11.4	38.8
2000[4]	*	*	*	*	13.9	39.8
1999[4]	*	*	*	*	*	51.2
Male						
2001	*	*	*	*	*	43.8
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	51.8
Female						
2001	*	*	*	*	*	36.3
2000[4]	*	*	*	*	*	41.5
1999[4]	*	*	*	*	*	51.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Nutritional deficiencies (E40-E64)														
Non-Hispanic[5]														
Both sexes														
2001	1.4	*	*	*	*	*	*	*	*	*	0.1	0.2	0.4	0.5
2000[4]	1.7	*	*	*	*	*	*	*	*	0.1	0.1	0.2	0.3	0.5
1999[4]	1.7	*	*	*	*	*	*	*	*	*	0.2	0.2	0.4	0.6
Male														
2001	1.0	*	*	*	*	*	*	*	*	*	*	0.3	0.3	0.5
2000[4]	1.1	*	*	*	*	*	*	*	*	*	*	0.3	*	0.7
1999[4]	1.2	*	*	*	*	*	*	*	*	*	*	0.2	0.3	0.7
Female														
2001	1.9	*	*	*	*	*	*	*	*	*	*	0.2	0.4	0.4
2000[4]	2.2	*	*	*	*	*	*	*	*	*	*	0.2	0.4	0.4
1999[4]	2.2	*	*	*	*	*	*	*	*	*	0.2	0.3	0.4	0.5
Non-Hispanic White														
Both sexes														
2001	1.5	*	*	*	*	*	*	*	*	*	*	0.2	0.3	0.4
2000[4]	1.8	*	*	*	*	*	*	*	*	*	*	0.2	0.2	0.5
1999[4]	1.8	*	*	*	*	*	*	*	*	*	0.2	0.2	0.3	0.4
Male														
2001	1.0	*	*	*	*	*	*	*	*	*	*	*	*	0.4
2000[4]	1.2	*	*	*	*	*	*	*	*	*	*	*	*	0.5
1999[4]	1.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	2.0	*	*	*	*	*	*	*	*	*	*	*	0.3	0.4
2000[4]	2.3	*	*	*	*	*	*	*	*	*	*	*	0.3	0.4
1999[4]	2.3	*	*	*	*	*	*	*	*	*	*	*	0.4	0.4
Non-Hispanic Black														
Both sexes														
2001	1.3	*	*	*	*	*	*	*	*	*	*	*	1.1	*
2000[4]	1.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.7	*	*	*	*	*	*	*	*	*	*	*	*	2.0
Male														
2001	1.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	1.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.8	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nutritional deficiencies (E40-E64)						
Non-Hispanic[5]						
Both sexes						
2001	0.8	1.4	2.9	5.9	12.9	43.0
2000[4]	1.0	2.0	3.1	6.8	15.8	50.3
1999[4]	0.9	1.6	3.2	6.5	16.1	53.9
Male						
2001	0.9	1.5	3.0	5.5	13.2	34.8
2000[4]	1.1	2.0	3.4	6.7	14.3	43.9
1999[4]	1.2	1.7	3.8	6.7	16.4	48.1
Female						
2001	0.8	1.3	2.7	6.2	12.7	46.4
2000[4]	0.9	2.0	2.8	6.9	16.6	52.8
1999[4]	0.7	1.5	2.6	6.3	16.0	56.2
Non-Hispanic White						
Both sexes						
2001	0.8	1.2	2.7	5.4	12.0	42.2
2000[4]	0.9	1.9	2.7	6.4	15.3	49.0
1999[4]	0.8	1.4	2.6	6.0	15.1	52.8
Male						
2001	0.8	1.2	2.7	5.1	12.3	33.3
2000[4]	0.9	1.8	2.8	6.1	13.8	43.0
1999[4]	1.0	1.4	3.0	5.9	15.0	46.2
Female						
2001	0.8	1.3	2.7	5.7	11.8	45.9
2000[4]	0.9	1.9	2.6	6.5	16.1	51.5
1999[4]	0.6	1.5	2.3	6.0	15.1	55.4
Non-Hispanic Black						
Both sexes						
2001	*	3.1	4.5	11.0	24.0	55.2
2000[4]	2.2	3.3	6.7	12.1	23.0	67.4
1999[4]	2.0	3.3	8.4	12.5	31.1	70.8
Male						
2001	*	*	6.9	11.9	24.5	58.9
2000[4]	*	*	9.9	14.9	23.2	64.6
1999[4]	*	*	13.2	16.4	35.4	76.1
Female						
2001	*	*	*	10.5	23.7	53.9
2000[4]	*	*	4.5	10.5	22.9	68.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Nutritional deficiencies (E40-E64)														
Non-Hispanic Black														
Female														
1999[4]	1.8	*	*	*	*	*	*	*	*	*	*	*	*	*
Malnutrition (E40-E46)														
All origins[3]														
Both sexes														
2001	1.2	*	*	*	*	*	*	*	*	*	0.1	0.2	0.3	0.4
2000[4]	1.4	*	*	*	*	*	*	*	*	0.1	0.1	0.2	0.3	0.5
1999[4]	1.4	*	*	*	*	*	*	*	*	0.1	0.2	0.2	0.3	0.5
Male														
2001	0.8	*	*	*	*	*	*	*	*	*	*	0.2	0.3	0.5
2000[4]	0.9	*	*	*	*	*	*	*	*	*	*	0.2	*	0.6
1999[4]	1.0	*	*	*	*	*	*	*	*	*	*	0.2	0.3	0.6
Female														
2001	1.6	*	*	*	*	*	*	*	*	*	*	0.2	0.3	0.4
2000[4]	1.8	*	*	*	*	*	*	*	*	0.2	*	0.2	0.3	0.3
1999[4]	1.8	*	*	*	*	*	*	*	*	*	0.2	0.2	0.3	0.4
Hispanic														
Both sexes														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	1.3	*	*	*	*	*	*	*	*	*	0.1	0.2	0.3	0.5
2000[4]	1.5	*	*	*	*	*	*	*	*	0.1	0.1	0.2	0.3	0.5
1999[4]	1.6	*	*	*	*	*	*	*	*	*	0.2	0.2	0.3	0.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nutritional deficiencies (E40-E64)						
Non-Hispanic Black						
Female						
1999[4]	*	*	5.3	10.2	29.0	68.8
Malnutrition (E40-E46)						
All origins[3]						
Both sexes						
2001	0.8	1.3	2.6	5.5	12.0	40.2
2000[4]	0.9	1.9	2.8	6.4	14.6	46.3
1999[4]	0.8	1.5	3.0	6.0	14.7	50.0
Male						
2001	0.8	1.3	2.6	5.1	12.6	33.6
2000[4]	0.9	1.8	3.3	6.4	13.1	40.7
1999[4]	1.1	1.5	3.6	6.2	14.8	44.4
Female						
2001	0.7	1.3	2.5	5.7	11.6	43.0
2000[4]	0.9	1.9	2.4	6.5	15.5	48.6
1999[4]	0.6	1.4	2.5	5.8	14.7	52.3
Hispanic						
Both sexes						
2001	*	*	*	6.2	10.9	38.1
2000[4]	*	*	*	*	12.8	38.5
1999[4]	*	*	*	*	*	47.7
Male						
2001	*	*	*	*	*	41.9
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	45.3
Female						
2001	*	*	*	*	*	36.3
2000[4]	*	*	*	*	*	41.5
1999[4]	*	*	*	*	*	48.9
Non-Hispanic[5]						
Both sexes						
2001	0.8	1.3	2.6	5.4	12.1	40.2
2000[4]	0.9	1.9	2.8	6.4	14.6	46.5
1999[4]	0.9	1.5	3.0	6.0	14.9	50.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malnutrition (E40-E46)														
Non-Hispanic[5]														
Male														
2001	0.9	*	*	*	*	*	*	*	*	*	*	0.2	0.3	0.5
2000[4]	1.0	*	*	*	*	*	*	*	*	*	*	0.2	*	0.6
1999[4]	1.1	*	*	*	*	*	*	*	*	*	*	0.2	0.3	0.6
Female														
2001	1.7	*	*	*	*	*	*	*	*	*	*	*	0.4	0.4
2000[4]	2.0	*	*	*	*	*	*	*	*	*	*	0.2	0.3	0.3
1999[4]	2.0	*	*	*	*	*	*	*	*	*	0.2	0.2	0.4	0.4
Non-Hispanic White														
Both sexes														
2001	1.4	*	*	*	*	*	*	*	*	*	*	0.2	0.2	0.4
2000[4]	1.6	*	*	*	*	*	*	*	*	*	*	0.2	0.2	0.4
1999[4]	1.6	*	*	*	*	*	*	*	*	*	0.2	0.2	0.3	0.3
Male														
2001	0.9	*	*	*	*	*	*	*	*	*	*	*	*	0.4
2000[4]	1.1	*	*	*	*	*	*	*	*	*	*	*	*	0.4
1999[4]	1.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	1.9	*	*	*	*	*	*	*	*	*	*	*	0.3	0.4
2000[4]	2.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	2.2	*	*	*	*	*	*	*	*	*	*	*	0.4	*
Non-Hispanic Black														
Both sexes														
2001	1.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.6	*	*	*	*	*	*	*	*	*	*	*	*	1.6
Male														
2001	1.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	1.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.6	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.7	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malnutrition (E40-E46)						
Non-Hispanic[5]						
Male						
2001	0.8	1.4	2.7	4.9	12.7	33.2
2000[4]	1.0	1.8	3.2	6.3	13.0	40.9
1999[4]	1.1	1.6	3.6	6.1	15.1	44.3
Female						
2001	0.8	1.3	2.6	5.7	11.7	43.2
2000[4]	0.9	1.9	2.5	6.6	15.5	48.7
1999[4]	0.6	1.5	2.5	6.0	14.9	52.2
Non-Hispanic White						
Both sexes						
2001	0.8	1.2	2.5	5.0	11.2	39.4
2000[4]	0.8	1.7	2.4	6.1	14.1	45.3
1999[4]	0.8	1.3	2.5	5.6	13.9	48.8
Male						
2001	0.7	1.1	2.5	4.5	11.8	31.7
2000[4]	0.8	1.7	2.6	5.8	12.3	39.8
1999[4]	1.0	1.3	2.8	5.5	13.7	42.0
Female						
2001	0.8	1.2	2.5	5.3	10.9	42.7
2000[4]	0.9	1.8	2.2	6.3	15.1	47.4
1999[4]	0.5	1.4	2.2	5.6	14.1	51.6
Non-Hispanic Black						
Both sexes						
2001	*	2.9	4.1	10.3	22.3	52.1
2000[4]	2.0	3.2	6.7	11.2	21.3	64.2
1999[4]	2.0	3.3	8.1	11.3	29.1	66.0
Male						
2001	*	*	*	11.4	23.7	56.6
2000[4]	*	*	9.9	14.0	23.2	62.2
1999[4]	*	*	12.5	13.5	33.7	74.9
Female						
2001	*	*	*	9.6	21.7	50.5
2000[4]	*	*	4.5	9.6	20.3	65.0
1999[4]	*	*	5.3	9.9	26.8	62.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other nutritional deficiencies (E50-E64)														
All origins[3]														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other nutritional deficiencies (E50-E64)						
All origins[3]						
Both sexes						
2001	*	*	*	0.4	0.8	2.7
2000[4]	*	*	0.3	0.4	1.2	3.7
1999[4]	*	*	*	0.4	1.1	3.9
Male						
2001	*	*	*	*	*	1.6
2000[4]	*	*	*	*	1.3	3.0
1999[4]	*	*	*	*	1.3	3.9
Female						
2001	*	*	*	*	1.0	3.1
2000[4]	*	*	*	*	1.1	4.0
1999[4]	*	*	*	*	1.1	3.9
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	0.5	0.8	2.7
2000[4]	*	*	0.3	0.4	1.2	3.8
1999[4]	*	*	*	0.5	1.2	3.9
Male						
2001	*	*	*	*	*	1.6
2000[4]	*	*	*	*	1.4	3.0
1999[4]	*	*	*	*	1.3	3.8
Female						
2001	*	*	*	*	1.0	3.2
2000[4]	*	*	*	*	1.1	4.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other nutritional deficiencies (E50-E64)														
Non-Hispanic[5]														
Female														
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Meningitis (G00,G03)														
All origins[3]														
Both sexes														
2001	0.3	1.9	0.2	*	*	*	0.1	*	0.1	0.1	0.2	0.3	0.3	0.4
2000[4]	0.3	2.4	0.2	*	*	*	*	*	0.1	0.2	0.2	0.2	0.3	0.3
1999[4]	0.3	3.1	0.3	*	*	*	0.1	0.1	0.1	0.1	0.2	0.3	0.3	0.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other nutritional deficiencies (E50-E64)						
Non-Hispanic[5]						
Female						
1999[4]	*	*	*	*	1.1	4.0
Non-Hispanic White						
Both sexes						
2001	*	*	*	0.4	0.8	2.8
2000[4]	*	*	0.3	*	1.2	3.8
1999[4]	*	*	*	0.4	1.1	3.9
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	1.5	3.1
1999[4]	*	*	*	*	1.3	4.2
Female						
2001	*	*	*	*	0.9	3.3
2000[4]	*	*	*	*	1.0	4.0
1999[4]	*	*	*	*	1.0	3.9
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Meningitis (G00,G03)						
All origins[3]						
Both sexes						
2001	0.5	0.5	0.5	0.6	1.0	0.9
2000[4]	0.4	0.5	0.7	0.9	1.1	1.2
1999[4]	0.5	0.5	0.7	0.9	1.2	1.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Meningitis (G00,G03)														
All origins[3]														
Male														
2001	0.3	1.9	*	*	*	*	*	*	*	0.2	*	0.3	0.3	0.4
2000[4]	0.3	2.5	0.3	*	*	*	*	*	*	0.2	0.2	0.3	0.4	0.4
1999[4]	0.3	3.1	0.3	*	*	*	*	*	*	*	0.2	0.4	0.4	0.4
Female														
2001	0.2	2.0	*	*	*	*	*	*	*	*	*	0.2	0.2	0.4
2000[4]	0.2	2.3	*	*	*	*	*	*	*	*	*	0.2	0.3	*
1999[4]	0.3	3.0	*	*	*	*	*	*	*	*	*	0.2	*	*
Hispanic														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.3	2.2	0.2	*	*	*	*	*	0.1	0.1	0.2	0.3	0.3	0.4
2000[4]	0.3	2.7	0.3	*	*	*	*	*	0.1	0.2	0.2	0.2	0.3	0.3
1999[4]	0.3	3.2	0.3	*	*	*	*	*	0.1	0.2	0.2	0.3	0.3	0.3
Male														
2001	0.3	2.1	*	*	*	*	*	*	*	*	*	0.3	0.4	0.4
2000[4]	0.3	2.7	*	*	*	*	*	*	*	*	0.2	0.3	0.3	0.4
1999[4]	0.3	3.1	0.3	*	*	*	*	*	*	*	0.2	0.4	0.4	0.3
Female														
2001	0.2	2.3	*	*	*	*	*	*	*	*	*	0.2	*	0.4
2000[4]	0.3	2.7	*	*	*	*	*	*	*	*	*	0.2	0.3	*
1999[4]	0.3	3.3	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Meningitis (G00,G03)						
All origins[3]						
Male						
2001	0.6	0.5	0.7	0.9	1.1	*
2000[4]	0.5	0.6	0.9	1.3	1.1	*
1999[4]	0.4	0.6	0.8	0.9	1.3	1.7
Female						
2001	0.4	0.4	0.4	0.5	0.9	0.8
2000[4]	*	0.4	0.5	0.6	1.1	1.0
1999[4]	0.5	0.4	0.7	0.9	1.1	1.4
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	0.5	0.5	0.5	0.6	0.9	0.9
2000[4]	0.4	0.5	0.7	0.9	1.1	1.2
1999[4]	0.5	0.5	0.7	0.9	1.2	1.4
Male						
2001	0.6	0.5	0.7	0.9	*	*
2000[4]	0.5	0.6	0.9	1.3	1.1	*
1999[4]	0.4	0.6	0.8	0.9	1.3	*
Female						
2001	0.4	0.5	0.4	*	0.9	0.7
2000[4]	*	0.4	0.5	0.6	1.1	1.1
1999[4]	0.5	*	0.7	0.8	1.1	1.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Meningitis (G00,G03)														
Non-Hispanic White														
Both sexes														
2001	0.2	1.6	*	*	*	*	*	*	*	*	0.1	0.2	0.2	0.3
2000[4]	0.2	2.1	*	*	*	*	*	*	*	*	*	0.2	0.2	0.3
1999[4]	0.3	2.2	0.3	*	*	*	*	*	*	0.1	0.2	0.2	0.2	0.3
Male														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	0.3	*
2000[4]	0.3	2.3	*	*	*	*	*	*	*	*	*	0.3	*	*
1999[4]	0.3	2.8	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.2	1.9	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	1.8	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	0.5	4.7	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.6	5.9	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.6	7.0	*	*	*	*	*	*	*	*	*	1.0	*	*
Male														
2001	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.7	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	9.6	*	*	*	*	*	*	*	*	*	*	*	*
Parkinson's disease (G20-G21)														
All origins[3]														
Both sexes														
2001	5.8	*	*	*	*	*	*	*	*	*	*	*	0.2	0.7
2000[4]	5.6	*	*	*	*	*	*	*	*	*	*	*	0.2	0.6
1999[4]	5.2	*	*	*	*	*	*	*	*	*	*	*	0.2	0.6
Male														
2001	6.7	*	*	*	*	*	*	*	*	*	*	*	0.3	1.0
2000[4]	6.4	*	*	*	*	*	*	*	*	*	*	*	*	0.8
1999[4]	6.0	*	*	*	*	*	*	*	*	*	*	*	0.3	0.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Meningitis (G00,G03)						
Non-Hispanic White						
Both sexes						
2001	0.5	0.4	0.6	0.6	0.8	0.7
2000[4]	0.4	0.4	0.6	0.8	1.1	1.1
1999[4]	0.4	0.4	0.6	0.7	1.1	1.3
Male						
2001	0.5	*	0.7	0.8	*	*
2000[4]	0.5	*	0.9	1.1	*	*
1999[4]	*	*	0.7	*	1.3	*
Female						
2001	*	*	*	*	0.8	*
2000[4]	*	*	*	0.6	1.1	1.0
1999[4]	*	*	0.6	0.7	1.0	1.2
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Parkinson's disease (G20-G21)						
All origins[3]						
Both sexes						
2001	1.8	5.8	18.1	46.4	90.9	134.2
2000[4]	1.7	6.6	16.7	44.8	87.6	131.9
1999[4]	1.6	5.6	16.7	41.8	82.8	124.4
Male						
2001	2.7	8.2	26.5	68.8	141.7	224.4
2000[4]	2.2	9.0	24.1	67.4	135.0	222.3
1999[4]	2.3	7.7	24.8	64.4	131.3	205.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Parkinson's disease (G20-G21)														
All origins[3]														
Female														
2001	4.9	*	*	*	*	*	*	*	*	*	*	*	*	0.4
2000[4]	4.8	*	*	*	*	*	*	*	*	*	*	*	*	0.4
1999[4]	4.5	*	*	*	*	*	*	*	*	*	*	*	*	0.4
Hispanic														
Both sexes														
2001	1.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	1.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	1.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	6.5	*	*	*	*	*	*	*	*	*	*	*	0.2	0.7
2000[4]	6.2	*	*	*	*	*	*	*	*	*	*	*	0.2	0.6
1999[4]	5.8	*	*	*	*	*	*	*	*	*	*	*	0.2	0.6
Male														
2001	7.5	*	*	*	*	*	*	*	*	*	*	*	0.3	1.1
2000[4]	7.1	*	*	*	*	*	*	*	*	*	*	*	*	0.9
1999[4]	6.7	*	*	*	*	*	*	*	*	*	*	*	0.3	0.7
Female														
2001	5.4	*	*	*	*	*	*	*	*	*	*	*	*	0.4
2000[4]	5.3	*	*	*	*	*	*	*	*	*	*	*	*	0.4
1999[4]	4.9	*	*	*	*	*	*	*	*	*	*	*	*	0.4
Non-Hispanic White														
Both sexes														
2001	7.7	*	*	*	*	*	*	*	*	*	*	*	0.2	0.8
2000[4]	7.3	*	*	*	*	*	*	*	*	*	*	*	0.2	0.7
1999[4]	6.8	*	*	*	*	*	*	*	*	*	*	*	0.2	0.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Parkinson's disease (G20-G21)						
All origins[3]						
Female						
2001	1.0	3.7	11.5	30.7	60.4	96.4
2000[4]	1.3	4.4	10.9	29.1	59.5	95.0
1999[4]	0.9	3.9	10.4	26.0	54.4	91.7
Hispanic						
Both sexes						
2001	*	4.8	13.5	30.6	60.6	86.3
2000[4]	*	5.7	10.5	27.2	56.8	86.3
1999[4]	*	4.1	13.1	31.3	51.1	77.2
Male						
2001	*	*	16.9	43.0	101.3	127.6
2000[4]	*	*	13.1	39.1	86.9	143.1
1999[4]	*	*	16.8	46.8	83.3	105.7
Female						
2001	*	*	10.8	21.8	35.4	66.1
2000[4]	*	*	8.5	18.8	38.5	58.4
1999[4]	*	*	10.3	20.4	31.9	63.4
Non-Hispanic[5]						
Both sexes						
2001	1.9	5.9	18.4	47.1	91.8	135.6
2000[4]	1.8	6.6	17.1	45.5	88.5	133.3
1999[4]	1.6	5.7	16.9	42.2	83.8	125.8
Male						
2001	2.8	8.3	27.1	69.9	142.9	227.8
2000[4]	2.3	9.1	24.7	68.5	136.7	225.2
1999[4]	2.4	7.7	25.2	65.1	132.9	209.4
Female						
2001	1.0	3.8	11.5	31.1	61.3	97.3
2000[4]	1.3	4.4	11.0	29.4	60.2	96.1
1999[4]	1.0	4.0	10.4	26.2	55.1	92.3
Non-Hispanic White						
Both sexes						
2001	2.0	6.3	19.7	50.4	97.5	143.5
2000[4]	1.9	7.1	18.2	48.4	93.7	141.2
1999[4]	1.7	6.1	18.0	44.6	88.6	132.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Parkinson's disease (G20-G21)														
Non-Hispanic White														
Male														
2001	9.0	*	*	*	*	*	*	*	*	*	*	*	0.3	1.1
2000[4]	8.4	*	*	*	*	*	*	*	*	*	*	*	*	0.9
1999[4]	7.9	*	*	*	*	*	*	*	*	*	*	*	0.3	0.8
Female														
2001	6.5	*	*	*	*	*	*	*	*	*	*	*	*	0.4
2000[4]	6.3	*	*	*	*	*	*	*	*	*	*	*	*	0.5
1999[4]	5.8	*	*	*	*	*	*	*	*	*	*	*	*	0.4
Non-Hispanic Black														
Both sexes														
2001	1.5	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	1.7	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.8	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.6	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	1.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Alzheimer's disease (G30)														
All origins[3]														
Both sexes														
2001	18.9	*	*	*	*	*	*	*	*	*	*	*	0.3	1.2
2000[4]	17.6	*	*	*	*	*	*	*	*	*	*	*	0.3	1.1
1999[4]	16.0	*	*	*	*	*	*	*	*	*	*	0.1	0.2	1.0
Male														
2001	11.3	*	*	*	*	*	*	*	*	*	*	*	0.3	0.9
2000[4]	10.5	*	*	*	*	*	*	*	*	*	*	*	0.3	0.9
1999[4]	9.8	*	*	*	*	*	*	*	*	*	*	*	*	1.1
Female														
2001	26.3	*	*	*	*	*	*	*	*	*	*	*	0.3	1.4
2000[4]	24.5	*	*	*	*	*	*	*	*	*	*	*	0.3	1.2
1999[4]	21.9	*	*	*	*	*	*	*	*	*	*	*	0.3	0.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Parkinson's disease (G20-G21)						
Non-Hispanic White						
Male						
2001	3.1	8.9	28.7	74.7	151.0	241.7
2000[4]	2.5	9.8	26.2	72.0	143.4	238.6
1999[4]	2.5	8.1	26.9	68.5	140.4	221.0
Female						
2001	0.9	4.0	12.4	33.1	65.2	102.7
2000[4]	1.4	4.8	11.8	31.8	64.1	101.9
1999[4]	1.0	4.3	10.8	27.8	58.0	97.7
Non-Hispanic Black						
Both sexes						
2001	*	2.8	9.1	19.3	35.6	59.0
2000[4]	*	3.2	8.9	21.0	37.4	52.5
1999[4]	*	3.6	8.7	19.8	32.9	51.7
Male						
2001	*	*	15.1	26.1	58.9	99.3
2000[4]	*	*	14.0	37.6	65.2	88.1
1999[4]	*	5.8	12.5	32.4	49.6	87.9
Female						
2001	*	*	5.0	15.3	23.7	44.1
2000[4]	*	*	5.4	11.0	23.3	39.2
1999[4]	*	*	6.2	12.2	24.6	38.2
Alzheimer's disease (G30)						
All origins[3]						
Both sexes						
2001	3.3	9.0	29.2	88.5	232.5	710.3
2000[4]	3.2	9.1	28.9	85.2	221.1	667.7
1999[4]	3.0	8.7	26.7	78.6	206.2	601.3
Male						
2001	3.2	9.0	27.7	85.6	216.6	556.1
2000[4]	2.9	9.5	29.1	79.9	206.8	522.1
1999[4]	3.0	8.8	26.9	77.4	198.2	485.7
Female						
2001	3.4	9.0	30.4	90.5	242.1	774.9
2000[4]	3.5	8.7	28.8	88.9	229.5	727.0
1999[4]	3.0	8.6	26.6	79.4	210.9	648.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Alzheimer's disease (G30)														
Hispanic														
Both sexes														
2001	3.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	3.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	2.9	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	2.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	2.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.8	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	4.7	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	4.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	4.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	21.2	*	*	*	*	*	*	*	*	*	*	*	0.3	1.2
2000[4]	19.6	*	*	*	*	*	*	*	*	*	*	*	0.3	1.1
1999[4]	17.7	*	*	*	*	*	*	*	*	*	*	0.1	0.2	1.0
Male														
2001	12.7	*	*	*	*	*	*	*	*	*	*	*	0.3	1.0
2000[4]	11.7	*	*	*	*	*	*	*	*	*	*	*	0.3	1.0
1999[4]	10.9	*	*	*	*	*	*	*	*	*	*	*	*	1.2
Female														
2001	29.3	*	*	*	*	*	*	*	*	*	*	*	0.3	1.4
2000[4]	27.2	*	*	*	*	*	*	*	*	*	*	*	0.3	1.2
1999[4]	24.2	*	*	*	*	*	*	*	*	*	*	*	*	0.9
Non-Hispanic White														
Both sexes														
2001	24.8	*	*	*	*	*	*	*	*	*	*	*	0.4	1.4
2000[4]	22.9	*	*	*	*	*	*	*	*	*	*	*	0.3	1.2
1999[4]	20.7	*	*	*	*	*	*	*	*	*	*	*	0.3	1.1
Male														
2001	14.7	*	*	*	*	*	*	*	*	*	*	*	0.3	1.0
2000[4]	13.6	*	*	*	*	*	*	*	*	*	*	*	*	1.0
1999[4]	12.7	*	*	*	*	*	*	*	*	*	*	*	*	1.2
Female														
2001	34.4	*	*	*	*	*	*	*	*	*	*	*	0.4	1.7
2000[4]	31.9	*	*	*	*	*	*	*	*	*	*	*	*	1.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alzheimer's disease (G30)						
Hispanic						
Both sexes						
2001	*	5.1	15.8	54.9	134.6	399.5
2000[4]	*	7.2	17.8	55.1	123.1	380.2
1999[4]	*	4.3	16.4	53.6	115.1	351.5
Male						
2001	*	*	12.8	60.7	126.0	306.6
2000[4]	*	*	22.8	55.4	134.0	284.2
1999[4]	*	*	17.3	52.3	95.9	278.2
Female						
2001	*	*	18.1	50.8	139.9	444.9
2000[4]	*	8.2	14.0	54.9	116.5	427.3
1999[4]	*	*	15.7	54.5	126.6	386.8
Non-Hispanic[5]						
Both sexes						
2001	3.4	9.3	30.0	90.0	236.3	720.5
2000[4]	3.3	9.2	29.5	86.5	224.5	676.7
1999[4]	3.1	9.0	27.3	79.5	209.1	608.9
Male						
2001	3.3	9.2	28.5	86.6	220.3	565.2
2000[4]	3.0	9.8	29.3	81.0	209.5	531.0
1999[4]	3.1	9.0	27.4	78.5	201.6	492.8
Female						
2001	3.5	9.3	31.1	92.3	245.8	785.1
2000[4]	3.6	8.7	29.6	90.3	233.3	735.6
1999[4]	3.1	9.0	27.2	80.3	213.5	655.4
Non-Hispanic White						
Both sexes						
2001	3.7	9.6	31.5	92.5	244.5	749.2
2000[4]	3.5	9.7	30.5	89.5	232.7	705.1
1999[4]	3.4	9.4	28.1	82.4	216.8	636.6
Male						
2001	3.5	9.4	29.7	89.2	226.7	587.2
2000[4]	3.2	10.3	30.2	83.8	217.3	551.7
1999[4]	3.4	9.3	28.0	81.3	209.4	515.1
Female						
2001	3.9	9.8	32.9	94.9	255.3	816.6
2000[4]	3.9	9.2	30.8	93.4	241.9	767.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Alzheimer's disease (G30)														
Non-Hispanic White														
Female														
1999[4]	28.4	*	*	*	*	*	*	*	*	*	*	*	*	0.9
Non-Hispanic Black														
Both sexes														
2001	8.7	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	7.7	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	6.7	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	5.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	4.6	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	4.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	11.8	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	10.6	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	9.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Major cardiovascular diseases (I00-I78)														
All origins[3]														
Both sexes														
2001	323.9	14.9	1.9	0.7	1.1	2.2	4.3	7.1	13.0	25.0	48.8	85.6	144.9	234.9
2000[4]	332.9	16.7	1.5	0.7	1.1	2.4	4.3	6.7	12.2	24.2	49.6	88.5	146.7	248.4
1999[4]	340.6	17.0	1.5	0.7	1.0	2.7	4.1	6.5	12.6	25.9	50.0	88.3	148.7	254.8
Male														
2001	306.2	15.3	1.9	0.6	1.1	2.7	5.3	8.8	16.5	33.1	66.5	121.4	204.7	327.6
2000[4]	315.8	17.5	1.6	0.7	1.2	2.8	5.0	8.2	15.3	31.3	68.0	126.8	209.0	350.6
1999[4]	323.0	17.9	1.6	0.8	1.1	3.3	4.8	7.8	16.2	34.2	68.9	125.7	212.5	358.4
Female														
2001	340.9	14.5	1.8	0.8	1.1	1.7	3.2	5.5	9.5	17.0	31.5	50.7	87.6	148.1
2000[4]	349.5	15.9	1.4	0.6	1.0	2.0	3.5	5.2	9.0	17.1	31.5	51.4	87.0	152.8
1999[4]	357.5	16.0	1.4	0.6	1.0	2.1	3.4	5.2	9.1	17.6	31.4	52.1	87.5	158.0
Hispanic														
Both sexes														
2001	96.2	12.0	1.5	0.7	0.9	2.1	2.8	4.4	8.6	14.1	26.7	54.0	103.2	165.7
2000[4]	96.5	13.7	1.3	0.6	0.8	1.9	3.2	4.8	8.2	14.5	29.7	56.1	99.6	178.2
1999[4]	99.2	15.2	1.4	0.6	0.8	2.2	2.9	4.6	8.5	15.6	28.6	59.1	108.5	186.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alzheimer's disease (G30)						
Non-Hispanic White						
Female						
1999[4]	3.4	9.5	28.1	83.2	221.1	685.0
Non-Hispanic Black						
Both sexes						
2001	1.8	8.9	23.8	88.4	186.8	513.2
2000[4]	2.2	7.6	26.5	72.3	173.9	450.9
1999[4]	*	7.0	25.8	65.1	154.1	374.7
Male						
2001	*	9.9	24.8	86.3	188.1	420.4
2000[4]	*	8.0	28.4	65.5	157.9	378.1
1999[4]	*	8.2	28.0	62.8	141.7	320.9
Female						
2001	*	8.1	23.2	89.7	186.2	547.5
2000[4]	*	7.3	25.2	76.4	181.9	478.0
1999[4]	*	6.1	24.4	66.4	160.2	394.9
Major cardiovascular diseases (I00-I78)						
All origins[3]						
Both sexes						
2001	385.7	619.6	1,017.5	1,759.4	3,133.9	7,673.7
2000[4]	407.5	647.9	1,066.0	1,824.4	3,253.7	8,030.3
1999[4]	416.0	678.9	1,111.3	1,888.7	3,356.4	8,197.1
Male						
2001	526.3	821.4	1,325.6	2,223.2	3,774.2	7,936.0
2000[4]	556.9	857.2	1,403.8	2,305.7	3,956.7	8,448.1
1999[4]	568.4	908.7	1,461.9	2,392.7	4,078.9	8,603.2
Female						
2001	258.3	445.9	772.7	1,434.7	2,750.2	7,563.9
2000[4]	272.0	468.5	799.9	1,489.2	2,839.0	7,860.1
1999[4]	278.2	482.2	837.3	1,538.3	2,933.7	8,033.5
Hispanic						
Both sexes						
2001	283.6	490.6	837.6	1,459.3	2,475.8	6,036.2
2000[4]	310.5	501.0	861.4	1,451.4	2,583.3	6,168.2
1999[4]	313.9	543.7	875.1	1,519.7	2,691.9	6,335.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Major cardiovascular diseases (I00-I78)														
Hispanic														
Male														
2001	95.4	12.7	1.4	*	*	2.4	3.5	5.6	11.2	19.0	36.5	77.7	146.6	233.8
2000[4]	96.0	13.7	1.4	*	1.2	2.4	3.7	6.0	10.4	18.2	40.0	80.2	144.5	257.4
1999[4]	99.2	12.7	1.4	*	*	2.5	3.5	6.1	11.4	20.8	40.8	83.2	160.7	259.8
Female														
2001	97.1	11.2	1.6	*	*	1.7	1.8	3.0	5.6	8.7	16.5	30.2	61.7	104.1
2000[4]	96.9	13.8	*	*	*	*	2.6	3.4	5.7	10.5	18.9	32.0	56.8	106.6
1999[4]	99.1	17.9	1.4	*	*	1.8	2.1	2.8	5.2	10.0	15.9	35.3	58.9	119.4
Non-Hispanic[5]														
Both sexes														
2001	356.7	15.5	1.9	0.7	1.1	2.2	4.5	7.7	13.8	26.6	51.3	88.3	147.8	239.3
2000[4]	365.6	17.3	1.6	0.7	1.1	2.5	4.5	7.0	12.8	25.4	51.5	91.1	149.9	252.6
1999[4]	372.7	17.4	1.5	0.7	1.1	2.8	4.4	6.8	13.3	27.1	52.0	90.6	151.3	258.7
Male														
2001	338.1	15.8	2.0	0.6	1.2	2.7	5.6	9.5	17.4	35.2	69.8	125.4	208.7	333.1
2000[4]	347.7	18.3	1.7	0.8	1.2	2.9	5.4	8.6	16.2	33.0	70.8	130.7	213.2	355.8
1999[4]	354.2	19.1	1.7	0.8	1.1	3.5	5.1	8.1	17.0	35.9	71.6	129.0	215.8	363.7
Female														
2001	374.3	15.2	1.8	0.8	1.1	1.7	3.4	6.0	10.1	18.2	33.1	52.6	89.5	151.1
2000[4]	382.6	16.3	1.5	0.6	1.1	2.1	3.7	5.5	9.5	18.0	32.7	53.0	89.2	155.8
1999[4]	390.2	15.5	1.4	0.6	1.0	2.1	3.7	5.6	9.7	18.5	33.0	53.4	89.4	160.3
Non-Hispanic White														
Both sexes														
2001	385.8	12.1	1.5	0.6	1.0	1.9	3.5	6.2	11.3	22.5	43.3	76.0	129.1	215.0
2000[4]	395.5	14.6	1.4	0.6	0.9	2.1	3.5	5.6	10.4	21.4	43.9	78.1	132.4	228.1
1999[4]	402.7	14.2	1.3	0.7	0.9	2.4	3.5	5.5	10.9	22.8	43.8	77.8	133.6	232.2
Male														
2001	363.0	13.0	1.6	0.6	1.1	2.2	4.5	7.8	14.6	30.4	61.8	111.7	188.0	305.3
2000[4]	373.6	15.5	1.6	0.7	1.1	2.4	4.3	7.0	13.7	28.9	62.9	117.3	193.1	326.7
1999[4]	379.8	16.1	1.5	0.8	0.9	2.9	4.0	6.5	14.3	30.9	62.8	115.3	195.1	333.2
Female														
2001	407.6	11.0	1.5	0.7	0.9	1.5	2.5	4.6	8.0	14.6	24.8	40.6	71.4	128.4
2000[4]	416.5	13.6	1.1	0.4	0.8	1.7	2.8	4.2	7.0	13.8	25.0	39.4	73.0	133.8
1999[4]	424.7	12.1	1.1	0.6	0.9	1.9	3.1	4.4	7.4	14.6	24.9	40.8	73.5	135.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Major cardiovascular diseases (I00-I78)						
Hispanic						
Male						
2001	383.5	672.6	1,084.7	1,808.6	3,004.8	6,155.6
2000[4]	426.9	658.5	1,145.9	1,849.2	3,116.9	6,304.3
1999[4]	427.7	730.3	1,182.4	1,968.4	3,137.4	6,595.2
Female						
2001	197.2	343.2	649.1	1,211.3	2,148.2	5,977.9
2000[4]	210.2	373.5	645.9	1,169.6	2,258.6	6,101.4
1999[4]	215.9	392.8	643.9	1,203.1	2,426.1	6,210.9
Non-Hispanic[5]						
Both sexes						
2001	391.7	626.6	1,025.2	1,768.6	3,152.1	7,715.2
2000[4]	412.4	655.5	1,073.4	1,835.5	3,270.0	8,076.4
1999[4]	421.2	684.9	1,119.5	1,898.7	3,370.5	8,239.8
Male						
2001	533.8	828.4	1,335.1	2,236.3	3,795.4	7,990.6
2000[4]	563.0	866.4	1,411.9	2,318.5	3,979.0	8,513.1
1999[4]	574.7	916.0	1,469.7	2,402.8	4,101.3	8,657.6
Female						
2001	262.3	452.1	778.4	1,441.4	2,767.2	7,600.7
2000[4]	275.5	473.9	806.2	1,499.3	2,852.2	7,899.8
1999[4]	281.8	486.4	845.2	1,548.4	2,943.2	8,072.5
Non-Hispanic White						
Both sexes						
2001	359.1	584.0	984.9	1,727.4	3,139.5	7,811.4
2000[4]	378.9	617.5	1,032.1	1,791.9	3,258.1	8,179.0
1999[4]	388.2	648.2	1,074.2	1,856.0	3,358.3	8,342.9
Male						
2001	498.8	785.3	1,292.9	2,203.7	3,797.2	8,116.3
2000[4]	527.5	829.2	1,372.2	2,284.2	3,984.9	8,650.5
1999[4]	537.4	879.0	1,426.0	2,370.6	4,111.9	8,776.1
Female						
2001	229.5	406.1	734.6	1,390.3	2,742.5	7,684.6
2000[4]	241.2	431.4	758.3	1,445.9	2,826.1	7,988.9
1999[4]	249.9	445.8	793.8	1,495.0	2,913.7	8,170.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Major cardiovascular diseases (I00-I78)														
Non-Hispanic Black														
Both sexes														
2001	289.7	30.1	3.7	1.0	1.6	4.1	9.8	17.1	29.5	53.3	108.4	185.9	313.3	478.6
2000[4]	294.2	28.8	2.5	1.0	1.9	4.8	9.5	15.7	28.8	52.8	106.7	195.9	310.0	497.1
1999[4]	298.5	30.2	2.6	1.1	1.9	5.0	9.3	15.1	28.0	56.8	112.2	192.5	317.0	517.6
Male														
2001	281.2	29.7	3.8	*	1.6	5.2	11.7	21.0	36.4	67.9	132.7	242.7	413.4	637.0
2000[4]	284.3	30.0	2.3	*	1.9	5.3	10.8	18.9	33.9	63.5	133.3	252.1	415.3	680.4
1999[4]	291.1	31.5	2.7	*	2.2	6.5	11.2	18.6	34.5	71.6	141.1	251.7	431.3	697.1
Female														
2001	297.4	30.6	3.6	1.3	1.5	2.9	7.9	13.6	23.3	40.4	86.9	136.8	228.2	348.9
2000[4]	303.1	27.5	2.9	*	2.0	4.4	8.4	12.9	24.2	43.3	83.2	147.3	220.3	347.1
1999[4]	305.1	28.9	2.4	*	1.6	3.3	7.4	11.9	22.2	43.8	86.9	141.5	220.2	371.1
Diseases of heart (I00-I09,I11,I13, I20-I51)														
All origins[3]														
Both sexes														
2001	245.8	11.9	1.5	0.5	0.8	1.7	3.3	5.6	10.2	19.9	39.1	69.3	119.5	194.2
2000[4]	252.6	13.0	1.2	0.5	0.8	2.0	3.3	5.2	9.5	19.1	39.4	71.2	120.4	205.5
1999[4]	259.9	13.8	1.2	0.6	0.8	2.3	3.3	5.1	10.1	20.6	40.1	71.7	123.5	211.0
Male														
2001	242.5	11.8	1.5	0.4	0.9	2.1	4.2	7.0	13.4	27.4	55.8	102.6	175.2	279.7
2000[4]	249.8	13.3	1.4	0.6	0.9	2.4	4.0	6.5	12.4	26.0	57.0	107.1	178.1	298.9
1999[4]	257.0	13.9	1.3	0.6	0.9	2.9	3.9	6.3	13.2	28.5	58.1	107.0	182.9	306.5
Female														
2001	249.0	12.0	1.4	0.6	0.8	1.3	2.4	4.1	6.9	12.4	22.6	37.0	66.1	114.2
2000[4]	255.3	12.5	1.0	0.4	0.7	1.6	2.6	3.9	6.5	12.2	22.2	36.4	65.0	118.1
1999[4]	262.6	13.7	1.1	0.5	0.7	1.7	2.6	3.9	6.9	12.6	22.4	37.5	66.6	121.7
Hispanic														
Both sexes														
2001	73.3	9.7	1.1	*	0.6	1.4	1.9	3.2	6.0	10.1	18.1	38.3	77.1	129.3
2000[4]	73.1	11.2	1.0	*	0.7	1.5	2.2	3.3	5.4	10.1	20.7	39.9	76.8	141.1
1999[4]	76.2	12.7	1.2	*	*	1.7	2.0	2.9	6.3	11.7	20.0	42.6	84.9	146.0
Male														
2001	74.6	10.3	*	*	*	1.6	2.5	4.4	8.0	14.6	26.6	58.7	115.3	191.0
2000[4]	74.7	11.2	1.4	*	*	2.1	2.7	4.3	7.2	13.4	29.9	60.8	115.6	208.7
1999[4]	77.7	10.3	1.4	*	*	1.9	2.4	4.4	8.7	16.5	30.0	63.4	130.7	213.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Major cardiovascular diseases (I00-I78)						
Non-Hispanic Black						
Both sexes						
2001	725.1	1,073.1	1,567.7	2,444.2	3,650.2	7,366.6
2000[4]	748.1	1,078.7	1,639.8	2,519.2	3,750.0	7,576.5
1999[4]	753.4	1,106.8	1,719.8	2,571.4	3,809.5	7,624.4
Male						
2001	947.4	1,361.6	2,006.9	2,966.0	4,316.4	7,413.8
2000[4]	967.2	1,346.6	2,057.4	3,018.8	4,434.6	7,755.5
1999[4]	1,003.1	1,415.5	2,159.6	3,060.6	4,425.3	7,952.2
Female						
2001	550.3	860.0	1,275.0	2,132.8	3,310.8	7,349.1
2000[4]	575.2	881.0	1,362.6	2,218.0	3,405.4	7,510.0
1999[4]	557.6	878.0	1,427.9	2,276.7	3,504.3	7,501.9
Diseases of heart (I00-I09,I11,I13, I20-I51)						
All origins[3]						
Both sexes						
2001	314.1	494.4	787.7	1,321.2	2,309.0	5,664.2
2000[4]	330.6	515.8	826.7	1,372.3	2,392.1	5,926.1
1999[4]	341.9	547.1	867.9	1,427.8	2,484.7	6,063.0
Male						
2001	440.9	672.2	1,054.4	1,732.2	2,882.7	6,040.5
2000[4]	463.9	702.0	1,119.7	1,787.5	3,012.3	6,430.0
1999[4]	478.2	752.8	1,174.1	1,865.5	3,123.3	6,561.6
Female						
2001	199.0	341.4	575.8	1,033.5	1,965.3	5,506.8
2000[4]	209.8	356.3	596.0	1,083.0	2,026.2	5,720.9
1999[4]	218.6	371.2	628.7	1,123.5	2,111.0	5,862.1
Hispanic						
Both sexes						
2001	225.5	391.0	646.2	1,097.2	1,866.0	4,607.5
2000[4]	237.3	388.4	654.8	1,096.6	1,940.5	4,701.8
1999[4]	251.5	425.7	676.2	1,167.9	2,013.5	4,888.6
Male						
2001	311.8	547.1	860.1	1,385.1	2,307.5	4,784.3
2000[4]	338.5	525.4	892.6	1,411.5	2,374.9	4,897.5
1999[4]	347.6	588.1	933.4	1,526.8	2,398.7	5,012.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Diseases of heart (I00-I09,I11,I13, I20-I51)														
Hispanic														
Female														
2001	71.8	9.0	*	*	*	*	*	1.9	3.7	5.2	9.2	17.8	40.7	73.5
2000[4]	71.5	11.2	*	*	*	*	1.6	2.1	3.3	6.4	11.0	19.1	39.8	80.0
1999[4]	74.6	15.2	*	*	*	1.5	1.5	*	3.6	6.5	9.6	21.9	41.5	84.9
Non-Hispanic[5]														
Both sexes														
2001	270.7	12.3	1.5	0.5	0.9	1.7	3.6	6.1	10.9	21.3	41.4	72.1	122.6	198.4
2000[4]	277.3	13.3	1.2	0.5	0.8	2.1	3.5	5.6	10.2	20.3	41.3	73.9	123.4	209.4
1999[4]	284.3	14.0	1.2	0.6	0.8	2.4	3.5	5.5	10.7	21.6	42.1	74.1	126.1	214.7
Male														
2001	268.0	12.0	1.7	0.4	0.9	2.2	4.6	7.6	14.4	29.3	59.2	106.6	179.5	285.1
2000[4]	275.2	13.7	1.4	0.6	0.9	2.4	4.3	7.0	13.4	27.7	59.7	111.1	182.4	304.2
1999[4]	282.0	14.8	1.3	0.6	0.9	3.0	4.2	6.6	14.0	30.0	60.9	110.6	186.3	311.6
Female														
2001	273.3	12.7	1.4	0.6	0.8	1.3	2.6	4.5	7.5	13.4	24.2	38.7	68.0	117.0
2000[4]	279.3	12.8	1.1	0.4	0.7	1.7	2.7	4.2	7.0	12.9	23.3	37.9	66.9	120.5
1999[4]	286.4	13.3	1.1	0.5	0.8	1.7	2.8	4.3	7.4	13.4	23.7	38.8	68.3	124.0
Non-Hispanic White														
Both sexes														
2001	294.1	9.8	1.2	0.4	0.8	1.4	2.7	4.8	9.0	18.4	36.2	64.0	110.9	181.6
2000[4]	301.3	11.4	1.1	0.4	0.7	1.7	2.7	4.3	8.3	17.6	36.5	65.8	112.1	192.8
1999[4]	308.2	11.3	1.0	0.5	0.7	2.1	2.7	4.3	8.8	18.6	36.5	65.8	114.4	196.2
Male														
2001	289.5	10.3	1.3	0.4	0.8	1.7	3.5	6.3	12.1	26.0	53.9	97.9	166.9	265.9
2000[4]	297.5	12.0	1.2	0.5	0.8	1.9	3.3	5.5	11.4	24.9	54.8	102.8	169.3	284.7
1999[4]	304.0	12.4	1.1	0.5	0.7	2.5	3.2	5.3	11.9	26.4	54.9	101.8	172.6	289.8
Female														
2001	298.4	9.2	1.1	0.5	0.7	1.1	1.9	3.4	5.9	10.8	18.5	30.5	56.0	100.8
2000[4]	304.9	10.8	0.9	0.3	0.5	1.4	2.0	3.1	5.2	10.2	18.2	29.1	56.0	105.0
1999[4]	312.3	10.2	0.8	0.5	0.7	1.6	2.2	3.4	5.7	10.7	18.2	30.3	57.3	106.9
Non-Hispanic Black														
Both sexes														
2001	215.5	22.9	3.1	0.8	1.3	3.5	7.8	13.5	23.5	41.3	81.6	140.3	236.6	375.4
2000[4]	218.6	20.5	2.0	0.9	1.4	4.2	7.8	13.4	22.3	39.4	78.8	145.2	236.1	386.8
1999[4]	223.8	24.8	2.1	1.0	1.4	4.3	8.0	12.3	22.5	43.8	84.7	145.9	244.9	407.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13, I20-I51)						
Hispanic						
Female						
2001	150.9	264.6	483.0	892.8	1,592.5	4,521.1
2000[4]	150.1	277.5	474.6	873.7	1,676.2	4,605.8
1999[4]	168.6	294.4	482.7	914.6	1,783.7	4,828.9
Non-Hispanic[5]						
Both sexes						
2001	319.3	499.9	793.7	1,327.9	2,320.3	5,688.6
2000[4]	335.7	522.3	833.0	1,380.1	2,402.3	5,953.6
1999[4]	346.6	552.7	874.4	1,434.0	2,493.7	6,086.3
Male						
2001	448.0	678.1	1,061.9	1,743.4	2,898.2	6,077.8
2000[4]	470.1	710.1	1,127.0	1,798.4	3,028.8	6,474.6
1999[4]	484.3	759.8	1,180.7	1,873.1	3,140.2	6,602.0
Female						
2001	202.2	345.9	580.0	1,037.2	1,974.4	5,526.8
2000[4]	213.4	360.8	600.9	1,089.1	2,033.1	5,742.9
1999[4]	221.4	374.9	634.5	1,129.0	2,115.7	5,879.8
Non-Hispanic White						
Both sexes						
2001	298.4	471.8	769.0	1,302.6	2,318.8	5,776.0
2000[4]	313.1	497.3	805.7	1,353.2	2,402.0	6,047.9
1999[4]	324.5	528.0	844.9	1,407.9	2,492.7	6,176.6
Male						
2001	426.1	650.7	1,037.0	1,726.7	2,911.3	6,195.4
2000[4]	446.8	686.5	1,102.9	1,780.9	3,046.4	6,606.6
1999[4]	459.4	735.6	1,154.2	1,858.9	3,162.2	6,713.7
Female						
2001	180.0	313.7	551.2	1,002.4	1,961.1	5,601.6
2000[4]	189.2	330.9	566.6	1,052.6	2,019.1	5,822.7
1999[4]	199.5	346.0	598.4	1,091.5	2,097.7	5,962.4
Non-Hispanic Black						
Both sexes						
2001	554.8	820.9	1,173.2	1,795.1	2,634.8	5,307.7
2000[4]	579.6	826.1	1,242.2	1,852.8	2,693.6	5,424.6
1999[4]	587.8	861.8	1,297.7	1,889.6	2,762.0	5,514.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Diseases of heart (I00-I09,I11,I13, I20-I51)														
Non-Hispanic Black														
Male														
2001	215.7	20.7	3.4	*	1.4	4.5	10.0	16.7	30.3	54.1	104.2	188.8	320.2	510.4
2000[4]	217.6	20.1	2.0	*	1.5	4.9	9.2	16.4	27.6	50.0	102.5	195.0	325.5	540.6
1999[4]	224.9	24.6	2.3	*	1.7	6.0	9.7	15.6	28.4	57.9	111.5	197.0	341.6	565.4
Female														
2001	215.4	25.2	2.9	*	*	2.4	5.8	10.7	17.5	30.0	61.6	98.4	165.5	264.8
2000[4]	219.5	20.8	2.0	*	1.4	3.4	6.5	10.6	17.5	30.0	58.0	102.0	159.9	260.9
1999[4]	222.8	25.0	1.9	*	*	2.6	6.5	9.4	17.2	31.4	61.1	101.9	163.0	278.2
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)														
All origins[3]														
Both sexes														
2001	1.2	*	*	*	*	*	*	*	0.1	0.2	0.4	0.5	0.7	1.2
2000[4]	1.3	*	*	*	*	*	*	*	0.1	0.2	0.4	0.6	0.8	1.4
1999[4]	1.3	*	*	*	*	*	*	*	0.1	0.2	0.3	0.5	0.9	1.4
Male														
2001	0.7	*	*	*	*	*	*	*	*	*	0.3	0.4	0.5	1.1
2000[4]	0.8	*	*	*	*	*	*	*	*	0.2	0.3	0.5	0.6	1.1
1999[4]	0.8	*	*	*	*	*	*	*	*	*	0.3	0.4	0.6	1.0
Female														
2001	1.7	*	*	*	*	*	*	*	*	0.2	0.4	0.6	0.9	1.3
2000[4]	1.8	*	*	*	*	*	*	*	*	0.2	0.5	0.7	0.9	1.6
1999[4]	1.9	*	*	*	*	*	*	*	*	0.2	0.4	0.6	1.1	1.8
Hispanic														
Both sexes														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13, I20-I51)						
Non-Hispanic Black						
Male						
2001	740.5	1,053.7	1,531.1	2,230.3	3,202.0	5,495.4
2000[4]	763.9	1,047.1	1,582.4	2,260.1	3,247.6	5,677.2
1999[4]	797.1	1,122.5	1,658.3	2,283.9	3,268.7	5,877.1
Female						
2001	408.8	649.0	934.8	1,535.3	2,345.9	5,238.2
2000[4]	434.1	663.2	1,016.4	1,607.3	2,414.7	5,330.8
1999[4]	423.7	668.6	1,058.5	1,652.1	2,510.8	5,379.3
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)						
All origins[3]						
Both sexes						
2001	1.7	3.2	5.0	8.1	11.4	17.7
2000[4]	2.2	3.4	5.6	8.1	11.6	16.8
1999[4]	2.1	3.5	5.8	8.4	12.6	18.0
Male						
2001	1.2	2.4	3.2	6.0	7.7	12.8
2000[4]	1.8	2.2	3.6	5.8	7.8	12.2
1999[4]	1.4	2.4	4.7	5.4	8.9	10.3
Female						
2001	2.2	3.9	6.4	9.5	13.7	19.8
2000[4]	2.5	4.5	7.1	9.7	13.9	18.7
1999[4]	2.7	4.5	6.7	10.4	14.7	21.1
Hispanic						
Both sexes						
2001	*	3.6	4.9	*	*	*
2000[4]	*	*	5.9	*	*	*
1999[4]	*	4.3	*	7.1	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	7.3	*	*	*
2000[4]	*	*	7.4	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)														
Hispanic														
Female														
1999[4]	0.8	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	1.3	*	*	*	*	*	*	*	0.1	0.2	0.4	0.5	0.7	1.2
2000[4]	1.4	*	*	*	*	*	*	*	*	0.2	0.4	0.6	0.8	1.4
1999[4]	1.4	*	*	*	*	*	*	*	0.2	0.2	0.3	0.5	0.8	1.4
Male														
2001	0.8	*	*	*	*	*	*	*	*	*	0.3	0.5	0.5	1.1
2000[4]	0.9	*	*	*	*	*	*	*	*	0.2	0.3	0.5	0.6	1.1
1999[4]	0.8	*	*	*	*	*	*	*	*	*	0.3	0.4	0.6	1.0
Female														
2001	1.8	*	*	*	*	*	*	*	*	0.2	0.4	0.6	0.9	1.3
2000[4]	1.9	*	*	*	*	*	*	*	*	*	0.5	0.7	0.9	1.6
1999[4]	2.0	*	*	*	*	*	*	*	*	0.2	0.4	0.6	1.0	1.8
Non-Hispanic White														
Both sexes														
2001	1.5	*	*	*	*	*	*	*	*	0.2	0.3	0.4	0.6	1.1
2000[4]	1.6	*	*	*	*	*	*	*	*	0.2	0.3	0.5	0.7	1.3
1999[4]	1.6	*	*	*	*	*	*	*	*	0.1	0.3	0.5	0.7	1.4
Male														
2001	0.9	*	*	*	*	*	*	*	*	*	0.3	0.4	0.4	1.0
2000[4]	0.9	*	*	*	*	*	*	*	*	*	0.3	0.5	0.6	1.1
1999[4]	0.9	*	*	*	*	*	*	*	*	*	0.2	0.4	0.6	1.0
Female														
2001	2.0	*	*	*	*	*	*	*	*	*	0.3	0.4	0.7	1.1
2000[4]	2.1	*	*	*	*	*	*	*	*	*	0.4	0.6	0.8	1.4
1999[4]	2.2	*	*	*	*	*	*	*	*	*	0.3	0.5	0.9	1.8
Non-Hispanic Black														
Both sexes														
2001	0.8	*	*	*	*	*	*	*	*	*	0.9	1.1	1.7	2.2
2000[4]	0.7	*	*	*	*	*	*	*	*	*	0.9	1.0	1.1	2.0
1999[4]	0.8	*	*	*	*	*	*	*	*	*	*	0.9	1.7	2.3
Male														
2001	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)						
Hispanic						
Female						
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	1.7	3.2	5.0	8.2	11.6	18.0
2000[4]	2.2	3.4	5.5	8.3	11.8	17.1
1999[4]	2.1	3.5	5.9	8.4	12.7	18.3
Male						
2001	1.2	2.3	3.3	6.0	7.8	13.1
2000[4]	1.8	2.3	3.5	5.9	8.1	12.5
1999[4]	1.3	2.4	4.8	5.6	9.1	10.7
Female						
2001	2.1	4.0	6.3	9.8	13.9	20.0
2000[4]	2.5	4.5	7.1	10.0	13.9	19.0
1999[4]	2.8	4.4	6.7	10.3	14.8	21.3
Non-Hispanic White						
Both sexes						
2001	1.7	3.1	5.0	8.6	12.4	19.0
2000[4]	2.2	3.6	5.6	8.9	12.5	18.0
1999[4]	2.0	3.6	6.1	8.8	13.3	19.2
Male						
2001	1.2	2.2	3.4	6.3	8.2	13.7
2000[4]	1.9	2.3	3.5	6.3	8.7	13.0
1999[4]	1.3	2.6	5.0	5.7	9.7	11.6
Female						
2001	2.1	4.0	6.3	10.2	14.9	21.2
2000[4]	2.4	4.7	7.3	10.7	14.8	20.0
1999[4]	2.7	4.4	7.0	11.0	15.5	22.3
Non-Hispanic Black						
Both sexes						
2001	2.0	3.4	5.2	5.0	*	6.9
2000[4]	2.1	3.5	3.7	3.8	*	8.6
1999[4]	2.1	2.7	4.1	4.2	*	8.4
Male						
2001	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)														
Non-Hispanic Black														
Male														
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	1.2	*	*	*	*	*	*	*	*	*	*	*	2.3	2.8
2000[4]	1.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.0	*	*	*	*	*	*	*	*	*	*	*	*	2.8
Hypertensive heart disease (I11)														
All origins[3]														
Both sexes														
2001	8.7	*	*	*	*	*	*	0.3	0.9	1.6	3.2	5.1	7.7	9.7
2000[4]	8.4	*	*	*	*	*	0.1	0.4	0.7	1.4	2.9	4.7	6.6	9.3
1999[4]	8.1	*	*	*	*	*	0.1	0.3	0.8	1.5	2.8	4.5	6.2	9.6
Male														
2001	7.5	*	*	*	*	*	*	0.4	1.2	2.1	4.5	7.0	10.3	12.8
2000[4]	7.2	*	*	*	*	*	0.2	0.4	0.9	1.9	3.9	6.3	9.4	12.1
1999[4]	6.8	*	*	*	*	*	*	0.4	1.0	2.0	3.9	6.1	8.1	12.7
Female														
2001	9.8	*	*	*	*	*	*	0.3	0.5	1.1	2.0	3.2	5.1	6.8
2000[4]	9.7	*	*	*	*	*	*	0.3	0.5	1.0	2.0	3.2	3.8	6.6
1999[4]	9.4	*	*	*	*	*	*	*	0.5	1.0	1.7	3.0	4.3	6.8
Hispanic														
Both sexes														
2001	3.3	*	*	*	*	*	*	*	*	1.4	2.1	3.1	7.2	8.5
2000[4]	2.8	*	*	*	*	*	*	*	*	0.9	1.6	3.1	4.6	5.8
1999[4]	2.7	*	*	*	*	*	*	*	*	1.1	1.4	2.6	4.4	8.4
Male														
2001	3.4	*	*	*	*	*	*	*	*	2.4	3.0	4.5	10.4	11.4
2000[4]	2.7	*	*	*	*	*	*	*	*	*	2.1	4.7	6.3	8.1
1999[4]	2.6	*	*	*	*	*	*	*	*	1.7	1.9	4.1	5.6	11.8
Female														
2001	3.1	*	*	*	*	*	*	*	*	*	*	*	4.1	6.0
2000[4]	2.9	*	*	*	*	*	*	*	*	*	*	*	3.0	*
1999[4]	2.9	*	*	*	*	*	*	*	*	*	*	*	3.2	5.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)						
Non-Hispanic Black						
Male						
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	3.8	7.3	6.5	*	*
2000[4]	*	4.5	*	*	*	*
1999[4]	*	*	4.8	*	*	11.1
Hypertensive heart disease (I11)						
All origins[3]						
Both sexes						
2001	12.5	16.3	23.5	36.6	66.0	189.5
2000[4]	13.2	16.5	23.0	36.3	66.1	193.1
1999[4]	11.9	16.3	22.1	36.4	64.7	184.9
Male						
2001	16.5	20.5	27.1	38.9	63.5	134.4
2000[4]	17.4	20.7	27.2	37.3	61.8	139.1
1999[4]	15.0	19.5	24.6	39.4	59.2	133.6
Female						
2001	8.9	12.7	20.6	35.1	67.5	212.6
2000[4]	9.3	12.9	19.6	35.7	68.6	215.1
1999[4]	9.2	13.5	20.2	34.3	67.9	205.5
Hispanic						
Both sexes						
2001	11.8	16.8	26.3	37.7	63.1	151.9
2000[4]	11.3	13.3	24.7	35.5	65.2	143.3
1999[4]	9.5	16.4	17.3	33.2	65.8	152.2
Male						
2001	17.1	22.3	31.6	38.9	66.2	144.7
2000[4]	14.1	16.4	30.1	43.6	64.8	112.9
1999[4]	13.4	20.1	21.9	34.3	59.7	107.8
Female						
2001	7.3	12.3	22.3	36.8	61.1	155.5
2000[4]	8.9	10.9	20.6	29.8	65.4	158.3
1999[4]	6.2	13.4	13.8	32.5	69.4	173.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Hypertensive heart disease (I11)														
Non-Hispanic[5]														
Both sexes														
2001	9.4	*	*	*	*	*	*	0.4	1.0	1.6	3.4	5.2	7.6	9.7
2000[4]	9.2	*	*	*	*	*	0.2	0.4	0.8	1.5	3.0	4.8	6.7	9.5
1999[4]	8.8	*	*	*	*	*	0.2	0.3	0.8	1.6	2.9	4.7	6.3	9.6
Male														
2001	8.1	*	*	*	*	*	*	0.4	1.4	2.1	4.6	7.2	10.2	12.7
2000[4]	7.8	*	*	*	*	*	0.3	0.4	1.0	1.9	4.0	6.4	9.6	12.3
1999[4]	7.4	*	*	*	*	*	*	0.4	1.1	2.1	4.1	6.2	8.3	12.6
Female														
2001	10.7	*	*	*	*	*	*	0.3	0.6	1.1	2.1	3.3	5.2	6.8
2000[4]	10.6	*	*	*	*	*	*	0.3	0.5	1.0	2.1	3.3	3.8	6.8
1999[4]	10.2	*	*	*	*	*	*	*	0.6	1.1	1.8	3.2	4.3	6.8
Non-Hispanic White														
Both sexes														
2001	8.6	*	*	*	*	*	*	0.2	0.6	1.0	2.1	3.3	5.0	6.7
2000[4]	8.3	*	*	*	*	*	*	0.2	0.4	0.9	2.0	3.1	4.3	6.2
1999[4]	7.9	*	*	*	*	*	*	*	0.5	0.9	1.8	2.9	4.0	6.3
Male														
2001	6.9	*	*	*	*	*	*	*	0.9	1.5	3.1	4.8	7.1	9.4
2000[4]	6.6	*	*	*	*	*	*	*	0.6	1.3	2.9	4.5	6.5	8.5
1999[4]	6.2	*	*	*	*	*	*	*	0.8	1.2	2.7	4.1	5.7	8.6
Female														
2001	10.2	*	*	*	*	*	*	*	*	0.6	1.0	1.8	2.9	4.2
2000[4]	9.9	*	*	*	*	*	*	*	*	0.5	1.2	1.7	2.1	4.0
1999[4]	9.6	*	*	*	*	*	*	*	*	0.5	0.8	1.6	2.4	4.0
Non-Hispanic Black														
Both sexes														
2001	16.5	*	*	*	*	*	*	1.1	3.3	5.2	11.8	18.9	28.5	35.7
2000[4]	16.6	*	*	*	*	*	*	1.4	2.9	5.1	9.9	18.1	25.9	38.4
1999[4]	16.0	*	*	*	*	*	*	1.0	2.7	5.9	10.4	18.2	25.3	38.4
Male														
2001	16.3	*	*	*	*	*	*	*	4.2	6.0	15.1	25.5	36.5	45.2
2000[4]	16.1	*	*	*	*	*	*	1.6	3.7	6.0	12.6	22.0	36.5	48.8
1999[4]	15.6	*	*	*	*	*	*	*	2.9	7.6	13.3	22.9	32.0	50.0
Female														
2001	16.7	*	*	*	*	*	*	*	2.4	4.6	8.9	13.2	21.7	28.0
2000[4]	17.1	*	*	*	*	*	*	*	2.2	4.3	7.6	14.8	16.9	29.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart disease (I11)						
Non-Hispanic[5]						
Both sexes						
2001	12.4	16.3	23.2	36.4	66.0	190.3
2000[4]	13.3	16.5	22.7	36.1	66.0	194.3
1999[4]	12.0	16.1	22.2	36.4	64.3	185.3
Male						
2001	16.2	20.4	26.5	38.6	63.4	133.4
2000[4]	17.6	20.6	26.7	36.6	61.5	139.6
1999[4]	14.8	19.3	24.4	39.4	58.8	134.2
Female						
2001	9.0	12.7	20.5	34.8	67.6	214.0
2000[4]	9.3	13.0	19.5	35.8	68.6	216.5
1999[4]	9.4	13.4	20.4	34.3	67.6	205.8
Non-Hispanic White						
Both sexes						
2001	8.8	11.8	17.9	30.4	58.9	180.4
2000[4]	9.1	12.2	17.0	29.7	58.0	182.5
1999[4]	8.5	11.7	16.7	29.6	56.8	175.2
Male						
2001	12.0	15.4	21.1	32.5	55.5	122.0
2000[4]	12.4	15.6	20.7	30.2	52.9	129.5
1999[4]	10.9	13.9	18.7	32.1	51.2	123.8
Female						
2001	5.8	8.6	15.3	28.9	60.9	204.7
2000[4]	6.0	9.2	14.0	29.4	61.0	203.9
1999[4]	6.3	9.7	15.1	27.9	60.1	195.7
Non-Hispanic Black						
Both sexes						
2001	44.1	56.4	78.2	106.1	155.9	328.7
2000[4]	48.9	55.6	82.1	111.8	165.8	344.1
1999[4]	42.2	57.4	79.5	114.9	161.4	308.5
Male						
2001	56.4	71.1	91.2	116.6	173.3	288.7
2000[4]	65.1	71.2	95.7	117.1	179.4	275.9
1999[4]	52.5	72.8	90.7	130.5	166.5	266.2
Female						
2001	34.3	45.6	69.5	99.9	147.0	343.5
2000[4]	36.1	44.0	73.1	108.6	159.0	369.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Hypertensive heart disease (I11)														
Non-Hispanic Black														
Female														
1999[4]	16.4	*	*	*	*	*	*	*	2.6	4.5	7.9	14.1	19.7	29.0
Hypertensive heart and renal disease (I13)														
All origins[3]														
Both sexes														
2001	1.0	*	*	*	*	*	*	*	*	0.2	0.3	0.3	0.4	0.8
2000[4]	1.0	*	*	*	*	*	*	*	*	0.1	0.3	0.3	0.4	0.8
1999[4]	1.2	*	*	*	*	*	*	*	*	0.2	0.3	0.4	0.4	0.9
Male														
2001	0.8	*	*	*	*	*	*	*	*	0.2	0.4	0.4	0.6	0.9
2000[4]	0.8	*	*	*	*	*	*	*	*	*	0.3	0.4	0.5	0.9
1999[4]	1.0	*	*	*	*	*	*	*	*	0.2	0.4	0.5	0.6	1.0
Female														
2001	1.1	*	*	*	*	*	*	*	*	0.2	*	*	0.3	0.7
2000[4]	1.1	*	*	*	*	*	*	*	*	*	0.2	0.2	0.4	0.7
1999[4]	1.4	*	*	*	*	*	*	*	*	*	0.3	0.3	0.3	0.7
Hispanic														
Both sexes														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	1.1	*	*	*	*	*	*	*	*	0.2	0.3	0.3	0.4	0.8
2000[4]	1.1	*	*	*	*	*	*	*	*	0.1	0.3	0.3	0.4	0.8
1999[4]	1.3	*	*	*	*	*	*	*	*	0.2	0.3	0.4	0.5	0.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart disease (I11)						
Non-Hispanic Black						
Female						
1999[4]	34.1	46.1	72.1	105.4	158.9	324.2
Hypertensive heart and renal disease (I13)						
All origins[3]						
Both sexes						
2001	1.2	1.8	2.8	5.0	9.3	23.3
2000[4]	1.0	1.6	3.0	5.1	9.8	24.3
1999[4]	1.4	2.1	3.0	6.6	12.3	29.1
Male						
2001	1.2	2.1	2.9	5.2	9.4	24.2
2000[4]	1.0	1.7	3.4	5.0	10.8	27.8
1999[4]	1.6	2.3	3.2	7.3	13.3	31.3
Female						
2001	1.3	1.5	2.8	4.9	9.3	22.9
2000[4]	1.0	1.5	2.7	5.1	9.3	22.9
1999[4]	1.2	2.0	2.9	6.1	11.7	28.2
Hispanic						
Both sexes						
2001	*	*	*	*	11.4	26.3
2000[4]	*	*	4.2	*	*	25.2
1999[4]	*	*	*	6.5	*	23.2
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	24.2
2000[4]	*	*	*	*	*	26.7
1999[4]	*	*	*	*	*	20.8
Non-Hispanic[5]						
Both sexes						
2001	1.2	1.7	2.7	5.0	9.2	23.1
2000[4]	1.0	1.6	3.0	5.0	9.8	24.2
1999[4]	1.4	2.0	3.0	6.6	12.3	29.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Hypertensive heart and renal disease (I13)														
Non-Hispanic[5]														
Male														
2001	0.9	*	*	*	*	*	*	*	*	*	0.4	0.5	0.6	0.9
2000[4]	0.9	*	*	*	*	*	*	*	*	*	0.3	0.4	0.5	0.9
1999[4]	1.1	*	*	*	*	*	*	*	*	0.2	0.4	0.5	0.6	1.0
Female														
2001	1.2	*	*	*	*	*	*	*	*	*	*	*	0.3	0.7
2000[4]	1.2	*	*	*	*	*	*	*	*	*	0.2	0.2	0.4	0.7
1999[4]	1.5	*	*	*	*	*	*	*	*	*	0.3	0.3	0.3	0.7
Non-Hispanic White														
Both sexes														
2001	0.9	*	*	*	*	*	*	*	*	*	*	*	0.2	0.3
2000[4]	0.9	*	*	*	*	*	*	*	*	*	0.1	*	0.2	0.4
1999[4]	1.1	*	*	*	*	*	*	*	*	*	*	*	0.2	0.4
Male														
2001	0.7	*	*	*	*	*	*	*	*	*	*	*	*	0.4
2000[4]	0.7	*	*	*	*	*	*	*	*	*	*	*	*	0.4
1999[4]	0.9	*	*	*	*	*	*	*	*	*	*	*	*	0.4
Female														
2001	1.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	2.3	*	*	*	*	*	*	*	*	1.0	1.3	1.9	2.4	4.4
2000[4]	2.3	*	*	*	*	*	*	*	*	*	1.2	2.0	2.5	4.1
1999[4]	2.8	*	*	*	*	*	*	*	*	1.0	2.0	2.3	2.3	5.2
Male														
2001	2.2	*	*	*	*	*	*	*	*	*	2.0	3.0	3.6	5.6
2000[4]	2.1	*	*	*	*	*	*	*	*	*	1.6	2.8	3.0	5.1
1999[4]	2.6	*	*	*	*	*	*	*	*	*	2.5	3.1	3.6	6.4
Female														
2001	2.4	*	*	*	*	*	*	*	*	*	*	*	*	3.4
2000[4]	2.4	*	*	*	*	*	*	*	*	*	*	*	2.1	3.3
1999[4]	3.1	*	*	*	*	*	*	*	*	*	1.6	1.7	*	4.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart and renal disease (I13)						
Non-Hispanic[5]						
Male						
2001	1.1	2.0	2.8	5.3	9.0	23.8
2000[4]	1.0	1.7	3.1	4.8	10.7	27.9
1999[4]	1.5	2.2	3.1	7.4	13.4	31.4
Female						
2001	1.3	1.4	2.7	4.8	9.3	22.8
2000[4]	0.9	1.5	2.8	5.2	9.3	22.8
1999[4]	1.2	1.9	2.9	6.1	11.7	28.4
Non-Hispanic White						
Both sexes						
2001	0.6	0.9	1.8	3.8	8.1	20.7
2000[4]	0.4	0.8	2.1	3.9	8.0	21.8
1999[4]	0.6	1.1	1.8	5.1	10.4	26.1
Male						
2001	0.5	1.1	1.9	4.2	8.3	22.2
2000[4]	*	1.0	2.2	4.2	9.0	24.3
1999[4]	0.7	1.3	1.8	5.9	12.2	28.7
Female						
2001	0.7	0.8	1.8	3.6	7.9	20.1
2000[4]	*	0.7	2.0	3.7	7.4	20.8
1999[4]	0.6	0.9	1.7	4.6	9.3	25.1
Non-Hispanic Black						
Both sexes						
2001	5.9	8.6	11.8	18.4	22.9	50.6
2000[4]	5.4	8.4	11.9	17.0	31.3	50.9
1999[4]	7.4	10.3	15.7	22.0	35.8	66.0
Male						
2001	6.5	10.9	13.4	18.5	18.8	45.0
2000[4]	5.7	9.3	13.3	12.5	33.5	64.6
1999[4]	9.6	10.9	17.7	23.7	30.1	60.6
Female						
2001	5.4	6.9	10.8	18.4	25.0	52.6
2000[4]	5.2	7.7	10.9	19.7	30.2	45.8
1999[4]	5.6	9.8	14.5	21.0	38.6	68.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Ischemic heart diseases (I20-I25)														
All origins[3]														
Both sexes														
2001	176.3	0.6	*	*	*	0.2	0.7	1.5	4.0	9.9	23.4	45.6	84.8	142.8
2000[4]	183.1	0.8	*	*	*	0.2	0.5	1.2	3.4	9.8	23.6	48.0	86.7	153.9
1999[4]	189.8	0.7	*	*	*	0.3	0.5	1.3	3.7	10.4	24.2	48.5	90.1	157.2
Male														
2001	181.7	*	*	*	*	0.2	0.9	2.2	5.9	15.0	35.8	71.6	131.1	215.5
2000[4]	188.7	*	*	*	*	0.2	0.6	1.8	5.1	14.7	37.0	76.9	134.2	232.8
1999[4]	195.4	*	*	*	*	0.3	0.6	1.7	5.5	15.8	37.9	77.0	140.3	237.7
Female														
2001	171.2	*	*	*	*	*	0.4	0.7	2.1	4.9	11.1	20.3	40.3	74.8
2000[4]	177.6	*	*	*	*	*	0.3	0.6	1.7	5.0	10.3	20.0	41.3	80.0
1999[4]	184.5	*	*	*	*	*	0.3	0.8	1.9	5.1	10.7	20.8	42.1	82.1
Hispanic														
Both sexes														
2001	55.9	*	*	*	*	*	*	0.9	2.3	4.3	9.7	25.8	55.5	98.1
2000[4]	55.9	*	*	*	*	*	*	0.6	1.6	4.3	11.8	27.4	55.4	109.9
1999[4]	58.7	*	*	*	*	*	*	0.7	2.1	4.7	10.8	29.9	64.4	111.6
Male														
2001	57.9	*	*	*	*	*	*	1.4	3.6	6.7	15.0	41.3	85.6	152.6
2000[4]	58.2	*	*	*	*	*	*	*	2.6	6.6	19.0	44.8	86.7	167.7
1999[4]	60.8	*	*	*	*	*	*	1.3	3.0	7.9	17.6	46.7	104.2	171.1
Female														
2001	53.7	*	*	*	*	*	*	*	*	1.6	4.1	10.1	26.6	48.7
2000[4]	53.6	*	*	*	*	*	*	*	*	1.7	4.2	10.0	25.5	57.6
1999[4]	56.6	*	*	*	*	*	*	*	*	*	3.7	13.2	26.6	58.0
Non-Hispanic[5]														
Both sexes														
2001	193.6	0.7	*	*	*	0.2	0.7	1.6	4.4	10.8	24.9	47.4	86.9	145.7
2000[4]	200.5	0.8	*	*	*	0.2	0.5	1.3	3.7	10.6	24.7	49.7	88.9	156.4
1999[4]	207.1	0.8	*	*	*	0.3	0.5	1.3	4.0	11.1	25.5	50.0	91.8	159.8
Male														
2001	200.4	*	*	*	*	0.2	1.0	2.4	6.4	16.3	38.2	74.5	134.5	219.2
2000[4]	207.6	*	*	*	*	*	0.7	2.1	5.5	15.8	38.9	79.6	137.3	236.5
1999[4]	214.1	*	*	*	*	0.4	0.7	1.8	6.0	16.8	40.0	79.5	142.5	241.2
Female														
2001	187.2	*	*	*	*	*	0.4	0.8	2.3	5.3	11.9	21.2	41.4	76.6
2000[4]	193.8	*	*	*	*	*	0.3	0.6	1.9	5.4	10.9	20.8	42.4	81.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Ischemic heart diseases (I20-I25)						
All origins[3]						
Both sexes						
2001	236.6	372.4	591.8	984.3	1,685.2	3,958.2
2000[4]	250.0	390.8	628.0	1,032.5	1,761.7	4,182.5
1999[4]	260.0	420.1	661.9	1,081.1	1,841.7	4,318.1
Male						
2001	344.4	525.7	820.3	1,337.5	2,180.0	4,395.8
2000[4]	362.9	550.5	879.6	1,394.4	2,305.0	4,701.7
1999[4]	376.1	595.2	924.6	1,462.5	2,400.4	4,837.5
Female						
2001	138.8	240.5	410.2	737.0	1,388.7	3,775.1
2000[4]	147.7	253.9	429.9	780.5	1,441.2	3,971.0
1999[4]	155.0	270.2	456.7	815.9	1,514.7	4,108.8
Hispanic						
Both sexes						
2001	172.8	306.5	518.7	881.4	1,481.0	3,620.2
2000[4]	186.8	312.5	521.9	873.5	1,544.0	3,697.2
1999[4]	199.2	340.5	551.1	948.1	1,596.5	3,876.2
Male						
2001	245.2	441.9	712.3	1,147.8	1,860.8	3,828.2
2000[4]	276.6	435.1	726.3	1,161.2	1,936.1	3,865.6
1999[4]	279.1	483.8	773.0	1,270.9	1,949.1	3,987.7
Female						
2001	110.0	196.9	371.0	692.3	1,245.8	3,518.6
2000[4]	109.4	213.2	367.1	669.8	1,305.3	3,614.6
1999[4]	130.4	224.7	384.2	720.2	1,386.1	3,822.4
Non-Hispanic[5]						
Both sexes						
2001	240.2	375.6	594.1	985.9	1,688.1	3,959.4
2000[4]	253.1	394.4	631.3	1,036.0	1,764.3	4,187.0
1999[4]	262.9	423.4	664.7	1,082.6	1,844.1	4,319.6
Male						
2001	349.7	529.1	823.6	1,342.0	2,185.7	4,408.5
2000[4]	366.8	555.3	883.8	1,399.7	2,312.5	4,721.1
1999[4]	380.4	599.4	927.6	1,464.8	2,409.4	4,854.5
Female						
2001	140.6	242.9	411.4	736.7	1,390.3	3,772.7
2000[4]	149.8	255.9	431.9	782.9	1,441.2	3,971.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Ischemic heart diseases (I20-I25)														
Non-Hispanic[5]														
Female														
1999[4]	200.6	*	*	*	*	*	0.3	0.9	2.1	5.6	11.4	21.4	43.1	83.6
Non-Hispanic White														
Both sexes														
2001	212.6	*	*	*	*	*	0.6	1.4	4.1	10.4	23.6	44.8	82.1	137.8
2000[4]	220.2	*	*	*	*	0.2	0.5	1.1	3.5	10.2	23.4	46.9	84.3	148.3
1999[4]	227.2	*	*	*	*	0.2	0.4	1.2	3.7	10.7	23.9	47.3	86.6	151.0
Male														
2001	219.6	*	*	*	*	*	0.9	2.2	6.0	15.8	37.2	72.2	129.4	210.4
2000[4]	227.6	*	*	*	*	*	0.6	1.8	5.3	15.5	37.8	77.2	132.8	226.8
1999[4]	234.2	*	*	*	*	0.3	0.6	1.6	5.5	16.2	38.3	77.1	136.7	231.0
Female														
2001	205.8	*	*	*	*	*	*	0.6	2.1	4.9	10.1	17.7	35.8	68.1
2000[4]	213.1	*	*	*	*	*	*	0.5	1.6	4.9	9.0	17.0	36.7	73.2
1999[4]	220.5	*	*	*	*	*	*	0.9	1.9	5.1	9.6	17.9	37.6	74.7
Non-Hispanic Black														
Both sexes														
2001	141.2	*	*	*	*	*	1.5	2.9	6.6	15.4	37.5	74.3	141.7	239.4
2000[4]	144.4	*	*	*	*	*	0.8	2.6	6.2	15.0	37.7	79.8	142.9	254.5
1999[4]	147.9	*	*	*	*	*	1.2	2.0	6.1	16.4	40.7	78.7	152.1	262.9
Male														
2001	143.2	*	*	*	*	*	2.0	4.0	9.5	22.5	52.0	104.5	203.8	337.8
2000[4]	145.7	*	*	*	*	*	*	3.9	8.2	20.8	52.5	113.9	201.0	371.1
1999[4]	150.0	*	*	*	*	*	*	3.1	8.8	23.8	58.1	112.9	221.3	378.6
Female														
2001	139.4	*	*	*	*	*	*	1.8	4.0	9.2	24.7	48.2	88.8	158.8
2000[4]	143.2	*	*	*	*	*	*	*	4.4	9.9	24.7	50.2	93.4	159.1
1999[4]	146.0	*	*	*	*	*	*	*	3.6	9.9	25.4	49.3	93.5	168.6
Acute myocardial infarction (I21-I22)														
All origins[3]														
Both sexes														
2001	64.9	*	*	*	*	*	0.2	0.6	1.8	4.2	9.9	19.7	36.5	61.3
2000[4]	68.5	0.5	*	*	*	*	0.2	0.6	1.3	4.0	9.9	21.1	37.8	67.0
1999[4]	71.5	*	*	*	*	0.1	0.2	0.5	1.6	4.1	10.4	21.5	39.0	69.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Ischemic heart diseases (I20-I25)						
Non-Hispanic[5]						
Female						
1999[4]	156.2	272.3	459.0	817.1	1,513.5	4,105.4
Non-Hispanic White						
Both sexes						
2001	228.7	359.7	581.5	973.2	1,692.1	4,022.8
2000[4]	240.8	380.8	617.2	1,021.9	1,770.0	4,261.3
1999[4]	251.6	410.9	649.3	1,071.3	1,851.7	4,390.2
Male						
2001	338.5	514.8	812.4	1,338.8	2,207.1	4,506.9
2000[4]	355.2	545.0	873.4	1,394.6	2,335.6	4,833.4
1999[4]	368.1	589.6	915.9	1,466.0	2,439.8	4,954.1
Female						
2001	126.7	222.7	393.8	714.3	1,381.2	3,821.4
2000[4]	134.8	236.5	410.9	759.9	1,433.9	4,030.7
1999[4]	143.8	254.3	436.8	794.3	1,504.7	4,165.3
Non-Hispanic Black						
Both sexes						
2001	381.7	570.0	820.0	1,257.4	1,844.5	3,628.7
2000[4]	398.2	573.6	874.6	1,307.5	1,894.4	3,701.7
1999[4]	400.2	602.8	911.2	1,327.7	1,932.5	3,817.9
Male						
2001	522.1	752.5	1,102.1	1,591.8	2,254.3	3,791.8
2000[4]	536.0	735.3	1,148.8	1,639.7	2,329.3	3,911.2
1999[4]	558.1	795.8	1,191.3	1,633.7	2,312.3	4,082.5
Female						
2001	271.3	435.3	632.0	1,057.8	1,635.8	3,568.3
2000[4]	289.5	454.4	692.6	1,107.3	1,675.5	3,623.9
1999[4]	276.3	459.7	725.3	1,143.3	1,744.2	3,719.0
Acute myocardial infarction (I21-I22)						
All origins[3]						
Both sexes						
2001	100.1	154.1	235.6	374.0	611.5	1,265.4
2000[4]	110.9	166.8	254.7	405.3	650.1	1,338.5
1999[4]	114.9	181.7	271.6	422.9	689.4	1,379.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute myocardial infarction (I21-I22)														
All origins[3]														
Male														
2001	68.6	*	*	*	*	*	0.3	0.7	2.5	6.2	15.0	30.8	55.6	91.3
2000[4]	72.7	*	*	*	*	*	0.2	0.8	1.9	5.8	15.5	33.6	57.5	99.7
1999[4]	76.1	*	*	*	*	*	0.3	0.7	2.3	6.2	16.0	33.7	60.9	105.3
Female														
2001	61.3	*	*	*	*	*	*	0.4	1.0	2.1	4.9	9.0	18.2	33.2
2000[4]	64.6	*	*	*	*	*	*	0.3	0.7	2.2	4.4	8.9	18.9	36.5
1999[4]	67.0	*	*	*	*	*	*	0.3	0.9	2.0	4.9	9.7	18.0	36.5
Hispanic														
Both sexes														
2001	20.0	*	*	*	*	*	*	*	1.0	1.5	3.8	10.5	20.5	36.2
2000[4]	20.1	*	*	*	*	*	*	*	*	1.7	4.6	11.0	21.3	41.7
1999[4]	21.1	*	*	*	*	*	*	*	0.7	1.7	4.4	12.9	24.3	38.6
Male														
2001	20.9	*	*	*	*	*	*	*	1.5	2.2	6.0	17.2	30.1	53.3
2000[4]	21.1	*	*	*	*	*	*	*	*	2.4	7.6	17.6	31.2	62.7
1999[4]	22.0	*	*	*	*	*	*	*	*	2.7	6.9	19.0	37.1	57.3
Female														
2001	19.1	*	*	*	*	*	*	*	*	*	*	3.7	11.3	20.6
2000[4]	19.1	*	*	*	*	*	*	*	*	*	*	4.5	11.9	22.6
1999[4]	20.1	*	*	*	*	*	*	*	*	*	*	6.8	12.2	21.7
Non-Hispanic[5]														
Both sexes														
2001	71.4	*	*	*	*	*	0.3	0.6	1.9	4.6	10.6	20.6	37.8	63.2
2000[4]	75.3	*	*	*	*	*	0.2	0.6	1.4	4.3	10.5	22.0	39.1	68.7
1999[4]	78.2	*	*	*	*	0.1	0.2	0.6	1.8	4.4	11.0	22.2	40.1	71.7
Male														
2001	75.9	*	*	*	*	*	0.4	0.8	2.7	6.9	16.0	32.1	57.7	94.0
2000[4]	80.3	*	*	*	*	*	*	0.9	2.0	6.3	16.4	35.1	59.5	102.2
1999[4]	83.8	*	*	*	*	*	0.3	0.7	2.5	6.6	17.0	35.0	62.6	108.3
Female														
2001	67.1	*	*	*	*	*	*	0.4	1.1	2.3	5.3	9.5	18.7	34.2
2000[4]	70.6	*	*	*	*	*	*	0.3	0.8	2.3	4.7	9.3	19.4	37.4
1999[4]	72.9	*	*	*	*	*	*	0.4	1.0	2.2	5.3	9.9	18.4	37.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute myocardial infarction (I21-I22)						
All origins[3]						
Male						
2001	143.5	214.2	322.0	495.6	776.1	1,461.3
2000[4]	158.6	231.3	349.6	535.2	835.8	1,568.3
1999[4]	164.9	253.5	373.8	564.2	885.1	1,622.1
Female						
2001	60.7	102.4	167.0	288.8	512.9	1,183.5
2000[4]	67.7	111.5	179.9	314.9	540.6	1,244.8
1999[4]	69.6	120.2	191.7	324.7	574.9	1,281.5
Hispanic						
Both sexes						
2001	67.2	113.4	195.2	328.7	511.1	1,185.5
2000[4]	72.1	118.8	198.8	331.8	537.5	1,177.1
1999[4]	78.2	129.6	207.5	365.1	558.6	1,227.8
Male						
2001	91.3	162.3	264.4	408.5	641.5	1,331.3
2000[4]	102.8	162.9	265.9	435.5	672.9	1,273.8
1999[4]	106.9	181.6	284.1	511.0	655.5	1,276.8
Female						
2001	46.3	73.7	142.4	272.1	430.3	1,114.2
2000[4]	45.7	83.0	148.0	258.3	455.1	1,129.7
1999[4]	53.4	87.5	149.8	262.1	500.8	1,204.1
Non-Hispanic[5]						
Both sexes						
2001	102.1	156.7	237.5	375.3	614.1	1,265.3
2000[4]	113.4	169.5	257.1	407.8	652.8	1,340.8
1999[4]	117.2	184.5	274.1	424.3	692.4	1,381.0
Male						
2001	146.7	217.3	324.5	498.6	779.7	1,463.9
2000[4]	162.2	235.0	353.2	538.5	840.4	1,576.2
1999[4]	168.4	257.4	377.0	564.9	891.3	1,631.5
Female						
2001	61.6	104.3	168.3	288.9	515.1	1,182.7
2000[4]	69.0	113.2	181.3	316.8	542.3	1,245.6
1999[4]	70.6	121.9	193.5	326.6	576.1	1,280.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute myocardial infarction (I21-I22)														
Non-Hispanic White														
Both sexes														
2001	78.4	*	*	*	*	*	0.2	0.5	1.9	4.5	10.3	20.2	36.7	61.6
2000[4]	82.7	*	*	*	*	*	0.2	0.5	1.4	4.4	10.1	21.6	37.9	67.3
1999[4]	86.0	*	*	*	*	*	*	0.5	1.7	4.4	10.7	21.8	39.3	70.2
Male														
2001	83.7	*	*	*	*	*	0.3	0.6	2.7	6.8	16.1	32.4	57.3	93.3
2000[4]	88.5	*	*	*	*	*	*	0.7	2.0	6.5	16.2	35.3	59.3	101.2
1999[4]	92.4	*	*	*	*	*	*	0.7	2.3	6.7	16.8	35.2	62.6	107.0
Female														
2001	73.4	*	*	*	*	*	*	0.4	1.1	2.2	4.5	8.1	16.5	31.1
2000[4]	77.2	*	*	*	*	*	*	*	0.7	2.2	3.9	8.0	17.0	34.9
1999[4]	79.8	*	*	*	*	*	*	0.4	1.0	2.0	4.6	8.6	16.4	35.0
Non-Hispanic Black														
Both sexes														
2001	51.6	*	*	*	*	*	*	1.1	2.6	6.0	14.6	28.6	54.3	90.9
2000[4]	54.0	*	*	*	*	*	*	1.2	2.3	5.1	15.3	30.2	56.7	96.2
1999[4]	55.1	*	*	*	*	*	*	0.9	2.5	5.6	15.8	30.9	56.3	98.8
Male														
2001	50.9	*	*	*	*	*	*	*	3.5	8.3	18.8	37.5	72.9	122.5
2000[4]	53.7	*	*	*	*	*	*	1.7	2.7	6.7	20.9	41.9	74.6	133.6
1999[4]	55.2	*	*	*	*	*	*	*	3.6	7.8	21.6	42.3	78.8	141.5
Female														
2001	52.3	*	*	*	*	*	*	*	1.8	3.9	10.9	20.8	38.5	65.0
2000[4]	54.2	*	*	*	*	*	*	*	1.9	3.7	10.3	20.1	41.4	65.7
1999[4]	54.9	*	*	*	*	*	*	*	1.5	3.6	10.6	21.0	37.3	64.0
Other acute ischemic heart diseases (I24)														
All origins[3]														
Both sexes														
2001	1.2	*	*	*	*	*	*	*	*	0.1	0.3	0.6	1.0	1.4
2000[4]	1.2	*	*	*	*	*	*	*	*	0.1	0.3	0.5	1.1	1.6
1999[4]	1.2	*	*	*	*	*	*	*	*	0.1	0.3	0.5	0.9	1.6
Male														
2001	1.2	*	*	*	*	*	*	*	*	*	0.4	0.9	1.6	2.0
2000[4]	1.3	*	*	*	*	*	*	*	*	0.2	0.4	0.8	1.6	2.5
1999[4]	1.2	*	*	*	*	*	*	*	*	*	0.4	0.7	1.3	2.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute myocardial infarction (I21-I22)						
Non-Hispanic White						
Both sexes						
2001	100.5	153.1	234.9	372.6	614.8	1,276.8
2000[4]	111.3	166.6	253.9	403.8	654.9	1,355.3
1999[4]	115.6	183.3	270.8	422.0	695.3	1,394.2
Male						
2001	147.1	216.2	324.4	500.3	786.9	1,483.2
2000[4]	161.8	234.8	352.6	539.5	847.9	1,600.7
1999[4]	168.7	258.1	377.4	569.2	903.1	1,653.8
Female						
2001	57.2	97.4	162.3	282.2	510.9	1,190.9
2000[4]	64.5	106.7	174.5	308.4	540.2	1,256.3
1999[4]	66.5	117.7	185.8	318.7	572.7	1,290.7
Non-Hispanic Black						
Both sexes						
2001	137.8	214.1	308.1	458.0	677.1	1,247.0
2000[4]	154.4	222.8	333.5	495.3	705.5	1,281.8
1999[4]	153.1	227.4	349.5	499.9	735.3	1,328.6
Male						
2001	176.6	267.8	390.8	556.0	802.9	1,389.4
2000[4]	200.3	275.0	426.1	596.8	865.1	1,426.6
1999[4]	202.7	295.6	436.9	597.4	863.5	1,509.4
Female						
2001	107.3	174.5	253.0	399.5	613.0	1,194.3
2000[4]	118.3	184.2	272.0	434.1	625.2	1,228.0
1999[4]	114.2	176.8	291.6	441.2	671.7	1,261.0
Other acute ischemic heart diseases (I24)						
All origins[3]						
Both sexes						
2001	2.4	2.9	4.1	5.3	9.2	21.5
2000[4]	2.5	3.1	4.0	5.9	9.4	21.9
1999[4]	2.2	3.1	4.3	5.5	9.2	21.9
Male						
2001	3.5	4.1	5.5	6.7	11.3	20.7
2000[4]	3.5	4.4	5.2	7.6	11.7	20.7
1999[4]	3.1	4.3	5.6	7.6	10.8	22.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other acute ischemic heart diseases (I24)														
All origins[3]														
Female														
2001	1.1	*	*	*	*	*	*	*	*	*	*	0.3	0.5	0.9
2000[4]	1.1	*	*	*	*	*	*	*	*	*	*	*	0.5	0.8
1999[4]	1.1	*	*	*	*	*	*	*	*	*	*	0.2	0.6	0.7
Hispanic														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	1.3	*	*	*	*	*	*	*	*	0.2	0.3	0.6	1.1	1.5
2000[4]	1.3	*	*	*	*	*	*	*	*	0.2	0.3	0.5	1.1	1.7
1999[4]	1.3	*	*	*	*	*	*	*	*	0.1	0.3	0.5	1.0	1.7
Male														
2001	1.4	*	*	*	*	*	*	*	*	*	0.5	1.0	1.7	2.0
2000[4]	1.4	*	*	*	*	*	*	*	*	0.2	0.5	0.8	1.7	2.6
1999[4]	1.4	*	*	*	*	*	*	*	*	*	0.4	0.8	1.4	2.6
Female														
2001	1.3	*	*	*	*	*	*	*	*	*	*	0.3	0.5	1.0
2000[4]	1.2	*	*	*	*	*	*	*	*	*	*	*	0.6	0.8
1999[4]	1.2	*	*	*	*	*	*	*	*	*	*	0.3	0.6	0.8
Non-Hispanic White														
Both sexes														
2001	1.4	*	*	*	*	*	*	*	*	0.2	0.2	0.6	0.9	1.3
2000[4]	1.4	*	*	*	*	*	*	*	*	0.1	0.3	0.4	1.0	1.4
1999[4]	1.4	*	*	*	*	*	*	*	*	*	0.2	0.4	0.9	1.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other acute ischemic heart diseases (I24)						
All origins[3]						
Female						
2001	1.3	2.0	3.1	4.4	8.0	21.8
2000[4]	1.5	1.9	2.9	4.8	8.1	22.4
1999[4]	1.5	2.0	3.2	4.1	8.3	21.7
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	2.5	3.0	4.3	5.5	9.5	22.0
2000[4]	2.6	3.2	4.0	6.1	9.6	22.2
1999[4]	2.3	3.2	4.5	5.6	9.5	22.2
Male						
2001	3.7	4.2	5.6	6.9	11.6	20.9
2000[4]	3.7	4.6	5.3	7.8	11.9	21.0
1999[4]	3.2	4.5	5.9	7.7	11.0	22.9
Female						
2001	1.4	2.1	3.2	4.5	8.2	22.4
2000[4]	1.6	2.1	3.0	4.9	8.2	22.7
1999[4]	1.5	2.1	3.3	4.2	8.5	21.9
Non-Hispanic White						
Both sexes						
2001	2.2	2.7	4.2	5.3	9.4	22.0
2000[4]	2.4	3.0	3.8	5.7	9.5	22.5
1999[4]	2.2	3.1	4.2	5.4	9.2	22.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other acute ischemic heart diseases (I24)														
Non-Hispanic White														
Male														
2001	1.4	*	*	*	*	*	*	*	*	*	0.4	0.9	1.4	1.9
2000[4]	1.5	*	*	*	*	*	*	*	*	*	0.4	0.8	1.6	2.3
1999[4]	1.4	*	*	*	*	*	*	*	*	*	0.4	0.6	1.3	2.4
Female														
2001	1.4	*	*	*	*	*	*	*	*	*	*	*	0.5	0.8
2000[4]	1.3	*	*	*	*	*	*	*	*	*	*	*	0.5	0.6
1999[4]	1.3	*	*	*	*	*	*	*	*	*	*	*	0.5	0.7
Non-Hispanic Black														
Both sexes														
2001	1.3	*	*	*	*	*	*	*	*	*	0.8	1.1	2.2	2.9
2000[4]	1.3	*	*	*	*	*	*	*	*	*	*	1.0	2.1	3.9
1999[4]	1.2	*	*	*	*	*	*	*	*	*	*	1.5	2.3	3.3
Male														
2001	1.5	*	*	*	*	*	*	*	*	*	*	*	3.7	3.5
2000[4]	1.5	*	*	*	*	*	*	*	*	*	*	*	3.5	5.9
1999[4]	1.5	*	*	*	*	*	*	*	*	*	*	2.5	2.9	5.7
Female														
2001	1.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Other forms of chronic ischemic heart disease (I20,I25)														
All origins[3]														
Both sexes														
2001	110.3	*	*	*	*	*	0.4	0.9	2.2	5.6	13.2	25.2	47.3	80.1
2000[4]	113.3	*	*	*	*	*	0.3	0.6	2.0	5.7	13.4	26.5	47.9	85.2
1999[4]	117.2	*	*	*	*	0.1	0.3	0.7	2.1	6.2	13.5	26.5	50.2	85.9
Male														
2001	111.9	*	*	*	*	*	0.5	1.5	3.3	8.6	20.4	40.0	74.0	122.2
2000[4]	114.8	*	*	*	*	*	0.4	1.1	3.1	8.7	21.1	42.5	75.1	130.6
1999[4]	118.0	*	*	*	*	0.2	0.3	1.0	3.2	9.4	21.5	42.6	78.1	129.9
Female														
2001	108.8	*	*	*	*	*	0.2	0.3	1.1	2.7	6.1	11.0	21.7	40.6
2000[4]	111.9	*	*	*	*	*	*	0.2	1.0	2.8	5.8	10.9	21.8	42.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other acute ischemic heart diseases (I24)						
Non-Hispanic White						
Male						
2001	3.3	3.6	5.5	6.7	11.9	20.9
2000[4]	3.4	4.2	5.0	7.4	11.8	21.3
1999[4]	2.9	4.3	5.5	7.7	10.5	23.6
Female						
2001	1.2	2.0	3.1	4.4	7.9	22.4
2000[4]	1.4	2.0	2.9	4.6	8.2	23.0
1999[4]	1.5	2.0	3.1	3.9	8.4	22.4
Non-Hispanic Black						
Both sexes						
2001	5.3	5.4	6.3	6.9	11.0	24.3
2000[4]	4.9	5.2	6.1	10.1	11.8	21.9
1999[4]	3.9	4.4	7.5	8.9	14.1	17.5
Male						
2001	8.4	9.4	7.9	11.4	*	24.3
2000[4]	6.8	8.3	9.2	13.5	*	*
1999[4]	6.5	6.9	10.4	10.2	17.7	*
Female						
2001	*	*	5.3	*	12.1	24.4
2000[4]	3.4	*	*	8.1	9.9	22.2
1999[4]	*	*	5.5	8.2	12.3	16.4
Other forms of chronic ischemic heart disease (I20,I25)						
All origins[3]						
Both sexes						
2001	134.1	215.4	352.0	605.0	1,064.5	2,671.3
2000[4]	136.6	221.0	369.4	621.2	1,102.2	2,822.1
1999[4]	142.9	235.3	386.1	652.6	1,143.1	2,917.0
Male						
2001	197.4	307.4	492.9	835.3	1,392.6	2,913.8
2000[4]	200.8	314.8	524.7	851.5	1,457.6	3,112.7
1999[4]	208.1	337.3	545.2	890.7	1,504.6	3,193.1
Female						
2001	76.8	136.2	240.1	443.8	867.9	2,569.8
2000[4]	78.5	140.5	247.0	460.8	892.5	2,703.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other forms of chronic ischemic heart disease (I20,I25)														
All origins[3]														
Female														
1999[4]	116.4	*	*	*	*	*	*	0.4	1.0	3.1	5.6	10.9	23.4	44.9
Hispanic														
Both sexes														
2001	35.7	*	*	*	*	*	*	*	1.3	2.8	5.9	15.3	34.8	61.3
2000[4]	35.6	*	*	*	*	*	*	*	1.0	2.6	7.1	16.1	33.9	67.6
1999[4]	37.5	*	*	*	*	*	*	*	1.3	3.0	6.4	17.0	39.6	72.6
Male														
2001	36.8	*	*	*	*	*	*	*	2.0	4.6	8.9	24.1	55.3	98.3
2000[4]	36.9	*	*	*	*	*	*	*	1.5	4.3	11.3	27.0	55.4	104.3
1999[4]	38.7	*	*	*	*	*	*	*	2.0	5.2	10.6	27.6	66.6	113.3
Female														
2001	34.5	*	*	*	*	*	*	*	*	*	2.7	6.4	15.2	27.9
2000[4]	34.2	*	*	*	*	*	*	*	*	*	2.7	5.3	13.5	34.3
1999[4]	36.3	*	*	*	*	*	*	*	*	*	2.0	6.4	14.0	35.9
Non-Hispanic[5]														
Both sexes														
2001	120.9	*	*	*	*	*	0.4	1.0	2.4	6.1	14.0	26.1	48.0	81.1
2000[4]	123.9	*	*	*	*	*	0.3	0.7	2.2	6.1	14.0	27.2	48.7	86.0
1999[4]	127.6	*	*	*	*	0.1	0.3	0.8	2.2	6.6	14.2	27.2	50.7	86.5
Male														
2001	123.1	*	*	*	*	*	0.6	1.6	3.6	9.3	21.8	41.4	75.1	123.2
2000[4]	126.0	*	*	*	*	*	0.4	1.2	3.4	9.3	22.0	43.7	76.1	131.7
1999[4]	128.9	*	*	*	*	*	0.4	1.1	3.4	10.0	22.6	43.7	78.5	130.3
Female														
2001	118.8	*	*	*	*	*	*	0.4	1.1	2.9	6.5	11.4	22.1	41.5
2000[4]	122.0	*	*	*	*	*	*	*	1.1	3.0	6.1	11.3	22.4	43.2
1999[4]	126.4	*	*	*	*	*	*	0.5	1.1	3.4	6.0	11.3	24.1	45.4
Non-Hispanic White														
Both sexes														
2001	132.7	*	*	*	*	*	0.3	0.9	2.1	5.7	13.1	24.0	44.5	74.9
2000[4]	136.1	*	*	*	*	*	0.2	0.6	2.0	5.7	13.1	24.9	45.3	79.5
1999[4]	139.9	*	*	*	*	*	0.2	0.7	2.0	6.2	13.0	25.2	46.5	79.3
Male														
2001	134.4	*	*	*	*	*	0.5	1.6	3.2	8.8	20.7	39.0	70.6	115.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)						
All origins[3]						
Female						
1999[4]	83.9	148.1	261.7	487.1	931.6	2,805.7
Hispanic						
Both sexes						
2001	104.8	191.7	321.5	549.9	968.0	2,426.6
2000[4]	114.2	193.0	320.8	539.0	1,002.0	2,508.8
1999[4]	119.8	210.1	343.0	579.7	1,036.7	2,640.7
Male						
2001	152.9	277.5	444.2	735.9	1,216.7	2,479.8
2000[4]	173.3	270.7	457.0	722.0	1,257.4	2,577.7
1999[4]	171.0	301.0	488.3	754.4	1,292.1	2,706.7
Female						
2001	63.2	122.2	227.9	417.8	813.9	2,400.7
2000[4]	63.3	130.1	217.6	409.4	846.6	2,475.0
1999[4]	75.7	136.6	233.7	456.5	884.3	2,608.9
Non-Hispanic[5]						
Both sexes						
2001	135.6	215.9	352.3	605.2	1,064.5	2,672.2
2000[4]	137.1	221.6	370.1	622.1	1,101.9	2,824.0
1999[4]	143.5	235.7	386.2	652.7	1,142.2	2,916.3
Male						
2001	199.4	307.7	493.5	836.5	1,394.4	2,923.7
2000[4]	200.9	315.7	525.3	853.4	1,460.2	3,123.9
1999[4]	208.8	337.5	544.7	892.2	1,507.1	3,200.1
Female						
2001	77.6	136.6	239.9	443.3	867.0	2,567.6
2000[4]	79.1	140.7	247.6	461.2	890.7	2,702.7
1999[4]	84.1	148.3	262.1	486.4	928.9	2,802.7
Non-Hispanic White						
Both sexes						
2001	126.0	203.8	342.4	595.3	1,067.8	2,724.1
2000[4]	127.1	211.2	359.4	612.4	1,105.6	2,883.6
1999[4]	133.8	224.5	374.3	643.9	1,147.2	2,973.2
Male						
2001	188.2	295.0	482.6	831.9	1,408.3	3,002.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other forms of chronic ischemic heart disease (I20,I25)														
Non-Hispanic White														
Male														
2000[4]	137.7	*	*	*	*	*	0.4	1.0	3.2	8.8	21.2	41.1	72.0	123.2
1999[4]	140.4	*	*	*	*	*	*	0.9	3.1	9.3	21.1	41.4	72.8	121.6
Female														
2001	131.1	*	*	*	*	*	*	*	1.0	2.6	5.5	9.3	18.8	36.2
2000[4]	134.6	*	*	*	*	*	*	*	0.9	2.6	4.9	8.9	19.2	37.8
1999[4]	139.3	*	*	*	*	*	*	0.5	0.9	3.1	4.8	9.2	20.7	38.9
Non-Hispanic Black														
Both sexes														
2001	88.2	*	*	*	*	*	0.8	1.8	3.9	9.2	22.2	44.6	85.2	145.6
2000[4]	89.1	*	*	*	*	*	*	1.3	3.7	9.7	21.9	48.5	84.1	154.4
1999[4]	91.6	*	*	*	*	*	*	1.2	3.5	10.4	24.4	46.4	93.5	160.9
Male														
2001	90.8	*	*	*	*	*	*	2.5	6.0	13.8	32.1	65.5	127.3	211.7
2000[4]	90.4	*	*	*	*	*	*	2.2	5.1	13.7	30.8	70.5	122.9	231.6
1999[4]	93.3	*	*	*	*	*	*	1.8	5.2	15.3	35.6	68.1	139.6	231.4
Female														
2001	86.0	*	*	*	*	*	*	*	2.0	5.0	13.4	26.6	49.4	91.4
2000[4]	88.0	*	*	*	*	*	*	*	2.5	6.0	14.1	29.5	51.1	91.2
1999[4]	90.1	*	*	*	*	*	*	*	2.0	6.0	14.6	27.6	54.5	103.3
Atherosclerotic cardiovascular disease, so described (I25.0)														
All origins[3]														
Both sexes														
2001	23.9	*	*	*	*	*	*	0.2	0.7	1.8	4.6	8.8	17.4	27.7
2000[4]	24.7	*	*	*	*	*	*	0.2	0.5	1.7	4.4	9.4	17.5	28.5
1999[4]	25.8	*	*	*	*	*	*	0.2	0.6	2.0	4.7	9.1	17.4	28.6
Male														
2001	25.3	*	*	*	*	*	*	0.3	1.0	2.8	7.2	14.6	28.3	43.4
2000[4]	26.0	*	*	*	*	*	*	0.3	0.9	2.8	7.1	15.5	28.4	45.8
1999[4]	26.8	*	*	*	*	*	*	0.2	0.9	3.1	7.6	14.9	28.0	45.0
Female														
2001	22.5	*	*	*	*	*	*	*	0.4	0.8	2.1	3.3	7.0	12.9
2000[4]	23.5	*	*	*	*	*	*	*	0.2	0.7	1.8	3.5	7.2	12.4
1999[4]	24.8	*	*	*	*	*	*	*	0.3	0.9	1.8	3.5	7.3	13.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)						
Non-Hispanic White						
Male						
2000[4]	189.9	306.1	515.8	847.8	1,475.9	3,211.3
1999[4]	196.5	327.2	533.0	889.2	1,526.1	3,276.7
Female						
2001	68.2	123.3	228.4	427.7	862.3	2,608.1
2000[4]	68.9	127.8	233.5	447.0	885.5	2,751.4
1999[4]	75.8	134.6	247.9	471.7	923.6	2,852.2
Non-Hispanic Black						
Both sexes						
2001	238.5	350.5	505.5	792.5	1,156.4	2,357.3
2000[4]	238.9	345.6	535.0	802.0	1,177.1	2,398.0
1999[4]	243.2	371.0	554.2	818.9	1,183.2	2,471.9
Male						
2001	337.1	475.2	703.4	1,024.4	1,442.4	2,378.1
2000[4]	328.9	452.0	713.5	1,029.4	1,448.7	2,463.5
1999[4]	348.9	493.3	744.0	1,026.2	1,431.1	2,552.9
Female						
2001	161.0	258.3	373.7	654.1	1,010.7	2,349.6
2000[4]	167.9	267.2	416.5	665.0	1,040.4	2,373.7
1999[4]	160.3	280.3	428.3	694.0	1,060.2	2,441.6
Atherosclerotic cardiovascular disease, so described (I25.0)						
All origins[3]						
Both sexes						
2001	40.6	55.3	76.7	121.8	197.2	493.6
2000[4]	40.7	55.7	81.9	125.0	207.8	534.6
1999[4]	43.4	59.6	86.8	131.6	218.2	563.4
Male						
2001	61.8	79.0	107.6	160.2	247.4	502.0
2000[4]	61.3	80.8	115.6	167.1	252.8	550.8
1999[4]	64.6	87.3	120.8	172.7	264.3	578.6
Female						
2001	21.4	34.9	52.1	94.9	167.1	490.1
2000[4]	22.1	34.1	55.4	95.7	181.2	528.0
1999[4]	24.1	36.0	60.2	103.1	191.3	557.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Atherosclerotic cardiovascular disease, so described (I25.0)														
Hispanic														
Both sexes														
2001	8.5	*	*	*	*	*	*	*	*	0.9	2.2	6.2	15.1	23.5
2000[4]	8.4	*	*	*	*	*	*	*	*	1.0	2.4	6.3	13.8	24.0
1999[4]	9.0	*	*	*	*	*	*	*	*	0.8	2.3	6.5	15.4	27.1
Male														
2001	10.1	*	*	*	*	*	*	*	*	1.4	3.2	10.0	26.6	39.2
2000[4]	9.9	*	*	*	*	*	*	*	*	1.7	3.9	10.9	24.1	39.9
1999[4]	10.6	*	*	*	*	*	*	*	*	1.5	3.6	10.6	27.4	42.5
Female														
2001	6.8	*	*	*	*	*	*	*	*	*	*	2.5	4.1	9.2
2000[4]	6.8	*	*	*	*	*	*	*	*	*	*	*	4.0	9.5
1999[4]	7.4	*	*	*	*	*	*	*	*	*	*	2.5	4.1	13.1
Non-Hispanic[5]														
Both sexes														
2001	26.0	*	*	*	*	*	*	0.2	0.7	1.9	4.9	9.1	17.5	27.7
2000[4]	26.9	*	*	*	*	*	*	0.2	0.6	1.8	4.6	9.6	17.7	28.6
1999[4]	27.9	*	*	*	*	*	*	0.2	0.7	2.1	4.9	9.3	17.4	28.5
Male														
2001	27.5	*	*	*	*	*	*	0.3	1.0	3.1	7.6	14.9	28.2	43.2
2000[4]	28.2	*	*	*	*	*	*	0.3	1.0	2.9	7.4	15.8	28.4	45.7
1999[4]	28.9	*	*	*	*	*	*	*	1.0	3.3	8.0	15.2	27.7	44.8
Female														
2001	24.6	*	*	*	*	*	*	*	0.4	0.8	2.2	3.4	7.2	13.1
2000[4]	25.6	*	*	*	*	*	*	*	0.3	0.7	1.9	3.7	7.4	12.5
1999[4]	27.0	*	*	*	*	*	*	*	0.3	1.0	1.9	3.6	7.5	13.3
Non-Hispanic White														
Both sexes														
2001	26.8	*	*	*	*	*	*	0.2	0.7	1.8	4.4	8.1	15.4	23.6
2000[4]	27.6	*	*	*	*	*	*	*	0.6	1.6	4.2	8.5	15.7	24.5
1999[4]	28.6	*	*	*	*	*	*	*	0.6	1.9	4.4	8.0	14.9	24.2
Male														
2001	27.8	*	*	*	*	*	*	*	1.0	2.9	7.0	13.6	25.3	37.7
2000[4]	28.4	*	*	*	*	*	*	*	0.9	2.7	6.9	14.2	25.6	39.5
1999[4]	29.0	*	*	*	*	*	*	*	0.9	3.0	7.3	13.4	24.2	38.7
Female														
2001	25.8	*	*	*	*	*	*	*	0.3	0.7	1.8	2.6	5.7	10.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)						
Hispanic						
Both sexes						
2001	37.9	54.9	79.6	112.1	168.4	417.0
2000[4]	38.9	54.1	80.2	101.3	177.1	452.5
1999[4]	40.4	62.9	83.7	124.0	190.3	469.4
Male						
2001	59.7	87.2	117.1	159.6	222.0	440.0
2000[4]	64.5	74.9	115.2	158.7	225.3	493.8
1999[4]	59.6	103.6	131.9	171.6	238.9	485.3
Female						
2001	19.0	28.7	51.0	78.4	135.1	405.8
2000[4]	16.9	37.1	53.8	60.6	147.8	432.3
1999[4]	23.9	30.0	47.5	90.3	161.3	461.7
Non-Hispanic[5]						
Both sexes						
2001	40.5	54.8	75.9	121.3	197.5	494.8
2000[4]	40.3	55.3	81.3	125.0	207.9	535.9
1999[4]	43.0	58.9	86.0	130.8	217.9	564.9
Male						
2001	61.4	77.6	106.0	158.8	247.2	503.0
2000[4]	60.2	80.4	114.3	165.7	252.5	551.0
1999[4]	63.8	85.4	118.6	170.8	263.4	580.2
Female						
2001	21.5	35.1	51.9	95.0	167.7	491.4
2000[4]	22.3	33.8	55.2	96.7	181.6	529.8
1999[4]	24.0	36.1	60.5	103.0	191.3	558.8
Non-Hispanic White						
Both sexes						
2001	34.4	46.7	68.0	111.2	190.8	490.7
2000[4]	34.3	48.6	72.3	115.5	198.2	530.2
1999[4]	36.6	51.0	76.8	120.3	209.3	558.8
Male						
2001	52.9	66.9	95.2	146.4	239.6	499.4
2000[4]	52.4	71.8	102.1	152.6	240.8	544.7
1999[4]	55.0	75.0	106.5	157.5	252.7	572.8
Female						
2001	17.3	28.8	45.9	86.3	161.3	487.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Atherosclerotic cardiovascular disease, so described (I25.0)														
Non-Hispanic White														
Female														
2000[4]	26.9	*	*	*	*	*	*	*	*	0.6	1.5	2.8	5.9	10.2
1999[4]	28.2	*	*	*	*	*	*	*	*	0.8	1.4	2.8	5.9	10.3
Non-Hispanic Black														
Both sexes														
2001	28.5	*	*	*	*	*	*	*	1.0	3.2	8.5	17.1	36.3	64.5
2000[4]	29.3	*	*	*	*	*	*	*	0.9	3.0	7.6	19.7	36.9	66.3
1999[4]	30.7	*	*	*	*	*	*	*	1.2	3.9	9.3	19.5	39.1	68.6
Male														
2001	32.3	*	*	*	*	*	*	*	*	4.8	12.8	26.3	56.7	97.3
2000[4]	33.2	*	*	*	*	*	*	*	*	4.8	11.2	30.5	57.2	107.3
1999[4]	34.5	*	*	*	*	*	*	*	1.8	6.0	13.9	30.5	60.8	105.2
Female														
2001	25.0	*	*	*	*	*	*	*	*	1.7	4.7	9.2	18.9	37.6
2000[4]	25.8	*	*	*	*	*	*	*	*	1.5	4.4	10.4	19.5	32.7
1999[4]	27.1	*	*	*	*	*	*	*	*	2.0	5.2	10.1	20.7	38.7
All other forms of chronic ischemic heart disease (I20, I25.1-I25.9)														
All origins[3]														
Both sexes														
2001	86.4	*	*	*	*	*	0.3	0.7	1.5	3.8	8.6	16.4	29.8	52.4
2000[4]	88.6	*	*	*	*	*	0.2	0.5	1.5	4.0	9.0	17.0	30.3	56.7
1999[4]	91.4	*	*	*	*	0.1	0.2	0.6	1.5	4.3	8.8	17.4	32.8	57.3
Male														
2001	86.5	*	*	*	*	*	0.4	1.1	2.3	5.8	13.2	25.4	45.7	78.8
2000[4]	88.8	*	*	*	*	*	0.2	0.8	2.2	5.9	14.0	27.0	46.7	84.9
1999[4]	91.2	*	*	*	*	0.2	0.2	0.8	2.2	6.3	13.9	27.7	50.2	84.9
Female														
2001	86.2	*	*	*	*	*	*	0.2	0.7	1.9	4.0	7.6	14.7	27.7
2000[4]	88.4	*	*	*	*	*	*	*	0.8	2.1	4.0	7.3	14.6	30.3
1999[4]	91.6	*	*	*	*	*	*	0.3	0.7	2.2	3.9	7.4	16.1	31.5
Hispanic														
Both sexes														
2001	27.2	*	*	*	*	*	*	*	0.8	1.9	3.6	9.0	19.7	37.9
2000[4]	27.2	*	*	*	*	*	*	*	0.8	1.6	4.7	9.9	20.1	43.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)						
Non-Hispanic White						
Female						
2000[4]	17.4	28.2	48.3	89.5	172.9	524.4
1999[4]	19.7	30.0	53.1	94.2	183.6	553.2
Non-Hispanic Black						
Both sexes						
2001	95.5	129.4	164.1	244.8	300.8	595.9
2000[4]	94.4	120.8	178.9	240.8	342.8	652.2
1999[4]	98.4	135.0	187.7	260.8	339.9	670.3
Male						
2001	145.3	185.0	241.7	332.8	385.9	609.8
2000[4]	137.8	170.0	259.7	339.1	429.1	711.6
1999[4]	146.7	198.2	267.3	346.6	436.6	714.3
Female						
2001	56.4	88.3	112.4	192.4	257.4	590.7
2000[4]	60.2	84.5	125.2	181.5	299.4	630.2
1999[4]	60.4	88.2	134.9	209.1	291.9	653.8
All other forms of chronic ischemic heart disease (I20, I25.1-I25.9)						
All origins[3]						
Both sexes						
2001	93.5	160.1	275.3	483.2	867.3	2,177.7
2000[4]	95.9	165.3	287.4	496.2	894.4	2,287.5
1999[4]	99.5	175.7	299.3	521.0	924.9	2,353.6
Male						
2001	135.6	228.4	385.2	675.1	1,145.1	2,411.8
2000[4]	139.5	234.0	409.1	684.5	1,204.8	2,561.9
1999[4]	143.5	250.0	424.4	718.0	1,240.3	2,614.4
Female						
2001	55.4	101.2	188.0	348.9	700.7	2,079.8
2000[4]	56.4	106.3	191.6	365.1	711.3	2,175.8
1999[4]	59.7	112.1	201.5	384.0	740.3	2,248.4
Hispanic						
Both sexes						
2001	66.9	136.8	241.9	437.7	799.6	2,009.6
2000[4]	75.3	139.0	240.5	437.7	824.9	2,056.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
All other forms of chronic ischemic heart disease (I20, I25.1-I25.9)														
Hispanic														
Both sexes														
1999[4]	28.5	*	*	*	*	*	*	*	1.0	2.2	4.1	10.4	24.2	45.5
Male														
2001	26.7	*	*	*	*	*	*	*	1.3	3.1	5.7	14.1	28.7	59.1
2000[4]	27.0	*	*	*	*	*	*	*	1.2	2.6	7.4	16.0	31.3	64.5
1999[4]	28.1	*	*	*	*	*	*	*	1.4	3.7	6.9	17.0	39.2	70.7
Female														
2001	27.7	*	*	*	*	*	*	*	*	*	*	3.9	11.1	18.6
2000[4]	27.5	*	*	*	*	*	*	*	*	*	1.9	3.7	9.5	24.8
1999[4]	28.9	*	*	*	*	*	*	*	*	*	*	3.9	9.9	22.7
Non-Hispanic[5]														
Both sexes														
2001	94.9	*	*	*	*	*	0.3	0.8	1.6	4.1	9.2	17.1	30.6	53.4
2000[4]	97.0	*	*	*	*	*	0.2	0.5	1.6	4.3	9.4	17.6	31.0	57.4
1999[4]	99.7	*	*	*	*	0.1	0.2	0.6	1.6	4.5	9.3	17.9	33.3	57.9
Male														
2001	95.6	*	*	*	*	*	0.4	1.3	2.5	6.2	14.1	26.5	46.9	79.9
2000[4]	97.8	*	*	*	*	*	0.3	0.9	2.4	6.4	14.7	27.9	47.7	86.0
1999[4]	100.0	*	*	*	*	*	*	0.8	2.4	6.6	14.6	28.6	50.8	85.6
Female														
2001	94.2	*	*	*	*	*	*	0.3	0.8	2.1	4.3	8.0	14.9	28.4
2000[4]	96.3	*	*	*	*	*	*	*	0.8	2.3	4.2	7.6	15.0	30.7
1999[4]	99.4	*	*	*	*	*	*	0.4	0.7	2.4	4.1	7.7	16.6	32.1
Non-Hispanic White														
Both sexes														
2001	106.0	*	*	*	*	*	0.2	0.7	1.5	3.9	8.7	16.0	29.1	51.3
2000[4]	108.4	*	*	*	*	*	*	0.5	1.5	4.0	8.9	16.4	29.6	55.0
1999[4]	111.3	*	*	*	*	*	*	0.6	1.5	4.3	8.6	17.1	31.5	55.1
Male														
2001	106.6	*	*	*	*	*	0.4	1.3	2.3	5.9	13.7	25.4	45.4	77.5
2000[4]	109.2	*	*	*	*	*	*	0.8	2.3	6.1	14.3	26.9	46.3	83.7
1999[4]	111.4	*	*	*	*	*	*	0.7	2.3	6.3	13.8	28.0	48.6	82.9
Female														
2001	105.3	*	*	*	*	*	*	*	0.7	1.9	3.7	6.7	13.1	26.2
2000[4]	107.7	*	*	*	*	*	*	*	0.7	2.0	3.5	6.1	13.2	27.6
1999[4]	111.1	*	*	*	*	*	*	0.4	0.6	2.2	3.4	6.4	14.8	28.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20, I25.1-I25.9)						
Hispanic						
Both sexes						
1999[4]	79.4	147.2	259.3	455.8	846.4	2,171.4
Male						
2001	93.2	190.3	327.1	576.3	994.7	2,039.8
2000[4]	108.8	195.8	341.8	563.3	1,032.1	2,084.0
1999[4]	111.4	197.4	356.4	582.8	1,053.1	2,221.4
Female						
2001	44.2	93.5	176.9	339.4	678.8	1,994.8
2000[4]	46.4	93.0	163.9	348.7	698.8	2,042.7
1999[4]	51.8	106.6	186.2	366.2	723.0	2,147.2
Non-Hispanic[5]						
Both sexes						
2001	95.1	161.1	276.4	483.9	867.0	2,177.4
2000[4]	96.8	166.3	288.8	497.1	894.0	2,288.1
1999[4]	100.5	176.8	300.2	522.0	924.3	2,351.4
Male						
2001	138.0	230.1	387.4	677.7	1,147.3	2,420.7
2000[4]	140.8	235.3	411.0	687.7	1,207.7	2,572.9
1999[4]	145.0	252.0	426.1	721.4	1,243.7	2,619.9
Female						
2001	56.1	101.5	188.0	348.3	699.3	2,076.2
2000[4]	56.9	106.9	192.4	364.5	709.1	2,172.9
1999[4]	60.1	112.2	201.6	383.4	737.6	2,243.9
Non-Hispanic White						
Both sexes						
2001	91.6	157.1	274.4	484.0	877.0	2,233.4
2000[4]	92.8	162.6	287.1	496.9	907.4	2,353.3
1999[4]	97.1	173.5	297.5	523.6	937.9	2,414.4
Male						
2001	135.4	228.0	387.4	685.5	1,168.6	2,503.4
2000[4]	137.5	234.2	413.6	695.2	1,235.1	2,666.6
1999[4]	141.5	252.2	426.5	731.7	1,273.5	2,703.9
Female						
2001	51.0	94.5	182.5	341.4	701.0	2,121.0
2000[4]	51.5	99.6	185.3	357.5	712.6	2,227.0
1999[4]	56.1	104.6	194.7	377.5	739.9	2,299.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
All other forms of chronic ischemic heart disease (I20, I25.1-I25.9)														
Non-Hispanic Black														
Both sexes														
2001	59.7	*	*	*	*	*	*	1.3	2.9	6.0	13.7	27.5	48.9	81.1
2000[4]	59.8	*	*	*	*	*	*	0.9	2.8	6.6	14.3	28.8	47.3	88.1
1999[4]	61.0	*	*	*	*	*	*	0.9	2.3	6.5	15.1	26.8	54.4	92.3
Male														
2001	58.4	*	*	*	*	*	*	1.9	4.6	9.0	19.4	39.2	70.5	114.5
2000[4]	57.3	*	*	*	*	*	*	*	3.9	8.9	19.6	40.0	65.6	124.3
1999[4]	58.8	*	*	*	*	*	*	*	3.4	9.3	21.6	37.7	78.8	126.1
Female														
2001	60.9	*	*	*	*	*	*	*	*	3.4	8.7	17.4	30.5	53.8
2000[4]	62.1	*	*	*	*	*	*	*	1.9	4.6	9.7	19.2	31.6	58.5
1999[4]	63.0	*	*	*	*	*	*	*	*	4.0	9.4	17.5	33.7	64.7
Other heart diseases (I26-I51)														
All origins[3]														
Both sexes														
2001	58.6	11.1	1.4	0.5	0.7	1.5	2.5	3.6	5.1	8.0	11.9	17.9	25.9	39.7
2000[4]	58.8	12.0	1.1	0.5	0.7	1.8	2.6	3.5	5.2	7.5	12.3	17.6	25.9	40.2
1999[4]	59.4	12.9	1.2	0.5	0.7	2.0	2.6	3.5	5.3	8.2	12.5	17.8	25.9	41.8
Male														
2001	51.8	10.8	1.5	0.4	0.8	1.9	3.1	4.3	6.0	9.9	14.9	23.1	32.7	49.4
2000[4]	52.2	12.2	1.2	0.6	0.9	2.1	3.1	4.2	6.2	9.1	15.4	23.0	33.4	52.0
1999[4]	53.1	13.1	1.3	0.5	0.8	2.4	3.0	4.0	6.4	10.3	15.6	23.0	33.3	54.1
Female														
2001	65.2	11.5	1.3	0.5	0.7	1.1	1.9	2.9	4.1	6.0	8.9	12.8	19.5	30.6
2000[4]	65.1	11.8	0.9	0.4	0.6	1.4	2.1	2.9	4.1	5.9	9.2	12.3	18.7	29.2
1999[4]	65.5	12.7	1.0	0.4	0.7	1.5	2.2	2.9	4.2	6.1	9.4	12.8	18.9	30.2
Hispanic														
Both sexes														
2001	13.3	9.2	1.0	*	*	1.2	1.3	2.1	3.2	4.1	5.9	8.9	13.5	21.3
2000[4]	13.6	10.2	1.0	*	0.6	1.5	1.9	2.3	3.3	4.6	6.7	8.5	15.6	23.3
1999[4]	13.8	12.2	1.2	*	*	1.5	1.7	1.9	3.7	5.4	7.4	9.1	15.0	23.6
Male														
2001	12.6	9.6	*	*	*	1.3	1.8	2.6	3.8	5.3	8.0	12.5	18.3	25.4
2000[4]	13.2	9.9	1.3	*	*	2.0	2.2	3.0	4.1	5.4	8.3	10.7	20.9	31.3
1999[4]	13.7	10.3	1.4	*	*	1.5	2.0	2.8	4.9	6.7	10.0	11.6	20.5	29.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20, I25.1-I25.9)						
Non-Hispanic Black						
Both sexes						
2001	143.0	221.1	341.4	547.6	855.6	1,761.4
2000[4]	144.5	224.8	356.1	561.3	834.3	1,745.8
1999[4]	144.8	236.0	366.5	558.1	843.3	1,801.6
Male						
2001	191.9	290.3	461.7	691.6	1,056.4	1,768.3
2000[4]	191.1	282.0	453.8	690.3	1,019.6	1,751.9
1999[4]	202.2	295.1	476.7	679.6	994.5	1,838.6
Female						
2001	104.7	170.0	261.3	461.7	753.4	1,758.9
2000[4]	107.7	182.6	291.3	483.5	741.0	1,743.5
1999[4]	99.9	192.1	293.4	484.9	768.3	1,787.8
Other heart diseases (I26-I51)						
All origins[3]						
Both sexes						
2001	62.0	100.7	164.7	287.2	537.0	1,475.5
2000[4]	64.3	103.6	167.1	290.2	542.8	1,509.4
1999[4]	66.5	105.2	175.0	295.4	553.5	1,512.9
Male						
2001	77.7	121.5	200.9	344.7	622.1	1,473.4
2000[4]	80.8	126.9	205.9	345.1	626.8	1,549.1
1999[4]	84.3	133.4	217.0	350.9	641.4	1,548.9
Female						
2001	47.8	82.8	135.9	247.0	486.0	1,476.4
2000[4]	49.3	83.6	136.6	252.1	493.2	1,493.2
1999[4]	50.5	81.0	142.2	256.8	502.0	1,498.5
Hispanic						
Both sexes						
2001	36.8	61.7	92.5	168.2	304.0	798.5
2000[4]	35.8	58.2	98.1	178.7	314.7	829.4
1999[4]	39.4	61.5	100.3	173.0	332.5	827.9
Male						
2001	45.9	75.5	111.2	190.3	359.7	775.2
2000[4]	44.9	70.8	124.5	194.9	362.2	890.8
1999[4]	51.2	79.6	132.4	213.0	377.2	888.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other heart diseases (I26-I51)														
Hispanic														
Female														
2001	13.9	8.7	*	*	*	*	*	1.4	2.5	2.8	3.6	5.3	8.9	17.6
2000[4]	14.0	10.6	*	*	*	*	1.4	1.5	2.3	3.7	5.0	6.3	10.5	16.1
1999[4]	13.9	14.1	*	*	*	1.4	*	*	2.4	4.0	4.5	6.6	9.7	18.6
Non-Hispanic[5]														
Both sexes														
2001	65.2	11.6	1.5	0.5	0.8	1.5	2.7	4.0	5.4	8.5	12.5	18.7	26.9	41.1
2000[4]	65.1	12.4	1.1	0.5	0.8	1.8	2.8	3.8	5.5	7.9	12.8	18.4	26.7	41.3
1999[4]	65.6	13.1	1.1	0.5	0.8	2.1	2.8	3.8	5.6	8.5	13.0	18.6	26.7	43.0
Male														
2001	57.8	11.0	1.6	0.4	0.8	1.9	3.3	4.7	6.4	10.6	15.6	24.0	33.8	51.2
2000[4]	58.0	12.7	1.2	0.6	0.9	2.1	3.3	4.4	6.6	9.6	16.1	24.1	34.3	53.3
1999[4]	58.6	13.8	1.3	0.5	0.8	2.6	3.2	4.3	6.7	10.8	16.2	24.0	34.2	55.7
Female														
2001	72.3	12.2	1.3	0.5	0.7	1.1	2.1	3.3	4.4	6.5	9.5	13.4	20.3	31.5
2000[4]	71.8	12.0	1.0	0.4	0.7	1.5	2.3	3.2	4.4	6.2	9.6	12.8	19.3	30.1
1999[4]	72.2	12.3	1.0	0.5	0.7	1.5	2.4	3.2	4.5	6.4	9.9	13.3	19.6	31.0
Non-Hispanic White														
Both sexes														
2001	70.6	9.1	1.1	0.4	0.7	1.3	2.1	3.1	4.3	6.8	10.1	15.4	23.0	35.7
2000[4]	70.3	10.5	0.9	0.4	0.6	1.4	2.1	2.9	4.4	6.2	10.6	15.2	22.6	36.7
1999[4]	70.4	10.5	0.9	0.4	0.6	1.8	2.2	2.9	4.4	6.8	10.5	15.1	22.8	37.2
Male														
2001	61.4	9.4	1.2	0.3	0.8	1.5	2.6	3.7	5.0	8.5	13.2	20.3	29.7	44.8
2000[4]	61.6	10.9	1.1	0.5	0.7	1.6	2.5	3.4	5.4	7.8	13.6	20.5	29.2	47.9
1999[4]	61.7	11.6	1.1	0.5	0.6	2.2	2.5	3.3	5.4	8.7	13.6	20.1	29.4	48.8
Female														
2001	79.4	8.8	1.0	0.5	0.6	1.0	1.6	2.6	3.5	5.1	7.0	10.6	16.5	27.1
2000[4]	78.7	10.1	0.8	*	0.5	1.2	1.6	2.4	3.4	4.7	7.6	9.9	16.2	26.1
1999[4]	78.8	9.4	0.8	0.4	0.6	1.4	2.0	2.4	3.5	4.9	7.5	10.1	16.3	26.2
Non-Hispanic Black														
Both sexes														
2001	54.7	21.3	3.0	0.8	1.2	3.0	5.8	9.1	12.8	19.1	30.0	44.0	62.3	93.6
2000[4]	54.6	19.6	2.0	0.8	1.4	3.8	6.3	9.1	12.4	18.5	29.1	44.3	63.6	87.8
1999[4]	56.2	23.2	2.0	0.8	1.3	3.7	5.9	9.0	13.0	20.1	30.8	45.8	63.5	98.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other heart diseases (I26-I51)						
Hispanic						
Female						
2001	28.9	50.5	78.2	152.5	269.4	809.8
2000[4]	28.0	48.0	78.1	167.3	285.8	799.3
1999[4]	29.3	46.9	76.2	144.8	305.7	798.6
Non-Hispanic[5]						
Both sexes						
2001	63.8	103.1	168.7	292.4	545.3	1,497.8
2000[4]	66.1	106.4	170.6	294.7	550.4	1,530.9
1999[4]	68.2	107.7	178.5	300.0	560.2	1,533.9
Male						
2001	79.7	124.2	205.6	351.5	632.2	1,499.0
2000[4]	83.0	130.2	209.8	351.4	636.0	1,573.6
1999[4]	86.2	136.5	220.7	356.0	649.4	1,571.2
Female						
2001	49.3	84.9	139.2	251.1	493.4	1,497.3
2000[4]	50.8	86.0	139.6	255.3	500.0	1,513.7
1999[4]	51.9	83.0	145.5	261.1	508.0	1,519.0
Non-Hispanic White						
Both sexes						
2001	58.7	96.3	162.8	286.7	547.4	1,533.2
2000[4]	60.6	99.8	163.9	288.8	553.5	1,564.2
1999[4]	61.7	100.8	171.0	293.1	560.5	1,565.9
Male						
2001	73.9	117.3	198.2	344.9	632.2	1,530.7
2000[4]	76.8	122.7	203.1	345.6	640.2	1,606.4
1999[4]	78.5	128.2	212.7	349.2	649.4	1,595.5
Female						
2001	44.6	77.7	134.0	245.4	496.2	1,534.2
2000[4]	45.6	79.8	132.3	248.8	502.0	1,547.2
1999[4]	46.2	76.7	137.8	253.7	508.0	1,554.1
Non-Hispanic Black						
Both sexes						
2001	121.1	182.4	258.0	408.2	608.1	1,292.9
2000[4]	125.0	185.1	270.0	412.7	597.8	1,319.4
1999[4]	135.9	188.6	287.1	420.9	627.5	1,314.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other heart diseases (I26-I51)														
Non-Hispanic Black														
Male														
2001	53.5	18.8	3.3	*	1.3	3.9	7.3	11.1	15.5	24.1	34.6	54.7	75.2	120.1
2000[4]	53.3	19.1	2.0	*	1.5	4.5	7.1	10.5	14.9	22.4	35.3	55.6	84.4	114.4
1999[4]	56.2	22.9	2.3	*	1.5	5.1	7.1	10.8	15.9	25.1	37.0	57.3	83.5	128.8
Female														
2001	55.8	23.8	2.7	*	*	2.1	4.3	7.2	10.4	14.7	26.0	34.8	51.3	71.9
2000[4]	55.8	20.1	1.9	*	1.3	3.1	5.5	7.8	10.2	15.0	23.6	34.5	45.9	66.0
1999[4]	56.3	23.6	1.8	*	*	2.2	4.7	7.4	10.4	15.7	25.4	35.8	46.5	73.6
Acute and subacute endocarditis (I33)														
All origins[3]														
Both sexes														
2001	0.4	*	*	*	*	*	*	*	0.1	0.2	0.3	0.4	0.5	0.5
2000[4]	0.4	*	*	*	*	*	*	*	0.1	0.2	0.3	0.4	0.6	0.7
1999[4]	0.4	*	*	*	*	*	*	*	*	0.2	0.3	0.3	0.4	0.7
Male														
2001	0.5	*	*	*	*	*	*	*	*	0.2	0.3	0.7	0.7	0.6
2000[4]	0.5	*	*	*	*	*	*	*	*	0.2	0.4	0.6	0.7	0.8
1999[4]	0.5	*	*	*	*	*	*	*	*	0.3	0.4	0.5	0.7	1.0
Female														
2001	0.4	*	*	*	*	*	*	*	*	0.2	0.3	0.2	0.3	0.5
2000[4]	0.4	*	*	*	*	*	*	*	*	*	0.2	0.3	0.4	0.5
1999[4]	0.3	*	*	*	*	*	*	*	*	0.2	*	*	0.2	0.5
Hispanic														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other heart diseases (I26-I51)						
Non-Hispanic Black						
Male						
2001	154.0	216.3	322.3	501.0	753.1	1,362.9
2000[4]	156.1	229.2	321.9	487.5	704.6	1,416.1
1999[4]	175.5	241.5	355.6	491.1	756.3	1,466.6
Female						
2001	95.3	157.3	215.2	352.8	534.3	1,267.0
2000[4]	100.5	152.6	235.5	367.6	544.0	1,283.4
1999[4]	105.0	149.4	241.7	378.6	563.7	1,257.0
Acute and subacute endocarditis (I33)						
All origins[3]						
Both sexes						
2001	0.8	1.1	1.7	2.2	3.0	2.5
2000[4]	0.7	1.1	1.3	2.2	2.8	2.8
1999[4]	0.7	1.1	1.5	2.1	2.7	2.9
Male						
2001	1.1	1.1	2.1	3.1	4.3	3.8
2000[4]	0.9	1.3	1.6	3.1	4.0	4.2
1999[4]	0.8	1.2	1.7	2.8	3.9	4.7
Female						
2001	0.5	1.0	1.3	1.6	2.2	2.0
2000[4]	0.6	0.9	1.1	1.6	2.0	2.3
1999[4]	0.7	1.0	1.4	1.7	2.0	2.1
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute and subacute endocarditis (I33)														
Non-Hispanic[5]														
Both sexes														
2001	0.5	*	*	*	*	*	*	*	*	0.2	0.3	0.5	0.5	0.5
2000[4]	0.4	*	*	*	*	*	*	*	0.1	0.2	0.3	0.5	0.5	0.7
1999[4]	0.4	*	*	*	*	*	*	*	*	0.2	0.3	0.4	0.5	0.7
Male														
2001	0.5	*	*	*	*	*	*	*	*	0.2	0.3	0.7	0.7	0.6
2000[4]	0.5	*	*	*	*	*	*	*	*	0.2	0.5	0.6	0.7	0.8
1999[4]	0.5	*	*	*	*	*	*	*	*	0.3	0.4	0.5	0.7	1.0
Female														
2001	0.4	*	*	*	*	*	*	*	*	0.2	0.3	0.3	0.3	0.4
2000[4]	0.4	*	*	*	*	*	*	*	*	*	0.2	0.3	0.4	0.5
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	0.3	0.5
Non-Hispanic White														
Both sexes														
2001	0.4	*	*	*	*	*	*	*	*	0.2	0.2	0.4	0.4	0.4
2000[4]	0.4	*	*	*	*	*	*	*	*	0.1	0.3	0.4	0.4	0.5
1999[4]	0.4	*	*	*	*	*	*	*	*	0.2	0.2	0.3	0.3	0.5
Male														
2001	0.5	*	*	*	*	*	*	*	*	*	0.3	0.5	0.5	0.5
2000[4]	0.5	*	*	*	*	*	*	*	*	*	0.4	0.5	0.5	0.7
1999[4]	0.5	*	*	*	*	*	*	*	*	*	0.2	0.4	0.4	0.7
Female														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	0.4
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	0.4
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	0.7	*	*	*	*	*	*	*	*	*	0.9	1.1	1.5	*
2000[4]	0.6	*	*	*	*	*	*	*	*	*	0.7	1.3	1.5	2.0
1999[4]	0.6	*	*	*	*	*	*	*	*	0.7	0.8	1.1	1.3	2.3
Male														
2001	0.8	*	*	*	*	*	*	*	*	*	*	1.9	*	*
2000[4]	0.7	*	*	*	*	*	*	*	*	*	*	1.9	*	*
1999[4]	0.7	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute and subacute endocarditis (I33)						
Non-Hispanic[5]						
Both sexes						
2001	0.8	1.1	1.7	2.2	3.1	2.6
2000[4]	0.7	1.1	1.3	2.2	2.8	2.9
1999[4]	0.7	1.1	1.5	2.1	2.8	2.9
Male						
2001	1.1	1.2	2.2	3.1	4.5	3.9
2000[4]	0.9	1.3	1.5	3.2	4.1	4.2
1999[4]	0.8	1.2	1.6	2.7	4.0	4.7
Female						
2001	0.6	1.1	1.3	1.5	2.2	2.0
2000[4]	0.6	0.9	1.1	1.6	2.0	2.3
1999[4]	0.7	1.0	1.4	1.7	2.1	2.2
Non-Hispanic White						
Both sexes						
2001	0.7	0.9	1.5	2.0	3.1	2.7
2000[4]	0.7	1.1	1.2	2.2	2.8	2.8
1999[4]	0.7	1.0	1.4	2.1	2.8	3.0
Male						
2001	0.9	0.9	1.8	3.0	4.5	4.0
2000[4]	0.9	1.2	1.6	3.3	4.0	4.3
1999[4]	0.8	1.2	1.5	2.9	4.0	5.1
Female						
2001	0.4	0.9	1.3	1.3	2.2	2.1
2000[4]	0.5	1.0	1.0	1.5	2.0	2.2
1999[4]	0.6	0.8	1.3	1.6	2.0	2.2
Non-Hispanic Black						
Both sexes						
2001	2.0	2.3	3.0	4.4	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	2.4	2.9	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute and subacute endocarditis (I33)														
Non-Hispanic Black														
Female														
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Diseases of pericardium and acute myocarditis (I30-I31,I40)														
All origins[3]														
Both sexes														
2001	0.3	0.8	0.2	*	*	*	0.1	0.1	0.1	0.2	0.2	0.3	0.3	0.4
2000[4]	0.3	0.5	0.2	*	*	*	0.1	0.1	0.1	0.2	0.2	0.2	0.2	0.3
1999[4]	0.3	0.7	0.2	*	*	*	0.1	*	0.2	0.2	0.3	0.3	0.3	0.5
Male														
2001	0.3	*	*	*	*	*	*	*	*	0.2	*	0.3	0.3	0.4
2000[4]	0.3	*	*	*	*	*	*	*	*	0.3	0.3	0.3	*	0.5
1999[4]	0.3	*	*	*	*	*	*	*	0.2	0.3	0.3	0.3	0.3	0.6
Female														
2001	0.3	*	*	*	*	*	*	*	*	*	0.3	0.3	0.3	0.4
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	0.2	*
1999[4]	0.3	*	*	*	*	*	*	*	*	0.2	0.3	0.3	0.2	0.3
Hispanic														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.3	0.9	0.2	*	*	*	0.1	*	0.1	0.2	0.2	0.3	0.3	0.4
2000[4]	0.3	*	*	*	*	*	*	0.1	0.1	0.2	0.2	0.2	0.2	0.3
1999[4]	0.3	0.8	0.2	*	*	*	*	*	0.1	0.2	0.3	0.3	0.3	0.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute and subacute endocarditis (I33)						
Non-Hispanic Black						
Female						
1999[4]	*	*	*	*	*	*
Diseases of pericardium and acute myocarditis (I30-I31,I40)						
All origins[3]						
Both sexes						
2001	0.5	0.5	0.8	1.2	1.3	1.9
2000[4]	0.4	0.5	0.9	1.1	1.3	1.8
1999[4]	0.6	0.7	0.9	1.2	1.5	2.0
Male						
2001	0.6	0.6	0.8	1.3	1.1	1.8
2000[4]	0.4	0.5	1.1	1.4	1.7	2.1
1999[4]	0.6	0.8	0.9	1.3	1.5	2.5
Female						
2001	*	*	0.9	1.1	1.4	2.0
2000[4]	0.4	0.5	0.7	0.9	1.0	1.7
1999[4]	0.6	0.7	0.8	1.2	1.5	1.9
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	0.5	0.5	0.8	1.2	1.3	1.9
2000[4]	0.4	0.5	0.9	1.1	1.2	1.8
1999[4]	0.6	0.7	0.9	1.2	1.4	2.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)														
Non-Hispanic[5]														
Male														
2001	0.3	*	*	*	*	*	*	*	*	0.2	*	0.3	0.3	0.4
2000[4]	0.3	*	*	*	*	*	*	*	*	0.3	0.3	0.3	*	0.5
1999[4]	0.3	*	*	*	*	*	*	*	*	0.3	0.3	0.3	0.3	0.7
Female														
2001	0.3	*	*	*	*	*	*	*	*	*	0.3	0.3	0.3	0.3
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	0.3	*
1999[4]	0.3	*	*	*	*	*	*	*	*	0.2	0.3	0.2	0.3	0.4
Non-Hispanic White														
Both sexes														
2001	0.3	*	*	*	*	*	*	*	*	0.2	0.2	0.3	0.2	0.3
2000[4]	0.3	*	*	*	*	*	*	*	*	0.1	0.2	0.2	0.2	0.3
1999[4]	0.3	*	*	*	*	*	*	*	*	0.2	0.2	0.3	0.3	0.5
Male														
2001	0.3	*	*	*	*	*	*	*	*	*	*	0.3	0.3	0.4
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	0.5
1999[4]	0.3	*	*	*	*	*	*	*	*	*	0.3	0.3	*	0.7
Female														
2001	0.3	*	*	*	*	*	*	*	*	*	*	0.3	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)						
Non-Hispanic[5]						
Male						
2001	0.6	0.7	0.8	1.3	1.2	1.8
2000[4]	*	0.6	1.2	1.4	1.5	2.1
1999[4]	0.6	0.8	0.9	1.3	1.5	2.6
Female						
2001	*	*	0.9	1.1	1.4	1.9
2000[4]	*	0.5	0.7	0.9	1.1	1.7
1999[4]	0.6	0.7	0.8	1.1	1.4	1.9
Non-Hispanic White						
Both sexes						
2001	0.5	0.4	0.9	1.2	1.3	1.9
2000[4]	0.3	0.5	0.9	1.0	1.3	1.8
1999[4]	0.6	0.7	0.8	1.2	1.4	2.0
Male						
2001	0.5	*	0.9	1.3	1.2	2.0
2000[4]	*	*	1.1	1.2	1.7	2.1
1999[4]	0.7	0.7	0.9	1.3	1.5	2.6
Female						
2001	*	*	0.9	1.2	1.4	1.8
2000[4]	*	0.6	0.7	0.9	1.1	1.7
1999[4]	0.6	0.7	0.8	1.1	1.4	1.8
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Heart failure (I50)														
All origins[3]														
Both sexes														
2001	20.0	0.6	*	*	*	*	*	0.2	0.2	0.5	0.8	1.7	3.3	6.5
2000[4]	19.8	0.8	*	*	*	0.1	*	0.1	0.2	0.3	0.8	1.4	3.0	6.1
1999[4]	19.7	*	*	*	*	*	*	0.1	0.2	0.4	0.8	1.5	2.9	6.5
Male														
2001	15.5	*	*	*	*	*	*	0.3	0.3	0.7	0.9	2.2	4.0	8.1
2000[4]	15.3	1.1	*	*	*	*	*	*	0.2	0.4	1.1	1.9	3.5	7.4
1999[4]	15.2	*	*	*	*	*	*	*	*	0.5	1.0	1.9	3.6	8.0
Female														
2001	24.3	*	*	*	*	*	*	*	0.2	0.4	0.7	1.2	2.6	5.1
2000[4]	24.1	*	*	*	*	*	*	*	*	0.3	0.6	0.9	2.5	5.0
1999[4]	24.0	*	*	*	*	*	*	*	*	0.3	0.6	1.1	2.1	5.1
Hispanic														
Both sexes														
2001	3.7	*	*	*	*	*	*	*	*	*	*	*	1.7	2.9
2000[4]	3.6	*	*	*	*	*	*	*	*	*	*	*	*	4.0
1999[4]	3.7	*	*	*	*	*	*	*	*	*	*	*	*	3.5
Male														
2001	3.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	2.9	*	*	*	*	*	*	*	*	*	*	*	*	5.3
1999[4]	3.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	4.5	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	4.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	4.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	22.4	*	*	*	*	*	*	0.2	0.3	0.5	0.9	1.8	3.4	6.8
2000[4]	22.1	0.8	*	*	*	0.1	*	0.2	0.2	0.4	0.9	1.5	3.2	6.3
1999[4]	21.8	*	*	*	*	*	*	0.2	0.2	0.4	0.9	1.6	3.0	6.7
Male														
2001	17.4	*	*	*	*	*	*	0.3	0.3	0.7	1.0	2.3	4.2	8.4
2000[4]	17.2	*	*	*	*	*	*	*	0.2	0.4	1.1	2.1	3.8	7.5
1999[4]	16.9	*	*	*	*	*	*	*	*	0.5	1.1	2.0	3.8	8.2
Female														
2001	27.1	*	*	*	*	*	*	*	*	0.4	0.8	1.3	2.7	5.2
2000[4]	26.7	*	*	*	*	*	*	*	*	0.3	0.6	0.9	2.6	5.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Heart failure (I50)						
All origins[3]						
Both sexes						
2001	13.0	23.8	45.6	92.6	199.2	674.6
2000[4]	12.8	24.0	46.5	93.5	199.9	684.9
1999[4]	11.7	23.5	48.3	94.3	201.6	681.9
Male						
2001	15.0	28.0	54.2	107.9	226.6	646.4
2000[4]	15.3	28.2	55.6	107.9	229.5	672.2
1999[4]	14.2	28.0	58.9	108.4	224.9	667.0
Female						
2001	11.3	20.2	38.7	81.9	182.7	686.4
2000[4]	10.5	20.3	39.3	83.5	182.4	690.0
1999[4]	9.5	19.6	39.9	84.5	188.0	687.9
Hispanic						
Both sexes						
2001	9.1	14.3	24.4	50.4	112.2	349.5
2000[4]	8.3	12.2	27.2	52.3	108.1	369.6
1999[4]	7.6	12.3	28.6	53.9	109.3	377.4
Male						
2001	9.5	19.5	24.7	47.1	129.9	333.3
2000[4]	10.9	16.4	31.6	52.4	117.8	380.9
1999[4]	8.3	15.5	36.7	67.9	113.2	358.0
Female						
2001	8.7	10.0	24.1	52.8	101.3	357.4
2000[4]	6.0	8.8	23.9	52.3	102.1	364.0
1999[4]	6.9	9.7	22.6	44.0	106.9	386.8
Non-Hispanic[5]						
Both sexes						
2001	13.3	24.4	46.8	94.5	202.2	685.6
2000[4]	13.1	24.7	47.5	95.2	202.8	694.8
1999[4]	11.9	24.1	49.2	95.8	204.5	691.0
Male						
2001	15.3	28.5	55.8	110.7	230.4	658.0
2000[4]	15.5	28.9	56.8	110.3	233.4	683.3
1999[4]	14.5	28.7	60.0	109.9	228.4	677.1
Female						
2001	11.5	20.8	39.6	83.2	185.4	697.0
2000[4]	10.9	21.1	40.1	84.7	184.8	699.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Heart failure (I50)														
Non-Hispanic[5]														
Female														
1999[4]	26.5	*	*	*	*	*	*	*	*	0.3	0.7	1.2	2.2	5.3
Non-Hispanic White														
Both sexes														
2001	25.1	*	*	*	*	*	*	0.2	0.2	0.4	0.6	1.4	2.8	5.9
2000[4]	24.7	*	*	*	*	*	*	*	0.2	0.3	0.7	1.1	2.7	5.5
1999[4]	24.3	*	*	*	*	*	*	*	*	0.3	0.7	1.2	2.3	5.7
Male														
2001	19.3	*	*	*	*	*	*	*	*	0.5	0.8	1.9	3.5	7.5
2000[4]	19.0	*	*	*	*	*	*	*	*	0.3	1.0	1.7	3.2	6.9
1999[4]	18.6	*	*	*	*	*	*	*	*	0.4	0.9	1.5	3.0	7.3
Female														
2001	30.7	*	*	*	*	*	*	*	*	0.3	0.5	0.8	2.1	4.3
2000[4]	30.2	*	*	*	*	*	*	*	*	*	0.4	0.6	2.1	4.2
1999[4]	29.9	*	*	*	*	*	*	*	*	0.3	0.4	0.8	1.7	4.2
Non-Hispanic Black														
Both sexes														
2001	14.3	*	*	*	*	*	*	*	*	1.2	2.8	4.7	8.9	16.0
2000[4]	14.1	*	*	*	*	*	*	*	*	1.1	2.3	4.3	7.9	14.4
1999[4]	14.3	*	*	*	*	*	*	*	*	1.0	2.4	4.9	8.6	16.6
Male														
2001	12.0	*	*	*	*	*	*	*	*	2.0	2.7	5.2	10.7	19.3
2000[4]	11.6	*	*	*	*	*	*	*	*	*	2.6	5.7	9.3	15.2
1999[4]	12.2	*	*	*	*	*	*	*	*	*	2.7	6.5	11.3	19.3
Female														
2001	16.4	*	*	*	*	*	*	*	*	*	2.9	4.2	7.3	13.3
2000[4]	16.3	*	*	*	*	*	*	*	*	*	2.0	3.1	6.6	13.7
1999[4]	16.2	*	*	*	*	*	*	*	*	*	2.2	3.5	6.3	14.4
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)														
All origins[3]														
Both sexes														
2001	37.9	9.7	1.1	0.4	0.7	1.3	2.3	3.3	4.6	7.0	10.5	15.4	21.9	32.3
2000[4]	38.3	10.6	0.9	0.4	0.6	1.5	2.4	3.2	4.8	6.8	10.9	15.5	22.1	33.1
1999[4]	39.0	11.9	0.9	0.4	0.7	1.8	2.4	3.2	4.9	7.4	11.1	15.7	22.3	34.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Heart failure (I50)						
Non-Hispanic[5]						
Female						
1999[4]	9.6	20.2	40.8	86.1	190.5	696.5
Non-Hispanic White						
Both sexes						
2001	12.2	22.9	45.5	93.9	204.3	706.8
2000[4]	12.0	23.3	45.8	94.5	204.9	714.3
1999[4]	11.1	22.6	47.7	94.0	205.5	709.1
Male						
2001	14.4	26.9	54.4	110.5	232.1	679.1
2000[4]	14.2	27.8	55.5	110.1	235.8	701.6
1999[4]	13.5	26.8	59.1	108.8	229.9	690.8
Female						
2001	10.2	19.4	38.2	82.2	187.4	718.4
2000[4]	9.9	19.3	38.1	83.6	186.6	719.4
1999[4]	8.8	18.9	38.7	83.6	191.1	716.4
Non-Hispanic Black						
Both sexes						
2001	25.1	42.1	72.3	122.2	216.6	551.2
2000[4]	24.8	42.0	73.4	121.4	211.8	564.8
1999[4]	22.4	43.0	74.9	131.6	218.4	557.4
Male						
2001	26.8	50.5	87.1	141.7	260.8	547.5
2000[4]	30.0	46.4	84.1	137.3	246.3	589.4
1999[4]	27.6	52.6	84.5	146.9	241.8	609.7
Female						
2001	23.8	35.8	62.4	110.6	194.1	552.6
2000[4]	20.6	38.7	66.2	111.8	194.4	555.6
1999[4]	18.3	35.8	68.5	122.3	206.8	537.9
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)						
All origins[3]						
Both sexes						
2001	47.8	75.3	116.6	191.2	333.6	796.5
2000[4]	50.4	78.0	118.4	193.4	338.9	820.0
1999[4]	53.5	79.9	124.4	197.7	347.6	826.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)														
All origins[3]														
Male														
2001	35.6	9.1	1.2	0.3	0.7	1.7	2.8	3.8	5.4	8.8	13.4	19.9	27.7	40.3
2000[4]	36.1	10.5	1.0	0.5	0.8	1.9	2.8	3.8	5.7	8.3	13.6	20.2	28.9	43.4
1999[4]	37.1	12.1	1.0	0.4	0.8	2.2	2.7	3.7	5.9	9.3	13.9	20.2	28.7	44.5
Female														
2001	40.2	10.4	1.0	0.4	0.6	1.0	1.7	2.7	3.7	5.3	7.6	11.1	16.3	24.8
2000[4]	40.4	10.7	0.7	0.3	0.5	1.2	2.0	2.6	3.8	5.3	8.2	11.0	15.6	23.4
1999[4]	40.9	11.6	0.8	0.4	0.5	1.4	2.0	2.7	3.9	5.4	8.3	11.2	16.2	24.3
Hispanic														
Both sexes														
2001	9.3	8.0	0.8	*	*	1.2	1.2	1.8	2.7	3.6	5.1	7.5	11.2	17.8
2000[4]	9.6	8.7	0.7	*	0.6	1.4	1.8	2.1	3.0	4.1	5.9	7.8	13.4	18.4
1999[4]	9.8	11.5	0.9	*	*	1.3	1.4	1.8	3.3	4.7	6.5	7.9	13.1	19.3
Male														
2001	9.3	7.9	*	*	*	1.3	1.6	2.2	3.3	4.4	7.3	10.4	15.5	22.0
2000[4]	9.9	8.1	*	*	*	2.0	2.2	2.8	3.8	4.9	7.3	10.0	18.7	25.0
1999[4]	10.3	9.8	*	*	*	1.3	1.7	2.5	4.3	5.8	8.9	10.5	18.1	23.6
Female														
2001	9.2	8.0	*	*	*	*	*	1.3	1.9	2.7	2.9	4.7	7.2	13.9
2000[4]	9.3	9.3	*	*	*	*	1.4	*	2.1	3.2	4.5	5.5	8.5	12.5
1999[4]	9.2	13.3	*	*	*	*	*	*	2.1	3.5	4.0	5.2	8.4	15.4
Non-Hispanic[5]														
Both sexes														
2001	42.1	10.1	1.2	0.3	0.7	1.3	2.4	3.6	4.9	7.5	11.1	16.2	22.7	33.4
2000[4]	42.3	11.0	0.9	0.4	0.6	1.6	2.5	3.4	5.0	7.2	11.4	16.2	22.7	34.0
1999[4]	43.0	11.9	0.9	0.4	0.7	1.9	2.6	3.4	5.2	7.7	11.6	16.3	23.0	35.0
Male														
2001	39.6	9.3	1.4	0.3	0.7	1.7	3.1	4.2	5.8	9.4	14.1	20.8	28.6	41.7
2000[4]	40.0	10.9	1.0	0.5	0.7	1.9	3.0	3.9	6.1	8.8	14.3	21.1	29.7	44.5
1999[4]	40.9	12.7	1.0	0.4	0.8	2.4	2.9	3.9	6.2	9.7	14.4	21.1	29.5	45.8
Female														
2001	44.5	11.0	1.0	0.4	0.6	1.0	1.8	3.0	4.0	5.7	8.1	11.7	17.1	25.6
2000[4]	44.5	11.0	0.8	0.2	0.5	1.3	2.1	2.9	4.0	5.6	8.6	11.4	16.1	24.2
1999[4]	44.9	11.1	0.9	0.4	0.6	1.5	2.2	3.0	4.2	5.7	8.7	11.7	16.9	24.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)						
All origins[3]						
Male						
2001	61.1	91.7	143.7	232.4	390.0	821.3
2000[4]	64.2	96.8	147.6	232.6	391.5	870.7
1999[4]	68.7	103.5	155.5	238.5	411.1	874.7
Female						
2001	35.7	61.2	95.0	162.4	299.7	786.1
2000[4]	37.8	61.9	95.4	166.0	307.9	799.3
1999[4]	39.8	59.7	100.1	169.4	310.5	806.5
Hispanic						
Both sexes						
2001	26.6	46.5	66.3	114.7	188.7	444.6
2000[4]	26.5	44.2	68.3	123.7	203.9	457.2
1999[4]	31.2	47.7	69.3	114.9	219.1	447.6
Male						
2001	34.5	55.3	83.7	139.1	229.8	438.1
2000[4]	33.4	52.6	89.9	140.3	237.0	505.9
1999[4]	42.6	63.0	92.2	139.7	259.4	526.2
Female						
2001	19.7	39.3	53.0	97.3	163.3	447.7
2000[4]	20.6	37.4	51.9	111.9	183.7	433.3
1999[4]	21.3	35.3	52.1	97.5	195.1	409.7
Non-Hispanic[5]						
Both sexes						
2001	49.2	77.1	119.3	194.6	338.8	807.8
2000[4]	52.0	80.1	120.9	196.2	343.5	831.4
1999[4]	54.9	81.7	126.9	200.9	351.5	837.9
Male						
2001	62.8	93.9	146.8	236.4	396.2	835.3
2000[4]	66.2	99.4	150.3	236.4	396.9	883.9
1999[4]	70.3	105.8	158.2	242.1	415.5	886.8
Female						
2001	36.9	62.6	97.4	165.3	304.4	796.3
2000[4]	39.0	63.5	97.7	168.2	312.1	810.2
1999[4]	41.0	61.1	102.4	172.3	314.0	818.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)														
Non-Hispanic White														
Both sexes														
2001	44.7	8.1	0.9	0.3	0.6	1.1	1.9	2.8	3.9	6.1	9.1	13.4	19.6	29.1
2000[4]	44.9	9.6	0.8	0.3	0.5	1.2	1.9	2.6	4.1	5.7	9.5	13.5	19.3	30.4
1999[4]	45.4	9.8	0.8	0.4	0.6	1.7	2.1	2.6	4.1	6.1	9.5	13.4	19.9	30.5
Male														
2001	41.3	8.1	1.0	*	0.7	1.3	2.4	3.4	4.5	7.7	12.0	17.6	25.4	36.4
2000[4]	41.7	9.6	0.8	0.4	0.6	1.4	2.3	3.1	5.0	7.1	12.0	18.2	25.2	39.8
1999[4]	42.4	10.8	0.9	0.4	0.6	2.0	2.3	3.1	5.0	7.9	12.2	17.9	25.7	40.1
Female														
2001	47.9	8.1	0.7	*	0.5	0.8	1.4	2.2	3.2	4.4	6.1	9.2	13.9	22.1
2000[4]	47.9	9.6	0.7	*	0.3	1.0	1.5	2.1	3.1	4.3	6.9	8.9	13.6	21.4
1999[4]	48.2	8.7	0.6	0.4	0.6	1.4	1.9	2.2	3.3	4.3	6.7	9.0	14.2	21.3
Non-Hispanic Black														
Both sexes														
2001	39.3	18.2	2.4	*	1.1	2.7	5.1	8.3	11.7	17.0	25.8	37.7	51.0	75.4
2000[4]	39.5	16.1	1.5	0.7	1.1	3.4	5.7	8.3	11.3	16.5	25.7	38.1	53.9	70.9
1999[4]	40.9	20.9	1.5	0.6	1.1	3.4	5.4	8.2	11.9	18.0	26.9	39.3	52.9	78.7
Male														
2001	40.3	16.0	2.7	*	1.2	3.5	6.6	9.7	14.1	21.1	30.8	46.8	61.6	98.2
2000[4]	40.5	15.0	1.7	*	1.2	4.0	6.4	9.5	13.7	20.0	31.2	47.4	72.9	96.8
1999[4]	42.8	20.8	*	*	1.5	4.8	6.5	9.7	14.5	22.5	32.6	48.9	68.9	105.2
Female														
2001	38.5	20.5	2.1	*	*	1.8	3.7	6.9	9.6	13.3	21.4	29.7	42.1	56.7
2000[4]	38.6	17.3	*	*	*	2.7	5.1	7.1	9.0	13.4	20.9	30.0	37.6	49.6
1999[4]	39.2	21.1	*	*	*	2.0	4.4	6.8	9.6	14.1	22.0	31.1	39.3	57.1
Essential (primary) hypertension and hypertensive renal disease (I10,I12)														
All origins[3]														
Both sexes														
2001	6.8	*	*	*	*	*	*	0.2	0.3	0.5	1.0	1.9	3.0	4.4
2000[4]	6.4	*	*	*	*	*	*	0.1	0.2	0.5	1.1	1.8	3.0	4.4
1999[4]	6.1	*	*	*	*	*	*	0.1	0.3	0.5	0.9	1.7	2.8	4.1
Male														
2001	5.2	*	*	*	*	*	*	0.2	0.4	0.5	1.3	2.3	3.7	5.2
2000[4]	5.0	*	*	*	*	*	*	*	0.3	0.5	1.4	2.3	3.8	5.4
1999[4]	4.7	*	*	*	*	*	*	*	0.3	0.6	1.1	2.0	3.5	5.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)						
Non-Hispanic White						
Both sexes						
2001	45.3	72.0	114.9	189.5	338.8	821.8
2000[4]	47.6	75.0	115.9	191.0	344.5	845.3
1999[4]	49.4	76.5	121.0	195.8	350.7	851.7
Male						
2001	58.0	88.9	141.1	230.1	394.4	845.6
2000[4]	61.4	93.3	144.9	231.0	398.7	898.3
1999[4]	63.6	99.5	151.1	236.2	413.9	897.0
Female						
2001	33.5	57.0	93.6	160.7	305.2	811.9
2000[4]	34.9	58.9	92.6	162.9	312.3	824.0
1999[4]	36.2	56.3	97.1	167.5	313.5	833.7
Non-Hispanic Black						
Both sexes						
2001	93.3	137.0	182.1	280.5	386.8	737.9
2000[4]	98.3	140.3	193.5	287.0	382.5	748.9
1999[4]	111.2	142.1	208.0	285.7	405.3	752.7
Male						
2001	123.3	161.0	229.6	352.7	484.9	813.1
2000[4]	124.0	177.8	235.1	344.9	454.0	821.9
1999[4]	145.4	185.7	266.2	342.7	509.2	855.7
Female						
2001	69.7	119.3	150.5	237.4	336.9	710.1
2000[4]	78.0	112.7	165.8	252.1	346.5	721.7
1999[4]	84.3	109.9	169.3	251.3	353.9	714.2
Essential (primary) hypertension and hypertensive renal disease (I10,I12)						
All origins[3]						
Both sexes						
2001	7.5	12.2	19.0	35.5	65.2	171.9
2000[4]	7.8	11.5	19.1	33.8	63.0	162.9
1999[4]	7.3	12.0	18.6	32.7	59.9	152.1
Male						
2001	8.1	13.7	20.0	36.0	61.3	150.8
2000[4]	9.1	11.5	20.6	34.5	62.9	142.4
1999[4]	8.2	13.4	20.1	33.1	59.4	132.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Essential (primary) hypertension and hypertensive renal disease (I10,I12)														
All origins[3]														
Female														
2001	8.3	*	*	*	*	*	*	*	0.3	0.4	0.7	1.4	2.3	3.8
2000[4]	7.8	*	*	*	*	*	*	*	0.2	0.4	0.8	1.3	2.2	3.5
1999[4]	7.4	*	*	*	*	*	*	*	*	0.3	0.7	1.3	2.1	3.3
Hispanic														
Both sexes														
2001	2.4	*	*	*	*	*	*	*	*	*	*	1.5	2.2	2.9
2000[4]	2.3	*	*	*	*	*	*	*	*	*	*	*	2.7	3.5
1999[4]	2.2	*	*	*	*	*	*	*	*	*	*	1.7	2.7	3.2
Male														
2001	1.9	*	*	*	*	*	*	*	*	*	*	2.2	*	*
2000[4]	2.1	*	*	*	*	*	*	*	*	*	*	*	4.2	4.4
1999[4]	1.9	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	2.9	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	2.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	2.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	7.4	*	*	*	*	*	*	0.2	0.4	0.5	1.1	1.9	3.0	4.5
2000[4]	7.0	*	*	*	*	*	*	*	0.3	0.5	1.1	1.8	3.0	4.5
1999[4]	6.6	*	*	*	*	*	*	0.1	0.3	0.5	1.0	1.7	2.8	4.2
Male														
2001	5.7	*	*	*	*	*	*	*	0.4	0.6	1.3	2.3	3.8	5.3
2000[4]	5.4	*	*	*	*	*	*	*	0.3	0.5	1.4	2.3	3.8	5.4
1999[4]	5.1	*	*	*	*	*	*	*	0.4	0.7	1.2	2.0	3.5	5.1
Female														
2001	9.0	*	*	*	*	*	*	*	0.4	0.4	0.8	1.4	2.3	3.8
2000[4]	8.5	*	*	*	*	*	*	*	0.2	0.4	0.8	1.3	2.2	3.5
1999[4]	8.0	*	*	*	*	*	*	*	*	0.4	0.8	1.3	2.1	3.2
Non-Hispanic White														
Both sexes														
2001	7.0	*	*	*	*	*	*	*	0.2	0.3	0.5	1.0	1.8	2.9
2000[4]	6.5	*	*	*	*	*	*	*	*	0.2	0.6	1.0	1.7	2.9
1999[4]	6.1	*	*	*	*	*	*	*	0.1	0.2	0.5	1.0	1.5	2.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Essential (primary) hypertension and hypertensive renal disease (I10,I12)						
All origins[3]						
Female						
2001	6.9	11.0	18.2	35.2	67.5	180.8
2000[4]	6.7	11.5	17.9	33.4	63.0	171.3
1999[4]	6.5	10.8	17.5	32.4	60.2	160.0
Hispanic						
Both sexes						
2001	6.6	11.4	22.8	38.2	67.5	160.1
2000[4]	10.0	10.8	20.3	36.1	63.5	148.0
1999[4]	7.3	12.3	19.0	35.5	71.7	129.8
Male						
2001	5.7	10.6	24.7	39.6	72.7	129.5
2000[4]	13.0	9.7	22.4	42.1	73.6	141.1
1999[4]	8.6	16.2	21.4	43.7	67.6	131.6
Female						
2001	7.3	12.1	21.3	37.3	64.3	175.0
2000[4]	7.4	11.8	18.8	31.9	57.3	151.3
1999[4]	6.2	9.1	17.2	29.7	74.1	128.9
Non-Hispanic[5]						
Both sexes						
2001	7.5	12.3	18.7	35.2	65.0	172.0
2000[4]	7.6	11.5	18.9	33.5	62.9	162.9
1999[4]	7.3	11.9	18.6	32.5	59.1	152.3
Male						
2001	8.3	13.8	19.6	35.6	60.6	151.1
2000[4]	8.7	11.5	20.4	33.9	62.4	142.0
1999[4]	8.1	13.2	20.0	32.6	58.5	131.9
Female						
2001	6.8	10.9	18.0	35.0	67.6	180.7
2000[4]	6.6	11.5	17.8	33.3	63.2	171.3
1999[4]	6.5	10.9	17.4	32.5	59.5	160.5
Non-Hispanic White						
Both sexes						
2001	5.2	8.9	14.7	30.3	58.1	162.8
2000[4]	5.0	8.4	14.9	28.3	57.0	152.7
1999[4]	4.8	8.8	14.9	27.1	52.2	141.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Essential (primary) hypertension and hypertensive renal disease (I10,I12)														
Non-Hispanic White														
Male														
2001	5.1	*	*	*	*	*	*	*	*	0.3	0.7	1.3	2.2	3.5
2000[4]	4.9	*	*	*	*	*	*	*	*	0.3	0.8	1.3	2.3	3.7
1999[4]	4.6	*	*	*	*	*	*	*	*	0.3	0.6	1.3	1.9	3.6
Female														
2001	8.8	*	*	*	*	*	*	*	*	*	0.3	0.7	1.3	2.3
2000[4]	8.1	*	*	*	*	*	*	*	*	*	0.4	0.6	1.1	2.2
1999[4]	7.6	*	*	*	*	*	*	*	*	*	0.3	0.7	1.1	1.8
Non-Hispanic Black														
Both sexes														
2001	11.3	*	*	*	*	*	*	*	1.4	1.9	4.5	7.9	12.9	18.3
2000[4]	11.0	*	*	*	*	*	*	*	1.1	2.1	4.2	7.3	13.1	18.3
1999[4]	10.7	*	*	*	*	*	*	*	1.0	2.3	4.1	6.5	13.2	17.3
Male														
2001	10.0	*	*	*	*	*	*	*	*	2.3	5.5	10.2	16.7	21.9
2000[4]	9.5	*	*	*	*	*	*	*	*	2.2	5.4	9.0	15.9	22.9
1999[4]	9.4	*	*	*	*	*	*	*	*	2.8	4.8	8.1	17.1	19.9
Female														
2001	12.4	*	*	*	*	*	*	*	1.4	1.5	3.5	5.9	9.6	15.4
2000[4]	12.4	*	*	*	*	*	*	*	*	2.1	3.2	5.9	10.6	14.5
1999[4]	11.8	*	*	*	*	*	*	*	*	1.9	3.5	5.1	9.8	15.2
Cerebrovascular diseases (I60-I69)														
All origins[3]														
Both sexes														
2001	57.4	2.7	0.4	0.2	0.2	0.3	0.7	1.1	1.9	3.8	7.2	12.1	18.5	28.8
2000[4]	59.6	3.3	0.3	0.1	0.2	0.3	0.7	1.1	1.9	3.9	7.7	13.1	19.3	30.8
1999[4]	60.0	2.7	0.3	0.1	0.2	0.3	0.6	1.0	1.8	4.0	7.5	12.5	18.4	31.7
Male														
2001	45.2	3.1	0.3	0.2	0.2	0.3	0.7	1.1	2.0	3.9	7.4	13.2	20.5	32.8
2000[4]	46.9	3.8	*	*	0.2	0.4	0.7	1.1	1.9	3.8	7.8	14.2	21.3	35.2
1999[4]	47.1	3.3	0.3	*	*	0.3	0.6	1.0	1.9	3.9	7.8	13.3	20.7	35.7
Female														
2001	69.2	2.3	0.4	*	0.2	0.3	0.6	1.0	1.9	3.6	7.1	10.9	16.5	25.1
2000[4]	71.8	2.7	0.4	*	0.3	0.3	0.7	1.0	1.9	4.0	7.5	12.0	17.3	26.6
1999[4]	72.3	2.1	0.3	*	0.2	0.3	0.6	1.0	1.8	4.1	7.1	11.7	16.1	28.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Essential (primary) hypertension and hypertensive renal disease (I10,I12)						
Non-Hispanic White						
Male						
2001	5.9	10.2	15.6	30.8	53.4	142.1
2000[4]	5.9	8.7	16.3	29.1	56.5	131.8
1999[4]	5.6	10.0	16.2	27.5	51.2	119.8
Female						
2001	4.5	7.8	14.0	29.9	60.9	171.5
2000[4]	4.2	8.0	13.8	27.8	57.2	161.1
1999[4]	4.1	7.8	13.7	26.9	52.8	150.5
Non-Hispanic Black						
Both sexes						
2001	27.6	41.2	57.4	90.7	150.1	291.5
2000[4]	29.5	41.0	59.9	93.8	132.8	285.9
1999[4]	27.3	38.7	58.3	96.0	140.3	272.2
Male						
2001	30.9	49.5	63.4	95.8	161.1	277.2
2000[4]	35.3	40.8	65.3	93.9	139.9	268.9
1999[4]	30.9	43.8	63.4	98.6	156.8	268.6
Female						
2001	25.0	35.0	53.4	87.7	144.5	296.9
2000[4]	25.0	41.1	56.3	93.8	129.2	292.2
1999[4]	24.4	34.8	54.9	94.4	132.1	273.6
Cerebrovascular diseases (I60-I69)						
All origins[3]						
Both sexes						
2001	49.6	85.7	164.3	320.2	622.4	1,500.2
2000[4]	53.8	91.0	169.1	332.8	654.0	1,589.2
1999[4]	51.4	91.1	173.5	340.6	664.0	1,614.8
Male						
2001	57.2	98.5	187.9	349.0	654.6	1,380.2
2000[4]	62.3	102.6	192.9	367.6	695.1	1,484.3
1999[4]	60.0	102.8	197.7	376.9	706.8	1,512.8
Female						
2001	42.7	74.7	145.6	300.0	603.2	1,550.4
2000[4]	46.0	81.0	150.3	308.5	629.7	1,632.0
1999[4]	43.7	81.1	154.5	315.3	639.0	1,656.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Cerebrovascular diseases (I60-I69)														
Hispanic														
Both sexes														
2001	17.4	*	*	*	*	*	0.8	1.0	2.0	3.4	7.0	12.4	20.3	28.6
2000[4]	17.5	*	*	*	*	*	0.8	1.1	2.3	3.4	7.5	13.8	17.0	28.2
1999[4]	17.4	*	*	*	*	*	0.7	1.4	1.8	3.3	6.9	13.6	18.3	31.4
Male														
2001	15.7	*	*	*	*	*	*	1.0	2.3	3.7	7.3	14.6	24.1	34.2
2000[4]	15.8	*	*	*	*	*	*	1.2	2.6	3.6	7.8	16.2	20.5	37.7
1999[4]	16.1	*	*	*	*	*	*	1.4	2.3	3.4	8.6	16.2	23.8	36.3
Female														
2001	19.1	*	*	*	*	*	*	*	1.6	3.1	6.6	10.2	16.7	23.5
2000[4]	19.4	*	*	*	*	*	*	*	1.9	3.1	7.1	11.4	13.8	19.6
1999[4]	18.8	*	*	*	*	*	*	1.4	*	3.1	5.1	11.0	13.1	26.9
Non-Hispanic[5]														
Both sexes														
2001	63.2	2.8	0.3	0.2	0.2	0.3	0.6	1.1	1.9	3.8	7.3	12.0	18.2	28.8
2000[4]	65.4	3.6	0.3	0.1	0.3	0.4	0.7	1.1	1.8	3.9	7.6	12.9	19.4	30.9
1999[4]	65.7	2.9	0.3	0.1	0.2	0.3	0.6	1.0	1.8	4.1	7.5	12.3	18.3	31.6
Male														
2001	49.7	3.3	0.3	*	*	0.3	0.6	1.1	1.9	3.9	7.3	13.1	20.2	32.5
2000[4]	51.5	4.2	*	*	*	0.4	0.7	1.1	1.8	3.8	7.7	13.9	21.3	34.9
1999[4]	51.5	3.7	0.3	*	*	0.3	0.6	0.9	1.8	4.0	7.7	12.9	20.4	35.5
Female														
2001	76.1	2.4	0.4	*	*	0.3	0.6	1.0	2.0	3.7	7.2	10.9	16.4	25.2
2000[4]	78.7	2.8	0.4	*	0.3	0.3	0.7	1.0	1.9	4.1	7.6	12.0	17.6	27.1
1999[4]	79.2	2.0	*	*	0.2	0.3	0.6	1.0	1.8	4.2	7.3	11.7	16.3	28.0
Non-Hispanic White														
Both sexes														
2001	67.6	1.9	0.3	0.2	0.2	0.3	0.5	0.9	1.6	3.0	5.4	8.8	13.0	23.4
2000[4]	70.0	2.6	0.3	*	0.2	0.4	0.6	1.0	1.4	2.9	5.6	9.2	15.1	25.0
1999[4]	70.4	2.4	0.3	*	0.2	0.3	0.6	0.8	1.5	3.2	5.5	8.8	14.0	25.6
Male														
2001	52.0	2.2	*	*	*	0.3	0.6	0.8	1.7	2.9	5.6	9.6	14.4	26.0
2000[4]	53.9	3.3	*	*	*	0.5	0.6	1.1	1.4	2.9	5.7	10.0	16.4	27.5
1999[4]	53.9	3.1	*	*	*	0.3	0.5	0.8	1.5	3.1	5.7	9.0	15.3	28.7
Female														
2001	82.6	*	*	*	*	0.3	0.4	0.9	1.6	3.0	5.1	8.0	11.6	20.8
2000[4]	85.5	1.9	*	*	*	*	0.6	0.9	1.4	3.0	5.5	8.3	13.8	22.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cerebrovascular diseases (I60-I69)						
Hispanic						
Both sexes						
2001	44.5	74.7	145.5	271.5	463.4	1,042.9
2000[4]	54.6	83.1	151.1	262.9	491.8	1,082.9
1999[4]	45.6	90.5	150.2	265.7	498.7	1,077.6
Male						
2001	56.2	95.3	168.4	311.0	529.8	1,005.6
2000[4]	63.3	99.6	183.8	310.8	556.5	1,029.9
1999[4]	58.7	105.5	187.4	330.0	526.6	1,136.6
Female						
2001	34.3	58.0	128.1	243.5	422.2	1,061.2
2000[4]	47.1	69.8	126.3	229.0	452.4	1,108.9
1999[4]	34.4	78.4	122.2	220.2	482.0	1,049.2
Non-Hispanic[5]						
Both sexes						
2001	49.8	86.3	165.1	321.7	627.7	1,514.0
2000[4]	53.5	91.3	169.6	335.1	658.3	1,604.5
1999[4]	51.6	90.9	174.2	343.0	668.9	1,630.4
Male						
2001	56.9	98.5	188.7	350.0	658.1	1,392.9
2000[4]	62.0	102.6	192.7	369.1	698.8	1,499.9
1999[4]	59.7	102.3	197.6	377.8	712.2	1,524.8
Female						
2001	43.2	75.8	146.3	302.0	609.6	1,564.4
2000[4]	45.7	81.7	151.4	311.4	634.5	1,646.8
1999[4]	44.2	81.1	155.9	318.9	643.6	1,672.7
Non-Hispanic White						
Both sexes						
2001	41.1	74.9	153.2	310.4	621.9	1,525.0
2000[4]	45.6	81.4	158.8	323.1	651.1	1,616.8
1999[4]	43.0	81.8	161.2	331.1	662.9	1,648.0
Male						
2001	46.9	85.6	174.7	337.7	651.5	1,399.0
2000[4]	52.8	90.8	179.5	355.4	689.2	1,505.9
1999[4]	49.6	91.7	182.4	363.1	705.6	1,535.1
Female						
2001	35.7	65.4	135.7	291.1	604.0	1,577.4
2000[4]	38.8	73.2	142.2	300.4	628.4	1,661.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Cerebrovascular diseases (I60-I69)														
Non-Hispanic White														
Female														
1999[4]	86.1	*	*	*	*	*	0.7	0.8	1.4	3.2	5.3	8.6	12.7	22.6
Non-Hispanic Black														
Both sexes														
2001	52.9	6.9	*	*	*	*	1.4	2.3	3.6	8.3	18.9	33.1	56.2	71.9
2000[4]	54.3	8.0	*	*	*	*	1.1	1.6	4.5	9.9	20.2	38.0	52.4	78.6
1999[4]	53.9	4.7	*	*	*	*	0.8	1.9	3.8	9.4	20.3	35.1	51.3	79.6
Male														
2001	46.2	8.3	*	*	*	*	*	2.8	3.4	8.9	18.2	37.3	65.4	89.2
2000[4]	47.7	9.2	*	*	*	*	*	*	4.2	9.5	21.0	41.5	61.7	99.4
1999[4]	47.5	*	*	*	*	*	*	2.0	3.8	9.2	21.1	39.8	62.3	94.3
Female														
2001	59.0	*	*	*	*	*	1.8	1.8	3.9	7.7	19.6	29.6	48.4	57.8
2000[4]	60.3	*	*	*	*	*	*	1.7	4.7	10.2	19.5	34.9	44.5	61.7
1999[4]	59.8	*	*	*	*	*	*	1.8	3.7	9.6	19.5	31.1	42.0	67.7
Atherosclerosis (I70)														
All origins[3]														
Both sexes														
2001	4.9	*	*	*	*	*	*	*	*	*	0.1	0.3	0.7	1.4
2000[4]	5.1	*	*	*	*	*	*	*	*	*	0.1	0.4	0.6	1.3
1999[4]	5.4	*	*	*	*	*	*	*	*	*	0.1	0.3	0.8	1.3
Male														
2001	3.7	*	*	*	*	*	*	*	*	*	*	0.4	0.8	1.7
2000[4]	3.9	*	*	*	*	*	*	*	*	*	0.2	0.5	0.8	1.7
1999[4]	4.0	*	*	*	*	*	*	*	*	*	0.2	0.4	1.0	1.8
Female														
2001	6.2	*	*	*	*	*	*	*	*	*	*	0.2	0.5	1.1
2000[4]	6.3	*	*	*	*	*	*	*	*	*	*	0.2	0.4	0.9
1999[4]	6.7	*	*	*	*	*	*	*	*	*	*	*	0.5	0.9
Hispanic														
Both sexes														
2001	1.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.2	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cerebrovascular diseases (I60-I69)						
Non-Hispanic White						
Female						
1999[4]	37.0	73.2	144.3	308.7	637.7	1,693.0
Non-Hispanic Black						
Both sexes						
2001	119.3	181.5	281.4	465.4	725.7	1,465.5
2000[4]	115.6	178.0	284.5	479.0	775.9	1,545.8
1999[4]	119.8	172.9	312.5	493.4	762.8	1,501.5
Male						
2001	145.9	222.5	340.9	530.4	776.0	1,359.4
2000[4]	138.8	216.4	343.4	548.2	872.8	1,479.5
1999[4]	151.1	207.7	372.2	565.0	824.5	1,455.9
Female						
2001	98.4	151.2	241.8	426.6	700.1	1,504.8
2000[4]	97.3	149.6	245.5	437.3	727.1	1,570.4
1999[4]	95.2	147.1	272.9	450.2	732.3	1,518.6
Atherosclerosis (I70)						
All origins[3]						
Both sexes						
2001	2.5	4.9	9.8	21.9	49.2	179.1
2000[4]	2.6	5.2	10.5	22.7	52.4	188.8
1999[4]	2.9	5.1	11.2	24.1	53.6	202.0
Male						
2001	3.2	6.2	12.1	25.0	53.2	167.2
2000[4]	3.6	6.6	13.1	27.5	59.8	176.5
1999[4]	3.8	6.3	13.3	28.1	60.0	189.0
Female						
2001	1.8	3.9	8.1	19.7	46.8	184.1
2000[4]	1.7	4.0	8.4	19.4	48.1	193.8
1999[4]	2.2	4.0	9.5	21.4	49.8	207.2
Hispanic						
Both sexes						
2001	*	*	6.3	17.0	33.8	130.7
2000[4]	*	4.0	8.2	16.2	39.5	132.7
1999[4]	*	*	8.5	19.4	41.1	129.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Atherosclerosis (I70)														
Hispanic														
Male														
2001	0.9	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	1.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	5.5	*	*	*	*	*	*	*	*	*	0.1	0.3	0.6	1.4
2000[4]	5.7	*	*	*	*	*	*	*	*	*	0.2	0.4	0.6	1.3
1999[4]	5.9	*	*	*	*	*	*	*	*	*	0.1	0.3	0.8	1.3
Male														
2001	4.1	*	*	*	*	*	*	*	*	*	*	0.5	0.8	1.7
2000[4]	4.3	*	*	*	*	*	*	*	*	*	0.3	0.5	0.8	1.8
1999[4]	4.4	*	*	*	*	*	*	*	*	*	0.2	0.4	1.0	1.8
Female														
2001	6.8	*	*	*	*	*	*	*	*	*	*	0.2	0.5	1.1
2000[4]	7.0	*	*	*	*	*	*	*	*	*	*	0.2	0.4	0.9
1999[4]	7.4	*	*	*	*	*	*	*	*	*	*	*	0.5	0.9
Non-Hispanic White														
Both sexes														
2001	6.3	*	*	*	*	*	*	*	*	*	0.1	0.3	0.6	1.4
2000[4]	6.5	*	*	*	*	*	*	*	*	*	0.1	0.3	0.5	1.3
1999[4]	6.7	*	*	*	*	*	*	*	*	*	*	0.3	0.7	1.2
Male														
2001	4.7	*	*	*	*	*	*	*	*	*	*	0.4	0.7	1.7
2000[4]	4.9	*	*	*	*	*	*	*	*	*	*	0.4	0.7	1.8
1999[4]	4.9	*	*	*	*	*	*	*	*	*	*	0.4	1.0	1.6
Female														
2001	7.9	*	*	*	*	*	*	*	*	*	*	*	0.5	1.1
2000[4]	8.0	*	*	*	*	*	*	*	*	*	*	*	0.3	0.8
1999[4]	8.5	*	*	*	*	*	*	*	*	*	*	*	0.4	0.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerosis (I70)						
Hispanic						
Male						
2001	*	*	*	21.1	39.0	127.6
2000[4]	*	*	12.6	25.1	45.6	104.8
1999[4]	*	*	11.2	23.4	47.2	151.0
Female						
2001	*	*	*	14.0	30.6	132.2
2000[4]	*	*	*	*	35.8	146.4
1999[4]	*	*	*	16.5	37.5	118.5
Non-Hispanic[5]						
Both sexes						
2001	2.6	5.1	10.0	22.0	49.6	180.5
2000[4]	2.7	5.2	10.6	23.0	52.8	190.3
1999[4]	3.0	5.2	11.3	24.3	53.8	203.9
Male						
2001	3.4	6.4	12.4	25.0	53.6	168.2
2000[4]	3.7	6.7	13.1	27.5	60.2	178.7
1999[4]	3.8	6.4	13.4	28.2	60.2	189.9
Female						
2001	1.9	4.0	8.2	19.9	47.3	185.5
2000[4]	1.7	4.0	8.6	19.8	48.5	195.1
1999[4]	2.2	4.1	9.7	21.5	50.1	209.6
Non-Hispanic White						
Both sexes						
2001	2.4	5.1	10.0	22.4	50.8	187.4
2000[4]	2.5	5.3	10.5	23.3	53.4	196.8
1999[4]	2.9	5.1	11.4	24.3	54.9	209.9
Male						
2001	3.2	6.4	12.4	25.1	54.4	175.2
2000[4]	3.6	6.6	13.3	28.3	60.8	183.6
1999[4]	3.7	6.3	13.3	27.9	61.0	194.6
Female						
2001	1.6	4.0	8.1	20.4	48.6	192.6
2000[4]	1.5	4.1	8.3	19.8	49.1	202.1
1999[4]	2.1	4.0	9.8	21.9	51.4	215.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Atherosclerosis (I70)														
Non-Hispanic Black														
Both sexes														
2001	2.7	*	*	*	*	*	*	*	*	*	*	*	1.0	2.0
2000[4]	3.0	*	*	*	*	*	*	*	*	*	*	*	1.5	2.1
1999[4]	3.1	*	*	*	*	*	*	*	*	*	*	*	1.3	2.0
Male														
2001	2.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	2.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	2.6	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	3.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	3.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	3.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Other diseases of circulatory system (I71-I78)														
All origins[3]														
Both sexes														
2001	8.9	*	*	*	*	0.1	0.2	0.4	0.5	0.9	1.3	2.0	3.3	6.0
2000[4]	9.3	*	*	*	*	*	0.2	0.3	0.5	0.7	1.3	2.1	3.5	6.4
1999[4]	9.2	*	*	*	*	*	0.2	0.3	0.4	0.8	1.4	2.1	3.3	6.7
Male														
2001	9.6	*	*	*	*	*	0.2	0.4	0.7	1.2	1.8	2.8	4.4	8.2
2000[4]	10.2	*	*	*	*	*	0.2	0.4	0.6	0.8	1.7	2.8	4.9	9.3
1999[4]	10.1	*	*	*	*	*	0.2	0.4	0.6	1.0	1.6	3.0	4.4	9.4
Female														
2001	8.2	*	*	*	*	*	*	0.3	0.3	0.6	0.9	1.2	2.2	3.9
2000[4]	8.3	*	*	*	*	*	*	*	0.4	0.6	0.9	1.5	2.1	3.7
1999[4]	8.4	*	*	*	*	*	*	*	0.2	0.5	1.1	1.3	2.1	4.1
Hispanic														
Both sexes														
2001	2.0	*	*	*	*	*	*	*	*	*	1.1	1.6	2.8	3.7
2000[4]	2.3	*	*	*	*	*	*	*	*	*	0.9	*	2.7	4.7
1999[4]	2.1	*	*	*	*	*	*	*	*	*	0.9	*	1.8	4.3
Male														
2001	2.2	*	*	*	*	*	*	*	*	*	1.6	2.1	3.6	4.4
2000[4]	2.5	*	*	*	*	*	*	*	*	*	*	*	3.6	6.1
1999[4]	2.4	*	*	*	*	*	*	*	*	*	*	*	*	5.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerosis (I70)						
Non-Hispanic Black						
Both sexes						
2001	5.2	6.3	12.1	23.2	42.5	126.1
2000[4]	4.8	5.8	13.1	23.6	54.6	140.2
1999[4]	4.8	7.0	12.2	27.6	47.0	155.8
Male						
2001	6.1	7.8	15.1	30.8	48.2	113.2
2000[4]	6.2	8.8	14.0	25.0	67.8	150.3
1999[4]	5.9	9.0	15.9	37.7	56.7	166.4
Female						
2001	4.4	5.2	10.1	18.7	39.6	130.9
2000[4]	3.7	*	12.5	22.7	48.0	136.5
1999[4]	3.9	5.5	9.6	21.5	42.1	151.9
Other diseases of circulatory system (I71-I78)						
All origins[3]						
Both sexes						
2001	12.1	22.3	36.6	60.6	88.1	158.2
2000[4]	12.7	24.5	40.6	62.9	92.2	163.2
1999[4]	12.5	23.5	40.1	63.5	94.3	165.2
Male						
2001	16.8	30.8	51.2	81.0	122.5	197.3
2000[4]	18.0	34.6	57.6	88.6	126.6	214.9
1999[4]	18.2	33.5	56.7	89.1	129.3	207.5
Female						
2001	7.8	15.0	25.1	46.3	67.4	141.8
2000[4]	7.9	15.7	27.2	45.0	71.9	142.2
1999[4]	7.3	15.0	27.2	45.7	73.7	148.2
Hispanic						
Both sexes						
2001	6.2	11.3	16.8	35.4	45.2	95.0
2000[4]	7.2	14.7	27.0	39.5	47.9	102.8
1999[4]	7.6	12.6	21.2	31.3	67.0	110.8
Male						
2001	9.0	16.7	24.2	51.8	55.8	108.6
2000[4]	10.4	19.0	34.5	59.8	66.3	131.0
1999[4]	10.1	17.0	29.0	44.5	97.5	163.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other diseases of circulatory system (I71-I78)														
Hispanic														
Female														
2001	1.8	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	2.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.9	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	9.9	*	*	*	*	*	0.2	0.4	0.5	0.9	1.4	2.1	3.3	6.2
2000[4]	10.2	*	*	*	*	*	0.2	0.2	0.5	0.7	1.3	2.2	3.5	6.5
1999[4]	10.2	*	*	*	*	*	0.2	0.3	0.5	0.8	1.4	2.2	3.4	6.9
Male														
2001	10.7	*	*	*	*	*	0.3	0.5	0.7	1.3	1.8	2.9	4.4	8.5
2000[4]	11.3	*	*	*	*	*	0.3	0.4	0.7	0.9	1.7	2.9	5.0	9.5
1999[4]	11.2	*	*	*	*	*	*	0.4	0.7	1.1	1.7	3.1	4.6	9.7
Female														
2001	9.1	*	*	*	*	*	*	0.3	0.3	0.6	0.9	1.2	2.2	3.9
2000[4]	9.2	*	*	*	*	*	*	*	0.4	0.5	1.0	1.5	2.1	3.7
1999[4]	9.3	*	*	*	*	*	*	*	0.2	0.6	1.1	1.4	2.2	4.2
Non-Hispanic White														
Both sexes														
2001	10.8	*	*	*	*	*	0.2	0.4	0.5	0.8	1.1	1.8	2.9	5.7
2000[4]	11.2	*	*	*	*	*	0.2	0.2	0.5	0.6	1.0	1.9	3.0	6.1
1999[4]	11.3	*	*	*	*	*	0.2	0.3	0.5	0.8	1.2	1.9	3.0	6.5
Male														
2001	11.8	*	*	*	*	*	*	0.5	0.6	1.1	1.4	2.5	3.8	8.2
2000[4]	12.6	*	*	*	*	*	*	*	0.7	0.7	1.3	2.7	4.4	9.0
1999[4]	12.5	*	*	*	*	*	*	0.4	0.7	1.0	1.5	2.8	4.1	9.5
Female														
2001	9.9	*	*	*	*	*	*	*	0.3	0.5	0.8	1.1	1.9	3.4
2000[4]	9.9	*	*	*	*	*	*	*	*	0.5	0.8	1.1	1.7	3.3
1999[4]	10.1	*	*	*	*	*	*	*	*	0.5	0.9	1.1	2.0	3.6
Non-Hispanic Black														
Both sexes														
2001	7.3	*	*	*	*	*	*	*	0.9	1.8	3.3	4.1	6.6	10.9
2000[4]	7.3	*	*	*	*	*	*	*	1.0	1.2	3.2	4.6	6.9	11.3
1999[4]	7.0	*	*	*	*	*	*	*	*	1.2	2.9	4.7	6.3	11.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of circulatory system (I71-I78)						
Hispanic						
Female						
2001	*	6.9	11.2	23.7	38.6	88.4
2000[4]	*	11.2	21.4	25.1	36.7	89.0
1999[4]	5.4	9.1	15.3	22.0	48.8	85.3
Non-Hispanic[5]						
Both sexes						
2001	12.5	23.0	37.7	61.7	89.6	160.1
2000[4]	13.0	25.0	41.3	63.8	93.7	165.0
1999[4]	12.8	24.1	41.1	64.8	95.0	166.8
Male						
2001	17.3	31.6	52.6	82.2	124.9	200.5
2000[4]	18.5	35.6	58.7	89.6	128.8	217.9
1999[4]	18.7	34.3	58.0	91.1	130.2	209.0
Female						
2001	8.1	15.6	25.9	47.3	68.4	143.3
2000[4]	8.1	16.0	27.5	45.8	73.0	143.6
1999[4]	7.5	15.3	27.8	46.6	74.5	149.9
Non-Hispanic White						
Both sexes						
2001	12.1	23.3	38.0	61.8	90.0	160.1
2000[4]	12.7	25.2	42.1	64.0	94.6	164.8
1999[4]	12.9	24.4	41.8	65.5	95.6	166.7
Male						
2001	16.9	32.5	53.3	83.4	126.6	204.7
2000[4]	18.4	36.6	60.3	90.5	132.0	222.7
1999[4]	19.1	35.4	59.8	93.2	131.9	212.8
Female						
2001	7.7	15.3	25.5	46.5	67.9	141.6
2000[4]	7.5	15.2	27.5	45.3	72.3	141.5
1999[4]	7.2	14.8	27.5	46.1	74.1	148.3
Non-Hispanic Black						
Both sexes						
2001	18.2	23.2	43.5	69.8	97.1	175.7
2000[4]	18.6	27.8	40.1	69.9	93.1	180.0
1999[4]	13.8	26.5	39.1	64.9	97.4	180.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other diseases of circulatory system (I71-I78)														
Non-Hispanic Black														
Male														
2001	7.2	*	*	*	*	*	*	*	*	2.5	4.6	5.7	9.6	13.2
2000[4]	7.2	*	*	*	*	*	*	*	*	1.7	4.0	5.6	10.2	15.2
1999[4]	6.7	*	*	*	*	*	*	*	*	1.6	3.2	6.3	8.9	14.3
Female														
2001	7.4	*	*	*	*	*	*	*	*	*	2.1	2.7	4.1	9.0
2000[4]	7.5	*	*	*	*	*	*	*	*	*	2.4	3.8	4.2	8.0
1999[4]	7.2	*	*	*	*	*	*	*	*	*	2.6	3.3	4.1	9.1
Aortic aneurysm and dissection (I71)														
All origins[3]														
Both sexes														
2001	5.3	*	*	*	*	*	0.1	0.2	0.3	0.6	0.9	1.4	2.0	3.7
2000[4]	5.6	*	*	*	*	*	0.1	0.2	0.3	0.5	0.9	1.3	2.2	4.3
1999[4]	5.7	*	*	*	*	*	0.1	0.2	0.3	0.6	0.9	1.4	2.1	4.4
Male														
2001	6.5	*	*	*	*	*	*	0.4	0.5	0.9	1.4	2.1	3.2	5.8
2000[4]	7.0	*	*	*	*	*	*	0.3	0.5	0.6	1.3	1.9	3.5	6.7
1999[4]	7.0	*	*	*	*	*	*	0.3	0.5	0.9	1.2	2.2	3.3	6.8
Female														
2001	4.2	*	*	*	*	*	*	*	*	0.3	0.3	0.6	0.9	1.8
2000[4]	4.3	*	*	*	*	*	*	*	0.2	0.3	0.5	0.8	1.0	2.1
1999[4]	4.4	*	*	*	*	*	*	*	*	0.3	0.5	0.7	1.0	2.1
Hispanic														
Both sexes														
2001	1.1	*	*	*	*	*	*	*	*	*	*	*	1.6	1.9
2000[4]	1.3	*	*	*	*	*	*	*	*	*	*	*	2.1	2.7
1999[4]	1.1	*	*	*	*	*	*	*	*	*	*	*	*	2.6
Male														
2001	1.5	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.6	*	*	*	*	*	*	*	*	*	*	*	3.0	*
1999[4]	1.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.7	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.8	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of circulatory system (I71-I78)						
Non-Hispanic Black						
Male						
2001	24.0	28.1	56.5	78.7	129.2	168.6
2000[4]	23.0	33.6	52.3	91.5	106.4	179.7
1999[4]	18.1	32.4	49.9	75.4	118.7	184.2
Female						
2001	13.6	19.6	34.9	64.5	80.8	178.4
2000[4]	15.1	23.6	32.0	56.9	86.4	180.1
1999[4]	10.4	22.0	31.9	58.5	86.9	178.6
Aortic aneurysm and dissection (I71)						
All origins[3]						
Both sexes						
2001	7.8	15.0	23.8	39.0	55.4	77.5
2000[4]	8.4	16.3	27.8	41.0	56.5	79.2
1999[4]	8.1	16.5	27.3	42.2	58.2	80.3
Male						
2001	12.0	22.2	36.0	56.3	82.5	114.8
2000[4]	13.1	24.9	41.8	61.9	84.9	127.3
1999[4]	13.0	25.3	41.0	63.3	88.6	125.3
Female						
2001	3.9	8.8	14.0	26.9	39.2	61.9
2000[4]	4.1	8.9	16.7	26.5	39.7	59.6
1999[4]	3.7	9.0	16.6	27.5	40.4	62.2
Hispanic						
Both sexes						
2001	3.4	7.1	9.5	21.2	25.8	31.9
2000[4]	4.3	7.7	15.7	25.4	30.1	41.8
1999[4]	4.3	7.3	11.6	18.4	34.1	40.7
Male						
2001	5.7	12.4	16.5	35.5	33.8	53.3
2000[4]	6.6	12.3	20.9	42.1	42.7	64.5
1999[4]	7.1	11.6	18.8	24.2	53.4	79.8
Female						
2001	*	*	*	11.1	20.9	21.4
2000[4]	*	*	11.8	13.6	22.4	30.7
1999[4]	*	*	*	14.3	22.5	21.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Aortic aneurysm and dissection (I71)														
Non-Hispanic[5]														
Both sexes														
2001	6.0	*	*	*	*	*	0.2	0.3	0.4	0.7	0.9	1.4	2.1	3.9
2000[4]	6.2	*	*	*	*	*	0.1	0.2	0.4	0.5	0.9	1.4	2.2	4.5
1999[4]	6.3	*	*	*	*	*	0.1	0.2	0.3	0.6	0.9	1.5	2.2	4.5
Male														
2001	7.3	*	*	*	*	*	*	0.4	0.5	1.0	1.4	2.2	3.2	6.0
2000[4]	7.8	*	*	*	*	*	*	0.3	0.5	0.7	1.4	2.0	3.5	6.9
1999[4]	7.8	*	*	*	*	*	*	0.3	0.6	0.9	1.3	2.3	3.4	7.0
Female														
2001	4.7	*	*	*	*	*	*	*	*	0.3	0.4	0.7	0.9	1.9
2000[4]	4.7	*	*	*	*	*	*	*	*	0.3	0.5	0.8	1.0	2.2
1999[4]	4.8	*	*	*	*	*	*	*	*	0.3	0.6	0.7	1.0	2.1
Non-Hispanic White														
Both sexes														
2001	6.7	*	*	*	*	*	0.2	0.3	0.3	0.6	0.7	1.2	1.7	3.7
2000[4]	7.0	*	*	*	*	*	*	*	0.3	0.4	0.7	1.2	2.0	4.2
1999[4]	7.1	*	*	*	*	*	*	0.2	0.3	0.6	0.8	1.2	1.9	4.3
Male														
2001	8.2	*	*	*	*	*	*	0.4	0.5	0.9	1.1	1.9	2.8	5.9
2000[4]	8.8	*	*	*	*	*	*	*	0.5	0.6	1.1	1.9	3.2	6.7
1999[4]	8.9	*	*	*	*	*	*	*	0.5	0.9	1.0	2.0	3.0	7.0
Female														
2001	5.2	*	*	*	*	*	*	*	*	*	0.3	0.5	0.7	1.6
2000[4]	5.2	*	*	*	*	*	*	*	*	*	0.4	0.5	0.8	1.9
1999[4]	5.3	*	*	*	*	*	*	*	*	0.3	0.5	0.5	0.9	1.8
Non-Hispanic Black														
Both sexes														
2001	3.5	*	*	*	*	*	*	*	*	1.2	2.1	2.8	4.5	6.2
2000[4]	3.6	*	*	*	*	*	*	*	*	0.9	2.2	2.9	3.7	7.2
1999[4]	3.4	*	*	*	*	*	*	*	*	0.8	2.1	3.6	4.1	6.4
Male														
2001	3.9	*	*	*	*	*	*	*	*	1.8	3.5	4.1	7.0	8.5
2000[4]	4.0	*	*	*	*	*	*	*	*	*	3.3	3.6	6.1	10.3
1999[4]	3.7	*	*	*	*	*	*	*	*	*	2.8	5.5	6.6	9.1
Female														
2001	3.1	*	*	*	*	*	*	*	*	*	*	1.7	2.3	4.4
2000[4]	3.3	*	*	*	*	*	*	*	*	*	*	2.4	*	4.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Aortic aneurysm and dissection (I71)						
Non-Hispanic[5]						
Both sexes						
2001	8.1	15.5	24.5	39.8	56.4	79.0
2000[4]	8.7	16.8	28.4	41.6	57.3	80.3
1999[4]	8.4	17.0	28.1	43.2	58.9	81.7
Male						
2001	12.4	22.8	37.0	57.2	84.2	117.0
2000[4]	13.5	25.7	42.9	62.6	86.5	129.6
1999[4]	13.3	26.1	42.1	65.0	89.7	127.0
Female						
2001	4.1	9.2	14.6	27.7	39.8	63.2
2000[4]	4.3	9.2	16.9	27.0	40.2	60.4
1999[4]	3.9	9.3	17.1	28.0	40.9	63.5
Non-Hispanic White						
Both sexes						
2001	8.1	16.1	25.5	40.7	58.1	81.3
2000[4]	8.8	17.3	29.7	42.6	59.1	82.0
1999[4]	8.8	17.7	29.2	44.7	60.7	83.7
Male						
2001	12.3	23.9	38.5	58.7	87.4	122.9
2000[4]	13.8	27.0	45.0	64.3	90.3	135.3
1999[4]	14.1	27.4	44.2	68.3	92.7	131.8
Female						
2001	4.1	9.2	14.9	27.8	40.5	64.0
2000[4]	4.1	8.8	17.4	27.4	40.5	60.5
1999[4]	3.8	9.2	17.3	28.1	41.8	64.5
Non-Hispanic Black						
Both sexes						
2001	9.5	11.6	18.6	34.5	43.3	55.9
2000[4]	9.5	14.6	20.5	34.1	41.1	63.0
1999[4]	6.7	14.2	20.0	29.8	40.8	61.1
Male						
2001	14.8	14.9	26.5	42.7	55.6	49.7
2000[4]	13.2	18.1	28.4	47.7	46.3	70.5
1999[4]	9.6	19.4	25.6	30.0	54.9	77.3
Female						
2001	5.3	9.2	13.3	29.7	37.1	58.2
2000[4]	6.5	12.0	15.2	25.8	38.5	60.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Aortic aneurysm and dissection (I71)														
Non-Hispanic Black														
Female														
1999[4]	3.1	*	*	*	*	*	*	*	*	*	1.5	2.0	*	4.3
Other diseases of arteries, arterioles and capillaries (I72-I78)														
All origins[3]														
Both sexes														
2001	3.5	*	*	*	*	*	*	0.1	0.2	0.3	0.5	0.6	1.2	2.2
2000[4]	3.6	*	*	*	*	*	*	*	0.2	0.2	0.4	0.8	1.3	2.1
1999[4]	3.6	*	*	*	*	*	*	*	0.1	0.2	0.5	0.7	1.2	2.3
Male														
2001	3.1	*	*	*	*	*	*	*	*	0.2	0.4	0.7	1.2	2.4
2000[4]	3.2	*	*	*	*	*	*	*	*	0.2	0.3	0.8	1.4	2.6
1999[4]	3.1	*	*	*	*	*	*	*	*	*	0.4	0.8	1.2	2.6
Female														
2001	4.0	*	*	*	*	*	*	*	*	0.3	0.5	0.6	1.3	2.0
2000[4]	4.0	*	*	*	*	*	*	*	*	0.3	0.5	0.7	1.1	1.6
1999[4]	4.1	*	*	*	*	*	*	*	*	0.2	0.5	0.6	1.2	2.1
Hispanic														
Both sexes														
2001	0.9	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.8	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.9	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.9	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	1.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	3.9	*	*	*	*	*	*	*	0.2	0.3	0.5	0.6	1.2	2.3
2000[4]	4.0	*	*	*	*	*	*	*	0.2	0.2	0.4	0.8	1.3	2.1
1999[4]	3.9	*	*	*	*	*	*	*	0.2	0.2	0.5	0.7	1.2	2.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Aortic aneurysm and dissection (I71)						
Non-Hispanic Black						
Female						
1999[4]	4.4	10.4	16.3	29.7	33.8	55.1
Other diseases of arteries, arterioles and capillaries (I72-I78)						
All origins[3]						
Both sexes						
2001	4.3	7.3	12.9	21.5	32.7	80.7
2000[4]	4.3	8.2	12.8	21.8	35.8	84.0
1999[4]	4.4	7.0	12.9	21.3	36.1	84.9
Male						
2001	4.8	8.5	15.2	24.7	40.0	82.5
2000[4]	4.9	9.7	15.7	26.6	41.7	87.6
1999[4]	5.2	8.1	15.7	25.9	40.8	82.2
Female						
2001	3.9	6.3	11.1	19.3	28.3	79.9
2000[4]	3.7	6.9	10.5	18.5	32.2	82.6
1999[4]	3.6	6.0	10.6	18.2	33.3	86.0
Hispanic						
Both sexes						
2001	2.8	4.1	7.3	14.2	19.4	63.2
2000[4]	2.9	7.0	11.3	14.1	17.8	61.0
1999[4]	3.3	5.4	9.6	12.9	32.9	70.2
Male						
2001	*	*	*	16.4	*	55.2
2000[4]	*	*	13.6	17.7	*	66.5
1999[4]	*	*	10.2	20.3	44.0	84.1
Female						
2001	*	*	7.0	12.6	17.7	67.0
2000[4]	*	7.2	9.6	11.5	*	58.4
1999[4]	*	*	9.2	*	26.3	63.4
Non-Hispanic[5]						
Both sexes						
2001	4.4	7.5	13.2	21.9	33.1	81.1
2000[4]	4.4	8.2	12.9	22.2	36.3	84.7
1999[4]	4.4	7.1	13.0	21.7	36.1	85.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other diseases of arteries, arterioles and capillaries (I72-I78)														
Non-Hispanic[5]														
Male														
2001	3.4	*	*	*	*	*	*	*	*	0.3	0.4	0.7	1.2	2.5
2000[4]	3.5	*	*	*	*	*	*	*	*	*	0.3	0.9	1.5	2.6
1999[4]	3.4	*	*	*	*	*	*	*	*	*	0.4	0.8	1.3	2.7
Female														
2001	4.4	*	*	*	*	*	*	*	*	0.3	0.6	0.6	1.2	2.1
2000[4]	4.4	*	*	*	*	*	*	*	*	0.3	0.5	0.7	1.1	1.6
1999[4]	4.5	*	*	*	*	*	*	*	*	0.2	0.6	0.6	1.2	2.1
Non-Hispanic White														
Both sexes														
2001	4.2	*	*	*	*	*	*	*	*	0.2	0.4	0.6	1.1	2.1
2000[4]	4.3	*	*	*	*	*	*	*	0.2	0.2	0.3	0.7	1.0	1.9
1999[4]	4.2	*	*	*	*	*	*	*	0.1	0.2	0.5	0.7	1.1	2.1
Male														
2001	3.6	*	*	*	*	*	*	*	*	*	0.3	0.6	1.0	2.3
2000[4]	3.8	*	*	*	*	*	*	*	*	*	*	0.8	1.2	2.4
1999[4]	3.6	*	*	*	*	*	*	*	*	*	0.4	0.9	1.2	2.5
Female														
2001	4.7	*	*	*	*	*	*	*	*	0.3	0.5	0.5	1.2	1.8
2000[4]	4.7	*	*	*	*	*	*	*	*	0.3	0.4	0.6	0.9	1.3
1999[4]	4.8	*	*	*	*	*	*	*	*	*	0.5	0.6	1.1	1.8
Non-Hispanic Black														
Both sexes														
2001	3.8	*	*	*	*	*	*	*	*	*	1.2	1.3	2.2	4.6
2000[4]	3.7	*	*	*	*	*	*	*	*	*	0.9	1.7	3.2	4.0
1999[4]	3.6	*	*	*	*	*	*	*	*	*	0.8	1.0	2.2	5.0
Male														
2001	3.3	*	*	*	*	*	*	*	*	*	*	*	2.6	4.7
2000[4]	3.2	*	*	*	*	*	*	*	*	*	*	2.0	4.1	4.9
1999[4]	3.0	*	*	*	*	*	*	*	*	*	*	*	*	5.2
Female														
2001	4.4	*	*	*	*	*	*	*	*	*	*	*	*	4.6
2000[4]	4.2	*	*	*	*	*	*	*	*	*	*	*	2.4	3.3
1999[4]	4.2	*	*	*	*	*	*	*	*	*	*	*	*	4.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)						
Non-Hispanic[5]						
Male						
2001	4.9	8.8	15.6	25.0	40.7	83.5
2000[4]	4.9	9.9	15.8	27.0	42.3	88.2
1999[4]	5.3	8.3	15.9	26.1	40.5	82.0
Female						
2001	4.0	6.4	11.3	19.6	28.6	80.1
2000[4]	3.8	6.8	10.6	18.8	32.8	83.3
1999[4]	3.6	6.0	10.7	18.6	33.6	86.4
Non-Hispanic White						
Both sexes						
2001	4.0	7.3	12.5	21.1	31.9	78.8
2000[4]	3.9	7.8	12.4	21.3	35.5	82.8
1999[4]	4.2	6.7	12.6	20.9	34.8	83.0
Male						
2001	4.5	8.6	14.8	24.6	39.2	81.8
2000[4]	4.5	9.5	15.3	26.2	41.7	87.4
1999[4]	5.1	8.0	15.6	25.0	39.2	81.0
Female						
2001	3.6	6.1	10.6	18.7	27.5	77.6
2000[4]	3.4	6.4	10.1	17.9	31.8	81.0
1999[4]	3.4	5.5	10.2	18.0	32.2	83.9
Non-Hispanic Black						
Both sexes						
2001	8.7	11.6	24.9	35.3	53.8	119.9
2000[4]	9.1	13.2	19.6	35.9	52.0	117.0
1999[4]	7.1	12.2	19.1	35.1	56.6	119.0
Male						
2001	9.2	13.3	30.0	36.0	73.6	119.0
2000[4]	9.8	15.5	23.9	43.8	60.1	109.2
1999[4]	8.5	13.0	24.2	45.4	63.8	107.0
Female						
2001	8.4	10.4	21.6	34.8	43.7	120.2
2000[4]	8.6	11.6	16.8	31.1	48.0	119.9
1999[4]	6.0	11.6	15.6	28.8	53.1	123.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other disorders of circulatory system (I80-I99)														
All origins[3]														
Both sexes														
2001	1.6	0.5	*	*	*	*	0.1	0.2	0.4	0.4	0.8	1.1	1.3	1.9
2000[4]	1.6	0.7	*	*	*	*	0.1	0.3	0.4	0.6	0.9	1.2	1.5	1.7
1999[4]	1.4	0.6	*	*	*	*	*	0.2	0.3	0.4	0.8	1.0	1.0	1.6
Male														
2001	1.4	*	*	*	*	*	*	*	0.3	0.5	0.9	1.2	1.5	2.2
2000[4]	1.4	*	*	*	*	*	*	*	0.4	0.6	1.1	1.5	1.7	1.9
1999[4]	1.3	*	*	*	*	*	*	*	0.3	0.5	0.9	1.1	1.2	1.7
Female														
2001	1.9	*	*	*	*	*	*	0.3	0.4	0.4	0.8	0.9	1.0	1.7
2000[4]	1.8	*	*	*	*	*	*	0.4	0.4	0.5	0.7	1.0	1.3	1.5
1999[4]	1.6	*	*	*	*	*	*	*	0.3	0.4	0.6	1.0	0.9	1.5
Hispanic														
Both sexes														
2001	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	1.5	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	1.2	*	*
Male														
2001	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.7	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	1.8	*	*	*	*	*	0.2	0.3	0.4	0.5	0.9	1.1	1.3	2.0
2000[4]	1.8	0.7	*	*	*	*	0.2	0.4	0.4	0.6	0.9	1.3	1.5	1.8
1999[4]	1.6	0.7	*	*	*	*	*	0.2	0.3	0.5	0.8	1.0	1.0	1.6
Male														
2001	1.5	*	*	*	*	*	*	*	0.4	0.5	0.9	1.2	1.5	2.3
2000[4]	1.5	*	*	*	*	*	*	*	0.4	0.6	1.1	1.5	1.7	2.0
1999[4]	1.4	*	*	*	*	*	*	*	0.3	0.5	1.0	1.0	1.2	1.7
Female														
2001	2.0	*	*	*	*	*	*	0.3	0.5	0.5	0.8	0.9	1.0	1.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other disorders of circulatory system (I80-I99)						
All origins[3]						
Both sexes						
2001	2.6	3.8	5.4	7.7	11.9	26.1
2000[4]	2.6	3.4	5.3	7.7	11.9	25.6
1999[4]	2.2	3.3	4.3	6.8	12.7	23.4
Male						
2001	2.9	3.9	5.9	8.3	12.1	22.0
2000[4]	2.7	3.9	5.6	8.0	12.4	22.2
1999[4]	2.3	3.7	4.8	7.5	13.2	20.8
Female						
2001	2.3	3.7	5.0	7.3	11.7	27.8
2000[4]	2.6	3.0	5.0	7.5	11.6	26.9
1999[4]	2.1	2.9	3.9	6.3	12.4	24.5
Hispanic						
Both sexes						
2001	*	3.3	5.3	5.9	9.9	32.5
2000[4]	*	*	*	8.6	*	24.6
1999[4]	*	*	5.9	*	*	24.6
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	33.5
2000[4]	*	*	*	*	*	23.7
1999[4]	*	*	*	*	*	23.9
Non-Hispanic[5]						
Both sexes						
2001	2.7	3.8	5.4	7.8	11.9	25.8
2000[4]	2.7	3.4	5.4	7.7	12.0	25.6
1999[4]	2.2	3.3	4.2	6.9	12.7	23.3
Male						
2001	3.0	3.9	6.1	8.3	12.0	21.6
2000[4]	2.7	3.9	5.7	8.0	12.3	22.1
1999[4]	2.3	3.8	4.7	7.6	13.3	20.5
Female						
2001	2.4	3.8	4.9	7.4	11.8	27.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other disorders of circulatory system (I80-I99)														
Non-Hispanic[5]														
Female														
2000[4]	2.0	*	*	*	*	*	*	0.5	0.4	0.5	0.6	1.1	1.3	1.6
1999[4]	1.8	*	*	*	*	*	*	*	0.3	0.4	0.7	1.0	0.9	1.5
Non-Hispanic White														
Both sexes														
2001	1.8	*	*	*	*	*	*	0.2	0.3	0.4	0.7	0.9	1.1	1.8
2000[4]	1.8	*	*	*	*	*	*	0.3	0.3	0.5	0.8	1.2	1.4	1.6
1999[4]	1.6	*	*	*	*	*	*	0.2	0.3	0.4	0.7	0.8	0.9	1.6
Male														
2001	1.5	*	*	*	*	*	*	*	0.3	0.4	0.8	1.2	1.3	2.1
2000[4]	1.5	*	*	*	*	*	*	*	*	0.5	1.0	1.4	1.6	1.7
1999[4]	1.4	*	*	*	*	*	*	*	*	0.4	0.8	0.9	1.0	1.7
Female														
2001	2.1	*	*	*	*	*	*	*	0.4	0.4	0.7	0.7	1.0	1.6
2000[4]	2.0	*	*	*	*	*	*	0.4	0.3	0.4	0.5	1.0	1.1	1.4
1999[4]	1.8	*	*	*	*	*	*	*	0.3	0.3	0.6	0.8	0.8	1.5
Non-Hispanic Black														
Both sexes														
2001	2.0	*	*	*	*	*	*	*	1.1	1.2	1.9	2.2	2.4	4.0
2000[4]	2.1	*	*	*	*	*	*	0.9	1.0	1.3	1.8	2.5	3.2	4.0
1999[4]	1.8	*	*	*	*	*	*	*	*	1.0	1.9	2.4	2.4	2.1
Male														
2001	2.0	*	*	*	*	*	*	*	*	*	2.3	1.8	3.1	4.8
2000[4]	2.0	*	*	*	*	*	*	*	*	1.7	2.0	3.2	3.8	4.9
1999[4]	1.7	*	*	*	*	*	*	*	*	*	2.6	2.7	3.0	*
Female														
2001	2.1	*	*	*	*	*	*	*	*	*	1.5	2.5	*	3.3
2000[4]	2.3	*	*	*	*	*	*	*	*	*	1.7	2.0	2.7	3.3
1999[4]	1.8	*	*	*	*	*	*	*	*	*	*	2.2	*	*
Influenza and pneumonia (J10-J18)														
All origins[3]														
Both sexes														
2001	21.8	7.4	0.7	0.2	0.2	0.3	0.6	0.7	1.0	1.8	2.6	3.9	5.4	8.1
2000[4]	23.2	7.6	0.7	0.2	0.2	0.3	0.7	0.7	1.1	1.8	3.0	3.9	5.6	8.9
1999[4]	22.8	8.4	0.8	0.2	0.2	0.4	0.6	0.7	1.0	2.0	2.7	4.1	5.3	8.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other disorders of circulatory system (I80-I99)						
Non-Hispanic[5]						
Female						
2000[4]	2.7	3.0	5.1	7.5	11.8	27.0
1999[4]	2.1	2.9	3.8	6.5	12.4	24.4
Non-Hispanic White						
Both sexes						
2001	2.4	3.5	5.1	7.3	11.8	25.7
2000[4]	2.4	2.9	5.1	7.3	11.9	25.5
1999[4]	2.1	2.9	4.0	6.6	12.4	23.0
Male						
2001	2.5	3.6	5.7	7.8	11.9	21.2
2000[4]	2.4	3.4	5.6	7.8	12.1	21.8
1999[4]	2.3	3.3	4.5	7.1	12.8	20.3
Female						
2001	2.3	3.5	4.7	6.9	11.8	27.6
2000[4]	2.4	2.5	4.7	7.0	11.8	27.0
1999[4]	1.9	2.6	3.6	6.3	12.2	24.1
Non-Hispanic Black						
Both sexes						
2001	6.2	7.1	8.9	14.0	14.9	29.7
2000[4]	5.3	8.7	9.3	14.5	14.4	29.9
1999[4]	4.0	7.0	6.5	11.6	17.9	28.5
Male						
2001	8.8	8.1	10.7	16.1	*	28.9
2000[4]	5.7	9.3	8.9	13.0	*	30.5
1999[4]	*	9.6	6.9	15.0	21.3	*
Female						
2001	4.1	6.3	7.8	12.7	14.6	29.9
2000[4]	4.9	8.3	9.5	15.4	14.7	29.7
1999[4]	4.3	5.1	6.2	9.6	16.2	30.6
Influenza and pneumonia (J10-J18)						
All origins[3]						
Both sexes						
2001	13.9	24.4	49.3	101.1	216.9	685.6
2000[4]	15.6	26.6	52.6	106.7	240.8	744.1
1999[4]	14.3	25.0	50.5	105.9	233.8	751.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Influenza and pneumonia (J10-J18)														
All origins[3]														
Male														
2001	19.6	8.1	0.7	0.2	0.2	0.3	0.7	0.7	1.2	2.0	3.0	4.9	6.7	10.0
2000[4]	20.8	8.4	0.7	0.3	0.2	0.3	0.7	0.8	1.2	1.9	3.8	4.7	7.0	10.9
1999[4]	20.3	8.8	0.8	0.3	0.3	0.3	0.6	0.8	1.2	2.2	3.1	5.2	6.8	9.7
Female														
2001	23.9	6.7	0.7	0.2	0.2	0.3	0.5	0.6	0.9	1.6	2.2	3.0	4.1	6.4
2000[4]	25.6	6.8	0.7	0.2	*	0.3	0.6	0.6	1.0	1.7	2.2	3.1	4.3	7.0
1999[4]	25.3	8.0	0.9	*	0.2	0.4	0.5	0.6	0.8	1.8	2.3	3.0	3.9	7.2
Hispanic														
Both sexes														
2001	7.4	7.0	1.1	*	*	*	*	*	*	1.1	1.7	2.9	4.4	6.8
2000[4]	7.4	7.8	*	*	*	*	*	*	0.9	1.2	2.1	3.0	4.8	8.5
1999[4]	6.6	8.5	*	*	*	*	*	*	*	1.7	1.3	2.9	4.4	6.4
Male														
2001	6.9	7.4	*	*	*	*	*	*	*	1.3	2.2	4.9	5.3	10.0
2000[4]	7.2	7.6	*	*	*	*	*	*	*	1.4	3.0	4.6	6.2	11.0
1999[4]	6.3	10.3	*	*	*	*	*	*	*	2.4	1.8	3.8	6.2	7.4
Female														
2001	7.9	6.5	*	*	*	*	*	*	*	*	*	*	3.6	4.0
2000[4]	7.7	8.0	*	*	*	*	*	*	*	*	*	*	3.6	6.4
1999[4]	7.0	6.5	*	*	*	*	*	*	*	*	*	*	*	5.4
Non-Hispanic[5]														
Both sexes														
2001	23.8	7.5	0.6	0.2	0.2	0.3	0.6	0.7	1.2	1.9	2.7	4.0	5.4	8.2
2000[4]	25.4	7.5	0.7	0.2	0.2	0.4	0.7	0.8	1.1	1.8	3.0	3.9	5.6	8.8
1999[4]	25.0	8.4	0.9	0.2	0.2	0.4	0.6	0.8	1.1	2.0	2.8	4.1	5.3	8.5
Male														
2001	21.5	8.2	0.6	0.2	0.2	0.3	0.8	0.7	1.4	2.1	3.1	4.8	6.7	9.9
2000[4]	22.7	8.5	0.7	0.2	*	0.4	0.7	0.9	1.2	1.9	3.8	4.7	7.0	10.8
1999[4]	22.2	8.4	0.9	0.3	0.3	0.3	0.6	0.9	1.3	2.1	3.2	5.2	6.7	9.7
Female														
2001	26.1	6.7	0.6	0.3	0.2	0.3	0.5	0.7	0.9	1.7	2.3	3.2	4.1	6.5
2000[4]	27.9	6.5	0.6	*	*	0.3	0.6	0.6	1.0	1.8	2.2	3.3	4.4	7.0
1999[4]	27.6	8.4	0.9	*	*	0.4	0.5	0.7	0.9	2.0	2.5	3.1	4.0	7.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Influenza and pneumonia (J10-J18)						
All origins[3]						
Male						
2001	17.0	30.3	61.6	130.7	274.7	808.4
2000[4]	18.9	32.9	65.7	134.1	308.8	892.0
1999[4]	16.8	31.7	64.2	133.5	301.7	894.7
Female						
2001	11.1	19.2	39.5	80.5	182.3	634.2
2000[4]	12.5	21.2	42.2	87.6	200.7	683.9
1999[4]	12.0	19.2	39.7	86.7	194.1	694.2
Hispanic						
Both sexes						
2001	12.1	22.2	51.9	98.0	206.1	660.3
2000[4]	14.5	23.0	55.9	95.8	210.0	643.6
1999[4]	13.4	24.7	40.4	81.3	186.2	614.6
Male						
2001	13.3	25.2	60.8	122.8	270.1	765.6
2000[4]	19.9	28.0	69.5	122.5	295.9	763.9
1999[4]	14.3	31.7	53.5	100.6	243.6	720.3
Female						
2001	11.0	19.8	45.0	80.4	166.5	608.8
2000[4]	9.9	19.0	45.7	76.9	157.7	584.6
1999[4]	12.6	19.1	30.6	67.7	151.9	563.6
Non-Hispanic[5]						
Both sexes						
2001	14.0	24.4	48.8	101.0	216.7	684.8
2000[4]	15.6	26.6	52.2	106.7	241.0	745.7
1999[4]	14.2	24.8	50.8	106.6	234.7	754.0
Male						
2001	17.2	30.5	61.1	130.6	274.2	808.2
2000[4]	18.7	32.9	65.2	134.0	307.9	894.9
1999[4]	16.7	31.5	64.5	134.4	302.7	897.9
Female						
2001	11.1	19.1	39.0	80.2	182.3	633.5
2000[4]	12.7	21.2	41.9	87.7	201.6	685.4
1999[4]	11.9	19.1	40.1	87.2	194.9	696.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Influenza and pneumonia (J10-J18)														
Non-Hispanic White														
Both sexes														
2001	26.2	4.9	0.5	0.2	0.2	0.3	0.6	0.6	1.0	1.6	2.2	3.3	4.8	7.5
2000[4]	27.9	5.0	0.5	0.2	0.2	0.3	0.6	0.6	1.0	1.7	2.5	3.3	4.9	7.9
1999[4]	27.5	5.3	0.6	0.2	0.2	0.3	0.4	0.6	0.9	1.8	2.2	3.4	4.6	7.7
Male														
2001	23.1	5.8	0.6	*	*	*	0.7	0.6	1.2	1.7	2.5	4.0	5.7	8.9
2000[4]	24.5	5.8	0.5	*	*	0.4	0.7	0.7	1.0	1.7	3.2	3.7	5.7	9.2
1999[4]	23.9	5.5	0.5	*	*	0.3	0.5	0.7	1.1	1.8	2.3	4.2	5.5	8.6
Female														
2001	29.2	3.9	0.5	*	*	0.3	0.4	0.7	0.8	1.5	2.0	2.6	3.9	6.1
2000[4]	31.3	4.2	0.6	*	*	*	0.5	0.6	1.0	1.7	1.9	2.9	4.0	6.7
1999[4]	31.0	5.1	0.8	*	*	0.4	0.4	0.5	0.7	1.7	2.1	2.6	3.7	6.8
Non-Hispanic Black														
Both sexes														
2001	16.0	16.9	1.0	*	*	*	1.0	1.2	2.1	3.9	6.0	9.2	10.8	15.4
2000[4]	16.9	17.5	1.2	*	*	*	1.1	1.4	2.1	2.9	6.3	8.6	12.4	18.1
1999[4]	16.7	20.0	1.7	*	*	*	1.3	1.8	2.3	4.1	7.1	9.9	11.9	16.3
Male														
2001	16.3	16.6	*	*	*	*	*	*	2.8	4.7	7.1	11.6	15.9	20.8
2000[4]	17.3	18.8	1.7	*	*	*	*	2.1	2.6	3.2	8.6	11.7	17.6	26.2
1999[4]	16.9	20.1	2.1	*	*	*	*	2.0	2.6	4.3	9.2	13.6	17.4	21.8
Female														
2001	15.6	17.1	*	*	*	*	*	*	1.5	3.2	4.9	7.1	6.6	11.1
2000[4]	16.5	16.2	*	*	*	*	*	*	1.7	2.7	4.3	6.0	8.1	11.4
1999[4]	16.5	20.0	*	*	*	*	*	1.6	1.9	3.8	5.2	6.7	7.3	11.8
Influenza (J10-J11)														
All origins[3]														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.6	*	*	*	*	*	*	*	*	*	*	0.1	0.1	0.3
1999[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	*	0.2
Male														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	0.3
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Influenza and pneumonia (J10-J18)						
Non-Hispanic White						
Both sexes						
2001	12.9	23.2	47.4	98.2	216.2	693.9
2000[4]	14.2	24.4	49.7	105.5	240.7	759.8
1999[4]	12.9	23.8	48.9	106.1	236.5	766.9
Male						
2001	15.3	28.3	58.5	125.1	271.6	813.9
2000[4]	16.9	29.5	61.4	131.6	305.1	908.1
1999[4]	15.0	29.7	61.0	132.9	302.9	909.3
Female						
2001	10.6	18.7	38.3	79.2	182.8	644.0
2000[4]	11.7	20.0	40.2	87.2	202.5	700.0
1999[4]	11.0	18.6	39.3	87.3	197.2	710.2
Non-Hispanic Black						
Both sexes						
2001	23.7	34.9	66.2	133.9	224.9	583.1
2000[4]	27.6	46.2	78.5	124.8	247.4	604.5
1999[4]	26.6	37.0	76.4	119.8	220.4	634.7
Male						
2001	34.3	50.8	95.4	203.4	309.9	718.4
2000[4]	35.5	65.0	108.7	173.4	345.9	763.2
1999[4]	34.2	52.6	110.8	162.4	311.7	803.4
Female						
2001	15.4	23.1	46.8	92.5	181.6	533.0
2000[4]	21.3	32.2	58.5	95.5	197.9	545.6
1999[4]	20.7	25.4	53.5	94.1	175.2	571.6
Influenza (J10-J11)						
All origins[3]						
Both sexes						
2001	*	*	*	*	0.7	2.2
2000[4]	0.5	0.8	1.4	2.5	6.1	20.2
1999[4]	0.4	0.6	1.2	2.1	6.1	21.2
Male						
2001	*	*	*	*	*	2.2
2000[4]	0.5	1.0	1.6	2.7	6.3	20.7
1999[4]	0.5	0.6	1.2	2.4	7.3	19.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Influenza (J10-J11)														
All origins[3]														
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.8	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.7	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.7	*	*	*	*	*	*	*	*	*	*	0.1	0.1	0.3
1999[4]	0.7	*	*	*	*	*	*	*	*	*	*	*	*	0.2
Male														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.9	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.8	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.8	*	*	*	*	*	*	*	*	*	*	*	0.2	0.3
1999[4]	0.8	*	*	*	*	*	*	*	*	*	*	*	*	0.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Influenza (J10-J11)						
All origins[3]						
Female						
2001	*	*	*	*	0.7	2.2
2000[4]	0.5	0.6	1.2	2.5	6.0	19.9
1999[4]	*	0.6	1.1	1.9	5.4	21.7
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	*	0.7	2.3
2000[4]	0.5	0.8	1.5	2.6	6.2	20.5
1999[4]	0.4	0.6	1.2	2.1	6.2	21.6
Male						
2001	*	*	*	*	*	2.3
2000[4]	0.5	1.0	1.7	2.8	6.2	21.1
1999[4]	0.5	0.6	1.2	2.4	7.5	20.7
Female						
2001	*	*	*	*	0.8	2.2
2000[4]	0.6	0.7	1.3	2.5	6.2	20.3
1999[4]	*	0.7	1.2	1.9	5.4	22.0
Non-Hispanic White						
Both sexes						
2001	*	*	*	*	0.7	2.4
2000[4]	0.6	0.9	1.5	2.8	6.5	21.7
1999[4]	0.4	0.7	1.2	2.2	6.5	23.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Influenza (J10-J11)														
Non-Hispanic White														
Male														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Pneumonia (J12-J18)														
All origins[3]														
Both sexes														
2001	21.7	7.2	0.7	0.2	0.2	0.3	0.6	0.6	1.0	1.8	2.6	3.9	5.3	8.1
2000[4]	22.6	7.4	0.6	0.2	0.2	0.3	0.6	0.7	1.1	1.7	2.9	3.8	5.5	8.6
1999[4]	22.2	8.1	0.8	0.2	0.2	0.3	0.5	0.7	1.0	2.0	2.7	4.0	5.2	8.2
Male														
2001	19.5	7.8	0.7	0.2	0.2	0.3	0.7	0.6	1.2	2.0	3.0	4.8	6.6	9.9
2000[4]	20.3	8.1	0.6	0.2	*	0.3	0.7	0.8	1.2	1.8	3.7	4.6	6.8	10.6
1999[4]	19.8	8.5	0.7	0.2	0.2	0.3	0.6	0.8	1.1	2.2	3.1	5.1	6.7	9.5
Female														
2001	23.8	6.7	0.7	*	0.2	0.3	0.5	0.6	0.8	1.6	2.2	2.9	4.1	6.3
2000[4]	24.8	6.6	0.6	*	*	0.3	0.6	0.6	0.9	1.7	2.1	3.0	4.2	6.7
1999[4]	24.6	7.7	0.8	*	*	0.3	0.5	0.6	0.8	1.8	2.3	2.9	3.9	7.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Influenza (J10-J11)						
Non-Hispanic White						
Male						
2001	*	*	*	*	*	2.4
2000[4]	0.5	1.1	1.8	2.8	6.7	22.0
1999[4]	0.5	0.6	1.2	2.7	7.8	22.5
Female						
2001	*	*	*	*	0.7	2.4
2000[4]	0.6	0.7	1.2	2.8	6.4	21.6
1999[4]	*	0.8	1.2	1.9	5.7	23.5
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	9.5
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Pneumonia (J12-J18)						
All origins[3]						
Both sexes						
2001	13.8	24.3	49.2	100.9	216.1	683.4
2000[4]	15.1	25.8	51.2	104.1	234.7	723.9
1999[4]	13.9	24.3	49.3	103.8	227.7	730.6
Male						
2001	16.9	30.3	61.4	130.5	273.9	806.2
2000[4]	18.4	31.9	64.1	131.5	302.6	871.3
1999[4]	16.3	31.1	63.0	131.1	294.5	874.7
Female						
2001	11.0	19.1	39.4	80.2	181.5	632.0
2000[4]	12.0	20.5	41.0	85.1	194.6	663.9
1999[4]	11.7	18.6	38.6	84.8	188.7	672.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Pneumonia (J12-J18)														
Hispanic														
Both sexes														
2001	7.3	7.0	1.0	*	*	*	*	*	*	1.1	1.6	2.9	4.4	6.8
2000[4]	7.3	7.5	*	*	*	*	*	*	0.9	1.1	2.1	2.9	4.8	8.4
1999[4]	6.5	7.9	*	*	*	*	*	*	*	1.7	1.2	2.9	4.2	6.1
Male														
2001	6.8	7.4	*	*	*	*	*	*	*	1.3	2.2	4.8	5.3	10.0
2000[4]	7.1	7.6	*	*	*	*	*	*	*	1.4	3.0	4.5	6.2	10.7
1999[4]	6.1	10.3	*	*	*	*	*	*	*	2.3	*	3.8	5.9	6.9
Female														
2001	7.8	6.5	*	*	*	*	*	*	*	*	*	*	3.6	4.0
2000[4]	7.6	7.4	*	*	*	*	*	*	*	*	*	*	3.6	6.4
1999[4]	6.8	5.4	*	*	*	*	*	*	*	*	*	*	*	5.4
Non-Hispanic[5]														
Both sexes														
2001	23.8	7.2	0.6	0.2	0.2	0.3	0.6	0.7	1.1	1.9	2.7	4.0	5.4	8.1
2000[4]	24.7	7.3	0.6	0.2	0.1	0.3	0.7	0.7	1.1	1.8	2.9	3.8	5.5	8.5
1999[4]	24.3	8.1	0.8	0.2	0.2	0.3	0.6	0.8	1.0	2.0	2.8	4.1	5.3	8.2
Male														
2001	21.4	7.8	0.6	*	0.2	0.3	0.7	0.7	1.3	2.1	3.0	4.8	6.7	9.8
2000[4]	22.2	8.2	0.7	*	*	0.4	0.7	0.8	1.2	1.9	3.7	4.5	6.8	10.5
1999[4]	21.7	8.0	0.8	*	0.3	0.3	0.6	0.9	1.2	2.1	3.2	5.1	6.6	9.5
Female														
2001	26.0	6.6	0.6	*	*	0.3	0.5	0.7	0.9	1.7	2.3	3.1	4.1	6.5
2000[4]	27.1	6.4	0.5	*	*	0.3	0.6	0.6	1.0	1.8	2.2	3.1	4.3	6.8
1999[4]	26.8	8.2	0.9	*	*	0.3	0.5	0.6	0.9	1.9	2.4	3.0	4.0	7.0
Non-Hispanic White														
Both sexes														
2001	26.1	4.6	0.5	0.2	0.2	0.3	0.5	0.6	1.0	1.6	2.2	3.3	4.7	7.4
2000[4]	27.1	4.8	0.5	0.2	0.2	0.3	0.6	0.6	1.0	1.7	2.4	3.2	4.7	7.6
1999[4]	26.8	5.2	0.6	0.2	0.2	0.3	0.4	0.6	0.9	1.7	2.1	3.3	4.5	7.5
Male														
2001	23.0	5.3	0.6	*	*	*	0.7	0.5	1.1	1.7	2.5	4.0	5.6	8.8
2000[4]	23.8	5.4	0.4	*	*	0.3	0.7	0.6	1.0	1.7	3.1	3.7	5.6	8.9
1999[4]	23.4	5.4	*	*	*	0.3	0.5	0.7	1.0	1.8	2.3	4.1	5.5	8.4
Female														
2001	29.1	3.8	0.4	*	*	0.3	0.4	0.7	0.8	1.5	1.9	2.6	3.9	6.0
2000[4]	30.3	4.2	0.5	*	*	*	0.5	0.6	0.9	1.6	1.8	2.8	3.9	6.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonia (J12-J18)						
Hispanic						
Both sexes						
2001	12.1	22.2	51.7	97.1	205.1	660.3
2000[4]	14.4	22.5	55.7	94.9	206.1	633.0
1999[4]	13.0	24.0	39.6	79.7	183.8	607.6
Male						
2001	13.3	25.2	60.8	122.8	267.5	765.6
2000[4]	19.6	27.2	69.5	121.8	290.1	751.8
1999[4]	13.7	30.1	51.9	99.9	242.1	718.2
Female						
2001	11.0	19.8	44.7	78.9	166.5	608.8
2000[4]	9.9	18.7	45.3	75.8	155.0	574.7
1999[4]	12.3	19.1	30.3	65.5	149.1	554.2
Non-Hispanic[5]						
Both sexes						
2001	13.9	24.3	48.7	100.8	216.0	682.5
2000[4]	15.1	25.8	50.7	104.1	234.8	725.2
1999[4]	13.8	24.2	49.6	104.4	228.5	732.4
Male						
2001	17.1	30.4	60.9	130.4	273.6	805.9
2000[4]	18.2	31.9	63.5	131.3	301.6	873.8
1999[4]	16.3	30.9	63.3	132.0	295.2	877.2
Female						
2001	10.9	19.0	38.9	80.0	181.6	631.3
2000[4]	12.2	20.5	40.6	85.2	195.4	665.2
1999[4]	11.6	18.4	38.9	85.3	189.6	674.4
Non-Hispanic White						
Both sexes						
2001	12.8	23.1	47.3	98.0	215.5	691.5
2000[4]	13.7	23.6	48.2	102.7	234.2	738.1
1999[4]	12.5	23.1	47.7	103.9	230.0	743.7
Male						
2001	15.3	28.3	58.4	124.9	270.9	811.5
2000[4]	16.4	28.5	59.6	128.7	298.4	886.1
1999[4]	14.5	29.1	59.8	130.3	295.2	886.8
Female						
2001	10.5	18.6	38.2	78.9	182.1	641.7
2000[4]	11.1	19.3	39.0	84.5	196.0	678.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Pneumonia (J12-J18)														
Non-Hispanic White														
Female														
1999[4]	30.0	4.9	0.7	*	*	*	*	0.5	0.7	1.7	2.0	2.5	3.6	6.6
Non-Hispanic Black														
Both sexes														
2001	15.9	16.9	1.0	*	*	*	1.0	1.2	2.1	3.9	5.9	9.2	10.8	15.4
2000[4]	16.6	17.2	1.2	*	*	*	1.1	1.4	2.1	2.9	6.2	8.4	12.3	17.9
1999[4]	16.5	19.3	1.7	*	*	*	1.3	1.8	2.3	4.0	7.0	9.9	11.8	16.0
Male														
2001	16.3	16.6	*	*	*	*	*	*	2.7	4.7	7.1	11.6	15.9	20.6
2000[4]	17.1	18.8	1.7	*	*	*	*	2.1	2.6	3.2	8.3	11.4	17.4	25.9
1999[4]	16.8	19.1	2.0	*	*	*	*	2.0	2.6	4.3	9.1	13.5	17.3	21.8
Female														
2001	15.6	17.1	*	*	*	*	*	*	1.5	3.2	4.8	7.1	6.6	11.1
2000[4]	16.3	15.5	*	*	*	*	*	*	1.7	2.7	4.3	5.8	8.1	11.4
1999[4]	16.3	19.6	*	*	*	*	*	1.5	1.9	3.7	5.1	6.7	7.2	11.4
Other acute lower respiratory infections (J20-J22)														
All origins[3]														
Both sexes														
2001	0.1	1.2	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	1.1	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	1.7	0.2	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.1	1.3	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	1.3	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	1.9	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.2	1.2	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	1.5	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonia (J12-J18)						
Non-Hispanic White						
Female						
1999[4]	10.6	17.8	38.1	85.3	191.5	686.6
Non-Hispanic Black						
Both sexes						
2001	23.5	34.6	66.2	133.9	223.8	581.5
2000[4]	27.2	45.6	77.0	123.9	245.1	595.0
1999[4]	26.4	36.7	75.0	118.8	217.5	628.9
Male						
2001	33.9	50.3	95.4	203.4	309.9	716.1
2000[4]	35.3	64.5	107.6	171.5	345.0	750.3
1999[4]	34.0	52.1	109.4	161.9	307.3	799.9
Female						
2001	15.4	23.1	46.8	92.5	179.9	531.7
2000[4]	20.8	31.7	56.7	95.2	194.9	537.3
1999[4]	20.5	25.4	52.2	92.9	173.0	565.0
Other acute lower respiratory infections (J20-J22)						
All origins[3]						
Both sexes						
2001	*	*	*	0.3	0.7	3.3
2000[4]	*	*	0.3	0.6	0.8	4.5
1999[4]	*	*	0.2	0.6	1.3	6.0
Male						
2001	*	*	*	*	*	2.3
2000[4]	*	*	*	0.7	*	3.8
1999[4]	*	*	*	0.7	1.6	4.0
Female						
2001	*	*	*	*	0.7	3.7
2000[4]	*	*	*	0.5	0.9	4.8
1999[4]	*	*	*	0.5	1.1	6.8
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other acute lower respiratory infections (J20-J22)														
Hispanic														
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.1	1.3	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	1.1	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	1.7	0.2	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.1	1.2	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	2.0	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.2	1.4	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	1.5	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	0.1	1.0	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	1.1	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other acute lower respiratory infections (J20-J22)						
Hispanic						
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	0.3	0.6	3.3
2000[4]	*	*	0.3	0.6	0.8	4.5
1999[4]	*	*	0.2	0.6	1.3	6.1
Male						
2001	*	*	*	*	*	2.4
2000[4]	*	*	*	0.7	*	3.7
1999[4]	*	*	*	0.8	1.7	4.1
Female						
2001	*	*	*	*	0.7	3.7
2000[4]	*	*	*	0.5	0.9	4.8
1999[4]	*	*	*	0.5	1.1	6.9
Non-Hispanic White						
Both sexes						
2001	*	*	*	0.3	0.7	3.5
2000[4]	*	*	0.3	0.6	0.9	4.7
1999[4]	*	*	*	0.7	1.4	6.4
Male						
2001	*	*	*	*	*	2.5
2000[4]	*	*	*	0.8	*	3.7
1999[4]	*	*	*	0.8	1.7	4.1
Female						
2001	*	*	*	*	0.8	3.9
2000[4]	*	*	*	*	1.0	5.1
1999[4]	*	*	*	0.5	1.2	7.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other acute lower respiratory infections (J20-J22)														
Non-Hispanic Black														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	4.2	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Acute bronchitis and bronchiolitis (J20-J21)														
All origins[3]														
Both sexes														
2001	0.1	1.2	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	1.1	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	1.6	0.2	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.1	1.2	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	1.2	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	1.9	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	1.2	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	1.4	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other acute lower respiratory infections (J20-J22)						
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Acute bronchitis and bronchiolitis (J20-J21)						
All origins[3]						
Both sexes						
2001	*	*	*	*	0.4	2.1
2000[4]	*	*	*	0.4	0.5	2.6
1999[4]	*	*	*	0.4	0.8	3.6
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	2.4
1999[4]	*	*	*	*	*	2.6
Female						
2001	*	*	*	*	*	2.4
2000[4]	*	*	*	*	*	2.7
1999[4]	*	*	*	*	0.7	4.0
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute bronchitis and bronchiolitis (J20-J21)														
Hispanic														
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.1	1.3	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	1.1	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	1.7	0.2	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	2.0	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	1.4	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	1.4	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	0.1	1.0	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	1.1	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	4.2	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute bronchitis and bronchiolitis (J20-J21)						
Hispanic						
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	*	0.4	2.1
2000[4]	*	*	*	0.4	0.5	2.6
1999[4]	*	*	*	0.4	0.9	3.6
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	2.2
1999[4]	*	*	*	*	*	2.6
Female						
2001	*	*	*	*	*	2.4
2000[4]	*	*	*	*	*	2.7
1999[4]	*	*	*	*	0.7	4.0
Non-Hispanic White						
Both sexes						
2001	*	*	*	*	0.4	2.2
2000[4]	*	*	*	0.4	0.5	2.7
1999[4]	*	*	*	0.4	0.9	3.9
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	2.3
1999[4]	*	*	*	*	*	2.6
Female						
2001	*	*	*	*	*	2.5
2000[4]	*	*	*	*	*	2.8
1999[4]	*	*	*	*	0.7	4.4
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute bronchitis and bronchiolitis (J20-J21)														
Non-Hispanic Black														
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Unspecified acute lower respiratory infection (J22)														
All origins[3]														
Both sexes														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute bronchitis and bronchiolitis (J20-J21)						
Non-Hispanic Black						
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Unspecified acute lower respiratory infection (J22)						
All origins[3]						
Both sexes						
2001	*	*	*	*	*	1.2
2000[4]	*	*	*	*	*	1.9
1999[4]	*	*	*	*	0.5	2.4
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	1.3
2000[4]	*	*	*	*	*	2.1
1999[4]	*	*	*	*	*	2.8
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Unspecified acute lower respiratory infection (J22)														
Hispanic														
Female														
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Unspecified acute lower respiratory infection (J22)						
Hispanic						
Female						
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	*	*	1.2
2000[4]	*	*	*	*	*	1.9
1999[4]	*	*	*	*	0.5	2.4
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	1.3
2000[4]	*	*	*	*	*	2.1
1999[4]	*	*	*	*	*	2.8
Non-Hispanic White						
Both sexes						
2001	*	*	*	*	*	1.3
2000[4]	*	*	*	*	*	2.0
1999[4]	*	*	*	*	0.5	2.5
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	1.4
2000[4]	*	*	*	*	*	2.2
1999[4]	*	*	*	*	*	3.0
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Unspecified acute lower respiratory infection (J22)														
Non-Hispanic Black														
Male														
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Chronic lower respiratory diseases (J40-J47)														
All origins[3]														
Both sexes														
2001	43.2	1.0	0.3	0.2	0.3	0.4	0.5	0.5	0.9	1.6	2.7	5.4	11.9	29.4
2000[4]	43.4	0.9	0.3	0.2	0.4	0.4	0.5	0.6	0.8	1.4	2.8	5.8	11.8	29.6
1999[4]	44.5	0.9	0.4	0.2	0.4	0.5	0.5	0.7	0.9	1.3	2.6	5.5	12.0	31.9
Male														
2001	42.7	1.1	0.4	0.2	0.3	0.4	0.6	0.6	0.9	1.5	2.5	5.8	12.3	31.4
2000[4]	43.5	1.2	0.4	0.2	0.5	0.6	0.6	0.7	0.9	1.3	2.6	6.2	12.1	31.8
1999[4]	45.6	*	0.4	0.3	0.6	0.6	0.7	0.7	0.8	1.2	2.4	5.7	12.1	34.3
Female														
2001	43.7	*	*	*	0.2	0.3	0.4	0.5	0.9	1.7	2.9	5.1	11.5	27.5
2000[4]	43.2	*	0.3	0.2	0.4	0.3	0.5	0.6	0.8	1.4	2.9	5.5	11.5	27.5
1999[4]	43.4	*	0.3	0.2	0.3	0.5	0.4	0.8	1.0	1.4	2.9	5.3	11.9	29.7
Hispanic														
Both sexes														
2001	7.7	*	*	*	*	*	*	*	*	0.8	1.1	2.4	5.2	9.5
2000[4]	7.6	*	*	*	*	*	*	*	*	0.7	1.5	2.5	4.4	9.1
1999[4]	8.4	*	*	*	*	*	*	*	*	1.2	2.0	3.1	4.9	10.4
Male														
2001	7.8	*	*	*	*	*	*	*	*	*	*	2.3	4.4	10.6
2000[4]	8.0	*	*	*	*	*	*	*	*	*	*	2.4	4.8	8.8
1999[4]	8.9	*	*	*	*	*	*	*	*	*	2.0	2.5	5.0	11.6
Female														
2001	7.5	*	*	*	*	*	*	*	*	*	*	2.6	6.0	8.5
2000[4]	7.2	*	*	*	*	*	*	*	*	*	1.9	2.7	4.0	9.3
1999[4]	8.0	*	*	*	*	*	*	*	*	1.5	2.1	3.7	4.9	9.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Unspecified acute lower respiratory infection (J22)						
Non-Hispanic Black						
Male						
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Chronic lower respiratory diseases (J40-J47)						
All origins[3]						
Both sexes						
2001	62.9	124.2	215.4	329.2	452.9	644.7
2000[4]	62.5	124.1	218.2	327.8	473.4	648.6
1999[4]	66.6	130.4	227.6	340.2	484.5	646.0
Male						
2001	67.0	137.1	252.9	406.7	583.5	916.9
2000[4]	68.0	139.7	257.8	404.2	628.5	967.9
1999[4]	74.6	151.0	278.3	432.4	648.3	996.7
Female						
2001	59.1	113.1	185.6	275.0	374.6	530.8
2000[4]	57.5	110.7	187.0	274.6	381.9	518.5
1999[4]	59.3	112.8	188.0	276.1	388.7	504.6
Hispanic						
Both sexes						
2001	19.9	42.0	73.1	145.3	212.1	480.2
2000[4]	20.9	40.0	69.6	143.2	252.3	481.1
1999[4]	19.9	45.3	84.6	155.9	265.5	536.0
Male						
2001	23.6	57.1	99.3	195.1	279.2	676.1
2000[4]	30.5	50.7	100.6	191.9	366.6	671.1
1999[4]	25.3	52.5	120.7	216.9	385.1	836.8
Female						
2001	16.7	29.8	53.0	109.9	170.5	384.4
2000[4]	12.7	31.4	46.0	108.8	182.8	387.8
1999[4]	15.1	39.4	57.5	112.9	194.1	391.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Chronic lower respiratory diseases (J40-J47)														
Non-Hispanic[5]														
Both sexes														
2001	48.4	1.1	0.3	0.2	0.3	0.4	0.6	0.6	1.0	1.7	2.9	5.7	12.4	30.9
2000[4]	48.3	1.0	0.3	0.2	0.5	0.5	0.6	0.7	0.9	1.4	2.9	6.1	12.3	31.0
1999[4]	49.4	1.0	0.4	0.2	0.5	0.6	0.6	0.8	1.0	1.3	2.7	5.7	12.5	33.5
Male														
2001	48.0	*	0.4	0.3	0.4	0.4	0.7	0.6	1.0	1.7	2.7	6.1	12.9	32.9
2000[4]	48.7	*	0.4	0.3	0.5	0.6	0.7	0.8	1.0	1.4	2.8	6.5	12.6	33.4
1999[4]	50.8	*	0.4	0.3	0.6	0.6	0.8	0.8	0.9	1.3	2.5	5.9	12.6	35.9
Female														
2001	48.7	*	*	*	0.3	0.3	0.5	0.6	1.0	1.7	3.0	5.3	12.0	29.0
2000[4]	48.0	*	*	*	0.4	0.3	0.5	0.6	0.8	1.5	3.0	5.7	12.1	28.8
1999[4]	48.0	*	*	*	0.3	0.5	0.5	0.8	1.1	1.4	2.9	5.5	12.4	31.2
Non-Hispanic White														
Both sexes														
2001	55.9	*	*	*	0.2	0.3	0.3	0.4	0.7	1.2	2.6	5.4	12.3	31.8
2000[4]	55.7	*	*	*	0.2	0.3	0.4	0.5	0.6	1.1	2.4	5.5	12.0	31.7
1999[4]	56.6	*	0.2	0.2	0.2	0.5	0.4	0.5	0.6	1.0	2.3	5.0	12.3	34.3
Male														
2001	54.6	*	*	*	*	*	0.4	0.4	0.6	1.3	2.4	5.9	12.5	33.4
2000[4]	55.1	*	*	*	*	0.3	0.5	0.5	0.6	0.9	2.4	5.9	12.1	33.6
1999[4]	57.3	*	*	*	0.3	0.5	0.5	0.4	0.6	1.0	2.0	5.2	12.1	36.1
Female														
2001	57.2	*	*	*	*	*	*	0.3	0.8	1.2	2.8	4.8	12.1	30.3
2000[4]	56.2	*	*	*	*	*	0.4	0.5	0.7	1.3	2.5	5.1	11.9	29.9
1999[4]	55.9	*	*	*	*	0.4	0.4	0.5	0.7	1.1	2.5	4.8	12.6	32.5
Non-Hispanic Black														
Both sexes														
2001	21.1	*	0.9	0.7	0.9	1.0	1.7	1.9	3.1	4.7	5.1	9.3	16.8	32.3
2000[4]	21.5	*	0.9	0.7	1.4	1.5	1.4	1.9	2.5	3.6	6.3	11.3	18.5	33.8
1999[4]	22.6	*	*	*	1.5	1.2	1.6	2.6	2.8	3.3	6.2	11.5	17.5	36.9
Male														
2001	24.5	*	*	*	*	1.4	1.9	2.2	3.4	4.4	5.1	9.1	19.0	37.8
2000[4]	25.1	*	*	*	1.8	2.1	1.6	2.4	3.3	3.8	6.1	11.8	19.6	40.3
1999[4]	27.0	*	*	*	2.1	1.5	2.3	2.8	2.6	3.3	6.2	12.3	20.0	44.3
Female														
2001	18.0	*	*	*	*	*	1.5	1.6	2.7	4.9	5.2	9.5	14.9	27.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)						
Non-Hispanic[5]						
Both sexes						
2001	65.9	129.7	223.3	337.5	461.7	649.4
2000[4]	65.3	129.3	226.0	335.4	480.7	653.2
1999[4]	69.7	135.5	234.6	347.3	491.4	648.0
Male						
2001	69.9	142.2	261.1	416.4	594.6	924.8
2000[4]	70.3	144.8	265.6	412.8	637.1	978.0
1999[4]	77.7	156.5	285.5	440.8	656.5	1,000.2
Female						
2001	62.3	118.9	193.3	282.4	382.2	534.8
2000[4]	60.8	115.9	194.7	281.5	388.6	521.9
1999[4]	62.4	117.4	194.8	282.3	394.8	507.0
Non-Hispanic White						
Both sexes						
2001	69.9	137.9	236.4	355.6	483.3	676.9
2000[4]	68.8	136.5	238.6	351.8	503.7	680.8
1999[4]	73.2	143.2	246.1	361.3	512.6	675.6
Male						
2001	72.5	147.9	270.1	429.6	611.8	951.5
2000[4]	72.5	148.9	274.6	425.1	658.0	1,008.6
1999[4]	79.7	162.0	293.9	450.4	673.4	1,033.5
Female						
2001	67.4	129.1	209.0	303.2	405.8	562.7
2000[4]	65.3	125.7	209.7	300.2	412.0	548.7
1999[4]	67.2	126.8	208.0	298.8	417.7	532.8
Non-Hispanic Black						
Both sexes						
2001	52.7	95.9	150.7	208.7	279.2	379.3
2000[4]	58.1	98.7	152.8	213.1	273.6	376.2
1999[4]	58.6	101.6	164.5	245.5	297.3	377.0
Male						
2001	66.7	129.2	224.1	333.2	475.9	682.6
2000[4]	74.2	140.2	228.6	328.0	451.4	696.3
1999[4]	80.5	141.6	244.1	391.5	524.3	707.2
Female						
2001	41.7	71.4	101.9	134.4	179.1	266.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Chronic lower respiratory diseases (J40-J47)														
Non-Hispanic Black														
Female														
2000[4]	18.2	*	*	*	*	*	*	1.5	1.7	3.4	6.4	10.8	17.4	28.4
1999[4]	18.6	*	*	*	*	*	*	2.4	2.9	3.3	6.2	10.9	15.3	31.0
Bronchitis, chronic and unspecified (J40-J42)														
All origins[3]														
Both sexes														
2001	0.3	0.5	*	*	*	*	*	*	*	*	*	0.1	0.1	0.2
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	0.1
1999[4]	0.4	0.5	*	*	*	*	*	*	*	*	*	*	0.1	0.3
Male														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	0.4
Female														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	0.1	0.2
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)						
Non-Hispanic Black						
Female						
2000[4]	45.5	68.0	102.5	143.7	184.1	257.3
1999[4]	41.5	72.1	111.7	157.5	184.8	253.6
Bronchitis, chronic and unspecified (J40-J42)						
All origins[3]						
Both sexes						
2001	0.3	0.6	0.9	1.7	2.6	9.1
2000[4]	0.4	0.7	1.0	2.1	3.6	11.8
1999[4]	0.6	0.7	1.2	1.9	4.1	10.7
Male						
2001	0.4	0.8	1.1	2.2	2.5	9.2
2000[4]	0.4	0.9	1.0	2.8	4.9	12.9
1999[4]	0.5	0.8	1.5	2.4	4.7	11.9
Female						
2001	*	0.4	0.7	1.4	2.6	9.1
2000[4]	0.4	0.5	1.1	1.6	2.8	11.4
1999[4]	0.6	0.5	1.0	1.5	3.8	10.3
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	0.3	0.6	0.9	1.7	2.6	9.1
2000[4]	0.5	0.7	1.0	2.1	3.5	11.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Bronchitis, chronic and unspecified (J40-J42)														
Non-Hispanic[5]														
Both sexes														
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	0.2	0.3
Male														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	0.4
Female														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	0.1	0.2
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	0.3
Male														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	0.4
Female														
2001	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Bronchitis, chronic and unspecified (J40-J42)						
Non-Hispanic[5]						
Both sexes						
1999[4]	0.6	0.7	1.3	1.9	4.1	10.7
Male						
2001	0.4	0.8	1.1	2.1	2.6	9.3
2000[4]	0.5	1.0	0.9	2.9	4.8	12.8
1999[4]	0.6	0.8	1.5	2.5	4.7	11.9
Female						
2001	*	0.4	0.8	1.4	2.6	9.1
2000[4]	0.5	0.5	1.1	1.7	2.8	11.4
1999[4]	0.6	0.6	1.1	1.5	3.7	10.2
Non-Hispanic White						
Both sexes						
2001	0.4	0.6	0.9	1.7	2.7	9.7
2000[4]	0.5	0.7	1.0	2.2	3.6	12.6
1999[4]	0.6	0.7	1.3	1.9	4.1	11.2
Male						
2001	*	0.9	1.1	2.1	2.6	9.6
2000[4]	0.5	1.0	0.9	2.9	5.1	13.8
1999[4]	0.6	0.9	1.5	2.5	4.5	12.4
Female						
2001	*	*	0.7	1.5	2.8	9.7
2000[4]	0.5	0.5	1.2	1.7	2.8	12.2
1999[4]	0.7	0.6	1.1	1.5	3.8	10.7
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Emphysema (J43)														
All origins[3]														
Both sexes														
2001	5.7	*	*	*	*	*	*	*	*	0.1	0.2	0.7	1.7	5.0
2000[4]	5.9	*	*	*	*	*	*	*	*	0.1	0.3	0.8	1.8	5.0
1999[4]	6.4	*	*	*	*	*	*	*	*	*	0.4	0.7	2.0	5.5
Male														
2001	6.0	*	*	*	*	*	*	*	*	*	0.3	0.9	2.0	5.8
2000[4]	6.3	*	*	*	*	*	*	*	*	*	0.3	1.0	2.1	5.6
1999[4]	6.9	*	*	*	*	*	*	*	*	*	0.4	0.9	2.2	6.2
Female														
2001	5.4	*	*	*	*	*	*	*	*	*	*	0.5	1.4	4.2
2000[4]	5.6	*	*	*	*	*	*	*	*	*	0.2	0.6	1.5	4.5
1999[4]	5.8	*	*	*	*	*	*	*	*	*	0.3	0.5	1.7	4.9
Hispanic														
Both sexes														
2001	0.8	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.8	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.9	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.9	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.9	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.7	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	6.4	*	*	*	*	*	*	*	*	0.1	0.2	0.7	1.8	5.3
2000[4]	6.7	*	*	*	*	*	*	*	*	0.1	0.3	0.8	1.9	5.4
1999[4]	7.1	*	*	*	*	*	*	*	*	*	0.4	0.7	2.0	5.9
Male														
2001	6.8	*	*	*	*	*	*	*	*	*	0.3	0.9	2.1	6.0
2000[4]	7.1	*	*	*	*	*	*	*	*	*	0.3	1.1	2.2	6.0
1999[4]	7.8	*	*	*	*	*	*	*	*	*	0.4	0.9	2.3	6.6
Female														
2001	6.1	*	*	*	*	*	*	*	*	*	*	0.5	1.5	4.5
2000[4]	6.3	*	*	*	*	*	*	*	*	*	0.3	0.6	1.6	4.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Emphysema (J43)						
All origins[3]						
Both sexes						
2001	10.1	19.8	33.2	46.9	55.5	62.1
2000[4]	10.4	20.7	35.5	47.5	60.4	63.3
1999[4]	12.1	23.5	38.4	51.3	65.3	62.1
Male						
2001	10.8	22.4	40.3	59.2	74.9	101.2
2000[4]	11.2	24.5	42.6	60.4	83.6	104.2
1999[4]	13.6	28.0	48.9	67.3	91.5	105.4
Female						
2001	9.5	17.6	27.5	38.3	43.8	45.8
2000[4]	9.7	17.5	29.9	38.5	46.8	46.7
1999[4]	10.8	19.8	30.2	40.2	49.9	44.7
Hispanic						
Both sexes						
2001	*	5.2	8.3	17.3	25.8	39.4
2000[4]	*	4.3	8.2	18.4	29.5	31.2
1999[4]	*	4.8	10.1	18.1	33.5	44.2
Male						
2001	*	*	12.4	28.0	45.4	60.9
2000[4]	*	*	12.6	25.8	45.6	58.4
1999[4]	*	*	16.8	26.5	55.0	77.6
Female						
2001	*	*	*	9.7	*	28.9
2000[4]	*	*	*	13.1	19.7	*
1999[4]	*	*	*	12.1	20.6	28.1
Non-Hispanic[5]						
Both sexes						
2001	10.7	20.8	34.6	48.2	56.6	62.8
2000[4]	11.0	21.8	36.9	48.7	61.5	64.3
1999[4]	12.8	24.7	39.9	52.6	66.3	62.5
Male						
2001	11.4	23.4	41.8	60.6	76.0	102.7
2000[4]	11.7	25.6	44.0	61.9	85.1	105.9
1999[4]	14.2	29.2	50.5	68.9	92.7	106.1
Female						
2001	10.1	18.6	28.9	39.6	44.9	46.3
2000[4]	10.3	18.5	31.3	39.5	47.7	47.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Emphysema (J43)														
Non-Hispanic[5]														
Female														
1999[4]	6.5	*	*	*	*	*	*	*	*	*	0.3	0.5	1.8	5.2
Non-Hispanic White														
Both sexes														
2001	7.5	*	*	*	*	*	*	*	*	0.1	0.3	0.8	1.8	5.5
2000[4]	7.8	*	*	*	*	*	*	*	*	*	0.3	0.8	1.9	5.6
1999[4]	8.3	*	*	*	*	*	*	*	*	*	0.3	0.7	2.1	6.3
Male														
2001	7.8	*	*	*	*	*	*	*	*	*	0.3	1.0	2.0	6.1
2000[4]	8.1	*	*	*	*	*	*	*	*	*	0.3	0.9	2.1	6.2
1999[4]	8.9	*	*	*	*	*	*	*	*	*	0.3	0.8	2.1	6.8
Female														
2001	7.2	*	*	*	*	*	*	*	*	*	*	0.5	1.5	4.9
2000[4]	7.4	*	*	*	*	*	*	*	*	*	0.3	0.6	1.6	5.1
1999[4]	7.7	*	*	*	*	*	*	*	*	*	0.3	0.5	2.0	5.7
Non-Hispanic Black														
Both sexes														
2001	2.2	*	*	*	*	*	*	*	*	*	*	*	2.4	5.0
2000[4]	2.5	*	*	*	*	*	*	*	*	*	*	1.6	2.6	4.8
1999[4]	2.7	*	*	*	*	*	*	*	*	*	0.7	1.4	2.5	4.7
Male														
2001	2.8	*	*	*	*	*	*	*	*	*	*	*	3.0	6.9
2000[4]	3.3	*	*	*	*	*	*	*	*	*	*	2.5	3.6	6.1
1999[4]	3.6	*	*	*	*	*	*	*	*	*	*	2.0	4.2	6.6
Female														
2001	1.7	*	*	*	*	*	*	*	*	*	*	*	*	3.4
2000[4]	1.8	*	*	*	*	*	*	*	*	*	*	*	*	3.7
1999[4]	1.8	*	*	*	*	*	*	*	*	*	*	*	*	3.1
Asthma (J45-J46)														
All origins[3]														
Both sexes														
2001	1.5	*	0.2	0.2	0.3	0.3	0.4	0.5	0.7	0.9	1.1	1.3	1.8	1.9
2000[4]	1.6	*	0.2	0.2	0.4	0.4	0.5	0.5	0.7	0.8	1.2	1.6	1.7	2.0
1999[4]	1.7	*	0.2	0.2	0.4	0.5	0.5	0.6	0.7	0.9	1.1	1.6	1.7	2.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Emphysema (J43)						
Non-Hispanic[5]						
Female						
1999[4]	11.5	20.8	31.6	41.3	50.9	45.0
Non-Hispanic White						
Both sexes						
2001	11.6	22.7	36.9	51.4	59.9	65.9
2000[4]	11.9	23.2	39.5	51.6	64.9	66.9
1999[4]	13.8	26.8	42.6	55.3	69.9	65.2
Male						
2001	12.0	25.1	43.3	63.4	79.2	106.6
2000[4]	12.3	26.3	46.3	64.8	88.6	107.3
1999[4]	14.9	30.8	52.4	71.4	95.7	110.0
Female						
2001	11.3	20.6	31.7	42.9	48.3	48.9
2000[4]	11.5	20.5	34.1	42.3	50.8	50.6
1999[4]	12.9	23.2	34.7	43.9	54.8	47.3
Non-Hispanic Black						
Both sexes						
2001	6.3	11.7	20.4	21.8	26.5	35.9
2000[4]	7.8	15.3	19.2	24.5	31.0	37.5
1999[4]	8.2	14.2	21.1	31.8	31.7	37.8
Male						
2001	9.2	15.9	32.4	34.6	49.9	69.3
2000[4]	10.8	25.9	29.4	34.7	53.2	89.2
1999[4]	12.2	23.1	37.4	49.8	66.4	73.7
Female						
2001	4.1	8.7	12.4	14.1	14.6	23.5
2000[4]	5.4	7.5	12.5	18.3	19.9	18.3
1999[4]	5.1	7.7	10.3	21.0	14.5	24.4
Asthma (J45-J46)						
All origins[3]						
Both sexes						
2001	2.6	3.0	3.9	5.2	8.1	16.1
2000[4]	2.5	3.4	4.6	5.7	8.7	16.7
1999[4]	2.2	3.7	4.7	6.2	9.6	17.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Asthma (J45-J46)														
All origins[3]														
Male														
2001	1.1	*	0.3	0.2	0.3	0.3	0.4	0.5	0.6	0.8	0.9	1.0	1.3	1.2
2000[4]	1.2	*	0.3	0.2	0.5	0.5	0.5	0.7	0.8	0.7	1.0	1.3	1.3	1.6
1999[4]	1.2	*	*	0.2	0.6	0.5	0.6	0.6	0.6	0.8	0.8	1.0	1.2	1.9
Female														
2001	1.9	*	*	*	0.2	0.3	0.4	0.4	0.8	1.0	1.3	1.7	2.2	2.5
2000[4]	2.0	*	*	0.2	0.3	0.3	0.4	0.4	0.6	0.9	1.4	1.9	2.1	2.4
1999[4]	2.1	*	*	*	0.3	0.4	0.4	0.6	0.8	1.0	1.3	2.2	2.2	3.3
Hispanic														
Both sexes														
2001	0.7	*	*	*	*	*	*	*	*	*	*	1.0	2.0	2.5
2000[4]	0.8	*	*	*	*	*	*	*	*	*	*	*	1.6	*
1999[4]	0.9	*	*	*	*	*	*	*	*	0.9	1.0	1.5	1.8	2.5
Male														
2001	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.7	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.7	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	1.0	*	*	*	*	*	*	*	*	*	*	*	2.8	*
2000[4]	1.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	1.6	*	0.2	0.2	0.3	0.3	0.5	0.5	0.8	1.0	1.1	1.4	1.7	1.8
2000[4]	1.7	*	0.2	0.2	0.4	0.4	0.5	0.6	0.8	0.9	1.3	1.6	1.7	2.0
1999[4]	1.8	*	0.2	0.2	0.4	0.5	0.5	0.7	0.7	0.9	1.1	1.6	1.7	2.6
Male														
2001	1.1	*	*	0.3	0.3	0.4	0.5	0.6	0.7	0.9	1.0	1.0	1.3	1.1
2000[4]	1.2	*	*	0.2	0.5	0.5	0.6	0.7	0.8	0.8	1.1	1.3	1.2	1.6
1999[4]	1.3	*	*	0.3	0.6	0.6	0.7	0.7	0.7	0.9	0.8	1.0	1.2	1.9
Female														
2001	2.1	*	*	*	0.3	0.3	0.4	0.5	0.8	1.1	1.3	1.7	2.1	2.5
2000[4]	2.1	*	*	*	0.4	0.3	0.5	0.5	0.7	1.0	1.5	1.9	2.1	2.4
1999[4]	2.2	*	*	*	0.3	0.5	0.4	0.6	0.8	1.0	1.4	2.2	2.3	3.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Asthma (J45-J46)						
All origins[3]						
Male						
2001	1.9	2.4	3.0	3.8	5.6	12.1
2000[4]	1.8	2.7	3.4	3.9	6.7	12.9
1999[4]	1.7	2.6	3.9	4.8	7.2	12.7
Female						
2001	3.3	3.6	4.7	6.1	9.6	17.8
2000[4]	3.1	4.0	5.5	6.9	9.8	18.2
1999[4]	2.7	4.6	5.4	7.2	11.1	19.2
Hispanic						
Both sexes						
2001	*	*	4.8	7.4	*	13.8
2000[4]	4.4	5.7	5.4	7.7	*	16.6
1999[4]	*	5.5	6.1	8.7	13.5	17.5
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	18.6
2000[4]	*	6.0	*	*	*	*
1999[4]	*	6.9	*	11.0	*	*
Non-Hispanic[5]						
Both sexes						
2001	2.6	3.0	3.9	5.0	8.1	16.1
2000[4]	2.3	3.3	4.5	5.5	8.7	16.6
1999[4]	2.2	3.6	4.6	6.1	9.5	17.3
Male						
2001	1.9	2.3	2.9	3.7	5.6	12.2
2000[4]	1.6	2.6	3.2	3.8	6.7	12.7
1999[4]	1.8	2.5	3.8	4.8	7.1	12.6
Female						
2001	3.2	3.6	4.7	6.0	9.6	17.7
2000[4]	3.0	3.9	5.5	6.7	9.8	18.2
1999[4]	2.7	4.5	5.3	7.0	10.9	19.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Asthma (J45-J46)														
Non-Hispanic White														
Both sexes														
2001	1.4	*	*	*	0.2	0.2	0.3	0.3	0.5	0.6	0.8	0.9	1.3	1.3
2000[4]	1.4	*	*	*	0.2	0.2	0.4	0.4	0.5	0.5	0.9	1.0	1.2	1.4
1999[4]	1.5	*	*	*	0.2	0.4	0.4	0.4	0.5	0.6	0.8	1.1	1.2	2.0
Male														
2001	0.9	*	*	*	*	*	*	0.3	0.3	0.5	0.6	0.7	0.9	0.8
2000[4]	1.0	*	*	*	*	*	0.4	0.4	0.5	0.4	0.7	0.9	0.9	1.0
1999[4]	1.0	*	*	*	0.3	0.4	0.4	0.3	0.4	0.5	0.4	0.6	0.8	1.4
Female														
2001	1.9	*	*	*	*	*	*	*	0.6	0.6	1.0	1.2	1.7	1.8
2000[4]	1.9	*	*	*	*	*	0.3	*	0.5	0.7	1.0	1.2	1.4	1.7
1999[4]	2.1	*	*	*	*	0.4	*	0.4	0.5	0.7	1.1	1.6	1.6	2.5
Non-Hispanic Black														
Both sexes														
2001	3.1	*	*	0.7	0.9	0.9	1.4	1.8	2.7	3.5	3.1	4.4	5.0	6.1
2000[4]	3.3	*	*	0.7	1.4	1.3	1.2	1.8	2.1	2.7	3.7	5.6	5.3	6.7
1999[4]	3.3	*	*	*	1.5	1.1	1.5	2.4	2.4	2.7	3.3	5.0	5.9	7.5
Male														
2001	2.7	*	*	*	*	*	1.6	2.0	3.0	3.1	2.9	3.8	4.6	3.7
2000[4]	2.8	*	*	*	1.7	1.8	*	2.3	2.7	2.9	3.2	4.1	3.2	6.3
1999[4]	2.9	*	*	*	2.0	1.3	2.1	2.8	2.6	2.7	2.9	3.7	4.3	5.2
Female														
2001	3.4	*	*	*	*	*	*	1.5	2.4	3.9	3.3	5.0	5.3	8.0
2000[4]	3.7	*	*	*	*	*	*	*	1.5	2.7	4.1	6.9	7.1	7.1
1999[4]	3.6	*	*	*	*	*	*	2.1	2.3	2.7	3.7	6.0	7.2	9.4
Other chronic lower respiratory diseases (J44,J47)														
All origins[3]														
Both sexes														
2001	35.7	*	*	*	*	*	*	*	0.2	0.5	1.3	3.3	8.4	22.4
2000[4]	35.4	*	*	*	*	*	*	*	*	0.4	1.2	3.4	8.2	22.4
1999[4]	36.0	*	*	*	*	*	*	0.1	0.1	0.3	1.2	3.1	8.2	23.5
Male														
2001	35.3	*	*	*	*	*	*	*	0.2	0.5	1.3	3.8	9.0	24.2
2000[4]	35.6	*	*	*	*	*	*	*	*	0.4	1.2	3.8	8.7	24.4
1999[4]	37.1	*	*	*	*	*	*	*	*	0.3	1.2	3.7	8.6	25.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Asthma (J45-J46)						
Non-Hispanic White						
Both sexes						
2001	1.8	2.6	3.3	4.5	7.7	15.7
2000[4]	1.8	2.5	3.9	5.0	8.4	15.8
1999[4]	1.7	2.9	3.9	5.6	9.2	16.5
Male						
2001	1.2	1.9	2.3	3.2	5.0	11.4
2000[4]	1.1	1.8	2.9	3.5	6.3	11.1
1999[4]	1.3	1.9	2.9	4.2	6.9	10.6
Female						
2001	2.4	3.2	4.1	5.5	9.3	17.5
2000[4]	2.3	3.2	4.8	6.1	9.7	17.7
1999[4]	2.1	3.8	4.6	6.7	10.5	18.8
Non-Hispanic Black						
Both sexes						
2001	7.9	6.3	9.5	10.3	11.3	16.5
2000[4]	7.2	9.1	9.8	10.9	9.8	22.3
1999[4]	6.3	8.6	11.2	10.0	11.4	20.7
Male						
2001	6.1	*	9.6	*	*	*
2000[4]	6.0	8.8	6.8	*	*	24.7
1999[4]	5.9	6.9	12.1	9.7	*	*
Female						
2001	9.4	7.5	9.4	11.6	12.1	17.1
2000[4]	8.2	9.2	11.8	14.5	9.9	21.4
1999[4]	6.7	9.8	10.6	10.2	14.0	21.3
Other chronic lower respiratory diseases (J44,J47)						
All origins[3]						
Both sexes						
2001	49.8	100.7	177.4	275.4	386.8	557.4
2000[4]	49.1	99.2	177.1	272.5	400.7	556.7
1999[4]	51.6	102.5	183.2	280.8	405.5	555.8
Male						
2001	54.0	111.6	208.6	341.5	500.4	794.5
2000[4]	54.5	111.6	210.9	337.1	533.3	838.0
1999[4]	58.7	119.6	223.9	357.9	544.9	866.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other chronic lower respiratory diseases (J44,J47)														
All origins[3]														
Female														
2001	36.0	*	*	*	*	*	*	*	*	0.5	1.4	2.8	7.8	20.7
2000[4]	35.2	*	*	*	*	*	*	*	*	0.5	1.2	2.9	7.8	20.5
1999[4]	35.0	*	*	*	*	*	*	*	*	0.3	1.1	2.6	7.8	21.4
Hispanic														
Both sexes														
2001	6.0	*	*	*	*	*	*	*	*	*	*	1.1	3.0	5.9
2000[4]	5.9	*	*	*	*	*	*	*	*	*	*	1.2	2.2	6.6
1999[4]	6.5	*	*	*	*	*	*	*	*	*	*	1.3	2.6	7.2
Male														
2001	6.3	*	*	*	*	*	*	*	*	*	*	*	2.9	6.6
2000[4]	6.2	*	*	*	*	*	*	*	*	*	*	*	*	7.0
1999[4]	7.0	*	*	*	*	*	*	*	*	*	*	*	*	8.8
Female														
2001	5.8	*	*	*	*	*	*	*	*	*	*	*	3.2	5.2
2000[4]	5.6	*	*	*	*	*	*	*	*	*	*	*	*	6.2
1999[4]	5.9	*	*	*	*	*	*	*	*	*	*	*	*	5.8
Non-Hispanic[5]														
Both sexes														
2001	40.0	*	*	*	*	*	*	*	0.2	0.6	1.4	3.5	8.8	23.6
2000[4]	39.5	*	*	*	*	*	*	*	*	0.5	1.3	3.5	8.7	23.5
1999[4]	40.0	*	*	*	*	*	*	*	0.1	0.3	1.2	3.3	8.6	24.7
Male														
2001	39.8	*	*	*	*	*	*	*	*	0.6	1.4	4.0	9.4	25.5
2000[4]	40.0	*	*	*	*	*	*	*	*	0.4	1.3	4.1	9.2	25.7
1999[4]	41.4	*	*	*	*	*	*	*	*	0.3	1.3	3.9	9.0	27.0
Female														
2001	40.1	*	*	*	*	*	*	*	*	0.5	1.5	3.0	8.2	21.9
2000[4]	39.1	*	*	*	*	*	*	*	*	0.5	1.2	3.0	8.2	21.5
1999[4]	38.7	*	*	*	*	*	*	*	*	0.3	1.1	2.7	8.2	22.5
Non-Hispanic White														
Both sexes														
2001	46.6	*	*	*	*	*	*	*	0.2	0.5	1.5	3.6	9.1	24.8
2000[4]	46.0	*	*	*	*	*	*	*	*	0.4	1.3	3.6	8.9	24.5
1999[4]	46.3	*	*	*	*	*	*	*	*	0.3	1.2	3.2	8.9	25.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)						
All origins[3]						
Female						
2001	46.1	91.4	152.7	229.1	318.6	458.2
2000[4]	44.3	88.7	150.4	227.6	322.5	442.2
1999[4]	45.2	87.9	151.4	227.2	324.0	430.5
Hispanic						
Both sexes						
2001	15.5	33.6	59.6	118.4	176.8	418.9
2000[4]	14.1	29.7	54.5	116.0	209.4	422.7
1999[4]	15.3	34.6	68.0	127.8	213.2	462.3
Male						
2001	19.8	47.1	82.3	157.5	227.2	605.7
2000[4]	22.2	38.8	79.7	159.5	307.7	584.5
1999[4]	20.6	41.0	98.8	184.1	319.1	735.4
Female						
2001	11.8	22.7	42.2	90.5	145.6	327.7
2000[4]	7.2	22.3	35.3	85.2	149.6	343.3
1999[4]	10.8	29.4	44.8	88.1	150.0	330.7
Non-Hispanic[5]						
Both sexes						
2001	52.3	105.2	184.0	282.5	394.4	561.3
2000[4]	51.5	103.5	183.6	279.1	407.0	560.5
1999[4]	54.0	106.6	188.8	286.7	411.5	557.6
Male						
2001	56.2	115.6	215.3	350.0	510.4	800.6
2000[4]	56.4	115.7	217.5	344.3	540.5	846.6
1999[4]	61.1	124.1	229.6	364.7	552.0	869.6
Female						
2001	48.7	96.2	159.0	235.3	325.1	461.7
2000[4]	47.0	93.0	156.8	233.7	328.4	444.8
1999[4]	47.5	91.6	156.9	232.5	329.4	432.7
Non-Hispanic White						
Both sexes						
2001	56.0	112.0	195.4	298.0	413.0	585.7
2000[4]	54.6	110.1	194.1	292.9	426.8	585.5
1999[4]	57.0	112.9	198.4	298.5	429.4	582.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other chronic lower respiratory diseases (J44,J47)														
Non-Hispanic White														
Male														
2001	45.6	*	*	*	*	*	*	*	*	0.5	1.4	4.1	9.5	26.2
2000[4]	45.7	*	*	*	*	*	*	*	*	0.4	1.3	4.0	9.0	26.3
1999[4]	47.1	*	*	*	*	*	*	*	*	0.3	1.2	3.8	9.1	27.5
Female														
2001	47.6	*	*	*	*	*	*	*	*	0.5	1.5	3.0	8.7	23.5
2000[4]	46.3	*	*	*	*	*	*	*	*	0.5	1.2	3.2	8.7	22.9
1999[4]	45.5	*	*	*	*	*	*	*	*	0.3	1.1	2.6	8.8	24.1
Non-Hispanic Black														
Both sexes														
2001	15.6	*	*	*	*	*	*	*	*	1.1	1.6	4.2	9.4	21.2
2000[4]	15.4	*	*	*	*	*	*	*	*	0.7	1.9	4.0	10.3	22.2
1999[4]	16.4	*	*	*	*	*	*	*	*	*	2.0	5.0	8.6	24.5
Male														
2001	18.8	*	*	*	*	*	*	*	*	*	1.6	4.5	11.3	27.2
2000[4]	18.7	*	*	*	*	*	*	*	*	*	1.8	5.2	12.7	27.9
1999[4]	20.3	*	*	*	*	*	*	*	*	*	2.6	6.2	11.0	32.1
Female														
2001	12.7	*	*	*	*	*	*	*	*	*	1.6	3.9	7.8	16.2
2000[4]	12.4	*	*	*	*	*	*	*	*	*	1.9	2.9	8.3	17.5
1999[4]	12.9	*	*	*	*	*	*	*	*	*	1.5	3.9	6.6	18.3
Pneumoconioses and chemical effects (J60-J66,J68)														
All origins[3]														
Both sexes														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	0.2
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	0.2
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.8	*	*	*	*	*	*	*	*	*	*	*	*	0.4
2000[4]	0.8	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.8	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)						
Non-Hispanic White						
Male						
2001	58.9	120.0	223.5	361.0	525.0	823.9
2000[4]	58.6	119.8	224.5	353.8	558.0	876.5
1999[4]	63.0	128.5	237.0	372.4	566.3	900.5
Female						
2001	53.4	104.9	172.5	253.3	345.3	486.6
2000[4]	51.0	101.5	169.7	250.1	348.8	468.2
1999[4]	51.5	99.2	167.5	246.7	348.6	456.1
Non-Hispanic Black						
Both sexes						
2001	38.2	77.1	119.8	175.2	240.6	323.4
2000[4]	42.9	73.5	122.6	175.9	229.9	312.6
1999[4]	43.7	78.4	130.6	201.5	250.1	314.0
Male						
2001	51.0	107.6	180.8	288.7	414.6	592.5
2000[4]	56.8	105.0	191.0	286.6	386.2	578.9
1999[4]	62.0	111.3	192.5	329.2	442.8	610.9
Female						
2001	28.1	54.6	79.1	107.5	152.0	223.7
2000[4]	31.9	50.3	77.1	109.2	151.2	213.7
1999[4]	29.4	53.9	89.6	124.6	154.5	203.0
Pneumoconioses and chemical effects (J60-J66,J68)						
All origins[3]						
Both sexes						
2001	0.3	1.0	2.0	3.4	5.1	6.2
2000[4]	0.3	0.9	2.2	3.4	4.8	7.5
1999[4]	0.4	1.1	1.9	3.4	5.2	6.1
Male						
2001	0.7	2.0	4.4	7.8	13.2	20.2
2000[4]	0.7	1.8	4.8	7.9	12.3	24.9
1999[4]	0.7	2.3	4.4	8.1	13.5	20.4
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Pneumoconioses and chemical effects (J60-J66,J68)														
All origins[3]														
Female														
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	0.2
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	0.2
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.9	*	*	*	*	*	*	*	*	*	*	*	*	0.4
2000[4]	0.9	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.9	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	0.5	*	*	*	*	*	*	*	*	*	*	*	*	0.2
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	0.2
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	1.0	*	*	*	*	*	*	*	*	*	*	*	*	0.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumoconioses and chemical effects (J60-J66,J68)						
All origins[3]						
Female						
1999[4]	*	*	*	*	*	*
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	0.3	1.0	2.0	3.5	5.3	6.2
2000[4]	0.4	0.9	2.3	3.5	4.7	7.6
1999[4]	0.4	1.1	2.0	3.5	5.3	6.2
Male						
2001	0.7	2.2	4.5	8.1	13.6	20.5
2000[4]	0.7	1.8	5.0	8.1	12.2	25.7
1999[4]	0.8	2.3	4.5	8.4	14.0	21.1
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic White						
Both sexes						
2001	0.3	1.1	2.2	3.8	5.5	6.6
2000[4]	0.4	1.0	2.5	3.6	4.9	8.1
1999[4]	0.3	1.2	2.1	3.8	5.7	6.4
Male						
2001	0.7	2.4	4.8	8.7	14.2	21.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Pneumoconioses and chemical effects (J60-J66,J68)														
Non-Hispanic White														
Male														
2000[4]	1.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Pneumonitis due to solids and liquids (J69)														
All origins[3]														
Both sexes														
2001	6.1	*	*	*	*	*	0.1	0.2	0.2	0.3	0.5	0.7	1.3	1.8
2000[4]	5.9	*	*	*	*	*	0.1	0.1	0.2	0.3	0.5	0.8	1.2	1.9
1999[4]	5.5	*	*	*	*	*	0.1	0.2	0.2	0.3	0.5	0.7	0.9	2.0
Male														
2001	6.1	*	*	*	*	*	*	*	0.2	0.4	0.7	0.8	1.7	2.3
2000[4]	6.1	*	*	*	*	*	*	*	0.2	0.4	0.6	1.0	1.5	2.4
1999[4]	5.7	*	*	*	*	*	*	0.2	0.3	0.3	0.6	0.9	1.0	2.4
Female														
2001	6.0	*	*	*	*	*	*	*	0.2	*	0.4	0.6	0.9	1.4
2000[4]	5.8	*	*	*	*	*	*	*	*	0.2	0.4	0.6	0.9	1.5
1999[4]	5.3	*	*	*	*	*	*	*	*	0.2	0.4	0.5	0.8	1.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumoconioses and chemical effects (J60-J66,J68)						
Non-Hispanic White						
Male						
2000[4]	0.8	1.9	5.3	8.5	12.7	27.2
1999[4]	0.7	2.5	4.8	8.8	14.9	21.8
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Pneumonitis due to solids and liquids (J69)						
All origins[3]						
Both sexes						
2001	3.6	6.2	14.1	30.4	68.0	189.4
2000[4]	3.2	6.7	14.2	30.4	65.7	187.6
1999[4]	3.1	6.1	13.2	28.2	60.7	175.6
Male						
2001	4.2	8.2	19.1	41.0	98.1	270.3
2000[4]	3.9	9.2	20.2	42.4	95.2	276.0
1999[4]	4.2	8.3	18.4	41.6	90.1	258.4
Female						
2001	3.0	4.4	10.2	23.0	49.9	155.6
2000[4]	2.5	4.6	9.4	22.1	48.2	151.6
1999[4]	2.0	4.2	9.2	18.8	43.5	142.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Pneumonitis due to solids and liquids (J69)														
Hispanic														
Both sexes														
2001	1.5	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	1.5	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	1.5	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	6.7	*	*	*	*	*	0.1	0.2	0.3	0.3	0.5	0.7	1.3	1.8
2000[4]	6.5	*	*	*	*	*	0.1	*	0.2	0.3	0.5	0.8	1.2	1.9
1999[4]	6.0	*	*	*	*	*	0.1	0.2	0.3	0.3	0.5	0.7	1.0	2.1
Male														
2001	6.8	*	*	*	*	*	*	*	0.3	0.4	0.7	0.8	1.7	2.3
2000[4]	6.8	*	*	*	*	*	*	*	0.3	0.4	0.7	1.0	1.5	2.3
1999[4]	6.3	*	*	*	*	*	*	*	0.3	0.3	0.6	0.9	1.0	2.5
Female														
2001	6.6	*	*	*	*	*	*	*	0.3	*	0.4	0.6	0.9	1.4
2000[4]	6.3	*	*	*	*	*	*	*	*	*	0.4	0.6	1.0	1.5
1999[4]	5.8	*	*	*	*	*	*	*	*	0.2	0.4	0.5	0.9	1.7
Non-Hispanic White														
Both sexes														
2001	7.5	*	*	*	*	*	*	0.2	0.2	0.3	0.5	0.7	1.2	1.8
2000[4]	7.3	*	*	*	*	*	*	*	0.2	0.2	0.6	0.7	1.1	1.8
1999[4]	6.7	*	*	*	*	*	*	0.2	0.2	0.3	0.5	0.6	0.9	1.9
Male														
2001	7.6	*	*	*	*	*	*	*	*	0.4	0.7	0.9	1.6	2.3
2000[4]	7.5	*	*	*	*	*	*	*	*	0.4	0.7	0.9	1.3	2.3
1999[4]	6.9	*	*	*	*	*	*	*	*	0.3	0.6	0.8	0.9	2.2
Female														
2001	7.4	*	*	*	*	*	*	*	*	*	0.4	0.5	0.9	1.4
2000[4]	7.1	*	*	*	*	*	*	*	*	*	0.4	0.5	0.9	1.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)						
Hispanic						
Both sexes						
2001	*	4.3	8.3	19.0	42.2	161.3
2000[4]	*	3.5	9.0	20.8	40.7	137.4
1999[4]	*	3.6	7.9	18.4	35.8	130.5
Male						
2001	*	*	11.4	25.2	57.1	220.9
2000[4]	*	*	12.2	32.5	63.3	169.3
1999[4]	*	*	*	26.5	48.7	159.6
Female						
2001	*	*	*	14.5	33.0	132.2
2000[4]	*	*	*	12.5	26.9	121.7
1999[4]	*	*	*	12.7	28.1	116.5
Non-Hispanic[5]						
Both sexes						
2001	3.7	6.3	14.4	30.9	68.8	190.0
2000[4]	3.2	7.0	14.5	30.8	66.4	189.0
1999[4]	3.1	6.2	13.4	28.5	61.4	176.8
Male						
2001	4.4	8.4	19.5	41.7	99.6	271.6
2000[4]	4.0	9.6	20.6	42.6	96.0	279.7
1999[4]	4.3	8.5	18.7	42.1	91.4	261.6
Female						
2001	3.1	4.5	10.4	23.4	50.4	156.0
2000[4]	2.5	4.7	9.6	22.6	48.9	152.3
1999[4]	2.0	4.3	9.3	19.0	43.8	142.9
Non-Hispanic White						
Both sexes						
2001	3.6	6.1	13.9	30.7	69.0	191.2
2000[4]	3.0	6.6	13.6	30.5	66.3	192.5
1999[4]	3.0	5.9	12.7	27.9	60.9	179.3
Male						
2001	4.1	8.1	18.8	41.3	100.4	276.9
2000[4]	3.6	8.8	19.0	41.7	96.9	288.1
1999[4]	4.0	7.9	17.4	40.9	90.2	267.4
Female						
2001	3.1	4.3	9.8	23.2	50.1	155.5
2000[4]	2.3	4.6	9.3	22.6	48.2	153.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Pneumonitis due to solids and liquids (J69)														
Non-Hispanic White														
Female														
1999[4]	6.5	*	*	*	*	*	*	*	*	*	0.4	0.4	0.8	1.5
Non-Hispanic Black														
Both sexes														
2001	4.4	*	*	*	*	*	*	*	*	*	0.8	1.0	2.5	2.5
2000[4]	4.3	*	*	*	*	*	*	*	*	0.7	*	1.7	2.3	3.0
1999[4]	4.3	*	*	*	*	*	*	*	*	*	1.0	1.3	1.8	4.0
Male														
2001	4.3	*	*	*	*	*	*	*	*	*	*	*	3.5	3.2
2000[4]	4.5	*	*	*	*	*	*	*	*	*	*	2.4	3.2	3.4
1999[4]	4.7	*	*	*	*	*	*	*	*	*	*	*	*	5.2
Female														
2001	4.6	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	4.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	4.0	*	*	*	*	*	*	*	*	*	*	*	*	3.0
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)														
All origins[3]														
Both sexes														
2001	9.2	7.3	0.6	0.1	0.2	0.3	0.4	0.5	0.6	1.0	1.5	2.6	4.4	7.3
2000[4]	9.1	7.2	0.6	0.2	0.2	0.3	0.4	0.5	0.8	1.0	1.7	2.6	4.6	7.8
1999[4]	9.0	6.6	0.6	0.2	0.2	0.3	0.3	0.5	0.8	1.0	1.6	2.7	4.1	6.9
Male														
2001	8.9	9.0	0.6	*	0.2	0.3	0.5	0.5	0.6	0.9	1.8	2.9	4.9	7.7
2000[4]	9.0	7.9	0.6	0.2	*	0.3	0.4	0.5	0.8	1.2	1.8	2.8	5.0	8.8
1999[4]	8.9	7.5	0.7	0.2	*	0.4	0.3	0.5	0.8	1.1	1.7	2.8	4.4	7.4
Female														
2001	9.4	5.5	0.6	*	0.2	0.3	0.4	0.6	0.6	1.0	1.3	2.3	4.0	6.8
2000[4]	9.2	6.4	0.6	*	0.2	0.3	0.3	0.5	0.7	0.9	1.6	2.4	4.2	6.8
1999[4]	9.1	5.8	0.5	*	0.2	0.2	0.3	0.5	0.8	0.9	1.4	2.6	3.8	6.5
Hispanic														
Both sexes														
2001	3.4	5.0	*	*	*	*	*	*	*	*	1.0	2.2	4.3	7.9
2000[4]	3.5	5.6	*	*	*	*	*	*	0.6	1.0	1.3	2.2	3.1	8.6
1999[4]	3.4	5.0	*	*	*	*	*	*	*	*	1.5	2.3	4.3	5.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)						
Non-Hispanic White						
Female						
1999[4]	2.0	4.2	8.9	18.8	43.7	144.2
Non-Hispanic Black						
Both sexes						
2001	6.1	9.5	22.7	39.2	76.2	187.3
2000[4]	5.9	11.5	25.5	39.1	72.4	160.6
1999[4]	5.2	10.2	24.0	40.9	71.6	168.5
Male						
2001	8.4	12.8	31.3	56.9	107.9	243.7
2000[4]	7.4	17.9	42.0	62.1	97.8	210.2
1999[4]	8.3	15.7	37.7	65.3	111.6	229.4
Female						
2001	4.3	7.1	17.0	28.6	60.0	166.4
2000[4]	4.7	6.9	14.5	25.3	59.6	142.2
1999[4]	*	6.1	14.9	26.2	51.8	145.7
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)						
All origins[3]						
Both sexes						
2001	12.6	22.8	38.0	61.8	89.6	142.7
2000[4]	12.7	22.9	36.9	60.5	89.2	141.7
1999[4]	12.7	22.5	36.8	59.5	87.6	147.6
Male						
2001	15.0	27.1	47.0	76.2	110.0	169.1
2000[4]	14.6	26.8	45.7	77.3	112.4	179.7
1999[4]	14.7	27.3	46.4	74.9	115.0	182.3
Female						
2001	10.5	19.1	30.8	51.7	77.3	131.6
2000[4]	10.8	19.6	29.9	48.8	75.6	126.2
1999[4]	10.9	18.4	29.3	48.8	71.6	133.6
Hispanic						
Both sexes						
2001	10.3	19.8	32.1	52.7	87.9	134.4
2000[4]	11.2	20.5	31.4	53.3	84.1	135.4
1999[4]	14.3	22.6	30.2	56.2	78.1	134.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)														
Hispanic														
Male														
2001	3.4	5.0	*	*	*	*	*	*	*	*	*	2.5	4.7	8.6
2000[4]	3.3	5.8	*	*	*	*	*	*	*	*	*	2.5	3.6	9.6
1999[4]	3.3	5.2	*	*	*	*	*	*	*	*	2.0	2.6	3.7	7.2
Female														
2001	3.5	5.0	*	*	*	*	*	*	*	*	*	*	4.0	7.2
2000[4]	3.6	5.3	*	*	*	*	*	*	*	*	*	*	*	7.7
1999[4]	3.5	*	*	*	*	*	*	*	*	*	*	*	4.9	4.6
Non-Hispanic[5]														
Both sexes														
2001	10.0	7.8	0.6	0.2	0.2	0.3	0.4	0.6	0.6	1.1	1.6	2.6	4.4	7.2
2000[4]	9.8	7.5	0.6	0.2	0.2	0.3	0.4	0.5	0.8	1.0	1.8	2.7	4.7	7.6
1999[4]	9.7	6.9	0.6	0.2	0.2	0.3	0.3	0.5	0.8	1.1	1.6	2.7	4.1	6.9
Male														
2001	9.8	10.0	0.6	*	0.2	0.3	0.5	0.5	0.6	1.0	1.8	2.9	4.9	7.6
2000[4]	9.8	8.4	0.6	0.2	*	0.3	0.5	0.5	0.8	1.1	1.9	2.9	5.1	8.7
1999[4]	9.6	7.9	0.8	0.2	*	0.4	0.4	0.5	0.9	1.2	1.7	2.8	4.5	7.3
Female														
2001	10.2	5.5	0.6	*	0.2	0.3	0.4	0.7	0.6	1.1	1.4	2.4	3.9	6.8
2000[4]	9.9	6.6	0.5	*	*	0.3	0.3	0.6	0.7	0.9	1.6	2.4	4.3	6.6
1999[4]	9.8	5.9	0.5	*	0.2	*	0.3	0.6	0.8	1.0	1.4	2.6	3.7	6.6
Non-Hispanic White														
Both sexes														
2001	11.0	4.8	0.5	0.2	0.2	0.3	0.4	0.5	0.5	0.8	1.4	2.2	3.8	6.9
2000[4]	10.8	5.4	0.5	0.2	0.2	0.3	0.3	0.5	0.7	0.8	1.6	2.2	4.4	7.4
1999[4]	10.6	4.9	0.6	0.2	0.2	0.3	0.3	0.4	0.7	0.9	1.3	2.3	3.7	6.4
Male														
2001	10.8	6.2	0.4	*	*	0.3	0.4	0.4	0.4	0.8	1.7	2.5	4.4	7.5
2000[4]	10.7	6.6	0.5	*	*	*	0.4	0.4	0.7	0.9	1.7	2.4	4.7	8.4
1999[4]	10.5	5.6	0.6	*	*	0.4	0.4	0.5	0.8	1.0	1.5	2.3	4.2	6.5
Female														
2001	11.3	3.4	0.5	*	*	*	0.3	0.6	0.5	0.8	1.1	2.0	3.3	6.3
2000[4]	11.0	4.1	0.5	*	*	*	*	0.5	0.6	0.7	1.5	2.1	4.0	6.4
1999[4]	10.7	4.0	0.5	*	*	*	*	0.4	0.7	0.8	1.1	2.2	3.3	6.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)						
Hispanic						
Male						
2001	13.3	23.4	38.9	64.8	103.9	167.6
2000[4]	15.0	22.7	37.9	67.9	107.5	133.0
1999[4]	17.3	26.7	35.6	71.0	105.3	161.8
Female						
2001	7.8	16.9	26.9	44.1	78.0	118.2
2000[4]	7.9	18.7	26.5	42.9	69.9	136.5
1999[4]	11.8	19.4	26.0	45.7	61.9	120.6
Non-Hispanic[5]						
Both sexes						
2001	12.8	23.0	38.2	62.1	89.4	142.7
2000[4]	12.7	23.0	37.0	60.7	89.1	141.6
1999[4]	12.5	22.4	37.0	59.5	87.7	147.7
Male						
2001	15.1	27.3	47.2	76.5	110.0	169.1
2000[4]	14.6	27.0	46.0	77.5	112.3	181.3
1999[4]	14.5	27.2	46.8	74.8	114.8	182.7
Female						
2001	10.7	19.2	31.0	52.0	77.1	131.8
2000[4]	11.1	19.6	30.0	49.0	75.5	125.5
1999[4]	10.8	18.2	29.4	48.8	71.8	133.7
Non-Hispanic White						
Both sexes						
2001	12.5	23.2	39.0	63.8	92.2	145.9
2000[4]	12.5	22.9	37.9	62.1	91.0	145.7
1999[4]	12.4	22.5	37.2	60.6	89.6	150.7
Male						
2001	14.5	27.7	48.6	78.9	113.3	173.4
2000[4]	14.2	27.0	46.6	79.0	114.9	185.9
1999[4]	14.4	27.6	47.4	76.5	117.4	186.1
Female						
2001	10.6	19.1	31.2	53.1	79.4	134.5
2000[4]	10.8	19.3	30.8	50.3	76.9	129.5
1999[4]	10.6	18.0	29.1	49.4	73.2	136.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)														
Non-Hispanic Black														
Both sexes														
2001	6.9	19.0	1.1	*	*	*	0.8	1.2	1.4	2.6	3.3	5.9	9.5	11.2
2000[4]	6.7	17.7	0.8	*	*	*	0.8	0.9	1.4	2.4	3.1	6.1	7.9	11.8
1999[4]	6.7	15.8	1.1	*	*	*	*	1.1	1.7	2.4	3.4	6.0	7.3	12.1
Male														
2001	6.6	24.3	*	*	*	*	*	*	1.6	2.0	3.2	6.4	9.6	10.8
2000[4]	6.8	17.1	*	*	*	*	*	*	*	3.0	3.5	7.4	8.7	13.7
1999[4]	6.6	18.0	*	*	*	*	*	*	1.8	2.5	3.2	6.8	7.9	14.1
Female														
2001	7.2	13.4	*	*	*	*	*	*	*	3.0	3.4	5.5	9.4	11.5
2000[4]	6.6	18.3	*	*	*	*	*	*	*	1.9	2.7	5.1	7.2	10.3
1999[4]	6.9	13.6	*	*	*	*	*	1.8	1.6	2.3	3.7	5.3	6.8	10.4
Peptic ulcer (K25-K28)														
All origins[3]														
Both sexes														
2001	1.6	*	*	*	*	*	*	*	*	0.2	0.4	0.7	0.8	1.3
2000[4]	1.6	*	*	*	*	*	*	*	*	0.2	0.3	0.6	1.0	1.5
1999[4]	1.6	*	*	*	*	*	*	*	*	0.2	0.4	0.6	0.8	1.4
Male														
2001	1.5	*	*	*	*	*	*	*	*	0.2	0.5	0.9	1.1	1.8
2000[4]	1.6	*	*	*	*	*	*	*	*	0.3	0.4	0.9	1.5	2.1
1999[4]	1.6	*	*	*	*	*	*	*	*	0.2	0.4	0.7	1.2	1.8
Female														
2001	1.7	*	*	*	*	*	*	*	*	*	0.2	0.4	0.6	0.8
2000[4]	1.6	*	*	*	*	*	*	*	*	*	0.2	0.4	0.5	0.9
1999[4]	1.7	*	*	*	*	*	*	*	*	*	0.3	0.5	0.5	1.1
Hispanic														
Both sexes														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)						
Non-Hispanic Black						
Both sexes						
2001	17.4	24.3	36.1	50.3	67.0	110.8
2000[4]	15.9	28.2	33.5	50.2	71.8	99.8
1999[4]	15.6	25.2	38.7	53.6	66.9	112.8
Male						
2001	22.2	27.9	39.6	59.7	85.0	131.7
2000[4]	18.1	32.2	41.3	68.9	83.3	135.0
1999[4]	18.1	28.7	44.7	67.2	85.0	146.2
Female						
2001	13.6	21.7	33.7	44.7	57.9	103.1
2000[4]	14.3	25.2	28.4	38.9	66.1	86.8
1999[4]	13.7	22.6	34.7	45.4	58.0	100.4
Peptic ulcer (K25-K28)						
All origins[3]						
Both sexes						
2001	2.1	3.1	4.8	8.3	16.0	33.4
2000[4]	2.2	3.1	5.1	8.7	15.6	35.6
1999[4]	2.2	3.3	5.9	9.1	17.1	34.6
Male						
2001	2.8	4.1	5.6	10.3	17.3	34.9
2000[4]	2.9	3.9	6.9	11.2	17.8	40.7
1999[4]	3.0	4.5	7.9	11.0	21.1	37.5
Female						
2001	1.5	2.2	4.2	6.9	15.2	32.7
2000[4]	1.6	2.5	3.6	6.9	14.4	33.5
1999[4]	1.5	2.3	4.3	7.7	14.7	33.5
Hispanic						
Both sexes						
2001	*	*	4.6	6.8	14.9	23.8
2000[4]	*	*	4.8	*	17.3	30.5
1999[4]	*	*	4.6	6.5	14.1	26.7
Male						
2001	*	*	*	*	26.0	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Peptic ulcer (K25-K28)														
Hispanic														
Female														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	1.7	*	*	*	*	*	*	*	*	0.2	0.4	0.7	0.8	1.3
2000[4]	1.8	*	*	*	*	*	*	*	*	0.2	0.3	0.7	1.0	1.5
1999[4]	1.8	*	*	*	*	*	*	*	*	0.2	0.4	0.6	0.9	1.4
Male														
2001	1.6	*	*	*	*	*	*	*	*	0.3	0.6	0.9	1.1	1.8
2000[4]	1.8	*	*	*	*	*	*	*	*	0.3	0.4	0.9	1.6	2.1
1999[4]	1.7	*	*	*	*	*	*	*	*	0.3	0.5	0.7	1.2	1.7
Female														
2001	1.8	*	*	*	*	*	*	*	*	*	0.2	0.5	0.6	0.8
2000[4]	1.8	*	*	*	*	*	*	*	*	*	0.2	0.5	0.5	1.0
1999[4]	1.9	*	*	*	*	*	*	*	*	*	0.3	0.5	0.5	1.1
Non-Hispanic White														
Both sexes														
2001	1.9	*	*	*	*	*	*	*	*	*	0.4	0.6	0.7	1.1
2000[4]	1.9	*	*	*	*	*	*	*	*	0.2	0.3	0.6	0.9	1.4
1999[4]	2.0	*	*	*	*	*	*	*	*	0.1	0.3	0.5	0.7	1.3
Male														
2001	1.7	*	*	*	*	*	*	*	*	*	0.5	0.8	0.9	1.5
2000[4]	1.9	*	*	*	*	*	*	*	*	*	0.4	0.8	1.3	2.0
1999[4]	1.8	*	*	*	*	*	*	*	*	*	0.3	0.6	1.1	1.5
Female														
2001	2.1	*	*	*	*	*	*	*	*	*	*	0.5	0.6	0.8
2000[4]	2.0	*	*	*	*	*	*	*	*	*	*	0.4	0.5	0.9
1999[4]	2.1	*	*	*	*	*	*	*	*	*	0.3	0.4	0.4	1.1
Non-Hispanic Black														
Both sexes														
2001	1.1	*	*	*	*	*	*	*	*	*	*	1.1	1.6	2.5
2000[4]	1.2	*	*	*	*	*	*	*	*	*	*	1.4	2.1	2.6
1999[4]	1.3	*	*	*	*	*	*	*	*	*	1.0	1.3	1.9	2.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Peptic ulcer (K25-K28)						
Hispanic						
Female						
2001	*	*	*	*	*	25.1
2000[4]	*	*	*	*	22.4	29.7
1999[4]	*	*	*	*	*	22.9
Non-Hispanic[5]						
Both sexes						
2001	2.2	3.1	4.8	8.3	16.0	33.6
2000[4]	2.2	3.2	5.1	8.8	15.5	35.7
1999[4]	2.2	3.3	5.9	9.1	17.2	34.8
Male						
2001	2.8	4.1	5.6	10.2	16.8	35.5
2000[4]	2.8	3.9	6.9	11.3	18.1	41.0
1999[4]	3.0	4.5	8.0	10.9	21.1	37.7
Female						
2001	1.6	2.3	4.3	7.0	15.5	32.8
2000[4]	1.7	2.5	3.6	7.0	14.0	33.5
1999[4]	1.5	2.3	4.3	7.9	14.9	33.7
Non-Hispanic White						
Both sexes						
2001	2.1	3.0	4.8	8.3	16.3	34.7
2000[4]	2.2	3.1	5.1	8.7	15.9	36.3
1999[4]	2.2	3.1	5.8	9.2	17.2	35.6
Male						
2001	2.7	3.9	5.3	9.9	16.9	36.3
2000[4]	2.8	3.8	6.7	11.0	18.1	41.0
1999[4]	2.9	4.1	7.9	11.0	21.1	38.6
Female						
2001	1.5	2.3	4.5	7.1	15.9	34.0
2000[4]	1.7	2.4	3.7	7.0	14.5	34.4
1999[4]	1.6	2.2	4.3	7.9	14.9	34.4
Non-Hispanic Black						
Both sexes						
2001	3.1	3.5	5.6	8.0	10.5	20.9
2000[4]	2.7	4.2	5.7	9.4	11.2	26.7
1999[4]	3.2	5.1	6.5	7.6	16.4	25.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Peptic ulcer (K25-K28)														
Non-Hispanic Black														
Male														
2001	1.4	*	*	*	*	*	*	*	*	*	*	1.9	2.8	4.0
2000[4]	1.4	*	*	*	*	*	*	*	*	*	*	*	3.4	3.6
1999[4]	1.5	*	*	*	*	*	*	*	*	*	*	1.9	2.9	3.8
Female														
2001	0.9	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Diseases of appendix (K35-K38)														
All origins[3]														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	0.1	*	0.2
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	0.1	0.2
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	0.1	0.1	0.2
Male														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	0.3
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Peptic ulcer (K25-K28)						
Non-Hispanic Black						
Male						
2001	4.6	5.2	9.3	10.9	*	24.3
2000[4]	*	*	10.3	12.5	*	31.7
1999[4]	5.2	7.7	9.0	*	18.6	28.5
Female						
2001	*	*	*	6.2	9.2	19.7
2000[4]	*	*	*	7.6	8.6	24.9
1999[4]	*	*	4.8	7.0	15.4	24.4
Diseases of appendix (K35-K38)						
All origins[3]						
Both sexes						
2001	0.2	0.4	0.6	0.5	1.0	2.1
2000[4]	0.3	0.3	0.5	0.7	1.4	1.6
1999[4]	0.2	0.3	0.4	0.6	1.1	1.5
Male						
2001	*	0.5	0.9	0.7	1.6	2.8
2000[4]	0.4	*	0.6	1.0	1.6	2.0
1999[4]	*	*	0.5	0.9	*	2.9
Female						
2001	*	*	0.4	*	0.7	1.8
2000[4]	*	*	0.4	0.5	1.2	1.5
1999[4]	*	*	*	*	1.1	0.9
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Diseases of appendix (K35-K38)														
Non-Hispanic[5]														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	0.1	*	0.2
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	0.2	0.2
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	0.1	0.1	0.2
Male														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	0.2	*
Male														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of appendix (K35-K38)						
Non-Hispanic[5]						
Both sexes						
2001	0.2	0.4	0.6	0.5	1.0	2.2
2000[4]	0.3	0.3	0.5	0.7	1.4	1.7
1999[4]	0.2	0.3	0.4	0.6	1.1	1.5
Male						
2001	*	0.5	0.8	0.7	1.5	3.0
2000[4]	0.5	*	0.7	1.0	1.6	2.0
1999[4]	*	*	*	0.9	*	3.0
Female						
2001	*	*	0.5	*	0.7	1.8
2000[4]	*	*	*	0.6	1.2	1.5
1999[4]	*	*	*	*	1.1	0.9
Non-Hispanic White						
Both sexes						
2001	*	0.3	0.7	0.5	0.9	2.1
2000[4]	0.3	0.3	0.4	0.6	1.3	1.8
1999[4]	*	0.3	0.4	0.6	1.1	1.6
Male						
2001	*	*	0.9	*	1.4	2.6
2000[4]	0.5	*	*	1.0	1.5	2.1
1999[4]	*	*	*	0.9	*	3.1
Female						
2001	*	*	*	*	*	2.0
2000[4]	*	*	*	*	1.2	1.6
1999[4]	*	*	*	*	1.1	1.0
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Diseases of appendix (K35-K38)														
Non-Hispanic Black														
Female														
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Hernia (K40-K46)														
All origins[3]														
Both sexes														
2001	0.5	1.1	*	*	*	*	*	*	*	*	0.1	0.1	0.3	0.4
2000[4]	0.5	1.2	*	*	*	*	*	*	*	*	*	0.1	0.3	0.4
1999[4]	0.5	1.2	*	*	*	*	*	*	*	*	0.1	*	0.2	0.3
Male														
2001	0.4	1.4	*	*	*	*	*	*	*	*	*	*	0.3	0.3
2000[4]	0.4	1.4	*	*	*	*	*	*	*	*	*	*	0.4	0.3
1999[4]	0.4	1.3	*	*	*	*	*	*	*	*	*	*	0.2	0.3
Female														
2001	0.6	*	*	*	*	*	*	*	*	*	*	*	0.3	0.5
2000[4]	0.7	*	*	*	*	*	*	*	*	*	*	*	0.3	0.4
1999[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.6	1.0	*	*	*	*	*	*	*	*	*	0.1	0.3	0.4
2000[4]	0.6	1.2	*	*	*	*	*	*	*	*	*	0.1	0.3	0.4
1999[4]	0.6	1.3	*	*	*	*	*	*	*	*	0.1	*	0.2	0.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of appendix (K35-K38)						
Non-Hispanic Black						
Female						
1999[4]	*	*	*	*	*	*
Hernia (K40-K46)						
All origins[3]						
Both sexes						
2001	0.7	0.8	1.5	2.5	5.0	12.6
2000[4]	0.6	0.9	1.4	3.0	4.9	13.1
1999[4]	0.7	0.8	1.6	2.5	5.4	12.7
Male						
2001	0.8	0.9	1.4	2.3	5.3	12.0
2000[4]	0.6	0.9	1.4	3.0	5.7	12.6
1999[4]	0.8	0.8	1.5	2.5	6.5	13.2
Female						
2001	0.6	0.8	1.6	2.6	4.8	12.8
2000[4]	0.6	0.9	1.5	3.0	4.4	13.3
1999[4]	0.6	0.8	1.6	2.6	4.8	12.6
Hispanic						
Both sexes						
2001	*	*	*	*	*	12.5
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	0.6	0.9	1.5	2.4	5.0	12.5
2000[4]	0.6	0.9	1.4	2.9	4.9	13.1
1999[4]	0.7	0.8	1.5	2.5	5.4	12.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Hernia (K40-K46)														
Non-Hispanic[5]														
Male														
2001	0.5	1.2	*	*	*	*	*	*	*	*	*	*	0.3	0.4
2000[4]	0.5	1.4	*	*	*	*	*	*	*	*	*	*	0.4	*
1999[4]	0.5	1.5	*	*	*	*	*	*	*	*	*	*	*	0.4
Female														
2001	0.7	*	*	*	*	*	*	*	*	*	*	*	0.3	0.5
2000[4]	0.7	*	*	*	*	*	*	*	*	*	*	*	0.3	0.5
1999[4]	0.7	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	0.6	0.9	*	*	*	*	*	*	*	*	*	*	0.3	0.4
2000[4]	0.6	1.2	*	*	*	*	*	*	*	*	*	*	0.3	0.4
1999[4]	0.6	1.4	*	*	*	*	*	*	*	*	*	*	0.2	0.3
Male														
2001	0.5	*	*	*	*	*	*	*	*	*	*	*	0.3	*
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	0.3	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	0.4
Female														
2001	0.8	*	*	*	*	*	*	*	*	*	*	*	0.3	0.5
2000[4]	0.8	*	*	*	*	*	*	*	*	*	*	*	0.3	0.5
1999[4]	0.7	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hernia (K40-K46)						
Non-Hispanic[5]						
Male						
2001	0.8	0.9	1.4	2.4	5.2	12.1
2000[4]	0.6	0.9	1.3	2.9	5.6	12.8
1999[4]	0.7	0.8	1.4	2.5	6.5	13.3
Female						
2001	0.6	0.8	1.5	2.5	4.8	12.7
2000[4]	0.5	0.8	1.5	3.0	4.5	13.2
1999[4]	0.6	0.8	1.6	2.5	4.7	12.4
Non-Hispanic White						
Both sexes						
2001	0.6	0.9	1.5	2.4	5.0	12.8
2000[4]	0.6	0.8	1.5	2.9	5.0	13.3
1999[4]	0.7	0.7	1.5	2.5	5.5	13.0
Male						
2001	0.7	0.9	1.4	2.4	5.2	11.9
2000[4]	0.6	0.9	1.4	3.0	5.7	13.0
1999[4]	0.8	0.8	1.5	2.5	6.6	13.0
Female						
2001	0.5	0.8	1.5	2.4	4.9	13.1
2000[4]	0.5	0.7	1.5	2.8	4.6	13.4
1999[4]	0.6	0.6	1.5	2.4	4.8	12.9
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	11.2
2000[4]	*	*	*	3.8	*	12.7
1999[4]	*	*	*	*	*	10.0
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	9.0
2000[4]	*	*	*	*	*	12.6
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Chronic liver disease and cirrhosis (K70,K73-K74)														
All origins[3]														
Both sexes														
2001	9.5	*	*	*	*	*	0.1	0.4	1.5	5.0	9.8	17.1	20.1	21.1
2000[4]	9.4	*	*	*	*	*	0.2	0.5	1.6	4.8	10.2	16.8	18.7	21.9
1999[4]	9.4	*	*	*	*	*	*	0.4	1.6	4.8	9.9	16.4	18.6	21.4
Male														
2001	12.4	*	*	*	*	*	*	0.6	2.0	6.4	13.2	25.2	30.1	31.5
2000[4]	12.5	*	*	*	*	*	*	0.6	2.1	6.3	14.4	25.4	28.6	31.7
1999[4]	12.5	*	*	*	*	*	*	0.4	2.1	6.3	14.5	25.3	28.8	32.5
Female														
2001	6.7	*	*	*	*	*	*	0.3	1.0	3.5	6.4	9.3	10.5	11.4
2000[4]	6.5	*	*	*	*	*	*	0.3	1.1	3.2	6.1	8.4	9.1	12.7
1999[4]	6.4	*	*	*	*	*	*	0.3	1.2	3.4	5.4	7.8	8.8	11.1
Hispanic														
Both sexes														
2001	8.9	*	*	*	*	*	*	*	2.0	6.3	13.9	25.0	31.9	35.9
2000[4]	9.0	*	*	*	*	*	*	*	2.4	6.0	14.4	24.8	30.4	39.6
1999[4]	8.8	*	*	*	*	*	*	*	2.4	7.0	14.9	25.2	28.4	35.6
Male														
2001	12.7	*	*	*	*	*	*	*	2.9	9.6	21.7	41.9	52.5	58.9
2000[4]	12.7	*	*	*	*	*	*	*	4.1	9.1	22.5	41.2	51.5	64.7
1999[4]	12.3	*	*	*	*	*	*	*	3.4	10.8	23.7	42.3	46.9	59.4
Female														
2001	5.0	*	*	*	*	*	*	*	*	2.6	5.6	8.0	12.2	15.0
2000[4]	5.1	*	*	*	*	*	*	*	*	2.7	5.9	8.4	10.3	16.9
1999[4]	5.0	*	*	*	*	*	*	*	*	2.8	5.7	8.2	10.8	14.2
Non-Hispanic[5]														
Both sexes														
2001	9.5	*	*	*	*	*	*	0.4	1.4	4.8	9.2	16.3	19.0	19.8
2000[4]	9.4	*	*	*	*	*	0.1	0.5	1.4	4.6	9.7	15.9	17.5	20.4
1999[4]	9.5	*	*	*	*	*	*	0.4	1.5	4.5	9.3	15.5	17.6	20.3
Male														
2001	12.3	*	*	*	*	*	*	0.5	1.8	5.9	12.1	23.4	28.0	29.1
2000[4]	12.4	*	*	*	*	*	*	0.6	1.7	5.9	13.3	23.7	26.4	29.1
1999[4]	12.5	*	*	*	*	*	*	0.4	1.8	5.6	13.4	23.6	27.1	30.3
Female														
2001	6.9	*	*	*	*	*	*	0.3	1.0	3.7	6.5	9.3	10.3	11.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic liver disease and cirrhosis (K70,K73-K74)						
All origins[3]						
Both sexes						
2001	24.7	28.9	31.1	31.3	28.7	22.2
2000[4]	26.2	28.9	30.8	31.9	29.6	23.1
1999[4]	26.5	29.7	31.6	32.6	30.8	23.2
Male						
2001	35.3	39.7	40.8	37.6	35.3	32.0
2000[4]	37.7	39.3	40.7	40.7	34.6	34.3
1999[4]	37.2	41.8	41.5	40.1	38.7	33.1
Female						
2001	15.2	19.6	23.5	26.9	24.7	18.1
2000[4]	15.7	19.9	23.1	25.8	26.6	18.5
1999[4]	16.8	19.4	23.9	27.4	26.2	19.3
Hispanic						
Both sexes						
2001	43.2	51.1	58.6	62.0	57.6	36.9
2000[4]	43.7	55.6	61.8	65.8	60.2	48.4
1999[4]	43.2	53.6	59.7	60.4	57.6	51.2
Male						
2001	70.6	78.0	75.9	81.8	64.9	41.9
2000[4]	67.9	86.9	78.8	89.3	61.8	60.5
1999[4]	66.7	76.5	82.5	72.6	67.6	66.9
Female						
2001	19.5	29.3	45.4	47.9	53.1	34.4
2000[4]	22.8	30.2	49.0	49.1	59.1	42.5
1999[4]	22.8	35.0	42.5	51.8	51.6	43.7
Non-Hispanic[5]						
Both sexes						
2001	23.2	27.2	29.4	29.7	27.4	21.6
2000[4]	24.7	26.9	29.0	30.3	28.3	22.1
1999[4]	25.2	28.0	30.0	31.3	29.8	22.1
Male						
2001	32.4	36.8	38.6	35.4	34.1	31.4
2000[4]	35.3	36.0	38.5	38.4	33.4	33.0
1999[4]	35.0	39.4	39.2	38.5	37.6	31.6
Female						
2001	14.8	18.8	22.0	25.7	23.4	17.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Chronic liver disease and cirrhosis (K70,K73-K74)														
Non-Hispanic[5]														
Female														
2000[4]	6.7	*	*	*	*	*	*	0.3	1.2	3.3	6.1	8.4	9.0	12.3
1999[4]	6.6	*	*	*	*	*	*	0.3	1.2	3.4	5.3	7.8	8.5	10.9
Non-Hispanic White														
Both sexes														
2001	10.1	*	*	*	*	*	*	0.3	1.4	4.8	9.0	16.0	18.1	19.1
2000[4]	10.0	*	*	*	*	*	*	0.3	1.3	4.4	9.4	15.7	16.7	19.9
1999[4]	9.9	*	*	*	*	*	*	0.4	1.4	4.4	8.8	14.6	17.2	19.4
Male														
2001	13.0	*	*	*	*	*	*	0.4	1.8	5.8	11.7	23.2	26.7	27.7
2000[4]	13.1	*	*	*	*	*	*	0.5	1.6	5.7	13.1	23.8	25.1	28.5
1999[4]	13.1	*	*	*	*	*	*	0.5	1.8	5.5	12.7	22.3	26.4	29.0
Female														
2001	7.4	*	*	*	*	*	*	*	1.0	3.8	6.3	9.0	9.7	10.8
2000[4]	7.0	*	*	*	*	*	*	*	1.0	3.1	5.8	7.7	8.4	11.8
1999[4]	6.9	*	*	*	*	*	*	*	1.1	3.3	5.0	7.0	8.1	10.3
Non-Hispanic Black														
Both sexes														
2001	7.7	*	*	*	*	*	*	*	1.0	4.2	10.9	19.4	27.2	28.3
2000[4]	7.7	*	*	*	*	*	*	0.8	1.4	5.0	11.2	19.8	26.3	26.0
1999[4]	8.0	*	*	*	*	*	*	*	1.4	4.5	12.1	22.8	23.2	28.4
Male														
2001	10.7	*	*	*	*	*	*	*	*	5.8	15.1	28.9	42.8	44.8
2000[4]	10.3	*	*	*	*	*	*	*	*	6.8	14.8	27.2	41.2	38.9
1999[4]	11.0	*	*	*	*	*	*	*	1.8	5.6	17.5	34.8	37.1	44.3
Female														
2001	4.9	*	*	*	*	*	*	*	*	2.7	7.3	11.3	14.0	14.8
2000[4]	5.4	*	*	*	*	*	*	*	1.5	3.5	8.0	13.3	13.6	15.4
1999[4]	5.4	*	*	*	*	*	*	*	*	3.6	7.5	12.6	11.4	15.5
Alcoholic liver disease (K70)														
All origins[3]														
Both sexes														
2001	4.3	*	*	*	*	*	*	0.2	1.1	3.3	6.1	10.2	11.3	11.1
2000[4]	4.3	*	*	*	*	*	*	0.3	1.1	3.3	6.6	10.0	10.7	11.7
1999[4]	4.3	*	*	*	*	*	*	0.2	1.1	3.3	6.4	10.0	10.9	11.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic liver disease and cirrhosis (K70,K73-K74)						
Non-Hispanic[5]						
Female						
2000[4]	15.1	19.1	21.5	24.7	25.3	17.7
1999[4]	16.3	18.3	22.9	26.3	25.3	18.3
Non-Hispanic White						
Both sexes						
2001	23.5	27.7	30.2	30.9	28.6	22.5
2000[4]	24.7	27.2	29.7	31.6	29.1	22.8
1999[4]	24.7	28.3	30.8	32.2	30.9	22.6
Male						
2001	32.3	36.8	39.6	36.6	35.8	32.9
2000[4]	35.4	36.4	39.2	39.7	34.7	34.1
1999[4]	34.7	39.5	40.0	39.3	39.1	32.7
Female						
2001	15.3	19.6	22.6	26.8	24.3	18.2
2000[4]	14.8	19.0	22.1	26.0	25.9	18.2
1999[4]	15.5	18.5	23.5	27.2	26.1	18.6
Non-Hispanic Black						
Both sexes						
2001	23.8	27.1	23.0	20.6	14.6	10.6
2000[4]	26.3	27.8	25.0	18.3	17.8	14.0
1999[4]	31.4	28.3	25.1	23.3	19.1	16.8
Male						
2001	38.5	43.8	34.4	27.0	*	*
2000[4]	37.8	38.4	37.9	27.9	19.7	*
1999[4]	43.3	44.1	37.0	32.4	25.7	*
Female						
2001	12.3	14.8	15.4	16.7	14.6	9.8
2000[4]	17.3	20.1	16.3	12.5	16.9	11.8
1999[4]	22.0	16.5	17.2	17.8	15.8	15.5
Alcoholic liver disease (K70)						
All origins[3]						
Both sexes						
2001	12.0	11.1	9.0	6.9	3.9	2.4
2000[4]	12.9	11.4	9.1	7.2	4.4	2.2
1999[4]	12.6	11.6	9.4	7.3	5.2	2.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Alcoholic liver disease (K70)														
All origins[3]														
Male														
2001	6.4	*	*	*	*	*	*	0.3	1.4	4.4	8.4	15.3	17.0	18.0
2000[4]	6.5	*	*	*	*	*	*	0.5	1.5	4.3	9.4	15.3	16.9	18.0
1999[4]	6.5	*	*	*	*	*	*	0.2	1.4	4.3	9.4	15.5	17.2	18.7
Female														
2001	2.3	*	*	*	*	*	*	*	0.7	2.3	3.9	5.3	5.8	4.7
2000[4]	2.2	*	*	*	*	*	*	*	0.6	2.2	3.9	4.8	4.8	5.8
1999[4]	2.1	*	*	*	*	*	*	*	0.8	2.4	3.5	4.6	4.8	4.9
Hispanic														
Both sexes														
2001	4.7	*	*	*	*	*	*	*	1.5	4.1	8.8	16.0	18.6	20.9
2000[4]	4.8	*	*	*	*	*	*	*	1.7	4.3	10.0	15.4	18.6	22.7
1999[4]	4.6	*	*	*	*	*	*	*	1.5	4.7	9.7	16.1	17.8	21.5
Male														
2001	7.7	*	*	*	*	*	*	*	2.4	6.9	13.8	27.5	31.7	37.8
2000[4]	7.9	*	*	*	*	*	*	*	3.0	6.4	16.0	26.7	32.8	41.9
1999[4]	7.4	*	*	*	*	*	*	*	2.3	7.1	15.6	27.7	31.0	40.2
Female														
2001	1.5	*	*	*	*	*	*	*	*	*	3.6	4.5	6.0	5.6
2000[4]	1.4	*	*	*	*	*	*	*	*	2.1	3.8	4.1	5.0	5.4
1999[4]	1.5	*	*	*	*	*	*	*	*	2.1	3.6	4.5	5.3	4.6
Non-Hispanic[5]														
Both sexes														
2001	4.2	*	*	*	*	*	*	0.2	1.0	3.2	5.8	9.6	10.6	10.3
2000[4]	4.2	*	*	*	*	*	*	0.3	1.0	3.1	6.2	9.4	10.0	10.8
1999[4]	4.2	*	*	*	*	*	*	0.2	1.0	3.1	6.0	9.4	10.2	10.8
Male														
2001	6.2	*	*	*	*	*	*	*	1.2	4.0	7.7	14.0	15.7	16.3
2000[4]	6.2	*	*	*	*	*	*	0.4	1.2	4.0	8.6	14.0	15.5	16.1
1999[4]	6.4	*	*	*	*	*	*	0.2	1.2	3.9	8.6	14.3	16.0	17.0
Female														
2001	2.4	*	*	*	*	*	*	*	0.7	2.4	3.9	5.3	5.8	4.6
2000[4]	2.3	*	*	*	*	*	*	*	0.7	2.3	3.9	4.8	4.8	5.8
1999[4]	2.2	*	*	*	*	*	*	*	0.8	2.4	3.5	4.6	4.7	4.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alcoholic liver disease (K70)						
All origins[3]						
Male						
2001	19.0	17.9	15.2	12.1	7.0	5.5
2000[4]	20.3	18.7	15.7	13.8	7.7	5.3
1999[4]	20.0	19.7	16.2	12.1	9.7	6.1
Female						
2001	5.6	5.3	4.0	3.3	2.1	1.0
2000[4]	6.2	5.1	3.9	2.6	2.4	1.0
1999[4]	5.9	4.6	4.0	3.9	2.7	1.5
Hispanic						
Both sexes						
2001	22.2	22.2	19.8	19.3	12.9	*
2000[4]	24.3	25.5	19.7	19.0	*	*
1999[4]	22.1	21.4	19.5	16.5	*	*
Male						
2001	40.5	43.9	39.3	34.1	*	*
2000[4]	44.0	51.8	39.9	41.3	*	*
1999[4]	39.6	39.8	39.2	29.6	*	*
Female						
2001	6.3	*	*	*	*	*
2000[4]	7.2	*	*	*	*	*
1999[4]	6.9	6.6	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	11.1	10.3	8.2	6.3	3.6	2.2
2000[4]	12.0	10.4	8.5	6.6	4.2	2.2
1999[4]	11.9	10.9	8.8	6.8	5.1	2.5
Male						
2001	17.3	15.9	13.6	11.0	6.3	5.1
2000[4]	18.4	16.4	14.3	12.4	7.3	5.0
1999[4]	18.5	18.3	14.8	11.2	9.2	5.5
Female						
2001	5.5	5.4	4.0	3.0	1.9	1.0
2000[4]	6.1	5.2	3.9	2.6	2.4	1.0
1999[4]	5.8	4.5	4.0	3.8	2.6	1.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Alcoholic liver disease (K70)														
Non-Hispanic White														
Both sexes														
2001	4.4	*	*	*	*	*	*	*	1.0	3.2	5.7	9.7	10.1	9.9
2000[4]	4.3	*	*	*	*	*	*	0.2	0.9	3.0	6.0	9.2	9.7	10.4
1999[4]	4.3	*	*	*	*	*	*	0.2	0.9	3.0	5.7	9.0	10.0	10.3
Male														
2001	6.4	*	*	*	*	*	*	*	1.2	4.0	7.6	14.3	15.1	15.6
2000[4]	6.5	*	*	*	*	*	*	0.4	1.2	3.8	8.3	14.1	15.0	15.8
1999[4]	6.5	*	*	*	*	*	*	*	1.1	3.8	8.1	13.9	15.7	16.2
Female														
2001	2.4	*	*	*	*	*	*	*	0.7	2.5	3.9	5.2	5.3	4.5
2000[4]	2.2	*	*	*	*	*	*	*	0.6	2.2	3.7	4.4	4.4	5.3
1999[4]	2.2	*	*	*	*	*	*	*	0.7	2.3	3.2	4.2	4.4	4.6
Non-Hispanic Black														
Both sexes														
2001	3.7	*	*	*	*	*	*	*	*	2.6	6.1	9.1	14.9	14.4
2000[4]	3.9	*	*	*	*	*	*	*	1.0	3.2	7.1	11.3	13.6	14.0
1999[4]	4.2	*	*	*	*	*	*	*	0.9	3.0	7.4	12.8	12.9	15.0
Male														
2001	5.5	*	*	*	*	*	*	*	*	3.8	8.5	13.3	22.5	24.0
2000[4]	5.5	*	*	*	*	*	*	*	*	4.4	9.9	15.5	21.0	20.5
1999[4]	6.1	*	*	*	*	*	*	*	*	3.7	10.3	19.2	20.4	25.3
Female														
2001	2.2	*	*	*	*	*	*	*	*	1.6	4.1	5.6	8.4	6.5
2000[4]	2.5	*	*	*	*	*	*	*	*	2.2	4.6	7.7	7.3	8.7
1999[4]	2.4	*	*	*	*	*	*	*	*	2.3	4.8	7.4	6.6	6.7
Other chronic liver disease and cirrhosis (K73-K74)														
All origins[3]														
Both sexes														
2001	5.2	*	*	*	*	*	*	0.2	0.4	1.6	3.6	6.9	8.8	10.0
2000[4]	5.1	*	*	*	*	*	0.1	0.2	0.5	1.5	3.6	6.8	7.9	10.2
1999[4]	5.1	*	*	*	*	*	*	0.2	0.6	1.5	3.5	6.4	7.7	9.9
Male														
2001	6.1	*	*	*	*	*	*	0.2	0.6	2.0	4.8	10.0	13.1	13.5
2000[4]	6.0	*	*	*	*	*	*	*	0.6	2.0	5.0	10.2	11.6	13.7
1999[4]	6.0	*	*	*	*	*	*	*	0.7	2.0	5.2	9.8	11.6	13.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alcoholic liver disease (K70)						
Non-Hispanic White						
Both sexes						
2001	11.3	10.1	8.4	6.4	3.6	2.1
2000[4]	11.9	10.4	8.6	6.7	4.1	2.2
1999[4]	11.5	10.7	8.8	6.7	5.0	2.4
Male						
2001	17.4	15.4	13.6	11.2	6.4	4.8
2000[4]	18.4	16.5	14.3	12.6	7.3	5.1
1999[4]	18.3	17.8	14.7	10.9	9.3	5.1
Female						
2001	5.6	5.4	4.1	3.0	1.9	1.0
2000[4]	5.8	5.1	4.0	2.5	2.3	1.0
1999[4]	5.3	4.5	4.0	3.7	2.5	1.3
Non-Hispanic Black						
Both sexes						
2001	11.6	13.1	8.0	5.7	*	*
2000[4]	12.8	12.0	8.6	6.3	*	*
1999[4]	15.9	13.2	10.2	8.2	*	*
Male						
2001	18.8	23.4	15.5	*	*	*
2000[4]	18.5	19.5	17.1	11.6	*	*
1999[4]	23.7	24.7	18.3	14.5	*	*
Female						
2001	5.9	5.4	*	*	*	*
2000[4]	8.4	6.5	*	*	*	*
1999[4]	9.7	4.7	4.8	*	*	*
Other chronic liver disease and cirrhosis (K73-K74)						
All origins[3]						
Both sexes						
2001	12.8	17.7	22.2	24.4	24.7	19.9
2000[4]	13.3	17.5	21.7	24.7	25.2	20.9
1999[4]	13.9	18.1	22.3	25.4	25.6	20.4
Male						
2001	16.3	21.8	25.6	25.5	28.4	26.5
2000[4]	17.4	20.6	25.0	27.0	26.8	29.0
1999[4]	17.3	22.0	25.3	28.0	29.0	27.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other chronic liver disease and cirrhosis (K73-K74)														
All origins[3]														
Female														
2001	4.4	*	*	*	*	*	*	*	0.3	1.3	2.5	4.0	4.7	6.7
2000[4]	4.3	*	*	*	*	*	*	*	0.4	1.0	2.2	3.6	4.3	6.9
1999[4]	4.3	*	*	*	*	*	*	*	0.4	1.0	1.8	3.2	4.0	6.2
Hispanic														
Both sexes														
2001	4.3	*	*	*	*	*	*	*	*	2.1	5.1	9.0	13.3	15.0
2000[4]	4.3	*	*	*	*	*	*	*	0.7	1.7	4.3	9.4	11.8	16.9
1999[4]	4.2	*	*	*	*	*	*	*	0.9	2.3	5.2	9.1	10.5	14.1
Male														
2001	5.0	*	*	*	*	*	*	*	*	2.8	7.9	14.3	20.8	21.2
2000[4]	4.8	*	*	*	*	*	*	*	*	2.7	6.5	14.4	18.7	22.8
1999[4]	4.9	*	*	*	*	*	*	*	*	3.7	8.2	14.6	15.8	19.2
Female														
2001	3.4	*	*	*	*	*	*	*	*	1.5	2.1	3.5	6.2	9.4
2000[4]	3.7	*	*	*	*	*	*	*	*	*	2.0	4.4	5.3	11.5
1999[4]	3.5	*	*	*	*	*	*	*	*	*	2.0	3.7	5.5	9.6
Non-Hispanic[5]														
Both sexes														
2001	5.3	*	*	*	*	*	*	0.2	0.4	1.5	3.4	6.7	8.4	9.6
2000[4]	5.2	*	*	*	*	*	*	0.2	0.5	1.5	3.5	6.6	7.5	9.6
1999[4]	5.2	*	*	*	*	*	*	0.2	0.5	1.4	3.3	6.1	7.4	9.5
Male														
2001	6.2	*	*	*	*	*	*	0.3	0.6	1.9	4.4	9.5	12.4	12.9
2000[4]	6.1	*	*	*	*	*	*	*	0.5	1.9	4.7	9.7	11.0	13.0
1999[4]	6.1	*	*	*	*	*	*	*	0.6	1.7	4.8	9.3	11.1	13.3
Female														
2001	4.5	*	*	*	*	*	*	*	0.3	1.2	2.5	4.0	4.6	6.5
2000[4]	4.4	*	*	*	*	*	*	*	0.4	1.0	2.2	3.5	4.3	6.5
1999[4]	4.4	*	*	*	*	*	*	*	0.4	1.0	1.8	3.1	3.9	5.9
Non-Hispanic White														
Both sexes														
2001	5.8	*	*	*	*	*	*	*	0.5	1.6	3.3	6.3	8.0	9.2
2000[4]	5.7	*	*	*	*	*	*	*	0.4	1.4	3.4	6.5	7.0	9.5
1999[4]	5.6	*	*	*	*	*	*	*	0.5	1.4	3.2	5.6	7.2	9.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)						
All origins[3]						
Female						
2001	9.6	14.2	19.5	23.6	22.6	17.1
2000[4]	9.5	14.8	19.2	23.2	24.2	17.5
1999[4]	10.8	14.8	19.9	23.5	23.6	17.7
Hispanic						
Both sexes						
2001	21.0	28.9	38.8	42.8	44.7	30.6
2000[4]	19.5	30.0	42.1	46.8	51.2	44.5
1999[4]	21.1	32.1	40.2	43.9	47.0	40.0
Male						
2001	30.1	34.0	36.6	47.7	41.6	*
2000[4]	23.9	35.1	38.9	48.0	42.7	50.4
1999[4]	27.1	36.7	43.3	42.9	45.6	45.3
Female						
2001	13.2	24.7	40.5	39.2	46.6	33.5
2000[4]	15.6	26.0	44.6	46.0	56.4	41.5
1999[4]	15.9	28.4	37.9	44.6	47.8	37.4
Non-Hispanic[5]						
Both sexes						
2001	12.1	16.9	21.1	23.4	23.8	19.4
2000[4]	12.7	16.5	20.5	23.7	24.1	19.9
1999[4]	13.3	17.2	21.3	24.5	24.8	19.6
Male						
2001	15.1	20.8	25.0	24.4	27.7	26.4
2000[4]	16.9	19.6	24.1	26.0	26.1	28.0
1999[4]	16.5	21.1	24.4	27.3	28.4	26.2
Female						
2001	9.3	13.5	18.0	22.7	21.5	16.5
2000[4]	9.0	14.0	17.6	22.1	22.9	16.7
1999[4]	10.5	13.8	18.9	22.5	22.6	17.0
Non-Hispanic White						
Both sexes						
2001	12.3	17.5	21.9	24.5	25.0	20.4
2000[4]	12.8	16.7	21.1	25.0	25.0	20.6
1999[4]	13.2	17.6	22.1	25.5	25.9	20.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other chronic liver disease and cirrhosis (K73-K74)														
Non-Hispanic White														
Male														
2001	6.6	*	*	*	*	*	*	*	0.6	1.8	4.2	8.9	11.6	12.1
2000[4]	6.6	*	*	*	*	*	*	*	0.4	1.9	4.8	9.7	10.1	12.7
1999[4]	6.6	*	*	*	*	*	*	*	0.6	1.7	4.6	8.5	10.7	12.8
Female														
2001	5.0	*	*	*	*	*	*	*	0.4	1.3	2.4	3.8	4.4	6.3
2000[4]	4.8	*	*	*	*	*	*	*	0.4	1.0	2.1	3.3	4.0	6.5
1999[4]	4.8	*	*	*	*	*	*	*	0.4	1.0	1.7	2.8	3.7	5.7
Non-Hispanic Black														
Both sexes														
2001	3.9	*	*	*	*	*	*	*	*	1.5	4.8	10.3	12.3	13.9
2000[4]	3.8	*	*	*	*	*	*	*	*	1.8	4.1	8.5	12.7	12.0
1999[4]	3.9	*	*	*	*	*	*	*	*	1.6	4.8	10.0	10.2	13.4
Male														
2001	5.3	*	*	*	*	*	*	*	*	2.0	6.6	15.6	20.2	20.8
2000[4]	4.8	*	*	*	*	*	*	*	*	2.3	4.9	11.8	20.2	18.5
1999[4]	4.9	*	*	*	*	*	*	*	*	1.9	7.1	15.6	16.6	19.0
Female														
2001	2.7	*	*	*	*	*	*	*	*	*	3.2	5.7	5.6	8.3
2000[4]	2.9	*	*	*	*	*	*	*	*	*	3.3	5.6	6.2	6.7
1999[4]	2.9	*	*	*	*	*	*	*	*	*	2.7	5.2	4.8	8.8
Cholelithiasis and other disorders of gallbladder (K80-K82)														
All origins[3]														
Both sexes														
2001	1.0	*	*	*	*	*	*	*	*	0.1	0.1	0.3	0.4	0.7
2000[4]	1.0	*	*	*	*	*	*	*	*	0.1	0.1	0.2	0.4	0.6
1999[4]	1.0	*	*	*	*	*	*	*	*	0.1	0.2	0.3	0.4	0.7
Male														
2001	1.0	*	*	*	*	*	*	*	*	*	*	0.4	0.4	0.9
2000[4]	0.9	*	*	*	*	*	*	*	*	*	*	0.2	0.3	0.7
1999[4]	0.9	*	*	*	*	*	*	*	*	*	*	0.3	0.4	0.9
Female														
2001	1.1	*	*	*	*	*	*	*	*	*	*	0.3	0.3	0.5
2000[4]	1.1	*	*	*	*	*	*	*	*	*	*	0.2	0.4	0.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)						
Non-Hispanic White						
Male						
2001	14.9	21.4	25.9	25.4	29.3	28.1
2000[4]	16.9	20.0	24.8	27.1	27.4	29.0
1999[4]	16.4	21.6	25.2	28.4	29.8	27.5
Female						
2001	9.8	14.2	18.5	23.8	22.4	17.2
2000[4]	9.0	13.8	18.1	23.5	23.6	17.2
1999[4]	10.3	14.0	19.5	23.5	23.5	17.3
Non-Hispanic Black						
Both sexes						
2001	12.2	14.1	15.0	14.9	11.6	7.8
2000[4]	13.5	15.8	16.4	12.0	13.5	12.4
1999[4]	15.5	15.0	14.9	15.1	13.5	12.3
Male						
2001	19.6	20.3	18.9	18.0	*	*
2000[4]	19.3	18.9	20.8	16.4	*	*
1999[4]	19.6	19.4	18.7	17.9	*	*
Female						
2001	6.4	9.4	12.4	13.0	12.9	*
2000[4]	8.9	13.6	13.4	9.3	14.3	11.3
1999[4]	12.3	11.8	12.4	13.4	12.3	12.9
Cholelithiasis and other disorders of gallbladder (K80-K82)						
All origins[3]						
Both sexes						
2001	1.3	2.0	3.8	5.8	10.0	23.7
2000[4]	1.0	1.9	3.0	5.7	10.3	24.8
1999[4]	1.0	1.8	3.4	5.7	10.9	23.9
Male						
2001	1.5	2.2	4.4	7.6	12.1	26.9
2000[4]	1.2	2.3	3.6	7.4	12.9	27.6
1999[4]	1.3	2.0	4.2	6.9	13.3	27.7
Female						
2001	1.1	1.9	3.3	4.6	8.7	22.4
2000[4]	0.8	1.6	2.5	4.5	8.7	23.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Cholelithiasis and other disorders of gallbladder (K80-K82)														
All origins[3]														
Female														
1999[4]	1.1	*	*	*	*	*	*	*	*	*	*	0.3	0.5	0.6
Hispanic														
Both sexes														
2001	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	1.1	*	*	*	*	*	*	*	*	*	0.1	0.3	0.4	0.6
2000[4]	1.1	*	*	*	*	*	*	*	*	0.1	0.1	0.2	0.3	0.6
1999[4]	1.1	*	*	*	*	*	*	*	*	*	0.1	0.3	0.4	0.7
Male														
2001	1.0	*	*	*	*	*	*	*	*	*	*	0.4	0.4	0.8
2000[4]	1.0	*	*	*	*	*	*	*	*	*	*	*	0.3	0.7
1999[4]	1.0	*	*	*	*	*	*	*	*	*	*	0.3	0.3	0.9
Female														
2001	1.2	*	*	*	*	*	*	*	*	*	*	0.3	0.3	0.4
2000[4]	1.2	*	*	*	*	*	*	*	*	*	*	*	0.4	0.5
1999[4]	1.2	*	*	*	*	*	*	*	*	*	*	0.2	0.5	0.6
Non-Hispanic White														
Both sexes														
2001	1.2	*	*	*	*	*	*	*	*	*	*	0.3	0.3	0.5
2000[4]	1.2	*	*	*	*	*	*	*	*	*	*	0.1	0.3	0.5
1999[4]	1.2	*	*	*	*	*	*	*	*	*	*	0.2	0.4	0.6
Male														
2001	1.1	*	*	*	*	*	*	*	*	*	*	0.3	0.4	0.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cholelithiasis and other disorders of gallbladder (K80-K82)						
All origins[3]						
Female						
1999[4]	0.7	1.6	2.8	4.8	9.5	22.4
Hispanic						
Both sexes						
2001	*	*	6.7	6.2	12.4	31.3
2000[4]	*	3.7	*	8.0	13.9	29.2
1999[4]	*	*	4.4	9.7	15.9	32.3
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	30.7
2000[4]	*	*	*	*	*	31.7
1999[4]	*	*	*	*	*	32.2
Non-Hispanic[5]						
Both sexes						
2001	1.3	2.0	3.6	5.8	9.8	23.4
2000[4]	1.0	1.8	2.9	5.5	10.1	24.6
1999[4]	1.0	1.8	3.3	5.5	10.7	23.6
Male						
2001	1.5	2.1	4.2	7.5	11.9	26.6
2000[4]	1.1	2.1	3.5	7.0	12.6	27.7
1999[4]	1.2	2.0	4.2	6.6	13.1	27.5
Female						
2001	1.2	1.8	3.1	4.6	8.6	22.1
2000[4]	0.9	1.5	2.5	4.5	8.6	23.4
1999[4]	0.8	1.6	2.7	4.7	9.2	22.0
Non-Hispanic White						
Both sexes						
2001	1.2	1.7	3.4	5.7	9.6	23.6
2000[4]	0.9	1.7	2.6	5.5	9.9	25.3
1999[4]	0.8	1.5	3.2	5.5	10.7	24.0
Male						
2001	1.4	1.8	4.1	7.5	11.8	26.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Cholelithiasis and other disorders of gallbladder (K80-K82)														
Non-Hispanic White														
Male														
2000[4]	1.0	*	*	*	*	*	*	*	*	*	*	*	*	0.7
1999[4]	1.1	*	*	*	*	*	*	*	*	*	*	*	0.3	0.7
Female														
2001	1.3	*	*	*	*	*	*	*	*	*	*	*	*	0.4
2000[4]	1.3	*	*	*	*	*	*	*	*	*	*	*	0.3	0.4
1999[4]	1.3	*	*	*	*	*	*	*	*	*	*	*	0.4	0.5
Non-Hispanic Black														
Both sexes														
2001	0.9	*	*	*	*	*	*	*	*	*	*	*	*	1.5
2000[4]	0.8	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.7	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.8	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	1.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.8	*	*	*	*	*	*	*	*	*	*	*	*	*
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)														
All origins[3]														
Both sexes														
2001	13.9	3.3	*	*	*	0.1	0.4	0.4	0.7	1.3	2.1	3.6	5.8	9.9
2000[4]	13.2	4.3	*	*	*	0.2	0.3	0.4	0.8	1.2	2.1	3.4	5.5	9.4
1999[4]	12.7	4.4	*	*	*	0.1	0.3	0.5	0.8	1.2	2.0	3.1	5.2	9.0
Male														
2001	13.5	3.7	*	*	*	*	0.3	0.5	0.6	1.4	2.3	4.4	6.7	10.9
2000[4]	12.9	5.3	*	*	*	*	0.3	0.4	0.8	1.3	2.5	4.4	6.4	10.7
1999[4]	12.4	4.6	*	*	*	*	0.2	0.4	0.8	1.3	2.3	3.8	5.8	9.9
Female														
2001	14.2	2.9	*	*	*	*	0.4	0.4	0.7	1.3	1.9	2.9	4.9	8.9
2000[4]	13.6	3.2	*	*	*	*	0.3	0.4	0.8	1.1	1.6	2.5	4.6	8.2
1999[4]	13.0	4.1	*	*	*	*	0.3	0.6	0.7	1.0	1.7	2.4	4.5	8.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cholelithiasis and other disorders of gallbladder (K80-K82)						
Non-Hispanic White						
Male						
2000[4]	1.0	2.1	3.2	7.1	12.8	28.9
1999[4]	1.1	1.8	4.1	6.7	13.6	27.8
Female						
2001	1.1	1.6	2.9	4.4	8.3	22.4
2000[4]	0.8	1.3	2.2	4.3	8.2	23.9
1999[4]	0.6	1.3	2.5	4.7	9.0	22.5
Non-Hispanic Black						
Both sexes						
2001	2.0	3.5	6.3	6.2	11.6	20.0
2000[4]	*	2.8	4.9	6.5	11.2	18.1
1999[4]	2.0	3.3	4.1	4.7	11.4	15.5
Male						
2001	*	*	*	*	*	27.7
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	6.4	7.1	12.9	17.1
2000[4]	*	*	4.8	6.4	11.7	19.6
1999[4]	*	*	*	*	12.3	15.5
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)						
All origins[3]						
Both sexes						
2001	16.9	31.7	49.3	82.2	135.8	287.7
2000[4]	16.9	29.5	47.2	79.5	132.8	277.8
1999[4]	15.8	28.2	46.7	77.7	127.4	268.9
Male						
2001	18.2	36.5	57.0	100.5	176.8	393.7
2000[4]	18.6	32.8	54.3	99.8	173.0	380.7
1999[4]	17.8	31.0	54.9	99.2	168.9	371.1
Female						
2001	15.7	27.6	43.2	69.4	111.2	243.3
2000[4]	15.5	26.6	41.6	65.3	109.0	235.9
1999[4]	14.0	25.8	40.3	62.8	103.1	227.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)														
Hispanic														
Both sexes														
2001	5.0	*	*	*	*	*	*	*	*	0.8	2.0	2.7	5.5	11.1
2000[4]	4.7	3.2	*	*	*	*	*	*	0.6	*	1.5	2.5	4.7	9.1
1999[4]	4.7	2.6	*	*	*	*	*	*	*	0.8	1.8	3.0	4.4	10.5
Male														
2001	4.7	*	*	*	*	*	*	*	*	*	2.5	2.8	5.1	12.6
2000[4]	4.4	*	*	*	*	*	*	*	*	*	2.3	3.6	5.6	9.9
1999[4]	4.7	*	*	*	*	*	*	*	*	*	*	3.6	5.4	13.4
Female														
2001	5.4	*	*	*	*	*	*	*	*	*	*	2.5	5.8	9.8
2000[4]	4.9	*	*	*	*	*	*	*	*	*	*	*	3.9	8.3
1999[4]	4.7	*	*	*	*	*	*	*	*	*	1.9	2.4	3.5	7.9
Non-Hispanic[5]														
Both sexes														
2001	15.1	3.5	*	*	*	*	0.4	0.5	0.7	1.4	2.1	3.7	5.8	9.8
2000[4]	14.4	4.4	*	*	*	0.1	0.3	0.5	0.8	1.3	2.1	3.5	5.5	9.4
1999[4]	13.8	4.8	*	*	*	0.1	0.3	0.5	0.8	1.2	2.0	3.1	5.2	8.8
Male														
2001	14.8	3.7	*	*	*	*	0.3	0.5	0.7	1.5	2.2	4.5	6.8	10.8
2000[4]	14.1	5.4	*	*	*	*	0.3	0.5	0.9	1.5	2.5	4.5	6.4	10.8
1999[4]	13.5	4.8	*	*	*	*	0.3	0.5	0.9	1.4	2.3	3.8	5.8	9.6
Female														
2001	15.4	3.3	*	*	*	*	0.4	0.4	0.8	1.3	1.9	2.9	4.8	8.8
2000[4]	14.7	3.4	*	*	*	*	0.4	0.4	0.7	1.2	1.7	2.6	4.7	8.2
1999[4]	14.1	4.8	*	*	*	*	0.3	0.6	0.8	1.0	1.7	2.4	4.6	8.1
Non-Hispanic White														
Both sexes														
2001	14.9	2.4	*	*	*	*	0.3	0.3	0.5	0.9	1.3	2.3	3.9	6.8
2000[4]	14.2	2.7	*	*	*	*	0.3	0.4	0.5	0.8	1.4	2.2	3.6	6.6
1999[4]	13.4	3.5	*	*	*	*	0.2	0.3	0.5	0.8	1.2	1.8	3.4	6.2
Male														
2001	14.8	2.5	*	*	*	*	*	*	0.4	1.2	1.5	2.7	4.8	7.8
2000[4]	14.1	3.5	*	*	*	*	*	0.4	0.6	0.9	1.7	2.8	4.2	7.7
1999[4]	13.3	3.9	*	*	*	*	*	0.3	0.6	0.9	1.4	2.2	3.7	7.2
Female														
2001	15.0	2.3	*	*	*	*	0.3	*	0.6	0.7	1.2	1.8	3.1	5.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)						
Hispanic						
Both sexes						
2001	15.1	34.6	49.9	83.8	113.2	222.0
2000[4]	19.3	32.5	43.8	71.9	108.1	229.6
1999[4]	18.5	29.7	43.1	77.1	109.8	232.2
Male						
2001	18.7	35.4	53.5	102.3	131.2	281.9
2000[4]	20.1	33.6	50.6	85.6	132.5	292.2
1999[4]	23.5	32.5	47.9	103.8	124.2	336.4
Female						
2001	12.0	33.9	47.1	70.7	102.1	192.7
2000[4]	18.6	31.7	38.7	62.2	93.2	198.8
1999[4]	14.1	27.5	39.5	58.4	101.3	182.0
Non-Hispanic[5]						
Both sexes						
2001	17.0	31.5	49.1	82.0	136.3	289.6
2000[4]	16.6	29.2	47.2	79.6	133.4	279.0
1999[4]	15.5	28.0	46.8	77.6	127.7	269.2
Male						
2001	18.1	36.5	57.0	100.3	178.2	397.6
2000[4]	18.3	32.7	54.3	100.3	174.1	384.0
1999[4]	17.3	30.8	55.1	98.8	170.2	370.9
Female						
2001	16.0	27.1	42.8	69.1	111.2	244.6
2000[4]	15.1	26.2	41.6	65.2	109.4	236.6
1999[4]	14.0	25.7	40.3	62.9	102.8	228.4
Non-Hispanic White						
Both sexes						
2001	12.5	25.0	42.4	74.4	128.0	279.7
2000[4]	12.5	23.0	40.2	72.7	125.6	269.4
1999[4]	11.6	22.4	40.1	70.5	118.6	255.3
Male						
2001	13.9	29.9	50.9	94.0	170.0	392.6
2000[4]	14.3	27.3	48.4	93.5	167.6	378.9
1999[4]	13.2	25.5	48.8	90.8	160.8	359.6
Female						
2001	11.2	20.6	35.5	60.5	102.6	232.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)														
Non-Hispanic White														
Female														
2000[4]	14.2	1.8	*	*	*	*	*	*	0.4	0.7	1.2	1.7	3.1	5.6
1999[4]	13.5	3.1	*	*	*	*	*	0.4	0.5	0.6	1.0	1.5	3.0	5.3
Non-Hispanic Black														
Both sexes														
2001	20.2	8.5	*	*	*	*	0.8	1.4	2.2	4.1	6.9	12.9	20.3	33.3
2000[4]	19.5	12.0	*	*	*	*	*	1.1	2.7	4.5	6.5	13.1	20.1	33.0
1999[4]	19.2	10.2	*	*	*	*	0.8	1.5	2.4	4.2	7.2	12.1	20.1	29.6
Male														
2001	18.6	8.9	*	*	*	*	*	1.8	2.6	3.4	7.2	17.0	23.6	38.6
2000[4]	18.3	13.6	*	*	*	*	*	*	3.0	5.3	8.2	18.0	24.3	38.3
1999[4]	18.1	8.7	*	*	*	*	*	*	2.4	5.0	8.5	15.9	24.1	29.8
Female														
2001	21.7	8.1	*	*	*	*	*	*	1.9	4.6	6.7	9.3	17.5	28.9
2000[4]	20.7	10.2	*	*	*	*	*	*	2.5	3.9	5.0	8.9	16.4	28.7
1999[4]	20.2	11.8	*	*	*	*	*	1.7	2.4	3.4	6.1	8.9	16.8	29.4
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01, N04)														
All origins[3]														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)						
Non-Hispanic White						
Female						
2000[4]	10.9	19.3	33.6	58.2	100.6	225.2
1999[4]	10.3	19.8	33.1	56.2	93.7	213.8
Non-Hispanic Black						
Both sexes						
2001	54.2	90.4	121.3	174.0	250.8	432.9
2000[4]	51.3	84.6	121.2	167.2	245.1	410.2
1999[4]	47.9	78.9	118.2	165.6	246.3	436.8
Male						
2001	56.2	103.4	132.6	190.6	312.3	517.4
2000[4]	54.9	87.9	128.5	201.4	292.7	496.7
1999[4]	55.5	82.3	131.9	210.3	325.9	532.4
Female						
2001	52.6	80.8	113.8	164.1	219.5	401.7
2000[4]	48.5	82.2	116.4	146.6	221.2	378.1
1999[4]	42.0	76.4	109.1	138.6	206.8	401.1
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01, N04)						
All origins[3]						
Both sexes						
2001	*	*	*	0.4	0.4	0.9
2000[4]	*	*	*	*	0.7	1.2
1999[4]	*	*	0.3	0.5	0.5	0.8
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	1.8
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	0.7
2000[4]	*	*	*	*	0.7	0.9
1999[4]	*	*	*	0.5	*	0.7
Hispanic						
Both sexes						
2001	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01, N04)														
Hispanic														
Both sexes														
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01, N04)						
Hispanic						
Both sexes						
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	0.4	*	0.8
2000[4]	*	*	*	*	0.7	1.2
1999[4]	*	*	0.3	0.5	0.5	0.8
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	1.8
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	0.7
2000[4]	*	*	*	*	0.7	0.9
1999[4]	*	*	*	0.5	*	0.7
Non-Hispanic White						
Both sexes						
2001	*	*	*	0.4	*	0.8
2000[4]	*	*	*	*	0.8	1.2
1999[4]	*	*	0.3	0.5	0.5	0.8
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01, N04)														
Non-Hispanic White														
Female														
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Chronic glomerulonephritis, nephritis and nephropathy, not specified as acute or chronic, and renal sclerosis, unspecified (N02-N03,N05-N07, N26)														
All origins[3]														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	0.1	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	0.1	*	0.1	0.2
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	0.2	0.2
Male														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01, N04)						
Non-Hispanic White						
Female						
2000[4]	*	*	*	*	0.7	1.0
1999[4]	*	*	*	*	*	*
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Chronic glomerulonephritis, nephritis and nephropathy, not specified as acute or chronic, and renal sclerosis, unspecified (N02-N03,N05-N07, N26)						
All origins[3]						
Both sexes						
2001	0.2	0.5	0.7	0.9	1.5	2.8
2000[4]	0.3	0.5	0.7	1.1	1.9	2.8
1999[4]	0.3	0.4	0.7	1.4	2.0	3.2
Male						
2001	*	0.5	1.0	1.3	2.3	3.7
2000[4]	*	0.5	0.8	1.5	2.1	4.5
1999[4]	*	0.5	0.6	1.9	2.7	5.0
Female						
2001	*	0.5	0.5	0.7	1.0	2.4
2000[4]	*	*	0.6	0.8	1.8	2.1
1999[4]	*	0.4	0.7	1.1	1.5	2.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Chronic glomerulonephritis, nephritis and nephropathy, not specified as acute or chronic, and renal sclerosis, unspecified (N02-N03,N05-N07, N26)														
Hispanic														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	0.1	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	0.2
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	0.2	0.2
Male														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	0.2
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic glomerulonephritis, nephritis and nephropathy, not specified as acute or chronic, and renal sclerosis, unspecified (N02-N03,N05-N07, N26)						
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	0.2	0.5	0.7	0.9	1.5	2.8
2000[4]	0.3	0.5	0.7	1.1	1.9	2.8
1999[4]	0.3	0.4	0.7	1.5	2.0	3.2
Male						
2001	*	0.5	1.1	1.3	2.3	3.7
2000[4]	*	0.6	0.8	1.5	2.1	4.2
1999[4]	*	*	0.7	2.0	2.8	5.1
Female						
2001	*	0.5	0.5	0.7	1.1	2.4
2000[4]	*	*	0.6	0.8	1.8	2.2
1999[4]	*	*	0.7	1.1	1.5	2.4
Non-Hispanic White						
Both sexes						
2001	*	0.4	0.7	0.9	1.5	2.8
2000[4]	*	0.4	0.7	1.0	1.8	2.4
1999[4]	0.2	0.3	0.6	1.4	1.9	2.9
Male						
2001	*	*	1.1	1.2	2.3	4.0
2000[4]	*	*	0.8	1.4	2.2	3.7
1999[4]	*	*	0.6	2.0	2.7	4.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Chronic glomerulonephritis, nephritis and nephropathy, not specified as acute or chronic, and renal sclerosis, unspecified (N02-N03,N05-N07, N26)														
Non-Hispanic White														
Female														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Renal failure (N17-N19)														
All origins[3]														
Both sexes														
2001	13.6	3.1	*	*	*	*	0.3	0.4	0.6	1.2	2.0	3.5	5.7	9.7
2000[4]	13.0	4.0	*	*	*	0.1	0.3	0.4	0.7	1.2	2.0	3.3	5.3	9.2
1999[4]	12.4	4.3	*	*	*	0.1	0.2	0.4	0.7	1.1	1.9	3.0	4.9	8.7
Male														
2001	13.2	3.5	*	*	*	*	0.3	0.4	0.6	1.3	2.2	4.2	6.5	10.7
2000[4]	12.6	5.1	*	*	*	*	0.2	0.4	0.7	1.3	2.4	4.2	6.1	10.4
1999[4]	12.1	4.5	*	*	*	*	*	0.3	0.7	1.3	2.2	3.6	5.6	9.7
Female														
2001	14.0	2.7	*	*	*	*	0.4	0.4	0.7	1.2	1.8	2.8	4.8	8.8
2000[4]	13.3	2.8	*	*	*	*	0.3	0.3	0.7	1.0	1.6	2.4	4.5	8.0
1999[4]	12.7	4.1	*	*	*	*	0.3	0.5	0.7	0.9	1.6	2.3	4.3	7.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic glomerulonephritis, nephritis and nephropathy, not specified as acute or chronic, and renal sclerosis, unspecified (N02-N03,N05-N07, N26)						
Non-Hispanic White						
Female						
2001	*	*	*	0.7	1.0	2.4
2000[4]	*	*	0.6	0.7	1.6	1.9
1999[4]	*	*	0.7	1.0	1.4	2.2
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Renal failure (N17-N19)						
All origins[3]						
Both sexes						
2001	16.6	31.1	48.4	80.9	133.8	284.0
2000[4]	16.6	28.9	46.2	78.1	130.0	273.7
1999[4]	15.3	27.7	45.7	75.7	124.8	264.8
Male						
2001	17.9	35.9	55.8	98.9	173.9	388.7
2000[4]	18.1	32.1	53.3	97.9	170.1	374.2
1999[4]	17.3	30.5	53.9	96.7	165.1	364.6
Female						
2001	15.5	27.0	42.5	68.3	109.8	240.1
2000[4]	15.2	26.1	40.6	64.4	106.4	232.8
1999[4]	13.5	25.3	39.3	61.1	101.2	224.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Renal failure (N17-N19)														
Hispanic														
Both sexes														
2001	4.9	*	*	*	*	*	*	*	*	*	1.9	2.4	5.3	10.8
2000[4]	4.5	2.9	*	*	*	*	*	*	*	*	1.4	2.4	4.4	8.9
1999[4]	4.5	*	*	*	*	*	*	*	*	0.8	1.7	2.8	4.4	10.2
Male														
2001	4.6	*	*	*	*	*	*	*	*	*	2.4	2.8	4.8	12.2
2000[4]	4.3	*	*	*	*	*	*	*	*	*	2.2	3.4	5.3	9.6
1999[4]	4.5	*	*	*	*	*	*	*	*	*	*	3.5	5.4	12.9
Female														
2001	5.2	*	*	*	*	*	*	*	*	*	*	2.1	5.7	9.6
2000[4]	4.8	*	*	*	*	*	*	*	*	*	*	*	3.6	8.1
1999[4]	4.5	*	*	*	*	*	*	*	*	*	1.9	*	3.5	7.7
Non-Hispanic[5]														
Both sexes														
2001	14.9	3.3	*	*	*	*	0.3	0.4	0.7	1.3	2.0	3.6	5.6	9.6
2000[4]	14.1	4.2	*	*	*	0.1	0.3	0.4	0.7	1.2	2.0	3.4	5.3	9.2
1999[4]	13.5	4.7	*	*	*	0.1	0.2	0.5	0.8	1.1	1.9	3.0	5.0	8.6
Male														
2001	14.6	3.5	*	*	*	*	0.3	0.5	0.7	1.4	2.2	4.4	6.6	10.6
2000[4]	13.8	5.2	*	*	*	*	*	0.4	0.8	1.4	2.4	4.3	6.1	10.5
1999[4]	13.2	4.6	*	*	*	*	*	0.4	0.8	1.3	2.2	3.6	5.5	9.4
Female														
2001	15.2	3.1	*	*	*	*	0.3	0.4	0.7	1.3	1.8	2.8	4.7	8.6
2000[4]	14.4	3.1	*	*	*	*	0.3	0.4	0.7	1.1	1.7	2.5	4.6	8.0
1999[4]	13.8	4.8	*	*	*	*	0.3	0.5	0.7	0.9	1.6	2.4	4.4	7.8
Non-Hispanic White														
Both sexes														
2001	14.6	2.2	*	*	*	*	0.2	0.3	0.4	0.9	1.3	2.2	3.8	6.7
2000[4]	13.9	2.5	*	*	*	*	0.2	0.3	0.4	0.8	1.3	2.1	3.6	6.4
1999[4]	13.1	3.5	*	*	*	*	*	0.3	0.5	0.7	1.2	1.7	3.2	6.0
Male														
2001	14.5	2.3	*	*	*	*	*	*	0.4	1.1	1.5	2.7	4.6	7.6
2000[4]	13.7	3.3	*	*	*	*	*	0.3	0.5	0.8	1.6	2.6	4.1	7.4
1999[4]	13.0	3.8	*	*	*	*	*	*	0.5	0.8	1.4	2.1	3.5	7.1
Female														
2001	14.7	2.1	*	*	*	*	*	*	0.5	0.7	1.1	1.8	3.0	5.8
2000[4]	14.0	*	*	*	*	*	*	*	0.4	0.7	1.1	1.6	3.0	5.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Renal failure (N17-N19)						
Hispanic						
Both sexes						
2001	14.6	33.9	48.7	82.7	111.7	217.6
2000[4]	18.9	32.0	43.0	70.7	105.3	225.6
1999[4]	17.8	28.2	42.2	75.5	107.5	228.0
Male						
2001	17.9	34.7	52.6	100.9	128.6	278.1
2000[4]	19.3	33.6	49.6	84.2	129.6	280.1
1999[4]	22.6	30.9	47.9	102.2	122.6	330.0
Female						
2001	11.8	33.3	45.7	69.7	101.3	188.0
2000[4]	18.6	30.8	37.9	61.2	90.5	198.8
1999[4]	13.6	25.9	37.9	56.7	98.5	178.8
Non-Hispanic[5]						
Both sexes						
2001	16.7	30.8	48.2	80.6	134.3	285.9
2000[4]	16.3	28.6	46.2	78.3	130.6	274.9
1999[4]	15.1	27.6	45.7	75.6	125.0	265.0
Male						
2001	17.8	35.9	55.8	98.7	175.3	392.7
2000[4]	17.9	31.9	53.2	98.4	171.3	377.7
1999[4]	16.9	30.3	54.0	96.3	166.3	364.4
Female						
2001	15.8	26.5	42.1	68.0	109.7	241.5
2000[4]	14.8	25.8	40.7	64.3	106.7	233.4
1999[4]	13.5	25.2	39.2	61.2	100.9	225.2
Non-Hispanic White						
Both sexes						
2001	12.3	24.4	41.4	73.1	126.1	276.1
2000[4]	12.2	22.6	39.2	71.5	122.8	265.7
1999[4]	11.3	22.0	39.1	68.5	116.2	251.6
Male						
2001	13.7	29.4	49.7	92.4	167.2	387.6
2000[4]	14.0	26.6	47.3	91.7	164.6	373.4
1999[4]	12.8	25.0	47.9	88.3	157.2	353.8
Female						
2001	11.1	20.0	34.7	59.4	101.3	229.7
2000[4]	10.6	19.0	32.7	57.3	98.0	222.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Renal failure (N17-N19)														
Non-Hispanic White														
Female														
1999[4]	13.2	3.1	*	*	*	*	*	*	0.5	0.6	1.0	1.4	2.9	5.1
Non-Hispanic Black														
Both sexes														
2001	20.0	8.5	*	*	*	*	0.8	1.3	2.2	3.9	6.8	12.6	19.8	33.0
2000[4]	19.2	11.4	*	*	*	*	*	1.0	2.6	4.2	6.3	12.7	19.2	32.6
1999[4]	18.8	10.2	*	*	*	*	*	1.5	2.3	3.9	6.9	12.0	19.4	29.0
Male														
2001	18.4	8.9	*	*	*	*	*	1.7	2.6	3.3	7.0	16.6	22.8	38.2
2000[4]	17.9	13.6	*	*	*	*	*	*	2.8	5.1	8.0	17.6	22.9	37.9
1999[4]	17.7	8.7	*	*	*	*	*	*	2.3	4.7	8.3	15.7	23.3	29.3
Female														
2001	21.4	8.1	*	*	*	*	*	*	1.8	4.5	6.6	9.2	17.2	28.8
2000[4]	20.3	9.2	*	*	*	*	*	*	2.4	3.5	4.9	8.5	16.0	28.3
1999[4]	19.8	11.8	*	*	*	*	*	1.7	2.3	3.3	5.8	8.8	16.1	28.7
Other disorders of kidney (N25,N27)														
All origins[3]														
Both sexes														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Renal failure (N17-N19)						
Non-Hispanic White						
Female						
1999[4]	10.0	19.4	32.1	54.6	92.0	210.8
Non-Hispanic Black						
Both sexes						
2001	53.6	88.9	120.5	172.2	248.1	428.9
2000[4]	50.6	83.2	120.1	164.9	241.7	404.2
1999[4]	46.9	77.7	116.8	162.8	241.9	428.7
Male						
2001	55.4	101.4	131.5	189.1	309.9	514.0
2000[4]	53.8	85.8	127.8	198.9	290.1	487.3
1999[4]	54.4	81.3	130.2	206.9	319.7	519.4
Female						
2001	52.2	79.6	113.1	162.1	216.6	397.4
2000[4]	48.1	81.2	115.0	144.3	217.3	373.3
1999[4]	41.0	75.0	108.0	136.3	203.3	394.9
Other disorders of kidney (N25,N27)						
All origins[3]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other disorders of kidney (N25,N27)														
Hispanic														
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other disorders of kidney (N25,N27)						
Hispanic						
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic White						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other disorders of kidney (N25,N27)														
Non-Hispanic Black														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Infections of kidney (N10-N12,N13.6, N15.1)														
All origins[3]														
Both sexes														
2001	0.3	*	*	*	*	*	*	*	*	*	*	0.1	0.1	0.2
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	0.1	0.1	0.2
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	0.1	0.2	0.2
Male														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	0.2	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	0.3
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	0.3	*
Hispanic														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other disorders of kidney (N25,N27)						
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Infections of kidney (N10-N12,N13.6, N15.1)						
All origins[3]						
Both sexes						
2001	0.2	0.4	0.6	1.7	2.4	6.0
2000[4]	0.3	0.4	0.7	1.5	2.7	7.5
1999[4]	0.3	0.5	0.9	1.5	2.7	7.5
Male						
2001	*	*	0.5	1.2	2.1	4.3
2000[4]	*	*	*	1.1	2.2	5.6
1999[4]	*	*	0.6	1.2	2.6	6.8
Female						
2001	*	*	0.6	2.1	2.6	6.7
2000[4]	0.4	0.5	1.1	1.8	3.0	8.3
1999[4]	*	0.6	1.1	1.7	2.8	7.8
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Infections of kidney (N10-N12,N13.6, N15.1)														
Hispanic														
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	0.1	0.2
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	0.1	0.2
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	0.2	0.2
Male														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	0.3	*
Non-Hispanic White														
Both sexes														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	0.2
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	0.2	0.2
Male														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Infections of kidney (N10-N12,N13.6, N15.1)						
Hispanic						
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	0.2	0.3	0.6	1.7	2.4	6.1
2000[4]	0.3	0.4	0.8	1.5	2.7	7.6
1999[4]	0.3	0.5	0.9	1.4	2.6	7.5
Male						
2001	*	*	*	1.2	2.2	4.3
2000[4]	*	*	*	1.2	2.1	5.8
1999[4]	*	*	0.6	1.2	2.4	6.9
Female						
2001	*	*	0.6	2.1	2.5	6.9
2000[4]	0.4	0.5	1.1	1.8	3.0	8.3
1999[4]	*	0.7	1.1	1.6	2.8	7.7
Non-Hispanic White						
Both sexes						
2001	*	0.3	0.6	1.8	2.2	6.2
2000[4]	0.3	0.3	0.7	1.4	2.5	7.6
1999[4]	*	0.5	0.8	1.4	2.7	7.7
Male						
2001	*	*	*	1.1	2.1	4.2
2000[4]	*	*	*	1.1	2.0	6.0
1999[4]	*	*	*	1.2	2.4	7.3
Female						
2001	*	*	0.7	2.2	2.3	7.1
2000[4]	*	0.5	1.0	1.6	2.8	8.2
1999[4]	*	0.6	1.1	1.6	2.9	7.9
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	7.9
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Infections of kidney (N10-N12,N13.6, N15.1)														
Non-Hispanic Black														
Male														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Hyperplasia of prostate (N40)														
All origins[3]														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Hispanic														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Infections of kidney (N10-N12,N13.6, N15.1)						
Non-Hispanic Black						
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	9.2
1999[4]	*	*	*	*	*	*
Hyperplasia of prostate (N40)						
All origins[3]						
Both sexes						
2001	*	*	0.4	0.8	1.6	5.5
2000[4]	*	*	0.3	0.9	1.9	5.2
1999[4]	*	*	0.4	0.8	2.1	5.0
Male						
2001	*	*	0.8	2.0	4.2	18.6
2000[4]	*	*	0.7	2.1	5.2	18.1
1999[4]	*	*	0.8	2.0	5.6	17.5
Female						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Hyperplasia of prostate (N40)														
Non-Hispanic[5]														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic White														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic Black														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hyperplasia of prostate (N40)						
Non-Hispanic[5]						
Both sexes						
2001	*	*	0.3	0.8	1.6	5.7
2000[4]	*	*	0.3	0.9	2.0	5.2
1999[4]	*	*	0.4	0.8	2.1	5.1
Male						
2001	*	*	0.8	2.0	4.2	19.3
2000[4]	*	*	0.7	2.1	5.4	18.0
1999[4]	*	*	0.8	2.0	5.6	18.0
Female						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic White						
Both sexes						
2001	*	*	0.3	0.8	1.6	5.9
2000[4]	*	*	0.3	0.9	2.0	5.2
1999[4]	*	*	0.3	0.9	2.0	5.2
Male						
2001	*	*	0.8	2.0	4.2	20.1
2000[4]	*	*	0.7	2.1	5.5	18.0
1999[4]	*	*	0.8	2.2	5.5	18.4
Female						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Hyperplasia of prostate (N40)														
Non-Hispanic Black														
Female														
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Inflammatory diseases of female pelvic organs (N70-N76)														
All origins[3]														
Both sexes														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hyperplasia of prostate (N40)						
Non-Hispanic Black						
Female						
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Inflammatory diseases of female pelvic organs (N70-N76)						
All origins[3]						
Both sexes						
2001	*	*	*	*	*	0.9
2000[4]	*	*	*	*	*	0.6
1999[4]	*	*	*	*	*	*
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	*	*	*	*	*	1.2
2000[4]	*	*	*	*	*	0.9
1999[4]	*	*	*	*	*	*
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	*	*	0.9
2000[4]	*	*	*	*	*	0.6
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Inflammatory diseases of female pelvic organs (N70-N76)														
Non-Hispanic[5]														
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Inflammatory diseases of female pelvic organs (N70-N76)						
Non-Hispanic[5]						
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	*	*	*	*	*	1.2
2000[4]	*	*	*	*	*	0.9
1999[4]	*	*	*	*	*	*
Non-Hispanic White						
Both sexes						
2001	*	*	*	*	*	0.9
2000[4]	*	*	*	*	*	0.6
1999[4]	*	*	*	*	*	*
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	*	*	*	*	*	1.2
2000[4]	*	*	*	*	*	0.9
1999[4]	*	*	*	*	*	*
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Pregnancy, childbirth and the puerperium (O00-O99)														
All origins[3]														
Both sexes														
2001	0.1	. . .	. . .	. . .	*	0.2	0.4	0.5	0.5	0.3	0.1	*	*	*
2000[4]	0.1	. . .	. . .	. . .	*	*	0.4	0.4	0.5	0.4	0.2	*	*	*
1999[4]	0.1	. . .	. . .	. . .	*	0.2	0.3	0.5	0.4	0.4	0.1	*	*	*
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	0.3	. . .	. . .	. . .	*	0.4	0.8	1.0	0.9	0.7	0.3	*	*	*
2000[4]	0.3	. . .	. . .	. . .	*	*	0.8	0.9	0.9	0.8	0.3	*	*	*
1999[4]	0.3	. . .	. . .	. . .	*	0.3	0.7	0.9	0.9	0.8	0.3	*	*	*
Hispanic														
Both sexes														
2001	0.2	. . .	. . .	. . .	*	*	*	0.6	*	*	*	*	*	*
2000[4]	0.2	. . .	. . .	. . .	*	*	0.6	0.6	*	*	*	*	*	*
1999[4]	0.2	. . .	. . .	. . .	*	*	*	*	*	0.8	*	*	*	*
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	0.5	. . .	. . .	. . .	*	*	*	1.3	*	*	*	*	*	*
2000[4]	0.5	. . .	. . .	. . .	*	*	1.4	1.3	*	*	*	*	*	*
1999[4]	0.4	. . .	. . .	. . .	*	*	*	*	*	1.6	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.1	. . .	. . .	. . .	*	0.2	0.3	0.5	0.5	0.3	0.1	*	*	*
2000[4]	0.1	. . .	. . .	. . .	*	*	0.4	0.4	0.5	0.3	0.2	*	*	*
1999[4]	0.1	. . .	. . .	. . .	*	0.2	0.3	0.4	0.5	0.3	0.1	*	*	*
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	0.3	. . .	. . .	. . .	*	0.3	0.7	0.9	0.9	0.6	0.3	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pregnancy, childbirth and the puerperium (O00-O99)						
All origins[3]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Pregnancy, childbirth and the puerperium (O00-O99)														
Non-Hispanic[5]														
Female														
2000[4]	0.3	. . .	. . .	. . .	*	*	0.7	0.8	0.9	0.7	0.3	*	*	*
1999[4]	0.3	. . .	. . .	. . .	*	0.3	0.6	0.9	1.0	0.7	0.3	*	*	*
Non-Hispanic White														
Both sexes														
2001	0.1	. . .	. . .	. . .	*	*	0.2	0.3	0.3	0.2	*	*	*	*
2000[4]	0.1	. . .	. . .	. . .	*	*	0.2	0.3	0.3	0.2	*	*	*	*
1999[4]	0.1	. . .	. . .	. . .	*	*	*	0.2	0.3	0.2	*	*	*	*
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	0.2	. . .	. . .	. . .	*	*	0.4	0.5	0.7	0.3	*	*	*	*
2000[4]	0.2	. . .	. . .	. . .	*	*	0.4	0.5	0.6	0.5	*	*	*	*
1999[4]	0.2	. . .	. . .	. . .	*	*	*	0.5	0.7	0.4	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	0.4	. . .	. . .	. . .	*	*	1.1	1.4	1.1	1.0	*	*	*	*
2000[4]	0.4	. . .	. . .	. . .	*	*	1.1	1.2	1.3	0.9	*	*	*	*
1999[4]	0.5	. . .	. . .	. . .	*	*	1.1	1.4	1.2	1.0	*	*	*	*
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	0.8	. . .	. . .	. . .	*	*	2.2	2.7	2.2	1.9	*	*	*	*
2000[4]	0.8	. . .	. . .	. . .	*	*	2.2	2.3	2.5	1.7	*	*	*	*
1999[4]	0.9	. . .	. . .	. . .	*	*	2.1	2.7	2.4	1.8	*	*	*	*
Pregnancy with abortive outcome (O00-O07)														
All origins[3]														
Both sexes														
2001	0.0	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pregnancy, childbirth and the puerperium (O00-O99)						
Non-Hispanic[5]						
Female						
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic White						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Pregnancy with abortive outcome (O00-O07)						
All origins[3]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Pregnancy with abortive outcome (O00-O07)														
All origins[3]														
Both sexes														
1999[4]	0.0	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	0.0	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	*	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
2000[4]	*	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
1999[4]	*	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	*	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
2000[4]	*	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
1999[4]	*	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.0	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	0.0	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pregnancy with abortive outcome (O00-O07)						
All origins[3]						
Both sexes						
1999[4]	*	*	*	*	*	*
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Pregnancy with abortive outcome (O00-O07)														
Non-Hispanic White														
Both sexes														
2001	*	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
2000[4]	*	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
1999[4]	*	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	*	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
2000[4]	*	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
1999[4]	*	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	*	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
2000[4]	*	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
1999[4]	*	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	*	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
2000[4]	*	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
1999[4]	*	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
Other complications of pregnancy, childbirth and the puerperium (O10-O99)														
All origins[3]														
Both sexes														
2001	0.1	. . .	. . .	. . .	*	0.2	0.3	0.5	0.4	0.3	0.1	*	*	*
2000[4]	0.1	. . .	. . .	. . .	*	*	0.4	0.4	0.4	0.3	0.1	*	*	*
1999[4]	0.1	. . .	. . .	. . .	*	0.1	0.3	0.4	0.4	0.4	0.1	*	*	*
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pregnancy with abortive outcome (O00-O07)						
Non-Hispanic White						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Other complications of pregnancy, childbirth and the puerperium (O10-O99)						
All origins[3]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other complications of pregnancy, childbirth and the puerperium (O10-O99)														
All origins[3]														
Male														
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	0.3	. . .	. . .	. . .	*	0.3	0.7	0.9	0.9	0.6	0.3	*	*	*
2000[4]	0.3	. . .	. . .	. . .	*	*	0.8	0.8	0.9	0.7	0.3	*	*	*
1999[4]	0.3	. . .	. . .	. . .	*	0.3	0.6	0.9	0.8	0.7	0.3	*	*	*
Hispanic														
Both sexes														
2001	0.2	. . .	. . .	. . .	*	*	*	0.6	*	*	*	*	*	*
2000[4]	0.2	. . .	. . .	. . .	*	*	0.6	*	*	*	*	*	*	*
1999[4]	0.2	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	0.5	. . .	. . .	. . .	*	*	*	1.3	*	*	*	*	*	*
2000[4]	0.5	. . .	. . .	. . .	*	*	1.3	*	*	*	*	*	*	*
1999[4]	0.4	. . .	. . .	. . .	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.1	. . .	. . .	. . .	*	0.1	0.3	0.4	0.4	0.3	0.1	*	*	*
2000[4]	0.1	. . .	. . .	. . .	*	*	0.3	0.4	0.4	0.3	0.1	*	*	*
1999[4]	0.1	. . .	. . .	. . .	*	0.2	0.3	0.4	0.4	0.3	0.1	*	*	*
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	0.2	. . .	. . .	. . .	*	0.3	0.6	0.9	0.8	0.6	0.2	*	*	*
2000[4]	0.2	. . .	. . .	. . .	*	*	0.7	0.7	0.8	0.6	0.3	*	*	*
1999[4]	0.2	. . .	. . .	. . .	*	0.3	0.6	0.8	0.8	0.6	0.3	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other complications of pregnancy, childbirth and the puerperium (O10-O99)						
All origins[3]						
Male						
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other complications of pregnancy, childbirth and the puerperium (O10-O99)														
Non-Hispanic White														
Both sexes														
2001	0.1	. . .	. . .	. . .	*	*	0.2	0.2	0.3	0.1	*	*	*	*
2000[4]	0.1	. . .	. . .	. . .	*	*	0.2	0.2	0.3	0.2	*	*	*	*
1999[4]	0.1	. . .	. . .	. . .	*	*	*	0.2	0.3	0.2	*	*	*	*
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	0.1	. . .	. . .	. . .	*	*	0.3	0.5	0.6	0.3	*	*	*	*
2000[4]	0.1	. . .	. . .	. . .	*	*	0.4	0.5	0.6	0.4	*	*	*	*
1999[4]	0.1	. . .	. . .	. . .	*	*	*	0.5	0.6	0.4	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	0.4	. . .	. . .	. . .	*	*	1.0	1.3	1.0	0.9	*	*	*	*
2000[4]	0.3	. . .	. . .	. . .	*	*	1.0	1.0	1.1	0.7	*	*	*	*
1999[4]	0.4	. . .	. . .	. . .	*	*	1.0	1.3	1.1	0.8	*	*	*	*
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2001	0.7	. . .	. . .	. . .	*	*	2.0	2.5	1.9	1.6	*	*	*	*
2000[4]	0.7	. . .	. . .	. . .	*	*	1.9	1.9	2.2	1.4	*	*	*	*
1999[4]	0.8	. . .	. . .	. . .	*	*	2.0	2.4	2.1	1.6	*	*	*	*
Certain conditions originating in the perinatal period (P00-P96)														
All origins[3]														
Both sexes														
2001	4.9	340.5	0.5	0.1	*	*	*	*	*	*	*	*	*	*
2000[4]	5.0	365.7	0.5	0.1	*	*	*	*	*	*	*	*	*	*
1999[4]	5.1	371.0	0.6	0.1	*	*	*	*	*	*	*	*	*	*
Male														
2001	5.6	376.6	0.6	*	*	*	*	*	*	*	*	*	*	*
2000[4]	5.8	403.9	0.7	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other complications of pregnancy, childbirth and the puerperium (O10-O99)						
Non-Hispanic White						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	. . .	. . .	. . .	. . .	. . .	. . .
2000[4]	. . .	. . .	. . .	. . .	. . .	. . .
1999[4]	. . .	. . .	. . .	. . .	. . .	. . .
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Certain conditions originating in the perinatal period (P00-P96)						
All origins[3]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Certain conditions originating in the perinatal period (P00-P96)														
All origins[3]														
Male														
1999[4]	5.9	408.6	0.7	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	4.2	302.6	0.4	*	*	*	*	*	*	*	*	*	*	*
2000[4]	4.3	325.6	0.4	*	*	*	*	*	*	*	*	*	*	*
1999[4]	4.4	331.7	0.5	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	6.0	269.6	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	6.1	274.9	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	6.3	281.4	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	6.7	302.0	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	6.6	299.0	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	6.8	305.4	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	5.3	235.9	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	5.5	249.7	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	5.8	256.4	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	4.6	353.5	0.5	*	*	*	*	*	*	*	*	*	*	*
2000[4]	4.7	380.5	0.5	*	*	*	*	*	*	*	*	*	*	*
1999[4]	4.8	383.8	0.6	0.1	*	*	*	*	*	*	*	*	*	*
Male														
2001	5.4	390.1	0.6	*	*	*	*	*	*	*	*	*	*	*
2000[4]	5.5	421.7	0.7	*	*	*	*	*	*	*	*	*	*	*
1999[4]	5.6	423.2	0.8	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	3.9	315.1	0.3	*	*	*	*	*	*	*	*	*	*	*
2000[4]	4.0	337.2	0.4	*	*	*	*	*	*	*	*	*	*	*
1999[4]	4.1	342.4	0.4	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	3.2	260.3	0.4	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Certain conditions originating in the perinatal period (P00-P96)						
All origins[3]						
Male						
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic White						
Both sexes						
2001	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Certain conditions originating in the perinatal period (P00-P96)														
Non-Hispanic White														
Both sexes														
2000[4]	3.2	275.5	0.4	*	*	*	*	*	*	*	*	*	*	*
1999[4]	3.2	275.1	0.4	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	3.7	286.6	0.5	*	*	*	*	*	*	*	*	*	*	*
2000[4]	3.7	305.5	0.5	*	*	*	*	*	*	*	*	*	*	*
1999[4]	3.8	304.4	0.5	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	2.7	232.8	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	2.7	243.9	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	2.7	244.2	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	13.2	765.0	0.9	*	*	*	*	*	*	*	*	*	*	*
2000[4]	13.9	838.3	1.1	*	*	*	*	*	*	*	*	*	*	*
1999[4]	14.4	872.7	1.2	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	15.8	846.6	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	16.6	932.9	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	17.1	968.0	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	10.9	679.1	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	11.5	740.7	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	12.0	774.4	*	*	*	*	*	*	*	*	*	*	*	*
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)														
All origins[3]														
Both sexes														
2001	3.7	136.7	3.6	0.9	0.9	1.3	1.3	1.2	1.1	1.2	1.4	1.5	1.8	2.1
2000[4]	3.8	150.9	3.2	1.0	1.0	1.1	1.1	1.1	1.3	1.2	1.3	1.4	1.8	2.2
1999[4]	3.7	144.2	3.6	1.0	1.1	1.1	1.1	1.1	1.2	1.2	1.2	1.7	1.8	2.3
Male														
2001	3.9	143.1	3.6	1.0	1.0	1.5	1.4	1.3	1.2	1.2	1.6	1.5	1.9	2.1
2000[4]	4.0	155.4	3.3	1.0	1.0	1.3	1.4	1.2	1.5	1.4	1.5	1.6	2.2	2.0
1999[4]	3.9	147.9	4.0	1.0	1.1	1.3	1.3	1.1	1.3	1.3	1.3	1.9	2.0	2.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Certain conditions originating in the perinatal period (P00-P96)						
Non-Hispanic White						
Both sexes						
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)						
All origins[3]						
Both sexes						
2001	2.5	2.2	2.7	3.3	4.3	7.2
2000[4]	2.5	2.6	2.4	3.3	5.0	7.5
1999[4]	2.4	2.6	2.5	3.4	5.0	7.8
Male						
2001	2.5	2.5	3.1	3.6	4.8	7.7
2000[4]	2.6	2.7	2.6	3.3	5.3	10.6
1999[4]	2.5	2.4	2.6	3.4	5.7	8.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)														
All origins[3]														
Female														
2001	3.4	129.9	3.7	0.8	0.8	1.0	1.1	1.1	1.0	1.1	1.3	1.4	1.8	2.1
2000[4]	3.5	146.2	3.1	0.9	1.0	0.9	0.9	0.9	1.1	1.1	1.1	1.3	1.5	2.4
1999[4]	3.5	140.3	3.2	1.0	1.1	0.9	1.0	1.1	1.2	1.1	1.2	1.6	1.7	2.2
Hispanic														
Both sexes														
2001	4.4	146.2	3.4	0.8	0.9	1.0	1.2	0.7	*	*	0.9	1.2	*	*
2000[4]	4.6	153.3	3.8	0.9	*	0.8	0.9	*	0.9	0.8	*	1.1	*	*
1999[4]	4.4	141.8	3.5	0.9	0.9	0.9	0.9	0.9	0.9	*	*	1.5	*	*
Male														
2001	4.7	153.4	3.5	*	*	*	1.1	*	*	*	*	*	*	*
2000[4]	4.7	153.3	4.0	*	*	*	1.1	*	*	*	*	*	*	*
1999[4]	4.3	138.1	4.3	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	4.1	138.8	3.3	*	*	*	*	*	*	*	*	*	*	*
2000[4]	4.4	153.3	3.5	*	*	*	*	*	*	*	*	*	*	*
1999[4]	4.5	145.8	2.7	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	3.5	132.8	3.7	0.9	0.9	1.3	1.3	1.3	1.2	1.2	1.5	1.5	1.9	2.2
2000[4]	3.6	148.8	3.1	1.0	1.0	1.2	1.2	1.2	1.4	1.3	1.4	1.5	1.9	2.2
1999[4]	3.6	143.1	3.6	1.0	1.1	1.1	1.2	1.1	1.3	1.3	1.3	1.7	1.8	2.4
Male														
2001	3.7	139.4	3.6	1.0	1.0	1.6	1.5	1.3	1.4	1.3	1.6	1.5	1.9	2.2
2000[4]	3.9	154.4	3.1	1.0	1.0	1.4	1.4	1.3	1.7	1.4	1.5	1.6	2.2	2.0
1999[4]	3.8	148.4	3.9	1.0	1.1	1.4	1.4	1.2	1.4	1.4	1.4	1.9	2.0	2.5
Female														
2001	3.3	125.8	3.7	0.9	0.9	0.9	1.1	1.2	1.1	1.2	1.4	1.5	1.8	2.2
2000[4]	3.4	142.8	3.0	0.9	1.1	1.0	1.0	1.0	1.1	1.1	1.2	1.3	1.6	2.4
1999[4]	3.4	137.4	3.3	1.1	1.1	0.9	1.0	1.1	1.2	1.2	1.2	1.6	1.7	2.2
Non-Hispanic White														
Both sexes														
2001	3.4	127.4	3.1	0.9	0.9	1.2	1.2	1.3	1.3	1.2	1.5	1.4	2.0	2.3
2000[4]	3.4	141.2	2.6	0.9	1.0	1.0	1.2	1.2	1.5	1.3	1.3	1.5	2.0	2.2
1999[4]	3.4	136.8	3.0	0.9	1.0	1.1	1.1	1.2	1.3	1.3	1.3	1.8	1.9	2.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)						
All origins[3]						
Female						
2001	2.6	2.0	2.4	3.1	4.0	7.0
2000[4]	2.5	2.5	2.3	3.3	4.8	6.3
1999[4]	2.3	2.8	2.3	3.4	4.5	7.6
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	2.6	2.3	2.7	3.2	4.3	7.2
2000[4]	2.5	2.6	2.5	3.2	4.9	7.5
1999[4]	2.4	2.6	2.4	3.4	4.9	7.9
Male						
2001	2.4	2.5	3.0	3.5	4.8	7.6
2000[4]	2.5	2.8	2.6	3.1	5.2	10.4
1999[4]	2.6	2.4	2.5	3.4	5.7	8.5
Female						
2001	2.8	2.1	2.4	3.0	4.1	7.0
2000[4]	2.5	2.5	2.4	3.3	4.8	6.4
1999[4]	2.2	2.8	2.3	3.5	4.5	7.6
Non-Hispanic White						
Both sexes						
2001	2.8	2.4	2.7	3.3	4.4	7.4
2000[4]	2.6	2.7	2.5	3.3	5.1	7.8
1999[4]	2.5	2.7	2.4	3.4	5.0	8.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)														
Non-Hispanic White														
Male														
2001	3.6	135.5	3.1	1.0	1.0	1.5	1.3	1.5	1.5	1.3	1.6	1.4	2.0	2.3
2000[4]	3.6	147.9	2.6	0.8	0.9	1.1	1.5	1.3	1.8	1.4	1.5	1.6	2.3	2.0
1999[4]	3.6	144.4	3.2	0.8	1.0	1.4	1.4	1.3	1.4	1.5	1.4	1.9	2.0	2.7
Female														
2001	3.2	118.8	3.2	0.8	0.7	0.9	1.0	1.1	1.2	1.1	1.4	1.5	1.9	2.3
2000[4]	3.2	134.2	2.6	0.9	1.0	0.9	1.0	1.0	1.2	1.1	1.2	1.4	1.8	2.4
1999[4]	3.3	128.9	2.9	1.0	1.0	0.8	0.9	1.2	1.2	1.2	1.2	1.6	1.8	2.4
Non-Hispanic Black														
Both sexes														
2001	4.7	163.5	5.4	1.0	1.2	2.0	1.9	1.4	1.2	1.5	1.7	1.8	1.6	2.0
2000[4]	5.0	184.1	5.2	1.3	1.4	1.9	1.3	1.4	1.4	1.6	1.7	1.7	1.5	2.4
1999[4]	4.8	174.8	5.8	1.5	1.8	1.4	1.5	1.1	1.2	1.5	1.3	1.8	1.9	2.0
Male														
2001	5.1	167.5	5.5	*	*	2.7	2.3	*	*	*	2.2	*	*	*
2000[4]	5.5	186.0	5.6	1.7	1.3	2.6	*	1.8	1.7	1.7	2.0	2.1	*	*
1999[4]	5.3	179.5	6.6	1.4	1.8	1.7	1.9	*	*	*	1.8	2.4	*	*
Female														
2001	4.4	159.2	5.4	*	1.4	1.4	1.5	1.8	*	1.7	*	2.0	*	*
2000[4]	4.5	182.2	4.8	*	1.5	*	*	*	*	1.5	*	*	*	2.9
1999[4]	4.4	170.0	5.0	1.5	1.8	*	*	*	*	1.5	*	*	*	*
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)														
All origins[3]														
Both sexes														
2001	11.4	86.5	1.8	0.4	0.5	1.8	3.0	3.5	4.2	6.2	7.7	9.1	8.8	8.4
2000[4]	11.3	95.6	1.6	0.4	0.4	1.6	2.9	3.4	4.3	5.7	7.6	8.6	8.0	8.6
1999[4]	9.6	93.2	1.5	0.3	0.4	1.1	2.0	2.3	3.1	4.1	5.1	5.9	5.8	6.9
Male														
2001	11.3	98.9	2.1	0.4	0.5	2.4	4.0	4.8	5.6	8.0	9.9	11.9	11.8	11.0
2000[4]	11.5	110.6	1.7	0.5	0.5	1.9	4.0	4.7	5.9	7.6	10.2	11.7	11.2	11.4
1999[4]	9.4	106.6	1.7	0.3	0.4	1.5	2.8	2.9	4.2	5.3	6.7	8.2	7.9	9.1
Female														
2001	11.4	73.6	1.5	0.4	0.5	1.1	1.9	2.2	2.8	4.4	5.5	6.3	5.9	5.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)						
Non-Hispanic White						
Male						
2001	2.6	2.7	3.1	3.8	4.5	7.4
2000[4]	2.7	2.6	2.5	3.2	5.5	10.8
1999[4]	2.7	2.4	2.5	3.0	5.9	8.6
Female						
2001	3.0	2.1	2.4	3.0	4.3	7.4
2000[4]	2.5	2.7	2.5	3.4	4.9	6.6
1999[4]	2.2	2.9	2.3	3.6	4.5	7.8
Non-Hispanic Black						
Both sexes						
2001	1.9	*	2.9	*	*	*
2000[4]	2.4	2.8	*	*	*	*
1999[4]	2.0	2.4	3.5	4.0	*	7.4
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	3.4	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)						
All origins[3]						
Both sexes						
2001	9.9	12.7	17.8	29.8	54.3	206.4
2000[4]	10.2	14.2	19.0	30.1	56.5	208.1
1999[4]	8.9	11.9	17.6	26.5	50.5	187.7
Male						
2001	13.2	15.5	20.9	33.2	60.6	175.0
2000[4]	13.4	18.9	22.4	36.1	62.3	182.7
1999[4]	11.9	15.1	21.8	31.5	54.4	166.3
Female						
2001	6.9	10.2	15.4	27.4	50.4	219.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)														
All origins[3]														
Female														
2000[4]	11.2	79.9	1.5	0.4	0.4	1.2	1.8	2.2	2.8	3.7	5.0	5.6	5.0	5.9
1999[4]	9.7	79.2	1.3	0.3	0.3	0.8	1.2	1.6	2.1	3.0	3.4	3.7	3.9	4.9
Hispanic														
Both sexes														
2001	5.9	60.7	1.6	*	0.6	2.3	3.2	3.7	4.7	5.9	6.1	9.0	7.4	6.7
2000[4]	5.9	65.6	1.7	*	*	1.6	2.6	3.5	4.5	6.3	7.2	8.1	7.9	7.5
1999[4]	3.8	54.9	1.0	*	*	1.0	1.6	1.8	2.6	2.9	3.1	3.8	2.3	3.2
Male														
2001	7.4	67.2	1.8	*	*	3.4	4.3	5.4	6.6	8.3	8.6	14.0	10.7	10.0
2000[4]	7.5	78.6	2.1	*	*	1.7	3.9	5.1	6.5	9.3	10.6	12.4	12.3	9.0
1999[4]	4.4	61.3	*	*	*	1.3	2.4	2.4	3.5	4.2	4.5	6.2	*	*
Female														
2001	4.3	53.9	*	*	*	*	1.8	1.7	2.5	3.2	3.6	4.0	4.4	3.8
2000[4]	4.1	52.1	1.4	*	*	1.4	*	1.6	2.3	3.0	3.6	3.8	3.6	6.2
1999[4]	3.1	48.2	*	*	*	*	*	*	1.6	1.6	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	12.1	92.6	1.8	0.4	0.5	1.7	2.9	3.4	4.1	6.2	7.8	9.0	8.8	8.5
2000[4]	12.0	102.4	1.6	0.4	0.4	1.6	2.9	3.4	4.3	5.5	7.5	8.6	7.9	8.5
1999[4]	10.3	101.9	1.6	0.3	0.4	1.2	2.1	2.4	3.2	4.2	5.2	6.0	6.0	7.1
Male														
2001	11.8	106.2	2.1	0.4	0.5	2.2	3.9	4.6	5.3	7.9	9.9	11.6	11.8	11.1
2000[4]	11.9	118.0	1.7	0.5	0.5	1.9	3.9	4.5	5.6	7.2	10.0	11.5	10.9	11.3
1999[4]	10.0	116.7	1.9	0.3	0.4	1.5	2.9	3.0	4.3	5.3	6.9	8.3	8.1	9.3
Female														
2001	12.3	78.3	1.5	0.3	0.4	1.1	1.9	2.2	2.9	4.6	5.7	6.4	6.0	6.1
2000[4]	12.1	86.1	1.5	0.3	0.4	1.2	1.9	2.2	2.9	3.8	5.1	5.7	5.0	5.8
1999[4]	10.5	86.3	1.4	0.3	0.3	0.8	1.3	1.7	2.2	3.1	3.6	3.9	4.0	5.1
Non-Hispanic White														
Both sexes														
2001	12.4	73.1	1.5	0.3	0.4	1.7	2.8	3.3	3.9	6.0	7.7	8.7	8.3	7.9
2000[4]	12.2	79.7	1.4	0.3	0.4	1.5	2.8	3.1	4.0	5.4	7.2	8.0	7.1	7.7
1999[4]	10.2	79.3	1.4	0.3	0.3	1.0	1.8	2.0	3.0	3.8	4.5	5.2	5.2	6.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)						
All origins[3]						
Female						
2000[4]	7.3	10.1	16.4	26.0	53.0	218.5
1999[4]	6.2	9.2	14.2	23.0	48.3	196.4
Hispanic						
Both sexes						
2001	7.1	8.7	11.3	14.4	27.8	108.2
2000[4]	5.6	8.2	12.2	16.5	27.8	102.2
1999[4]	4.7	6.2	8.1	13.9	29.4	87.0
Male						
2001	11.4	12.8	16.0	18.4	37.7	87.6
2000[4]	8.9	13.1	16.0	19.9	36.8	106.8
1999[4]	6.9	8.5	*	21.8	*	77.6
Female						
2001	*	*	7.7	11.6	21.7	118.2
2000[4]	*	*	9.2	14.1	22.4	99.9
1999[4]	*	*	8.0	*	30.0	91.5
Non-Hispanic[5]						
Both sexes						
2001	10.0	12.9	18.2	30.4	55.3	209.6
2000[4]	10.5	14.5	19.3	30.6	57.4	211.6
1999[4]	9.1	12.2	17.9	27.0	51.2	190.9
Male						
2001	13.2	15.7	21.1	33.8	61.6	178.2
2000[4]	13.5	19.0	22.6	36.6	63.0	185.4
1999[4]	12.1	15.4	22.3	31.8	55.2	169.5
Female						
2001	7.1	10.5	15.8	28.0	51.6	222.7
2000[4]	7.7	10.5	16.7	26.5	54.1	222.3
1999[4]	6.5	9.4	14.5	23.6	48.9	199.5
Non-Hispanic White						
Both sexes						
2001	9.5	12.1	17.5	30.1	55.0	213.3
2000[4]	9.7	13.2	18.7	30.0	57.0	215.8
1999[4]	8.4	11.3	17.3	25.9	50.1	193.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)														
Non-Hispanic White														
Male														
2001	11.8	85.2	1.7	*	0.5	2.1	3.8	4.6	5.1	7.5	9.9	11.2	11.1	10.0
2000[4]	11.8	91.8	1.7	0.4	0.4	1.9	3.8	4.2	5.4	7.2	9.8	10.9	9.7	10.0
1999[4]	9.6	92.3	1.7	0.3	0.4	1.3	2.6	2.7	3.8	4.9	6.0	6.9	7.0	8.2
Female														
2001	13.1	60.4	1.3	*	*	1.2	1.9	1.9	2.8	4.5	5.4	6.2	5.5	5.9
2000[4]	12.6	66.9	1.2	*	0.4	1.0	1.7	2.0	2.6	3.5	4.7	5.2	4.5	5.6
1999[4]	10.9	65.5	1.0	0.4	*	0.7	1.0	1.4	2.1	2.6	3.0	3.5	3.5	4.6
Non-Hispanic Black														
Both sexes														
2001	12.9	176.9	3.0	0.6	0.7	1.7	3.8	4.8	5.5	8.5	10.1	13.2	14.9	14.6
2000[4]	13.9	205.6	2.3	0.6	0.8	2.2	4.3	5.5	6.4	7.3	11.3	13.9	16.0	15.8
1999[4]	13.0	202.0	2.9	*	*	1.8	3.8	4.4	5.5	7.9	10.5	13.2	13.2	15.0
Male														
2001	14.2	195.9	3.7	*	*	2.4	5.3	5.6	6.6	10.9	12.0	17.4	19.2	22.2
2000[4]	15.7	236.8	2.2	*	*	2.1	5.7	7.3	8.2	8.5	13.3	18.3	23.5	24.7
1999[4]	14.9	228.0	2.9	*	*	2.5	4.9	5.5	8.4	9.5	13.8	20.0	18.7	22.3
Female														
2001	11.7	156.8	2.2	*	*	*	2.4	4.1	4.5	6.4	8.4	9.6	11.2	8.4
2000[4]	12.4	173.4	2.4	*	*	2.2	3.0	3.8	4.8	6.3	9.4	10.1	9.7	8.5
1999[4]	11.2	175.3	2.8	*	*	*	2.7	3.5	2.9	6.5	7.5	7.5	8.5	9.1
All other diseases (Residual)														
All origins[3]														
Both sexes														
2001	65.5	28.0	4.2	1.9	2.3	3.8	5.2	6.5	8.9	14.1	21.7	32.1	40.9	53.2
2000[4]	63.4	29.0	4.0	1.8	2.5	3.6	5.4	6.5	9.0	13.9	22.1	31.0	40.4	53.1
1999[4]	61.5	30.2	4.2	2.1	2.4	3.9	5.3	6.3	8.9	14.0	21.6	29.9	38.4	51.0
Male														
2001	53.8	31.4	4.6	1.8	2.5	4.4	6.0	7.3	9.9	16.8	25.7	38.8	49.1	59.9
2000[4]	52.3	32.3	4.3	1.9	2.6	4.2	6.5	7.5	10.4	16.5	26.3	38.5	47.6	60.9
1999[4]	51.0	35.0	4.5	2.2	2.5	4.7	5.9	7.3	10.4	16.8	26.6	36.3	45.4	59.5
Female														
2001	76.7	24.5	3.8	2.1	2.0	3.1	4.4	5.6	8.0	11.5	17.7	25.6	33.0	46.9
2000[4]	74.0	25.6	3.6	1.8	2.3	3.1	4.4	5.4	7.5	11.3	17.9	23.7	33.4	45.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)						
Non-Hispanic White						
Male						
2001	12.1	14.4	20.5	33.9	61.6	180.4
2000[4]	12.5	17.1	21.6	35.9	62.6	189.6
1999[4]	11.2	14.2	21.5	30.5	53.3	171.1
Female						
2001	7.0	10.0	15.1	27.4	51.0	227.0
2000[4]	7.1	9.7	16.3	25.9	53.6	226.4
1999[4]	5.9	8.8	14.0	22.7	48.2	202.8
Non-Hispanic Black						
Both sexes						
2001	15.7	21.8	28.1	39.2	67.6	194.2
2000[4]	19.1	28.3	30.3	44.7	74.1	189.2
1999[4]	17.1	21.9	27.8	43.3	73.4	182.0
Male						
2001	25.9	29.7	32.0	41.2	71.1	186.0
2000[4]	25.5	41.0	38.6	55.4	85.0	163.2
1999[4]	23.9	30.0	36.4	54.6	89.4	180.7
Female						
2001	7.7	16.0	25.5	37.9	65.8	197.2
2000[4]	14.1	18.9	24.7	38.3	68.7	198.9
1999[4]	11.8	15.9	22.1	36.4	65.4	182.6
All other diseases (Residual)						
All origins[3]						
Both sexes						
2001	76.4	112.0	172.3	295.5	539.3	1,476.7
2000[4]	74.1	106.0	168.2	288.9	537.8	1,445.7
1999[4]	72.1	107.7	167.8	281.3	510.8	1,410.7
Male						
2001	85.8	124.3	186.5	307.6	539.6	1,271.6
2000[4]	83.1	117.6	183.1	303.4	537.6	1,266.2
1999[4]	80.9	117.1	184.3	298.1	528.1	1,225.4
Female						
2001	67.9	101.4	161.0	287.0	539.1	1,562.6
2000[4]	65.9	96.1	156.5	278.8	537.8	1,518.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
All other diseases (Residual)														
All origins[3]														
Female														
1999[4]	71.6	25.3	3.9	2.1	2.3	3.1	4.7	5.3	7.4	11.3	16.6	23.7	31.7	43.1
Hispanic														
Both sexes														
2001	21.2	22.9	3.7	1.7	1.8	3.2	4.1	4.6	5.8	9.7	15.4	22.8	30.7	42.2
2000[4]	20.7	25.4	3.7	1.6	1.8	2.6	3.9	5.0	6.8	11.0	17.1	24.8	31.0	45.2
1999[4]	21.5	26.5	3.7	1.4	1.9	3.1	3.8	5.1	8.0	11.2	17.4	26.0	31.2	43.3
Male														
2001	20.2	25.4	4.3	1.6	2.0	3.9	4.2	5.0	7.1	12.6	20.0	30.6	40.0	51.7
2000[4]	20.2	25.3	3.6	1.6	2.1	3.2	4.6	5.5	8.8	14.3	23.1	33.3	38.7	59.2
1999[4]	21.0	32.1	3.4	1.6	1.7	3.6	3.9	5.9	10.6	15.3	23.2	35.4	39.2	53.2
Female														
2001	22.3	20.2	3.1	1.8	1.5	2.5	3.9	4.0	4.3	6.5	10.6	14.9	21.9	33.5
2000[4]	21.4	25.5	3.8	1.6	1.4	1.9	3.0	4.4	4.6	7.4	10.9	16.3	23.7	32.5
1999[4]	22.0	20.6	4.0	1.3	2.1	2.5	3.7	4.2	5.0	6.7	11.4	16.7	23.5	34.4
Non-Hispanic[5]														
Both sexes														
2001	71.8	29.0	4.3	2.0	2.3	3.9	5.4	6.9	9.5	14.8	22.3	32.9	41.5	53.8
2000[4]	69.3	29.6	4.0	1.9	2.6	3.8	5.8	6.8	9.3	14.3	22.5	31.3	40.9	53.4
1999[4]	66.8	30.8	4.3	2.3	2.5	4.0	5.6	6.5	9.0	14.3	21.9	30.0	38.8	51.3
Male														
2001	58.8	32.7	4.6	1.8	2.6	4.5	6.4	7.8	10.4	17.4	26.2	39.4	49.5	60.0
2000[4]	56.9	33.6	4.4	1.9	2.7	4.3	6.9	7.9	10.7	16.8	26.5	38.7	48.0	60.8
1999[4]	55.2	35.1	4.7	2.3	2.6	4.9	6.3	7.6	10.3	16.8	26.7	36.1	45.6	59.6
Female														
2001	84.2	25.2	4.0	2.1	2.1	3.2	4.5	6.0	8.6	12.1	18.5	26.6	33.9	47.9
2000[4]	81.0	25.4	3.5	1.8	2.4	3.3	4.6	5.6	7.9	11.8	18.6	24.2	34.1	46.5
1999[4]	77.9	26.2	3.9	2.2	2.4	3.2	4.8	5.6	7.7	11.8	17.1	24.2	32.3	43.6
Non-Hispanic White														
Both sexes														
2001	77.3	23.2	3.9	2.0	2.2	3.6	5.0	6.0	8.2	13.0	19.6	29.2	37.5	49.8
2000[4]	74.3	24.0	3.4	1.7	2.4	3.7	5.4	5.8	8.1	12.3	19.9	27.3	37.3	49.3
1999[4]	71.5	24.2	4.0	2.1	2.3	3.9	5.0	6.0	8.0	12.1	19.0	26.0	34.8	47.4
Male														
2001	61.9	25.3	4.4	1.9	2.4	4.4	6.2	7.0	9.3	15.8	23.5	35.5	45.0	55.1
2000[4]	59.6	26.7	3.8	1.9	2.7	4.3	6.6	7.3	9.7	14.9	23.9	34.7	44.1	56.1

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other diseases (Residual)						
All origins[3]						
Female						
1999[4]	64.2	99.5	154.8	269.7	500.7	1,485.3
Hispanic						
Both sexes						
2001	59.7	83.9	136.4	209.8	396.3	991.0
2000[4]	50.5	84.3	127.4	217.6	404.9	913.0
1999[4]	61.6	88.6	135.5	202.4	374.7	1,039.7
Male						
2001	69.0	96.0	144.6	205.3	403.8	939.0
2000[4]	55.8	105.9	138.1	221.5	378.4	870.7
1999[4]	74.8	96.6	159.9	220.0	389.8	923.1
Female						
2001	51.7	74.0	130.2	213.0	391.7	1,016.5
2000[4]	45.9	66.7	119.3	214.9	421.1	933.8
1999[4]	50.3	82.2	117.2	190.0	365.7	1,096.0
Non-Hispanic[5]						
Both sexes						
2001	77.5	113.6	174.0	299.1	543.9	1,491.3
2000[4]	75.6	107.1	169.8	291.3	541.5	1,462.0
1999[4]	72.6	108.5	169.1	284.1	514.7	1,420.1
Male						
2001	86.7	125.7	188.5	311.8	544.1	1,282.3
2000[4]	84.6	117.8	184.8	306.1	542.6	1,279.8
1999[4]	80.8	117.9	185.0	300.8	532.1	1,234.0
Female						
2001	69.1	103.1	162.4	290.1	543.7	1,578.3
2000[4]	67.3	97.8	158.0	281.0	540.9	1,535.7
1999[4]	65.1	100.4	156.6	272.4	504.6	1,494.7
Non-Hispanic White						
Both sexes						
2001	73.3	110.2	170.7	296.0	545.5	1,512.9
2000[4]	71.2	103.2	165.9	287.8	540.7	1,481.3
1999[4]	68.6	103.6	164.6	279.7	516.0	1,443.8
Male						
2001	82.2	121.0	183.6	306.0	544.6	1,293.1
2000[4]	78.5	112.4	178.8	300.6	538.2	1,288.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
All other diseases (Residual)														
Non-Hispanic White														
Male														
1999[4]	57.6	27.4	4.5	2.1	2.3	5.0	5.9	7.2	9.4	14.8	24.0	31.8	41.2	55.0
Female														
2001	92.1	21.0	3.4	2.1	2.0	2.8	3.8	5.0	7.2	10.2	15.8	23.0	30.2	44.7
2000[4]	88.3	21.2	2.9	1.6	2.1	3.0	4.1	4.3	6.6	9.7	15.9	19.9	30.7	42.7
1999[4]	84.9	20.9	3.5	2.1	2.3	2.9	4.1	4.7	6.5	9.4	14.0	20.3	28.5	40.1
Non-Hispanic Black														
Both sexes														
2001	61.1	53.8	5.6	2.0	3.0	5.4	8.0	12.0	17.3	27.4	42.3	64.1	80.9	97.2
2000[4]	60.1	51.3	6.1	2.4	3.5	4.9	8.6	13.4	17.8	27.8	42.7	65.4	77.2	97.4
1999[4]	58.0	58.4	5.5	3.0	3.2	4.9	9.3	10.9	16.0	29.5	43.4	63.0	77.3	93.5
Male														
2001	56.4	61.3	5.2	1.6	3.4	5.6	8.1	12.9	18.2	30.5	48.2	75.8	97.7	116.3
2000[4]	56.1	61.4	6.4	2.1	3.2	5.1	9.4	13.6	18.9	30.9	48.8	75.8	90.8	116.7
1999[4]	54.4	66.5	5.6	3.3	3.9	5.0	10.1	11.0	16.7	32.5	48.5	74.5	91.6	112.6
Female														
2001	65.2	46.0	6.0	2.5	2.7	5.2	8.0	11.2	16.5	24.6	37.2	54.1	66.6	81.6
2000[4]	63.7	40.9	5.8	2.7	3.7	4.8	7.8	13.2	16.8	25.0	37.3	56.4	65.7	81.6
1999[4]	61.3	50.0	5.4	2.7	2.5	4.7	8.6	10.8	15.3	26.9	38.9	53.2	65.2	78.0
Accidents (unintentional injuries) (V01-X59,Y85-Y86)														
All origins[3]														
Both sexes														
2001	35.7	24.2	11.2	6.4	7.4	32.8	39.5	31.4	28.5	34.1	36.7	36.0	31.9	29.3
2000[4]	34.8	23.1	11.9	6.8	7.7	33.4	38.8	30.4	28.6	32.9	35.4	34.2	30.8	29.9
1999[4]	35.1	22.3	12.4	7.1	8.1	33.3	37.5	30.0	29.2	32.2	35.4	34.1	29.2	30.1
Male														
2001	47.2	26.2	13.3	7.6	9.5	45.3	60.7	48.0	43.1	49.5	54.1	54.0	46.9	41.8
2000[4]	46.2	25.7	14.0	7.8	9.9	45.8	58.9	47.0	43.1	48.9	53.1	51.6	46.0	43.1
1999[4]	46.4	25.4	14.4	8.5	10.4	44.8	57.1	46.1	43.9	47.4	53.1	51.7	43.6	43.6
Female														
2001	24.5	22.1	8.9	5.0	5.3	19.5	17.2	14.3	13.8	18.7	19.6	18.5	17.5	17.6
2000[4]	23.8	20.5	9.7	5.7	5.5	20.3	17.8	13.5	13.9	17.0	18.0	17.3	16.1	17.5
1999[4]	24.1	19.0	10.3	5.6	5.6	21.1	17.1	13.7	14.3	17.0	18.1	17.0	15.5	17.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other diseases (Residual)						
Non-Hispanic White						
Male						
1999[4]	76.3	112.0	179.6	293.3	529.5	1,239.8
Female						
2001	65.1	100.7	160.1	288.9	546.0	1,604.4
2000[4]	64.4	95.2	155.5	278.8	542.2	1,559.1
1999[4]	61.5	96.2	152.6	270.1	508.1	1,525.2
Non-Hispanic Black						
Both sexes						
2001	124.2	165.8	239.3	384.5	619.4	1,431.5
2000[4]	124.6	159.8	237.4	375.9	624.5	1,403.3
1999[4]	118.8	169.4	242.1	373.1	560.6	1,295.3
Male						
2001	144.0	199.8	276.5	449.4	669.7	1,329.4
2000[4]	153.5	191.1	284.0	432.1	698.6	1,340.9
1999[4]	135.8	192.6	274.5	439.4	638.5	1,303.8
Female						
2001	108.6	140.6	214.5	345.7	593.9	1,469.3
2000[4]	101.8	136.6	206.4	342.1	587.2	1,426.5
1999[4]	105.5	152.2	220.6	333.1	522.0	1,292.1
Accidents (unintentional injuries) (V01-X59,Y85-Y86)						
All origins[3]						
Both sexes						
2001	31.5	36.8	49.3	80.4	130.4	276.4
2000[4]	32.2	35.4	48.8	75.5	124.6	273.5
1999[4]	31.2	38.7	50.9	78.6	133.3	282.4
Male						
2001	44.5	49.9	65.6	108.3	173.2	358.5
2000[4]	45.3	48.1	64.6	99.8	164.8	359.7
1999[4]	44.5	53.1	68.2	102.8	181.8	375.4
Female						
2001	19.6	25.5	36.3	60.9	104.8	242.0
2000[4]	20.4	24.5	36.4	58.5	100.9	238.4
1999[4]	19.2	26.4	37.3	61.8	105.0	244.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Accidents (unintentional injuries) (V01-X59,Y85-Y86)														
Hispanic														
Both sexes														
2001	25.8	18.8	10.3	5.0	5.7	27.8	37.0	30.2	26.1	31.6	33.1	32.6	30.5	29.7
2000[4]	25.0	15.3	10.4	5.7	5.6	27.9	35.5	27.9	26.6	30.3	33.1	30.5	30.2	28.3
1999[4]	25.5	17.7	11.9	4.9	6.4	26.1	35.3	28.7	28.0	31.0	34.6	32.5	29.6	27.4
Male														
2001	37.6	18.0	12.7	6.0	6.4	40.2	56.9	47.0	40.2	48.9	51.7	50.0	47.3	42.6
2000[4]	36.9	17.0	11.6	5.7	6.5	40.8	54.8	44.0	41.7	46.3	53.9	49.1	47.4	42.3
1999[4]	37.7	20.7	13.5	6.0	7.8	37.3	54.5	44.7	43.4	48.7	54.7	52.9	46.6	42.3
Female														
2001	13.2	19.7	7.8	4.0	5.0	14.0	12.6	10.2	10.0	12.6	13.5	15.2	14.6	18.1
2000[4]	12.4	13.5	9.2	5.6	4.7	13.1	11.9	9.0	9.4	12.9	11.3	12.0	13.8	15.7
1999[4]	12.6	14.6	10.3	3.7	4.9	13.4	12.2	9.9	10.3	11.6	13.7	12.2	13.5	14.0
Non-Hispanic[5]														
Both sexes														
2001	36.9	25.4	11.3	6.6	7.7	33.6	39.8	31.5	28.8	34.3	36.9	36.0	31.8	29.0
2000[4]	36.0	25.0	12.2	7.0	8.1	34.3	39.3	30.8	28.8	33.1	35.4	34.3	30.5	29.8
1999[4]	36.2	23.4	12.4	7.5	8.3	34.5	37.8	30.1	29.2	32.2	35.3	33.9	28.9	30.0
Male														
2001	48.5	28.0	13.4	7.9	10.0	46.1	61.4	48.1	43.4	49.3	54.0	54.0	46.5	41.4
2000[4]	47.3	27.7	14.5	8.3	10.5	46.6	59.6	47.4	43.1	49.0	52.5	51.4	45.4	42.8
1999[4]	47.4	26.6	14.5	9.0	10.8	46.1	57.3	46.1	43.6	46.9	52.5	51.1	42.9	43.1
Female														
2001	26.0	22.6	9.2	5.3	5.3	20.5	18.1	15.1	14.4	19.6	20.2	18.7	17.8	17.4
2000[4]	25.2	22.2	9.7	5.6	5.6	21.5	18.9	14.3	14.6	17.5	18.6	17.7	16.2	17.5
1999[4]	25.5	20.0	10.2	6.0	5.8	22.4	17.9	14.3	14.9	17.6	18.4	17.4	15.6	17.7
Non-Hispanic White														
Both sexes														
2001	38.5	20.9	10.9	5.9	7.3	36.0	42.2	32.9	29.6	34.9	36.8	34.7	30.8	28.1
2000[4]	37.3	20.1	11.1	6.3	8.0	37.3	41.4	31.1	29.1	33.4	35.0	33.0	28.7	28.6
1999[4]	37.2	19.0	11.1	6.6	8.1	37.2	39.1	30.8	29.1	32.1	34.8	32.3	27.4	28.6
Male														
2001	49.7	22.9	13.2	6.7	9.5	48.8	65.0	49.7	44.2	49.7	53.7	51.3	44.3	39.2
2000[4]	48.1	22.9	13.4	7.5	10.2	50.0	62.6	47.2	43.4	49.3	51.5	49.1	42.2	40.2
1999[4]	47.8	22.0	12.9	7.7	10.4	48.8	59.1	46.9	43.0	46.6	51.5	47.9	40.1	40.3
Female														
2001	27.8	18.7	8.5	5.0	5.1	22.5	18.9	15.8	14.7	20.0	19.9	18.3	17.5	17.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)						
Hispanic						
Both sexes						
2001	31.6	32.5	49.3	62.9	92.9	166.3
2000[4]	29.3	31.2	48.4	59.1	95.8	184.5
1999[4]	29.8	42.3	42.6	59.1	88.7	182.4
Male						
2001	48.3	46.1	66.3	92.8	128.6	198.1
2000[4]	44.3	47.0	70.5	86.4	109.0	251.9
1999[4]	45.0	62.6	64.2	78.0	128.9	258.8
Female						
2001	17.2	21.5	36.3	41.6	70.8	150.8
2000[4]	16.4	18.4	31.7	39.7	87.8	151.3
1999[4]	16.7	25.9	26.4	45.7	64.7	145.6
Non-Hispanic[5]						
Both sexes						
2001	31.2	36.9	49.1	81.1	131.5	279.8
2000[4]	32.3	35.5	48.6	76.0	125.4	276.1
1999[4]	31.1	38.3	51.2	79.2	134.3	285.0
Male						
2001	43.9	50.0	65.2	108.8	174.4	364.5
2000[4]	45.0	47.9	64.0	100.1	166.5	363.1
1999[4]	44.1	52.2	68.2	103.5	182.5	378.7
Female						
2001	19.7	25.6	36.2	61.8	105.8	244.6
2000[4]	20.6	24.9	36.5	59.2	101.1	240.9
1999[4]	19.4	26.3	37.9	62.2	106.1	247.5
Non-Hispanic White						
Both sexes						
2001	30.1	36.5	48.5	81.9	134.4	289.3
2000[4]	31.7	34.5	47.8	76.2	128.1	285.6
1999[4]	29.2	37.1	49.7	78.5	136.1	293.3
Male						
2001	41.8	48.8	64.2	108.7	176.3	374.7
2000[4]	43.3	45.4	62.5	98.8	168.3	373.8
1999[4]	41.3	49.9	65.4	101.3	184.2	387.1
Female						
2001	19.4	25.6	35.9	63.0	109.1	253.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Accidents (unintentional injuries) (V01-X59,Y85-Y86)														
Non-Hispanic White														
Female														
2000[4]	26.9	17.1	8.7	5.0	5.6	24.0	19.6	14.7	14.6	17.5	18.5	17.1	15.5	17.6
1999[4]	27.0	15.9	9.3	5.4	5.7	25.1	18.6	14.5	15.0	17.6	18.2	16.9	14.9	17.5
Non-Hispanic Black														
Both sexes														
2001	34.4	44.5	14.2	9.8	9.7	25.3	34.2	31.3	30.9	36.5	45.3	50.8	45.7	41.2
2000[4]	34.4	46.5	16.8	10.0	9.1	24.3	35.5	34.2	32.8	36.7	44.6	47.9	47.6	42.7
1999[4]	36.1	44.1	18.2	11.8	9.9	25.8	35.8	32.5	34.0	37.4	44.3	51.3	44.5	44.0
Male														
2001	49.5	50.4	15.8	13.0	13.2	37.0	52.9	49.5	48.0	54.3	68.4	83.0	73.2	67.0
2000[4]	50.3	49.5	19.4	12.2	12.5	35.3	54.5	55.9	51.1	56.9	69.9	76.2	77.4	72.6
1999[4]	52.3	48.5	21.8	15.0	13.5	37.9	56.5	52.7	53.4	56.5	70.0	84.1	73.2	72.8
Female														
2001	20.6	38.3	12.4	6.4	6.0	13.3	16.3	14.6	15.5	20.7	24.9	23.0	22.3	20.0
2000[4]	20.1	43.3	14.2	7.8	5.5	13.0	17.5	14.6	16.5	18.8	22.3	23.5	22.3	18.3
1999[4]	21.4	39.6	14.5	8.4	6.3	13.4	16.3	14.3	16.6	20.5	21.7	23.0	20.2	20.5
Transport accidents (V01-V99,Y85)														
All origins[3]														
Both sexes														
2001	16.6	3.7	4.4	3.8	5.0	26.6	29.2	20.9	16.9	17.4	17.3	16.3	15.6	15.2
2000[4]	16.6	4.6	4.5	4.1	5.3	27.1	29.3	20.4	16.8	17.1	16.8	16.2	15.5	15.7
1999[4]	16.6	5.0	4.6	4.4	5.3	27.1	28.2	20.0	17.0	16.9	16.6	15.9	14.9	16.3
Male														
2001	23.5	3.3	5.0	4.3	6.2	35.4	43.9	31.4	25.4	24.7	25.4	24.0	22.5	21.4
2000[4]	23.3	4.8	5.0	4.4	6.4	35.5	43.3	30.8	24.7	24.8	24.8	23.6	22.3	22.4
1999[4]	23.1	5.2	5.1	5.0	6.5	34.8	41.8	29.9	24.7	24.0	24.0	23.1	21.5	23.0
Female														
2001	10.0	4.2	3.8	3.3	3.8	17.3	13.8	10.1	8.3	10.2	9.4	8.8	8.9	9.4
2000[4]	10.2	4.3	4.0	3.8	4.1	18.1	14.6	9.8	8.7	9.5	9.0	8.9	8.9	9.5
1999[4]	10.4	4.8	4.2	3.6	4.1	19.0	14.0	10.0	9.3	9.8	9.4	8.9	8.6	10.0
Hispanic														
Both sexes														
2001	15.6	5.3	5.3	3.3	4.3	22.7	29.0	21.6	15.9	17.3	17.3	16.3	16.5	17.1
2000[4]	14.8	5.4	5.2	4.0	4.3	21.9	27.0	19.4	15.0	15.6	16.9	15.8	16.7	15.9
1999[4]	14.4	5.3	5.5	3.1	4.7	20.0	25.9	19.6	15.9	14.4	16.6	14.6	15.5	16.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)						
Non-Hispanic White						
Female						
2000[4]	20.9	24.9	35.9	60.3	104.3	250.0
1999[4]	18.0	25.9	37.1	62.5	107.7	255.9
Non-Hispanic Black						
Both sexes						
2001	43.9	45.9	58.0	79.2	105.7	183.9
2000[4]	39.6	46.8	60.4	75.9	100.3	183.2
1999[4]	46.4	51.3	67.5	90.7	125.0	208.9
Male						
2001	69.8	69.8	81.6	121.4	167.6	256.4
2000[4]	65.3	75.2	87.1	120.4	156.2	277.1
1999[4]	70.5	78.9	103.5	136.3	178.0	318.5
Female						
2001	23.5	28.3	42.2	54.0	74.1	157.0
2000[4]	19.3	26.0	42.7	49.1	72.2	148.3
1999[4]	27.5	30.9	43.7	63.2	98.8	167.9
Transport accidents (V01-V99,Y85)						
All origins[3]						
Both sexes						
2001	15.4	16.4	19.5	26.2	30.4	28.9
2000[4]	16.4	16.8	19.4	25.1	30.0	31.5
1999[4]	15.7	18.1	20.9	25.9	32.1	32.0
Male						
2001	22.0	22.1	26.8	36.2	45.5	52.7
2000[4]	22.6	22.6	26.3	34.5	44.0	59.8
1999[4]	21.8	24.8	27.5	36.2	48.3	63.5
Female						
2001	9.3	11.6	13.8	19.2	21.3	18.9
2000[4]	10.7	11.7	13.9	18.5	21.7	19.9
1999[4]	10.2	12.4	15.7	18.8	22.6	19.3
Hispanic						
Both sexes						
2001	19.0	18.1	22.8	28.0	33.8	23.8
2000[4]	15.3	16.4	25.6	23.3	25.1	25.2
1999[4]	14.5	22.8	21.0	24.5	25.8	25.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Transport accidents (V01-V99,Y85)														
Hispanic														
Male														
2001	22.8	*	6.0	3.8	4.9	31.7	44.1	33.2	24.1	25.8	26.0	22.4	24.1	21.4
2000[4]	21.6	6.1	5.6	3.8	5.2	31.3	40.4	30.2	22.1	22.9	25.6	23.0	25.4	23.5
1999[4]	20.8	5.7	5.8	3.6	5.6	27.7	38.5	29.7	23.5	21.6	25.2	22.0	23.4	21.7
Female														
2001	8.0	6.7	4.6	2.7	3.7	12.7	10.5	7.8	6.4	7.9	8.2	10.1	9.3	13.2
2000[4]	7.6	*	4.8	4.1	3.3	11.3	10.7	6.8	6.8	7.8	7.7	8.7	8.3	9.1
1999[4]	7.6	*	5.2	2.5	3.8	11.3	10.7	7.9	7.2	6.5	7.7	7.3	8.1	11.3
Non-Hispanic[5]														
Both sexes														
2001	16.7	3.4	4.2	3.9	5.2	27.2	29.2	20.7	17.1	17.4	17.2	16.2	15.4	15.0
2000[4]	16.8	4.3	4.3	4.1	5.4	27.9	29.7	20.5	17.0	17.3	16.7	16.1	15.2	15.6
1999[4]	16.9	4.9	4.4	4.6	5.4	28.3	28.5	20.0	17.2	17.2	16.5	15.9	14.8	16.2
Male														
2001	23.5	3.2	4.7	4.4	6.5	36.0	43.7	30.9	25.6	24.4	25.2	24.0	22.3	21.3
2000[4]	23.4	4.5	4.8	4.5	6.6	36.2	43.9	30.8	25.1	24.9	24.5	23.6	21.9	22.2
1999[4]	23.3	5.1	4.9	5.3	6.6	36.0	42.3	29.7	24.8	24.3	23.7	23.1	21.3	22.9
Female														
2001	10.2	3.6	3.6	3.4	3.8	18.0	14.4	10.5	8.6	10.5	9.5	8.6	8.9	9.0
2000[4]	10.5	4.2	3.8	3.7	4.2	19.3	15.3	10.4	9.0	9.8	9.1	8.9	8.9	9.5
1999[4]	10.7	4.7	3.9	3.9	4.2	20.3	14.5	10.3	9.6	10.2	9.5	9.0	8.6	9.9
Non-Hispanic White														
Both sexes														
2001	17.0	3.1	3.8	3.6	5.2	29.6	30.3	21.3	17.1	17.5	17.3	15.7	15.4	14.8
2000[4]	17.1	3.5	3.8	4.0	5.6	30.9	30.9	20.2	17.0	17.4	16.6	15.9	14.8	15.4
1999[4]	17.1	4.3	3.7	4.4	5.6	31.1	29.5	20.2	17.2	16.9	16.5	15.6	14.5	15.8
Male														
2001	23.8	2.9	4.4	4.0	6.4	38.6	45.5	31.5	25.4	24.2	25.0	23.1	22.2	20.8
2000[4]	23.7	3.8	4.3	4.5	6.8	39.6	45.6	29.7	24.8	24.8	24.1	23.2	21.1	21.4
1999[4]	23.5	4.9	4.0	4.9	6.8	39.0	43.7	29.8	24.3	23.7	23.5	22.3	20.7	22.3
Female														
2001	10.6	3.3	3.2	3.2	3.8	20.0	14.8	10.9	8.7	10.8	9.6	8.5	8.8	9.0
2000[4]	10.9	3.2	3.4	3.5	4.4	21.8	15.9	10.5	9.0	10.0	9.1	8.7	8.5	9.6
1999[4]	11.0	3.6	3.4	3.7	4.4	22.8	15.0	10.5	9.9	10.2	9.6	8.9	8.4	9.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Transport accidents (V01-V99,Y85)						
Hispanic						
Male						
2001	29.1	25.9	31.1	45.7	61.0	43.8
2000[4]	23.6	23.5	37.4	39.1	30.9	44.3
1999[4]	20.0	32.8	30.0	35.9	48.7	*
Female						
2001	10.3	11.8	16.4	15.5	16.9	*
2000[4]	8.2	10.6	16.6	12.0	21.5	*
1999[4]	9.8	14.7	14.2	16.5	*	*
Non-Hispanic[5]						
Both sexes						
2001	15.0	16.3	19.2	26.0	30.1	29.0
2000[4]	16.3	16.7	18.9	25.1	30.0	31.7
1999[4]	15.7	17.7	20.8	25.9	32.0	32.2
Male						
2001	21.3	21.8	26.4	35.7	44.6	52.9
2000[4]	22.3	22.5	25.5	34.3	44.3	60.4
1999[4]	21.8	24.2	27.2	36.1	47.7	64.3
Female						
2001	9.3	11.5	13.5	19.3	21.4	19.1
2000[4]	10.9	11.8	13.8	18.8	21.7	20.1
1999[4]	10.2	12.2	15.7	18.8	22.9	19.3
Non-Hispanic White						
Both sexes						
2001	14.7	16.2	19.2	26.5	30.8	29.9
2000[4]	16.4	16.5	18.7	25.3	30.9	32.4
1999[4]	15.3	17.6	20.7	26.0	32.4	32.5
Male						
2001	20.7	21.4	26.2	35.7	45.7	54.3
2000[4]	21.9	21.6	24.9	34.0	44.8	62.3
1999[4]	21.0	23.8	26.7	35.7	47.8	64.6
Female						
2001	9.1	11.6	13.5	20.0	21.9	19.8
2000[4]	11.3	12.0	13.7	19.2	22.6	20.4
1999[4]	10.1	12.1	15.8	19.1	23.3	19.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Transport accidents (V01-V99,Y85)														
Non-Hispanic Black														
Both sexes														
2001	16.3	*	5.4	5.5	5.4	19.3	26.7	21.6	19.6	18.5	19.8	20.7	17.2	18.1
2000[4]	16.4	6.9	5.9	4.6	5.0	17.8	27.7	24.4	20.0	18.4	19.7	19.8	19.6	18.4
1999[4]	16.7	7.7	7.1	6.0	5.2	19.2	26.6	22.0	19.2	20.5	18.4	20.0	18.3	19.1
Male														
2001	24.5	*	5.8	6.9	7.3	27.2	40.5	34.1	31.1	27.9	31.5	34.5	26.1	29.3
2000[4]	24.8	7.2	6.4	5.0	6.7	24.6	41.7	39.5	31.7	29.5	31.6	30.0	30.3	31.5
1999[4]	24.9	*	8.0	7.3	6.6	26.7	40.3	35.5	30.5	30.7	28.8	31.8	29.2	29.8
Female														
2001	8.9	*	5.0	4.1	3.4	11.1	13.5	10.1	9.2	10.2	9.5	8.8	9.6	8.8
2000[4]	8.8	*	5.4	4.2	3.2	10.8	14.4	10.7	9.6	8.6	9.2	11.0	10.5	7.8
1999[4]	9.4	9.3	6.2	4.6	3.7	11.7	13.7	9.8	9.0	11.5	9.2	9.7	9.1	10.4
Motor vehicle accidents (V02-V04, V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5, V81.0-V81.1,V82.0-V82.1,V83-V86, V87.0-V87.8,V88.0-V88.8,V89.0, V89.2)														
All origins[3]														
Both sexes														
2001	15.4	3.6	4.1	3.6	4.6	25.7	28.0	19.6	15.6	15.9	15.5	14.5	13.9	13.7
2000[4]	15.4	4.4	4.2	3.8	4.8	26.0	28.0	19.2	15.5	15.6	15.1	14.6	13.8	14.2
1999[4]	15.2	4.8	4.2	3.9	4.8	25.9	26.5	18.4	15.4	15.1	14.8	14.0	13.1	14.4
Male														
2001	21.5	3.2	4.5	4.0	5.7	34.0	42.1	29.5	23.1	22.3	22.5	21.2	19.7	18.9
2000[4]	21.3	4.6	4.7	4.1	5.8	33.9	41.2	28.6	22.6	22.1	21.9	21.1	19.6	19.9
1999[4]	20.9	5.0	4.7	4.4	5.7	33.1	39.1	27.2	22.2	21.2	21.0	20.0	18.6	19.8
Female														
2001	9.5	4.0	3.6	3.0	3.5	16.9	13.4	9.5	8.0	9.6	8.7	8.0	8.3	8.8
2000[4]	9.7	4.2	3.8	3.5	3.8	17.5	14.2	9.5	8.2	9.1	8.4	8.3	8.2	9.0
1999[4]	9.7	4.7	3.8	3.3	3.8	18.3	13.4	9.4	8.5	9.0	8.6	8.1	8.0	9.4
Hispanic														
Both sexes														
2001	14.3	4.9	4.7	3.0	4.0	22.1	27.7	20.2	14.5	15.4	15.2	13.8	13.8	14.4
2000[4]	13.9	5.2	4.9	3.7	4.0	21.2	26.0	18.5	14.0	14.4	15.2	14.8	15.4	14.7
1999[4]	13.3	5.3	5.2	2.9	4.4	19.1	24.5	18.2	14.5	13.0	15.0	13.3	14.1	15.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Transport accidents (V01-V99,Y85)						
Non-Hispanic Black						
Both sexes						
2001	18.9	18.9	18.7	22.3	19.9	17.8
2000[4]	16.4	18.6	21.5	19.4	20.7	21.9
1999[4]	17.8	18.2	21.1	24.7	28.2	27.8
Male						
2001	31.3	28.9	25.5	39.8	31.9	35.8
2000[4]	27.2	31.7	33.8	34.7	39.5	45.8
1999[4]	28.7	27.4	32.2	40.6	48.7	68.9
Female						
2001	9.2	11.5	14.2	11.9	13.7	11.1
2000[4]	7.9	8.8	13.4	10.2	11.2	13.1
1999[4]	9.2	11.4	13.8	15.1	18.0	12.4
Motor vehicle accidents (V02-V04, V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5, V81.0-V81.1,V82.0-V82.1,V83-V86, V87.0-V87.8,V88.0-V88.8,V89.0, V89.2)						
All origins[3]						
Both sexes						
2001	13.7	14.8	18.0	24.6	28.9	27.8
2000[4]	14.7	15.1	18.1	23.8	28.5	30.4
1999[4]	13.9	16.4	19.3	24.1	30.1	30.4
Male						
2001	19.2	19.2	24.2	33.6	43.1	51.1
2000[4]	19.7	19.8	23.9	32.2	41.4	57.5
1999[4]	18.7	22.0	24.7	33.6	45.2	59.8
Female						
2001	8.8	11.0	13.1	18.2	20.4	18.0
2000[4]	10.2	11.0	13.5	17.9	20.9	19.3
1999[4]	9.5	11.6	15.0	17.5	21.3	18.5
Hispanic						
Both sexes						
2001	16.0	15.2	20.8	23.5	31.3	20.0
2000[4]	14.0	14.2	23.7	21.7	23.4	22.6
1999[4]	13.4	20.6	18.6	23.2	24.1	23.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Motor vehicle accidents (V02-V04, V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5, V81.0-V81.1,V82.0-V82.1,V83-V86, V87.0-V87.8,V88.0-V88.8,V89.0, V89.2)														
Hispanic														
Male														
2001	21.0	*	5.3	3.6	4.5	30.8	42.1	31.1	22.2	22.9	23.4	19.9	20.5	18.6
2000[4]	20.1	6.1	5.2	3.5	4.8	30.2	38.7	28.6	20.7	20.6	22.9	21.0	23.3	21.5
1999[4]	19.1	5.7	5.6	3.3	5.0	26.4	36.4	27.3	21.4	19.4	22.7	19.7	20.6	20.1
Female														
2001	7.2	6.2	4.1	2.5	3.5	12.5	10.1	7.1	5.8	7.1	6.6	7.7	7.4	10.7
2000[4]	7.2	*	4.6	4.0	3.1	10.9	10.4	6.7	6.4	7.6	7.1	8.6	7.9	8.5
1999[4]	7.2	*	4.7	2.4	3.7	10.9	10.3	7.5	6.6	6.0	7.0	7.0	7.9	11.3
Non-Hispanic[5]														
Both sexes														
2001	15.5	3.2	3.9	3.7	4.7	26.3	28.0	19.4	15.7	16.0	15.5	14.5	13.8	13.6
2000[4]	15.6	4.2	4.1	3.8	5.0	26.8	28.3	19.2	15.6	15.7	15.0	14.5	13.6	14.1
1999[4]	15.4	4.7	4.0	4.1	4.9	27.0	26.8	18.3	15.5	15.3	14.7	14.0	13.0	14.2
Male														
2001	21.5	3.1	4.3	4.1	5.9	34.5	41.9	29.0	23.2	22.1	22.3	21.2	19.5	18.9
2000[4]	21.4	4.2	4.5	4.2	6.0	34.5	41.6	28.5	22.8	22.3	21.6	21.1	19.2	19.6
1999[4]	21.0	4.8	4.4	4.6	5.8	34.2	39.6	27.1	22.3	21.4	20.7	19.9	18.3	19.5
Female														
2001	9.8	3.4	3.5	3.2	3.5	17.6	14.0	9.9	8.3	9.9	8.9	8.0	8.3	8.6
2000[4]	10.0	4.2	3.6	3.4	3.9	18.7	14.9	10.0	8.5	9.3	8.5	8.2	8.2	9.0
1999[4]	10.0	4.7	3.6	3.5	3.8	19.5	13.9	9.7	8.8	9.4	8.8	8.2	7.9	9.2
Non-Hispanic White														
Both sexes														
2001	15.8	3.0	3.5	3.4	4.7	28.5	29.0	19.9	15.7	16.0	15.5	14.1	13.8	13.4
2000[4]	15.8	3.3	3.7	3.7	5.1	29.7	29.4	18.7	15.5	15.8	14.8	14.3	13.0	13.9
1999[4]	15.6	4.1	3.4	3.8	5.0	29.7	27.7	18.4	15.4	15.0	14.6	13.6	12.7	13.8
Male														
2001	21.7	2.9	3.9	3.7	5.9	36.9	43.5	29.3	22.9	21.8	22.0	20.3	19.4	18.5
2000[4]	21.5	3.4	4.1	4.2	6.1	37.9	43.0	27.2	22.4	22.1	21.1	20.7	18.3	18.9
1999[4]	21.1	4.7	3.6	4.3	5.9	37.0	40.8	26.9	21.8	20.7	20.4	19.1	17.7	18.8
Female														
2001	10.1	3.1	3.1	3.0	3.5	19.6	14.2	10.3	8.4	10.2	8.9	8.0	8.2	8.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Motor vehicle accidents (V02-V04, V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5, V81.0-V81.1,V82.0-V82.1,V83-V86, V87.0-V87.8,V88.0-V88.8,V89.0, V89.2)						
Hispanic						
Male						
2001	24.7	21.6	28.4	38.2	55.8	*
2000[4]	20.7	20.1	33.5	36.9	*	*
1999[4]	17.6	29.0	25.0	32.8	45.6	*
Female						
2001	8.5	10.0	15.0	13.1	16.1	*
2000[4]	8.2	9.4	16.2	11.0	21.5	*
1999[4]	9.8	13.8	13.8	16.5	*	*
Non-Hispanic[5]						
Both sexes						
2001	13.5	14.7	17.7	24.6	28.7	28.0
2000[4]	14.7	15.1	17.7	23.9	28.6	30.6
1999[4]	13.9	16.1	19.2	24.1	30.1	30.5
Male						
2001	18.6	19.0	23.8	33.3	42.4	51.6
2000[4]	19.5	19.7	23.2	32.0	41.7	58.3
1999[4]	18.7	21.5	24.6	33.5	44.7	60.7
Female						
2001	8.8	11.0	12.9	18.4	20.4	18.2
2000[4]	10.3	11.1	13.3	18.2	20.9	19.4
1999[4]	9.5	11.4	15.1	17.5	21.6	18.5
Non-Hispanic White						
Both sexes						
2001	13.1	14.7	17.6	25.0	29.4	29.0
2000[4]	14.7	14.9	17.4	24.1	29.3	31.3
1999[4]	13.5	15.9	19.1	24.2	30.6	30.9
Male						
2001	17.8	18.7	23.5	33.3	43.5	53.1
2000[4]	18.9	18.8	22.7	31.7	42.0	60.0
1999[4]	18.0	21.1	24.2	33.2	44.9	61.2
Female						
2001	8.8	11.1	12.8	19.2	21.0	19.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Motor vehicle accidents (V02-V04, V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5, V81.0-V81.1,V82.0-V82.1,V83-V86, V87.0-V87.8,V88.0-V88.8,V89.0, V89.2)														
Non-Hispanic White														
Female														
2000[4]	10.3	3.2	3.2	3.2	4.1	21.1	15.4	10.1	8.5	9.5	8.5	8.0	7.8	9.1
1999[4]	10.3	3.5	3.1	3.3	4.0	22.0	14.3	9.8	9.0	9.3	8.7	8.1	7.7	9.0
Non-Hispanic Black														
Both sexes														
2001	15.2	*	5.3	5.1	4.9	18.6	26.1	20.7	18.5	17.2	18.1	18.5	15.7	16.2
2000[4]	15.5	6.9	5.6	4.3	4.7	16.9	26.9	23.7	18.6	16.9	18.3	18.0	18.6	17.0
1999[4]	15.5	7.6	6.5	5.5	4.8	18.3	25.6	20.9	17.6	18.8	16.9	18.2	16.4	17.4
Male														
2001	22.7	*	5.6	6.5	6.5	26.1	39.4	32.7	29.3	25.6	28.2	30.6	23.7	25.3
2000[4]	23.1	7.2	6.0	4.5	6.4	23.2	40.3	38.0	29.3	26.9	29.0	27.0	28.5	28.6
1999[4]	22.8	*	7.0	6.5	6.0	25.3	38.8	33.6	28.0	27.8	26.0	28.8	25.8	26.7
Female														
2001	8.5	*	4.9	3.7	3.2	10.8	13.4	9.7	8.8	9.7	9.1	8.1	8.9	8.7
2000[4]	8.5	*	5.2	4.1	3.1	10.5	14.2	10.6	9.0	8.2	8.8	10.3	10.2	7.5
1999[4]	8.9	9.3	5.9	4.5	3.6	11.2	13.1	9.4	8.4	10.8	8.9	9.0	8.5	9.8
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9, V10-V11,V15-V18,V19.3, V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9, V82.2-V82.9,V87.9,V88.9,V89.1, V89.3,V89.9)														
All origins[3]														
Both sexes														
2001	0.5	*	0.2	0.1	0.3	0.5	0.5	0.4	0.4	0.6	0.6	0.7	0.6	0.5
2000[4]	0.5	*	0.2	0.2	0.3	0.6	0.6	0.5	0.6	0.7	0.7	0.6	0.6	0.5
1999[4]	0.8	*	0.3	0.3	0.3	0.9	1.0	0.8	0.8	0.9	0.9	0.9	0.7	0.8
Male														
2001	0.8	*	0.3	*	0.3	0.7	0.9	0.7	0.8	1.1	1.0	1.0	0.9	0.8
2000[4]	0.8	*	*	0.2	0.4	1.0	1.0	0.8	0.9	1.1	1.2	0.9	0.9	0.7
1999[4]	1.1	*	0.3	0.4	0.5	1.2	1.7	1.3	1.1	1.4	1.5	1.4	1.2	1.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Motor vehicle accidents (V02-V04, V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5, V81.0-V81.1,V82.0-V82.1,V83-V86, V87.0-V87.8,V88.0-V88.8,V89.0, V89.2)						
Non-Hispanic White						
Female						
2000[4]	10.7	11.4	13.2	18.7	21.8	19.8
1999[4]	9.3	11.4	15.1	17.9	22.1	18.8
Non-Hispanic Black						
Both sexes						
2001	17.2	16.8	18.0	20.4	19.0	16.9
2000[4]	15.5	16.9	20.3	18.1	19.8	20.7
1999[4]	15.8	16.7	19.7	22.9	25.8	25.5
Male						
2001	28.8	24.5	23.8	37.0	31.1	35.8
2000[4]	25.5	28.2	31.8	32.3	38.6	44.6
1999[4]	24.8	25.0	29.1	39.2	45.2	61.8
Female						
2001	8.1	11.2	14.2	10.5	12.9	9.8
2000[4]	7.5	8.5	12.7	9.6	10.4	11.8
1999[4]	8.7	10.6	13.6	13.1	16.2	12.0
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9, V10-V11,V15-V18,V19.3, V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9, V82.2-V82.9,V87.9,V88.9,V89.1, V89.3,V89.9)						
All origins[3]						
Both sexes						
2001	0.6	0.7	0.6	0.8	0.8	0.7
2000[4]	0.5	0.5	0.5	0.7	0.8	0.8
1999[4]	0.8	0.8	0.7	1.3	1.4	1.3
Male						
2001	0.9	1.1	0.9	1.2	*	*
2000[4]	1.0	0.9	0.9	1.1	1.3	*
1999[4]	1.1	1.2	1.2	1.7	2.1	2.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9, V10-V11,V15-V18,V19.3, V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9, V82.2-V82.9,V87.9,V88.9,V89.1, V89.3,V89.9)														
All origins[3]														
Female														
2001	0.2	*	*	*	*	0.2	*	*	*	0.2	0.2	0.4	0.2	*
2000[4]	0.3	*	*	*	*	0.3	0.2	*	0.3	0.2	0.3	0.3	0.4	*
1999[4]	0.4	*	0.3	*	*	0.5	0.3	0.3	0.5	0.4	0.4	0.4	0.3	0.4
Hispanic														
Both sexes														
2001	0.5	*	*	*	*	*	0.7	0.7	*	0.7	0.9	*	*	*
2000[4]	0.6	*	*	*	*	*	*	*	0.7	*	1.1	*	*	*
1999[4]	0.8	*	*	*	*	0.7	1.1	0.9	1.0	1.0	1.3	*	*	*
Male														
2001	0.9	*	*	*	*	*	1.3	1.2	*	*	1.6	*	*	*
2000[4]	0.9	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.3	*	*	*	*	*	1.6	1.6	1.5	1.7	2.0	*	*	*
Female														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.5	*	*	0.1	0.3	0.5	0.5	0.4	0.4	0.6	0.6	0.6	0.5	0.4
2000[4]	0.5	*	*	0.2	0.3	0.7	0.6	0.5	0.6	0.6	0.7	0.6	0.6	0.4
1999[4]	0.7	*	0.3	0.3	0.3	0.9	1.0	0.7	0.7	0.9	0.9	0.9	0.7	0.8
Male														
2001	0.7	*	*	*	0.4	0.7	0.8	0.6	0.8	1.0	1.0	1.0	0.9	0.8
2000[4]	0.8	*	*	*	0.4	1.0	1.0	0.8	0.8	1.0	1.2	0.9	0.8	0.7
1999[4]	1.1	*	0.4	0.4	0.5	1.2	1.7	1.2	1.0	1.3	1.4	1.4	1.1	1.3
Female														
2001	0.2	*	*	*	*	0.3	*	*	*	0.2	0.2	0.3	0.2	*
2000[4]	0.3	*	*	*	*	0.3	*	*	0.3	0.3	0.2	0.3	0.4	*
1999[4]	0.4	*	*	*	*	0.6	0.3	0.3	0.5	0.4	0.3	0.4	0.3	0.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9, V10-V11,V15-V18,V19.3, V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9, V82.2-V82.9,V87.9,V88.9,V89.1, V89.3,V89.9)						
All origins[3]						
Female						
2001	*	*	*	0.5	0.7	*
2000[4]	*	*	*	*	*	*
1999[4]	0.4	0.5	0.4	0.9	1.0	*
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	0.6	0.7	0.6	0.7	0.7	0.6
2000[4]	0.5	0.5	0.5	0.7	0.8	0.8
1999[4]	0.7	0.7	0.7	1.3	1.3	1.3
Male						
2001	0.8	1.2	0.9	1.0	*	*
2000[4]	0.9	0.8	0.9	1.1	1.2	*
1999[4]	1.1	1.0	1.0	1.6	2.0	3.0
Female						
2001	*	*	*	*	0.6	*
2000[4]	*	*	*	*	*	*
1999[4]	0.4	0.5	0.4	1.0	1.0	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9, V10-V11,V15-V18,V19.3, V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9, V82.2-V82.9,V87.9,V88.9,V89.1, V89.3,V89.9)														
Non-Hispanic White														
Both sexes														
2001	0.5	*	*	*	0.3	0.5	0.5	0.4	0.4	0.5	0.5	0.6	0.5	0.4
2000[4]	0.5	*	*	0.2	0.3	0.7	0.6	0.5	0.5	0.6	0.6	0.5	0.6	0.4
1999[4]	0.8	*	0.3	0.3	0.4	0.9	1.0	0.8	0.7	0.9	0.9	0.9	0.7	0.8
Male														
2001	0.7	*	*	*	0.3	0.7	0.8	0.7	0.7	0.9	0.8	0.9	0.9	0.7
2000[4]	0.8	*	*	*	0.4	1.1	1.0	0.8	0.8	1.0	1.1	0.8	0.8	0.5
1999[4]	1.1	*	*	0.4	0.5	1.2	1.6	1.2	1.0	1.3	1.4	1.3	1.0	1.3
Female														
2001	0.2	*	*	*	*	*	*	*	*	*	*	0.3	*	*
2000[4]	0.3	*	*	*	*	0.4	*	*	*	*	*	0.3	0.4	*
1999[4]	0.4	*	*	*	*	0.6	0.4	0.3	0.5	0.4	0.4	0.5	0.3	0.4
Non-Hispanic Black														
Both sexes														
2001	0.6	*	*	*	*	*	*	*	*	1.0	1.2	1.1	*	*
2000[4]	0.6	*	*	*	*	*	*	*	0.8	0.9	1.0	1.0	*	*
1999[4]	0.7	*	*	*	*	*	*	*	0.9	1.0	0.9	1.0	*	*
Male														
2001	0.9	*	*	*	*	*	*	*	*	1.6	2.1	1.8	*	*
2000[4]	0.9	*	*	*	*	*	*	*	*	1.6	1.7	*	*	*
1999[4]	1.2	*	*	*	*	*	*	*	*	1.6	*	*	*	*
Female														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)														
All origins[3]														
Both sexes														
2001	0.7	*	0.2	0.1	0.1	0.4	0.6	0.8	0.9	0.9	1.2	1.1	1.2	1.0
2000[4]	0.7	*	*	0.1	0.1	0.5	0.7	0.8	0.8	0.9	1.0	0.9	1.0	1.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9, V10-V11,V15-V18,V19.3, V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9, V82.2-V82.9,V87.9,V88.9,V89.1, V89.3,V89.9)						
Non-Hispanic White						
Both sexes						
2001	0.5	0.6	0.6	0.7	0.7	0.5
2000[4]	0.5	0.4	0.5	0.6	0.8	0.8
1999[4]	0.7	0.7	0.7	1.2	1.3	1.3
Male						
2001	0.8	1.0	0.9	1.0	*	*
2000[4]	1.0	0.6	0.9	1.1	1.3	*
1999[4]	1.1	1.0	0.9	1.6	1.9	2.8
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	0.5	*	*	0.9	1.0	*
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)						
All origins[3]						
Both sexes						
2001	1.1	1.0	0.9	0.8	0.7	0.5
2000[4]	1.1	1.2	0.7	0.6	0.7	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)														
All origins[3]														
Both sexes														
1999[4]	0.7	*	*	0.2	0.2	0.4	0.6	0.8	0.8	0.9	0.9	1.0	1.0	1.1
Male														
2001	1.2	*	*	*	0.2	0.7	1.0	1.2	1.5	1.4	1.8	1.8	1.9	1.7
2000[4]	1.2	*	*	*	0.2	0.6	1.2	1.3	1.3	1.5	1.7	1.6	1.8	1.8
1999[4]	1.1	*	*	0.2	0.3	0.6	0.9	1.4	1.3	1.4	1.5	1.7	1.8	2.0
Female														
2001	0.3	*	*	*	*	*	0.3	0.4	0.3	0.4	0.5	0.4	0.4	0.4
2000[4]	0.2	*	*	*	*	0.3	*	*	0.2	0.2	0.3	0.3	0.3	*
1999[4]	0.3	*	*	*	*	0.2	0.3	0.3	0.3	0.4	0.4	0.3	0.3	*
Hispanic														
Both sexes														
2001	0.8	*	*	*	*	*	0.6	0.8	0.9	1.2	1.2	1.5	2.0	2.0
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.9	*	*	*	*	*	*	*	1.2	1.7	*	*	*	*
2000[4]	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.7	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.7	*	*	0.1	0.1	0.5	0.7	0.8	0.9	0.8	1.1	1.1	1.1	1.0
2000[4]	0.7	*	*	0.1	0.2	0.5	0.7	0.8	0.9	0.9	1.0	1.0	1.1	1.1
1999[4]	0.7	*	*	0.2	0.2	0.4	0.7	0.9	0.9	1.0	1.0	1.0	1.1	1.2
Male														
2001	1.2	*	*	*	*	0.8	1.0	1.3	1.5	1.3	1.9	1.9	1.9	1.7
2000[4]	1.2	*	*	*	*	0.7	1.3	1.5	1.5	1.6	1.8	1.7	1.9	1.9
1999[4]	1.2	*	*	*	0.4	0.7	1.0	1.5	1.5	1.5	1.6	1.8	1.9	2.1
Female														
2001	0.2	*	*	*	*	*	0.3	0.4	*	0.3	0.4	0.3	0.3	*
2000[4]	0.2	*	*	*	*	0.3	*	*	0.2	0.2	0.3	0.3	0.3	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)						
All origins[3]						
Both sexes						
1999[4]	1.1	0.9	0.8	0.5	0.6	*
Male						
2001	1.9	1.8	1.7	1.3	1.4	*
2000[4]	1.9	2.0	1.4	1.2	1.4	*
1999[4]	1.9	1.6	1.6	0.9	*	*
Female						
2001	*	*	*	0.5	*	*
2000[4]	0.4	0.4	*	*	*	*
1999[4]	*	*	*	*	*	*
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	1.0	0.9	0.9	0.8	0.7	*
2000[4]	1.1	1.2	0.7	0.6	0.7	*
1999[4]	1.1	0.9	0.9	0.5	0.6	*
Male						
2001	1.9	1.6	1.7	1.3	1.3	*
2000[4]	1.9	2.1	1.4	1.2	1.3	*
1999[4]	2.0	1.7	1.6	0.9	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	0.4	0.5	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)														
Non-Hispanic[5]														
Female														
1999[4]	0.3	*	*	*	*	*	0.3	0.3	0.4	0.4	0.4	0.3	0.4	*
Non-Hispanic White														
Both sexes														
2001	0.8	*	*	*	0.2	0.6	0.8	1.0	1.0	0.9	1.3	1.1	1.1	0.9
2000[4]	0.8	*	*	*	0.2	0.5	0.9	1.0	0.9	1.0	1.2	1.1	1.2	1.1
1999[4]	0.8	*	*	0.2	0.3	0.5	0.8	1.0	1.0	1.1	1.1	1.1	1.1	1.2
Male														
2001	1.3	*	*	*	*	1.0	1.2	1.5	1.7	1.5	2.1	1.9	2.0	1.6
2000[4]	1.4	*	*	*	*	0.7	1.6	1.7	1.6	1.7	2.0	1.7	2.0	2.0
1999[4]	1.3	*	*	*	0.4	0.7	1.2	1.7	1.6	1.6	1.7	1.9	1.9	2.1
Female														
2001	0.3	*	*	*	*	*	0.3	0.5	*	0.4	0.4	*	0.3	*
2000[4]	0.3	*	*	*	*	0.3	*	*	*	*	0.4	0.4	*	*
1999[4]	0.3	*	*	*	*	*	0.3	0.4	0.4	0.5	0.5	0.4	0.3	*
Non-Hispanic Black														
Both sexes														
2001	0.5	*	*	*	*	*	*	*	*	*	*	1.1	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	*	*	*	*	*	*	*	0.7	*	*	*	*
Male														
2001	0.8	*	*	*	*	*	*	*	*	*	*	2.1	*	*
2000[4]	0.7	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.9	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Nontransport accidents (W00-X59,Y86)														
All origins[3]														
Both sexes														
2001	19.0	20.5	6.8	2.5	2.4	6.2	10.3	10.5	11.6	16.7	19.4	19.7	16.3	14.1
2000[4]	18.2	18.6	7.4	2.7	2.5	6.3	9.5	10.0	11.8	15.8	18.6	18.0	15.3	14.2
1999[4]	18.4	17.3	7.7	2.7	2.7	6.2	9.3	10.0	12.1	15.3	18.8	18.2	14.3	13.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)						
Non-Hispanic[5]						
Female						
1999[4]	*	*	*	*	*	*
Non-Hispanic White						
Both sexes						
2001	1.1	0.9	0.9	0.8	0.7	*
2000[4]	1.2	1.2	0.7	0.6	0.7	*
1999[4]	1.1	1.0	0.9	0.5	0.5	*
Male						
2001	2.0	1.6	1.8	1.3	1.4	*
2000[4]	2.0	2.1	1.4	1.2	1.4	*
1999[4]	2.0	1.7	1.6	1.0	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	0.5	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Nontransport accidents (W00-X59,Y86)						
All origins[3]						
Both sexes						
2001	16.1	20.3	29.8	54.3	100.1	247.5
2000[4]	15.9	18.6	29.4	50.4	94.6	242.0
1999[4]	15.5	20.6	30.0	52.7	101.3	250.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Nontransport accidents (W00-X59,Y86)														
All origins[3]														
Male														
2001	23.8	22.9	8.3	3.3	3.3	9.9	16.8	16.6	17.7	24.8	28.7	30.0	24.3	20.5
2000[4]	22.9	20.9	9.0	3.4	3.5	10.2	15.5	16.2	18.4	24.2	28.3	28.0	23.7	20.7
1999[4]	23.3	20.2	9.3	3.5	3.9	10.0	15.3	16.2	19.2	23.4	29.1	28.6	22.0	20.6
Female														
2001	14.5	17.9	5.1	1.8	1.5	2.3	3.4	4.2	5.5	8.5	10.2	9.7	8.6	8.2
2000[4]	13.6	16.2	5.7	1.9	1.4	2.2	3.2	3.7	5.2	7.4	9.0	8.4	7.3	8.0
1999[4]	13.7	14.2	6.1	2.0	1.5	2.1	3.1	3.7	5.0	7.2	8.7	8.1	6.9	7.6
Hispanic														
Both sexes														
2001	10.1	13.6	5.0	1.8	1.4	5.1	8.0	8.6	10.3	14.4	15.8	16.3	14.0	12.6
2000[4]	10.2	9.9	5.2	1.7	1.4	6.0	8.5	8.4	11.6	14.7	16.2	14.7	13.5	12.4
1999[4]	11.1	12.4	6.4	1.8	1.7	6.0	9.4	9.1	12.1	16.6	18.0	17.8	14.1	11.2
Male														
2001	14.8	14.2	6.7	2.3	1.5	8.5	12.8	13.8	16.1	23.1	25.7	27.5	23.1	21.2
2000[4]	15.3	10.9	5.9	1.8	1.3	9.5	14.5	13.8	19.5	23.4	28.3	26.1	22.0	18.9
1999[4]	16.9	15.0	7.7	2.4	2.2	9.6	15.9	15.1	20.0	27.1	29.5	30.9	23.2	20.6
Female														
2001	5.2	13.0	3.3	1.3	1.3	*	2.1	2.3	3.6	4.7	5.3	5.1	5.3	4.9
2000[4]	4.9	8.8	4.4	1.5	1.4	1.9	*	2.2	2.6	5.1	3.6	3.4	5.5	6.5
1999[4]	5.0	9.8	5.2	1.2	*	2.0	1.4	2.1	3.1	5.1	6.0	4.9	5.5	*
Non-Hispanic[5]														
Both sexes														
2001	20.2	22.0	7.2	2.7	2.6	6.4	10.7	10.9	11.8	16.9	19.7	19.8	16.4	14.1
2000[4]	19.2	20.7	7.9	2.9	2.7	6.4	9.7	10.3	11.8	15.8	18.7	18.1	15.2	14.2
1999[4]	19.3	18.5	8.0	2.9	2.9	6.1	9.3	10.1	12.0	15.0	18.7	18.0	14.1	13.8
Male														
2001	25.0	24.9	8.7	3.5	3.6	10.1	17.7	17.2	17.8	24.8	28.8	29.9	24.2	20.1
2000[4]	23.9	23.2	9.7	3.7	3.8	10.4	15.7	16.7	18.0	24.1	28.0	27.8	23.5	20.6
1999[4]	24.0	21.5	9.6	3.7	4.2	10.0	15.1	16.4	18.8	22.7	28.8	28.0	21.6	20.2
Female														
2001	15.7	19.1	5.6	1.9	1.5	2.5	3.6	4.6	5.8	9.1	10.7	10.1	8.9	8.4
2000[4]	14.7	18.0	6.0	1.9	1.4	2.2	3.6	3.9	5.6	7.7	9.6	8.8	7.3	8.1
1999[4]	14.8	15.2	6.3	2.1	1.6	2.1	3.4	3.9	5.3	7.4	8.9	8.4	6.9	7.9

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)						
All origins[3]						
Male						
2001	22.5	27.8	38.8	72.1	127.8	305.8
2000[4]	22.7	25.5	38.4	65.3	120.8	299.8
1999[4]	22.7	28.3	40.7	66.6	133.5	311.9
Female						
2001	10.3	13.9	22.6	41.8	83.5	223.1
2000[4]	9.7	12.8	22.4	40.0	79.2	218.4
1999[4]	9.0	14.0	21.7	43.0	82.4	225.5
Hispanic						
Both sexes						
2001	12.6	14.4	26.5	34.8	59.1	142.6
2000[4]	14.0	14.8	22.8	35.8	70.7	159.2
1999[4]	15.3	19.5	21.6	34.5	62.8	157.2
Male						
2001	19.3	20.2	35.2	47.1	67.5	154.3
2000[4]	20.7	23.5	33.1	47.2	78.0	207.6
1999[4]	25.0	29.8	34.1	42.1	80.2	217.8
Female						
2001	6.8	9.8	19.9	26.1	53.9	136.8
2000[4]	8.2	7.9	15.1	27.7	66.3	135.5
1999[4]	6.9	11.3	12.3	29.2	52.5	127.9
Non-Hispanic[5]						
Both sexes						
2001	16.2	20.6	29.8	55.1	101.4	250.8
2000[4]	15.9	18.8	29.7	50.8	95.3	244.4
1999[4]	15.4	20.5	30.4	53.3	102.3	252.8
Male						
2001	22.6	28.1	38.8	73.1	129.8	311.6
2000[4]	22.7	25.4	38.5	65.9	122.2	302.7
1999[4]	22.3	28.0	40.9	67.4	134.9	314.3
Female						
2001	10.5	14.1	22.7	42.5	84.4	225.5
2000[4]	9.7	13.1	22.7	40.4	79.5	220.8
1999[4]	9.1	14.1	22.1	43.4	83.2	228.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Nontransport accidents (W00-X59,Y86)														
Non-Hispanic White														
Both sexes														
2001	21.5	17.8	7.1	2.3	2.2	6.5	11.9	11.6	12.5	17.4	19.5	19.0	15.3	13.3
2000[4]	20.1	16.6	7.3	2.3	2.3	6.4	10.5	10.9	12.1	16.0	18.4	17.2	13.9	13.2
1999[4]	20.0	14.7	7.4	2.2	2.5	6.1	9.6	10.6	11.9	15.2	18.3	16.7	12.9	12.8
Male														
2001	25.9	20.0	8.8	2.8	3.0	10.2	19.6	18.2	18.9	25.6	28.7	28.2	22.0	18.4
2000[4]	24.5	19.1	9.1	3.0	3.4	10.4	17.0	17.5	18.6	24.5	27.4	25.9	21.1	18.8
1999[4]	24.3	17.1	8.9	2.7	3.6	9.8	15.4	17.1	18.7	22.9	28.0	25.6	19.4	18.0
Female														
2001	17.2	15.5	5.3	1.8	1.2	2.5	4.1	4.9	6.0	9.2	10.3	9.8	8.7	8.4
2000[4]	16.0	14.0	5.4	1.5	1.2	2.3	3.8	4.2	5.6	7.5	9.3	8.5	6.9	7.9
1999[4]	16.0	12.3	5.9	1.7	1.4	2.2	3.6	4.0	5.1	7.4	8.6	8.0	6.5	7.8
Non-Hispanic Black														
Both sexes														
2001	18.1	41.6	8.7	4.3	4.3	6.1	7.5	9.7	11.3	18.0	25.5	30.1	28.5	23.1
2000[4]	18.0	39.5	10.9	5.4	4.1	6.5	7.8	9.8	12.8	18.3	24.9	28.1	28.0	24.3
1999[4]	19.4	36.4	11.1	5.8	4.8	6.5	9.2	10.5	14.8	16.9	25.9	31.3	26.2	24.9
Male														
2001	25.0	48.2	10.0	6.2	5.9	9.8	12.4	15.4	16.9	26.4	36.9	48.6	47.1	37.7
2000[4]	25.5	42.3	12.9	7.2	5.8	10.7	12.8	16.4	19.4	27.5	38.2	46.2	47.1	41.1
1999[4]	27.5	42.3	13.7	7.7	6.8	11.2	16.2	17.2	22.8	25.9	41.2	52.3	44.0	43.0
Female														
2001	11.8	34.6	7.4	2.3	2.6	2.2	2.8	4.5	6.3	10.5	15.4	14.2	12.7	11.2
2000[4]	11.3	36.6	8.8	3.6	2.4	2.2	3.1	3.9	6.9	10.2	13.0	12.5	11.8	10.5
1999[4]	12.0	30.3	8.3	3.9	2.6	1.7	2.6	4.5	7.6	9.0	12.5	13.2	11.1	10.1
Falls (W00-W19)														
All origins[3]														
Both sexes														
2001	5.3	0.6	0.2	0.2	0.2	0.4	0.9	0.8	0.9	1.2	1.6	2.5	2.7	3.3
2000[4]	4.7	*	0.2	*	0.1	0.5	0.7	0.7	0.8	1.0	1.6	2.0	2.6	3.3
1999[4]	4.7	*	0.4	0.1	0.1	0.6	0.7	0.8	0.9	1.2	1.7	2.3	2.2	3.0
Male														
2001	5.8	*	0.3	0.2	0.2	0.8	1.5	1.5	1.5	2.1	2.8	4.2	4.4	5.0
2000[4]	5.2	*	0.3	*	*	0.8	1.2	1.2	1.5	1.8	2.7	3.4	4.2	5.0
1999[4]	5.2	*	0.5	*	0.2	0.9	1.2	1.4	1.5	2.0	2.8	3.8	3.6	4.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)						
Non-Hispanic White						
Both sexes						
2001	15.5	20.3	29.4	55.4	103.6	259.4
2000[4]	15.3	18.0	29.1	50.9	97.3	253.2
1999[4]	13.9	19.5	29.0	52.5	103.7	260.8
Male						
2001	21.1	27.5	38.0	73.0	130.7	320.4
2000[4]	21.4	23.8	37.6	64.9	123.5	311.5
1999[4]	20.3	26.0	38.7	65.5	136.4	322.6
Female						
2001	10.3	14.0	22.3	43.0	87.2	234.0
2000[4]	9.6	12.9	22.2	41.0	81.7	229.7
1999[4]	8.0	13.9	21.3	43.4	84.3	236.2
Non-Hispanic Black						
Both sexes						
2001	24.9	27.0	39.2	56.9	85.8	166.1
2000[4]	23.2	28.3	38.9	56.5	79.6	161.2
1999[4]	28.6	33.1	46.4	66.0	96.9	181.1
Male						
2001	38.5	40.9	56.1	81.5	135.7	220.6
2000[4]	38.1	43.4	53.3	85.7	116.7	231.3
1999[4]	41.8	51.5	71.3	95.7	129.3	249.6
Female						
2001	14.3	16.7	28.0	42.2	60.4	145.9
2000[4]	11.4	17.1	29.3	38.9	60.9	135.2
1999[4]	18.3	19.5	29.9	48.1	80.8	155.5
Falls (W00-W19)						
All origins[3]						
Both sexes						
2001	4.9	7.8	12.4	25.2	49.9	121.5
2000[4]	4.6	6.2	12.0	22.1	44.5	112.6
1999[4]	4.6	6.9	11.3	22.3	45.5	110.2
Male						
2001	7.1	10.6	16.0	34.0	64.1	155.7
2000[4]	6.9	8.6	15.7	29.4	57.3	145.4
1999[4]	7.0	9.8	15.3	27.9	61.6	142.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Falls (W00-W19)														
All origins[3]														
Female														
2001	4.8	*	*	*	*	*	*	*	0.3	0.4	0.6	0.9	1.2	1.6
2000[4]	4.3	*	*	*	*	*	0.2	*	*	0.3	0.6	0.8	1.2	1.8
1999[4]	4.3	*	*	*	*	*	*	*	0.3	0.4	0.6	0.8	0.9	1.5
Hispanic														
Both sexes														
2001	2.0	*	*	*	*	*	0.8	1.0	1.1	1.5	2.3	2.7	2.4	3.8
2000[4]	2.0	*	*	*	*	*	0.8	1.1	1.2	1.1	1.5	2.2	2.8	3.6
1999[4]	1.9	*	*	*	*	*	1.0	1.1	1.1	1.2	1.7	2.0	2.4	3.3
Male														
2001	2.8	*	*	*	*	*	1.5	1.8	2.1	2.6	4.2	5.1	3.6	6.4
2000[4]	2.7	*	*	*	*	*	1.4	1.9	2.2	1.8	3.0	3.9	5.1	5.9
1999[4]	2.7	*	*	*	*	*	1.7	1.9	1.9	2.1	2.8	3.3	4.5	6.5
Female														
2001	1.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	5.7	*	0.2	0.1	0.2	0.4	0.9	0.8	0.8	1.2	1.6	2.4	2.8	3.2
2000[4]	5.1	*	0.2	*	*	0.5	0.7	0.6	0.7	1.0	1.6	2.0	2.6	3.3
1999[4]	5.1	*	0.4	0.1	0.2	0.6	0.6	0.7	0.9	1.2	1.7	2.3	2.2	3.0
Male														
2001	6.2	*	0.3	*	0.2	0.7	1.5	1.4	1.3	2.0	2.6	4.0	4.5	4.9
2000[4]	5.5	*	*	*	*	0.8	1.1	1.1	1.3	1.7	2.6	3.3	4.0	4.9
1999[4]	5.5	*	0.5	*	*	1.0	1.1	1.3	1.4	1.9	2.7	3.8	3.5	4.5
Female														
2001	5.3	*	*	*	*	*	*	*	0.3	0.4	0.6	0.9	1.1	1.6
2000[4]	4.7	*	*	*	*	*	0.3	*	*	0.3	0.7	0.8	1.2	1.8
1999[4]	4.6	*	*	*	*	*	*	*	0.3	0.5	0.6	0.8	1.0	1.6
Non-Hispanic White														
Both sexes														
2001	6.6	*	0.2	*	0.2	0.5	1.0	0.9	0.9	1.2	1.5	2.4	2.8	3.1
2000[4]	5.8	*	*	*	*	0.6	0.8	0.7	0.8	1.0	1.7	2.1	2.5	3.2
1999[4]	5.7	*	0.4	*	0.2	0.7	0.7	0.8	0.8	1.3	1.7	2.2	2.1	2.8

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Falls (W00-W19)						
All origins[3]						
Female						
2001	2.9	5.4	9.5	19.0	41.3	107.2
2000[4]	2.6	4.2	9.1	17.1	36.9	99.2
1999[4]	2.4	4.5	8.1	18.4	36.0	97.2
Hispanic						
Both sexes						
2001	4.8	5.5	13.5	19.0	31.8	70.0
2000[4]	5.2	6.7	9.6	18.4	41.2	81.0
1999[4]	4.5	8.1	8.1	17.8	35.8	86.3
Male						
2001	8.1	7.4	16.5	22.5	37.7	72.4
2000[4]	7.5	10.4	15.6	25.1	41.2	104.8
1999[4]	8.0	10.8	12.2	22.6	48.7	120.8
Female						
2001	*	*	11.2	16.5	28.1	68.9
2000[4]	*	*	*	13.6	41.2	69.2
1999[4]	*	*	*	14.3	28.1	69.7
Non-Hispanic[5]						
Both sexes						
2001	4.8	7.9	12.3	25.5	50.4	123.1
2000[4]	4.6	6.2	12.1	22.2	44.5	113.5
1999[4]	4.5	6.8	11.4	22.3	45.6	110.7
Male						
2001	6.9	10.7	15.8	34.6	64.9	159.0
2000[4]	6.8	8.4	15.7	29.5	57.9	146.5
1999[4]	6.8	9.7	15.4	27.9	61.7	142.9
Female						
2001	2.9	5.5	9.4	19.2	41.7	108.1
2000[4]	2.5	4.3	9.3	17.2	36.7	100.1
1999[4]	2.4	4.4	8.3	18.5	36.2	97.8
Non-Hispanic White						
Both sexes						
2001	4.9	8.1	12.7	26.6	53.1	129.9
2000[4]	4.5	6.1	12.5	23.2	46.7	120.2
1999[4]	4.3	6.8	11.5	23.1	47.9	116.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Falls (W00-W19)														
Non-Hispanic White														
Male														
2001	7.0	*	*	*	*	0.8	1.9	1.6	1.4	2.0	2.5	3.8	4.4	4.7
2000[4]	6.2	*	*	*	*	1.0	1.3	1.3	1.3	1.7	2.7	3.4	3.7	4.6
1999[4]	6.1	*	0.5	*	*	1.1	1.2	1.4	1.4	2.0	2.7	3.6	3.2	4.1
Female														
2001	6.2	*	*	*	*	*	*	*	0.3	0.4	0.6	1.0	1.2	1.6
2000[4]	5.5	*	*	*	*	*	*	*	*	0.3	0.7	0.8	1.2	1.8
1999[4]	5.4	*	*	*	*	*	*	*	0.3	0.5	0.7	0.8	0.9	1.5
Non-Hispanic Black														
Both sexes														
2001	2.3	*	*	*	*	*	*	*	*	1.2	2.1	2.6	3.4	4.0
2000[4]	2.1	*	*	*	*	*	*	*	*	1.0	1.6	2.0	3.6	4.6
1999[4]	2.5	*	*	*	*	*	*	*	1.1	0.9	1.8	3.3	3.7	4.5
Male														
2001	3.2	*	*	*	*	*	*	*	*	2.0	3.5	4.9	6.1	6.4
2000[4]	2.8	*	*	*	*	*	*	*	*	1.7	2.5	3.3	6.3	8.0
1999[4]	3.4	*	*	*	*	*	*	*	1.8	1.7	3.4	5.7	6.1	7.5
Female														
2001	1.6	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.5	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.7	*	*	*	*	*	*	*	*	*	*	*	*	*
Accidental discharge of firearms (W32-W34)														
All origins[3]														
Both sexes														
2001	0.3	*	*	*	0.2	0.5	0.5	0.4	0.2	0.3	0.3	0.3	0.2	0.3
2000[4]	0.3	*	*	*	0.2	0.5	0.5	0.4	0.3	0.3	0.3	0.2	0.2	0.3
1999[4]	0.3	*	*	*	0.3	0.6	0.7	0.4	0.3	0.3	0.3	0.3	0.2	0.2
Male														
2001	0.5	*	*	*	0.3	1.0	0.9	0.7	0.4	0.5	0.6	0.5	0.4	0.5
2000[4]	0.5	*	*	*	0.4	0.9	0.9	0.7	0.4	0.5	0.6	0.3	0.4	0.5
1999[4]	0.5	*	*	*	0.5	1.1	1.2	0.6	0.6	0.5	0.5	0.5	0.4	0.3
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Falls (W00-W19)						
Non-Hispanic White						
Male						
2001	6.9	10.9	16.3	35.7	67.5	166.0
2000[4]	6.8	8.2	16.2	30.4	60.3	154.1
1999[4]	6.4	9.4	15.4	28.4	64.4	149.7
Female						
2001	3.0	5.6	9.7	20.1	44.4	114.8
2000[4]	2.5	4.3	9.5	18.2	38.6	106.5
1999[4]	2.3	4.5	8.4	19.4	38.1	103.4
Non-Hispanic Black						
Both sexes						
2001	4.9	8.0	9.8	16.7	21.8	44.0
2000[4]	5.1	7.1	9.7	11.8	20.7	39.1
1999[4]	6.7	9.0	9.7	13.3	23.5	45.3
Male						
2001	7.5	12.0	13.4	24.6	36.8	68.1
2000[4]	8.5	10.7	12.3	18.8	26.6	61.1
1999[4]	10.2	15.1	14.9	19.8	31.9	58.2
Female						
2001	*	5.0	7.3	11.9	14.2	35.1
2000[4]	*	4.5	7.9	7.6	17.7	31.0
1999[4]	3.9	4.5	6.2	9.3	19.3	40.4
Accidental discharge of firearms (W32-W34)						
All origins[3]						
Both sexes						
2001	0.2	*	*	0.3	*	*
2000[4]	*	0.3	*	*	*	*
1999[4]	0.2	*	*	*	*	*
Male						
2001	0.4	*	*	0.7	*	*
2000[4]	*	0.5	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Accidental discharge of firearms (W32-W34)														
Hispanic														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.3	*	*	*	0.2	0.5	0.5	0.4	0.2	0.3	0.3	0.3	0.2	0.3
2000[4]	0.3	*	*	*	0.3	0.6	0.5	0.4	0.3	0.3	0.4	0.2	0.2	0.3
1999[4]	0.3	*	*	*	0.3	0.7	0.7	0.4	0.3	0.3	0.3	0.3	0.3	0.2
Male														
2001	0.5	*	*	*	0.4	1.0	0.9	0.7	0.4	0.5	0.6	0.5	0.4	0.5
2000[4]	0.5	*	*	*	0.5	1.0	0.9	0.7	0.5	0.6	0.6	0.3	0.5	0.5
1999[4]	0.5	*	*	*	0.5	1.2	1.3	0.7	0.6	0.5	0.5	0.5	0.4	0.4
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	0.3	*	*	*	0.2	0.5	0.5	0.4	0.3	0.4	0.4	0.3	0.3	0.3
2000[4]	0.3	*	*	*	0.3	0.5	0.4	0.4	0.3	0.3	0.4	0.2	0.3	0.3
1999[4]	0.3	*	*	*	0.3	0.6	0.6	0.4	0.3	0.3	0.3	0.3	0.3	0.2
Male														
2001	0.5	*	*	*	0.4	0.8	0.9	0.6	0.5	0.6	0.6	0.6	0.5	0.5
2000[4]	0.5	*	*	*	0.5	0.9	0.8	0.6	0.4	0.6	0.7	0.3	0.5	0.5
1999[4]	0.5	*	*	*	0.6	1.0	1.1	0.7	0.5	0.5	0.5	0.5	0.4	*
Female														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental discharge of firearms (W32-W34)						
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	0.2	*	*	0.3	*	*
2000[4]	*	0.3	*	*	*	*
1999[4]	0.2	*	*	*	*	*
Male						
2001	0.4	*	*	0.7	*	*
2000[4]	*	0.5	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic White						
Both sexes						
2001	0.3	*	*	0.3	*	*
2000[4]	*	0.3	*	*	*	*
1999[4]	0.2	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Accidental discharge of firearms (W32-W34)														
Non-Hispanic White														
Female														
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	0.3	*	*	*	*	0.8	0.8	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	0.9	0.9	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	1.2	1.2	*	*	*	*	*	*	*
Male														
2001	0.6	*	*	*	*	1.6	1.6	*	*	*	*	*	*	*
2000[4]	0.7	*	*	*	*	1.7	1.6	*	*	*	*	*	*	*
1999[4]	0.8	*	*	*	*	2.2	2.2	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Accidental drowning and submersion (W65-W74)														
All origins[3]														
Both sexes														
2001	1.2	1.7	3.0	0.8	0.8	1.6	1.4	1.1	0.8	1.0	1.1	1.0	0.8	0.8
2000[4]	1.2	2.0	3.2	1.0	0.8	1.8	1.5	1.1	1.0	1.1	1.0	0.9	1.0	0.9
1999[4]	1.3	1.8	3.2	0.9	0.9	1.8	1.5	1.1	1.1	1.1	1.0	1.0	1.0	0.9
Male														
2001	1.8	1.5	3.7	1.2	1.1	2.8	2.5	1.9	1.4	1.6	1.7	1.8	1.3	1.2
2000[4]	2.0	2.2	4.0	1.3	1.2	3.3	2.5	1.9	1.7	1.9	1.7	1.6	1.7	1.6
1999[4]	2.0	2.0	4.0	1.3	1.3	3.2	2.8	2.0	2.0	1.9	1.7	1.7	1.8	1.5
Female														
2001	0.5	1.9	2.2	0.5	0.5	0.3	0.3	0.3	0.2	0.4	0.4	0.3	0.3	0.4
2000[4]	0.5	1.7	2.4	0.6	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	*
1999[4]	0.5	1.6	2.4	0.5	0.4	0.3	0.3	*	0.3	0.3	0.3	0.3	0.3	0.4
Hispanic														
Both sexes														
2001	1.1	*	2.6	0.7	*	1.3	1.7	1.1	1.1	1.2	0.8	1.1	*	*
2000[4]	1.2	*	2.5	0.8	*	2.3	2.2	1.0	0.9	0.9	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental discharge of firearms (W32-W34)						
Non-Hispanic White						
Female						
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Accidental drowning and submersion (W65-W74)						
All origins[3]						
Both sexes						
2001	0.9	0.8	0.9	1.1	1.4	1.7
2000[4]	0.9	0.8	1.1	1.0	1.7	1.5
1999[4]	0.9	0.9	1.1	1.2	1.4	1.6
Male						
2001	1.4	1.4	1.6	1.8	2.4	3.5
2000[4]	1.5	1.4	1.8	1.2	2.7	2.8
1999[4]	1.5	1.5	1.7	2.1	2.7	2.3
Female						
2001	0.4	*	*	0.7	0.8	1.0
2000[4]	0.4	*	0.6	0.8	1.1	1.0
1999[4]	0.4	0.4	0.7	0.6	0.7	1.4
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Accidental drowning and submersion (W65-W74)														
Hispanic														
Both sexes														
1999[4]	1.4	*	2.7	0.7	0.7	2.2	2.3	1.2	1.3	1.0	*	*	*	*
Male														
2001	1.8	*	3.5	1.1	*	2.5	2.9	1.7	1.8	1.9	*	*	*	*
2000[4]	1.9	*	2.9	*	*	3.8	3.7	1.8	1.7	1.4	*	*	*	*
1999[4]	2.3	*	2.9	*	*	4.1	4.1	2.1	2.3	1.7	*	*	*	*
Female														
2001	0.4	*	1.7	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.5	*	2.2	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.5	*	2.5	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	1.1	1.6	3.1	0.8	0.9	1.6	1.3	1.1	0.7	1.0	1.1	1.0	0.8	0.8
2000[4]	1.2	2.1	3.4	1.0	0.9	1.7	1.3	1.1	1.0	1.1	1.1	0.9	1.0	0.9
1999[4]	1.2	1.7	3.3	1.0	0.9	1.7	1.4	1.1	1.1	1.1	1.0	1.0	1.0	1.0
Male														
2001	1.8	1.3	3.8	1.1	1.2	2.9	2.4	2.0	1.3	1.6	1.8	1.8	1.4	1.3
2000[4]	2.0	2.4	4.3	1.4	1.4	3.1	2.2	1.9	1.8	1.9	1.8	1.6	1.7	1.6
1999[4]	2.0	1.7	4.2	1.4	1.3	3.0	2.4	1.9	1.9	1.9	1.8	1.7	1.7	1.6
Female														
2001	0.5	1.9	2.4	0.5	0.5	0.3	0.3	0.3	*	0.3	0.4	0.3	0.3	0.4
2000[4]	0.5	1.9	2.4	0.6	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	*
1999[4]	0.5	1.7	2.4	0.6	0.4	0.3	0.3	*	0.3	0.3	0.3	0.3	0.3	0.4
Non-Hispanic White														
Both sexes														
2001	1.1	1.5	3.5	0.6	0.5	1.3	1.3	1.0	0.7	0.9	1.0	0.9	0.8	0.7
2000[4]	1.1	1.7	3.7	0.8	0.6	1.4	1.1	1.0	1.0	1.1	1.0	0.9	0.9	0.9
1999[4]	1.1	1.3	3.6	0.6	0.5	1.4	1.2	0.9	1.0	1.1	1.1	1.1	1.0	0.9
Male														
2001	1.6	*	4.3	0.6	0.7	2.2	2.2	1.7	1.1	1.4	1.7	1.6	1.3	1.1
2000[4]	1.8	1.9	4.7	1.0	0.9	2.5	1.9	1.7	1.7	1.9	1.7	1.5	1.5	1.6
1999[4]	1.8	*	4.5	0.8	0.7	2.4	2.0	1.6	1.7	1.8	1.7	1.8	1.7	1.5
Female														
2001	0.5	1.7	2.6	0.5	*	*	0.4	*	*	0.4	0.4	0.3	0.4	0.4
2000[4]	0.5	*	2.6	0.5	*	*	0.4	*	*	0.4	0.3	0.3	0.3	*
1999[4]	0.5	*	2.6	0.5	*	*	*	*	0.3	0.4	0.4	0.4	0.3	0.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental drowning and submersion (W65-W74)						
Hispanic						
Both sexes						
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	0.9	0.8	1.0	1.2	1.4	1.7
2000[4]	0.9	0.9	1.1	1.0	1.7	1.5
1999[4]	0.9	0.9	1.2	1.2	1.5	1.7
Male						
2001	1.4	1.4	1.7	1.8	2.5	3.4
2000[4]	1.6	1.5	1.7	1.3	2.7	2.8
1999[4]	1.5	1.5	1.6	2.0	2.8	2.4
Female						
2001	0.5	*	*	0.7	0.8	1.0
2000[4]	*	*	0.6	0.8	1.1	1.0
1999[4]	0.4	0.4	0.8	0.6	0.7	1.4
Non-Hispanic White						
Both sexes						
2001	0.9	0.8	1.0	1.2	1.4	1.7
2000[4]	0.9	0.9	1.0	0.9	1.8	1.4
1999[4]	0.9	0.9	1.0	1.1	1.5	1.6
Male						
2001	1.3	1.4	1.7	1.8	2.4	3.5
2000[4]	1.7	1.4	1.6	1.1	3.0	2.7
1999[4]	1.4	1.4	1.5	1.9	2.8	2.4
Female						
2001	0.5	*	*	0.7	0.7	1.0
2000[4]	*	*	0.6	0.7	1.1	0.9
1999[4]	*	*	0.7	0.5	*	1.3

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Accidental drowning and submersion (W65-W74)														
Non-Hispanic Black														
Both sexes														
2001	1.6	*	2.0	1.7	2.4	3.0	1.6	1.3	1.1	1.2	1.3	1.4	*	1.5
2000[4]	1.7	4.3	2.3	2.0	2.1	2.8	1.7	1.3	0.9	1.2	1.5	1.0	1.6	*
1999[4]	1.8	3.7	2.5	2.3	2.6	2.8	2.0	1.8	1.5	1.1	1.0	*	*	*
Male														
2001	2.8	*	2.4	2.8	3.4	5.4	3.2	2.8	2.2	2.2	2.2	2.4	*	*
2000[4]	2.8	*	2.8	2.8	3.3	5.4	3.3	2.4	1.8	2.3	2.8	2.0	2.9	*
1999[4]	3.2	*	3.4	3.6	4.1	5.1	4.0	3.6	2.7	2.2	1.9	*	*	*
Female														
2001	0.5	*	*	*	1.4	*	*	*	*	*	*	*	*	*
2000[4]	0.6	*	1.8	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
Accidental exposure to smoke, fire and flames (X00-X09)														
All origins[3]														
Both sexes														
2001	1.2	1.2	1.5	0.8	0.4	0.4	0.7	0.6	0.6	0.9	1.1	1.0	1.1	1.5
2000[4]	1.2	1.0	1.9	0.9	0.4	0.4	0.6	0.6	0.6	0.8	1.0	1.1	1.3	1.4
1999[4]	1.2	1.1	2.0	0.8	0.5	0.4	0.6	0.7	0.7	0.9	1.0	1.0	1.0	1.2
Male														
2001	1.4	1.5	1.8	1.0	0.4	0.4	0.9	0.8	0.8	1.2	1.6	1.5	1.5	2.1
2000[4]	1.5	1.0	2.2	1.0	0.3	0.4	0.8	0.8	0.8	1.2	1.2	1.5	1.8	1.9
1999[4]	1.5	1.1	2.2	0.9	0.5	0.6	0.8	0.9	1.0	1.2	1.5	1.5	1.3	1.6
Female														
2001	0.9	1.0	1.1	0.6	0.4	0.3	0.4	0.4	0.4	0.5	0.7	0.6	0.7	1.0
2000[4]	0.9	*	1.5	0.7	0.5	0.4	0.4	0.4	0.4	0.5	0.7	0.7	0.7	0.9
1999[4]	0.9	1.1	1.7	0.7	0.4	0.3	0.4	0.5	0.4	0.5	0.5	0.5	0.7	0.9
Hispanic														
Both sexes														
2001	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.5	*	0.7	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.6	*	1.4	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.6	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.7	*	1.6	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental drowning and submersion (W65-W74)						
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Accidental exposure to smoke, fire and flames (X00-X09)						
All origins[3]						
Both sexes						
2001	1.5	2.0	2.5	3.0	3.8	5.7
2000[4]	1.7	2.0	2.4	3.5	4.3	6.2
1999[4]	1.5	2.0	2.5	3.5	4.7	5.9
Male						
2001	1.8	2.5	3.0	3.5	5.1	8.3
2000[4]	2.2	2.4	3.2	4.5	5.9	9.4
1999[4]	1.9	2.5	3.4	4.3	6.6	9.3
Female						
2001	1.2	1.5	2.1	2.7	3.1	4.6
2000[4]	1.2	1.6	1.8	2.8	3.4	4.9
1999[4]	1.2	1.6	1.8	3.0	3.6	4.5
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Accidental exposure to smoke, fire and flames (X00-X09)														
Hispanic														
Female														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	1.3	1.3	1.7	0.9	0.5	0.4	0.7	0.7	0.7	0.9	1.1	1.1	1.2	1.6
2000[4]	1.3	1.0	2.2	1.0	0.5	0.4	0.7	0.7	0.7	0.9	1.0	1.2	1.3	1.4
1999[4]	1.3	1.2	2.1	0.9	0.5	0.4	0.7	0.8	0.7	0.9	1.0	1.0	1.0	1.2
Male														
2001	1.6	1.5	2.0	1.1	0.5	0.5	1.0	0.8	0.9	1.3	1.6	1.5	1.6	2.1
2000[4]	1.6	*	2.6	1.2	0.4	0.4	0.9	0.9	0.9	1.3	1.2	1.6	1.9	2.0
1999[4]	1.6	*	2.3	1.0	0.6	0.6	0.8	1.1	1.1	1.3	1.5	1.5	1.4	1.6
Female														
2001	1.0	*	1.3	0.6	0.4	0.4	0.4	0.5	0.5	0.6	0.7	0.6	0.7	1.1
2000[4]	1.0	*	1.7	0.8	0.5	0.3	0.5	0.5	0.5	0.5	0.7	0.7	0.8	0.9
1999[4]	1.0	*	1.8	0.8	0.4	0.3	0.5	0.6	0.4	0.6	0.6	0.6	0.7	0.9
Non-Hispanic White														
Both sexes														
2001	1.1	1.1	1.3	0.7	0.4	0.4	0.7	0.5	0.7	0.8	1.0	0.9	1.0	1.3
2000[4]	1.1	*	1.5	0.7	0.4	0.4	0.6	0.6	0.6	0.8	0.9	1.0	1.2	1.1
1999[4]	1.1	0.9	1.5	0.6	0.4	0.4	0.6	0.8	0.7	0.8	1.0	0.9	0.9	1.1
Male														
2001	1.4	*	1.7	0.9	0.4	0.5	1.0	0.8	0.9	1.1	1.5	1.3	1.3	1.8
2000[4]	1.4	*	1.8	0.8	0.3	0.4	0.9	0.7	0.8	1.1	1.1	1.4	1.7	1.6
1999[4]	1.4	*	1.7	0.6	0.5	0.6	0.8	1.1	1.1	1.1	1.4	1.3	1.2	1.4
Female														
2001	0.9	*	0.9	0.5	0.4	0.3	0.3	*	0.5	0.5	0.6	0.6	0.7	0.9
2000[4]	0.9	*	1.1	0.6	0.4	*	0.4	0.4	0.4	0.4	0.6	0.6	0.7	0.7
1999[4]	0.8	*	1.3	0.6	0.3	*	0.4	0.5	0.3	0.5	0.5	0.5	0.6	0.9
Non-Hispanic Black														
Both sexes														
2001	2.3	*	3.4	1.8	0.8	*	1.0	1.4	1.0	1.6	2.1	2.3	2.6	3.8
2000[4]	2.6	*	5.1	2.2	0.8	*	1.0	1.2	1.3	1.6	2.0	2.2	2.5	3.9
1999[4]	2.6	*	4.7	2.2	0.9	*	0.9	1.1	0.9	1.9	1.7	2.1	2.3	2.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)						
Hispanic						
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	1.5	2.1	2.6	3.1	3.9	5.7
2000[4]	1.7	2.0	2.4	3.4	4.3	6.3
1999[4]	1.5	2.1	2.6	3.5	4.7	6.0
Male						
2001	1.8	2.6	3.1	3.5	5.2	8.3
2000[4]	2.3	2.5	3.2	4.4	5.9	9.3
1999[4]	1.8	2.5	3.5	4.2	6.6	9.2
Female						
2001	1.2	1.6	2.2	2.8	3.1	4.6
2000[4]	1.2	1.7	1.7	2.8	3.4	5.0
1999[4]	1.2	1.7	1.9	3.1	3.6	4.7
Non-Hispanic White						
Both sexes						
2001	1.3	1.7	2.2	2.6	3.3	4.7
2000[4]	1.4	1.7	2.0	2.8	3.5	5.1
1999[4]	1.2	1.7	2.1	2.8	3.9	4.7
Male						
2001	1.5	2.2	2.5	2.8	4.5	6.9
2000[4]	1.8	2.0	2.6	3.6	4.9	8.1
1999[4]	1.6	2.0	2.7	3.2	5.7	7.8
Female						
2001	1.1	1.4	1.9	2.5	2.6	3.8
2000[4]	1.0	1.5	1.4	2.3	2.6	3.9
1999[4]	0.8	1.4	1.5	2.6	2.8	3.5
Non-Hispanic Black						
Both sexes						
2001	3.3	5.1	7.3	8.7	11.9	17.8
2000[4]	4.6	5.1	7.2	11.1	16.1	20.7
1999[4]	4.4	5.9	8.4	12.2	15.6	22.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Accidental exposure to smoke, fire and flames (X00-X09)														
Non-Hispanic Black														
Male														
2001	3.0	*	3.7	2.3	*	*	*	*	*	2.6	3.0	3.6	4.4	5.2
2000[4]	3.2	*	5.8	2.7	*	*	*	*	1.6	2.3	2.6	3.2	4.0	5.6
1999[4]	3.1	*	5.3	2.5	*	*	*	*	*	2.8	2.5	3.1	3.4	4.2
Female														
2001	1.8	*	3.1	1.3	*	*	*	1.5	*	*	*	*	*	2.8
2000[4]	2.0	*	4.3	1.7	*	*	*	*	*	*	1.4	*	*	*
1999[4]	2.1	*	4.1	1.9	*	*	*	*	*	*	*	*	*	*
Accidental poisoning and exposure to noxious substances (X40-X49)														
All origins[3]														
Both sexes														
2001	4.9	*	0.2	*	0.2	2.0	4.9	5.7	6.9	10.4	12.0	10.9	7.0	3.7
2000[4]	4.5	*	0.2	*	0.1	1.7	4.3	5.1	6.8	9.5	11.2	9.9	6.1	3.3
1999[4]	4.4	*	0.2	*	0.1	1.3	3.8	4.9	6.8	9.1	11.1	9.8	5.4	3.4
Male														
2001	7.1	*	0.3	*	*	3.0	7.6	8.7	9.9	14.6	16.8	15.7	9.9	5.0
2000[4]	6.6	*	0.3	*	*	2.6	6.7	7.9	10.1	13.7	16.4	14.7	8.9	4.0
1999[4]	6.5	*	*	*	*	1.9	5.9	7.7	10.3	13.3	16.6	15.1	7.9	4.7
Female														
2001	2.9	*	*	*	*	1.0	1.9	2.6	3.9	6.1	7.3	6.2	4.1	2.6
2000[4]	2.5	*	*	*	*	0.8	1.7	2.1	3.5	5.3	6.0	5.3	3.4	2.7
1999[4]	2.3	*	*	*	*	0.7	1.6	2.0	3.2	4.9	5.7	4.7	3.1	2.2
Hispanic														
Both sexes														
2001	3.6	*	*	*	*	1.5	2.7	4.3	5.5	8.4	9.1	8.6	6.6	4.7
2000[4]	3.7	*	*	*	*	1.2	2.8	3.9	6.8	9.5	9.8	8.9	6.8	4.0
1999[4]	4.4	*	*	*	*	1.3	3.2	4.8	7.5	11.4	12.4	11.4	7.6	4.1
Male														
2001	5.6	*	*	*	*	2.1	3.9	6.8	8.1	13.1	14.2	14.0	11.5	8.0
2000[4]	6.0	*	*	*	*	1.8	4.4	6.1	11.0	14.5	17.1	15.7	10.8	5.3
1999[4]	7.3	*	*	*	*	1.7	5.3	7.6	12.2	18.4	20.5	19.7	12.0	7.6
Female														
2001	1.5	*	*	*	*	*	1.3	1.4	2.5	3.2	3.7	3.1	*	*
2000[4]	1.3	*	*	*	*	*	*	1.3	1.9	3.9	2.2	*	3.0	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)						
Non-Hispanic Black						
Male						
2001	5.0	7.3	11.4	12.3	*	26.6
2000[4]	7.2	7.7	10.6	15.9	21.5	28.2
1999[4]	4.4	8.0	13.5	17.4	20.4	29.7
Female						
2001	*	*	4.6	6.5	10.4	14.5
2000[4]	*	*	5.0	8.1	13.4	17.9
1999[4]	4.4	4.3	5.1	9.0	13.2	19.1
Accidental poisoning and exposure to noxious substances (X40-X49)						
All origins[3]						
Both sexes						
2001	2.4	1.7	1.5	2.1	2.4	3.5
2000[4]	2.2	1.5	1.5	1.6	2.6	4.3
1999[4]	2.0	1.7	1.6	1.9	2.6	3.4
Male						
2001	3.0	2.0	2.1	2.7	2.8	3.8
2000[4]	2.7	1.8	1.7	1.6	2.9	6.2
1999[4]	3.0	2.2	1.9	2.1	2.9	4.4
Female						
2001	1.9	1.4	1.0	1.6	2.2	3.3
2000[4]	1.8	1.3	1.3	1.6	2.4	3.6
1999[4]	1.2	1.2	1.4	1.7	2.4	3.0
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	2.9	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Accidental poisoning and exposure to noxious substances (X40-X49)														
Hispanic														
Female														
1999[4]	1.4	*	*	*	*	*	*	1.5	2.0	3.8	4.0	3.1	3.5	*
Non-Hispanic[5]														
Both sexes														
2001	5.1	*	0.2	*	0.1	2.1	5.3	6.0	7.1	10.6	12.2	11.0	6.9	3.6
2000[4]	4.6	*	0.2	*	0.1	1.8	4.6	5.3	6.8	9.4	11.2	9.9	5.9	3.2
1999[4]	4.3	*	0.2	*	0.1	1.3	3.9	4.8	6.6	8.7	10.8	9.5	5.1	3.3
Male														
2001	7.2	*	*	*	*	3.1	8.5	9.1	10.1	14.7	17.0	15.7	9.6	4.6
2000[4]	6.6	*	*	*	*	2.8	7.2	8.3	9.8	13.5	16.2	14.4	8.5	3.8
1999[4]	6.3	*	*	*	*	1.9	6.0	7.6	9.9	12.5	15.9	14.4	7.3	4.4
Female														
2001	3.1	*	*	*	*	1.0	2.1	2.9	4.1	6.5	7.6	6.5	4.3	2.6
2000[4]	2.7	*	*	*	*	0.8	1.9	2.2	3.8	5.5	6.3	5.5	3.3	2.6
1999[4]	2.4	*	*	*	*	0.6	1.8	2.1	3.4	5.0	5.8	4.8	3.0	2.3
Non-Hispanic White														
Both sexes														
2001	5.2	*	*	*	*	2.4	6.2	6.9	7.7	11.0	12.3	10.5	6.0	3.2
2000[4]	4.6	*	*	*	*	2.2	5.4	5.9	7.2	9.7	11.0	9.2	5.0	2.9
1999[4]	4.2	*	*	*	*	1.5	4.3	5.3	6.7	8.8	10.3	8.7	4.4	3.0
Male														
2001	7.3	*	*	*	*	3.8	9.9	10.3	11.2	15.3	17.2	14.7	8.0	3.8
2000[4]	6.6	*	*	*	*	3.3	8.6	9.2	10.4	14.0	15.8	13.2	7.0	3.0
1999[4]	6.1	*	*	*	*	2.2	6.6	8.3	10.0	12.6	15.3	12.9	5.9	3.6
Female														
2001	3.1	*	*	*	*	1.0	2.5	3.4	4.2	6.7	7.4	6.3	4.0	2.7
2000[4]	2.7	*	*	*	*	1.0	2.0	2.5	3.9	5.4	6.2	5.3	3.0	2.7
1999[4]	2.4	*	*	*	*	0.7	2.0	2.3	3.3	4.9	5.4	4.5	2.9	2.4
Non-Hispanic Black														
Both sexes														
2001	6.0	*	*	*	*	0.9	2.4	4.1	6.4	11.2	15.6	18.1	15.6	7.3
2000[4]	5.7	*	*	*	*	*	2.2	4.3	7.0	10.9	15.6	16.8	14.3	6.9
1999[4]	5.8	*	*	*	*	*	2.9	4.3	8.2	10.4	16.3	18.2	12.2	6.5
Male														
2001	8.6	*	*	*	*	*	3.9	6.8	8.4	15.5	20.9	28.4	25.2	12.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)						
Hispanic						
Female						
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	2.4	1.7	1.5	2.1	2.4	3.6
2000[4]	2.2	1.5	1.5	1.6	2.6	4.4
1999[4]	1.9	1.6	1.6	1.9	2.6	3.5
Male						
2001	2.9	1.9	2.0	2.7	2.9	3.9
2000[4]	2.7	1.7	1.7	1.7	2.9	6.2
1999[4]	2.7	2.1	1.9	2.1	3.0	4.4
Female						
2001	1.9	1.5	1.0	1.7	2.2	3.4
2000[4]	1.8	1.4	1.3	1.6	2.4	3.6
1999[4]	1.3	1.2	1.4	1.8	2.4	3.1
Non-Hispanic White						
Both sexes						
2001	2.1	1.6	1.4	2.0	2.3	3.4
2000[4]	2.1	1.5	1.4	1.7	2.7	4.1
1999[4]	1.6	1.5	1.5	2.0	2.7	3.3
Male						
2001	2.3	1.8	1.9	2.4	2.7	3.8
2000[4]	2.3	1.4	1.6	1.7	3.0	5.8
1999[4]	2.0	1.9	1.7	2.2	3.1	4.6
Female						
2001	1.9	1.5	0.9	1.8	2.0	3.2
2000[4]	1.9	1.5	1.3	1.7	2.5	3.4
1999[4]	1.2	1.2	1.2	1.8	2.4	2.8
Non-Hispanic Black						
Both sexes						
2001	5.2	2.5	*	*	*	6.6
2000[4]	3.8	2.6	*	*	*	8.3
1999[4]	5.4	2.8	3.3	*	*	*
Male						
2001	9.0	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Accidental poisoning and exposure to noxious substances (X40-X49)														
Non-Hispanic Black														
Male														
2000[4]	8.4	*	*	*	*	*	2.8	7.1	9.8	15.0	23.2	27.0	23.2	11.7
1999[4]	8.8	*	*	*	*	*	4.8	6.9	11.7	14.9	24.4	30.0	20.7	12.2
Female														
2001	3.6	*	*	*	*	*	*	1.7	4.7	7.4	10.9	9.2	7.5	3.0
2000[4]	3.2	*	*	*	*	*	1.5	1.8	4.4	7.2	8.8	8.1	6.7	3.0
1999[4]	3.1	*	*	*	*	*	*	2.0	5.0	6.4	9.2	8.0	4.9	*
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)														
All origins[3]														
Both sexes														
2001	6.2	16.6	1.8	0.6	0.7	1.2	2.0	1.8	2.2	2.9	3.2	3.9	4.5	4.6
2000[4]	6.2	15.0	1.7	0.6	0.7	1.4	2.0	2.1	2.3	3.0	3.4	3.9	4.1	5.0
1999[4]	6.6	13.8	1.9	0.7	0.9	1.4	2.0	2.2	2.3	2.7	3.7	3.8	4.4	5.1
Male														
2001	7.2	18.9	2.1	0.7	1.0	1.9	3.4	3.1	3.7	4.8	5.3	6.3	6.8	6.7
2000[4]	7.2	16.8	2.1	0.7	1.1	2.1	3.5	3.7	3.9	5.1	5.6	6.5	6.7	7.7
1999[4]	7.6	16.2	2.3	0.9	1.2	2.3	3.4	3.7	3.7	4.5	6.1	6.0	7.2	7.8
Female														
2001	5.3	14.2	1.5	0.4	0.3	0.6	0.6	0.6	0.6	1.1	1.2	1.6	2.2	2.5
2000[4]	5.2	13.2	1.4	0.4	0.3	0.5	0.4	0.6	0.7	0.8	1.3	1.4	1.7	2.4
1999[4]	5.6	11.3	1.5	0.5	0.4	0.5	0.5	0.6	0.8	0.9	1.4	1.7	1.7	2.5
Hispanic														
Both sexes														
2001	2.7	9.5	1.4	*	*	1.1	1.8	1.5	2.0	2.6	2.8	3.2	4.1	3.3
2000[4]	2.7	7.5	1.4	*	*	1.1	2.2	1.9	2.5	2.7	3.2	2.8	2.6	3.2
1999[4]	2.6	8.6	1.9	*	*	1.1	1.9	1.7	1.5	2.5	2.4	2.9	2.6	3.0
Male														
2001	3.7	9.6	1.8	*	*	1.9	3.2	2.6	3.4	4.5	4.7	5.3	7.1	5.4
2000[4]	3.8	7.9	1.5	*	*	1.8	3.9	3.2	4.3	5.0	5.6	4.8	4.1	5.3
1999[4]	3.6	9.0	2.5	*	*	1.7	3.4	2.9	2.4	4.2	3.6	5.3	4.2	5.3
Female														
2001	1.6	9.5	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)						
Non-Hispanic Black						
Male						
2000[4]	6.8	*	*	*	*	*
1999[4]	9.6	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)						
All origins[3]						
Both sexes						
2001	6.3	7.8	12.3	22.5	42.2	115.0
2000[4]	6.3	7.8	12.2	22.0	41.4	117.1
1999[4]	6.3	8.9	13.3	23.6	46.9	128.9
Male						
2001	8.9	10.9	15.9	29.3	52.7	134.0
2000[4]	9.1	10.8	15.6	28.1	51.6	135.2
1999[4]	9.1	12.0	18.0	29.9	59.3	152.7
Female						
2001	3.8	5.2	9.5	17.7	35.9	107.0
2000[4]	3.8	5.2	9.6	17.8	35.4	109.8
1999[4]	3.7	6.3	9.6	19.3	39.6	119.3
Hispanic						
Both sexes						
2001	4.4	5.9	10.1	12.7	21.9	63.8
2000[4]	4.9	5.3	7.3	13.5	22.3	70.3
1999[4]	5.8	7.1	10.7	11.3	21.7	65.9
Male						
2001	6.0	8.5	13.7	17.7	26.0	68.6
2000[4]	8.3	7.8	9.7	16.2	*	88.7
1999[4]	7.4	10.4	16.8	*	*	84.1
Female						
2001	*	*	7.3	*	19.3	61.4

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)														
Hispanic														
Female														
2000[4]	1.4	7.2	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.6	8.1	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	6.7	18.3	1.9	0.6	0.7	1.3	2.0	1.9	2.2	2.9	3.3	4.0	4.5	4.6
2000[4]	6.7	16.9	1.8	0.6	0.8	1.4	1.9	2.2	2.3	3.0	3.4	4.0	4.2	5.1
1999[4]	7.1	15.0	1.9	0.7	0.9	1.5	2.0	2.2	2.4	2.7	3.9	3.9	4.5	5.1
Male														
2001	7.7	21.1	2.1	0.8	1.1	1.9	3.4	3.2	3.8	4.8	5.3	6.4	6.8	6.7
2000[4]	7.7	19.0	2.2	0.8	1.3	2.2	3.4	3.8	3.8	5.1	5.6	6.6	6.8	7.9
1999[4]	8.1	17.9	2.3	0.9	1.4	2.4	3.5	3.8	4.0	4.5	6.3	6.1	7.3	7.8
Female														
2001	5.8	15.4	1.6	0.4	0.3	0.6	0.6	0.6	0.6	1.2	1.3	1.6	2.3	2.6
2000[4]	5.7	14.7	1.4	0.4	0.3	0.6	0.5	0.6	0.8	0.9	1.3	1.4	1.7	2.4
1999[4]	6.2	12.0	1.5	0.5	0.5	0.5	0.5	0.7	0.8	1.0	1.5	1.8	1.8	2.7
Non-Hispanic White														
Both sexes														
2001	7.2	14.5	1.7	0.7	0.7	1.4	2.2	2.0	2.2	3.1	3.2	3.8	4.4	4.5
2000[4]	7.1	13.7	1.7	0.6	0.8	1.4	2.0	2.3	2.4	3.0	3.4	3.8	4.1	4.9
1999[4]	7.6	12.1	1.7	0.8	1.0	1.6	2.2	2.3	2.4	2.9	3.9	3.6	4.3	4.7
Male														
2001	8.1	16.8	2.0	0.9	1.1	2.1	3.7	3.3	3.8	5.1	5.2	6.2	6.5	6.4
2000[4]	8.0	16.0	2.1	0.8	1.3	2.3	3.5	4.0	4.0	5.2	5.5	6.1	6.5	7.4
1999[4]	8.4	14.2	2.0	1.0	1.4	2.6	3.7	4.1	4.0	4.8	6.3	5.5	7.0	7.1
Female														
2001	6.4	12.1	1.5	0.5	0.3	0.7	0.7	0.6	0.7	1.1	1.2	1.5	2.4	2.7
2000[4]	6.3	11.2	1.3	0.4	*	0.5	0.5	0.7	0.7	0.9	1.3	1.5	1.7	2.4
1999[4]	6.8	9.8	1.5	0.5	0.5	0.6	0.6	0.6	0.8	1.0	1.6	1.7	1.7	2.5
Non-Hispanic Black														
Both sexes														
2001	5.5	35.5	2.6	*	0.8	0.8	1.4	1.9	2.0	2.6	4.0	5.6	6.0	6.1
2000[4]	5.6	31.7	2.3	0.7	0.8	1.4	1.8	2.1	2.4	3.2	4.0	5.8	5.8	8.1
1999[4]	6.3	29.5	2.6	0.7	0.8	1.0	1.7	2.3	2.8	2.3	4.8	6.7	6.6	9.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)						
Hispanic						
Female						
2000[4]	*	*	*	11.5	19.7	61.3
1999[4]	*	*	*	11.6	19.7	57.2
Non-Hispanic[5]						
Both sexes						
2001	6.4	7.9	12.4	22.9	42.9	116.6
2000[4]	6.4	7.9	12.4	22.3	42.0	118.5
1999[4]	6.3	9.0	13.4	24.1	47.7	130.7
Male						
2001	9.1	11.0	15.9	29.8	53.6	136.5
2000[4]	9.1	10.9	15.8	28.5	52.4	136.9
1999[4]	9.1	12.0	18.0	30.7	60.3	154.8
Female						
2001	3.9	5.2	9.6	18.1	36.5	108.4
2000[4]	3.9	5.4	9.8	18.0	35.8	111.1
1999[4]	3.7	6.4	9.8	19.5	40.3	121.1
Non-Hispanic White						
Both sexes						
2001	6.0	7.8	12.0	22.7	43.2	119.5
2000[4]	6.2	7.5	12.0	21.9	42.5	122.0
1999[4]	5.7	8.6	12.7	23.3	47.6	134.3
Male						
2001	8.7	10.8	15.3	29.6	52.7	139.7
2000[4]	8.6	10.2	15.3	27.5	52.0	139.7
1999[4]	8.5	11.2	16.9	29.4	60.1	157.3
Female						
2001	3.6	5.1	9.2	17.9	37.4	111.2
2000[4]	4.0	5.2	9.4	18.0	36.9	114.9
1999[4]	3.2	6.3	9.4	19.0	40.3	125.2
Non-Hispanic Black						
Both sexes						
2001	10.5	10.5	18.9	27.5	44.7	96.5
2000[4]	8.8	12.6	18.1	30.8	39.4	90.9
1999[4]	11.0	14.1	23.2	36.3	54.0	106.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)														
Non-Hispanic Black														
Male														
2001	7.0	41.8	2.9	*	*	*	2.1	2.9	3.7	3.8	6.7	8.9	9.7	10.1
2000[4]	7.5	32.8	2.7	*	1.2	1.9	3.4	3.7	3.9	5.6	6.7	10.4	10.3	14.2
1999[4]	8.3	35.0	3.5	*	1.3	1.9	3.1	3.6	4.6	3.6	8.4	11.3	11.0	15.9
Female														
2001	4.2	28.9	2.3	*	*	*	*	*	*	1.6	1.6	2.7	2.8	2.8
2000[4]	3.9	30.7	2.0	*	*	*	*	*	*	*	1.6	1.8	*	3.0
1999[4]	4.5	23.9	*	*	*	*	*	*	*	*	1.6	2.8	2.9	4.3
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)														
All origins[3]														
Both sexes														
2001[6]	10.8	. . .	. . .	*	1.3	7.9	12.0	12.6	13.0	14.3	15.2	15.7	14.6	14.0
2000[4]	10.4	. . .	. . .	*	1.5	8.0	12.5	12.1	12.0	13.7	15.4	14.9	13.9	12.7
1999[4]	10.5	. . .	. . .	*	1.2	8.0	12.3	12.3	13.1	14.4	14.3	14.3	13.4	12.8
Male														
2001[6]	17.6	. . .	. . .	*	1.9	12.9	20.5	21.0	21.0	22.7	23.6	24.3	22.4	22.3
2000[4]	17.1	. . .	. . .	*	2.3	13.0	21.4	20.0	19.2	21.7	23.9	23.3	21.4	20.1
1999[4]	17.1	. . .	. . .	*	1.9	13.0	20.8	20.3	21.1	22.7	22.1	22.3	20.6	20.3
Female														
2001[6]	4.1	. . .	. . .	*	0.6	2.7	3.1	4.0	4.8	5.9	6.9	7.4	7.0	6.2
2000[4]	4.0	. . .	. . .	*	0.6	2.7	3.2	4.0	4.6	5.8	7.0	6.7	6.8	5.8
1999[4]	4.0	. . .	. . .	*	0.5	2.7	3.4	4.1	5.1	6.1	6.7	6.7	6.5	5.8
Hispanic														
Both sexes														
2001[6]	5.0	. . .	. . .	*	1.1	5.2	7.0	7.3	7.0	7.5	7.0	6.8	7.4	6.2
2000[4]	5.1	. . .	. . .	*	0.9	5.6	7.9	6.7	6.2	6.9	8.3	7.9	7.6	6.5
1999[4]	5.0	. . .	. . .	*	1.0	5.7	6.7	6.7	6.7	8.4	6.4	7.2	6.9	7.1
Male														
2001[6]	8.3	. . .	. . .	*	1.5	7.8	11.0	11.6	11.1	12.8	11.8	11.0	11.9	11.2
2000[4]	8.4	. . .	. . .	*	1.2	8.5	13.0	11.1	10.0	10.7	13.7	12.6	12.3	11.8
1999[4]	8.2	. . .	. . .	*	1.4	8.9	10.8	10.6	10.9	13.2	10.6	12.4	10.4	12.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)						
Non-Hispanic Black						
Male						
2001	14.8	15.9	25.1	35.6	73.6	115.5
2000[4]	14.2	18.7	23.6	47.7	64.4	124.5
1999[4]	16.1	21.3	35.3	51.2	70.0	155.7
Female						
2001	7.1	6.5	14.7	22.6	30.0	89.4
2000[4]	4.5	8.1	14.5	20.6	26.8	78.5
1999[4]	7.0	8.9	15.2	27.4	46.1	88.4
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)						
All origins[3]						
Both sexes						
2001[6]	12.0	12.7	13.9	16.4	18.9	17.5
2000[4]	11.4	11.3	13.8	16.5	19.3	19.6
1999[4]	11.4	12.4	14.5	17.4	19.2	19.3
Male						
2001[6]	19.5	22.7	26.7	33.8	44.2	51.1
2000[4]	18.7	19.8	26.1	34.9	44.8	57.5
1999[4]	19.1	22.1	27.7	35.6	44.1	57.1
Female						
2001[6]	5.2	4.1	3.7	4.3	3.8	3.4
2000[4]	4.9	4.0	4.1	3.8	4.2	4.2
1999[4]	4.4	4.2	4.1	4.8	4.6	4.1
Hispanic						
Both sexes						
2001[6]	6.2	6.7	8.7	6.8	12.4	13.8
2000[4]	5.3	7.2	8.6	9.8	15.0	*
1999[4]	6.5	7.9	5.9	11.3	16.4	15.4
Male						
2001[6]	11.9	13.8	17.4	15.7	28.6	41.9
2000[4]	10.1	14.5	17.0	19.9	38.3	*
1999[4]	11.6	15.8	11.2	23.4	37.7	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)														
Hispanic														
Female														
2001[6]	1.5	. . .	. . .	*	*	2.5	2.2	2.2	2.3	1.7	1.9	2.5	3.0	*
2000[4]	1.5	. . .	. . .	*	*	2.4	1.6	1.5	1.8	2.7	2.6	3.3	3.0	*
1999[4]	1.6	. . .	. . .	*	*	2.0	1.6	2.2	1.8	3.1	2.0	*	3.6	*
Non-Hispanic[5]														
Both sexes														
2001[6]	11.6	. . .	. . .	*	1.3	8.4	13.0	13.8	14.1	15.3	16.1	16.5	15.1	14.5
2000[4]	11.1	. . .	. . .	*	1.6	8.4	13.5	13.1	12.9	14.6	16.1	15.4	14.4	13.1
1999[4]	11.2	. . .	. . .	*	1.2	8.4	13.5	13.3	14.2	15.1	15.1	14.9	13.8	13.2
Male														
2001[6]	19.0	. . .	. . .	*	2.0	13.8	22.7	23.3	22.9	24.2	24.9	25.6	23.2	23.1
2000[4]	18.3	. . .	. . .	*	2.4	13.8	23.3	21.9	20.9	23.2	24.9	24.2	22.0	20.6
1999[4]	18.4	. . .	. . .	*	1.9	13.8	23.1	22.3	23.0	24.0	23.2	23.1	21.2	20.8
Female														
2001[6]	4.5	. . .	. . .	*	0.6	2.7	3.3	4.4	5.3	6.5	7.5	7.8	7.4	6.5
2000[4]	4.3	. . .	. . .	*	0.6	2.8	3.5	4.4	5.0	6.1	7.5	7.0	7.1	6.1
1999[4]	4.3	. . .	. . .	*	0.5	2.8	3.7	4.4	5.6	6.4	7.2	7.1	6.7	6.1
Non-Hispanic White														
Both sexes														
2001[6]	13.0	. . .	. . .	*	1.3	9.3	13.9	15.1	15.7	17.2	18.2	18.8	17.0	16.0
2000[4]	12.5	. . .	. . .	*	1.7	9.0	14.3	14.0	14.4	16.4	18.0	17.3	15.9	14.4
1999[4]	12.4	. . .	. . .	*	1.3	9.0	14.2	14.2	15.6	16.7	17.0	16.6	15.3	14.5
Male														
2001[6]	21.4	. . .	. . .	*	2.1	15.4	24.1	25.3	25.2	26.8	27.9	28.8	25.8	25.2
2000[4]	20.4	. . .	. . .	*	2.6	14.8	24.8	23.2	22.9	25.7	27.6	26.8	24.0	22.4
1999[4]	20.3	. . .	. . .	*	2.1	14.7	24.3	23.5	25.0	25.9	26.0	25.5	23.2	22.5
Female														
2001[6]	5.0	. . .	. . .	*	0.6	3.0	3.6	4.8	6.1	7.6	8.5	8.8	8.3	7.2
2000[4]	4.9	. . .	. . .	*	0.7	3.0	3.7	4.7	5.7	7.1	8.5	8.0	7.9	6.8
1999[4]	4.9	. . .	. . .	*	0.5	3.0	3.9	4.8	6.2	7.5	8.1	7.9	7.6	6.8
Non-Hispanic Black														
Both sexes														
2001[6]	5.3	. . .	. . .	*	1.2	4.5	10.5	9.7	8.7	8.1	6.9	6.4	6.0	5.7
2000[4]	5.5	. . .	. . .	*	1.2	5.7	11.4	10.2	7.9	7.2	8.2	6.6	6.4	4.6
1999[4]	5.5	. . .	. . .	*	0.9	5.9	11.1	10.0	9.0	8.6	7.0	6.5	4.8	5.2

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)						
Hispanic						
Female						
2001[6]	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001[6]	12.3	13.1	14.2	16.9	19.1	17.5
2000[4]	11.8	11.5	14.0	16.8	19.4	19.9
1999[4]	11.7	12.6	14.9	17.6	19.2	19.3
Male						
2001[6]	19.9	23.1	27.2	34.6	44.6	51.3
2000[4]	19.1	20.0	26.5	35.5	44.9	58.4
1999[4]	19.5	22.3	28.5	36.0	44.2	57.6
Female						
2001[6]	5.5	4.3	3.8	4.4	3.8	3.5
2000[4]	5.2	4.1	4.1	3.8	4.4	4.3
1999[4]	4.6	4.4	4.2	4.8	4.6	4.0
Non-Hispanic White						
Both sexes						
2001[6]	13.6	14.2	15.2	18.2	20.3	18.7
2000[4]	13.1	12.6	15.0	18.0	20.6	21.3
1999[4]	12.7	13.6	16.1	18.8	20.4	20.8
Male						
2001[6]	21.5	25.1	28.8	37.0	47.4	55.2
2000[4]	21.1	21.6	28.0	37.9	47.9	62.7
1999[4]	21.0	23.8	30.5	38.4	46.9	62.2
Female						
2001[6]	6.2	4.6	4.1	4.8	3.9	3.5
2000[4]	5.8	4.6	4.5	4.0	4.4	4.6
1999[4]	5.0	4.6	4.6	5.1	4.8	4.2
Non-Hispanic Black						
Both sexes						
2001[6]	4.4	4.6	6.6	5.1	6.3	*
2000[4]	3.8	4.4	6.3	5.4	6.6	*
1999[4]	4.7	6.6	5.0	6.4	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)														
Non-Hispanic Black														
Male														
2001[6]	9.3	. . .	. . .	*	1.7	7.5	20.0	17.7	15.5	14.2	10.8	9.9	9.7	10.0
2000[4]	9.6	. . .	. . .	*	2.1	9.9	20.1	18.2	14.1	12.9	13.7	11.6	10.4	8.3
1999[4]	9.9	. . .	. . .	*	1.4	10.1	20.3	18.7	16.3	16.1	11.1	11.1	8.8	9.4
Female														
2001[6]	1.7	. . .	. . .	*	*	*	*	2.4	2.5	2.6	3.4	3.4	2.8	*
2000[4]	1.7	. . .	. . .	*	*	1.5	3.1	3.0	2.3	2.2	3.3	2.2	3.0	*
1999[4]	1.6	. . .	. . .	*	*	1.6	2.3	2.1	2.6	1.9	3.4	2.5	*	*
Intentional self-harm (suicide) by discharge of firearms (X72-X74)														
All origins[3]														
Both sexes														
2001	5.9	. . .	. . .	*	0.4	4.1	6.6	6.7	6.3	6.4	7.0	7.5	8.0	8.4
2000[4]	5.9	. . .	. . .	*	0.5	4.4	7.2	6.3	6.0	6.7	7.4	7.4	7.8	7.4
1999[4]	5.9	. . .	. . .	*	0.5	4.9	7.2	6.6	6.5	7.1	6.7	7.4	7.2	8.1
Male														
2001	10.6	. . .	. . .	*	0.6	7.1	11.8	11.7	10.8	10.8	12.0	12.8	13.8	14.9
2000[4]	10.5	. . .	. . .	*	0.9	7.6	12.9	11.1	10.1	11.1	12.6	12.8	13.3	13.1
1999[4]	10.6	. . .	. . .	*	0.8	8.4	12.8	11.5	10.9	11.8	11.2	12.6	12.4	14.0
Female														
2001	1.5	. . .	. . .	*	0.2	1.0	1.0	1.5	1.7	2.1	2.2	2.3	2.4	2.4
2000[4]	1.5	. . .	. . .	*	0.2	1.1	1.3	1.5	1.9	2.3	2.4	2.2	2.5	2.1
1999[4]	1.5	. . .	. . .	*	0.2	1.1	1.4	1.6	1.9	2.3	2.2	2.4	2.1	2.6
Hispanic														
Both sexes														
2001	2.2	. . .	. . .	*	*	2.1	3.6	3.6	2.9	2.5	2.4	2.8	3.3	3.0
2000[4]	2.3	. . .	. . .	*	*	2.8	4.0	3.1	2.5	2.3	3.4	4.1	3.8	2.9
1999[4]	2.3	. . .	. . .	*	*	3.1	3.8	3.2	2.6	3.3	2.6	3.8	2.5	3.2
Male														
2001	3.8	. . .	. . .	*	*	3.5	5.9	6.0	4.9	4.5	4.4	5.0	5.8	6.4
2000[4]	4.1	. . .	. . .	*	*	4.6	6.8	5.3	4.1	3.7	6.0	6.9	6.8	5.9
1999[4]	4.0	. . .	. . .	*	*	5.0	6.5	5.2	4.4	5.3	4.5	6.6	4.2	6.0
Female														
2001	0.4	. . .	. . .	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	. . .	. . .	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)						
Non-Hispanic Black						
Male						
2001[6]	8.4	8.1	14.5	11.9	*	*
2000[4]	6.6	9.6	13.7	12.0	*	*
1999[4]	8.9	12.5	10.7	14.5	*	*
Female						
2001[6]	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Intentional self-harm (suicide) by discharge of firearms (X72-X74)						
All origins[3]						
Both sexes						
2001	8.0	9.0	10.3	12.4	14.1	12.2
2000[4]	7.7	8.3	10.3	12.4	14.0	13.2
1999[4]	7.5	8.7	10.9	12.9	13.5	12.6
Male						
2001	14.2	17.3	21.4	27.7	35.5	38.9
2000[4]	13.9	15.9	21.3	28.1	35.4	42.9
1999[4]	14.0	16.9	22.4	29.1	34.6	41.6
Female						
2001	2.3	1.9	1.4	1.7	1.3	1.0
2000[4]	2.1	1.8	1.6	1.4	1.4	1.1
1999[4]	1.6	1.7	1.8	1.6	1.2	0.9
Hispanic						
Both sexes						
2001	3.4	3.8	4.2	*	*	*
2000[4]	*	4.0	5.2	*	*	*
1999[4]	*	4.1	*	7.7	*	*
Male						
2001	6.8	8.2	9.6	*	*	*
2000[4]	*	8.6	11.2	*	*	*
1999[4]	*	8.5	*	18.7	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Intentional self-harm (suicide) by discharge of firearms (X72-X74)														
Hispanic														
Female														
1999[4]	0.5	. . .	. . .	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	6.5	. . .	. . .	*	0.5	4.5	7.2	7.4	6.9	7.0	7.6	7.9	8.4	8.8
2000[4]	6.4	. . .	. . .	*	0.6	4.7	7.9	7.0	6.6	7.3	7.9	7.7	8.1	7.8
1999[4]	6.4	. . .	. . .	*	0.5	5.2	7.9	7.3	7.1	7.5	7.1	7.8	7.5	8.4
Male														
2001	11.6	. . .	. . .	*	0.7	7.8	13.2	13.1	12.0	11.8	12.9	13.6	14.5	15.4
2000[4]	11.4	. . .	. . .	*	0.9	8.2	14.3	12.3	11.2	12.1	13.3	13.3	13.8	13.6
1999[4]	11.5	. . .	. . .	*	0.8	9.0	14.3	12.8	12.1	12.7	12.0	13.2	13.1	14.5
Female														
2001	1.6	. . .	. . .	*	0.2	1.0	1.1	1.7	1.9	2.3	2.4	2.5	2.6	2.6
2000[4]	1.6	. . .	. . .	*	0.2	1.1	1.4	1.6	2.1	2.6	2.6	2.3	2.6	2.3
1999[4]	1.6	. . .	. . .	*	*	1.1	1.5	1.7	2.1	2.5	2.3	2.5	2.2	2.8
Non-Hispanic White														
Both sexes														
2001	7.4	. . .	. . .	*	0.5	5.0	7.5	8.1	7.9	8.0	8.6	9.1	9.5	9.9
2000[4]	7.3	. . .	. . .	*	0.7	5.0	8.5	7.4	7.5	8.3	9.0	8.7	9.1	8.6
1999[4]	7.3	. . .	. . .	*	0.6	5.6	8.2	7.8	8.0	8.4	8.1	8.7	8.5	9.4
Male														
2001	13.2	. . .	. . .	*	0.8	8.6	13.6	14.2	13.4	13.2	14.5	15.6	16.3	17.1
2000[4]	12.9	. . .	. . .	*	1.2	8.6	15.3	12.9	12.5	13.5	15.0	14.9	15.4	14.9
1999[4]	12.9	. . .	. . .	*	1.0	9.7	14.7	13.6	13.3	13.8	13.5	14.7	14.5	15.8
Female														
2001	1.9	. . .	. . .	*	*	1.2	1.2	1.9	2.3	2.7	2.7	2.8	2.9	3.0
2000[4]	1.9	. . .	. . .	*	*	1.2	1.5	1.8	2.3	3.0	3.0	2.7	2.9	2.6
1999[4]	1.9	. . .	. . .	*	*	1.3	1.5	2.0	2.5	2.9	2.8	2.8	2.6	3.2
Non-Hispanic Black														
Both sexes														
2001	3.0	. . .	. . .	*	*	2.9	7.4	5.9	4.2	3.8	3.3	2.6	3.1	3.0
2000[4]	3.1	. . .	. . .	*	*	3.9	7.1	6.2	4.1	3.5	3.4	3.2	3.2	2.9
1999[4]	3.2	. . .	. . .	*	*	4.0	7.9	5.9	4.4	4.3	2.8	3.4	2.4	3.6
Male														
2001	5.6	. . .	. . .	*	*	5.2	14.5	11.0	8.2	7.3	5.6	4.3	5.6	6.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)						
Hispanic						
Female						
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	8.3	9.3	10.6	12.8	14.4	12.4
2000[4]	8.0	8.6	10.5	12.7	14.3	13.4
1999[4]	7.8	8.9	11.3	13.1	13.7	12.7
Male						
2001	14.6	17.8	22.0	28.6	36.1	39.7
2000[4]	14.4	16.3	21.7	28.9	36.1	43.8
1999[4]	14.5	17.3	23.2	29.4	35.1	42.2
Female						
2001	2.5	2.0	1.5	1.8	1.4	1.0
2000[4]	2.2	1.9	1.6	1.4	1.5	1.1
1999[4]	1.7	1.7	1.9	1.7	1.2	0.9
Non-Hispanic White						
Both sexes						
2001	9.3	10.4	11.5	13.9	15.3	13.4
2000[4]	9.0	9.5	11.4	13.8	15.5	14.5
1999[4]	8.6	9.8	12.3	14.1	14.6	13.7
Male						
2001	16.2	19.7	23.7	30.8	38.5	43.0
2000[4]	15.9	17.8	23.2	31.3	39.0	47.6
1999[4]	15.9	18.8	24.9	31.6	37.3	45.7
Female						
2001	2.9	2.2	1.6	1.9	1.4	1.1
2000[4]	2.5	2.2	1.9	1.5	1.6	1.2
1999[4]	1.9	1.8	2.2	1.8	1.3	1.0
Non-Hispanic Black						
Both sexes						
2001	2.4	3.0	4.8	4.4	6.1	*
2000[4]	3.1	3.2	4.6	*	*	*
1999[4]	3.3	4.1	4.0	5.1	*	*
Male						
2001	5.0	5.7	10.7	10.4	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Intentional self-harm (suicide) by discharge of firearms (X72-X74)														
Non-Hispanic Black														
Male														
2000[4]	5.7	. . .	. . .	*	*	7.3	13.0	11.8	7.5	6.6	6.3	5.8	5.4	5.8
1999[4]	6.1	. . .	. . .	*	*	7.1	14.8	11.6	8.4	8.4	5.3	6.1	4.9	7.3
Female														
2001	0.6	. . .	. . .	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.7	. . .	. . .	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.6	. . .	. . .	*	*	*	*	*	*	*	*	*	*	*
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)														
All origins[3]														
Both sexes														
2001[6]	4.8	. . .	. . .	*	0.9	3.8	5.4	5.9	6.7	7.9	8.2	8.2	6.6	5.6
2000[4]	4.5	. . .	. . .	*	0.9	3.6	5.3	5.8	6.0	7.0	7.9	7.4	6.2	5.2
1999[4]	4.5	. . .	. . .	*	0.7	3.2	5.1	5.7	6.7	7.3	7.7	6.9	6.2	4.7
Male														
2001[6]	7.1	. . .	. . .	*	1.3	5.8	8.7	9.3	10.2	11.9	11.6	11.5	8.6	7.5
2000[4]	6.6	. . .	. . .	*	1.4	5.4	8.5	8.9	9.1	10.6	11.3	10.6	8.1	6.9
1999[4]	6.6	. . .	. . .	*	1.1	4.6	8.0	8.8	10.2	10.9	10.8	9.6	8.1	6.4
Female														
2001[6]	2.6	. . .	. . .	*	0.4	1.7	2.0	2.5	3.1	3.9	4.8	5.1	4.6	3.8
2000[4]	2.5	. . .	. . .	*	0.4	1.7	1.9	2.6	2.7	3.4	4.6	4.4	4.3	3.7
1999[4]	2.5	. . .	. . .	*	0.3	1.6	2.0	2.5	3.2	3.7	4.5	4.2	4.4	3.2
Hispanic														
Both sexes														
2001[6]	2.8	. . .	. . .	*	1.0	3.1	3.5	3.7	4.1	5.0	4.6	4.0	4.1	3.1
2000[4]	2.8	. . .	. . .	*	0.6	2.8	3.9	3.6	3.7	4.6	4.9	3.9	3.7	3.5
1999[4]	2.7	. . .	. . .	*	*	2.6	2.8	3.6	4.1	5.0	3.8	3.4	4.4	3.9
Male														
2001[6]	4.4	. . .	. . .	*	1.3	4.3	5.1	5.5	6.2	8.3	7.4	6.0	6.1	4.8
2000[4]	4.3	. . .	. . .	*	*	3.8	6.2	5.9	5.9	7.0	7.7	5.8	5.6	5.9
1999[4]	4.2	. . .	. . .	*	*	4.0	4.3	5.4	6.5	7.9	6.0	5.9	6.2	6.7
Female														
2001[6]	1.2	. . .	. . .	*	*	1.9	1.4	1.4	1.8	*	*	*	*	*
2000[4]	1.1	. . .	. . .	*	*	1.7	*	*	*	1.9	1.9	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)						
Non-Hispanic Black						
Male						
2000[4]	5.7	7.2	11.3	*	*	*
1999[4]	6.7	8.0	10.0	12.1	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)						
All origins[3]						
Both sexes						
2001[6]	4.0	3.7	3.6	4.0	4.8	5.3
2000[4]	3.7	3.0	3.5	4.2	5.3	6.5
1999[4]	3.9	3.8	3.6	4.5	5.7	6.7
Male						
2001[6]	5.3	5.4	5.4	6.1	8.7	12.2
2000[4]	4.8	3.9	4.8	6.8	9.4	14.6
1999[4]	5.1	5.2	5.3	6.6	9.5	15.5
Female						
2001[6]	2.8	2.2	2.3	2.6	2.4	2.4
2000[4]	2.8	2.1	2.5	2.4	2.8	3.2
1999[4]	2.8	2.5	2.3	3.1	3.4	3.1
Hispanic						
Both sexes						
2001[6]	2.8	*	4.6	*	*	*
2000[4]	2.8	*	*	*	*	*
1999[4]	3.9	3.8	*	*	*	*
Male						
2001[6]	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	6.0	*	*	*	*	*
Female						
2001[6]	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)														
Hispanic														
Female														
1999[4]	1.1	. . .	. . .	*	*	*	*	1.4	1.4	1.9	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001[6]	5.1	. . .	. . .	*	0.9	3.9	5.8	6.4	7.1	8.3	8.5	8.6	6.7	5.7
2000[4]	4.8	. . .	. . .	*	1.0	3.7	5.5	6.2	6.3	7.3	8.2	7.7	6.3	5.3
1999[4]	4.7	. . .	. . .	*	0.7	3.3	5.6	6.0	7.1	7.6	8.0	7.2	6.3	4.8
Male														
2001[6]	7.5	. . .	. . .	*	1.3	6.0	9.5	10.2	10.9	12.4	12.0	12.0	8.8	7.6
2000[4]	6.9	. . .	. . .	*	1.5	5.6	9.0	9.6	9.7	11.1	11.6	10.9	8.2	6.9
1999[4]	6.9	. . .	. . .	*	1.1	4.8	8.9	9.4	10.9	11.3	11.3	9.9	8.2	6.3
Female														
2001[6]	2.9	. . .	. . .	*	0.4	1.7	2.2	2.7	3.4	4.2	5.1	5.4	4.8	3.9
2000[4]	2.7	. . .	. . .	*	0.4	1.7	2.1	2.8	2.9	3.6	4.9	4.6	4.5	3.8
1999[4]	2.7	. . .	. . .	*	0.3	1.7	2.2	2.6	3.4	3.9	4.8	4.5	4.5	3.3
Non-Hispanic White														
Both sexes														
2001[6]	5.6	. . .	. . .	*	0.8	4.4	6.5	7.0	7.8	9.2	9.6	9.6	7.4	6.2
2000[4]	5.2	. . .	. . .	*	0.9	4.0	5.8	6.6	6.9	8.1	9.1	8.6	6.8	5.9
1999[4]	5.2	. . .	. . .	*	0.7	3.4	6.0	6.4	7.7	8.3	8.9	7.9	6.8	5.1
Male														
2001[6]	8.2	. . .	. . .	*	1.2	6.8	10.5	11.0	11.9	13.6	13.4	13.3	9.5	8.1
2000[4]	7.5	. . .	. . .	*	1.4	6.2	9.5	10.3	10.4	12.1	12.6	11.9	8.6	7.6
1999[4]	7.4	. . .	. . .	*	1.1	5.0	9.6	9.9	11.6	12.1	12.6	10.7	8.7	6.7
Female														
2001[6]	3.2	. . .	. . .	*	0.4	1.7	2.4	2.9	3.8	4.8	5.8	6.0	5.4	4.3
2000[4]	3.0	. . .	. . .	*	0.4	1.7	2.1	2.9	3.4	4.0	5.5	5.3	5.0	4.2
1999[4]	3.0	. . .	. . .	*	*	1.7	2.3	2.8	3.7	4.5	5.3	5.1	5.0	3.6
Non-Hispanic Black														
Both sexes														
2001[6]	2.3	. . .	. . .	*	0.8	1.5	3.1	3.8	4.5	4.3	3.6	3.8	2.9	2.7
2000[4]	2.4	. . .	. . .	*	1.1	1.8	4.3	4.0	3.7	3.7	4.8	3.4	3.2	1.7
1999[4]	2.3	. . .	. . .	*	0.8	2.0	3.2	4.1	4.7	4.3	4.2	3.1	2.3	1.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)						
Hispanic						
Female						
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001[6]	4.1	3.7	3.6	4.1	4.7	5.1
2000[4]	3.8	2.9	3.5	4.1	5.1	6.5
1999[4]	3.9	3.7	3.6	4.6	5.5	6.6
Male						
2001[6]	5.3	5.4	5.2	6.0	8.6	11.6
2000[4]	4.7	3.7	4.7	6.7	8.8	14.6
1999[4]	5.0	5.0	5.3	6.7	9.1	15.4
Female						
2001[6]	3.0	2.3	2.3	2.7	2.4	2.5
2000[4]	2.9	2.2	2.5	2.4	2.9	3.2
1999[4]	2.8	2.6	2.3	3.1	3.5	3.1
Non-Hispanic White						
Both sexes						
2001[6]	4.3	3.8	3.7	4.3	5.0	5.3
2000[4]	4.2	3.1	3.6	4.2	5.1	6.8
1999[4]	4.1	3.8	3.8	4.7	5.8	7.0
Male						
2001[6]	5.3	5.4	5.1	6.2	9.0	12.3
2000[4]	5.1	3.8	4.8	6.6	8.9	15.2
1999[4]	5.2	5.0	5.6	6.8	9.6	16.5
Female						
2001[6]	3.3	2.4	2.5	2.9	2.5	2.4
2000[4]	3.3	2.4	2.6	2.5	2.8	3.4
1999[4]	3.1	2.7	2.4	3.3	3.5	3.2
Non-Hispanic Black						
Both sexes						
2001[6]	2.0	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	2.5	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)														
Non-Hispanic Black														
Male														
2001[6]	3.7	. . .	. . .	*	1.2	2.3	5.6	6.8	7.4	7.0	5.2	5.5	4.1	4.0
2000[4]	3.9	. . .	. . .	*	1.9	2.6	7.1	6.4	6.6	6.3	7.4	5.7	5.0	*
1999[4]	3.8	. . .	. . .	*	*	3.0	5.5	7.1	7.9	7.7	5.8	5.0	3.8	*
Female														
2001[6]	1.1	. . .	. . .	*	*	*	*	*	1.9	1.9	2.2	2.3	*	*
2000[4]	1.1	. . .	. . .	*	*	*	1.7	1.9	*	1.4	2.4	*	*	*
1999[4]	1.0	. . .	. . .	*	*	*	*	*	1.9	*	2.8	*	*	*
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)														
All origins[3]														
Both sexes														
2001[6]	7.1	8.2	2.7	0.7	0.9	9.4	17.3	14.9	11.5	10.4	8.6	7.0	5.5	4.4
2000[4]	6.0	9.2	2.3	0.7	1.1	9.5	16.0	11.9	9.0	7.6	6.6	5.3	4.0	3.4
1999[4]	6.1	8.7	2.5	0.9	1.2	10.4	15.6	12.1	9.1	7.7	6.6	5.1	4.0	3.2
Male														
2001[6]	11.1	9.5	3.0	0.7	1.1	15.7	28.9	24.1	17.7	15.3	12.6	10.6	8.3	6.7
2000[4]	9.3	10.4	2.5	0.7	1.4	15.5	26.7	19.6	14.0	10.9	9.7	7.8	5.9	5.2
1999[4]	9.3	9.6	2.5	0.9	1.5	16.9	25.7	19.5	13.8	11.0	9.5	7.7	5.8	5.1
Female														
2001[6]	3.3	6.9	2.4	0.6	0.7	2.7	5.1	5.5	5.2	5.5	4.7	3.6	2.7	2.3
2000[4]	2.8	7.9	2.1	0.6	0.8	3.1	4.7	4.1	4.0	4.4	3.6	2.8	2.1	1.7
1999[4]	2.9	7.8	2.4	0.9	0.9	3.5	5.1	4.5	4.3	4.3	3.6	2.5	2.2	1.4
Hispanic														
Both sexes														
2001[6]	9.0	6.7	2.3	0.5	1.1	13.8	22.3	16.1	11.7	11.0	9.6	8.1	6.2	4.4
2000[4]	8.3	7.0	1.9	0.7	1.6	15.0	19.1	13.7	11.1	8.8	8.6	7.6	5.6	5.2
1999[4]	8.4	7.0	1.8	0.8	2.0	15.7	19.6	14.8	10.5	9.1	8.4	7.9	5.0	5.2
Male														
2001[6]	14.5	7.9	2.5	*	1.5	23.9	36.1	25.3	17.6	16.5	15.2	12.6	9.6	6.6
2000[4]	13.4	6.6	2.0	*	2.3	25.7	30.9	22.2	17.5	13.5	13.6	11.1	9.0	8.8
1999[4]	13.8	6.7	1.7	*	2.8	26.0	31.8	24.3	16.6	14.1	13.6	11.9	8.0	8.6
Female														
2001[6]	3.2	5.5	2.1	*	*	2.6	5.3	5.2	4.9	5.0	3.7	3.7	3.0	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)						
Non-Hispanic Black						
Male						
2001[6]	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001[6]	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)						
All origins[3]						
Both sexes						
2001[6]	3.5	3.3	2.5	2.5	2.5	2.4
2000[4]	2.6	2.4	2.4	2.3	2.6	2.4
1999[4]	2.8	2.7	2.4	2.5	2.4	2.4
Male						
2001[6]	5.0	4.6	3.3	3.2	3.0	3.2
2000[4]	3.9	3.3	3.4	2.8	3.8	3.3
1999[4]	4.3	4.2	3.1	3.4	3.4	3.4
Female						
2001[6]	2.2	2.1	1.9	1.9	2.2	2.0
2000[4]	1.5	1.7	1.6	2.0	1.9	2.0
1999[4]	1.5	1.5	1.9	1.9	1.9	2.0
Hispanic						
Both sexes						
2001[6]	4.5	4.9	*	*	*	*
2000[4]	3.6	*	4.4	*	*	*
1999[4]	3.0	3.5	*	*	*	*
Male						
2001[6]	7.1	8.5	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001[6]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)														
Hispanic														
Female														
2000[4]	2.8	7.4	1.9	*	*	2.8	4.6	3.7	3.8	3.6	3.5	4.2	*	*
1999[4]	2.8	7.3	1.9	*	*	4.0	4.9	3.6	3.4	3.5	3.0	3.9	*	*
Non-Hispanic[5]														
Both sexes														
2001[6]	6.7	8.4	2.8	0.7	0.9	8.5	16.0	14.4	11.3	10.1	8.4	6.8	5.3	4.3
2000[4]	5.6	9.6	2.4	0.7	1.0	8.4	15.1	11.5	8.6	7.4	6.4	5.0	3.8	3.2
1999[4]	5.7	8.9	2.6	0.9	1.1	9.4	14.7	11.5	8.7	7.4	6.3	4.8	3.8	3.0
Male														
2001[6]	10.4	9.8	3.2	0.7	1.0	14.0	26.8	23.4	17.5	14.9	12.1	10.2	8.0	6.6
2000[4]	8.6	11.2	2.6	0.7	1.3	13.4	25.5	18.9	13.2	10.4	9.2	7.4	5.6	4.9
1999[4]	8.6	10.1	2.6	0.9	1.3	15.1	24.3	18.3	13.1	10.4	9.0	7.3	5.5	4.8
Female														
2001[6]	3.2	6.9	2.4	0.7	0.7	2.7	5.0	5.4	5.2	5.5	4.7	3.5	2.7	2.2
2000[4]	2.7	7.9	2.2	0.7	0.8	3.0	4.7	4.2	4.0	4.5	3.6	2.6	2.0	1.6
1999[4]	2.9	7.6	2.5	0.9	0.9	3.4	5.1	4.7	4.4	4.4	3.6	2.4	2.2	1.4
Non-Hispanic White														
Both sexes														
2001[6]	4.0	5.8	1.7	0.7	0.6	2.9	5.5	6.5	6.6	7.0	6.2	5.1	4.1	3.7
2000[4]	2.8	6.2	1.6	0.6	0.7	2.7	4.4	4.0	3.9	4.1	3.8	3.2	2.6	2.4
1999[4]	2.9	6.3	1.7	0.6	0.7	3.1	4.7	4.2	4.4	4.3	3.9	3.1	2.7	2.3
Male														
2001[6]	5.6	6.8	2.0	0.6	0.6	3.9	7.5	9.3	9.4	10.3	8.7	7.4	6.0	5.3
2000[4]	3.6	8.3	1.8	0.6	0.8	3.5	6.0	5.2	5.2	5.1	5.3	4.3	3.5	3.5
1999[4]	3.9	8.0	1.7	0.6	0.8	4.2	6.1	5.7	5.7	5.6	5.2	4.5	3.9	3.4
Female														
2001[6]	2.4	4.7	1.5	0.7	0.6	1.9	3.5	3.6	3.8	3.7	3.7	2.9	2.3	2.0
2000[4]	1.9	4.1	1.3	0.5	0.6	1.9	2.8	2.8	2.6	3.1	2.3	2.0	1.7	1.4
1999[4]	2.0	4.4	1.6	0.6	0.7	2.0	3.3	2.8	3.1	3.0	2.5	1.7	1.6	1.2
Non-Hispanic Black														
Both sexes														
2001[6]	22.7	19.5	7.1	0.9	1.9	33.6	64.9	54.6	36.5	29.0	22.0	18.2	13.9	10.0
2000[4]	22.1	24.1	6.1	1.2	2.6	34.1	66.4	50.4	34.3	26.6	22.3	17.1	12.4	9.2
1999[4]	21.7	20.0	6.4	2.0	2.6	37.7	63.0	48.8	32.6	25.7	21.5	16.4	11.2	9.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)						
Hispanic						
Female						
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001[6]	3.3	3.1	2.5	2.5	2.4	2.4
2000[4]	2.5	2.4	2.3	2.3	2.6	2.3
1999[4]	2.8	2.6	2.4	2.5	2.4	2.4
Male						
2001[6]	4.7	4.3	3.2	3.2	2.9	3.1
2000[4]	3.7	3.2	3.2	2.7	3.8	3.1
1999[4]	4.1	3.9	3.0	3.3	3.3	3.4
Female						
2001[6]	2.1	2.1	1.9	2.0	2.2	2.1
2000[4]	1.4	1.6	1.5	2.0	1.9	2.0
1999[4]	1.6	1.5	1.9	2.0	1.9	2.0
Non-Hispanic White						
Both sexes						
2001[6]	2.8	2.5	2.1	2.1	2.1	2.2
2000[4]	1.8	1.7	1.8	2.0	2.0	2.1
1999[4]	2.1	2.2	2.0	2.0	2.1	2.1
Male						
2001[6]	3.8	3.2	2.5	2.6	2.4	2.8
2000[4]	2.6	2.1	2.4	2.3	2.6	2.6
1999[4]	3.0	3.1	2.3	2.5	2.8	2.6
Female						
2001[6]	1.8	1.9	1.7	1.7	1.9	2.0
2000[4]	1.2	1.4	1.3	1.8	1.6	1.9
1999[4]	1.4	1.5	1.7	1.7	1.6	1.9
Non-Hispanic Black						
Both sexes						
2001[6]	7.5	7.6	6.6	7.1	6.3	*
2000[4]	7.5	8.1	7.0	5.4	9.8	*
1999[4]	7.6	6.7	6.1	8.4	6.2	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)														
Non-Hispanic Black														
Male														
2001[6]	39.3	21.7	8.4	*	2.8	59.7	120.3	99.7	64.5	44.2	34.7	30.1	23.6	17.6
2000[4]	38.3	24.9	6.1	*	3.3	59.0	122.1	93.5	59.9	43.8	34.8	29.2	22.5	16.3
1999[4]	37.2	20.1	6.2	2.0	3.5	64.6	116.0	87.2	56.2	41.1	34.6	27.7	18.5	17.2
Female														
2001[6]	7.6	17.1	5.7	*	*	6.6	11.8	13.6	11.4	15.6	10.7	7.8	5.7	3.8
2000[4]	7.4	23.3	6.0	1.5	1.8	8.6	13.7	11.4	11.3	11.3	11.2	6.5	3.8	3.5
1999[4]	7.7	20.0	6.7	1.9	1.6	10.1	12.8	14.1	11.4	12.1	10.1	6.6	5.1	*
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)														
All origins[3]														
Both sexes														
2001	4.0	*	0.4	0.3	0.6	7.5	13.6	10.3	6.6	5.1	3.7	2.7	2.1	1.5
2000[4]	3.8	*	0.2	0.2	0.7	7.7	12.7	9.2	6.3	4.7	3.8	2.9	2.1	1.7
1999[4]	3.9	*	0.3	0.3	0.8	8.5	12.5	9.2	6.1	4.6	3.7	2.8	2.0	1.6
Male														
2001	6.8	*	0.3	0.3	0.8	13.2	24.0	17.8	11.1	8.0	5.6	4.2	3.2	2.3
2000[4]	6.5	*	*	0.3	1.0	13.3	22.4	16.1	10.4	7.4	5.8	4.5	3.3	2.6
1999[4]	6.5	*	0.3	0.4	1.1	14.5	21.8	15.9	10.1	7.1	5.5	4.4	3.0	2.6
Female														
2001	1.3	*	0.4	0.2	0.3	1.5	2.7	2.5	2.1	2.2	1.7	1.1	1.0	0.7
2000[4]	1.3	*	*	0.2	0.3	1.7	2.7	2.2	2.1	1.9	1.7	1.3	1.0	0.9
1999[4]	1.3	*	0.3	0.2	0.5	2.2	2.8	2.3	2.1	2.2	1.8	1.3	1.1	0.6
Hispanic														
Both sexes														
2001	5.7	*	*	*	0.7	11.7	17.1	11.7	7.2	5.8	5.4	3.4	2.4	*
2000[4]	5.5	*	*	*	1.0	12.6	14.8	10.3	7.4	4.8	4.9	4.5	3.4	2.3
1999[4]	5.7	*	*	*	1.4	13.4	15.3	10.5	6.8	5.4	5.0	4.8	2.5	2.5
Male														
2001	9.9	*	*	*	*	20.4	28.8	19.8	12.0	9.4	8.8	5.5	3.9	*
2000[4]	9.6	*	*	*	1.7	21.9	24.9	17.5	12.3	7.5	8.1	7.0	5.6	*
1999[4]	9.8	*	*	*	2.1	22.9	25.4	18.1	11.1	9.1	7.9	6.9	4.2	*
Female														
2001	1.3	*	*	*	*	2.1	2.8	2.0	1.6	1.8	1.9	*	*	*
2000[4]	1.3	*	*	*	*	2.0	2.5	2.0	1.8	1.9	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)						
Non-Hispanic Black						
Male						
2001[6]	11.7	13.0	12.1	11.9	*	*
2000[4]	13.4	14.9	12.6	*	19.7	*
1999[4]	13.1	13.0	10.4	14.0	*	*
Female						
2001[6]	4.1	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)						
All origins[3]						
Both sexes						
2001	1.4	1.2	0.9	0.8	0.7	*
2000[4]	1.2	1.0	1.0	0.7	0.5	0.7
1999[4]	1.6	1.1	1.0	0.8	0.7	0.6
Male						
2001	2.0	1.8	1.1	1.2	*	*
2000[4]	1.8	1.3	1.3	0.8	*	*
1999[4]	2.5	1.5	1.4	1.1	*	*
Female						
2001	0.8	0.7	0.7	0.5	0.7	*
2000[4]	0.7	0.8	0.7	0.7	*	*
1999[4]	0.7	0.7	0.7	0.6	0.6	*
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)														
Hispanic														
Female														
1999[4]	1.4	*	*	*	*	2.6	3.2	1.7	1.8	*	1.9	2.6	*	*
Non-Hispanic[5]														
Both sexes														
2001	3.7	*	0.3	0.3	0.5	6.7	12.7	9.8	6.5	5.0	3.4	2.6	2.0	1.5
2000[4]	3.6	*	0.2	0.2	0.6	6.7	12.2	9.0	6.0	4.6	3.6	2.7	2.0	1.7
1999[4]	3.6	*	0.4	0.3	0.7	7.6	11.9	8.8	5.9	4.5	3.5	2.6	2.0	1.5
Male														
2001	6.3	*	0.3	0.4	0.8	11.7	22.6	17.1	10.8	7.8	5.2	4.1	3.2	2.3
2000[4]	6.0	*	*	0.2	0.9	11.5	21.6	15.7	10.0	7.3	5.5	4.2	3.1	2.5
1999[4]	6.0	*	0.4	0.4	1.0	12.9	20.9	15.3	9.8	6.7	5.2	4.1	2.9	2.5
Female														
2001	1.2	*	0.3	0.3	0.3	1.4	2.6	2.6	2.1	2.3	1.7	1.1	0.9	0.8
2000[4]	1.2	*	*	*	0.3	1.7	2.7	2.3	2.1	2.0	1.8	1.3	1.0	0.9
1999[4]	1.3	*	0.4	*	0.4	2.0	2.8	2.5	2.1	2.2	1.8	1.1	1.1	0.5
Non-Hispanic White														
Both sexes														
2001	1.6	*	0.2	0.3	0.4	1.8	3.2	2.8	2.6	2.7	2.1	1.7	1.4	1.2
2000[4]	1.4	*	*	0.2	0.3	1.7	2.7	2.5	2.4	2.4	2.1	1.7	1.4	1.3
1999[4]	1.5	*	0.2	0.2	0.4	2.0	2.9	2.7	2.6	2.5	2.1	1.6	1.4	1.1
Male														
2001	2.2	*	*	0.4	0.5	2.6	4.7	4.3	3.6	3.9	2.9	2.5	2.1	1.5
2000[4]	2.0	*	*	*	0.5	2.5	4.0	3.5	3.4	3.4	3.0	2.5	1.9	1.9
1999[4]	2.2	*	*	*	0.5	3.0	4.2	4.0	3.8	3.4	2.9	2.4	2.1	1.7
Female														
2001	0.9	*	*	*	*	0.9	1.6	1.4	1.6	1.5	1.3	0.9	0.8	0.8
2000[4]	0.9	*	*	*	*	0.9	1.5	1.4	1.4	1.5	1.3	1.0	0.9	0.8
1999[4]	0.9	*	*	*	0.3	0.9	1.5	1.3	1.4	1.6	1.3	0.8	0.7	0.5
Non-Hispanic Black														
Both sexes														
2001	16.3	*	*	*	1.4	29.5	57.6	45.3	27.9	19.1	11.8	8.5	6.6	4.4
2000[4]	16.0	*	*	*	1.8	29.5	57.3	42.5	26.6	17.8	12.9	9.7	6.9	4.3
1999[4]	15.9	*	*	0.7	1.9	33.0	55.8	40.7	24.3	16.3	12.5	9.7	6.1	4.6
Male														
2001	30.6	*	*	*	2.2	54.2	110.3	86.3	53.2	32.6	20.7	15.6	12.1	8.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)						
Hispanic						
Female						
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	1.4	1.1	0.9	0.9	0.7	*
2000[4]	1.2	1.0	0.9	0.7	0.6	0.7
1999[4]	1.5	1.1	1.1	0.8	0.7	0.5
Male						
2001	2.0	1.6	1.1	1.3	*	*
2000[4]	1.8	1.4	1.2	0.8	*	*
1999[4]	2.5	1.5	1.5	1.0	*	*
Female						
2001	0.8	0.7	0.7	0.6	*	*
2000[4]	0.6	0.7	0.7	0.7	*	*
1999[4]	0.7	0.6	0.7	0.7	*	*
Non-Hispanic White						
Both sexes						
2001	1.2	1.0	0.7	0.7	0.6	*
2000[4]	0.9	0.8	0.7	0.7	*	0.6
1999[4]	1.2	0.9	0.9	0.6	0.7	0.6
Male						
2001	1.7	1.2	0.8	1.0	*	*
2000[4]	1.2	1.0	0.9	0.8	*	*
1999[4]	1.7	1.3	1.2	*	*	*
Female						
2001	0.8	0.8	0.6	0.5	*	*
2000[4]	0.7	0.7	0.7	0.7	*	*
1999[4]	0.7	0.7	0.7	0.6	*	*
Non-Hispanic Black						
Both sexes						
2001	2.9	2.8	*	*	*	*
2000[4]	2.7	2.8	2.7	*	*	*
1999[4]	4.3	2.3	*	*	*	*
Male						
2001	5.0	6.0	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)														
Non-Hispanic Black														
Male														
2000[4]	30.1	*	*	*	2.8	53.3	109.1	82.3	49.4	32.6	22.0	17.1	13.1	7.3
1999[4]	29.3	*	*	*	2.9	58.3	105.7	77.2	45.3	28.0	21.0	17.4	9.8	9.6
Female														
2001	3.2	*	*	*	*	4.0	7.2	8.0	5.2	7.0	4.0	2.4	2.0	*
2000[4]	3.3	*	*	*	*	5.2	8.4	6.5	6.2	4.6	4.9	3.3	*	*
1999[4]	3.7	*	*	*	*	7.2	8.5	7.8	5.5	5.9	5.0	3.1	2.9	*
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)														
All origins[3]														
Both sexes														
2001[6]	3.1	8.0	2.3	0.4	0.3	1.8	3.7	4.6	4.9	5.2	4.9	4.4	3.4	2.9
2000[4]	2.1	8.9	2.1	0.4	0.5	1.8	3.2	2.7	2.8	2.9	2.9	2.4	1.8	1.6
1999[4]	2.2	8.5	2.1	0.6	0.4	1.9	3.1	2.9	3.0	3.0	2.9	2.3	1.9	1.6
Male														
2001[6]	4.3	9.2	2.7	0.4	0.3	2.4	4.9	6.3	6.7	7.3	7.0	6.4	5.0	4.4
2000[4]	2.8	10.0	2.3	0.5	0.4	2.2	4.4	3.5	3.6	3.5	3.9	3.3	2.6	2.6
1999[4]	2.8	9.4	2.1	0.5	0.4	2.4	3.9	3.6	3.7	3.9	4.0	3.3	2.8	2.5
Female														
2001[6]	2.0	6.7	2.0	0.4	0.4	1.2	2.4	3.0	3.2	3.2	2.9	2.5	1.8	1.5
2000[4]	1.5	7.7	2.0	0.4	0.5	1.4	2.0	1.9	2.0	2.4	1.8	1.5	1.0	0.8
1999[4]	1.6	7.6	2.1	0.7	0.5	1.4	2.2	2.2	2.3	2.2	1.8	1.3	1.1	0.9
Hispanic														
Both sexes														
2001[6]	3.3	6.2	1.9	*	*	2.1	5.2	4.4	4.5	5.3	4.2	4.7	3.8	3.0
2000[4]	2.7	6.9	1.8	*	*	2.4	4.3	3.3	3.7	4.0	3.7	3.1	2.2	2.9
1999[4]	2.7	6.7	1.6	*	*	2.3	4.3	4.2	3.7	3.6	3.4	3.2	2.5	2.6
Male														
2001[6]	4.5	7.2	2.2	*	*	3.5	7.4	5.4	5.6	7.2	6.5	7.1	5.7	4.4
2000[4]	3.8	6.6	1.8	*	*	3.8	6.0	4.7	5.2	6.0	5.4	4.1	3.5	4.6
1999[4]	4.0	6.2	1.5	*	*	3.2	6.4	6.2	5.5	5.1	5.7	5.0	3.8	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)						
Non-Hispanic Black						
Male						
2000[4]	5.5	5.3	*	*	*	*
1999[4]	8.5	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)						
All origins[3]						
Both sexes						
2001[6]	2.1	2.1	1.6	1.6	1.8	2.0
2000[4]	1.4	1.4	1.4	1.6	2.1	1.7
1999[4]	1.3	1.6	1.4	1.7	1.7	1.8
Male						
2001[6]	3.0	2.8	2.2	2.0	2.1	2.7
2000[4]	2.0	2.0	2.1	2.0	3.0	2.5
1999[4]	1.8	2.6	1.7	2.2	2.6	2.6
Female						
2001[6]	1.4	1.4	1.2	1.4	1.6	1.7
2000[4]	0.8	0.9	0.9	1.3	1.5	1.4
1999[4]	0.8	0.8	1.2	1.3	1.2	1.5
Hispanic						
Both sexes						
2001[6]	3.8	3.2	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001[6]	6.0	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)														
Hispanic														
Female														
2001[6]	1.9	5.2	1.6	*	*	*	2.5	3.2	3.2	3.2	1.8	2.3	*	*
2000[4]	1.6	7.2	1.7	*	*	*	2.1	1.7	2.0	1.8	2.0	*	*	*
1999[4]	1.4	7.3	1.8	*	*	1.4	1.7	1.9	1.6	2.1	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001[6]	3.1	8.2	2.4	0.4	0.3	1.8	3.3	4.6	4.9	5.1	4.9	4.2	3.3	2.8
2000[4]	2.0	9.2	2.2	0.4	0.4	1.7	3.0	2.6	2.6	2.8	2.8	2.3	1.8	1.5
1999[4]	2.1	8.7	2.2	0.6	0.4	1.8	2.8	2.6	2.8	2.9	2.8	2.2	1.8	1.5
Male														
2001[6]	4.2	9.5	2.8	0.4	0.2	2.2	4.2	6.3	6.7	7.1	6.9	6.1	4.9	4.3
2000[4]	2.6	10.6	2.4	0.4	0.3	1.9	3.9	3.2	3.2	3.1	3.8	3.2	2.5	2.4
1999[4]	2.6	10.0	2.3	0.5	0.3	2.3	3.3	3.0	3.3	3.7	3.8	3.1	2.6	2.3
Female														
2001[6]	2.0	6.8	2.0	0.4	0.4	1.3	2.4	2.8	3.1	3.2	3.0	2.4	1.7	1.4
2000[4]	1.5	7.8	2.0	0.5	0.5	1.4	2.0	1.9	1.9	2.5	1.8	1.4	1.0	0.7
1999[4]	1.6	7.3	2.2	0.7	0.4	1.4	2.3	2.2	2.4	2.1	1.8	1.2	1.1	0.8
Non-Hispanic White														
Both sexes														
2001[6]	2.4	5.6	1.5	0.3	0.3	1.2	2.4	3.6	4.0	4.3	4.1	3.4	2.7	2.5
2000[4]	1.3	6.0	1.4	0.3	0.4	1.0	1.7	1.6	1.5	1.7	1.7	1.5	1.2	1.1
1999[4]	1.4	6.2	1.4	0.4	0.3	1.2	1.9	1.6	1.8	1.8	1.8	1.5	1.4	1.2
Male														
2001[6]	3.4	6.7	1.8	*	*	1.3	2.8	5.0	5.7	6.4	5.8	4.9	3.9	3.8
2000[4]	1.6	8.1	1.7	0.4	0.3	1.0	2.0	1.7	1.8	1.8	2.3	1.9	1.6	1.6
1999[4]	1.7	7.8	1.5	0.4	*	1.3	1.9	1.7	1.9	2.2	2.3	2.1	1.8	1.7
Female														
2001[6]	1.5	4.5	1.2	0.4	0.4	1.0	1.9	2.2	2.3	2.2	2.4	2.0	1.5	1.2
2000[4]	1.0	3.9	1.2	*	0.4	1.0	1.4	1.4	1.2	1.7	1.0	1.1	0.9	0.5
1999[4]	1.1	4.4	1.4	0.5	0.4	1.1	1.8	1.4	1.7	1.4	1.2	0.9	0.9	0.7
Non-Hispanic Black														
Both sexes														
2001[6]	6.4	19.1	6.3	*	*	4.1	7.3	9.3	8.6	10.0	10.2	9.6	7.3	5.7

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)						
Hispanic						
Female						
2001[6]	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001[6]	2.0	1.9	1.6	1.6	1.7	2.1
2000[4]	1.3	1.4	1.3	1.5	2.1	1.6
1999[4]	1.2	1.6	1.3	1.7	1.7	1.8
Male						
2001[6]	2.6	2.6	2.1	1.9	2.0	2.7
2000[4]	1.9	1.9	2.0	1.9	3.1	2.4
1999[4]	1.6	2.4	1.6	2.3	2.6	2.6
Female						
2001[6]	1.3	1.4	1.2	1.4	1.6	1.8
2000[4]	0.8	0.9	0.8	1.2	1.5	1.3
1999[4]	0.8	0.9	1.1	1.3	1.2	1.5
Non-Hispanic White						
Both sexes						
2001[6]	1.6	1.5	1.3	1.3	1.5	1.9
2000[4]	0.9	0.9	1.0	1.3	1.6	1.5
1999[4]	1.0	1.3	1.1	1.4	1.4	1.5
Male						
2001[6]	2.2	2.1	1.7	1.6	1.8	2.3
2000[4]	1.3	1.1	1.5	1.5	2.2	1.9
1999[4]	1.3	1.8	1.2	1.8	2.1	*
Female						
2001[6]	1.0	1.1	1.1	1.2	1.2	1.7
2000[4]	0.5	0.7	0.7	1.1	1.2	1.3
1999[4]	0.7	0.8	1.0	1.1	1.0	1.4
Non-Hispanic Black						
Both sexes						
2001[6]	4.5	4.9	4.1	4.8	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)														
Non-Hispanic Black														
Both sexes														
2000[4]	6.1	22.9	5.7	0.9	0.8	4.6	9.0	7.8	7.7	8.8	9.4	7.4	5.5	5.0
1999[4]	5.8	19.7	5.7	1.3	0.7	4.7	7.2	8.1	8.3	9.4	9.1	6.7	5.2	4.4
Male														
2001[6]	8.7	21.4	7.7	*	*	5.6	10.0	13.4	11.3	11.6	14.1	14.6	11.5	8.9
2000[4]	8.2	22.5	5.7	*	*	5.7	13.0	11.2	10.5	11.2	12.8	12.2	9.5	9.0
1999[4]	7.9	19.7	5.6	1.2	*	6.3	10.3	10.0	10.9	13.0	13.6	10.3	8.8	7.7
Female														
2001[6]	4.4	16.8	5.0	*	*	2.6	4.6	5.6	6.2	8.6	6.7	5.4	3.7	3.0
2000[4]	4.1	23.3	5.8	*	*	3.4	5.3	4.8	5.1	6.7	6.4	3.3	2.0	*
1999[4]	4.0	19.6	5.7	1.4	*	2.9	4.4	6.3	5.9	6.2	5.1	3.5	2.1	*
Legal intervention (Y35,Y89.0)														
All origins[3]														
Both sexes														
2001	0.1	*	*	*	*	0.1	0.3	0.2	0.3	0.3	0.2	0.2	*	*
2000[4]	0.1	*	*	*	*	0.1	0.2	0.3	0.2	0.3	0.2	0.2	*	*
1999[4]	0.1	*	*	*	*	0.1	0.3	0.3	0.3	0.3	0.2	*	0.2	*
Male														
2001	0.3	*	*	*	*	0.2	0.6	0.5	0.6	0.6	0.4	0.3	*	*
2000[4]	0.3	*	*	*	*	0.3	0.5	0.6	0.5	0.5	0.3	0.4	*	*
1999[4]	0.3	*	*	*	*	0.2	0.5	0.7	0.7	0.6	0.4	*	0.4	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)						
Non-Hispanic Black						
Both sexes						
2000[4]	4.8	5.3	4.2	4.2	6.9	*
1999[4]	3.3	4.4	3.5	5.3	*	*
Male						
2001[6]	6.7	7.0	7.2	*	*	*
2000[4]	7.9	9.6	6.8	*	*	*
1999[4]	4.6	8.5	*	*	*	*
Female						
2001[6]	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Legal intervention (Y35,Y89.0)						
All origins[3]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Legal intervention (Y35,Y89.0)														
Hispanic														
Male														
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.1	*	*	*	*	*	0.3	0.2	0.3	0.3	0.2	0.2	*	*
2000[4]	0.1	*	*	*	*	0.1	0.2	0.3	0.2	0.3	0.2	0.2	*	*
1999[4]	0.1	*	*	*	*	*	0.3	0.3	0.3	0.3	0.2	*	0.2	*
Male														
2001	0.3	*	*	*	*	*	0.5	0.4	0.6	0.6	0.4	0.3	*	*
2000[4]	0.2	*	*	*	*	0.3	0.4	0.7	0.5	0.5	0.3	0.4	*	*
1999[4]	0.3	*	*	*	*	*	0.5	0.7	0.6	0.6	0.4	*	0.3	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	0.1	*	*	*	*	*	0.2	0.2	0.3	0.3	0.2	0.1	*	*
2000[4]	0.1	*	*	*	*	*	0.2	0.2	0.2	0.2	0.1	0.2	*	*
1999[4]	0.1	*	*	*	*	*	*	0.2	0.2	0.2	0.2	*	0.2	*
Male														
2001	0.2	*	*	*	*	*	0.5	*	0.5	0.5	0.4	0.3	*	*
2000[4]	0.2	*	*	*	*	*	*	0.5	0.4	0.4	0.3	0.4	*	*
1999[4]	0.2	*	*	*	*	*	*	0.4	0.5	0.4	0.3	*	0.3	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	0.7	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Legal intervention (Y35,Y89.0)						
Hispanic						
Male						
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic White						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Legal intervention (Y35,Y89.0)														
Non-Hispanic Black														
Both sexes														
1999[4]	0.4	*	*	*	*	*	0.8	0.9	*	0.7	*	*	*	*
Male														
2001	0.5	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.6	*	*	*	*	*	*	*	*	1.5	*	*	*	*
1999[4]	0.8	*	*	*	*	*	1.7	2.0	*	1.6	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)														
All origins[3]														
Both sexes														
2001	1.5	2.1	0.3	*	0.2	0.7	1.4	1.6	2.0	3.0	3.2	3.0	2.0	1.1
2000[4]	1.4	1.7	0.3	*	0.2	0.6	1.3	1.6	1.8	2.6	3.1	2.7	1.7	1.1
1999[4]	1.4	1.4	0.3	*	0.2	0.8	1.5	1.6	2.1	2.8	2.9	2.7	1.6	1.0
Male														
2001	2.0	2.4	*	*	0.3	1.0	2.1	2.4	2.7	3.9	4.4	4.2	2.5	1.5
2000[4]	1.9	1.4	0.3	*	0.4	0.9	2.0	2.3	2.6	3.7	4.2	3.8	2.3	1.6
1999[4]	2.0	1.7	0.3	*	0.3	1.3	2.2	2.3	2.8	3.9	4.0	3.7	2.1	1.5
Female														
2001	0.9	1.7	0.3	*	*	0.3	0.7	0.9	1.3	2.1	2.0	1.8	1.5	0.6
2000[4]	0.8	2.0	*	*	*	0.3	0.6	0.9	1.1	1.6	1.9	1.7	1.1	0.7
1999[4]	0.9	1.2	0.3	*	*	0.3	0.6	0.9	1.3	1.7	1.9	1.7	1.0	0.6
Hispanic														
Both sexes														
2001	0.8	*	*	*	*	*	0.6	0.9	1.0	1.6	1.7	1.6	1.6	*
2000[4]	0.6	*	*	*	*	*	0.7	0.9	1.0	1.5	1.0	*	*	*
1999[4]	0.7	*	*	*	*	0.7	1.0	1.0	1.2	1.3	1.1	*	*	*
Male														
2001	1.1	*	*	*	*	*	*	1.3	1.5	2.2	3.0	*	*	*
2000[4]	0.9	*	*	*	*	*	1.1	1.2	1.6	2.1	*	*	*	*
1999[4]	1.1	*	*	*	*	*	1.6	1.5	1.7	2.0	1.8	*	*	*
Female														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Legal intervention (Y35,Y89.0)						
Non-Hispanic Black						
Both sexes						
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)						
All origins[3]						
Both sexes						
2001	0.7	0.6	0.5	0.7	1.1	1.0
2000[4]	0.7	0.6	0.6	0.6	0.7	1.3
1999[4]	0.8	0.8	0.5	0.7	0.9	1.4
Male						
2001	0.8	0.9	0.6	0.9	1.5	1.5
2000[4]	0.9	0.9	0.8	1.0	1.1	1.7
1999[4]	1.1	1.1	0.6	1.0	1.3	2.1
Female						
2001	0.5	*	*	0.6	0.8	0.8
2000[4]	0.5	*	*	*	*	1.2
1999[4]	0.6	0.5	*	0.5	0.7	1.1
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)														
Hispanic														
Female														
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	1.6	2.4	0.3	*	0.2	0.7	1.6	1.8	2.2	3.2	3.3	3.1	2.0	1.1
2000[4]	1.5	1.9	0.3	*	0.2	0.6	1.5	1.7	1.9	2.8	3.3	2.8	1.8	1.1
1999[4]	1.5	1.6	0.3	*	0.2	0.8	1.5	1.7	2.2	3.0	3.1	2.8	1.6	1.0
Male														
2001	2.1	2.7	*	*	0.3	1.0	2.5	2.7	2.9	4.2	4.5	4.5	2.5	1.5
2000[4]	2.0	1.6	0.4	*	0.4	1.0	2.2	2.5	2.7	3.9	4.5	4.0	2.4	1.6
1999[4]	2.1	1.9	*	*	0.3	1.3	2.4	2.5	3.0	4.2	4.2	3.9	2.1	1.4
Female														
2001	1.0	2.0	0.4	*	*	0.3	0.7	1.0	1.5	2.3	2.2	1.8	1.5	0.7
2000[4]	0.9	2.2	*	*	*	0.3	0.7	0.9	1.2	1.7	2.1	1.8	1.2	0.7
1999[4]	0.9	*	*	*	*	0.3	0.7	0.9	1.5	1.9	2.0	1.8	1.1	0.7
Non-Hispanic White														
Both sexes														
2001	1.6	1.7	0.3	*	0.2	0.8	1.8	2.0	2.4	3.3	3.4	3.1	1.9	1.1
2000[4]	1.4	1.5	0.2	*	0.2	0.7	1.5	1.8	2.0	2.8	3.2	2.8	1.7	1.0
1999[4]	1.4	1.3	*	*	0.2	0.8	1.6	1.7	2.3	3.0	3.0	2.6	1.4	1.0
Male														
2001	2.1	2.0	*	*	0.3	1.2	2.6	2.8	3.1	4.1	4.4	4.3	2.3	1.4
2000[4]	2.0	*	*	*	0.4	1.1	2.2	2.6	2.7	3.9	4.3	3.7	2.2	1.3
1999[4]	2.0	*	*	*	0.4	1.4	2.4	2.4	3.0	3.9	4.0	3.4	1.8	1.2
Female														
2001	1.1	*	*	*	*	0.4	0.9	1.1	1.6	2.5	2.4	1.9	1.6	0.8
2000[4]	0.9	*	*	*	*	0.4	0.8	1.1	1.3	1.8	2.0	1.8	1.2	0.7
1999[4]	1.0	*	*	*	*	*	0.7	1.1	1.6	2.0	1.9	1.9	1.1	0.8
Non-Hispanic Black														
Both sexes														
2001	1.8	5.9	*	*	*	*	1.1	1.5	2.0	3.9	4.0	4.1	3.3	1.5
2000[4]	1.9	*	*	*	*	*	1.4	1.7	2.2	3.4	4.9	4.4	2.9	2.4
1999[4]	2.0	*	*	*	*	0.8	1.7	2.0	2.6	4.0	4.7	4.8	3.4	1.6

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)						
Hispanic						
Female						
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	0.6	0.6	0.5	0.7	1.1	0.9
2000[4]	0.7	0.5	0.6	0.6	0.7	1.4
1999[4]	0.8	0.8	0.5	0.7	0.9	1.4
Male						
2001	0.8	0.9	0.6	0.9	1.5	*
2000[4]	0.9	0.8	0.8	1.0	*	1.7
1999[4]	1.1	1.1	0.6	1.0	1.4	2.1
Female						
2001	0.5	*	*	0.6	0.8	0.8
2000[4]	0.5	*	*	*	*	1.2
1999[4]	0.6	0.5	*	0.5	0.7	1.1
Non-Hispanic White						
Both sexes						
2001	0.6	0.5	0.5	0.7	1.1	1.0
2000[4]	0.7	0.5	0.5	0.6	0.7	1.3
1999[4]	0.8	0.8	0.4	0.7	1.0	1.4
Male						
2001	0.7	0.8	*	1.0	1.5	*
2000[4]	0.9	0.7	0.7	1.0	*	*
1999[4]	1.0	1.1	*	1.0	1.3	2.1
Female						
2001	0.6	*	*	0.6	0.8	0.8
2000[4]	0.5	*	*	*	*	1.2
1999[4]	0.7	0.5	*	*	0.7	1.1
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)														
Non-Hispanic Black														
Male														
2001	2.8	6.4	*	*	*	*	2.0	2.5	2.8	6.0	6.6	7.1	5.4	*
2000[4]	2.9	*	*	*	*	*	2.6	2.8	3.5	5.0	7.3	7.1	4.6	4.1
1999[4]	3.1	*	*	*	*	1.3	2.8	3.6	3.8	6.4	6.7	8.4	5.8	*
Female														
2001	0.9	*	*	*	*	*	*	*	*	2.1	1.8	*	*	*
2000[4]	1.0	*	*	*	*	*	*	*	*	2.0	2.8	2.0	*	*
1999[4]	0.9	*	*	*	*	*	*	*	1.5	1.8	3.0	1.8	*	*
Discharge of firearms, undetermined intent (Y22-Y24)														
All origins[3]														
Both sexes														
2001	0.1	*	*	*	*	0.1	0.2	0.2	*	0.1	*	*	*	*
2000[4]	0.1	*	*	*	*	0.1	0.2	0.2	0.1	*	0.1	*	*	*
1999[4]	0.1	*	*	*	*	0.3	0.3	0.2	*	0.1	0.1	*	*	*
Male														
2001	0.1	*	*	*	*	0.2	0.4	0.3	*	0.2	*	*	*	*
2000[4]	0.1	*	*	*	*	0.2	0.4	0.3	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	0.6	0.5	0.3	*	0.2	0.2	*	*	*
Female														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)						
Non-Hispanic Black						
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Discharge of firearms, undetermined intent (Y22-Y24)						
All origins[3]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Discharge of firearms, undetermined intent (Y22-Y24)														
Hispanic														
Female														
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	0.1	*	*	*	*	0.1	0.2	0.2	*	0.2	*	*	*	*
2000[4]	0.1	*	*	*	*	*	0.3	0.2	*	*	0.1	*	*	*
1999[4]	0.1	*	*	*	*	0.3	0.3	0.2	*	0.1	0.1	*	*	*
Male														
2001	0.1	*	*	*	*	*	0.4	0.3	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	0.4	0.3	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	0.6	0.5	0.3	*	0.2	0.2	*	*	*
Female														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	0.1	*	*	*	*	*	0.2	0.2	*	0.2	*	*	*	*
2000[4]	0.1	*	*	*	*	*	0.2	0.2	*	*	*	*	*	*
1999[4]	0.1	*	*	*	*	0.3	0.3	0.2	*	0.1	0.1	*	*	*
Male														
2001	0.1	*	*	*	*	*	0.3	0.3	*	*	*	*	*	*
2000[4]	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	0.5	0.5	*	*	*	0.2	*	*	*
Female														
2001	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	0.1	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.2	*	*	*	*	0.7	*	*	*	*	*	*	*	*
Male														
2001	0.2	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Discharge of firearms, undetermined intent (Y22-Y24)						
Hispanic						
Female						
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic White						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Discharge of firearms, undetermined intent (Y22-Y24)														
Non-Hispanic Black														
Male														
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)														
All origins[3]														
Both sexes														
2001	1.4	2.1	0.3	*	0.1	0.5	1.2	1.5	1.9	2.9	3.1	2.9	1.9	1.0
2000[4]	1.3	1.7	0.3	*	0.2	0.5	1.1	1.4	1.7	2.6	3.0	2.7	1.6	1.1
1999[4]	1.3	1.4	0.3	*	0.1	0.5	1.1	1.4	2.0	2.7	2.8	2.6	1.5	1.0
Male														
2001	1.9	2.4	*	*	0.2	0.8	1.8	2.1	2.6	3.8	4.2	4.1	2.4	1.4
2000[4]	1.8	1.4	0.3	*	0.3	0.7	1.6	2.0	2.4	3.6	4.1	3.7	2.1	1.5
1999[4]	1.8	1.7	*	*	0.2	0.7	1.7	2.0	2.7	3.7	3.8	3.5	2.0	1.4
Female														
2001	0.9	1.7	0.3	*	*	0.3	0.6	0.8	1.3	2.0	2.0	1.7	1.5	0.6
2000[4]	0.8	2.0	*	*	*	0.3	0.5	0.8	1.0	1.6	1.9	1.6	1.1	0.7
1999[4]	0.8	1.2	0.3	*	*	0.2	0.5	0.8	1.3	1.7	1.8	1.7	1.0	0.6
Hispanic														
Both sexes														
2001	0.7	*	*	*	*	*	*	0.8	0.9	1.5	1.7	1.5	1.5	*
2000[4]	0.6	*	*	*	*	*	*	0.8	1.0	1.4	1.0	*	*	*
1999[4]	0.6	*	*	*	*	*	0.7	0.9	1.2	1.2	1.1	*	*	*
Male														
2001	1.1	*	*	*	*	*	*	1.2	1.5	2.1	3.0	*	*	*
2000[4]	0.8	*	*	*	*	*	*	1.1	1.6	2.0	*	*	*	*
1999[4]	1.0	*	*	*	*	*	1.2	1.4	1.7	1.9	1.8	*	*	*
Female														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Discharge of firearms, undetermined intent (Y22-Y24)						
Non-Hispanic Black						
Male						
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)						
All origins[3]						
Both sexes						
2001	0.6	0.5	0.4	0.6	1.0	1.0
2000[4]	0.7	0.5	0.5	0.5	0.7	1.3
1999[4]	0.7	0.7	0.4	0.6	0.8	1.3
Male						
2001	0.7	0.8	0.6	0.7	1.3	*
2000[4]	0.9	0.7	0.7	0.8	*	*
1999[4]	0.9	0.9	*	0.9	*	1.8
Female						
2001	0.5	*	*	0.6	0.7	0.8
2000[4]	0.4	*	*	*	*	1.2
1999[4]	0.5	0.5	*	*	0.7	1.1
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)														
Hispanic														
Female														
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	1.5	2.4	0.3	*	0.2	0.6	1.4	1.6	2.1	3.1	3.2	3.0	1.9	1.0
2000[4]	1.4	1.9	0.3	*	0.2	0.5	1.2	1.5	1.8	2.7	3.2	2.8	1.7	1.0
1999[4]	1.4	1.6	0.3	*	0.1	0.5	1.2	1.5	2.1	2.9	3.0	2.7	1.5	1.0
Male														
2001	2.0	2.7	*	*	0.2	0.8	2.0	2.3	2.8	4.0	4.4	4.4	2.4	1.4
2000[4]	1.9	1.6	0.4	*	0.3	0.7	1.8	2.2	2.5	3.8	4.4	3.9	2.2	1.4
1999[4]	1.9	1.9	*	*	*	0.7	1.8	2.1	2.8	3.9	4.0	3.7	2.1	1.4
Female														
2001	1.0	2.0	0.4	*	*	0.3	0.7	0.9	1.5	2.2	2.1	1.8	1.5	0.7
2000[4]	0.9	2.2	*	*	*	0.3	0.6	0.9	1.1	1.7	2.0	1.7	1.2	0.7
1999[4]	0.9	*	*	*	*	*	0.6	0.9	1.4	1.8	2.0	1.8	1.0	0.6
Non-Hispanic White														
Both sexes														
2001	1.5	1.7	0.3	*	0.2	0.7	1.6	1.8	2.3	3.1	3.3	3.0	1.8	1.0
2000[4]	1.4	1.5	0.2	*	0.2	0.6	1.3	1.7	1.9	2.7	3.1	2.7	1.6	0.9
1999[4]	1.3	1.3	*	*	*	0.6	1.3	1.6	2.2	2.8	2.8	2.6	1.4	0.9
Male														
2001	2.0	2.0	*	*	*	1.0	2.2	2.5	3.0	3.9	4.3	4.1	2.2	1.3
2000[4]	1.8	*	*	*	0.3	0.9	1.9	2.3	2.5	3.8	4.1	3.6	2.0	1.2
1999[4]	1.8	*	*	*	*	0.9	1.9	2.1	2.8	3.7	3.8	3.2	1.7	1.2
Female														
2001	1.1	*	*	*	*	0.4	0.9	1.1	1.6	2.3	2.3	1.9	1.5	0.8
2000[4]	0.9	*	*	*	*	0.3	0.7	1.0	1.3	1.7	2.0	1.8	1.2	0.7
1999[4]	0.9	*	*	*	*	*	0.6	1.1	1.5	1.9	1.8	1.9	1.0	0.7
Non-Hispanic Black														
Both sexes														
2001	1.7	5.9	*	*	*	*	0.8	1.3	2.0	3.9	4.0	4.1	3.2	*
2000[4]	1.7	*	*	*	*	*	0.9	1.4	2.0	3.3	4.8	4.3	2.9	2.1
1999[4]	1.8	*	*	*	*	*	1.2	1.6	2.4	3.9	4.6	4.6	3.4	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)						
Hispanic						
Female						
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	0.6	0.5	0.4	0.6	1.0	0.9
2000[4]	0.6	0.5	0.5	0.5	0.7	1.3
1999[4]	0.7	0.7	0.4	0.6	0.8	1.3
Male						
2001	0.7	0.8	*	0.7	1.3	*
2000[4]	0.9	0.7	0.7	0.9	*	*
1999[4]	0.9	0.9	*	0.8	*	1.7
Female						
2001	0.5	*	*	0.6	0.8	0.8
2000[4]	0.4	*	*	*	*	1.2
1999[4]	0.6	0.5	*	*	0.7	1.1
Non-Hispanic White						
Both sexes						
2001	0.6	0.5	0.4	0.6	1.0	0.9
2000[4]	0.6	0.4	0.5	0.5	0.7	1.2
1999[4]	0.7	0.7	0.4	0.6	0.8	1.3
Male						
2001	0.6	0.6	*	*	1.4	*
2000[4]	0.8	0.6	0.6	0.9	*	*
1999[4]	0.8	0.9	*	0.8	*	*
Female						
2001	0.6	*	*	0.6	*	0.8
2000[4]	*	*	*	*	*	1.2
1999[4]	0.7	0.5	*	*	0.7	1.1
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)														
Non-Hispanic Black														
Male														
2001	2.6	6.4	*	*	*	*	1.5	2.3	2.7	6.0	6.5	7.1	5.1	*
2000[4]	2.6	*	*	*	*	*	1.6	2.2	3.1	4.7	7.2	7.0	4.4	3.6
1999[4]	2.8	*	*	*	*	*	1.8	2.9	3.5	6.2	6.6	8.0	5.8	*
Female														
2001	0.9	*	*	*	*	*	*	*	*	2.1	1.7	*	*	*
2000[4]	0.9	*	*	*	*	*	*	*	*	2.0	2.6	2.0	*	*
1999[4]	0.9	*	*	*	*	*	*	*	1.4	1.8	2.8	1.8	*	*
Operations of war and their sequelae (Y36,Y89.1)														
All origins[3]														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)						
Non-Hispanic Black						
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Operations of war and their sequelae (Y36,Y89.1)						
All origins[3]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Operations of war and their sequelae (Y36,Y89.1)														
Hispanic														
Female														
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic White														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic Black														
Both sexes														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Operations of war and their sequelae (Y36,Y89.1)						
Hispanic						
Female						
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic White						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Non-Hispanic Black						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Operations of war and their sequelae (Y36,Y89.1)														
Non-Hispanic Black														
Male														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Complications of medical and surgical care (Y40-Y84,Y88)														
All origins[3]														
Both sexes														
2001	1.1	0.6	0.2	*	*	*	0.1	0.1	0.3	0.3	0.4	0.7	0.8	1.3
2000[4]	1.1	*	0.1	*	*	*	0.1	0.2	0.2	0.3	0.4	0.6	0.9	1.2
1999[4]	1.0	1.4	0.2	*	*	*	*	0.1	0.2	0.3	0.4	0.5	0.7	1.1
Male														
2001	1.0	*	*	*	*	*	*	*	0.3	0.4	0.4	0.7	0.8	1.3
2000[4]	1.0	*	*	*	*	*	*	0.2	0.2	0.3	0.4	0.6	1.0	1.2
1999[4]	1.0	1.6	*	*	*	*	*	*	0.3	0.3	0.4	0.4	0.7	1.2
Female														
2001	1.1	*	*	*	*	*	*	*	0.2	0.3	0.5	0.6	0.8	1.2
2000[4]	1.1	*	*	*	*	*	*	*	0.2	0.3	0.4	0.7	0.9	1.1
1999[4]	1.1	1.2	*	*	*	*	*	0.2	*	0.3	0.4	0.5	0.7	0.9
Hispanic														
Both sexes														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Male														
2001	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	0.4	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Operations of war and their sequelae (Y36,Y89.1)						
Non-Hispanic Black						
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Complications of medical and surgical care (Y40-Y84,Y88)						
All origins[3]						
Both sexes						
2001	1.9	2.6	3.6	5.9	8.3	12.8
2000[4]	2.0	2.8	4.3	5.8	8.0	13.6
1999[4]	1.7	2.4	4.0	5.5	8.4	12.7
Male						
2001	2.0	2.6	3.8	6.3	9.9	14.5
2000[4]	2.0	2.8	5.3	7.0	10.4	16.0
1999[4]	1.7	2.3	4.5	6.4	10.1	16.5
Female						
2001	1.7	2.6	3.5	5.7	7.3	12.1
2000[4]	2.0	2.7	3.6	4.9	6.7	12.7
1999[4]	1.6	2.5	3.6	4.8	7.4	11.2
Hispanic						
Both sexes						
2001	*	*	*	*	*	*
2000[4]	*	3.7	*	*	*	*
1999[4]	*	*	*	*	*	*
Male						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*
1999[4]	*	*	*	*	*	*
Female						
2001	*	*	*	*	*	*
2000[4]	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Complications of medical and surgical care (Y40-Y84,Y88)														
Hispanic														
Female														
1999[4]	0.3	*	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic[5]														
Both sexes														
2001	1.2	0.7	*	*	*	*	0.1	*	0.3	0.3	0.4	0.7	0.8	1.3
2000[4]	1.2	*	*	*	*	*	*	0.2	0.3	0.3	0.4	0.7	1.0	1.2
1999[4]	1.1	1.6	0.2	*	*	*	*	0.2	0.2	0.3	0.4	0.5	0.7	1.1
Male														
2001	1.1	*	*	*	*	*	*	*	0.3	0.4	0.4	0.7	0.8	1.4
2000[4]	1.1	*	*	*	*	*	*	0.3	0.3	0.3	0.4	0.6	1.0	1.2
1999[4]	1.0	1.8	*	*	*	*	*	*	0.4	0.3	0.4	0.4	0.7	1.2
Female														
2001	1.2	*	*	*	*	*	*	*	0.3	0.3	0.5	0.6	0.8	1.2
2000[4]	1.2	*	*	*	*	*	*	*	0.3	0.3	0.4	0.7	0.9	1.2
1999[4]	1.2	1.4	*	*	*	*	*	*	*	0.3	0.4	0.6	0.8	0.9
Non-Hispanic White														
Both sexes														
2001	1.2	*	*	*	*	*	*	*	0.3	0.3	0.4	0.6	0.7	1.2
2000[4]	1.2	*	*	*	*	*	*	0.2	0.3	0.3	0.3	0.6	0.9	1.1
1999[4]	1.1	1.3	*	*	*	*	*	*	0.2	0.3	0.4	0.4	0.6	0.9
Male														
2001	1.0	*	*	*	*	*	*	*	0.3	0.3	0.3	0.6	0.7	1.2
2000[4]	1.2	*	*	*	*	*	*	*	*	0.3	0.4	0.6	1.0	1.2
1999[4]	1.1	*	*	*	*	*	*	*	0.4	0.3	0.3	0.3	0.6	1.0
Female														
2001	1.3	*	*	*	*	*	*	*	*	0.3	0.4	0.5	0.7	1.2
2000[4]	1.2	*	*	*	*	*	*	*	*	0.3	0.3	0.6	0.9	1.0
1999[4]	1.2	*	*	*	*	*	*	*	*	0.2	0.4	0.5	0.6	0.8
Non-Hispanic Black														
Both sexes														
2001	1.4	*	*	*	*	*	*	*	*	0.8	0.9	1.6	1.6	2.8
2000[4]	1.4	*	*	*	*	*	*	*	*	*	*	1.2	1.7	2.7
1999[4]	1.3	*	*	*	*	*	*	*	*	*	0.8	0.9	1.8	2.7
Male														
2001	1.5	*	*	*	*	*	*	*	*	*	*	1.8	*	3.5

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)						
Hispanic						
Female						
1999[4]	*	*	*	*	*	*
Non-Hispanic[5]						
Both sexes						
2001	1.9	2.6	3.7	5.9	8.3	12.9
2000[4]	2.1	2.7	4.4	5.8	8.1	13.7
1999[4]	1.7	2.5	4.2	5.6	8.6	12.8
Male						
2001	2.0	2.6	4.0	6.3	9.9	14.5
2000[4]	2.1	2.8	5.3	6.9	10.4	16.1
1999[4]	1.7	2.4	4.8	6.5	10.3	16.8
Female						
2001	1.8	2.5	3.5	5.6	7.4	12.2
2000[4]	2.1	2.6	3.7	5.0	6.8	12.7
1999[4]	1.6	2.5	3.7	5.0	7.6	11.2
Non-Hispanic White						
Both sexes						
2001	1.7	2.4	3.3	5.8	8.2	12.2
2000[4]	1.8	2.5	4.1	5.5	8.0	13.2
1999[4]	1.6	2.4	3.9	5.3	8.2	12.1
Male						
2001	1.7	2.5	3.3	6.2	10.1	13.5
2000[4]	1.8	2.5	5.0	6.8	10.6	15.9
1999[4]	1.6	2.3	4.5	6.3	9.9	15.8
Female						
2001	1.6	2.3	3.2	5.5	7.1	11.6
2000[4]	1.8	2.5	3.5	4.5	6.5	12.0
1999[4]	1.5	2.5	3.5	4.7	7.2	10.7
Non-Hispanic Black						
Both sexes						
2001	4.2	4.6	8.5	8.1	11.0	23.7
2000[4]	4.8	4.8	8.0	10.5	10.6	21.3
1999[4]	2.6	3.5	7.3	9.4	12.3	20.4
Male						
2001	5.2	5.2	11.4	*	*	30.0

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages[1]	Under 1 year[2]	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Complications of medical and surgical care (Y40-Y84,Y88)														
Non-Hispanic Black														
Male														
2000[4]	1.2	*	*	*	*	*	*	*	*	*	*	*	*	*
1999[4]	1.2	*	*	*	*	*	*	*	*	*	*	*	*	*
Female														
2001	1.4	*	*	*	*	*	*	*	*	*	*	*	*	*
2000[4]	1.6	*	*	*	*	*	*	*	*	*	*	*	*	3.2
1999[4]	1.4	*	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig291R. Deaths Rates for 113 Selected Causes, by 5-Year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)						
Non-Hispanic Black						
Male						
2000[4]	4.7	6.4	9.6	10.1	*	*
1999[4]	*	*	9.3	10.2	*	23.8
Female						
2001	3.5	4.2	6.7	8.2	11.2	21.4
2000[4]	4.9	*	7.0	10.7	10.8	22.2
1999[4]	*	*	6.0	9.0	11.0	19.1

0.0 Quantity more than zero but less than 0.05
* Figure does not meet standards of reliability or precision.
. . . Category not applicable.

[1] Figures for age not stated included in "All ages" but not distributed among age groups.
[2] Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).
[3] All origins include origin not stated; specified origins exclude origin not stated.
[4] Rates are revised and may differ from rates previously published; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."
[5] Includes races other than white and black.
[6] Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

Source: CDC/NCHS, National Vital Statistics System, Mortality.