Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
All causes													
All origins													
Both sexes													
2001	2,416,425	27,568	5,107	7,095	32,252	41,683	91,674	168,065	244,139	430,960	701,929	665,531	422
2000	2,403,351	28,035	4,979	7,413	31,307	40,451	89,798	160,341	240,846	441,209	700,445	658,171	356
1999	2,391,399	27,937	5,249	7,595	30,656	41,066	89,256	152,974	238,979	452,600	698,590	646,141	356
Male													
2001	1,183,421	15,477	2,899	4,168	23,963	28,757	58,164	104,848	144,958	241,581	340,742	217,533	331
2000	1,177,578	15,718	2,824	4,401	23,071	27,890	57,297	100,398	143,321	247,408	340,219	214,742	289
1999	1,175,460	15,646	2,975	4,492	22,414	28,276	57,118	95,659	142,724	254,920	340,970	209,989	277
Female													
2001	1,233,004	12,091	2,208	2,927	8,289	12,926	33,510	63,217	99,181	189,379	361,187	447,998	91
2000	1,225,773	12,317	2,155	3,012	8,236	12,561	32,501	59,943	97,525	193,801	360,226	443,429	67
1999	1,215,939	12,291	2,274	3,103	8,242	12,790	32,138	57,315	96,255	197,680	357,620	436,152	79
Hispanic													
Both sexes													
2001	113,413	4,680	917	1,038	5,058	5,797	8,441	11,362	13,337	20,172	23,637	18,907	67
2000	107,254	4,598	872	1,065	4,793	5,509	8,085	10,356	12,733	19,420	21,951	17,818	54
1999	103,740	4,412	882	1,014	4,505	5,397	7,798	9,876	12,319	19,201	20,912	17,358	66
Male													
2001	63,317	2,601	518	600	4,019	4,366	5,895	7,477	8,026	11,231	11,759	6,767	58
2000	60,172	2,514	475	622	3,837	4,200	5,597	6,808	7,760	10,842	10,973	6,493	51
1999	57,991	2,411	481	592	3,547	4,084	5,476	6,497	7,490	10,783	10,317	6,253	60
Female													
2001	50,096	2,079	399	438	1,039	1,431	2,546	3,885	5,311	8,941	11,878	12,140	9
2000	47,082	2,084	397	443	956	1,309	2,488	3,548	4,973	8,578	10,978	11,325	3

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
All causes													
Hispanic													
Female													
1999	45,749	2,001	401	422	958	1,313	2,322	3,379	4,829	8,418	10,595	11,105	6
Non-Hispanic [1]													
Both sexes													
2001	2,295,244	22,629	4,171	6,033	27,059	35,656	82,763	155,916	229,812	409,500	676,512	645,020	173
2000	2,287,846	23,074	4,086	6,316	26,374	34,741	81,182	149,141	227,118	420,369	676,632	638,673	140
1999	2,279,325	23,126	4,343	6,560	26,043	35,471	80,949	142,302	225,659	431,952	675,730	627,035	155
Male													
2001	1,115,683	12,739	2,371	3,554	19,836	24,225	51,918	96,815	136,214	229,539	328,079	210,256	137
2000	1,112,704	13,007	2,338	3,770	19,136	23,548	51,304	92,984	134,890	235,648	328,284	207,688	107
1999	1,112,718	12,998	2,479	3,885	18,797	24,047	51,272	88,567	134,540	243,210	329,682	203,128	113
Female													
2001	1,179,561	9,890	1,800	2,479	7,223	11,431	30,845	59,101	93,598	179,961	348,433	434,764	36
2000	1,175,142	10,067	1,748	2,546	7,238	11,193	29,878	56,157	92,228	184,721	348,348	430,985	33
1999	1,166,607	10,128	1,864	2,675	7,246	11,424	29,677	53,735	91,119	188,742	346,048	423,907	42
Non-Hispanic white													
Both sexes													
2001	1,962,810	13,311	2,772	4,151	18,730	24,191	59,546	117,086	184,109	346,389	603,143	589,269	113
2000	1,959,919	13,513	2,641	4,397	18,261	23,191	58,231	111,964	182,348	356,869	604,658	583,757	89
1999	1,953,197	13,553	2,819	4,488	17,868	23,986	57,630	106,808	180,793	367,490	604,344	573,327	91
Male													
2001	945,967	7,508	1,592	2,428	13,542	16,613	38,213	73,731	109,993	196,158	294,013	192,084	92
2000	944,781	7,660	1,541	2,652	13,066	15,934	37,622	70,788	109,031	202,038	294,842	189,537	70
1999	944,913	7,718	1,605	2,643	12,677	16,449	37,291	67,151	108,692	209,155	296,528	184,938	66

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
All causes													
Non-Hispanic white													
Female													
2001	1,016,843	5,803	1,180	1,723	5,188	7,578	21,333	43,355	74,116	150,231	309,130	397,185	21
2000	1,015,138	5,853	1,100	1,745	5,195	7,257	20,609	41,176	73,317	154,831	309,816	394,220	19
1999	1,008,284	5,835	1,214	1,845	5,191	7,537	20,339	39,657	72,101	158,335	307,816	388,389	25
Non-Hispanic black													
Both sexes													
2001	284,343	8,277	1,156	1,557	7,024	9,859	20,436	34,098	39,523	53,803	61,592	46,968	50
2000	282,676	8,525	1,217	1,620	6,926	10,103	20,279	32,911	38,867	54,645	60,874	46,660	49
1999	281,979	8,589	1,283	1,771	6,980	9,992	20,770	31,377	39,063	55,567	60,634	45,895	58
Male													
2001	143,971	4,651	655	941	5,377	6,548	11,971	20,307	22,600	28,446	28,083	14,352	40
2000	143,297	4,757	672	959	5,224	6,649	12,061	19,647	22,402	28,745	27,715	14,431	35
1999	143,883	4,765	734	1,081	5,286	6,633	12,384	19,026	22,493	29,259	27,552	14,628	42
Female													
2001	140,372	3,626	501	616	1,647	3,311	8,465	13,791	16,923	25,357	33,509	32,616	10
2000	139,379	3,768	545	661	1,702	3,454	8,218	13,264	16,465	25,900	33,159	32,229	14
1999	138,096	3,824	549	690	1,694	3,359	8,386	12,351	16,570	26,308	33,082	31,267	16
Origin not stated [2]													
Both sexes													
2001	7,768	259	19	24	135	230	470	787	990	1,288	1,780	1,604	182
2000	8,251	363	21	32	140	201	531	844	995	1,420	1,862	1,680	162
1999	8,334	399	24	21	108	198	509	796	1,001	1,447	1,948	1,748	135
Male													
2001	4,421	137	10	14	108	166	351	556	718	811	904	510	136

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
All causes													
Origin not stated [2]													
Male													
2000	4,702	197	11	9	98	142	396	606	671	918	962	561	131
1999	4,751	237	15	15	70	145	370	595	694	927	971	608	104
Female													
2001	3,347	122	9	10	27	64	119	231	272	477	876	1,094	46
2000	3,549	166	10	23	42	59	135	238	324	502	900	1,119	31
1999	3,583	162	9	6	38	53	139	201	307	520	977	1,140	31
Salmonella infections (A01-A02)													
All origins													
Both sexes													
2001	41	2	-	-	-	2	1	3	3	7	14	9	-
2000	28	3	2	-	1	1	-	4	2	7	5	3	-
1999	38	3	-	-	2	-	2	2	3	7	12	7	-
Male													
2001	21	-	-	-	-	1	1	3	2	6	7	1	-
2000	15	2	-	-	1	1	-	2	2	5	1	1	-
1999	21	2	-	-	2	-	1	2	2	5	5	2	-
Female													
2001	20	2	-	-	-	1	-	-	1	1	7	8	-
2000	13	1	2	-	-	-	-	2	-	2	4	2	-
1999	17	1	-	-	-	-	1	-	1	2	7	5	-
Hispanic													
Both sexes													
2001	5	-	-	-	-	-	-	1	1	1	2	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Salmonella infections (A01-A02)													
Hispanic													
Both sexes													
2000	2	2	-	-	-	-	-	-	-	-	-	-	-
1999	4	-	-	-	-	-	-	-	1	1	2	-	-
Male													
2001	5	-	-	-	-	-	-	1	1	1	2	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	-	-	-	-	-	-	1	1	1	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	1	-	-
Non-Hispanic [1]													
Both sexes													
2001	36	2	-	-	-	2	1	2	2	6	12	9	-
2000	26	1	2	-	1	1	-	4	2	7	5	3	-
1999	34	3	-	-	2	-	2	2	2	6	10	7	-
Male													
2001	16	-	-	-	-	1	1	2	1	5	5	1	-
2000	14	1	-	-	1	1	-	2	2	5	1	1	-
1999	18	2	-	-	2	-	1	2	1	4	4	2	-
Female													
2001	20	2	-	-	-	1	-	-	1	1	7	8	-
2000	12	-	2	-	-	-	-	2	-	2	4	2	-
1999	16	1	-	-	-	-	1	-	1	2	6	5	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Salmonella infections (A01-A02)													
Non-Hispanic white													
Both sexes													
2001	28	1	-	-	-	2	-	2	2	3	10	8	-
2000	23	-	2	-	-	1	-	4	2	6	5	3	-
1999	28	3	-	-	1	-	2	2	1	4	8	7	-
Male													
2001	13	-	-	-	-	1	-	2	1	3	5	1	-
2000	12	-	-	-	-	1	-	2	2	5	1	1	-
1999	13	2	-	-	1	-	1	2	-	2	3	2	-
Female													
2001	15	1	-	-	-	1	-	-	1	-	5	7	-
2000	11	-	2	-	-	-	-	2	-	1	4	2	-
1999	15	1	-	-	-	-	1	-	1	2	5	5	-
Non-Hispanic black													
Both sexes													
2001	5	1	-	-	-	-	-	-	-	1	2	1	-
2000	2	1	-	-	1	-	-	-	-	-	-	-	-
1999	4	-	-	-	1	-	-	-	-	1	2	-	-
Male													
2001	1	-	-	-	-	-	-	-	-	1	-	-	-
2000	2	1	-	-	1	-	-	-	-	-	-	-	-
1999	3	-	-	-	1	-	-	-	-	1	1	-	-
Female													
2001	4	1	-	-	-	-	-	-	-	-	2	1	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Salmonella infections (A01-A02)													
Non-Hispanic black													
Female													
1999	1	-	-	-	-	-	-	-	-	-	1	-	-
Origin not stated [2]													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Shigellosis and amebiasis (A03,A06)													
All origins													
Both sexes													
2001	7	1	1	1	-	-	-	3	-	-	1	-	-
2000	12	-	-	-	-	-	1	1	1	1	7	1	-
1999	10	-	1	1	-	-	1	-	1	1	1	4	-
Male													
2001	6	1	1	1	-	-	-	3	-	-	-	-	-
2000	5	-	-	-	-	-	1	-	1	-	3	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Shigellosis and amebiasis (A03,A06)													
All origins													
Male													
1999	6	-	1	1	-	-	1	-	1	1	-	1	-
Female													
2001	1	-	-	-	-	-	-	-	-	-	1	-	-
2000	7	-	-	-	-	-	-	1	-	1	4	1	-
1999	4	-	-	-	-	-	-	-	-	-	1	3	-
Hispanic													
Both sexes													
2001	3	-	1	1	-	-	-	1	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	4	-	1	1	-	-	1	-	-	-	1	-	-
Male													
2001	3	-	1	1	-	-	-	1	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	1	1	-	-	1	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	1	-	-
Non-Hispanic [1]													
Both sexes													
2001	4	1	-	-	-	-	-	2	-	-	1	-	-
2000	12	-	-	-	-	-	1	1	1	1	7	1	-
1999	6	-	-	-	-	-	-	-	1	1	-	4	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Shigellosis and amebiasis (A03,A06)													
Non-Hispanic [1]													
Male													
2001	3	1	-	-	-	-	-	2	-	-	-	-	-
2000	5	-	-	-	-	-	1	-	1	-	3	-	-
1999	3	-	-	-	-	-	-	-	1	1	-	1	-
Female													
2001	1	-	-	-	-	-	-	-	-	-	1	-	-
2000	7	-	-	-	-	-	-	1	-	1	4	1	-
1999	3	-	-	-	-	-	-	-	-	-	-	3	-
Non-Hispanic white													
Both sexes													
2001	4	1	-	-	-	-	-	2	-	-	1	-	-
2000	10	-	-	-	-	-	-	1	1	-	7	1	-
1999	6	-	-	-	-	-	-	-	1	1	-	4	-
Male													
2001	3	1	-	-	-	-	-	2	-	-	-	-	-
2000	4	-	-	-	-	-	-	-	1	-	3	-	-
1999	3	-	-	-	-	-	-	-	1	1	-	1	-
Female													
2001	1	-	-	-	-	-	-	-	-	-	1	-	-
2000	6	-	-	-	-	-	-	1	-	-	4	1	-
1999	3	-	-	-	-	-	-	-	-	-	-	3	-
Non-Hispanic black													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Shigellosis and amebiasis (A03,A06)													
Non-Hispanic black													
Both sexes													
2000	1	-	-	-	-	-	1	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	1	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated [2]													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Certain other intestinal infections (A04,A07-A09)													
All origins													
Both sexes													
2001	1,537	10	18	4	-	4	12	42	81	235	566	565	-
2000	1,328	13	10	-	1	3	19	29	70	222	462	499	-
1999	1,108	12	8	3	5	6	7	23	49	176	390	429	-
Male													
2001	572	7	11	2	-	2	8	15	35	108	227	157	-
2000	524	10	6	-	-	3	10	16	39	105	195	140	-
1999	420	8	4	3	3	5	4	11	26	74	169	113	-
Female													
2001	965	3	7	2	-	2	4	27	46	127	339	408	-
2000	804	3	4	-	1	-	9	13	31	117	267	359	-
1999	688	4	4	-	2	1	3	12	23	102	221	316	-
Hispanic													
Both sexes													
2001	53	1	1	-	-	-	4	2	4	8	16	17	-
2000	46	3	2	-	-	-	2	-	8	9	11	11	-
1999	34	3	1	-	2	1	-	4	5	7	4	7	-
Male													
2001	19	1	1	-	-	-	3	-	2	1	5	6	-
2000	23	2	1	-	-	-	1	-	5	6	4	4	-
1999	17	2	-	-	2	1	-	2	2	3	2	3	-
Female													
2001	34	-	-	-	-	-	1	2	2	7	11	11	-
2000	23	1	1	-	-	-	1	-	3	3	7	7	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Certain other intestinal infections (A04,A07-A09)													
Hispanic													
Female													
1999	17	1	1	-	-	-	-	2	3	4	2	4	-
Non-Hispanic [1]													
Both sexes													
2001	1,474	9	16	4	-	4	8	40	77	225	546	545	-
2000	1,277	10	8	-	1	3	17	29	62	211	449	487	-
1999	1,071	9	7	3	3	5	7	19	44	169	386	419	-
Male													
2001	550	6	10	2	-	2	5	15	33	105	222	150	-
2000	497	8	5	-	-	3	9	16	34	98	189	135	-
1999	403	6	4	3	1	4	4	9	24	71	167	110	-
Female													
2001	924	3	6	2	-	2	3	25	44	120	324	395	-
2000	780	2	3	-	1	-	8	13	28	113	260	352	-
1999	668	3	3	-	2	1	3	10	20	98	219	309	-
Non-Hispanic white													
Both sexes													
2001	1,357	5	13	3	-	2	6	35	62	202	509	520	-
2000	1,175	5	5	-	1	2	9	22	54	191	430	456	-
1999	990	5	6	1	2	5	4	15	36	155	360	401	-
Male													
2001	497	3	8	1	-	2	3	12	29	94	203	142	-
2000	453	4	4	-	-	2	6	12	29	89	180	127	-
1999	373	4	3	1	1	4	3	6	20	69	158	104	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Certain other intestinal infections (A04,A07-A09)													
Non-Hispanic white													
Female													
2001	860	2	5	2	-	-	3	23	33	108	306	378	-
2000	722	1	1	-	1	-	3	10	25	102	250	329	-
1999	617	1	3	-	1	1	1	9	16	86	202	297	-
Non-Hispanic black													
Both sexes													
2001	98	3	3	1	-	2	2	5	13	22	28	19	-
2000	83	4	2	-	-	1	6	7	6	15	16	26	-
1999	74	4	-	2	1	-	3	3	6	14	25	16	-
Male													
2001	42	2	2	1	-	-	2	3	4	10	12	6	-
2000	36	3	1	-	-	1	2	4	4	6	8	7	-
1999	25	2	-	2	-	-	1	2	2	2	9	5	-
Female													
2001	56	1	1	-	-	2	-	2	9	12	16	13	-
2000	47	1	1	-	-	-	4	3	2	9	8	19	-
1999	49	2	-	-	1	-	2	1	4	12	16	11	-
Origin not stated [2]													
Both sexes													
2001	10	-	1	-	-	-	-	-	-	2	4	3	-
2000	5	-	-	-	-	-	-	-	-	2	2	1	-
1999	3	-	-	-	-	-	-	-	-	-	-	3	-
Male													
2001	3	-	-	-	-	-	-	-	-	2	-	1	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Certain other intestinal infections (A04,A07-A09)													
Origin not stated [2]													
Male													
2000	4	-	-	-	-	-	-	-	-	1	2	1	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	7	-	1	-	-	-	-	-	-	-	4	2	-
2000	1	-	-	-	-	-	-	-	-	1	-	-	-
1999	3	-	-	-	-	-	-	-	-	-	-	3	-
Tuberculosis (A16-A19)													
All origins													
Both sexes													
2001	764	2	1	-	12	25	40	84	116	152	206	125	1
2000	776	1	6	1	11	20	42	91	81	155	252	115	1
1999	930	2	2	3	11	19	57	106	130	173	253	173	1
Male													
2001	453	-	-	-	8	14	27	63	85	98	100	57	1
2000	468	-	1	-	7	14	28	76	59	87	139	56	1
1999	584	-	2	2	4	11	43	87	96	116	137	85	1
Female													
2001	311	2	1	-	4	11	13	21	31	54	106	68	-
2000	308	1	5	1	4	6	14	15	22	68	113	59	-
1999	346	2	-	1	7	8	14	19	34	57	116	88	-
Hispanic													
Both sexes													
2001	97	-	-	-	6	9	9	9	18	18	20	8	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Tuberculosis (A16-A19)													
Hispanic													
Both sexes													
2000	83	-	2	1	3	7	3	9	12	13	23	10	-
1999	125	1	-	1	4	3	13	19	20	24	27	13	-
Male													
2001	67	-	-	-	6	7	5	9	13	14	10	3	-
2000	46	-	-	-	1	4	2	5	7	8	14	5	-
1999	78	-	-	1	3	3	9	16	11	16	13	6	-
Female													
2001	30	-	-	-	-	2	4	-	5	4	10	5	-
2000	37	-	2	1	2	3	1	4	5	5	9	5	-
1999	47	1	-	-	1	-	4	3	9	8	14	7	-
Non-Hispanic [1]													
Both sexes													
2001	660	2	1	-	6	16	30	73	96	134	186	116	-
2000	683	1	4	-	8	13	39	80	68	140	226	104	-
1999	788	1	2	2	6	16	42	82	109	148	222	158	-
Male													
2001	379	-	-	-	2	7	21	52	70	84	90	53	-
2000	414	-	1	-	6	10	26	69	51	78	122	51	-
1999	492	-	2	1	-	8	32	66	84	99	122	78	-
Female													
2001	281	2	1	-	4	9	9	21	26	50	96	63	-
2000	269	1	3	-	2	3	13	11	17	62	104	53	-
1999	296	1	-	1	6	8	10	16	25	49	100	80	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Tuberculosis (A16-A19)													
Non-Hispanic white													
Both sexes													
2001	368	1	1	-	2	2	12	32	48	72	120	78	-
2000	353	-	2	-	3	4	10	34	22	72	131	75	-
1999	430	-	-	-	-	2	11	30	53	84	142	108	-
Male													
2001	200	-	-	-	-	2	10	20	37	39	59	33	-
2000	205	-	-	-	3	3	9	28	17	44	63	38	-
1999	263	-	-	-	-	1	9	27	41	57	77	51	-
Female													
2001	168	1	1	-	2	-	2	12	11	33	61	45	-
2000	148	-	2	-	-	1	1	6	5	28	68	37	-
1999	167	-	-	-	-	1	2	3	12	27	65	57	-
Non-Hispanic black													
Both sexes													
2001	200	-	-	-	1	9	16	36	37	44	35	22	-
2000	238	-	2	-	2	7	25	41	33	52	58	18	-
1999	239	1	2	1	4	9	24	41	39	42	50	26	-
Male													
2001	124	-	-	-	-	3	10	29	26	30	14	12	-
2000	156	-	1	-	1	5	14	36	26	28	38	7	-
1999	154	-	2	-	-	4	18	33	31	28	25	13	-
Female													
2001	76	-	-	-	1	6	6	7	11	14	21	10	-
2000	82	-	1	-	1	2	11	5	7	24	20	11	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Tuberculosis (A16-A19)													
Non-Hispanic black													
Female													
1999	85	1	-	1	4	5	6	8	8	14	25	13	-
Origin not stated [2]													
Both sexes													
2001	7	-	-	-	-	-	1	2	2	-	-	1	1
2000	10	-	-	-	-	-	-	2	1	2	3	1	1
1999	17	-	-	-	1	-	2	5	1	1	4	2	1
Male													
2001	7	-	-	-	-	-	1	2	2	-	-	1	1
2000	8	-	-	-	-	-	-	2	1	1	3	-	1
1999	14	-	-	-	1	-	2	5	1	1	2	1	1
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	1	-	1	-
1999	3	-	-	-	-	-	-	-	-	-	2	1	-
Respiratory tuberculosis (A16)													
All origins													
Both sexes													
2001	600	2	1	-	7	17	28	61	96	115	166	106	1
2000	615	-	3	-	7	11	35	72	60	119	209	98	1
1999	764	1	1	1	8	14	45	76	109	139	216	153	1
Male													
2001	371	-	-	-	5	10	18	49	78	76	84	50	1
2000	393	-	1	-	4	8	26	64	50	73	116	50	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Respiratory tuberculosis (A16)													
All origins													
Male													
1999	486	-	1	1	2	8	33	64	83	97	120	76	1
Female													
2001	229	2	1	-	2	7	10	12	18	39	82	56	-
2000	222	-	2	-	3	3	9	8	10	46	93	48	-
1999	278	1	-	-	6	6	12	12	26	42	96	77	-
Hispanic													
Both sexes													
2001	77	-	-	-	3	8	5	7	15	17	17	5	-
2000	60	-	1	-	2	3	2	5	9	12	18	8	-
1999	93	-	-	-	2	1	10	10	16	18	24	12	-
Male													
2001	56	-	-	-	3	6	3	7	12	13	10	2	-
2000	40	-	-	-	1	3	2	3	6	8	12	5	-
1999	55	-	-	-	1	1	6	9	9	13	11	5	-
Female													
2001	21	-	-	-	-	2	2	-	3	4	7	3	-
2000	20	-	1	-	1	-	-	2	3	4	6	3	-
1999	38	-	-	-	1	-	4	1	7	5	13	7	-
Non-Hispanic [1]													
Both sexes													
2001	517	2	1	-	4	9	22	53	79	98	149	100	-
2000	546	-	2	-	5	8	33	65	50	105	188	90	-
1999	657	1	1	1	5	13	33	63	92	120	189	139	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Respiratory tuberculosis (A16)													
Non-Hispanic [1]													
Male													
2001	309	-	-	-	2	4	14	41	64	63	74	47	-
2000	345	-	1	-	3	5	24	59	43	64	101	45	-
1999	419	-	1	1	-	7	25	52	73	83	107	70	-
Female													
2001	208	2	1	-	2	5	8	12	15	35	75	53	-
2000	201	-	1	-	2	3	9	6	7	41	87	45	-
1999	238	1	-	-	5	6	8	11	19	37	82	69	-
Non-Hispanic white													
Both sexes													
2001	288	1	1	-	1	-	8	23	41	52	97	64	-
2000	288	-	-	-	2	2	9	29	17	52	112	65	-
1999	371	-	-	-	-	2	9	27	45	69	119	100	-
Male													
2001	162	-	-	-	-	-	6	16	35	29	49	27	-
2000	174	-	-	-	2	1	9	26	14	34	54	34	-
1999	234	-	-	-	-	1	8	24	36	48	67	50	-
Female													
2001	126	1	1	-	1	-	2	7	6	23	48	37	-
2000	114	-	-	-	-	1	-	3	3	18	58	31	-
1999	137	-	-	-	-	1	1	3	9	21	52	50	-
Non-Hispanic black													
Both sexes													
2001	157	-	-	-	1	6	12	28	30	32	28	20	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Respiratory tuberculosis (A16)													
Non-Hispanic black													
Both sexes													
2000	187	-	2	-	1	6	20	31	22	43	46	16	-
1999	194	1	1	-	4	8	19	29	35	32	43	22	-
Male													
2001	103	-	-	-	-	2	7	24	23	23	12	12	-
2000	129	-	1	-	-	4	12	28	22	26	30	6	-
1999	122	-	1	-	-	3	13	24	28	21	22	10	-
Female													
2001	54	-	-	-	1	4	5	4	7	9	16	8	-
2000	58	-	1	-	1	2	8	3	-	17	16	10	-
1999	72	1	-	-	4	5	6	5	7	11	21	12	-
Origin not stated [2]													
Both sexes													
2001	6	-	-	-	-	-	1	1	2	-	-	1	1
2000	9	-	-	-	-	-	-	2	1	2	3	-	1
1999	14	-	-	-	1	-	2	3	1	1	3	2	1
Male													
2001	6	-	-	-	-	-	1	1	2	-	-	1	1
2000	8	-	-	-	-	-	-	2	1	1	3	-	1
1999	12	-	-	-	1	-	2	3	1	1	2	1	1
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	1	-	-	-
1999	2	-	-	-	-	-	-	-	-	-	1	1	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other tuberculosis (A17-A19)													
All origins													
Both sexes													
2001	164	-	-	-	5	8	12	23	20	37	40	19	-
2000	161	1	3	1	4	9	7	19	21	36	43	17	-
1999	166	1	1	2	3	5	12	30	21	34	37	20	-
Male													
2001	82	-	-	-	3	4	9	14	7	22	16	7	-
2000	75	-	-	-	3	6	2	12	9	14	23	6	-
1999	98	-	1	1	2	3	10	23	13	19	17	9	-
Female													
2001	82	-	-	-	2	4	3	9	13	15	24	12	-
2000	86	1	3	1	1	3	5	7	12	22	20	11	-
1999	68	1	-	1	1	2	2	7	8	15	20	11	-
Hispanic													
Both sexes													
2001	20	-	-	-	3	1	4	2	3	1	3	3	-
2000	23	-	1	1	1	4	1	4	3	1	5	2	-
1999	32	1	-	1	2	2	3	9	4	6	3	1	-
Male													
2001	11	-	-	-	3	1	2	2	1	1	-	1	-
2000	6	-	-	-	-	1	-	2	1	-	2	-	-
1999	23	-	-	1	2	2	3	7	2	3	2	1	-
Female													
2001	9	-	-	-	-	-	2	-	2	-	3	2	-
2000	17	-	1	1	1	3	1	2	2	1	3	2	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other tuberculosis (A17-A19)													
Hispanic													
Female													
1999	9	1	-	-	-	-	-	2	2	3	1	-	-
Non-Hispanic [1]													
Both sexes													
2001	143	-	-	-	2	7	8	20	17	36	37	16	-
2000	137	1	2	-	3	5	6	15	18	35	38	14	-
1999	131	-	1	1	1	3	9	19	17	28	33	19	-
Male													
2001	70	-	-	-	-	3	7	11	6	21	16	6	-
2000	69	-	-	-	3	5	2	10	8	14	21	6	-
1999	73	-	1	-	-	1	7	14	11	16	15	8	-
Female													
2001	73	-	-	-	2	4	1	9	11	15	21	10	-
2000	68	1	2	-	-	-	4	5	10	21	17	8	-
1999	58	-	-	1	1	2	2	5	6	12	18	11	-
Non-Hispanic white													
Both sexes													
2001	80	-	-	-	1	2	4	9	7	20	23	14	-
2000	65	-	2	-	1	2	1	5	5	20	19	10	-
1999	59	-	-	-	-	-	2	3	8	15	23	8	-
Male													
2001	38	-	-	-	-	2	4	4	2	10	10	6	-
2000	31	-	-	-	1	2	-	2	3	10	9	4	-
1999	29	-	-	-	-	-	1	3	5	9	10	1	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other tuberculosis (A17-A19)													
Non-Hispanic white													
Female													
2001	42	-	-	-	1	-	-	5	5	10	13	8	-
2000	34	-	2	-	-	-	1	3	2	10	10	6	-
1999	30	-	-	-	-	-	1	-	3	6	13	7	-
Non-Hispanic black													
Both sexes													
2001	43	-	-	-	-	3	4	8	7	12	7	2	-
2000	51	-	-	-	1	1	5	10	11	9	12	2	-
1999	45	-	1	1	-	1	5	12	4	10	7	4	-
Male													
2001	21	-	-	-	-	1	3	5	3	7	2	-	-
2000	27	-	-	-	1	1	2	8	4	2	8	1	-
1999	32	-	1	-	-	1	5	9	3	7	3	3	-
Female													
2001	22	-	-	-	-	2	1	3	4	5	5	2	-
2000	24	-	-	-	-	-	3	2	7	7	4	1	-
1999	13	-	-	1	-	-	-	3	1	3	4	1	-
Origin not stated [2]													
Both sexes													
2001	1	-	-	-	-	-	-	1	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	1	-
1999	3	-	-	-	-	-	-	2	-	-	1	-	-
Male													
2001	1	-	-	-	-	-	-	1	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other tuberculosis (A17-A19)													
Origin not stated [2]													
Male													
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	-	-	-	-	-	-	2	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	1	-
1999	1	-	-	-	-	-	-	-	-	-	1	-	-
Whooping cough (A37)													
All origins													
Both sexes													
2001	17	16	1	-	-	-	-	-	-	-	-	-	-
2000	12	12	-	-	-	-	-	-	-	-	-	-	-
1999	7	7	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	7	6	1	-	-	-	-	-	-	-	-	-	-
2000	5	5	-	-	-	-	-	-	-	-	-	-	-
1999	1	1	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	10	10	-	-	-	-	-	-	-	-	-	-	-
2000	7	7	-	-	-	-	-	-	-	-	-	-	-
1999	6	6	-	-	-	-	-	-	-	-	-	-	-
Hispanic													
Both sexes													
2001	10	10	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Whooping cough (A37)													
Hispanic													
Both sexes													
2000	7	7	-	-	-	-	-	-	-	-	-	-	-
1999	3	3	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	4	4	-	-	-	-	-	-	-	-	-	-	-
2000	2	2	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	6	6	-	-	-	-	-	-	-	-	-	-	-
2000	5	5	-	-	-	-	-	-	-	-	-	-	-
1999	3	3	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic [1]													
Both sexes													
2001	7	6	1	-	-	-	-	-	-	-	-	-	-
2000	5	5	-	-	-	-	-	-	-	-	-	-	-
1999	4	4	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	3	2	1	-	-	-	-	-	-	-	-	-	-
2000	3	3	-	-	-	-	-	-	-	-	-	-	-
1999	1	1	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	4	4	-	-	-	-	-	-	-	-	-	-	-
2000	2	2	-	-	-	-	-	-	-	-	-	-	-
1999	3	3	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Whooping cough (A37)													
Non-Hispanic white													
Both sexes													
2001	6	5	1	-	-	-	-	-	-	-	-	-	-
2000	4	4	-	-	-	-	-	-	-	-	-	-	-
1999	3	3	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	3	2	1	-	-	-	-	-	-	-	-	-	-
2000	2	2	-	-	-	-	-	-	-	-	-	-	-
1999	1	1	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	3	3	-	-	-	-	-	-	-	-	-	-	-
2000	2	2	-	-	-	-	-	-	-	-	-	-	-
1999	2	2	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic black													
Both sexes													
2001	1	1	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-
1999	1	1	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	1	1	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Whooping cough (A37)													
Non-Hispanic black													
Female													
1999	1	1	-	-	-	-	-	-	-	-	-	-	-
Origin not stated [2]													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Scarlet fever and erysipelas (A38,A46)													
All origins													
Both sexes													
2001	3	-	-	1	-	-	-	-	-	1	-	1	-
2000	2	-	-	-	-	-	-	1	-	-	1	-	-
1999	5	-	-	-	-	-	-	-	-	2	1	2	-
Male													
2001	2	-	-	1	-	-	-	-	-	-	-	1	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Scarlet fever and erysipelas (A38,A46)													
All origins													
Male													
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	1	-	-	-	-	-	-	-	-	1	-	-	-
2000	1	-	-	-	-	-	-	1	-	-	-	-	-
1999	5	-	-	-	-	-	-	-	-	2	1	2	-
Hispanic													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic [1]													
Both sexes													
2001	3	-	-	1	-	-	-	-	-	1	-	1	-
2000	2	-	-	-	-	-	-	1	-	-	1	-	-
1999	5	-	-	-	-	-	-	-	-	2	1	2	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Scarlet fever and erysipelas (A38,A46)													
Non-Hispanic [1]													
Male													
2001	2	-	-	1	-	-	-	-	-	-	-	1	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	1	-	-	-	-	-	-	-	-	1	-	-	-
2000	1	-	-	-	-	-	-	1	-	-	-	-	-
1999	5	-	-	-	-	-	-	-	-	2	1	2	-
Non-Hispanic white													
Both sexes													
2001	3	-	-	1	-	-	-	-	-	1	-	1	-
2000	2	-	-	-	-	-	-	1	-	-	1	-	-
1999	4	-	-	-	-	-	-	-	-	1	1	2	-
Male													
2001	2	-	-	1	-	-	-	-	-	-	-	1	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	1	-	-	-	-	-	-	-	-	1	-	-	-
2000	1	-	-	-	-	-	-	1	-	-	-	-	-
1999	4	-	-	-	-	-	-	-	-	1	1	2	-
Non-Hispanic black													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Scarlet fever and erysipelas (A38,A46)													
Non-Hispanic black													
Both sexes													
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	1	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	1	-	-	-
Origin not stated [2]													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Meningococcal infection (A39)													
All origins													
Both sexes													
2001	199	16	19	26	55	10	13	13	15	15	8	9	-
2000	211	22	19	18	56	15	13	29	9	8	15	7	-
1999	227	32	30	25	39	23	27	17	4	9	13	8	-
Male													
2001	94	6	9	15	27	6	6	8	7	5	3	2	-
2000	106	14	12	7	35	6	9	11	2	2	7	1	-
1999	101	14	19	13	19	12	9	4	-	5	4	2	-
Female													
2001	105	10	10	11	28	4	7	5	8	10	5	7	-
2000	105	8	7	11	21	9	4	18	7	6	8	6	-
1999	126	18	11	12	20	11	18	13	4	4	9	6	-
Hispanic													
Both sexes													
2001	21	4	5	2	2	2	2	1	2	-	1	-	-
2000	34	9	3	2	9	5	-	3	1	-	2	-	-
1999	27	7	1	3	2	6	5	2	-	-	1	-	-
Male													
2001	11	-	3	2	2	1	1	1	-	-	1	-	-
2000	16	4	2	-	6	2	-	1	-	-	1	-	-
1999	11	5	1	2	1	2	-	-	-	-	-	-	-
Female													
2001	10	4	2	-	-	1	1	-	2	-	-	-	-
2000	18	5	1	2	3	3	-	2	1	-	1	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Meningococcal infection (A39)													
Hispanic													
Female													
1999	16	2	-	1	1	4	5	2	-	-	1	-	-
Non-Hispanic [1]													
Both sexes													
2001	178	12	14	24	53	8	11	12	13	15	7	9	-
2000	177	13	16	16	47	10	13	26	8	8	13	7	-
1999	198	25	28	22	37	17	22	14	4	9	12	8	-
Male													
2001	83	6	6	13	25	5	5	7	7	5	2	2	-
2000	90	10	10	7	29	4	9	10	2	2	6	1	-
1999	89	9	17	11	18	10	9	4	-	5	4	2	-
Female													
2001	95	6	8	11	28	3	6	5	6	10	5	7	-
2000	87	3	6	9	18	6	4	16	6	6	7	6	-
1999	109	16	11	11	19	7	13	10	4	4	8	6	-
Non-Hispanic white													
Both sexes													
2001	138	10	10	18	38	5	9	11	9	14	6	8	-
2000	129	10	11	8	37	6	7	16	7	7	13	7	-
1999	155	23	24	15	27	8	14	12	4	9	11	8	-
Male													
2001	62	5	5	10	18	3	3	6	5	5	1	1	-
2000	60	7	6	4	22	1	5	4	2	2	6	1	-
1999	67	9	15	7	12	4	6	3	-	5	4	2	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Meningococcal infection (A39)													
Non-Hispanic white													
Female													
2001	76	5	5	8	20	2	6	5	4	9	5	7	-
2000	69	3	5	4	15	5	2	12	5	5	7	6	-
1999	88	14	9	8	15	4	8	9	4	4	7	6	-
Non-Hispanic black													
Both sexes													
2001	36	2	4	6	13	2	1	1	4	1	1	1	-
2000	44	3	5	8	7	4	6	9	1	1	-	-	-
1999	39	2	4	6	9	8	8	2	-	-	-	-	-
Male													
2001	18	1	1	3	6	1	1	1	2	-	1	1	-
2000	27	3	4	3	5	3	4	5	-	-	-	-	-
1999	20	-	2	3	6	5	3	1	-	-	-	-	-
Female													
2001	18	1	3	3	7	1	-	-	2	1	-	-	-
2000	17	-	1	5	2	1	2	4	1	1	-	-	-
1999	19	2	2	3	3	3	5	1	-	-	-	-	-
Origin not stated [2]													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	-	1	-	-	-	-	1	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Meningococcal infection (A39)													
Origin not stated [2]													
Male													
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	1	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	1	-	-	-	-	-
Septicemia (A40-A41)													
All origins													
Both sexes													
2001	32,238	312	108	69	133	281	829	1,977	3,111	5,998	10,351	9,069	-
2000	31,224	274	99	63	100	280	877	1,845	2,899	5,704	9,938	9,144	1
1999	30,680	283	87	77	121	275	812	1,685	2,714	5,750	9,710	9,166	-
Male													
2001	14,307	174	60	39	75	139	451	1,043	1,606	3,044	4,762	2,914	-
2000	13,537	139	53	38	60	158	451	1,039	1,513	2,867	4,482	2,736	1
1999	13,395	175	48	36	67	147	460	919	1,412	2,919	4,426	2,786	-
Female													
2001	17,931	138	48	30	58	142	378	934	1,505	2,954	5,589	6,155	-
2000	17,687	135	46	25	40	122	426	806	1,386	2,837	5,456	6,408	-
1999	17,285	108	39	41	54	128	352	766	1,302	2,831	5,284	6,380	-
Hispanic													
Both sexes													
2001	1,386	50	18	18	13	23	72	133	151	288	346	274	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Septicemia (A40-A41)													
Hispanic													
Both sexes													
2000	1,226	62	15	7	12	31	79	122	145	235	300	218	-
1999	1,215	43	14	8	9	26	74	92	131	265	296	257	-
Male													
2001	684	29	5	14	9	15	43	80	86	149	170	84	-
2000	625	33	9	5	6	21	46	78	80	126	143	78	-
1999	605	27	8	4	5	14	51	56	78	134	144	84	-
Female													
2001	702	21	13	4	4	8	29	53	65	139	176	190	-
2000	601	29	6	2	6	10	33	44	65	109	157	140	-
1999	610	16	6	4	4	12	23	36	53	131	152	173	-
Non-Hispanic [1]													
Both sexes													
2001	30,754	261	90	50	120	257	755	1,837	2,941	5,690	9,977	8,776	-
2000	29,864	208	83	56	88	248	793	1,713	2,739	5,442	9,601	8,892	1
1999	29,351	239	73	69	112	248	736	1,585	2,570	5,466	9,378	8,875	-
Male													
2001	13,573	145	55	25	66	123	408	957	1,510	2,882	4,575	2,827	-
2000	12,842	104	44	33	54	137	401	957	1,424	2,724	4,316	2,647	1
1999	12,737	147	40	32	62	132	408	858	1,330	2,777	4,261	2,690	-
Female													
2001	17,181	116	35	25	54	134	347	880	1,431	2,808	5,402	5,949	-
2000	17,022	104	39	23	34	111	392	756	1,315	2,718	5,285	6,245	-
1999	16,614	92	33	37	50	116	328	727	1,240	2,689	5,117	6,185	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Septicemia (A40-A41)													
Non-Hispanic white													
Both sexes													
2001	24,389	122	63	35	69	145	458	1,181	2,081	4,385	8,352	7,498	-
2000	23,517	108	51	42	48	134	498	1,092	1,919	4,173	7,901	7,550	1
1999	23,023	121	47	47	71	155	439	1,015	1,792	4,140	7,717	7,479	-
Male													
2001	10,668	71	41	18	37	69	266	615	1,071	2,223	3,851	2,406	-
2000	10,056	55	26	25	33	74	262	613	1,015	2,131	3,579	2,242	1
1999	9,863	79	25	24	44	87	243	551	904	2,111	3,538	2,257	-
Female													
2001	13,721	51	22	17	32	76	192	566	1,010	2,162	4,501	5,092	-
2000	13,461	53	25	17	15	60	236	479	904	2,042	4,322	5,308	-
1999	13,160	42	22	23	27	68	196	464	888	2,029	4,179	5,222	-
Non-Hispanic black													
Both sexes													
2001	5,829	124	21	10	41	98	268	595	804	1,186	1,501	1,181	-
2000	5,863	96	25	11	33	107	269	572	756	1,173	1,576	1,245	-
1999	5,772	106	16	14	32	85	279	518	716	1,212	1,513	1,281	-
Male													
2001	2,611	67	12	3	25	46	128	304	411	586	656	373	-
2000	2,559	48	13	6	18	59	125	319	381	545	677	368	-
1999	2,594	60	10	7	14	41	154	279	392	609	648	380	-
Female													
2001	3,218	57	9	7	16	52	140	291	393	600	845	808	-
2000	3,304	48	12	5	15	48	144	253	375	628	899	877	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Septicemia (A40-A41)													
Non-Hispanic black													
Female													
1999	3,178	46	6	7	18	44	125	239	324	603	865	901	-
Origin not stated [2]													
Both sexes													
2001	98	1	-	1	-	1	2	7	19	20	28	19	-
2000	134	4	1	-	-	1	5	10	15	27	37	34	-
1999	114	1	-	-	-	1	2	8	13	19	36	34	-
Male													
2001	50	-	-	-	-	1	-	6	10	13	17	3	-
2000	70	2	-	-	-	-	4	4	9	17	23	11	-
1999	53	1	-	-	-	1	1	5	4	8	21	12	-
Female													
2001	48	1	-	1	-	-	2	1	9	7	11	16	-
2000	64	2	1	-	-	1	1	6	6	10	14	23	-
1999	61	-	-	-	-	-	1	3	9	11	15	22	-
Syphilis (A50-A53)													
All origins													
Both sexes													
2001	36	1	-	1	-	1	3	3	4	5	13	5	-
2000	41	1	-	-	-	1	-	6	3	7	13	10	-
1999	33	2	-	-	-	1	1	5	4	6	10	4	-
Male													
2001	19	1	-	1	-	-	2	3	3	3	5	1	-
2000	24	1	-	-	-	1	-	5	3	2	7	5	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Syphilis (A50-A53)													
All origins													
Male													
1999	13	-	-	-	-	1	-	3	3	5	-	1	-
Female													
2001	17	-	-	-	-	1	1	-	1	2	8	4	-
2000	17	-	-	-	-	-	-	1	-	5	6	5	-
1999	20	2	-	-	-	-	1	2	1	1	10	3	-
Hispanic													
Both sexes													
2001	7	1	-	-	-	-	-	-	2	-	3	1	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	5	2	-	-	-	-	-	2	-	-	1	-	-
Male													
2001	5	1	-	-	-	-	-	-	2	-	1	1	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	1	-	-	-	-	-
Female													
2001	2	-	-	-	-	-	-	-	-	-	2	-	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	4	2	-	-	-	-	-	1	-	-	1	-	-
Non-Hispanic [1]													
Both sexes													
2001	28	-	-	1	-	1	3	3	1	5	10	4	-
2000	39	1	-	-	-	1	-	6	3	7	11	10	-
1999	27	-	-	-	-	1	1	2	4	6	9	4	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Syphilis (A50-A53)													
Non-Hispanic [1]													
Male													
2001	14	-	-	1	-	-	2	3	1	3	4	-	-
2000	24	1	-	-	-	1	-	5	3	2	7	5	-
1999	11	-	-	-	-	1	-	1	3	5	-	1	-
Female													
2001	14	-	-	-	-	1	1	-	-	2	6	4	-
2000	15	-	-	-	-	-	-	1	-	5	4	5	-
1999	16	-	-	-	-	-	1	1	1	1	9	3	-
Non-Hispanic white													
Both sexes													
2001	14	-	-	-	-	-	1	1	1	2	7	2	-
2000	20	-	-	-	-	1	-	2	-	3	6	8	-
1999	14	-	-	-	-	1	1	2	3	3	3	1	-
Male													
2001	5	-	-	-	-	-	-	1	1	1	2	-	-
2000	7	-	-	-	-	1	-	1	-	-	2	3	-
1999	6	-	-	-	-	1	-	1	2	2	-	-	-
Female													
2001	9	-	-	-	-	-	1	-	-	1	5	2	-
2000	13	-	-	-	-	-	-	1	-	3	4	5	-
1999	8	-	-	-	-	-	1	1	1	1	3	1	-
Non-Hispanic black													
Both sexes													
2001	14	-	-	1	-	1	2	2	-	3	3	2	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Syphilis (A50-A53)													
Non-Hispanic black													
Both sexes													
2000	19	1	-	-	-	-	-	4	3	4	5	2	-
1999	12	-	-	-	-	-	-	-	1	3	5	3	-
Male													
2001	9	-	-	1	-	-	2	2	-	2	2	-	-
2000	17	1	-	-	-	-	-	4	3	2	5	2	-
1999	5	-	-	-	-	-	-	-	1	3	-	1	-
Female													
2001	5	-	-	-	-	1	-	-	-	1	1	2	-
2000	2	-	-	-	-	-	-	-	-	2	-	-	-
1999	7	-	-	-	-	-	-	-	-	-	5	2	-
Origin not stated [2]													
Both sexes													
2001	1	-	-	-	-	-	-	-	1	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	1	-	-	-	-	-	-	1	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	1	-	-	-	-	-
Female													
2001	1	-	-	-	-	-	-	-	1	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Acute poliomyelitis (A80)													
All origins													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Hispanic													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Acute poliomyelitis (A80)													
Hispanic													
Female													
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic [1]													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic white													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Acute poliomyelitis (A80)													
Non-Hispanic white													
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic black													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated [2]													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Acute poliomyelitis (A80)													
Origin not stated [2]													
Male													
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Arthropod-borne viral encephalitis (A83-A84,A85.2)													
All origins													
Both sexes													
2001	9	1	-	-	-	-	-	1	-	5	2	-	-
2000	2	-	-	-	-	-	1	-	-	-	1	-	-
1999	3	-	-	-	-	-	-	-	-	2	-	1	-
Male													
2001	6	-	-	-	-	-	-	-	-	4	2	-	-
2000	2	-	-	-	-	-	1	-	-	-	1	-	-
1999	3	-	-	-	-	-	-	-	-	2	-	1	-
Female													
2001	3	1	-	-	-	-	-	1	-	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Hispanic													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Arthropod-borne viral encephalitis (A83-A84,A85.2)													
Hispanic													
Both sexes													
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic [1]													
Both sexes													
2001	9	1	-	-	-	-	-	1	-	5	2	-	-
2000	2	-	-	-	-	-	1	-	-	-	1	-	-
1999	3	-	-	-	-	-	-	-	-	2	-	1	-
Male													
2001	6	-	-	-	-	-	-	-	-	4	2	-	-
2000	2	-	-	-	-	-	1	-	-	-	1	-	-
1999	3	-	-	-	-	-	-	-	-	2	-	1	-
Female													
2001	3	1	-	-	-	-	-	1	-	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Arthropod-borne viral encephalitis (A83-A84,A85.2)													
Non-Hispanic white													
Both sexes													
2001	6	-	-	-	-	-	-	1	-	4	1	-	-
2000	2	-	-	-	-	-	1	-	-	-	1	-	-
1999	3	-	-	-	-	-	-	-	-	2	-	1	-
Male													
2001	5	-	-	-	-	-	-	-	-	4	1	-	-
2000	2	-	-	-	-	-	1	-	-	-	1	-	-
1999	3	-	-	-	-	-	-	-	-	2	-	1	-
Female													
2001	1	-	-	-	-	-	-	1	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic black													
Both sexes													
2001	2	1	-	-	-	-	-	-	-	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	2	1	-	-	-	-	-	-	-	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Arthropod-borne viral encephalitis (A83-A84,A85.2)													
Non-Hispanic black													
Female													
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated [2]													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Measles (B05)													
All origins													
Both sexes													
2001	1	-	-	-	-	-	1	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-
1999	2	1	-	-	-	-	-	-	1	-	-	-	-
Male													
2001	1	-	-	-	-	-	1	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Measles (B05)													
All origins													
Male													
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-
1999	2	1	-	-	-	-	-	-	1	-	-	-	-
Hispanic													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic [1]													
Both sexes													
2001	1	-	-	-	-	-	1	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	1	-	-	-	-	-	-	1	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Measles (B05)													
Non-Hispanic [1]													
Male													
2001	1	-	-	-	-	-	1	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	1	-	-	-	-	-	-	1	-	-	-	-
Non-Hispanic white													
Both sexes													
2001	1	-	-	-	-	-	1	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	1	-	-	-	-
Male													
2001	1	-	-	-	-	-	1	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	1	-	-	-	-
Non-Hispanic black													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Measles (B05)													
Non-Hispanic black													
Both sexes													
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	1	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	1	-	-	-	-	-	-	-	-	-	-	-
Origin not stated [2]													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Viral hepatitis (B15-B19)													
All origins													
Both sexes													
2001	5,585	1	-	3	20	74	797	2,060	1,014	836	613	166	1
2000	5,357	6	1	2	18	79	779	1,894	904	845	660	168	1
1999	4,853	2	3	3	14	88	722	1,506	786	881	651	197	-
Male													
2001	3,539	1	-	2	11	42	553	1,570	643	396	263	57	1
2000	3,381	2	1	1	11	46	540	1,425	577	396	311	70	1
1999	3,043	2	2	2	10	53	520	1,138	491	426	317	82	-
Female													
2001	2,046	-	-	1	9	32	244	490	371	440	350	109	-
2000	1,976	4	-	1	7	33	239	469	327	449	349	98	-
1999	1,810	-	1	1	4	35	202	368	295	455	334	115	-
Hispanic													
Both sexes													
2001	791	-	-	2	6	10	126	266	192	113	61	15	-
2000	708	3	-	-	3	14	126	218	147	120	66	11	-
1999	656	-	1	-	3	16	104	203	127	129	65	8	-
Male													
2001	487	-	-	2	2	3	89	206	123	35	23	4	-
2000	456	1	-	-	2	11	100	165	94	52	25	6	-
1999	406	-	1	-	1	11	84	153	75	59	18	4	-
Female													
2001	304	-	-	-	4	7	37	60	69	78	38	11	-
2000	252	2	-	-	1	3	26	53	53	68	41	5	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Viral hepatitis (B15-B19)													
Hispanic													
Female													
1999	250	-	-	-	2	5	20	50	52	70	47	4	-
Non-Hispanic [1]													
Both sexes													
2001	4,776	1	-	1	14	64	669	1,785	817	721	552	151	1
2000	4,629	3	1	2	15	65	652	1,665	751	724	594	156	1
1999	4,176	2	2	3	11	71	613	1,294	656	750	585	189	-
Male													
2001	3,038	1	-	-	9	39	462	1,357	515	361	240	53	1
2000	2,909	1	1	1	9	35	440	1,250	479	343	286	63	1
1999	2,620	2	1	2	9	41	432	978	414	365	298	78	-
Female													
2001	1,738	-	-	1	5	25	207	428	302	360	312	98	-
2000	1,720	2	-	1	6	30	212	415	272	381	308	93	-
1999	1,556	-	1	1	2	30	181	316	242	385	287	111	-
Non-Hispanic white													
Both sexes													
2001	3,663	1	-	1	7	43	522	1,364	586	544	467	128	-
2000	3,513	1	-	2	5	37	501	1,243	520	559	506	138	1
1999	3,176	1	2	2	6	51	464	958	465	567	497	163	-
Male													
2001	2,410	1	-	-	4	28	368	1,062	384	296	218	49	-
2000	2,234	-	-	1	4	19	342	958	339	265	248	57	1
1999	2,026	1	1	1	6	29	333	717	308	295	268	67	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Viral hepatitis (B15-B19)													
Non-Hispanic white													
Female													
2001	1,253	-	-	1	3	15	154	302	202	248	249	79	-
2000	1,279	1	-	1	1	18	159	285	181	294	258	81	-
1999	1,150	-	1	1	-	22	131	241	157	272	229	96	-
Non-Hispanic black													
Both sexes													
2001	819	-	-	-	6	15	121	349	167	91	53	16	1
2000	777	2	-	-	10	21	117	329	138	93	57	10	-
1999	718	1	-	1	4	17	119	277	118	109	53	19	-
Male													
2001	479	-	-	-	4	7	79	240	99	34	13	2	1
2000	490	1	-	-	5	11	77	234	86	48	25	3	-
1999	433	1	-	1	2	9	79	215	65	38	15	8	-
Female													
2001	340	-	-	-	2	8	42	109	68	57	40	14	-
2000	287	1	-	-	5	10	40	95	52	45	32	7	-
1999	285	-	-	-	2	8	40	62	53	71	38	11	-
Origin not stated [2]													
Both sexes													
2001	18	-	-	-	-	-	2	9	5	2	-	-	-
2000	20	-	-	-	-	-	1	11	6	1	-	1	-
1999	21	-	-	-	-	1	5	9	3	2	1	-	-
Male													
2001	14	-	-	-	-	-	2	7	5	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Viral hepatitis (B15-B19)													
Origin not stated [2]													
Male													
2000	16	-	-	-	-	-	-	10	4	1	-	1	-
1999	17	-	-	-	-	1	4	7	2	2	1	-	-
Female													
2001	4	-	-	-	-	-	-	2	-	2	-	-	-
2000	4	-	-	-	-	-	1	1	2	-	-	-	-
1999	4	-	-	-	-	-	1	2	1	-	-	-	-
Human immunodeficiency virus (HIV) disease (B20-B24)													
All origins													
Both sexes													
2001	14,175	10	13	38	225	2,101	5,867	4,120	1,317	384	88	10	2
2000	14,478	10	14	46	179	2,437	5,919	4,142	1,239	397	85	9	1
1999	14,802	14	24	66	198	2,729	6,232	3,907	1,149	398	70	14	1
Male													
2001	10,590	7	8	23	105	1,421	4,360	3,241	1,051	292	73	7	2
2000	10,868	8	6	22	99	1,602	4,441	3,296	1,013	313	65	2	1
1999	11,234	9	16	36	95	1,914	4,706	3,134	948	317	50	8	1
Female													
2001	3,585	3	5	15	120	680	1,507	879	266	92	15	3	-
2000	3,610	2	8	24	80	835	1,478	846	226	84	20	7	-
1999	3,568	5	8	30	103	815	1,526	773	201	81	20	6	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Human immunodeficiency virus (HIV) disease (B20-B24)													
Hispanic													
Both sexes													
2001	1,853	1	-	3	32	300	759	526	169	51	10	2	-
2000	1,908	-	-	9	23	365	762	531	163	41	14	-	-
1999	1,958	3	5	9	33	407	855	455	137	49	4	1	-
Male													
2001	1,437	-	-	2	21	222	589	420	129	43	9	2	-
2000	1,493	-	-	5	18	274	603	426	123	34	10	-	-
1999	1,525	1	4	3	20	314	666	354	118	41	4	-	-
Female													
2001	416	1	-	1	11	78	170	106	40	8	1	-	-
2000	415	-	-	4	5	91	159	105	40	7	4	-	-
1999	433	2	1	6	13	93	189	101	19	8	-	1	-
Non-Hispanic [1]													
Both sexes													
2001	12,141	9	13	35	193	1,779	5,044	3,533	1,123	325	77	8	2
2000	12,389	10	14	37	156	2,053	5,077	3,562	1,058	343	70	9	-
1999	12,638	11	19	56	165	2,301	5,305	3,377	987	339	65	13	-
Male													
2001	9,011	7	8	21	84	1,187	3,725	2,766	902	241	63	5	2
2000	9,224	8	6	17	81	1,313	3,778	2,825	872	267	55	2	-
1999	9,538	8	12	33	75	1,584	3,984	2,715	806	268	45	8	-
Female													
2001	3,130	2	5	14	109	592	1,319	767	221	84	14	3	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Human immunodeficiency virus (HIV) disease (B20-B24)													
Non-Hispanic [1]													
Female													
2000	3,165	2	8	20	75	740	1,299	737	186	76	15	7	-
1999	3,100	3	7	23	90	717	1,321	662	181	71	20	5	-
Non-Hispanic white													
Both sexes													
2001	4,286	2	3	8	30	519	1,911	1,248	423	110	27	5	-
2000	4,562	1	1	13	30	619	2,000	1,325	420	122	28	3	-
1999	4,744	2	2	15	34	794	2,082	1,286	369	121	32	7	-
Male													
2001	3,669	2	-	5	15	410	1,639	1,106	370	95	24	3	-
2000	3,837	1	-	5	20	475	1,678	1,161	369	103	25	-	-
1999	4,043	2	1	10	22	643	1,768	1,136	331	104	24	2	-
Female													
2001	617	-	3	3	15	109	272	142	53	15	3	2	-
2000	725	-	1	8	10	144	322	164	51	19	3	3	-
1999	701	-	1	5	12	151	314	150	38	17	8	5	-
Non-Hispanic black													
Both sexes													
2001	7,710	7	10	26	161	1,234	3,072	2,249	690	208	49	2	2
2000	7,708	9	12	23	125	1,414	3,028	2,211	625	214	41	6	-
1999	7,751	9	17	41	130	1,479	3,154	2,057	613	213	33	5	-
Male													
2001	5,227	5	8	15	68	753	2,040	1,629	524	144	38	1	2

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Human immunodeficiency virus (HIV) disease (B20-B24)													
Non-Hispanic black													
Male													
2000	5,291	7	6	12	60	822	2,060	1,642	492	159	29	2	-
1999	5,377	6	11	23	52	919	2,157	1,552	471	160	21	5	-
Female													
2001	2,483	2	2	11	93	481	1,032	620	166	64	11	1	-
2000	2,417	2	6	11	65	592	968	569	133	55	12	4	-
1999	2,374	3	6	18	78	560	997	505	142	53	12	-	-
Origin not stated [2]													
Both sexes													
2001	181	-	-	-	-	22	64	61	25	8	1	-	-
2000	181	-	-	-	-	19	80	49	18	13	1	-	1
1999	206	-	-	1	-	21	72	75	25	10	1	-	1
Male													
2001	142	-	-	-	-	12	46	55	20	8	1	-	-
2000	151	-	-	-	-	15	60	45	18	12	-	-	1
1999	171	-	-	-	-	16	56	65	24	8	1	-	1
Female													
2001	39	-	-	-	-	10	18	6	5	-	-	-	-
2000	30	-	-	-	-	4	20	4	-	1	1	-	-
1999	35	-	-	1	-	5	16	10	1	2	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malaria (B50-B54)													
All origins													
Both sexes													
2001	9	-	-	1	-	-	2	2	2	1	1	-	-
2000	3	-	-	-	1	-	-	1	-	-	1	-	-
1999	7	1	-	-	-	2	1	-	1	2	-	-	-
Male													
2001	5	-	-	1	-	-	-	1	1	1	1	-	-
2000	2	-	-	-	-	-	-	1	-	-	1	-	-
1999	4	1	-	-	-	1	-	-	1	1	-	-	-
Female													
2001	4	-	-	-	-	-	2	1	1	-	-	-	-
2000	1	-	-	-	1	-	-	-	-	-	-	-	-
1999	3	-	-	-	-	1	1	-	-	1	-	-	-
Hispanic													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malaria (B50-B54)													
Hispanic													
Female													
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic [1]													
Both sexes													
2001	9	-	-	1	-	-	2	2	2	1	1	-	-
2000	3	-	-	-	1	-	-	1	-	-	1	-	-
1999	7	1	-	-	-	2	1	-	1	2	-	-	-
Male													
2001	5	-	-	1	-	-	-	1	1	1	1	-	-
2000	2	-	-	-	-	-	-	1	-	-	1	-	-
1999	4	1	-	-	-	1	-	-	1	1	-	-	-
Female													
2001	4	-	-	-	-	-	2	1	1	-	-	-	-
2000	1	-	-	-	1	-	-	-	-	-	-	-	-
1999	3	-	-	-	-	1	1	-	-	1	-	-	-
Non-Hispanic white													
Both sexes													
2001	5	-	-	-	-	-	2	1	-	1	1	-	-
2000	1	-	-	-	1	-	-	-	-	-	-	-	-
1999	4	-	-	-	-	1	1	-	1	1	-	-	-
Male													
2001	2	-	-	-	-	-	-	-	-	1	1	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	-	-	-	1	-	-	1	1	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malaria (B50-B54)													
Non-Hispanic white													
Female													
2001	3	-	-	-	-	-	2	1	-	-	-	-	-
2000	1	-	-	-	1	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	1	-	-	-	-	-	-
Non-Hispanic black													
Both sexes													
2001	3	-	-	1	-	-	-	1	1	-	-	-	-
2000	1	-	-	-	-	-	-	1	-	-	-	-	-
1999	3	1	-	-	-	1	-	-	-	1	-	-	-
Male													
2001	2	-	-	1	-	-	-	1	-	-	-	-	-
2000	1	-	-	-	-	-	-	1	-	-	-	-	-
1999	1	1	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	1	-	-	-	-	-	-	-	1	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	-	-	-	-	1	-	-	-	1	-	-	-
Origin not stated [2]													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malaria (B50-B54)													
Origin not stated [2]													
Male													
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44, A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8, A86-B04,B06-B09,B25-B49,B55-B99)													
All origins													
Both sexes													
2001	5,715	191	64	64	86	147	369	663	771	1,143	1,330	887	-
2000	5,532	167	71	67	96	141	350	585	707	1,074	1,375	899	-
1999	7,279	203	81	79	90	242	545	836	898	1,443	1,739	1,123	-
Male													
2001	2,931	101	37	29	51	83	227	426	449	565	656	307	-
2000	2,861	94	41	35	54	75	220	358	383	597	665	339	-
1999	3,655	119	51	44	49	131	333	478	482	760	827	381	-
Female													
2001	2,784	90	27	35	35	64	142	237	322	578	674	580	-
2000	2,671	73	30	32	42	66	130	227	324	477	710	560	-
1999	3,624	84	30	35	41	111	212	358	416	683	912	742	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44, A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8, A86-B04,B06-B09,B25-B49,B55-B99)													
Hispanic													
Both sexes													
2001	423	41	7	11	12	33	46	69	56	67	57	24	-
2000	412	23	18	13	19	26	50	61	54	62	57	29	-
1999	497	36	13	14	13	39	67	68	62	80	68	37	-
Male													
2001	250	21	4	5	9	22	31	47	38	38	27	8	-
2000	237	14	12	8	12	20	37	33	35	28	27	11	-
1999	260	22	10	7	6	22	49	40	29	36	29	10	-
Female													
2001	173	20	3	6	3	11	15	22	18	29	30	16	-
2000	175	9	6	5	7	6	13	28	19	34	30	18	-
1999	237	14	3	7	7	17	18	28	33	44	39	27	-
Non-Hispanic [1]													
Both sexes													
2001	5,271	149	57	53	73	114	319	589	711	1,073	1,272	861	-
2000	5,100	144	53	54	75	115	297	521	649	1,012	1,312	868	-
1999	6,753	166	68	63	77	202	475	763	831	1,358	1,666	1,084	-
Male													
2001	2,666	79	33	24	42	61	193	374	408	525	628	299	-
2000	2,613	80	29	27	42	55	182	322	347	569	633	327	-
1999	3,379	96	41	36	43	108	282	435	449	722	797	370	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44, A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8, A86-B04,B06-B09,B25-B49,B55-B99)													
Non-Hispanic [1]													
Female													
2001	2,605	70	24	29	31	53	126	215	303	548	644	562	-
2000	2,487	64	24	27	33	60	115	199	302	443	679	541	-
1999	3,374	70	27	27	34	94	193	328	382	636	869	714	-
Non-Hispanic white													
Both sexes													
2001	4,257	80	37	38	45	63	211	435	551	899	1,123	775	-
2000	4,131	79	33	40	51	58	189	383	505	851	1,165	777	-
1999	5,288	101	43	42	51	104	287	509	628	1,126	1,438	959	-
Male													
2001	2,148	40	21	17	24	39	129	281	320	443	562	272	-
2000	2,092	45	16	18	29	28	115	239	275	483	557	287	-
1999	2,644	64	27	24	28	55	190	303	345	608	681	319	-
Female													
2001	2,109	40	16	21	21	24	82	154	231	456	561	503	-
2000	2,039	34	17	22	22	30	74	144	230	368	608	490	-
1999	2,644	37	16	18	23	49	97	206	283	518	757	640	-
Non-Hispanic black													
Both sexes													
2001	815	61	16	13	23	45	91	130	129	135	105	67	-
2000	815	60	14	10	21	50	95	124	116	129	120	76	-
1999	1,247	57	23	17	20	89	169	219	169	192	186	106	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44, A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8, A86-B04,B06-B09,B25-B49,B55-B99)													
Non-Hispanic black													
Male													
2001	419	36	10	6	15	19	56	80	70	66	42	19	-
2000	428	31	10	5	10	22	58	75	62	70	58	27	-
1999	617	29	14	10	12	48	81	114	82	94	94	39	-
Female													
2001	396	25	6	7	8	26	35	50	59	69	63	48	-
2000	387	29	4	5	11	28	37	49	54	59	62	49	-
1999	630	28	9	7	8	41	88	105	87	98	92	67	-
Origin not stated [2]													
Both sexes													
2001	21	1	-	-	1	-	4	5	4	3	1	2	-
2000	20	-	-	-	2	-	3	3	4	-	6	2	-
1999	29	1	-	2	-	1	3	5	5	5	5	2	-
Male													
2001	15	1	-	-	-	-	3	5	3	2	1	-	-
2000	11	-	-	-	-	-	1	3	1	-	5	1	-
1999	16	1	-	1	-	1	2	3	4	2	1	1	-
Female													
2001	6	-	-	-	1	-	1	-	1	1	-	2	-
2000	9	-	-	-	2	-	2	-	3	-	1	1	-
1999	13	-	-	1	-	-	1	2	1	3	4	1	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms (C00-C97)													
All origins													
Both sexes													
2001	553,768	66	420	1,008	1,704	3,994	16,569	49,562	90,223	147,018	165,445	77,751	8
2000	553,091	92	420	1,014	1,713	3,916	16,520	48,034	89,005	150,131	165,099	77,136	11
1999	549,838	68	418	1,012	1,724	4,005	16,732	46,681	89,067	152,338	162,770	75,014	9
Male													
2001	287,075	31	229	533	1,024	1,860	7,297	25,071	49,244	81,699	86,126	33,954	7
2000	286,082	50	233	569	1,022	1,854	7,344	24,215	48,427	83,191	85,907	33,260	10
1999	285,832	36	227	555	1,023	1,959	7,471	23,634	48,833	84,544	85,060	32,484	6
Female													
2001	266,693	35	191	475	680	2,134	9,272	24,491	40,979	65,319	79,319	43,797	1
2000	267,009	42	187	445	691	2,062	9,176	23,819	40,578	66,940	79,192	43,876	1
1999	264,006	32	191	457	701	2,046	9,261	23,047	40,234	67,794	77,710	42,530	3
Hispanic													
Both sexes													
2001	22,371	14	84	177	289	497	1,396	2,815	4,126	5,724	5,068	2,181	-
2000	21,160	17	75	198	275	476	1,295	2,575	3,780	5,557	4,827	2,085	-
1999	20,233	16	72	176	259	466	1,204	2,516	3,686	5,394	4,449	1,995	-
Male													
2001	11,825	4	44	90	199	233	615	1,372	2,202	3,219	2,814	1,033	-
2000	11,138	10	43	122	162	241	534	1,232	2,034	3,086	2,657	1,017	-
1999	10,668	11	34	92	158	222	504	1,237	2,060	3,045	2,403	902	-
Female													
2001	10,546	10	40	87	90	264	781	1,443	1,924	2,505	2,254	1,148	-
2000	10,022	7	32	76	113	235	761	1,343	1,746	2,471	2,170	1,068	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms (C00-C97)													
Hispanic													
Female													
1999	9,565	5	38	84	101	244	700	1,279	1,626	2,349	2,046	1,093	-
Non-Hispanic [1]													
Both sexes													
2001	530,058	52	333	828	1,413	3,485	15,121	46,617	85,842	140,943	160,023	75,394	7
2000	530,531	75	341	812	1,432	3,431	15,170	45,305	84,977	144,200	159,906	74,871	11
1999	528,218	52	343	835	1,462	3,525	15,482	44,036	85,146	146,572	157,904	72,852	9
Male													
2001	274,492	27	183	442	824	1,623	6,652	23,631	46,871	78,266	83,121	32,846	6
2000	274,185	40	186	446	858	1,607	6,787	22,902	46,253	79,878	83,047	32,171	10
1999	274,419	25	190	463	864	1,732	6,946	22,323	46,644	81,271	82,443	31,512	6
Female													
2001	255,566	25	150	386	589	1,862	8,469	22,986	38,971	62,677	76,902	42,548	1
2000	256,346	35	155	366	574	1,824	8,383	22,403	38,724	64,322	76,859	42,700	1
1999	253,799	27	153	372	598	1,793	8,536	21,713	38,502	65,301	75,461	41,340	3
Non-Hispanic white													
Both sexes													
2001	456,709	42	268	607	1,076	2,594	11,551	36,836	71,389	121,529	142,760	68,053	4
2000	458,200	60	256	627	1,059	2,599	11,458	35,861	70,865	124,842	143,079	67,487	7
1999	456,315	37	281	608	1,070	2,682	11,848	34,944	70,788	127,119	141,295	65,636	7
Male													
2001	236,004	21	144	310	632	1,260	5,160	18,679	38,884	67,554	73,885	29,471	4
2000	235,932	34	146	338	636	1,259	5,201	18,092	38,368	69,085	73,952	28,814	7
1999	236,384	20	160	345	629	1,341	5,425	17,588	38,658	70,518	73,535	28,161	4

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms (C00-C97)													
Non-Hispanic white													
Female													
2001	220,705	21	124	297	444	1,334	6,391	18,157	32,505	53,975	68,875	38,582	-
2000	222,268	26	110	289	423	1,340	6,257	17,769	32,497	55,757	69,127	38,673	-
1999	219,931	17	121	263	441	1,341	6,423	17,356	32,130	56,601	67,760	37,475	3
Non-Hispanic black													
Both sexes													
2001	61,580	5	46	172	257	691	2,950	8,175	12,196	16,392	14,454	6,239	3
2000	61,390	11	66	147	297	641	3,072	8,068	12,019	16,485	14,244	6,337	3
1999	61,409	11	47	179	314	660	3,049	7,749	12,322	16,625	14,192	6,260	1
Male													
2001	32,335	2	29	100	148	267	1,201	4,196	6,752	9,134	7,700	2,804	2
2000	32,493	4	31	86	178	259	1,292	4,152	6,763	9,235	7,689	2,802	2
1999	32,542	3	24	94	181	294	1,255	4,098	6,933	9,228	7,603	2,828	1
Female													
2001	29,245	3	17	72	109	424	1,749	3,979	5,444	7,258	6,754	3,435	1
2000	28,897	7	35	61	119	382	1,780	3,916	5,256	7,250	6,555	3,535	1
1999	28,867	8	23	85	133	366	1,794	3,651	5,389	7,397	6,589	3,432	-
Origin not stated [2]													
Both sexes													
2001	1,339	-	3	3	2	12	52	130	255	351	354	176	1
2000	1,400	-	4	4	6	9	55	154	248	374	366	180	-
1999	1,387	-	3	1	3	14	46	129	235	372	417	167	-
Male													
2001	758	-	2	1	1	4	30	68	171	214	191	75	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms (C00-C97)													
Origin not stated [2]													
Male													
2000	759	-	4	1	2	6	23	81	140	227	203	72	-
1999	745	-	3	-	1	5	21	74	129	228	214	70	-
Female													
2001	581	-	1	2	1	8	22	62	84	137	163	101	-
2000	641	-	-	3	4	3	32	73	108	147	163	108	-
1999	642	-	-	1	2	9	25	55	106	144	203	97	-
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)													
All origins													
Both sexes													
2001	7,702	-	-	4	29	56	283	1,076	1,740	1,912	1,787	814	1
2000	7,492	-	-	4	26	50	293	1,068	1,642	1,959	1,626	824	-
1999	7,486	-	2	6	27	43	291	1,061	1,616	1,896	1,744	800	-
Male													
2001	5,156	-	-	1	12	35	202	846	1,372	1,327	1,025	335	1
2000	4,952	-	-	4	13	28	215	825	1,270	1,326	939	332	-
1999	4,944	-	1	3	20	27	204	820	1,231	1,308	1,007	323	-
Female													
2001	2,546	-	-	3	17	21	81	230	368	585	762	479	-
2000	2,540	-	-	-	13	22	78	243	372	633	687	492	-
1999	2,542	-	1	3	7	16	87	241	385	588	737	477	-
Hispanic													
Both sexes													
2001	311	-	-	-	4	1	14	56	69	78	58	31	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)													
Hispanic													
Both sexes													
2000	287	-	-	1	1	6	13	42	75	73	48	28	-
1999	282	-	-	1	-	1	18	41	66	72	63	20	-
Male													
2001	219	-	-	-	2	1	11	41	55	58	39	12	-
2000	218	-	-	1	-	4	11	34	62	58	28	20	-
1999	204	-	-	1	-	-	9	33	55	62	35	9	-
Female													
2001	92	-	-	-	2	-	3	15	14	20	19	19	-
2000	69	-	-	-	1	2	2	8	13	15	20	8	-
1999	78	-	-	-	-	1	9	8	11	10	28	11	-
Non-Hispanic [1]													
Both sexes													
2001	7,370	-	-	4	25	55	268	1,018	1,666	1,829	1,724	781	-
2000	7,181	-	-	3	24	44	280	1,022	1,560	1,877	1,575	796	-
1999	7,181	-	2	5	27	42	272	1,018	1,542	1,818	1,678	777	-
Male													
2001	4,920	-	-	1	10	34	190	804	1,313	1,264	982	322	-
2000	4,719	-	-	3	13	24	204	788	1,204	1,263	908	312	-
1999	4,725	-	1	2	20	27	194	785	1,172	1,241	970	313	-
Female													
2001	2,450	-	-	3	15	21	78	214	353	565	742	459	-
2000	2,462	-	-	-	11	20	76	234	356	614	667	484	-
1999	2,456	-	1	3	7	15	78	233	370	577	708	464	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)													
Non-Hispanic white													
Both sexes													
2001	6,052	-	-	2	18	36	165	736	1,288	1,524	1,560	723	-
2000	5,843	-	-	1	16	29	186	723	1,177	1,566	1,412	733	-
1999	5,826	-	2	3	14	29	176	677	1,180	1,527	1,497	721	-
Male													
2001	3,961	-	-	1	6	22	120	586	1,004	1,045	880	297	-
2000	3,726	-	-	1	7	16	128	546	904	1,036	797	291	-
1999	3,746	-	1	1	9	17	125	523	892	1,043	850	285	-
Female													
2001	2,091	-	-	1	12	14	45	150	284	479	680	426	-
2000	2,117	-	-	-	9	13	58	177	273	530	615	442	-
1999	2,080	-	1	2	5	12	51	154	288	484	647	436	-
Non-Hispanic black													
Both sexes													
2001	1,056	-	-	1	4	6	71	229	318	256	129	42	-
2000	1,089	-	-	2	7	7	60	249	327	254	136	47	-
1999	1,131	-	-	2	8	8	67	287	316	248	150	45	-
Male													
2001	773	-	-	-	2	5	45	179	263	184	78	17	-
2000	821	-	-	2	5	4	46	203	262	191	94	14	-
1999	819	-	-	1	6	6	47	224	243	169	102	21	-
Female													
2001	283	-	-	1	2	1	26	50	55	72	51	25	-
2000	268	-	-	-	2	3	14	46	65	63	42	33	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)													
Non-Hispanic black													
Female													
1999	312	-	-	1	2	2	20	63	73	79	48	24	-
Origin not stated [2]													
Both sexes													
2001	21	-	-	-	-	-	1	2	5	5	5	2	1
2000	24	-	-	-	1	-	-	4	7	9	3	-	-
1999	23	-	-	-	-	-	1	2	8	6	3	3	-
Male													
2001	17	-	-	-	-	-	1	1	4	5	4	1	1
2000	15	-	-	-	-	-	-	3	4	5	3	-	-
1999	15	-	-	-	-	-	1	2	4	5	2	1	-
Female													
2001	4	-	-	-	-	-	-	1	1	-	1	1	-
2000	9	-	-	-	1	-	-	1	3	4	-	-	-
1999	8	-	-	-	-	-	-	-	4	1	1	2	-
Malignant neoplasm of esophagus (C15)													
All origins													
Both sexes													
2001	12,530	-	-	-	6	29	309	1,408	2,554	3,779	3,307	1,137	1
2000	12,232	1	-	-	3	39	297	1,348	2,610	3,671	3,163	1,100	-
1999	11,918	-	-	-	-	37	293	1,262	2,531	3,694	2,986	1,114	1
Male													
2001	9,607	-	-	-	5	26	255	1,204	2,137	2,989	2,365	625	1
2000	9,279	1	-	-	2	31	265	1,161	2,169	2,831	2,233	586	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of esophagus (C15)													
All origins													
Male													
1999	9,094	-	-	-	-	29	255	1,094	2,081	2,925	2,136	573	1
Female													
2001	2,923	-	-	-	1	3	54	204	417	790	942	512	-
2000	2,953	-	-	-	1	8	32	187	441	840	930	514	-
1999	2,824	-	-	-	-	8	38	168	450	769	850	541	-
Hispanic													
Both sexes													
2001	401	-	-	-	-	2	12	63	96	95	101	32	-
2000	384	-	-	-	-	2	10	38	86	120	98	30	-
1999	367	-	-	-	-	2	16	44	85	95	87	38	-
Male													
2001	316	-	-	-	-	2	9	61	74	80	68	22	-
2000	292	-	-	-	-	2	8	34	64	92	72	20	-
1999	294	-	-	-	-	2	15	39	78	74	63	23	-
Female													
2001	85	-	-	-	-	-	3	2	22	15	33	10	-
2000	92	-	-	-	-	-	2	4	22	28	26	10	-
1999	73	-	-	-	-	-	1	5	7	21	24	15	-
Non-Hispanic [1]													
Both sexes													
2001	12,085	-	-	-	6	27	296	1,339	2,447	3,664	3,201	1,104	1
2000	11,819	1	-	-	3	37	287	1,308	2,521	3,538	3,058	1,066	-
1999	11,520	-	-	-	-	35	276	1,215	2,439	3,587	2,893	1,074	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of esophagus (C15)													
Non-Hispanic [1]													
Male													
2001	9,254	-	-	-	5	24	245	1,138	2,052	2,894	2,292	603	1
2000	8,964	1	-	-	2	29	257	1,126	2,102	2,728	2,156	563	-
1999	8,775	-	-	-	-	27	239	1,053	1,996	2,841	2,068	550	1
Female													
2001	2,831	-	-	-	1	3	51	201	395	770	909	501	-
2000	2,855	-	-	-	1	8	30	182	419	810	902	503	-
1999	2,745	-	-	-	-	8	37	162	443	746	825	524	-
Non-Hispanic white													
Both sexes													
2001	10,193	-	-	-	6	22	238	1,039	1,925	3,060	2,888	1,014	1
2000	10,005	-	-	-	3	36	239	1,013	1,997	3,017	2,714	986	-
1999	9,703	-	-	-	-	30	213	936	1,923	3,038	2,576	987	-
Male													
2001	7,863	-	-	-	5	19	200	897	1,645	2,458	2,086	552	1
2000	7,680	-	-	-	2	29	221	895	1,702	2,366	1,941	524	-
1999	7,440	-	-	-	-	22	188	827	1,598	2,451	1,857	497	-
Female													
2001	2,330	-	-	-	1	3	38	142	280	602	802	462	-
2000	2,325	-	-	-	1	7	18	118	295	651	773	462	-
1999	2,263	-	-	-	-	8	25	109	325	587	719	490	-
Non-Hispanic black													
Both sexes													
2001	1,676	-	-	-	-	3	54	264	466	534	278	77	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of esophagus (C15)													
Non-Hispanic black													
Both sexes													
2000	1,645	1	-	-	-	1	44	278	481	462	307	71	-
1999	1,629	-	-	-	-	3	53	261	472	486	280	73	1
Male													
2001	1,222	-	-	-	-	3	41	210	363	379	185	41	-
2000	1,162	1	-	-	-	-	32	218	369	318	191	33	-
1999	1,187	-	-	-	-	3	44	209	365	340	183	42	1
Female													
2001	454	-	-	-	-	-	13	54	103	155	93	36	-
2000	483	-	-	-	-	1	12	60	112	144	116	38	-
1999	442	-	-	-	-	-	9	52	107	146	97	31	-
Origin not stated [2]													
Both sexes													
2001	44	-	-	-	-	-	1	6	11	20	5	1	-
2000	29	-	-	-	-	-	-	2	3	13	7	4	-
1999	31	-	-	-	-	-	1	3	7	12	6	2	-
Male													
2001	37	-	-	-	-	-	1	5	11	15	5	-	-
2000	23	-	-	-	-	-	-	1	3	11	5	3	-
1999	25	-	-	-	-	-	1	2	7	10	5	-	-
Female													
2001	7	-	-	-	-	-	-	1	-	5	-	1	-
2000	6	-	-	-	-	-	-	1	-	2	2	1	-
1999	6	-	-	-	-	-	-	1	-	2	1	2	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of stomach (C16)													
All origins													
Both sexes													
2001	12,319	-	-	-	20	122	464	1,127	1,771	3,021	3,793	2,001	-
2000	12,645	-	-	1	14	124	484	1,097	1,804	3,054	3,833	2,234	-
1999	12,711	-	-	1	13	136	505	1,101	1,860	3,152	3,860	2,083	-
Male													
2001	7,191	-	-	-	9	71	253	731	1,238	1,928	2,148	813	-
2000	7,328	-	-	-	6	69	284	703	1,228	1,966	2,144	928	-
1999	7,459	-	-	1	6	76	283	746	1,252	2,005	2,239	851	-
Female													
2001	5,128	-	-	-	11	51	211	396	533	1,093	1,645	1,188	-
2000	5,317	-	-	1	8	55	200	394	576	1,088	1,689	1,306	-
1999	5,252	-	-	-	7	60	222	355	608	1,147	1,621	1,232	-
Hispanic													
Both sexes													
2001	1,189	-	-	-	4	41	98	182	209	300	253	102	-
2000	1,111	-	-	1	4	24	105	151	167	306	254	99	-
1999	1,055	-	-	-	4	38	85	121	192	271	239	105	-
Male													
2001	680	-	-	-	2	19	54	104	130	186	135	50	-
2000	655	-	-	-	1	16	52	78	110	205	144	49	-
1999	607	-	-	-	1	25	41	76	126	162	141	35	-
Female													
2001	509	-	-	-	2	22	44	78	79	114	118	52	-
2000	456	-	-	1	3	8	53	73	57	101	110	50	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of stomach (C16)													
Hispanic													
Female													
1999	448	-	-	-	3	13	44	45	66	109	98	70	-
Non-Hispanic [1]													
Both sexes													
2001	11,099	-	-	-	16	81	365	943	1,558	2,712	3,529	1,895	-
2000	11,491	-	-	-	10	99	378	943	1,627	2,739	3,568	2,127	-
1999	11,615	-	-	1	9	98	420	970	1,662	2,867	3,615	1,973	-
Male													
2001	6,492	-	-	-	7	52	199	625	1,105	1,737	2,007	760	-
2000	6,648	-	-	-	5	53	231	622	1,113	1,753	1,994	877	-
1999	6,827	-	-	1	5	51	242	662	1,122	1,834	2,096	814	-
Female													
2001	4,607	-	-	-	9	29	166	318	453	975	1,522	1,135	-
2000	4,843	-	-	-	5	46	147	321	514	986	1,574	1,250	-
1999	4,788	-	-	-	4	47	178	308	540	1,033	1,519	1,159	-
Non-Hispanic white													
Both sexes													
2001	8,308	-	-	-	6	43	211	615	1,110	2,010	2,779	1,534	-
2000	8,600	-	-	-	5	53	213	625	1,180	2,031	2,781	1,712	-
1999	8,712	-	-	-	4	50	237	646	1,156	2,181	2,859	1,579	-
Male													
2001	4,884	-	-	-	2	29	122	416	794	1,288	1,606	627	-
2000	5,033	-	-	-	2	29	129	424	825	1,330	1,580	714	-
1999	5,171	-	-	-	3	30	142	454	798	1,426	1,670	648	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of stomach (C16)													
Non-Hispanic white													
Female													
2001	3,424	-	-	-	4	14	89	199	316	722	1,173	907	-
2000	3,567	-	-	-	3	24	84	201	355	701	1,201	998	-
1999	3,541	-	-	-	1	20	95	192	358	755	1,189	931	-
Non-Hispanic black													
Both sexes													
2001	2,051	-	-	-	5	25	121	237	324	505	548	286	-
2000	2,156	-	-	-	4	36	117	239	335	523	579	323	-
1999	2,183	-	-	1	3	36	127	237	373	507	580	319	-
Male													
2001	1,177	-	-	-	2	14	63	161	220	332	293	92	-
2000	1,202	-	-	-	2	20	77	154	222	314	295	118	-
1999	1,252	-	-	1	1	19	72	158	241	309	325	126	-
Female													
2001	874	-	-	-	3	11	58	76	104	173	255	194	-
2000	954	-	-	-	2	16	40	85	113	209	284	205	-
1999	931	-	-	-	2	17	55	79	132	198	255	193	-
Origin not stated [2]													
Both sexes													
2001	31	-	-	-	-	-	1	2	4	9	11	4	-
2000	43	-	-	-	-	1	1	3	10	9	11	8	-
1999	41	-	-	-	-	-	-	10	6	14	6	5	-
Male													
2001	19	-	-	-	-	-	-	2	3	5	6	3	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of stomach (C16)													
Origin not stated [2]													
Male													
2000	25	-	-	-	-	-	1	3	5	8	6	2	-
1999	25	-	-	-	-	-	-	8	4	9	2	2	-
Female													
2001	12	-	-	-	-	-	1	-	1	4	5	1	-
2000	18	-	-	-	-	1	-	-	5	1	5	6	-
1999	16	-	-	-	-	-	-	2	2	5	4	3	-
Malignant neoplasms of colon, rectum and anus (C18-C21)													
All origins													
Both sexes													
2001	56,887	-	-	-	46	299	1,433	4,548	7,984	13,791	17,527	11,259	-
2000	57,477	-	-	3	40	272	1,369	4,237	7,940	14,101	18,053	11,462	-
1999	57,155	1	-	1	49	283	1,333	4,057	7,967	14,249	17,929	11,286	-
Male													
2001	28,225	-	-	-	24	157	781	2,512	4,652	7,773	8,451	3,875	-
2000	28,462	-	-	2	24	153	759	2,359	4,608	7,924	8,760	3,873	-
1999	28,240	-	-	1	27	160	678	2,250	4,621	7,883	8,754	3,866	-
Female													
2001	28,662	-	-	-	22	142	652	2,036	3,332	6,018	9,076	7,384	-
2000	29,015	-	-	1	16	119	610	1,878	3,332	6,177	9,293	7,589	-
1999	28,915	1	-	-	22	123	655	1,807	3,346	6,366	9,175	7,420	-
Hispanic													
Both sexes													
2001	2,250	-	-	-	6	20	127	245	415	592	535	310	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of colon, rectum and anus (C18-C21)													
Hispanic													
Both sexes													
2000	2,117	-	-	-	8	33	118	263	395	530	497	273	-
1999	2,031	-	-	-	7	31	86	245	344	581	485	252	-
Male													
2001	1,185	-	-	-	3	8	76	124	245	333	261	135	-
2000	1,124	-	-	-	3	17	52	140	227	301	267	117	-
1999	1,062	-	-	-	5	14	39	133	185	331	257	98	-
Female													
2001	1,065	-	-	-	3	12	51	121	170	259	274	175	-
2000	993	-	-	-	5	16	66	123	168	229	230	156	-
1999	969	-	-	-	2	17	47	112	159	250	228	154	-
Non-Hispanic [1]													
Both sexes													
2001	54,513	-	-	-	40	279	1,301	4,289	7,545	13,172	16,963	10,924	-
2000	55,244	-	-	3	32	238	1,247	3,963	7,523	13,544	17,527	11,167	-
1999	54,976	1	-	1	41	252	1,241	3,804	7,604	13,640	17,390	11,002	-
Male													
2001	26,978	-	-	-	21	149	701	2,382	4,390	7,419	8,181	3,735	-
2000	27,287	-	-	2	21	135	705	2,215	4,373	7,609	8,476	3,751	-
1999	27,103	-	-	1	22	146	636	2,113	4,424	7,538	8,467	3,756	-
Female													
2001	27,535	-	-	-	19	130	600	1,907	3,155	5,753	8,782	7,189	-
2000	27,957	-	-	1	11	103	542	1,748	3,150	5,935	9,051	7,416	-
1999	27,873	1	-	-	19	106	605	1,691	3,180	6,102	8,923	7,246	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of colon, rectum and anus (C18-C21)													
Non-Hispanic white													
Both sexes													
2001	46,450	-	-	-	28	203	966	3,284	6,092	11,124	14,903	9,850	-
2000	47,317	-	-	3	23	167	898	3,034	6,085	11,543	15,519	10,045	-
1999	47,275	-	-	1	26	182	933	2,956	6,184	11,643	15,427	9,923	-
Male													
2001	23,128	-	-	-	13	111	523	1,874	3,616	6,369	7,263	3,359	-
2000	23,369	-	-	2	13	101	515	1,732	3,561	6,553	7,539	3,353	-
1999	23,446	-	-	1	12	105	500	1,679	3,638	6,559	7,556	3,396	-
Female													
2001	23,322	-	-	-	15	92	443	1,410	2,476	4,755	7,640	6,491	-
2000	23,948	-	-	1	10	66	383	1,302	2,524	4,990	7,980	6,692	-
1999	23,829	-	-	-	14	77	433	1,277	2,546	5,084	7,871	6,527	-
Non-Hispanic black													
Both sexes													
2001	6,847	-	-	-	8	56	280	845	1,248	1,726	1,758	926	-
2000	6,820	-	-	-	6	54	290	819	1,215	1,720	1,732	984	-
1999	6,709	1	-	-	14	55	268	711	1,248	1,724	1,733	955	-
Male													
2001	3,221	-	-	-	7	30	152	428	652	881	766	305	-
2000	3,307	-	-	-	5	26	160	430	676	893	792	325	-
1999	3,160	-	-	-	9	30	123	374	692	850	786	296	-
Female													
2001	3,626	-	-	-	1	26	128	417	596	845	992	621	-
2000	3,513	-	-	-	1	28	130	389	539	827	940	659	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of colon, rectum and anus (C18-C21)													
Non-Hispanic black													
Female													
1999	3,549	1	-	-	5	25	145	337	556	874	947	659	-
Origin not stated [2]													
Both sexes													
2001	124	-	-	-	-	-	5	14	24	27	29	25	-
2000	116	-	-	-	-	1	4	11	22	27	29	22	-
1999	148	-	-	-	1	-	6	8	19	28	54	32	-
Male													
2001	62	-	-	-	-	-	4	6	17	21	9	5	-
2000	51	-	-	-	-	1	2	4	8	14	17	5	-
1999	75	-	-	-	-	-	3	4	12	14	30	12	-
Female													
2001	62	-	-	-	-	-	1	8	7	6	20	20	-
2000	65	-	-	-	-	-	2	7	14	13	12	17	-
1999	73	-	-	-	1	-	3	4	7	14	24	20	-
Malignant neoplasms of liver and intrahepatic bile ducts (C22)													
All origins													
Both sexes													
2001	13,351	5	18	15	29	75	401	1,868	2,351	3,440	3,649	1,500	-
2000	12,916	2	23	20	30	86	429	1,686	2,256	3,439	3,558	1,387	-
1999	12,382	8	29	11	43	78	444	1,492	2,091	3,491	3,425	1,270	-
Male													
2001	8,483	3	7	7	15	43	302	1,490	1,691	2,224	2,047	654	-
2000	8,176	1	14	8	20	59	297	1,324	1,607	2,246	2,052	548	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of liver and intrahepatic bile ducts (C22)													
All origins													
Male													
1999	7,689	5	20	9	24	48	318	1,161	1,465	2,284	1,877	478	-
Female													
2001	4,868	2	11	8	14	32	99	378	660	1,216	1,602	846	-
2000	4,740	1	9	12	10	27	132	362	649	1,193	1,506	839	-
1999	4,693	3	9	2	19	30	126	331	626	1,207	1,548	792	-
Hispanic													
Both sexes													
2001	1,256	1	3	3	2	14	46	226	261	327	289	84	-
2000	1,176	1	3	7	4	7	59	176	234	335	262	88	-
1999	1,084	1	4	2	4	7	43	170	214	324	239	76	-
Male													
2001	812	1	1	-	1	7	36	187	187	206	152	34	-
2000	762	1	2	4	3	5	45	141	165	208	152	36	-
1999	688	1	3	2	3	5	27	141	159	195	121	31	-
Female													
2001	444	-	2	3	1	7	10	39	74	121	137	50	-
2000	414	-	1	3	1	2	14	35	69	127	110	52	-
1999	396	-	1	-	1	2	16	29	55	129	118	45	-
Non-Hispanic [1]													
Both sexes													
2001	12,053	4	15	12	27	61	352	1,637	2,079	3,103	3,352	1,411	-
2000	11,702	1	20	13	26	78	368	1,502	2,018	3,091	3,287	1,298	-
1999	11,265	7	23	9	39	71	398	1,320	1,869	3,158	3,180	1,191	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of liver and intrahepatic bile ducts (C22)													
Non-Hispanic [1]													
Male													
2001	7,640	2	6	7	14	36	263	1,299	1,495	2,009	1,889	620	-
2000	7,384	-	12	4	17	53	251	1,175	1,440	2,028	1,893	511	-
1999	6,976	4	15	7	21	43	290	1,019	1,300	2,081	1,752	444	-
Female													
2001	4,413	2	9	5	13	25	89	338	584	1,094	1,463	791	-
2000	4,318	1	8	9	9	25	117	327	578	1,063	1,394	787	-
1999	4,289	3	8	2	18	28	108	301	569	1,077	1,428	747	-
Non-Hispanic white													
Both sexes													
2001	9,433	4	11	5	20	34	189	1,089	1,505	2,477	2,857	1,242	-
2000	9,121	-	14	8	16	37	201	987	1,447	2,448	2,824	1,139	-
1999	8,890	5	20	4	25	29	238	880	1,340	2,546	2,755	1,048	-
Male													
2001	5,956	2	4	4	11	17	137	859	1,088	1,651	1,632	551	-
2000	5,679	-	9	4	10	22	134	759	1,020	1,624	1,655	442	-
1999	5,455	2	13	3	14	16	173	660	934	1,702	1,534	404	-
Female													
2001	3,477	2	7	1	9	17	52	230	417	826	1,225	691	-
2000	3,442	-	5	4	6	15	67	228	427	824	1,169	697	-
1999	3,435	3	7	1	11	13	65	220	406	844	1,221	644	-
Non-Hispanic black													
Both sexes													
2001	1,649	-	3	5	4	15	95	385	373	357	299	113	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of liver and intrahepatic bile ducts (C22)													
Non-Hispanic black													
Both sexes													
2000	1,581	1	2	2	6	20	101	360	344	383	262	100	-
1999	1,498	1	3	4	7	21	91	291	312	395	265	108	-
Male													
2001	1,036	-	2	2	1	10	70	302	266	204	142	37	-
2000	1,047	-	1	-	5	14	65	285	256	249	131	41	-
1999	939	1	2	3	2	14	65	238	210	245	133	26	-
Female													
2001	613	-	1	3	3	5	25	83	107	153	157	76	-
2000	534	1	1	2	1	6	36	75	88	134	131	59	-
1999	559	-	1	1	5	7	26	53	102	150	132	82	-
Origin not stated [2]													
Both sexes													
2001	42	-	-	-	-	-	3	5	11	10	8	5	-
2000	38	-	-	-	-	1	2	8	4	13	9	1	-
1999	33	-	2	-	-	-	3	2	8	9	6	3	-
Male													
2001	31	-	-	-	-	-	3	4	9	9	6	-	-
2000	30	-	-	-	-	1	1	8	2	10	7	1	-
1999	25	-	2	-	-	-	1	1	6	8	4	3	-
Female													
2001	11	-	-	-	-	-	-	1	2	1	2	5	-
2000	8	-	-	-	-	-	1	-	2	3	2	-	-
1999	8	-	-	-	-	-	2	1	2	1	2	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of pancreas (C25)													
All origins													
Both sexes													
2001	29,803	-	1	2	2	55	574	2,410	4,868	8,158	9,322	4,410	1
2000	29,332	2	1	1	5	55	548	2,362	4,708	8,103	9,459	4,087	1
1999	29,082	1	-	-	3	55	594	2,353	4,707	8,293	8,933	4,142	1
Male													
2001	14,467	-	1	-	2	30	321	1,467	2,874	4,215	4,191	1,365	1
2000	14,238	1	1	-	3	33	324	1,486	2,749	4,189	4,189	1,262	1
1999	14,176	-	-	-	1	33	369	1,482	2,721	4,347	3,940	1,283	-
Female													
2001	15,336	-	-	2	-	25	253	943	1,994	3,943	5,131	3,045	-
2000	15,094	1	-	1	2	22	224	876	1,959	3,914	5,270	2,825	-
1999	14,906	1	-	-	2	22	225	871	1,986	3,946	4,993	2,859	1
Hispanic													
Both sexes													
2001	1,336	-	-	-	1	6	46	127	263	419	340	134	-
2000	1,243	1	-	-	2	6	37	128	234	397	313	125	-
1999	1,135	-	-	-	-	6	44	137	218	338	272	120	-
Male													
2001	643	-	-	-	1	3	26	73	141	212	154	33	-
2000	620	1	-	-	1	2	21	80	133	214	128	40	-
1999	563	-	-	-	-	2	30	84	117	182	108	40	-
Female													
2001	693	-	-	-	-	3	20	54	122	207	186	101	-
2000	623	-	-	-	1	4	16	48	101	183	185	85	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of pancreas (C25)													
Hispanic													
Female													
1999	572	-	-	-	-	4	14	53	101	156	164	80	-
Non-Hispanic [1]													
Both sexes													
2001	28,407	-	1	2	1	48	526	2,279	4,598	7,719	8,963	4,269	1
2000	28,026	1	1	1	3	49	510	2,228	4,465	7,692	9,124	3,951	1
1999	27,869	1	-	-	3	49	547	2,207	4,480	7,928	8,638	4,015	1
Male													
2001	13,792	-	1	-	1	27	294	1,393	2,728	3,989	4,031	1,327	1
2000	13,590	-	1	-	2	31	302	1,401	2,609	3,970	4,054	1,219	1
1999	13,572	-	-	-	1	31	338	1,391	2,598	4,150	3,822	1,241	-
Female													
2001	14,615	-	-	2	-	21	232	886	1,870	3,730	4,932	2,942	-
2000	14,436	1	-	1	1	18	208	827	1,856	3,722	5,070	2,732	-
1999	14,297	1	-	-	2	18	209	816	1,882	3,778	4,816	2,774	1
Non-Hispanic white													
Both sexes													
2001	24,424	-	1	2	1	36	415	1,816	3,847	6,564	7,931	3,810	1
2000	24,034	1	1	1	2	39	391	1,797	3,690	6,545	8,066	3,500	1
1999	23,903	-	-	-	2	34	421	1,771	3,675	6,784	7,625	3,590	1
Male													
2001	11,994	-	1	-	1	22	233	1,131	2,324	3,441	3,624	1,216	1
2000	11,764	-	1	-	2	25	236	1,141	2,183	3,444	3,638	1,093	1
1999	11,684	-	-	-	1	23	266	1,114	2,144	3,605	3,412	1,119	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of pancreas (C25)													
Non-Hispanic white													
Female													
2001	12,430	-	-	2	-	14	182	685	1,523	3,123	4,307	2,594	-
2000	12,270	1	-	1	-	14	155	656	1,507	3,101	4,428	2,407	-
1999	12,219	-	-	-	1	11	155	657	1,531	3,179	4,213	2,471	1
Non-Hispanic black													
Both sexes													
2001	3,352	-	-	-	-	9	89	399	638	961	865	391	-
2000	3,391	-	-	-	1	9	97	379	664	970	890	381	-
1999	3,368	1	-	-	-	12	112	388	669	972	846	368	-
Male													
2001	1,501	-	-	-	-	4	47	229	339	460	335	87	-
2000	1,546	-	-	-	-	5	54	229	367	447	350	94	-
1999	1,601	-	-	-	-	8	63	246	386	457	344	97	-
Female													
2001	1,851	-	-	-	-	5	42	170	299	501	530	304	-
2000	1,845	-	-	-	1	4	43	150	297	523	540	287	-
1999	1,767	1	-	-	-	4	49	142	283	515	502	271	-
Origin not stated [2]													
Both sexes													
2001	60	-	-	-	-	1	2	4	7	20	19	7	-
2000	63	-	-	-	-	-	1	6	9	14	22	11	-
1999	78	-	-	-	-	-	3	9	9	27	23	7	-
Male													
2001	32	-	-	-	-	-	1	1	5	14	6	5	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of pancreas (C25)													
Origin not stated [2]													
Male													
2000	28	-	-	-	-	-	1	5	7	5	7	3	-
1999	41	-	-	-	-	-	1	7	6	15	10	2	-
Female													
2001	28	-	-	-	-	1	1	3	2	6	13	2	-
2000	35	-	-	-	-	-	-	1	2	9	15	8	-
1999	37	-	-	-	-	-	2	2	3	12	13	5	-
Malignant neoplasm of larynx (C32)													
All origins													
Both sexes													
2001	3,797	-	-	-	-	3	86	419	913	1,170	910	296	-
2000	3,861	-	-	2	1	4	64	426	873	1,219	996	276	-
1999	3,815	-	-	-	-	6	78	405	904	1,239	892	291	-
Male													
2001	3,018	-	-	-	-	2	66	346	737	919	710	238	-
2000	3,041	-	-	-	-	2	49	340	726	945	765	214	-
1999	2,992	-	-	-	-	5	64	346	735	948	681	213	-
Female													
2001	779	-	-	-	-	1	20	73	176	251	200	58	-
2000	820	-	-	2	1	2	15	86	147	274	231	62	-
1999	823	-	-	-	-	1	14	59	169	291	211	78	-
Hispanic													
Both sexes													
2001	163	-	-	-	-	-	3	20	38	60	29	13	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of larynx (C32)													
Hispanic													
Both sexes													
2000	143	-	-	1	-	1	4	17	29	45	35	11	-
1999	171	-	-	-	-	-	5	25	29	55	40	17	-
Male													
2001	141	-	-	-	-	-	3	18	34	53	24	9	-
2000	124	-	-	-	-	-	3	16	28	37	30	10	-
1999	149	-	-	-	-	-	4	24	27	48	31	15	-
Female													
2001	22	-	-	-	-	-	-	2	4	7	5	4	-
2000	19	-	-	1	-	1	1	1	1	8	5	1	-
1999	22	-	-	-	-	-	1	1	2	7	9	2	-
Non-Hispanic [1]													
Both sexes													
2001	3,614	-	-	-	-	3	82	398	865	1,106	877	283	-
2000	3,699	-	-	1	1	3	59	406	840	1,166	959	264	-
1999	3,627	-	-	-	-	6	73	377	871	1,180	847	273	-
Male													
2001	2,858	-	-	-	-	2	62	327	694	862	682	229	-
2000	2,901	-	-	-	-	2	45	322	694	901	733	204	-
1999	2,832	-	-	-	-	5	60	320	706	897	646	198	-
Female													
2001	756	-	-	-	-	1	20	71	171	244	195	54	-
2000	798	-	-	1	1	1	14	84	146	265	226	60	-
1999	795	-	-	-	-	1	13	57	165	283	201	75	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of larynx (C32)													
Non-Hispanic white													
Both sexes													
2001	2,876	-	-	-	-	3	58	268	657	893	747	250	-
2000	2,940	-	-	1	1	3	44	281	612	936	822	240	-
1999	2,924	-	-	-	-	4	40	256	660	978	751	235	-
Male													
2001	2,267	-	-	-	-	2	47	216	529	697	570	206	-
2000	2,291	-	-	-	-	2	31	220	500	721	632	185	-
1999	2,263	-	-	-	-	3	34	219	534	739	568	166	-
Female													
2001	609	-	-	-	-	1	11	52	128	196	177	44	-
2000	649	-	-	1	1	1	13	61	112	215	190	55	-
1999	661	-	-	-	-	1	6	37	126	239	183	69	-
Non-Hispanic black													
Both sexes													
2001	668	-	-	-	-	-	23	124	195	188	111	27	-
2000	714	-	-	-	-	-	15	121	218	209	129	22	-
1999	657	-	-	-	-	2	32	117	203	188	89	26	-
Male													
2001	531	-	-	-	-	-	14	105	156	144	94	18	-
2000	574	-	-	-	-	-	14	99	186	164	94	17	-
1999	532	-	-	-	-	2	25	97	168	147	72	21	-
Female													
2001	137	-	-	-	-	-	9	19	39	44	17	9	-
2000	140	-	-	-	-	-	1	22	32	45	35	5	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of larynx (C32)													
Non-Hispanic black													
Female													
1999	125	-	-	-	-	-	7	20	35	41	17	5	-
Origin not stated [2]													
Both sexes													
2001	20	-	-	-	-	-	1	1	10	4	4	-	-
2000	19	-	-	-	-	-	1	3	4	8	2	1	-
1999	17	-	-	-	-	-	-	3	4	4	5	1	-
Male													
2001	19	-	-	-	-	-	1	1	9	4	4	-	-
2000	16	-	-	-	-	-	1	2	4	7	2	-	-
1999	11	-	-	-	-	-	-	2	2	3	4	-	-
Female													
2001	1	-	-	-	-	-	-	-	1	-	-	-	-
2000	3	-	-	-	-	-	-	1	-	1	-	1	-
1999	6	-	-	-	-	-	-	1	2	1	1	1	-
Malignant neoplasms of trachea, bronchus and lung (C33-C34)													
All origins													
Both sexes													
2001	156,058	-	5	2	20	157	2,796	12,041	29,777	51,225	46,700	13,331	4
2000	155,521	2	2	1	22	197	2,756	11,915	29,721	52,258	45,836	12,806	5
1999	152,156	-	-	5	20	179	2,724	11,434	29,344	51,848	44,261	12,339	2
Male													
2001	90,426	-	4	1	12	77	1,504	7,190	17,873	30,699	26,406	6,656	4
2000	90,469	2	1	-	14	102	1,542	7,123	17,937	31,382	25,966	6,395	5

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of trachea, bronchus and lung (C33-C34)													
All origins													
Male													
1999	89,453	-	-	4	13	97	1,474	6,851	18,013	31,287	25,397	6,315	2
Female													
2001	65,632	-	1	1	8	80	1,292	4,851	11,904	20,526	20,294	6,675	-
2000	65,052	-	1	1	8	95	1,214	4,792	11,784	20,876	19,870	6,411	-
1999	62,703	-	-	1	7	82	1,250	4,583	11,331	20,561	18,864	6,024	-
Hispanic													
Both sexes													
2001	3,816	-	2	1	7	16	106	364	758	1,230	1,011	321	-
2000	3,643	-	-	-	3	17	90	342	672	1,191	1,006	322	-
1999	3,545	-	-	-	3	17	98	308	724	1,170	901	324	-
Male													
2001	2,495	-	2	-	6	10	68	239	517	820	657	176	-
2000	2,411	-	-	-	3	10	56	229	471	793	666	183	-
1999	2,373	-	-	-	2	10	58	208	523	798	606	168	-
Female													
2001	1,321	-	-	1	1	6	38	125	241	410	354	145	-
2000	1,232	-	-	-	-	7	34	113	201	398	340	139	-
1999	1,172	-	-	-	1	7	40	100	201	372	295	156	-
Non-Hispanic [1]													
Both sexes													
2001	151,882	-	3	1	13	141	2,678	11,642	28,936	49,887	45,599	12,978	4
2000	151,483	2	2	1	19	180	2,653	11,525	28,965	50,940	44,742	12,449	5
1999	148,206	-	-	5	17	161	2,618	11,087	28,549	50,536	43,243	11,988	2

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of trachea, bronchus and lung (C33-C34)													
Non-Hispanic [1]													
Male													
2001	87,706	-	2	1	6	67	1,431	6,929	17,291	29,818	25,695	6,462	4
2000	87,816	2	1	-	11	92	1,479	6,866	17,409	30,511	25,245	6,195	5
1999	86,822	-	-	4	11	87	1,411	6,616	17,438	30,397	24,724	6,132	2
Female													
2001	64,176	-	1	-	7	74	1,247	4,713	11,645	20,069	19,904	6,516	-
2000	63,667	-	1	1	8	88	1,174	4,659	11,556	20,429	19,497	6,254	-
1999	61,384	-	-	1	6	74	1,207	4,471	11,111	20,139	18,519	5,856	-
Non-Hispanic white													
Both sexes													
2001	133,266	-	2	1	9	114	2,100	9,301	24,713	43,972	41,223	11,829	2
2000	133,061	1	2	1	11	139	1,996	9,176	24,777	44,952	40,666	11,337	3
1999	129,967	-	-	3	9	131	2,025	8,832	24,329	44,577	39,153	10,907	1
Male													
2001	76,066	-	1	1	4	55	1,095	5,421	14,552	26,056	23,022	5,857	2
2000	76,218	1	1	-	7	73	1,076	5,355	14,682	26,704	22,718	5,598	3
1999	75,338	-	-	2	6	71	1,078	5,155	14,654	26,596	22,219	5,556	1
Female													
2001	57,200	-	1	-	5	59	1,005	3,880	10,161	17,916	18,201	5,972	-
2000	56,843	-	1	1	4	66	920	3,821	10,095	18,248	17,948	5,739	-
1999	54,629	-	-	1	3	60	947	3,677	9,675	17,981	16,934	5,351	-
Non-Hispanic black													
Both sexes													
2001	15,883	-	1	-	3	16	515	2,060	3,685	5,076	3,623	902	2

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of trachea, bronchus and lung (C33-C34)													
Non-Hispanic black													
Both sexes													
2000	15,924	1	-	-	5	31	578	2,113	3,688	5,178	3,414	915	1
1999	15,834	-	-	2	8	20	531	2,029	3,743	5,111	3,483	907	-
Male													
2001	10,030	-	1	-	1	7	298	1,352	2,400	3,266	2,226	477	2
2000	10,056	1	-	-	2	14	359	1,372	2,410	3,309	2,105	483	1
1999	10,039	-	-	2	5	11	301	1,338	2,500	3,267	2,127	488	-
Female													
2001	5,853	-	-	-	2	9	217	708	1,285	1,810	1,397	425	-
2000	5,868	-	-	-	3	17	219	741	1,278	1,869	1,309	432	-
1999	5,795	-	-	-	3	9	230	691	1,243	1,844	1,356	419	-
Origin not stated [2]													
Both sexes													
2001	360	-	-	-	-	-	12	35	83	108	90	32	-
2000	395	-	-	-	-	-	13	48	84	127	88	35	-
1999	405	-	-	-	-	1	8	39	71	142	117	27	-
Male													
2001	225	-	-	-	-	-	5	22	65	61	54	18	-
2000	242	-	-	-	-	-	7	28	57	78	55	17	-
1999	258	-	-	-	-	-	5	27	52	92	67	15	-
Female													
2001	135	-	-	-	-	-	7	13	18	47	36	14	-
2000	153	-	-	-	-	-	6	20	27	49	33	18	-
1999	147	-	-	-	-	1	3	12	19	50	50	12	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant melanoma of skin (C43)													
All origins													
Both sexes													
2001	7,542	-	-	-	41	186	622	1,208	1,333	1,689	1,678	785	-
2000	7,420	-	-	1	41	186	626	1,199	1,303	1,637	1,624	803	-
1999	7,215	1	-	1	39	194	625	1,120	1,263	1,652	1,639	681	-
Male													
2001	4,763	-	-	-	24	103	380	756	893	1,156	1,063	388	-
2000	4,592	-	-	1	21	110	355	769	866	1,067	1,013	390	-
1999	4,529	1	-	-	23	110	397	717	851	1,096	1,006	328	-
Female													
2001	2,779	-	-	-	17	83	242	452	440	533	615	397	-
2000	2,828	-	-	-	20	76	271	430	437	570	611	413	-
1999	2,686	-	-	1	16	84	228	403	412	556	633	353	-
Hispanic													
Both sexes													
2001	145	-	-	-	2	6	24	23	25	28	28	9	-
2000	144	-	-	-	2	5	26	28	27	26	19	11	-
1999	133	1	-	1	2	5	14	22	30	27	20	11	-
Male													
2001	78	-	-	-	-	2	17	11	15	19	13	1	-
2000	92	-	-	-	1	4	17	19	19	14	13	5	-
1999	79	1	-	-	1	4	9	16	19	17	9	3	-
Female													
2001	67	-	-	-	2	4	7	12	10	9	15	8	-
2000	52	-	-	-	1	1	9	9	8	12	6	6	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant melanoma of skin (C43)													
Hispanic													
Female													
1999	54	-	-	1	1	1	5	6	11	10	11	8	-
Non-Hispanic [1]													
Both sexes													
2001	7,387	-	-	-	39	179	597	1,184	1,307	1,660	1,646	775	-
2000	7,263	-	-	1	39	181	599	1,167	1,276	1,608	1,602	790	-
1999	7,062	-	-	-	37	189	611	1,094	1,231	1,620	1,612	668	-
Male													
2001	4,678	-	-	-	24	100	363	744	877	1,136	1,047	387	-
2000	4,490	-	-	1	20	106	337	746	847	1,051	998	384	-
1999	4,438	-	-	-	22	106	388	698	831	1,076	994	323	-
Female													
2001	2,709	-	-	-	15	79	234	440	430	524	599	388	-
2000	2,773	-	-	-	19	75	262	421	429	557	604	406	-
1999	2,624	-	-	-	15	83	223	396	400	544	618	345	-
Non-Hispanic white													
Both sexes													
2001	7,250	-	-	-	38	177	590	1,168	1,286	1,616	1,614	761	-
2000	7,090	-	-	1	38	175	579	1,147	1,247	1,570	1,566	767	-
1999	6,936	-	-	-	34	183	604	1,081	1,213	1,593	1,577	651	-
Male													
2001	4,618	-	-	-	23	98	359	737	867	1,116	1,034	384	-
2000	4,414	-	-	1	20	104	327	737	833	1,032	981	379	-
1999	4,370	-	-	-	19	102	385	689	819	1,064	976	316	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant melanoma of skin (C43)													
Non-Hispanic white													
Female													
2001	2,632	-	-	-	15	79	231	431	419	500	580	377	-
2000	2,676	-	-	-	18	71	252	410	414	538	585	388	-
1999	2,566	-	-	-	15	81	219	392	394	529	601	335	-
Non-Hispanic black													
Both sexes													
2001	102	-	-	-	1	1	7	9	13	36	24	11	-
2000	134	-	-	-	1	3	15	13	19	30	32	21	-
1999	94	-	-	-	1	4	4	9	14	23	25	14	-
Male													
2001	36	-	-	-	1	1	4	3	4	16	6	1	-
2000	59	-	-	-	-	2	9	6	10	14	15	3	-
1999	47	-	-	-	1	2	1	6	10	9	12	6	-
Female													
2001	66	-	-	-	-	-	3	6	9	20	18	10	-
2000	75	-	-	-	1	1	6	7	9	16	17	18	-
1999	47	-	-	-	-	2	3	3	4	14	13	8	-
Origin not stated [2]													
Both sexes													
2001	10	-	-	-	-	1	1	1	1	1	4	1	-
2000	13	-	-	-	-	-	1	4	-	3	3	2	-
1999	20	-	-	-	-	-	-	4	2	5	7	2	-
Male													
2001	7	-	-	-	-	1	-	1	1	1	3	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant melanoma of skin (C43)													
Origin not stated [2]													
Male													
2000	10	-	-	-	-	-	1	4	-	2	2	1	-
1999	12	-	-	-	-	-	-	3	1	3	3	2	-
Female													
2001	3	-	-	-	-	-	1	-	-	-	1	1	-
2000	3	-	-	-	-	-	-	-	-	1	1	1	-
1999	8	-	-	-	-	-	-	1	1	2	4	-	-
Malignant neoplasm of breast (C50)													
All origins													
Both sexes													
2001	41,809	-	-	-	9	463	2,823	6,593	7,632	8,715	9,663	5,911	-
2000	42,300	-	-	-	11	467	2,841	6,385	7,573	9,024	9,750	6,249	-
1999	41,528	-	1	1	13	439	2,741	6,174	7,399	9,109	9,651	6,000	-
Male													
2001	415	-	-	-	-	2	14	46	72	120	115	46	-
2000	428	-	-	-	-	3	19	46	85	115	109	51	-
1999	384	-	-	-	-	3	15	41	70	103	98	54	-
Female													
2001	41,394	-	-	-	9	461	2,809	6,547	7,560	8,595	9,548	5,865	-
2000	41,872	-	-	-	11	464	2,822	6,339	7,488	8,909	9,641	6,198	-
1999	41,144	-	1	1	13	436	2,726	6,133	7,329	9,006	9,553	5,946	-
Hispanic													
Both sexes													
2001	1,760	-	-	-	1	53	244	408	407	318	219	110	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of breast (C50)													
Hispanic													
Both sexes													
2000	1,689	-	-	-	1	54	218	381	359	337	231	108	-
1999	1,551	-	-	-	2	47	197	362	321	290	209	123	-
Male													
2001	14	-	-	-	-	-	2	1	1	5	5	-	-
2000	19	-	-	-	-	-	2	2	4	6	4	1	-
1999	13	-	-	-	-	-	-	3	3	5	2	-	-
Female													
2001	1,746	-	-	-	1	53	242	407	406	313	214	110	-
2000	1,670	-	-	-	1	54	216	379	355	331	227	107	-
1999	1,538	-	-	-	2	47	197	359	318	285	207	123	-
Non-Hispanic [1]													
Both sexes													
2001	39,957	-	-	-	8	409	2,576	6,170	7,208	8,379	9,418	5,789	-
2000	40,506	-	-	-	10	413	2,615	5,989	7,192	8,670	9,494	6,123	-
1999	39,881	-	1	1	11	390	2,537	5,800	7,056	8,803	9,419	5,863	-
Male													
2001	399	-	-	-	-	2	12	45	70	114	110	46	-
2000	409	-	-	-	-	3	17	44	81	109	105	50	-
1999	370	-	-	-	-	3	15	38	67	97	96	54	-
Female													
2001	39,558	-	-	-	8	407	2,564	6,125	7,138	8,265	9,308	5,743	-
2000	40,097	-	-	-	10	410	2,598	5,945	7,111	8,561	9,389	6,073	-
1999	39,511	-	1	1	11	387	2,522	5,762	6,989	8,706	9,323	5,809	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of breast (C50)													
Non-Hispanic white													
Both sexes													
2001	33,629	-	-	-	3	255	1,828	4,737	5,895	7,185	8,472	5,254	-
2000	34,386	-	-	-	4	275	1,877	4,620	5,911	7,581	8,591	5,527	-
1999	33,790	-	-	1	6	257	1,802	4,470	5,779	7,623	8,488	5,364	-
Male													
2001	329	-	-	-	-	1	6	33	59	95	96	39	-
2000	341	-	-	-	-	1	9	35	63	97	94	42	-
1999	299	-	-	-	-	2	8	28	51	79	84	47	-
Female													
2001	33,300	-	-	-	3	254	1,822	4,704	5,836	7,090	8,376	5,215	-
2000	34,045	-	-	-	4	274	1,868	4,585	5,848	7,484	8,497	5,485	-
1999	33,491	-	-	1	6	255	1,794	4,442	5,728	7,544	8,404	5,317	-
Non-Hispanic black													
Both sexes													
2001	5,528	-	-	-	5	140	641	1,209	1,125	1,053	862	493	-
2000	5,384	-	-	-	5	120	641	1,167	1,103	961	825	562	-
1999	5,386	-	1	-	5	124	661	1,150	1,095	1,037	852	461	-
Male													
2001	64	-	-	-	-	1	5	12	10	17	13	6	-
2000	63	-	-	-	-	2	8	8	16	11	10	8	-
1999	67	-	-	-	-	1	7	9	16	16	11	7	-
Female													
2001	5,464	-	-	-	5	139	636	1,197	1,115	1,036	849	487	-
2000	5,321	-	-	-	5	118	633	1,159	1,087	950	815	554	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of breast (C50)													
Non-Hispanic black													
Female													
1999	5,319	-	1	-	5	123	654	1,141	1,079	1,021	841	454	-
Origin not stated [2]													
Both sexes													
2001	92	-	-	-	-	1	3	15	17	18	26	12	-
2000	105	-	-	-	-	-	8	15	22	17	25	18	-
1999	96	-	-	-	-	2	7	12	22	16	23	14	-
Male													
2001	2	-	-	-	-	-	-	-	1	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	1	-	-	-
Female													
2001	90	-	-	-	-	1	3	15	16	17	26	12	-
2000	105	-	-	-	-	-	8	15	22	17	25	18	-
1999	95	-	-	-	-	2	7	12	22	15	23	14	-
Malignant neoplasm of cervix uteri (C53)													
All origins													
Both sexes													
2001	4,092	-	-	1	12	209	711	953	723	585	583	315	-
2000	4,200	-	-	1	17	230	694	915	748	686	599	310	-
1999	4,205	-	-	-	12	217	699	914	769	696	568	329	1
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of cervix uteri (C53)													
All origins													
Male													
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	4,092	-	-	1	12	209	711	953	723	585	583	315	-
2000	4,200	-	-	1	17	230	694	915	748	686	599	310	-
1999	4,205	-	-	-	12	217	699	914	769	696	568	329	1
Hispanic													
Both sexes													
2001	412	-	-	-	3	36	102	102	75	39	39	16	-
2000	434	-	-	-	4	33	108	87	78	60	43	21	-
1999	378	-	-	-	2	31	87	99	64	51	31	13	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	412	-	-	-	3	36	102	102	75	39	39	16	-
2000	434	-	-	-	4	33	108	87	78	60	43	21	-
1999	378	-	-	-	2	31	87	99	64	51	31	13	-
Non-Hispanic [1]													
Both sexes													
2001	3,669	-	-	1	9	172	606	846	646	546	544	299	-
2000	3,755	-	-	1	13	197	583	825	668	625	554	289	-
1999	3,817	-	-	-	10	186	611	813	702	644	535	315	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of cervix uteri (C53)													
Non-Hispanic [1]													
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	3,669	-	-	1	9	172	606	846	646	546	544	299	-
2000	3,755	-	-	1	13	197	583	825	668	625	554	289	-
1999	3,817	-	-	-	10	186	611	813	702	644	535	315	1
Non-Hispanic white													
Both sexes													
2001	2,725	-	-	-	8	125	448	632	478	392	409	233	-
2000	2,725	-	-	-	9	141	403	600	479	438	434	221	-
1999	2,854	-	-	-	7	147	451	630	508	454	414	242	1
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	2,725	-	-	-	8	125	448	632	478	392	409	233	-
2000	2,725	-	-	-	9	141	403	600	479	438	434	221	-
1999	2,854	-	-	-	7	147	451	630	508	454	414	242	1
Non-Hispanic black													
Both sexes													
2001	778	-	-	1	1	42	137	165	131	129	111	61	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of cervix uteri (C53)													
Non-Hispanic black													
Both sexes													
2000	860	-	-	1	4	51	147	177	152	158	108	62	-
1999	825	-	-	-	3	34	140	140	167	173	101	67	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	778	-	-	1	1	42	137	165	131	129	111	61	-
2000	860	-	-	1	4	51	147	177	152	158	108	62	-
1999	825	-	-	-	3	34	140	140	167	173	101	67	-
Origin not stated [2]													
Both sexes													
2001	11	-	-	-	-	1	3	5	2	-	-	-	-
2000	11	-	-	-	-	-	3	3	2	1	2	-	-
1999	10	-	-	-	-	-	1	2	3	1	2	1	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	11	-	-	-	-	1	3	5	2	-	-	-	-
2000	11	-	-	-	-	-	3	3	2	1	2	-	-
1999	10	-	-	-	-	-	1	2	3	1	2	1	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)													
All origins													
Both sexes													
2001	6,784	-	-	-	-	38	139	566	1,105	1,784	2,063	1,088	1
2000	6,586	-	-	-	3	27	138	507	1,122	1,781	1,958	1,049	1
1999	6,468	-	-	-	2	23	142	468	1,019	1,807	2,008	999	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	6,784	-	-	-	-	38	139	566	1,105	1,784	2,063	1,088	1
2000	6,586	-	-	-	3	27	138	507	1,122	1,781	1,958	1,049	1
1999	6,468	-	-	-	2	23	142	468	1,019	1,807	2,008	999	-
Hispanic													
Both sexes													
2001	319	-	-	-	-	7	13	48	71	85	65	30	-
2000	277	-	-	-	-	2	19	35	62	75	54	30	-
1999	272	-	-	-	1	3	15	33	61	86	51	22	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	319	-	-	-	-	7	13	48	71	85	65	30	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)													
Hispanic													
Female													
2000	277	-	-	-	-	2	19	35	62	75	54	30	-
1999	272	-	-	-	1	3	15	33	61	86	51	22	-
Non-Hispanic [1]													
Both sexes													
2001	6,449	-	-	-	-	31	126	517	1,029	1,697	1,996	1,052	1
2000	6,294	-	-	-	3	25	119	469	1,057	1,702	1,901	1,017	1
1999	6,188	-	-	-	1	20	127	435	956	1,719	1,954	976	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	6,449	-	-	-	-	31	126	517	1,029	1,697	1,996	1,052	1
2000	6,294	-	-	-	3	25	119	469	1,057	1,702	1,901	1,017	1
1999	6,188	-	-	-	1	20	127	435	956	1,719	1,954	976	-
Non-Hispanic white													
Both sexes													
2001	5,283	-	-	-	-	19	94	398	802	1,319	1,712	939	-
2000	5,113	-	-	-	1	18	88	383	827	1,298	1,594	904	-
1999	5,114	-	-	-	1	15	96	353	753	1,351	1,687	858	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)													
Non-Hispanic white													
Male													
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	5,283	-	-	-	-	19	94	398	802	1,319	1,712	939	-
2000	5,113	-	-	-	1	18	88	383	827	1,298	1,594	904	-
1999	5,114	-	-	-	1	15	96	353	753	1,351	1,687	858	-
Non-Hispanic black													
Both sexes													
2001	1,037	-	-	-	-	9	22	93	190	352	265	105	1
2000	1,055	-	-	-	1	6	21	68	201	371	280	106	1
1999	980	-	-	-	-	5	27	57	189	346	245	111	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	1,037	-	-	-	-	9	22	93	190	352	265	105	1
2000	1,055	-	-	-	1	6	21	68	201	371	280	106	1
1999	980	-	-	-	-	5	27	57	189	346	245	111	-
Origin not stated [2]													
Both sexes													
2001	16	-	-	-	-	-	-	1	5	2	2	6	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)													
Origin not stated [2]													
Both sexes													
2000	15	-	-	-	-	-	-	3	3	4	3	2	-
1999	8	-	-	-	-	-	-	-	2	2	3	1	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	16	-	-	-	-	-	-	1	5	2	2	6	-
2000	15	-	-	-	-	-	-	3	3	4	3	2	-
1999	8	-	-	-	-	-	-	-	2	2	3	1	-
Malignant neoplasm of ovary (C56)													
All origins													
Both sexes													
2001	14,414	-	-	2	21	102	472	1,626	2,612	3,666	4,170	1,743	-
2000	14,060	-	-	2	23	101	472	1,583	2,444	3,601	4,098	1,736	-
1999	13,627	-	1	2	27	77	496	1,549	2,426	3,612	3,875	1,562	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	14,414	-	-	2	21	102	472	1,626	2,612	3,666	4,170	1,743	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of ovary (C56)													
All origins													
Female													
2000	14,060	-	-	2	23	101	472	1,583	2,444	3,601	4,098	1,736	-
1999	13,627	-	1	2	27	77	496	1,549	2,426	3,612	3,875	1,562	-
Hispanic													
Both sexes													
2001	603	-	-	1	-	13	43	91	131	164	111	49	-
2000	578	-	-	1	4	9	30	94	111	154	126	49	-
1999	478	-	1	-	2	6	29	80	100	126	100	34	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	603	-	-	1	-	13	43	91	131	164	111	49	-
2000	578	-	-	1	4	9	30	94	111	154	126	49	-
1999	478	-	1	-	2	6	29	80	100	126	100	34	-
Non-Hispanic [1]													
Both sexes													
2001	13,783	-	-	1	21	89	428	1,533	2,475	3,495	4,052	1,689	-
2000	13,454	-	-	1	19	92	441	1,484	2,329	3,440	3,964	1,684	-
1999	13,103	-	-	2	25	71	466	1,466	2,316	3,474	3,756	1,527	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of ovary (C56)													
Non-Hispanic [1]													
Male													
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	13,783	-	-	1	21	89	428	1,533	2,475	3,495	4,052	1,689	-
2000	13,454	-	-	1	19	92	441	1,484	2,329	3,440	3,964	1,684	-
1999	13,103	-	-	2	25	71	466	1,466	2,316	3,474	3,756	1,527	-
Non-Hispanic white													
Both sexes													
2001	12,338	-	-	-	14	68	347	1,335	2,151	3,125	3,741	1,557	-
2000	12,138	-	-	-	13	75	367	1,284	2,072	3,087	3,680	1,560	-
1999	11,795	-	-	2	20	51	394	1,268	2,061	3,119	3,456	1,424	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	12,338	-	-	-	14	68	347	1,335	2,151	3,125	3,741	1,557	-
2000	12,138	-	-	-	13	75	367	1,284	2,072	3,087	3,680	1,560	-
1999	11,795	-	-	2	20	51	394	1,268	2,061	3,119	3,456	1,424	-
Non-Hispanic black													
Both sexes													
2001	1,140	-	-	1	4	15	65	134	251	301	252	117	-
2000	1,065	-	-	1	2	12	59	138	205	295	244	109	-
1999	1,065	-	-	-	4	13	47	143	211	301	256	90	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of ovary (C56)													
Non-Hispanic black													
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	1,140	-	-	1	4	15	65	134	251	301	252	117	-
2000	1,065	-	-	1	2	12	59	138	205	295	244	109	-
1999	1,065	-	-	-	4	13	47	143	211	301	256	90	-
Origin not stated [2]													
Both sexes													
2001	28	-	-	-	-	-	1	2	6	7	7	5	-
2000	28	-	-	-	-	-	1	5	4	7	8	3	-
1999	46	-	-	-	-	-	1	3	10	12	19	1	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	28	-	-	-	-	-	1	2	6	7	7	5	-
2000	28	-	-	-	-	-	1	5	4	7	8	3	-
1999	46	-	-	-	-	-	1	3	10	12	19	1	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of prostate (C61)													
All origins													
Both sexes													
2001	30,719	-	-	-	1	1	30	412	1,907	6,543	12,976	8,849	-
2000	31,078	-	-	-	-	1	25	371	1,929	6,808	13,071	8,873	-
1999	31,729	1	-	-	1	2	33	374	1,946	7,316	13,435	8,620	1
Male													
2001	30,719	-	-	-	1	1	30	412	1,907	6,543	12,976	8,849	-
2000	31,078	-	-	-	-	1	25	371	1,929	6,808	13,071	8,873	-
1999	31,729	1	-	-	1	2	33	374	1,946	7,316	13,435	8,620	1
Female													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Hispanic													
Both sexes													
2001	1,226	-	-	-	-	-	-	27	108	332	487	272	-
2000	1,033	-	-	-	-	-	1	26	78	277	401	250	-
1999	1,050	-	-	-	1	1	2	22	92	295	386	251	-
Male													
2001	1,226	-	-	-	-	-	-	27	108	332	487	272	-
2000	1,033	-	-	-	-	-	1	26	78	277	401	250	-
1999	1,050	-	-	-	1	1	2	22	92	295	386	251	-
Female													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of prostate (C61)													
Hispanic													
Female													
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic [1]													
Both sexes													
2001	29,407	-	-	-	1	1	29	385	1,790	6,197	12,449	8,555	-
2000	29,964	-	-	-	-	1	24	344	1,846	6,512	12,634	8,603	-
1999	30,614	1	-	-	-	1	31	351	1,848	7,004	13,025	8,352	1
Male													
2001	29,407	-	-	-	1	1	29	385	1,790	6,197	12,449	8,555	-
2000	29,964	-	-	-	-	1	24	344	1,846	6,512	12,634	8,603	-
1999	30,614	1	-	-	-	1	31	351	1,848	7,004	13,025	8,352	1
Female													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic white													
Both sexes													
2001	23,811	-	-	-	1	-	18	258	1,285	4,755	10,230	7,264	-
2000	24,287	-	-	-	-	1	15	237	1,325	4,946	10,399	7,364	-
1999	24,936	1	-	-	-	1	23	244	1,310	5,413	10,850	7,093	1
Male													
2001	23,811	-	-	-	1	-	18	258	1,285	4,755	10,230	7,264	-
2000	24,287	-	-	-	-	1	15	237	1,325	4,946	10,399	7,364	-
1999	24,936	1	-	-	-	1	23	244	1,310	5,413	10,850	7,093	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of prostate (C61)													
Non-Hispanic white													
Female													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic black													
Both sexes													
2001	5,215	-	-	-	-	1	11	124	481	1,372	2,054	1,172	-
2000	5,292	-	-	-	-	-	9	100	494	1,481	2,077	1,131	-
1999	5,315	-	-	-	-	-	7	105	512	1,524	2,027	1,140	-
Male													
2001	5,215	-	-	-	-	1	11	124	481	1,372	2,054	1,172	-
2000	5,292	-	-	-	-	-	9	100	494	1,481	2,077	1,131	-
1999	5,315	-	-	-	-	-	7	105	512	1,524	2,027	1,140	-
Female													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Origin not stated [2]													
Both sexes													
2001	86	-	-	-	-	-	1	-	9	14	40	22	-
2000	81	-	-	-	-	-	-	1	5	19	36	20	-
1999	65	-	-	-	-	-	-	1	6	17	24	17	-
Male													
2001	86	-	-	-	-	-	1	-	9	14	40	22	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of prostate (C61)													
Origin not stated [2]													
Male													
2000	81	-	-	-	-	-	-	1	5	19	36	20	-
1999	65	-	-	-	-	-	-	1	6	17	24	17	-
Female													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Malignant neoplasms of kidney and renal pelvis (C64-C65)													
All origins													
Both sexes													
2001	12,078	1	22	35	20	52	358	1,311	2,328	3,221	3,306	1,424	-
2000	11,736	2	15	29	20	42	334	1,239	2,195	3,146	3,282	1,432	-
1999	11,116	2	19	33	17	54	342	1,147	2,111	3,076	3,010	1,305	-
Male													
2001	7,524	1	10	9	16	27	240	941	1,634	2,093	1,866	687	-
2000	7,249	1	8	15	10	26	228	880	1,530	2,010	1,865	676	-
1999	6,867	1	11	16	11	37	236	816	1,474	1,958	1,717	590	-
Female													
2001	4,554	-	12	26	4	25	118	370	694	1,128	1,440	737	-
2000	4,487	1	7	14	10	16	106	359	665	1,136	1,417	756	-
1999	4,249	1	8	17	6	17	106	331	637	1,118	1,293	715	-
Hispanic													
Both sexes													
2001	640	-	6	6	-	8	38	112	127	163	129	51	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of kidney and renal pelvis (C64-C65)													
Hispanic													
Both sexes													
2000	567	-	3	3	3	7	32	88	135	132	125	39	-
1999	547	-	1	4	1	4	31	84	129	144	108	41	-
Male													
2001	426	-	1	4	-	5	34	80	86	107	79	30	-
2000	372	-	1	3	2	4	27	60	92	85	78	20	-
1999	353	-	1	2	-	2	23	63	91	96	57	18	-
Female													
2001	214	-	5	2	-	3	4	32	41	56	50	21	-
2000	195	-	2	-	1	3	5	28	43	47	47	19	-
1999	194	-	-	2	1	2	8	21	38	48	51	23	-
Non-Hispanic [1]													
Both sexes													
2001	11,415	1	16	29	20	44	319	1,196	2,198	3,051	3,172	1,369	-
2000	11,124	2	12	26	17	35	302	1,147	2,053	2,994	3,150	1,386	-
1999	10,542	2	18	29	16	49	309	1,060	1,979	2,927	2,892	1,261	-
Male													
2001	7,084	1	9	5	16	22	205	859	1,546	1,981	1,784	656	-
2000	6,847	1	7	12	8	22	201	818	1,433	1,911	1,782	652	-
1999	6,501	1	10	14	11	35	212	751	1,381	1,858	1,657	571	-
Female													
2001	4,331	-	7	24	4	22	114	337	652	1,070	1,388	713	-
2000	4,277	1	5	14	9	13	101	329	620	1,083	1,368	734	-
1999	4,041	1	8	15	5	14	97	309	598	1,069	1,235	690	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of kidney and renal pelvis (C64-C65)													
Non-Hispanic white													
Both sexes													
2001	10,073	-	11	17	8	23	235	998	1,904	2,741	2,880	1,256	-
2000	9,884	-	7	16	11	24	239	977	1,794	2,665	2,871	1,280	-
1999	9,319	2	11	21	9	28	238	877	1,740	2,613	2,636	1,144	-
Male													
2001	6,263	-	6	1	5	11	157	726	1,340	1,790	1,616	611	-
2000	6,082	-	4	6	6	13	165	701	1,258	1,697	1,628	604	-
1999	5,754	1	7	9	7	18	165	618	1,220	1,662	1,528	519	-
Female													
2001	3,810	-	5	16	3	12	78	272	564	951	1,264	645	-
2000	3,802	-	3	10	5	11	74	276	536	968	1,243	676	-
1999	3,565	1	4	12	2	10	73	259	520	951	1,108	625	-
Non-Hispanic black													
Both sexes													
2001	1,105	-	3	12	10	19	72	163	240	249	241	96	-
2000	1,033	1	5	8	6	10	51	142	213	271	233	93	-
1999	1,028	-	4	6	7	20	61	154	202	263	213	98	-
Male													
2001	673	-	1	4	10	10	40	111	168	153	136	40	-
2000	625	-	3	4	2	8	27	94	142	179	124	42	-
1999	618	-	2	3	4	16	42	109	137	159	104	42	-
Female													
2001	432	-	2	8	-	9	32	52	72	96	105	56	-
2000	408	1	2	4	4	2	24	48	71	92	109	51	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of kidney and renal pelvis (C64-C65)													
Non-Hispanic black													
Female													
1999	410	-	2	3	3	4	19	45	65	104	109	56	-
Origin not stated [2]													
Both sexes													
2001	23	-	-	-	-	-	1	3	3	7	5	4	-
2000	45	-	-	-	-	-	-	4	7	20	7	7	-
1999	27	-	-	-	-	1	2	3	3	5	10	3	-
Male													
2001	14	-	-	-	-	-	1	2	2	5	3	1	-
2000	30	-	-	-	-	-	-	2	5	14	5	4	-
1999	13	-	-	-	-	-	1	2	2	4	3	1	-
Female													
2001	9	-	-	-	-	-	-	1	1	2	2	3	-
2000	15	-	-	-	-	-	-	2	2	6	2	3	-
1999	14	-	-	-	-	1	1	1	1	1	7	2	-
Malignant neoplasm of bladder (C67)													
All origins													
Both sexes													
2001	12,225	-	-	1	3	15	128	475	1,265	2,814	4,488	3,036	-
2000	12,002	-	-	1	3	10	111	440	1,223	2,855	4,407	2,952	-
1999	11,911	-	-	1	1	16	100	463	1,244	2,923	4,317	2,845	1
Male													
2001	8,343	-	-	-	3	8	70	351	956	2,088	3,116	1,751	-
2000	8,163	-	-	-	3	2	62	322	916	2,094	3,054	1,710	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of bladder (C67)													
All origins													
Male													
1999	8,105	-	-	1	-	9	63	350	933	2,132	3,008	1,608	1
Female													
2001	3,882	-	-	1	-	7	58	124	309	726	1,372	1,285	-
2000	3,839	-	-	1	-	8	49	118	307	761	1,353	1,242	-
1999	3,806	-	-	-	1	7	37	113	311	791	1,309	1,237	-
Hispanic													
Both sexes													
2001	336	-	-	-	2	-	4	21	41	90	104	74	-
2000	307	-	-	-	-	1	8	17	50	71	89	71	-
1999	301	-	-	-	-	2	4	22	51	88	79	55	-
Male													
2001	223	-	-	-	2	-	1	11	37	59	67	46	-
2000	216	-	-	-	-	-	6	12	37	50	69	42	-
1999	222	-	-	-	-	2	4	16	43	69	52	36	-
Female													
2001	113	-	-	-	-	-	3	10	4	31	37	28	-
2000	91	-	-	-	-	1	2	5	13	21	20	29	-
1999	79	-	-	-	-	-	-	6	8	19	27	19	-
Non-Hispanic [1]													
Both sexes													
2001	11,858	-	-	1	1	15	124	452	1,220	2,718	4,372	2,955	-
2000	11,665	-	-	1	3	9	102	423	1,169	2,780	4,304	2,874	-
1999	11,582	-	-	1	1	14	96	439	1,190	2,832	4,228	2,780	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of bladder (C67)													
Non-Hispanic [1]													
Male													
2001	8,100	-	-	-	1	8	69	339	917	2,025	3,041	1,700	-
2000	7,925	-	-	-	3	2	56	310	876	2,040	2,974	1,664	-
1999	7,865	-	-	1	-	7	59	332	888	2,060	2,950	1,567	1
Female													
2001	3,758	-	-	1	-	7	55	113	303	693	1,331	1,255	-
2000	3,740	-	-	1	-	7	46	113	293	740	1,330	1,210	-
1999	3,717	-	-	-	1	7	37	107	302	772	1,278	1,213	-
Non-Hispanic white													
Both sexes													
2001	10,823	-	-	1	1	14	100	386	1,076	2,479	4,026	2,740	-
2000	10,712	-	-	1	2	8	91	374	1,056	2,533	3,957	2,690	-
1999	10,548	-	-	1	-	8	77	375	1,042	2,556	3,921	2,567	1
Male													
2001	7,541	-	-	-	1	8	56	296	830	1,884	2,856	1,610	-
2000	7,390	-	-	-	2	2	52	279	803	1,897	2,784	1,571	-
1999	7,290	-	-	1	-	3	46	281	800	1,907	2,771	1,480	1
Female													
2001	3,282	-	-	1	-	6	44	90	246	595	1,170	1,130	-
2000	3,322	-	-	1	-	6	39	95	253	636	1,173	1,119	-
1999	3,258	-	-	-	-	5	31	94	242	649	1,150	1,087	-
Non-Hispanic black													
Both sexes													
2001	870	-	-	-	-	1	23	55	125	197	291	178	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasm of bladder (C67)													
Non-Hispanic black													
Both sexes													
2000	842	-	-	-	1	-	10	44	99	218	314	156	-
1999	906	-	-	-	1	4	17	60	130	244	262	188	-
Male													
2001	444	-	-	-	-	-	13	36	72	111	144	68	-
2000	460	-	-	-	1	-	4	27	63	119	171	75	-
1999	487	-	-	-	-	2	12	47	76	128	149	73	-
Female													
2001	426	-	-	-	-	1	10	19	53	86	147	110	-
2000	382	-	-	-	-	-	6	17	36	99	143	81	-
1999	419	-	-	-	1	2	5	13	54	116	113	115	-
Origin not stated [2]													
Both sexes													
2001	31	-	-	-	-	-	-	2	4	6	12	7	-
2000	30	-	-	-	-	-	1	-	4	4	14	7	-
1999	28	-	-	-	-	-	-	2	3	3	10	10	-
Male													
2001	20	-	-	-	-	-	-	1	2	4	8	5	-
2000	22	-	-	-	-	-	-	-	3	4	11	4	-
1999	18	-	-	-	-	-	-	2	2	3	6	5	-
Female													
2001	11	-	-	-	-	-	-	1	2	2	4	2	-
2000	8	-	-	-	-	-	1	-	1	-	3	3	-
1999	10	-	-	-	-	-	-	-	1	-	4	5	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)													
All origins													
Both sexes													
2001	12,609	16	100	300	224	416	1,028	1,951	2,464	2,823	2,566	721	-
2000	12,655	20	104	316	225	406	1,096	1,924	2,337	3,032	2,478	717	-
1999	12,765	10	113	308	218	444	1,065	1,880	2,453	3,119	2,492	663	-
Male													
2001	7,015	8	52	145	129	262	638	1,173	1,470	1,589	1,288	261	-
2000	7,011	10	58	176	132	264	704	1,148	1,379	1,678	1,187	275	-
1999	7,001	8	56	154	131	276	638	1,120	1,417	1,728	1,226	247	-
Female													
2001	5,594	8	48	155	95	154	390	778	994	1,234	1,278	460	-
2000	5,644	10	46	140	93	142	392	776	958	1,354	1,291	442	-
1999	5,764	2	57	154	87	168	427	760	1,036	1,391	1,266	416	-
Hispanic													
Both sexes													
2001	617	4	20	37	28	38	77	87	109	117	82	18	-
2000	631	4	16	63	28	45	76	92	106	105	72	24	-
1999	568	4	27	48	18	31	65	88	99	104	67	17	-
Male													
2001	335	1	10	16	19	21	41	50	59	72	40	6	-
2000	346	1	10	41	16	27	40	41	62	59	38	11	-
1999	293	4	11	22	10	17	45	42	60	48	31	3	-
Female													
2001	282	3	10	21	9	17	36	37	50	45	42	12	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)													
Hispanic													
Female													
2000	285	3	6	22	12	18	36	51	44	46	34	13	-
1999	275	-	16	26	8	14	20	46	39	56	36	14	-
Non-Hispanic [1]													
Both sexes													
2001	11,967	12	79	261	196	377	950	1,862	2,346	2,703	2,479	702	-
2000	11,995	16	87	253	197	361	1,019	1,828	2,222	2,923	2,398	691	-
1999	12,173	6	86	260	199	413	1,000	1,789	2,348	3,010	2,418	644	-
Male													
2001	6,667	7	41	128	110	241	596	1,122	1,406	1,516	1,245	255	-
2000	6,646	9	47	135	116	237	663	1,103	1,313	1,615	1,145	263	-
1999	6,696	4	45	132	121	259	593	1,076	1,353	1,678	1,191	244	-
Female													
2001	5,300	5	38	133	86	136	354	740	940	1,187	1,234	447	-
2000	5,349	7	40	118	81	124	356	725	909	1,308	1,253	428	-
1999	5,477	2	41	128	78	154	407	713	995	1,332	1,227	400	-
Non-Hispanic white													
Both sexes													
2001	11,004	8	63	205	155	335	864	1,695	2,155	2,509	2,356	659	-
2000	11,026	12	65	194	156	315	896	1,678	2,047	2,756	2,260	647	-
1999	11,238	3	70	195	152	372	902	1,635	2,171	2,839	2,293	606	-
Male													
2001	6,154	5	35	96	88	217	547	1,030	1,293	1,427	1,181	235	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)													
Non-Hispanic white													
Male													
2000	6,128	8	35	101	93	200	591	1,015	1,219	1,531	1,086	249	-
1999	6,193	2	39	104	91	233	533	980	1,252	1,591	1,136	232	-
Female													
2001	4,850	3	28	109	67	118	317	665	862	1,082	1,175	424	-
2000	4,898	4	30	93	63	115	305	663	828	1,225	1,174	398	-
1999	5,045	1	31	91	61	139	369	655	919	1,248	1,157	374	-
Non-Hispanic black													
Both sexes													
2001	736	1	10	44	29	28	63	133	145	153	92	38	-
2000	762	2	18	51	35	36	88	119	138	133	109	33	-
1999	754	3	11	54	37	36	76	128	142	132	104	31	-
Male													
2001	378	-	3	25	16	14	33	75	87	70	40	15	-
2000	407	1	10	30	21	28	52	71	76	66	43	9	-
1999	399	2	4	24	24	23	47	81	76	68	40	10	-
Female													
2001	358	1	7	19	13	14	30	58	58	83	52	23	-
2000	355	1	8	21	14	8	36	48	62	67	66	24	-
1999	355	1	7	30	13	13	29	47	66	64	64	21	-
Origin not stated [2]													
Both sexes													
2001	25	-	1	2	-	1	1	2	9	3	5	1	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)													
Origin not stated [2]													
Both sexes													
2000	29	-	1	-	-	-	1	4	9	4	8	2	-
1999	24	-	-	-	1	-	-	3	6	5	7	2	-
Male													
2001	13	-	1	1	-	-	1	1	5	1	3	-	-
2000	19	-	1	-	-	-	1	4	4	4	4	1	-
1999	12	-	-	-	-	-	-	2	4	2	4	-	-
Female													
2001	12	-	-	1	-	1	-	1	4	2	2	1	-
2000	10	-	-	-	-	-	-	-	5	-	4	1	-
1999	12	-	-	-	1	-	-	1	2	3	3	2	-
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)													
All origins													
Both sexes													
2001	56,341	27	157	367	705	988	1,857	4,064	7,455	13,659	18,257	8,805	-
2000	56,518	39	146	372	718	918	1,853	4,025	7,284	14,041	18,448	8,674	-
1999	56,318	30	137	351	720	960	1,913	3,950	7,554	14,332	18,089	8,282	-
Male													
2001	30,060	12	82	217	426	574	1,133	2,442	4,412	7,859	9,327	3,576	-
2000	29,888	21	76	223	455	539	1,075	2,399	4,205	8,038	9,399	3,458	-
1999	29,721	12	75	206	437	557	1,144	2,368	4,426	8,067	9,100	3,329	-
Female													
2001	26,281	15	75	150	279	414	724	1,622	3,043	5,800	8,930	5,229	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)													
All origins													
Female													
2000	26,630	18	70	149	263	379	778	1,626	3,079	6,003	9,049	5,216	-
1999	26,597	18	62	145	283	403	769	1,582	3,128	6,265	8,989	4,953	-
Hispanic													
Both sexes													
2001	2,747	7	38	87	144	154	211	279	442	597	573	215	-
2000	2,643	8	29	74	129	124	200	280	427	633	523	216	-
1999	2,538	8	28	75	128	135	190	299	396	606	485	188	-
Male													
2001	1,577	2	21	48	99	99	142	164	251	338	318	95	-
2000	1,460	5	14	47	83	83	115	172	247	338	258	98	-
1999	1,319	3	16	42	73	75	108	175	226	312	226	63	-
Female													
2001	1,170	5	17	39	45	55	69	115	191	259	255	120	-
2000	1,183	3	15	27	46	41	85	108	180	295	265	118	-
1999	1,219	5	12	33	55	60	82	124	170	294	259	125	-
Non-Hispanic [1]													
Both sexes													
2001	53,458	20	117	279	559	828	1,640	3,772	6,994	13,027	17,645	8,577	-
2000	53,749	31	116	295	586	793	1,645	3,735	6,833	13,379	17,893	8,443	-
1999	53,669	22	108	276	592	823	1,719	3,642	7,143	13,704	17,561	8,079	-
Male													
2001	28,410	10	60	169	326	473	987	2,269	4,151	7,505	8,985	3,475	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)													
Non-Hispanic [1]													
Male													
2000	28,358	16	61	175	371	455	956	2,219	3,941	7,682	9,127	3,355	-
1999	28,347	9	58	164	364	481	1,032	2,190	4,195	7,740	8,853	3,261	-
Female													
2001	25,048	10	57	110	233	355	653	1,503	2,843	5,522	8,660	5,102	-
2000	25,391	15	55	120	215	338	689	1,516	2,892	5,697	8,766	5,088	-
1999	25,322	13	50	112	228	342	687	1,452	2,948	5,964	8,708	4,818	-
Non-Hispanic white													
Both sexes													
2001	47,190	17	97	198	422	595	1,255	3,057	5,878	11,492	16,199	7,980	-
2000	47,695	30	92	232	420	585	1,283	3,017	5,812	11,911	16,528	7,785	-
1999	47,608	16	91	199	437	618	1,352	2,980	6,042	12,208	16,172	7,493	-
Male													
2001	25,162	8	47	111	252	352	748	1,873	3,519	6,693	8,307	3,252	-
2000	25,272	16	50	134	268	338	775	1,803	3,379	6,899	8,491	3,119	-
1999	25,219	9	48	120	277	357	816	1,812	3,591	6,967	8,202	3,020	-
Female													
2001	22,028	9	50	87	170	243	507	1,184	2,359	4,799	7,892	4,728	-
2000	22,423	14	42	98	152	247	508	1,214	2,433	5,012	8,037	4,666	-
1999	22,389	7	43	79	160	261	536	1,168	2,451	5,241	7,970	4,473	-
Non-Hispanic black													
Both sexes													
2001	5,162	3	13	62	107	181	322	584	915	1,288	1,183	504	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)													
Non-Hispanic black													
Both sexes													
2000	5,026	1	19	43	130	158	302	615	868	1,232	1,101	557	-
1999	5,038	4	14	61	133	157	307	554	917	1,247	1,142	502	-
Male													
2001	2,642	2	9	43	57	98	199	326	512	679	538	179	-
2000	2,535	-	8	26	79	90	149	346	481	653	510	193	-
1999	2,566	-	8	38	73	97	180	318	500	645	521	186	-
Female													
2001	2,520	1	4	19	50	83	123	258	403	609	645	325	-
2000	2,491	1	11	17	51	68	153	269	387	579	591	364	-
1999	2,472	4	6	23	60	60	127	236	417	602	621	316	-
Origin not stated [2]													
Both sexes													
2001	136	-	2	1	2	6	6	13	19	35	39	13	-
2000	126	-	1	3	3	1	8	10	24	29	32	15	-
1999	111	-	1	-	-	2	4	9	15	22	43	15	-
Male													
2001	73	-	1	-	1	2	4	9	10	16	24	6	-
2000	70	-	1	1	1	1	4	8	17	18	14	5	-
1999	55	-	1	-	-	1	4	3	5	15	21	5	-
Female													
2001	63	-	1	1	1	4	2	4	9	19	15	7	-
2000	56	-	-	2	2	-	4	2	7	11	18	10	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)													
Origin not stated [2]													
Female													
1999	56	-	-	-	-	1	-	6	10	7	22	10	-
Hodgkin's disease (C81)													
All origins													
Both sexes													
2001	1,323	-	-	6	78	167	161	182	143	211	278	97	-
2000	1,287	-	-	8	73	145	151	161	158	232	255	104	-
1999	1,403	-	1	9	94	148	165	165	178	283	262	98	-
Male													
2001	725	-	-	3	43	90	101	110	85	129	130	34	-
2000	686	-	-	4	35	79	82	99	87	134	121	45	-
1999	785	-	1	3	55	86	103	98	115	160	119	45	-
Female													
2001	598	-	-	3	35	77	60	72	58	82	148	63	-
2000	601	-	-	4	38	66	69	62	71	98	134	59	-
1999	618	-	-	6	39	62	62	67	63	123	143	53	-
Hispanic													
Both sexes													
2001	99	-	-	1	7	13	13	14	14	21	11	5	-
2000	95	-	-	-	9	14	8	13	13	11	23	4	-
1999	113	-	-	3	8	7	12	17	17	26	18	5	-
Male													
2001	66	-	-	1	4	8	8	11	11	15	5	3	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Hodgkin's disease (C81)													
Hispanic													
Male													
2000	63	-	-	-	7	7	6	11	8	10	12	2	-
1999	65	-	-	2	6	3	8	11	9	13	11	2	-
Female													
2001	33	-	-	-	3	5	5	3	3	6	6	2	-
2000	32	-	-	-	2	7	2	2	5	1	11	2	-
1999	48	-	-	1	2	4	4	6	8	13	7	3	-
Non-Hispanic [1]													
Both sexes													
2001	1,221	-	-	5	71	154	147	167	129	189	267	92	-
2000	1,189	-	-	8	64	131	143	148	143	221	231	100	-
1999	1,286	-	1	6	86	141	153	147	159	257	243	93	-
Male													
2001	657	-	-	2	39	82	92	98	74	114	125	31	-
2000	621	-	-	4	28	72	76	88	78	124	108	43	-
1999	719	-	1	1	49	83	95	87	105	147	108	43	-
Female													
2001	564	-	-	3	32	72	55	69	55	75	142	61	-
2000	568	-	-	4	36	59	67	60	65	97	123	57	-
1999	567	-	-	5	37	58	58	60	54	110	135	50	-
Non-Hispanic white													
Both sexes													
2001	1,069	-	-	4	55	113	124	146	110	178	249	90	-
2000	1,059	-	-	5	48	117	119	128	123	200	224	95	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Hodgkin's disease (C81)													
Non-Hispanic white													
Both sexes													
1999	1,149	-	1	5	67	116	134	129	144	234	235	84	-
Male													
2001	577	-	-	2	34	62	76	85	63	109	117	29	-
2000	551	-	-	2	23	63	64	76	65	112	105	41	-
1999	643	-	1	1	39	69	84	73	97	135	105	39	-
Female													
2001	492	-	-	2	21	51	48	61	47	69	132	61	-
2000	508	-	-	3	25	54	55	52	58	88	119	54	-
1999	506	-	-	4	28	47	50	56	47	99	130	45	-
Non-Hispanic black													
Both sexes													
2001	136	-	-	1	15	39	21	17	18	11	12	2	-
2000	109	-	-	2	14	12	24	17	17	15	5	3	-
1999	125	-	-	-	19	25	18	18	14	18	6	7	-
Male													
2001	72	-	-	-	4	20	15	11	10	5	5	2	-
2000	57	-	-	1	5	8	12	9	11	7	3	1	-
1999	74	-	-	-	10	14	11	14	8	12	3	2	-
Female													
2001	64	-	-	1	11	19	6	6	8	6	7	-	-
2000	52	-	-	1	9	4	12	8	6	8	2	2	-
1999	51	-	-	-	9	11	7	4	6	6	3	5	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Hodgkin's disease (C81)													
Origin not stated [2]													
Both sexes													
2001	3	-	-	-	-	-	1	1	-	1	-	-	-
2000	3	-	-	-	-	-	-	-	2	-	1	-	-
1999	4	-	-	-	-	-	-	1	2	-	1	-	-
Male													
2001	2	-	-	-	-	-	1	1	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	1	-	1	-	-
1999	1	-	-	-	-	-	-	-	1	-	-	-	-
Female													
2001	1	-	-	-	-	-	-	-	-	1	-	-	-
2000	1	-	-	-	-	-	-	-	1	-	-	-	-
1999	3	-	-	-	-	-	-	1	1	-	1	-	-
Non-Hodgkin's lymphoma (C82-C85)													
All origins													
Both sexes													
2001	22,305	1	1	38	148	298	756	1,732	3,098	5,358	7,369	3,506	-
2000	22,729	1	7	47	180	293	751	1,745	3,170	5,598	7,535	3,402	-
1999	22,802	2	16	37	144	289	789	1,741	3,154	5,848	7,554	3,228	-
Male													
2001	11,710	1	1	29	94	179	500	1,113	1,825	3,025	3,558	1,385	-
2000	11,812	-	5	28	126	172	482	1,112	1,849	3,105	3,634	1,299	-
1999	11,794	-	9	30	91	178	510	1,111	1,847	3,176	3,596	1,246	-
Female													
2001	10,595	-	-	9	54	119	256	619	1,273	2,333	3,811	2,121	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Non-Hodgkin's lymphoma (C82-C85)													
All origins													
Female													
2000	10,917	1	2	19	54	121	269	633	1,321	2,493	3,901	2,103	-
1999	11,008	2	7	7	53	111	279	630	1,307	2,672	3,958	1,982	-
Hispanic													
Both sexes													
2001	1,008	-	-	6	22	37	77	122	201	208	240	95	-
2000	1,026	-	2	2	21	38	83	113	190	277	210	90	-
1999	958	-	1	3	19	37	68	131	153	261	205	80	-
Male													
2001	579	-	-	6	15	24	56	74	115	122	131	36	-
2000	560	-	2	1	17	23	51	72	112	143	98	41	-
1999	493	-	-	2	10	18	43	84	91	125	92	28	-
Female													
2001	429	-	-	-	7	13	21	48	86	86	109	59	-
2000	466	-	-	1	4	15	32	41	78	134	112	49	-
1999	465	-	1	1	9	19	25	47	62	136	113	52	-
Non-Hispanic [1]													
Both sexes													
2001	21,246	1	1	32	125	258	678	1,605	2,892	5,134	7,113	3,407	-
2000	21,657	1	5	45	159	254	668	1,626	2,970	5,311	7,312	3,306	-
1999	21,792	2	15	34	125	252	718	1,606	2,996	5,575	7,327	3,142	-
Male													
2001	11,105	1	1	23	79	153	443	1,036	1,708	2,897	3,416	1,348	-
2000	11,225	-	3	27	109	148	431	1,036	1,728	2,954	3,533	1,256	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Non-Hodgkin's lymphoma (C82-C85)													
Non-Hispanic [1]													
Male													
1999	11,271	-	9	28	81	160	464	1,025	1,754	3,044	3,490	1,216	-
Female													
2001	10,141	-	-	9	46	105	235	569	1,184	2,237	3,697	2,059	-
2000	10,432	1	2	18	50	106	237	590	1,242	2,357	3,779	2,050	-
1999	10,521	2	6	6	44	92	254	581	1,242	2,531	3,837	1,926	-
Non-Hispanic white													
Both sexes													
2001	19,300	1	1	26	87	182	523	1,337	2,535	4,683	6,702	3,223	-
2000	19,843	-	2	38	110	181	531	1,367	2,655	4,878	6,938	3,143	-
1999	19,956	-	12	23	91	177	559	1,372	2,667	5,145	6,935	2,975	-
Male													
2001	10,052	1	1	17	54	115	339	873	1,497	2,655	3,226	1,274	-
2000	10,264	-	2	23	79	109	353	870	1,538	2,725	3,365	1,200	-
1999	10,259	-	8	21	61	112	362	879	1,554	2,822	3,302	1,138	-
Female													
2001	9,248	-	-	9	33	67	184	464	1,038	2,028	3,476	1,949	-
2000	9,579	-	-	15	31	72	178	497	1,117	2,153	3,573	1,943	-
1999	9,697	-	4	2	30	65	197	493	1,113	2,323	3,633	1,837	-
Non-Hispanic black													
Both sexes													
2001	1,478	-	-	5	32	62	129	221	269	348	284	128	-
2000	1,355	1	2	5	35	55	118	206	256	318	244	115	-
1999	1,427	2	2	9	32	58	135	189	259	330	286	125	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Non-Hodgkin's lymphoma (C82-C85)													
Non-Hispanic black													
Male													
2001	795	-	-	5	22	32	88	135	151	190	124	48	-
2000	719	-	-	3	19	30	68	133	157	169	103	37	-
1999	778	-	1	6	18	39	86	120	153	174	131	50	-
Female													
2001	683	-	-	-	10	30	41	86	118	158	160	80	-
2000	636	1	2	2	16	25	50	73	99	149	141	78	-
1999	649	2	1	3	14	19	49	69	106	156	155	75	-
Origin not stated [2]													
Both sexes													
2001	51	-	-	-	1	3	1	5	5	16	16	4	-
2000	46	-	-	-	-	1	-	6	10	10	13	6	-
1999	52	-	-	-	-	-	3	4	5	12	22	6	-
Male													
2001	26	-	-	-	-	2	1	3	2	6	11	1	-
2000	27	-	-	-	-	1	-	4	9	8	3	2	-
1999	30	-	-	-	-	-	3	2	2	7	14	2	-
Female													
2001	25	-	-	-	1	1	-	2	3	10	5	3	-
2000	19	-	-	-	-	-	-	2	1	2	10	4	-
1999	22	-	-	-	-	-	-	2	3	5	8	4	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Leukemia (C91-C95)													
All origins													
Both sexes													
2001	21,451	25	155	323	479	513	779	1,475	2,535	4,886	6,696	3,585	-
2000	21,339	38	137	315	464	473	801	1,411	2,414	5,010	6,733	3,543	-
1999	21,014	27	119	305	482	500	799	1,370	2,516	5,012	6,460	3,424	-
Male													
2001	11,850	10	81	185	289	300	432	813	1,552	2,962	3,699	1,527	-
2000	11,777	21	69	190	293	283	423	791	1,452	3,090	3,695	1,470	-
1999	11,519	11	65	173	291	276	435	777	1,489	3,054	3,525	1,423	-
Female													
2001	9,601	15	74	138	190	213	347	662	983	1,924	2,997	2,058	-
2000	9,562	17	68	125	171	190	378	620	962	1,920	3,038	2,073	-
1999	9,495	16	54	132	191	224	364	593	1,027	1,958	2,935	2,001	-
Hispanic													
Both sexes													
2001	1,114	7	38	80	115	101	103	100	120	197	185	68	-
2000	1,038	8	27	72	99	71	97	110	133	182	165	74	-
1999	994	8	26	69	101	86	98	100	115	179	149	63	-
Male													
2001	640	2	21	41	80	66	65	56	73	98	102	36	-
2000	595	5	12	46	59	52	52	64	82	98	87	38	-
1999	534	3	16	38	57	49	48	52	66	107	75	23	-
Female													
2001	474	5	17	39	35	35	38	44	47	99	83	32	-
2000	443	3	15	26	40	19	45	46	51	84	78	36	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Leukemia (C91-C95)													
Hispanic													
Female													
1999	460	5	10	31	44	37	50	48	49	72	74	40	-
Non-Hispanic [1]													
Both sexes													
2001	20,285	18	115	242	363	409	673	1,371	2,406	4,677	6,501	3,510	-
2000	20,248	30	109	240	362	402	697	1,299	2,274	4,817	6,555	3,463	-
1999	19,981	19	92	236	381	412	700	1,266	2,395	4,826	6,298	3,356	-
Male													
2001	11,180	8	59	144	208	234	365	754	1,474	2,857	3,590	1,487	-
2000	11,155	16	56	143	233	231	367	725	1,366	2,987	3,601	1,430	-
1999	10,968	8	48	135	234	226	386	724	1,421	2,940	3,446	1,400	-
Female													
2001	9,105	10	56	98	155	175	308	617	932	1,820	2,911	2,023	-
2000	9,093	14	53	97	129	171	330	574	908	1,830	2,954	2,033	-
1999	9,013	11	44	101	147	186	314	542	974	1,886	2,852	1,956	-
Non-Hispanic white													
Both sexes													
2001	18,105	15	95	168	280	298	524	1,123	2,084	4,185	6,031	3,302	-
2000	18,128	30	88	187	262	284	546	1,059	1,952	4,372	6,121	3,227	-
1999	17,885	15	78	171	279	315	564	1,040	2,075	4,348	5,848	3,152	-
Male													
2001	10,018	6	46	92	164	173	277	632	1,289	2,579	3,349	1,411	-
2000	10,009	16	46	108	166	164	298	593	1,183	2,726	3,377	1,332	-
1999	9,853	8	39	98	177	170	313	593	1,246	2,673	3,220	1,316	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Leukemia (C91-C95)													
Non-Hispanic white													
Female													
2001	8,087	9	49	76	116	125	247	491	795	1,606	2,682	1,891	-
2000	8,119	14	42	79	96	120	248	466	769	1,646	2,744	1,895	-
1999	8,032	7	39	73	102	145	251	447	829	1,675	2,628	1,836	-
Non-Hispanic black													
Both sexes													
2001	1,741	3	13	56	60	75	118	187	248	408	390	183	-
2000	1,729	-	17	36	80	88	116	204	258	373	349	208	-
1999	1,694	2	12	52	82	67	104	174	252	405	371	173	-
Male													
2001	915	2	9	38	31	44	67	93	147	232	187	65	-
2000	923	-	8	22	54	50	52	107	149	216	181	84	-
1999	887	-	7	32	45	38	54	103	142	222	178	66	-
Female													
2001	826	1	4	18	29	31	51	94	101	176	203	118	-
2000	806	-	9	14	26	38	64	97	109	157	168	124	-
1999	807	2	5	20	37	29	50	71	110	183	193	107	-
Origin not stated [2]													
Both sexes													
2001	52	-	2	1	1	3	3	4	9	12	10	7	-
2000	53	-	1	3	3	-	7	2	7	11	13	6	-
1999	39	-	1	-	-	2	1	4	6	7	13	5	-
Male													
2001	30	-	1	-	1	-	2	3	5	7	7	4	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Leukemia (C91-C95)													
Origin not stated [2]													
Male													
2000	27	-	1	1	1	-	4	2	4	5	7	2	-
1999	17	-	1	-	-	1	1	1	2	7	4	-	-
Female													
2001	22	-	1	1	-	3	1	1	4	5	3	3	-
2000	26	-	-	2	2	-	3	-	3	6	6	4	-
1999	22	-	-	-	-	1	-	3	4	-	9	5	-
Multiple myeloma and immunoproliferative neoplasms (C88,C90)													
All origins													
Both sexes													
2001	11,172	-	-	-	-	9	157	671	1,671	3,181	3,886	1,597	-
2000	11,063	-	-	1	-	7	146	700	1,530	3,179	3,896	1,604	-
1999	11,016	-	-	-	-	22	156	665	1,694	3,172	3,788	1,519	-
Male													
2001	5,724	-	-	-	-	4	97	403	945	1,725	1,926	624	-
2000	5,553	-	-	-	-	5	85	391	808	1,697	1,932	635	-
1999	5,584	-	-	-	-	16	95	378	968	1,669	1,846	612	-
Female													
2001	5,448	-	-	-	-	5	60	268	726	1,456	1,960	973	-
2000	5,510	-	-	1	-	2	61	309	722	1,482	1,964	969	-
1999	5,432	-	-	-	-	6	61	287	726	1,503	1,942	907	-
Hispanic													
Both sexes													
2001	524	-	-	-	-	3	18	43	107	169	137	47	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Multiple myeloma and immunoproliferative neoplasms (C88,C90)													
Hispanic													
Both sexes													
2000	478	-	-	-	-	1	12	43	90	161	124	47	-
1999	468	-	-	-	-	4	12	51	111	139	111	40	-
Male													
2001	291	-	-	-	-	1	13	23	52	102	80	20	-
2000	239	-	-	-	-	1	6	24	44	87	61	16	-
1999	225	-	-	-	-	4	9	28	60	67	47	10	-
Female													
2001	233	-	-	-	-	2	5	20	55	67	57	27	-
2000	239	-	-	-	-	-	6	19	46	74	63	31	-
1999	243	-	-	-	-	-	3	23	51	72	64	30	-
Non-Hispanic [1]													
Both sexes													
2001	10,618	-	-	-	-	6	138	625	1,559	3,006	3,736	1,548	-
2000	10,561	-	-	1	-	6	133	655	1,435	3,010	3,767	1,554	-
1999	10,532	-	-	-	-	18	144	614	1,581	3,030	3,670	1,475	-
Male													
2001	5,418	-	-	-	-	3	84	378	890	1,620	1,840	603	-
2000	5,300	-	-	-	-	4	79	365	761	1,605	1,868	618	-
1999	5,352	-	-	-	-	12	86	350	908	1,601	1,796	599	-
Female													
2001	5,200	-	-	-	-	3	54	247	669	1,386	1,896	945	-
2000	5,261	-	-	1	-	2	54	290	674	1,405	1,899	936	-
1999	5,180	-	-	-	-	6	58	264	673	1,429	1,874	876	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Multiple myeloma and immunoproliferative neoplasms (C88,C90)													
Non-Hispanic white													
Both sexes													
2001	8,641	-	-	-	-	2	81	447	1,144	2,427	3,194	1,346	-
2000	8,582	-	-	1	-	3	84	458	1,072	2,442	3,220	1,302	-
1999	8,547	-	-	-	-	10	92	432	1,145	2,465	3,133	1,270	-
Male													
2001	4,472	-	-	-	-	2	54	280	666	1,334	1,603	533	-
2000	4,400	-	-	-	-	2	58	261	586	1,325	1,628	540	-
1999	4,430	-	-	-	-	6	56	264	688	1,329	1,563	524	-
Female													
2001	4,169	-	-	-	-	-	27	167	478	1,093	1,591	813	-
2000	4,182	-	-	1	-	1	26	197	486	1,117	1,592	762	-
1999	4,117	-	-	-	-	4	36	168	457	1,136	1,570	746	-
Non-Hispanic black													
Both sexes													
2001	1,795	-	-	-	-	4	53	159	377	519	493	190	-
2000	1,824	-	-	-	-	3	44	186	336	526	500	229	-
1999	1,786	-	-	-	-	7	49	171	391	494	478	196	-
Male													
2001	853	-	-	-	-	1	28	87	203	251	220	63	-
2000	829	-	-	-	-	2	17	95	163	261	222	69	-
1999	824	-	-	-	-	6	29	80	196	237	208	68	-
Female													
2001	942	-	-	-	-	3	25	72	174	268	273	127	-
2000	995	-	-	-	-	1	27	91	173	265	278	160	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Multiple myeloma and immunoproliferative neoplasms (C88,C90)													
Non-Hispanic black													
Female													
1999	962	-	-	-	-	1	20	91	195	257	270	128	-
Origin not stated [2]													
Both sexes													
2001	30	-	-	-	-	-	1	3	5	6	13	2	-
2000	24	-	-	-	-	-	1	2	5	8	5	3	-
1999	16	-	-	-	-	-	-	-	2	3	7	4	-
Male													
2001	15	-	-	-	-	-	-	2	3	3	6	1	-
2000	14	-	-	-	-	-	-	2	3	5	3	1	-
1999	7	-	-	-	-	-	-	-	-	1	3	3	-
Female													
2001	15	-	-	-	-	-	1	1	2	3	7	1	-
2000	10	-	-	-	-	-	1	-	2	3	2	2	-
1999	9	-	-	-	-	-	-	-	2	2	4	1	-
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)													
All origins													
Both sexes													
2001	90	1	1	-	-	1	4	4	8	23	28	20	-
2000	100	-	2	1	1	-	4	8	12	22	29	21	-
1999	83	1	1	-	-	1	4	9	12	17	25	13	-
Male													
2001	51	1	-	-	-	1	3	3	5	18	14	6	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)													
All origins													
Male													
2000	60	-	2	1	1	-	3	6	9	12	17	9	-
1999	39	1	-	-	-	1	1	4	7	8	14	3	-
Female													
2001	39	-	1	-	-	-	1	1	3	5	14	14	-
2000	40	-	-	-	-	-	1	2	3	10	12	12	-
1999	44	-	1	-	-	-	3	5	5	9	11	10	-
Hispanic													
Both sexes													
2001	2	-	-	-	-	-	-	-	-	2	-	-	-
2000	6	-	-	-	-	-	-	1	1	2	1	1	-
1999	5	-	1	-	-	1	-	-	-	1	2	-	-
Male													
2001	1	-	-	-	-	-	-	-	-	1	-	-	-
2000	3	-	-	-	-	-	-	1	1	-	-	1	-
1999	2	-	-	-	-	1	-	-	-	-	1	-	-
Female													
2001	1	-	-	-	-	-	-	-	-	1	-	-	-
2000	3	-	-	-	-	-	-	-	-	2	1	-	-
1999	3	-	1	-	-	-	-	-	-	1	1	-	-
Non-Hispanic [1]													
Both sexes													
2001	88	1	1	-	-	1	4	4	8	21	28	20	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)													
Non-Hispanic [1]													
Both sexes													
2000	94	-	2	1	1	-	4	7	11	20	28	20	-
1999	78	1	-	-	-	-	4	9	12	16	23	13	-
Male													
2001	50	1	-	-	-	1	3	3	5	17	14	6	-
2000	57	-	2	1	1	-	3	5	8	12	17	8	-
1999	37	1	-	-	-	-	1	4	7	8	13	3	-
Female													
2001	38	-	1	-	-	-	1	1	3	4	14	14	-
2000	37	-	-	-	-	-	1	2	3	8	11	12	-
1999	41	-	-	-	-	-	3	5	5	8	10	10	-
Non-Hispanic white													
Both sexes													
2001	75	1	1	-	-	-	3	4	5	19	23	19	-
2000	83	-	2	1	-	-	3	5	10	19	25	18	-
1999	71	1	-	-	-	-	3	7	11	16	21	12	-
Male													
2001	43	1	-	-	-	-	2	3	4	16	12	5	-
2000	48	-	2	1	-	-	2	3	7	11	16	6	-
1999	34	1	-	-	-	-	1	3	6	8	12	3	-
Female													
2001	32	-	1	-	-	-	1	1	1	3	11	14	-
2000	35	-	-	-	-	-	1	2	3	8	9	12	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)													
Non-Hispanic white													
Female													
1999	37	-	-	-	-	-	2	4	5	8	9	9	-
Non-Hispanic black													
Both sexes													
2001	12	-	-	-	-	1	1	-	3	2	4	1	-
2000	9	-	-	-	1	-	-	2	1	-	3	2	-
1999	6	-	-	-	-	-	1	2	1	-	1	1	-
Male													
2001	7	-	-	-	-	1	1	-	1	1	2	1	-
2000	7	-	-	-	1	-	-	2	1	-	1	2	-
1999	3	-	-	-	-	-	-	1	1	-	1	-	-
Female													
2001	5	-	-	-	-	-	-	-	2	1	2	-	-
2000	2	-	-	-	-	-	-	-	-	-	2	-	-
1999	3	-	-	-	-	-	1	1	-	-	-	1	-
Origin not stated [2]													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)													
Origin not stated [2]													
Male													
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49, C51-C52,C57-C60,C62-C63,C66,C68-C69, C73-C80,C97)													
All origins													
Both sexes													
2001	62,708	17	117	279	516	728	2,055	5,506	9,441	15,023	18,700	10,326	-
2000	63,060	24	129	260	511	701	2,090	5,307	9,293	15,716	18,860	10,165	4
1999	66,251	14	116	291	519	762	2,314	5,477	9,863	16,834	19,656	10,403	2
Male													
2001	31,663	7	73	153	346	442	1,108	3,164	5,326	8,177	9,032	3,835	-
2000	31,728	13	75	140	319	432	1,141	2,959	5,223	8,572	9,161	3,689	4
1999	33,449	8	64	160	329	490	1,300	3,098	5,597	9,157	9,439	3,806	1
Female													
2001	31,045	10	44	126	170	286	947	2,342	4,115	6,846	9,668	6,491	-
2000	31,332	11	54	120	192	269	949	2,348	4,070	7,144	9,699	6,476	-
1999	32,802	6	52	131	190	272	1,014	2,379	4,266	7,677	10,217	6,597	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49, C51-C52,C57-C60,C62-C63,C66,C68-C69, C73-C80,C97)													
Hispanic													
Both sexes													
2001	2,844	2	15	42	85	82	188	334	481	690	615	310	-
2000	2,753	3	24	47	82	100	141	290	455	690	631	290	-
1999	2,747	2	11	45	84	99	175	314	471	671	587	288	-
Male													
2001	1,455	-	9	22	64	56	95	181	262	339	315	112	-
2000	1,394	2	16	26	49	67	78	148	235	349	309	115	-
1999	1,399	2	3	23	62	63	90	162	256	351	278	109	-
Female													
2001	1,389	2	6	20	21	26	93	153	219	351	300	198	-
2000	1,359	1	8	21	33	33	63	142	220	341	322	175	-
1999	1,348	-	8	22	22	36	85	152	215	320	309	179	-
Non-Hispanic [1]													
Both sexes													
2001	59,685	15	102	237	431	645	1,858	5,155	8,935	14,278	18,042	9,987	-
2000	60,117	21	103	212	427	596	1,939	4,997	8,813	14,980	18,172	9,853	4
1999	63,328	12	105	245	434	655	2,130	5,149	9,361	16,121	19,020	10,094	2
Male													
2001	30,107	7	64	131	282	385	1,006	2,971	5,046	7,800	8,701	3,714	-
2000	30,237	11	57	114	269	362	1,059	2,803	4,972	8,195	8,823	3,568	4
1999	31,956	6	61	137	266	423	1,206	2,928	5,325	8,779	9,132	3,692	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49, C51-C52,C57-C60,C62-C63,C66,C68-C69, C73-C80,C97)													
Non-Hispanic [1]													
Female													
2001	29,578	8	38	106	149	260	852	2,184	3,889	6,478	9,341	6,273	-
2000	29,880	10	46	98	158	234	880	2,194	3,841	6,785	9,349	6,285	-
1999	31,372	6	44	108	168	232	924	2,221	4,036	7,342	9,888	6,402	1
Non-Hispanic white													
Both sexes													
2001	51,581	13	83	176	338	492	1,430	4,024	7,342	12,292	16,233	9,158	-
2000	52,223	16	75	168	328	479	1,452	3,908	7,330	13,019	16,395	9,050	3
1999	54,977	10	87	178	324	513	1,626	4,077	7,722	14,076	17,158	9,204	2
Male													
2001	26,007	6	50	96	220	296	792	2,326	4,139	6,789	7,882	3,411	-
2000	26,258	9	46	89	204	303	797	2,213	4,111	7,208	7,989	3,286	3
1999	27,780	5	52	104	190	338	943	2,305	4,423	7,714	8,322	3,383	1
Female													
2001	25,574	7	33	80	118	196	638	1,698	3,203	5,503	8,351	5,747	-
2000	25,965	7	29	79	124	176	655	1,695	3,219	5,811	8,406	5,764	-
1999	27,197	5	35	74	134	175	683	1,772	3,299	6,362	8,836	5,821	1
Non-Hispanic black													
Both sexes													
2001	6,725	1	16	46	76	124	339	963	1,333	1,659	1,468	700	-
2000	6,617	4	22	39	83	87	427	927	1,255	1,636	1,472	664	1
1999	7,009	1	14	49	83	106	421	928	1,407	1,704	1,539	757	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49, C51-C52,C57-C60,C62-C63,C66,C68-C69, C73-C80,C97)													
Non-Hispanic black													
Male													
2001	3,392	-	13	26	51	69	166	543	759	866	650	249	-
2000	3,337	1	9	24	56	46	227	510	733	827	687	216	1
1999	3,514	-	8	22	56	60	219	539	801	895	667	247	-
Female													
2001	3,333	1	3	20	25	55	173	420	574	793	818	451	-
2000	3,280	3	13	15	27	41	200	417	522	809	785	448	-
1999	3,495	1	6	27	27	46	202	389	606	809	872	510	-
Origin not stated [2]													
Both sexes													
2001	179	-	-	-	-	1	9	17	25	55	43	29	-
2000	190	-	2	1	2	5	10	20	25	46	57	22	-
1999	176	-	-	1	1	8	9	14	31	42	49	21	-
Male													
2001	101	-	-	-	-	1	7	12	18	38	16	9	-
2000	97	-	2	-	1	3	4	8	16	28	29	6	-
1999	94	-	-	-	1	4	4	8	16	27	29	5	-
Female													
2001	78	-	-	-	-	-	2	5	7	17	27	20	-
2000	93	-	-	1	1	2	6	12	9	18	28	16	-
1999	82	-	-	1	-	4	5	6	15	15	20	16	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)													
All origins													
Both sexes													
2001	13,676	51	58	105	102	151	366	719	1,253	2,574	4,654	3,643	-
2000	13,546	60	53	99	95	165	368	690	1,224	2,546	4,645	3,600	1
1999	13,227	58	63	101	98	171	342	662	1,133	2,684	4,472	3,443	-
Male													
2001	6,735	26	30	52	53	77	193	391	673	1,439	2,403	1,398	-
2000	6,559	32	33	49	62	82	188	385	657	1,415	2,319	1,336	1
1999	6,520	30	33	49	54	96	197	362	656	1,522	2,218	1,303	-
Female													
2001	6,941	25	28	53	49	74	173	328	580	1,135	2,251	2,245	-
2000	6,987	28	20	50	33	83	180	305	567	1,131	2,326	2,264	-
1999	6,707	28	30	52	44	75	145	300	477	1,162	2,254	2,140	-
Hispanic													
Both sexes													
2001	543	5	13	21	8	20	39	50	67	100	128	92	-
2000	521	6	7	13	10	25	37	43	53	98	123	106	-
1999	497	8	12	17	12	25	30	39	60	109	111	74	-
Male													
2001	276	2	7	12	6	6	24	30	35	50	67	37	-
2000	237	4	2	8	8	15	14	18	25	46	54	43	-
1999	233	5	4	8	8	14	21	26	30	47	47	23	-
Female													
2001	267	3	6	9	2	14	15	20	32	50	61	55	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)													
Hispanic													
Female													
2000	284	2	5	5	2	10	23	25	28	52	69	63	-
1999	264	3	8	9	4	11	9	13	30	62	64	51	-
Non-Hispanic [1]													
Both sexes													
2001	13,098	46	45	84	94	131	327	667	1,182	2,468	4,510	3,544	-
2000	12,990	53	46	85	85	140	330	645	1,167	2,445	4,511	3,482	1
1999	12,699	50	51	83	85	146	311	620	1,070	2,572	4,349	3,362	-
Male													
2001	6,442	24	23	40	47	71	169	360	636	1,385	2,328	1,359	-
2000	6,302	28	31	41	54	67	173	365	629	1,367	2,258	1,288	1
1999	6,276	25	29	41	46	82	176	334	623	1,474	2,168	1,278	-
Female													
2001	6,656	22	22	44	47	60	158	307	546	1,083	2,182	2,185	-
2000	6,688	25	15	44	31	73	157	280	538	1,078	2,253	2,194	-
1999	6,423	25	22	42	39	64	135	286	447	1,098	2,181	2,084	-
Non-Hispanic white													
Both sexes													
2001	11,701	32	36	56	66	106	252	561	1,006	2,185	4,150	3,251	-
2000	11,636	41	37	63	63	105	255	516	1,004	2,144	4,171	3,237	-
1999	11,335	34	36	66	58	99	239	494	899	2,276	4,001	3,133	-
Male													
2001	5,801	18	18	25	31	58	135	312	547	1,250	2,152	1,255	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)													
Non-Hispanic white													
Male													
2000	5,653	21	23	31	40	55	132	301	539	1,202	2,108	1,201	-
1999	5,653	17	18	32	30	53	143	276	536	1,326	2,013	1,209	-
Female													
2001	5,900	14	18	31	35	48	117	249	459	935	1,998	1,996	-
2000	5,983	20	14	32	23	50	123	215	465	942	2,063	2,036	-
1999	5,682	17	18	34	28	46	96	218	363	950	1,988	1,924	-
Non-Hispanic black													
Both sexes													
2001	1,143	11	5	26	21	18	61	89	147	236	296	233	-
2000	1,122	9	9	18	16	33	62	112	133	245	282	202	1
1999	1,161	14	13	17	25	30	65	104	148	256	295	194	-
Male													
2001	522	5	3	13	13	11	30	38	73	119	141	76	-
2000	514	4	8	8	9	11	32	54	74	130	118	65	1
1999	516	7	10	9	15	20	30	44	75	128	124	54	-
Female													
2001	621	6	2	13	8	7	31	51	74	117	155	157	-
2000	608	5	1	10	7	22	30	58	59	115	164	137	-
1999	645	7	3	8	10	10	35	60	73	128	171	140	-
Origin not stated [2]													
Both sexes													
2001	35	-	-	-	-	-	-	2	4	6	16	7	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)													
Origin not stated [2]													
Both sexes													
2000	35	1	-	1	-	-	1	2	4	3	11	12	-
1999	31	-	-	1	1	-	1	3	3	3	12	7	-
Male													
2001	17	-	-	-	-	-	-	1	2	4	8	2	-
2000	20	-	-	-	-	-	1	2	3	2	7	5	-
1999	11	-	-	-	-	-	-	2	3	1	3	2	-
Female													
2001	18	-	-	-	-	-	-	1	2	2	8	5	-
2000	15	1	-	1	-	-	-	-	1	1	4	7	-
1999	20	-	-	1	1	-	1	1	-	2	9	5	-
Anemias (D50-D64)													
All origins													
Both sexes													
2001	4,627	16	22	55	109	160	215	232	249	492	1,226	1,851	-
2000	4,494	10	35	52	85	153	190	227	260	515	1,171	1,796	-
1999	4,503	12	30	53	121	160	194	232	231	486	1,164	1,820	-
Male													
2001	1,774	11	12	27	59	95	91	115	111	237	496	520	-
2000	1,669	6	16	29	39	70	78	108	130	236	465	492	-
1999	1,746	8	17	23	67	81	101	116	105	226	484	518	-
Female													
2001	2,853	5	10	28	50	65	124	117	138	255	730	1,331	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Anemias (D50-D64)													
All origins													
Female													
2000	2,825	4	19	23	46	83	112	119	130	279	706	1,304	-
1999	2,757	4	13	30	54	79	93	116	126	260	680	1,302	-
Hispanic													
Both sexes													
2001	185	-	3	5	11	12	12	11	17	25	43	46	-
2000	144	1	1	7	5	13	8	10	13	24	36	26	-
1999	164	-	6	3	6	8	10	9	10	27	35	50	-
Male													
2001	82	-	2	5	6	8	6	4	7	12	21	11	-
2000	55	1	-	3	1	5	1	4	4	12	15	9	-
1999	79	-	4	1	3	5	6	4	3	17	17	19	-
Female													
2001	103	-	1	-	5	4	6	7	10	13	22	35	-
2000	89	-	1	4	4	8	7	6	9	12	21	17	-
1999	85	-	2	2	3	3	4	5	7	10	18	31	-
Non-Hispanic [1]													
Both sexes													
2001	4,424	15	19	50	97	146	202	219	231	466	1,177	1,802	-
2000	4,336	8	34	44	79	140	181	215	246	489	1,132	1,768	-
1999	4,321	12	24	50	115	152	183	221	220	455	1,125	1,764	-
Male													
2001	1,687	10	10	22	53	86	84	110	104	225	474	509	-
2000	1,608	4	16	26	37	65	77	104	126	222	448	483	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Anemias (D50-D64)													
Non-Hispanic [1]													
Male													
1999	1,659	8	13	22	64	76	95	111	101	207	465	497	-
Female													
2001	2,737	5	9	28	44	60	118	109	127	241	703	1,293	-
2000	2,728	4	18	18	42	75	104	111	120	267	684	1,285	-
1999	2,662	4	11	28	51	76	88	110	119	248	660	1,267	-
Non-Hispanic white													
Both sexes													
2001	3,350	5	8	22	22	27	41	81	150	368	1,013	1,613	-
2000	3,299	3	14	14	16	16	36	82	155	384	994	1,585	-
1999	3,222	6	4	17	28	10	37	86	112	360	995	1,567	-
Male													
2001	1,220	4	3	11	12	12	15	37	68	175	419	464	-
2000	1,178	1	8	9	7	10	20	39	83	185	392	424	-
1999	1,162	4	2	6	11	2	18	46	45	163	416	449	-
Female													
2001	2,130	1	5	11	10	15	26	44	82	193	594	1,149	-
2000	2,121	2	6	5	9	6	16	43	72	199	602	1,161	-
1999	2,060	2	2	11	17	8	19	40	67	197	579	1,118	-
Non-Hispanic black													
Both sexes													
2001	993	6	11	26	72	117	154	135	75	90	149	158	-
2000	961	4	19	30	61	124	139	128	83	89	121	163	-
1999	1,010	3	20	33	84	139	142	125	100	81	119	164	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Anemias (D50-D64)													
Non-Hispanic black													
Male													
2001	443	4	7	10	41	73	66	73	35	46	51	37	-
2000	392	3	8	17	29	55	55	64	37	31	45	48	-
1999	453	2	11	16	50	72	73	61	53	37	41	37	-
Female													
2001	550	2	4	16	31	44	88	62	40	44	98	121	-
2000	569	1	11	13	32	69	84	64	46	58	76	115	-
1999	557	1	9	17	34	67	69	64	47	44	78	127	-
Origin not stated [2]													
Both sexes													
2001	18	1	-	-	1	2	1	2	1	1	6	3	-
2000	14	1	-	1	1	-	1	2	1	2	3	2	-
1999	18	-	-	-	-	-	1	2	1	4	4	6	-
Male													
2001	5	1	-	-	-	1	1	1	-	-	1	-	-
2000	6	1	-	-	1	-	-	-	-	2	2	-	-
1999	8	-	-	-	-	-	-	1	1	2	2	2	-
Female													
2001	13	-	-	-	1	1	-	1	1	1	5	3	-
2000	8	-	-	1	-	-	1	2	1	-	1	2	-
1999	10	-	-	-	-	-	1	1	-	2	2	4	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Diabetes mellitus (E10-E14)													
All origins													
Both sexes													
2001	71,372	2	6	39	151	595	1,958	5,343	9,570	16,731	22,805	14,171	1
2000	69,301	4	6	26	162	623	1,926	4,954	9,186	16,674	22,184	13,556	-
1999	68,399	2	8	32	155	582	1,942	4,735	9,097	16,908	21,757	13,178	3
Male													
2001	32,841	1	2	23	85	341	1,181	3,154	5,241	8,437	9,918	4,457	1
2000	31,602	4	2	13	94	376	1,151	2,821	4,943	8,226	9,738	4,234	-
1999	31,150	-	4	16	80	319	1,139	2,791	4,856	8,279	9,539	4,125	2
Female													
2001	38,531	1	4	16	66	254	777	2,189	4,329	8,294	12,887	9,714	-
2000	37,699	-	4	13	68	247	775	2,133	4,243	8,448	12,446	9,322	-
1999	37,249	2	4	16	75	263	803	1,944	4,241	8,629	12,218	9,053	1
Hispanic													
Both sexes													
2001	5,663	1	1	2	19	49	167	507	982	1,604	1,548	783	-
2000	5,328	1	-	4	13	64	154	439	950	1,592	1,410	701	-
1999	5,182	-	-	4	14	43	157	413	927	1,551	1,369	704	-
Male													
2001	2,590	-	1	2	11	28	109	311	510	758	638	222	-
2000	2,507	1	-	2	7	48	95	240	520	762	609	223	-
1999	2,335	-	-	2	8	21	84	252	480	709	562	217	-
Female													
2001	3,073	1	-	-	8	21	58	196	472	846	910	561	-
2000	2,821	-	-	2	6	16	59	199	430	830	801	478	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Diabetes mellitus (E10-E14)													
Hispanic													
Female													
1999	2,847	-	-	2	6	22	73	161	447	842	807	487	-
Non-Hispanic [1]													
Both sexes													
2001	65,498	1	5	37	131	542	1,787	4,810	8,562	15,066	21,198	13,358	1
2000	63,769	3	6	22	149	557	1,762	4,501	8,207	15,031	20,717	12,814	-
1999	63,017	2	8	28	140	534	1,781	4,305	8,138	15,318	20,325	12,436	2
Male													
2001	30,144	1	1	21	74	311	1,069	2,828	4,715	7,643	9,255	4,225	1
2000	28,994	3	2	11	87	328	1,048	2,574	4,408	7,436	9,098	3,999	-
1999	28,719	-	4	14	72	295	1,053	2,525	4,358	7,548	8,955	3,894	1
Female													
2001	35,354	-	4	16	57	231	718	1,982	3,847	7,423	11,943	9,133	-
2000	34,775	-	4	11	62	229	714	1,927	3,799	7,595	11,619	8,815	-
1999	34,298	2	4	14	68	239	728	1,780	3,780	7,770	11,370	8,542	1
Non-Hispanic white													
Both sexes													
2001	51,482	-	5	19	66	335	1,233	3,332	6,118	11,464	17,507	11,402	1
2000	50,163	2	6	13	92	357	1,260	3,078	5,923	11,475	17,040	10,917	-
1999	49,347	2	6	19	94	363	1,231	2,940	5,674	11,664	16,821	10,533	-
Male													
2001	24,302	-	1	13	42	184	767	2,014	3,524	6,142	7,874	3,740	1
2000	23,465	2	2	7	57	220	766	1,842	3,298	5,967	7,796	3,508	-
1999	23,177	-	3	9	52	203	747	1,801	3,200	6,066	7,722	3,374	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Diabetes mellitus (E10-E14)													
Non-Hispanic white													
Female													
2001	27,180	-	4	6	24	151	466	1,318	2,594	5,322	9,633	7,662	-
2000	26,698	-	4	6	35	137	494	1,236	2,625	5,508	9,244	7,409	-
1999	26,170	2	3	10	42	160	484	1,139	2,474	5,598	9,099	7,159	-
Non-Hispanic black													
Both sexes													
2001	12,167	1	-	15	56	198	506	1,306	2,133	3,110	3,132	1,710	-
2000	11,918	1	-	9	55	187	458	1,270	1,995	3,083	3,193	1,667	-
1999	11,830	-	2	9	43	160	502	1,219	2,136	3,120	2,995	1,642	2
Male													
2001	4,983	1	-	6	27	122	275	701	1,031	1,278	1,150	392	-
2000	4,720	1	-	4	28	98	252	648	952	1,248	1,100	389	-
1999	4,718	-	1	5	19	85	276	641	986	1,262	1,016	426	1
Female													
2001	7,184	-	-	9	29	76	231	605	1,102	1,832	1,982	1,318	-
2000	7,198	-	-	5	27	89	206	622	1,043	1,835	2,093	1,278	-
1999	7,112	-	1	4	24	75	226	578	1,150	1,858	1,979	1,216	1
Origin not stated [2]													
Both sexes													
2001	211	-	-	-	1	4	4	26	26	61	59	30	-
2000	204	-	-	-	-	2	10	14	29	51	57	41	-
1999	200	-	-	-	1	5	4	17	32	39	63	38	1
Male													
2001	107	-	-	-	-	2	3	15	16	36	25	10	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Diabetes mellitus (E10-E14)													
Origin not stated [2]													
Male													
2000	101	-	-	-	-	-	8	7	15	28	31	12	-
1999	96	-	-	-	-	3	2	14	18	22	22	14	1
Female													
2001	104	-	-	-	1	2	1	11	10	25	34	20	-
2000	103	-	-	-	-	2	2	7	14	23	26	29	-
1999	104	-	-	-	1	2	2	3	14	17	41	24	-
Nutritional deficiencies (E40-E64)													
All origins													
Both sexes													
2001	3,704	4	4	2	10	14	44	110	153	374	1,100	1,889	-
2000	4,242	14	2	5	10	18	56	99	173	460	1,285	2,120	-
1999	4,289	9	2	1	8	12	64	106	170	429	1,248	2,240	-
Male													
2001	1,233	4	1	2	7	4	21	56	81	174	426	457	-
2000	1,412	10	2	2	2	9	16	46	97	221	471	536	-
1999	1,486	4	2	1	5	4	28	52	98	221	494	577	-
Female													
2001	2,471	-	3	-	3	10	23	54	72	200	674	1,432	-
2000	2,830	4	-	3	8	9	40	53	76	239	814	1,584	-
1999	2,803	5	-	-	3	8	36	54	72	208	754	1,663	-
Hispanic													
Both sexes													
2001	136	-	1	1	-	-	7	3	5	12	45	62	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Nutritional deficiencies (E40-E64)													
Hispanic													
Both sexes													
2000	147	-	-	-	1	3	4	5	7	24	43	60	-
1999	138	-	-	-	-	-	4	6	7	17	31	73	-
Male													
2001	53	-	-	1	-	-	2	1	4	2	20	23	-
2000	57	-	-	-	-	2	1	3	3	13	17	18	-
1999	54	-	-	-	-	-	2	5	2	8	13	24	-
Female													
2001	83	-	1	-	-	-	5	2	1	10	25	39	-
2000	90	-	-	-	1	1	3	2	4	11	26	42	-
1999	84	-	-	-	-	-	2	1	5	9	18	49	-
Non-Hispanic [1]													
Both sexes													
2001	3,563	4	3	1	9	14	37	107	148	362	1,054	1,824	-
2000	4,080	14	2	5	9	15	52	94	166	435	1,233	2,055	-
1999	4,142	9	2	1	8	12	60	100	162	411	1,215	2,162	-
Male													
2001	1,178	4	1	1	6	4	19	55	77	172	405	434	-
2000	1,348	10	2	2	2	7	15	43	94	208	448	517	-
1999	1,428	4	2	1	5	4	26	47	95	213	479	552	-
Female													
2001	2,385	-	2	-	3	10	18	52	71	190	649	1,390	-
2000	2,732	4	-	3	7	8	37	51	72	227	785	1,538	-
1999	2,714	5	-	-	3	8	34	53	67	198	736	1,610	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Nutritional deficiencies (E40-E64)													
Non-Hispanic white													
Both sexes													
2001	3,015	3	3	-	7	10	22	70	113	291	878	1,618	-
2000	3,469	8	1	2	7	6	36	68	123	345	1,062	1,811	-
1999	3,486	5	1	-	5	9	45	70	110	307	1,018	1,916	-
Male													
2001	948	3	1	-	6	3	10	27	52	131	340	375	-
2000	1,112	6	1	2	2	2	10	30	62	159	382	456	-
1999	1,142	2	1	-	3	3	18	34	61	150	392	478	-
Female													
2001	2,067	-	2	-	1	7	12	43	61	160	538	1,243	-
2000	2,357	2	-	-	5	4	26	38	61	186	680	1,355	-
1999	2,344	3	-	-	2	6	27	36	49	157	626	1,438	-
Non-Hispanic black													
Both sexes													
2001	472	1	-	1	2	4	15	33	29	61	149	177	-
2000	537	5	1	3	2	8	16	24	41	78	147	212	-
1999	585	4	1	1	3	3	15	27	47	90	175	219	-
Male													
2001	202	1	-	1	-	1	9	25	21	38	55	51	-
2000	214	3	1	-	-	5	5	13	30	44	58	55	-
1999	250	2	1	1	2	1	8	12	30	55	74	64	-
Female													
2001	270	-	-	-	2	3	6	8	8	23	94	126	-
2000	323	2	-	3	2	3	11	11	11	34	89	157	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Nutritional deficiencies (E40-E64)													
Non-Hispanic black													
Female													
1999	335	2	-	-	1	2	7	15	17	35	101	155	-
Origin not stated [2]													
Both sexes													
2001	5	-	-	-	1	-	-	-	-	-	1	3	-
2000	15	-	-	-	-	-	-	-	-	1	9	5	-
1999	9	-	-	-	-	-	-	-	1	1	2	5	-
Male													
2001	2	-	-	-	1	-	-	-	-	-	1	-	-
2000	7	-	-	-	-	-	-	-	-	-	6	1	-
1999	4	-	-	-	-	-	-	-	1	-	2	1	-
Female													
2001	3	-	-	-	-	-	-	-	-	-	-	3	-
2000	8	-	-	-	-	-	-	-	-	1	3	4	-
1999	5	-	-	-	-	-	-	-	-	1	-	4	-
Malnutrition (E40-E46)													
All origins													
Both sexes													
2001	3,454	2	-	2	9	13	39	99	146	347	1,025	1,772	-
2000	3,926	11	1	3	8	16	53	90	158	423	1,199	1,964	-
1999	3,985	7	2	1	7	11	61	99	154	407	1,159	2,077	-
Male													
2001	1,148	2	-	2	6	4	19	47	75	157	400	436	-
2000	1,302	8	1	1	2	8	15	41	86	207	434	499	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malnutrition (E40-E46)													
All origins													
Male													
1999	1,371	4	2	1	4	4	27	48	92	207	452	530	-
Female													
2001	2,306	-	-	-	3	9	20	52	71	190	625	1,336	-
2000	2,624	3	-	2	6	8	38	49	72	216	765	1,465	-
1999	2,614	3	-	-	3	7	34	51	62	200	707	1,547	-
Hispanic													
Both sexes													
2001	132	-	-	1	-	-	6	3	5	12	44	61	-
2000	137	-	-	-	1	2	4	4	7	22	39	58	-
1999	132	-	-	-	-	-	4	6	7	17	30	68	-
Male													
2001	52	-	-	1	-	-	2	1	4	2	20	22	-
2000	52	-	-	-	-	2	1	2	3	12	16	16	-
1999	50	-	-	-	-	-	2	5	2	8	12	21	-
Female													
2001	80	-	-	-	-	-	4	2	1	10	24	39	-
2000	85	-	-	-	1	-	3	2	4	10	23	42	-
1999	82	-	-	-	-	-	2	1	5	9	18	47	-
Non-Hispanic [1]													
Both sexes													
2001	3,317	2	-	1	8	13	33	96	141	335	980	1,708	-
2000	3,774	11	1	3	7	14	49	86	151	400	1,151	1,901	-
1999	3,844	7	2	1	7	11	57	93	146	389	1,127	2,004	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malnutrition (E40-E46)													
Non-Hispanic [1]													
Male													
2001	1,094	2	-	1	5	4	17	46	71	155	379	414	-
2000	1,243	8	1	1	2	6	14	39	83	195	412	482	-
1999	1,317	4	2	1	4	4	25	43	89	199	438	508	-
Female													
2001	2,223	-	-	-	3	9	16	50	70	180	601	1,294	-
2000	2,531	3	-	2	5	8	35	47	68	205	739	1,419	-
1999	2,527	3	-	-	3	7	32	50	57	190	689	1,496	-
Non-Hispanic white													
Both sexes													
2001	2,806	1	-	-	6	9	19	64	109	270	816	1,512	-
2000	3,204	7	-	1	6	6	33	62	112	312	993	1,672	-
1999	3,232	4	1	-	5	8	43	67	99	287	945	1,773	-
Male													
2001	881	1	-	-	5	3	8	23	49	119	316	357	-
2000	1,021	5	-	1	2	2	9	27	55	147	350	423	-
1999	1,047	2	1	-	3	3	17	31	58	138	359	435	-
Female													
2001	1,925	-	-	-	1	6	11	41	60	151	500	1,155	-
2000	2,183	2	-	-	4	4	24	35	57	165	643	1,249	-
1999	2,185	2	-	-	2	5	26	36	41	149	586	1,338	-
Non-Hispanic black													
Both sexes													
2001	438	1	-	1	2	4	14	28	26	56	139	167	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Malnutrition (E40-E46)													
Non-Hispanic black													
Both sexes													
2000	505	3	1	2	1	7	16	23	37	77	136	202	-
1999	543	3	1	1	2	3	14	24	42	88	161	204	-
Male													
2001	186	1	-	1	-	1	9	20	18	34	53	49	-
2000	202	2	1	-	-	4	5	12	26	43	56	53	-
1999	234	2	1	1	1	1	8	11	27	53	66	63	-
Female													
2001	252	-	-	-	2	3	5	8	8	22	86	118	-
2000	303	1	-	2	1	3	11	11	11	34	80	149	-
1999	309	1	-	-	1	2	6	13	15	35	95	141	-
Origin not stated [2]													
Both sexes													
2001	5	-	-	-	1	-	-	-	-	-	1	3	-
2000	15	-	-	-	-	-	-	-	-	1	9	5	-
1999	9	-	-	-	-	-	-	-	1	1	2	5	-
Male													
2001	2	-	-	-	1	-	-	-	-	-	1	-	-
2000	7	-	-	-	-	-	-	-	-	-	6	1	-
1999	4	-	-	-	-	-	-	-	1	-	2	1	-
Female													
2001	3	-	-	-	-	-	-	-	-	-	-	3	-
2000	8	-	-	-	-	-	-	-	-	1	3	4	-
1999	5	-	-	-	-	-	-	-	-	1	-	4	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other nutritional deficiencies (E50-E64)													
All origins													
Both sexes													
2001	250	2	4	-	1	1	5	11	7	27	75	117	-
2000	316	3	1	2	2	2	3	9	15	37	86	156	-
1999	304	2	-	-	1	1	3	7	16	22	89	163	-
Male													
2001	85	2	1	-	1	-	2	9	6	17	26	21	-
2000	110	2	1	1	-	1	1	5	11	14	37	37	-
1999	115	-	-	-	1	-	1	4	6	14	42	47	-
Female													
2001	165	-	3	-	-	1	3	2	1	10	49	96	-
2000	206	1	-	1	2	1	2	4	4	23	49	119	-
1999	189	2	-	-	-	1	2	3	10	8	47	116	-
Hispanic													
Both sexes													
2001	4	-	1	-	-	-	1	-	-	-	1	1	-
2000	10	-	-	-	-	1	-	1	-	2	4	2	-
1999	6	-	-	-	-	-	-	-	-	-	1	5	-
Male													
2001	1	-	-	-	-	-	-	-	-	-	-	1	-
2000	5	-	-	-	-	-	-	1	-	1	1	2	-
1999	4	-	-	-	-	-	-	-	-	-	1	3	-
Female													
2001	3	-	1	-	-	-	1	-	-	-	1	-	-
2000	5	-	-	-	-	1	-	-	-	1	3	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other nutritional deficiencies (E50-E64)													
Hispanic													
Female													
1999	2	-	-	-	-	-	-	-	-	-	-	2	-
Non-Hispanic [1]													
Both sexes													
2001	246	2	3	-	1	1	4	11	7	27	74	116	-
2000	306	3	1	2	2	1	3	8	15	35	82	154	-
1999	298	2	-	-	1	1	3	7	16	22	88	158	-
Male													
2001	84	2	1	-	1	-	2	9	6	17	26	20	-
2000	105	2	1	1	-	1	1	4	11	13	36	35	-
1999	111	-	-	-	1	-	1	4	6	14	41	44	-
Female													
2001	162	-	2	-	-	1	2	2	1	10	48	96	-
2000	201	1	-	1	2	-	2	4	4	22	46	119	-
1999	187	2	-	-	-	1	2	3	10	8	47	114	-
Non-Hispanic white													
Both sexes													
2001	209	2	3	-	1	1	3	6	4	21	62	106	-
2000	265	1	1	1	1	-	3	6	11	33	69	139	-
1999	254	1	-	-	-	1	2	3	11	20	73	143	-
Male													
2001	67	2	1	-	1	-	2	4	3	12	24	18	-
2000	91	1	1	1	-	-	1	3	7	12	32	33	-
1999	95	-	-	-	-	-	1	3	3	12	33	43	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other nutritional deficiencies (E50-E64)													
Non-Hispanic white													
Female													
2001	142	-	2	-	-	1	1	2	1	9	38	88	-
2000	174	-	-	-	1	-	2	3	4	21	37	106	-
1999	159	1	-	-	-	1	1	-	8	8	40	100	-
Non-Hispanic black													
Both sexes													
2001	34	-	-	-	-	-	1	5	3	5	10	10	-
2000	32	2	-	1	1	1	-	1	4	1	11	10	-
1999	42	1	-	-	1	-	1	3	5	2	14	15	-
Male													
2001	16	-	-	-	-	-	-	5	3	4	2	2	-
2000	12	1	-	-	-	1	-	1	4	1	2	2	-
1999	16	-	-	-	1	-	-	1	3	2	8	1	-
Female													
2001	18	-	-	-	-	-	1	-	-	1	8	8	-
2000	20	1	-	1	1	-	-	-	-	-	9	8	-
1999	26	1	-	-	-	-	1	2	2	-	6	14	-
Origin not stated [2]													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other nutritional deficiencies (E50-E64)													
Origin not stated [2]													
Male													
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Meningitis (G00,G03)													
All origins													
Both sexes													
2001	727	78	26	24	40	41	69	105	113	94	96	41	-
2000	758	91	36	30	15	43	72	100	90	108	123	50	-
1999	850	116	45	27	41	48	76	108	90	115	122	62	-
Male													
2001	394	39	16	15	23	25	42	60	61	51	47	15	-
2000	405	48	20	16	4	20	46	56	54	60	62	19	-
1999	442	61	26	18	26	24	45	68	45	60	49	20	-
Female													
2001	333	39	10	9	17	16	27	45	52	43	49	26	-
2000	353	43	16	14	11	23	26	44	36	48	61	31	-
1999	408	55	19	9	15	24	31	40	45	55	73	42	-
Hispanic													
Both sexes													
2001	58	8	4	3	3	4	7	6	7	6	8	2	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Meningitis (G00,G03)													
Hispanic													
Both sexes													
2000	52	9	4	3	3	5	8	6	5	5	4	-	-
1999	76	17	7	1	5	6	4	12	7	9	4	4	-
Male													
2001	33	5	3	2	1	1	6	3	3	5	4	-	-
2000	32	6	2	1	2	3	5	5	3	1	4	-	-
1999	41	13	5	1	1	3	3	6	3	5	-	1	-
Female													
2001	25	3	1	1	2	3	1	3	4	1	4	2	-
2000	20	3	2	2	1	2	3	1	2	4	-	-	-
1999	35	4	2	-	4	3	1	6	4	4	4	3	-
Non-Hispanic [1]													
Both sexes													
2001	666	70	22	21	37	37	62	98	106	88	88	37	-
2000	705	82	32	27	12	37	64	94	85	103	119	50	-
1999	767	97	38	25	34	42	71	96	82	106	118	58	-
Male													
2001	361	34	13	13	22	24	36	57	58	46	43	15	-
2000	372	42	18	15	2	16	41	51	51	59	58	19	-
1999	396	48	21	16	23	21	41	62	41	55	49	19	-
Female													
2001	305	36	9	8	15	13	26	41	48	42	45	22	-
2000	333	40	14	12	10	21	23	43	34	44	61	31	-
1999	371	49	17	9	11	21	30	34	41	51	69	39	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Meningitis (G00,G03)													
Non-Hispanic white													
Both sexes													
2001	469	39	15	15	24	25	33	66	77	73	74	28	-
2000	486	47	19	18	7	20	36	65	60	74	100	40	-
1999	524	51	26	18	15	28	45	60	63	80	92	46	-
Male													
2001	243	17	10	8	12	17	18	38	42	37	35	9	-
2000	264	27	12	9	1	7	22	38	39	47	48	14	-
1999	265	33	15	11	10	10	24	35	33	41	38	15	-
Female													
2001	226	22	5	7	12	8	15	28	35	36	39	19	-
2000	222	20	7	9	6	13	14	27	21	27	52	26	-
1999	259	18	11	7	5	18	21	25	30	39	54	31	-
Non-Hispanic black													
Both sexes													
2001	172	29	5	4	10	11	26	30	25	13	11	8	-
2000	202	34	13	9	5	17	27	26	22	27	13	9	-
1999	217	40	11	7	17	11	21	35	18	21	24	12	-
Male													
2001	104	16	2	3	8	6	17	18	14	8	6	6	-
2000	100	15	6	6	1	9	18	11	11	11	7	5	-
1999	119	13	6	5	11	9	14	27	8	12	10	4	-
Female													
2001	68	13	3	1	2	5	9	12	11	5	5	2	-
2000	102	19	7	3	4	8	9	15	11	16	6	4	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Meningitis (G00,G03)													
Non-Hispanic black													
Female													
1999	98	27	5	2	6	2	7	8	10	9	14	8	-
Origin not stated [2]													
Both sexes													
2001	3	-	-	-	-	-	-	1	-	-	-	2	-
2000	1	-	-	-	-	1	-	-	-	-	-	-	-
1999	7	2	-	1	2	-	1	-	1	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	1	-	-	-	-	-	-	-
1999	5	-	-	1	2	-	1	-	1	-	-	-	-
Female													
2001	3	-	-	-	-	-	-	1	-	-	-	2	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	2	-	-	-	-	-	-	-	-	-	-	-
Parkinson's disease (G20-G21)													
All origins													
Both sexes													
2001	16,544	-	-	-	-	2	15	45	301	2,147	8,125	5,909	-
2000	15,682	-	1	-	-	4	7	49	270	2,108	7,652	5,590	1
1999	14,593	1	-	-	2	2	5	43	242	2,020	7,110	5,168	-
Male													
2001	9,405	-	-	-	-	1	11	33	214	1,394	4,837	2,915	-
2000	8,804	-	-	-	-	3	5	29	170	1,338	4,530	2,728	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Parkinson's disease (G20-G21)													
All origins													
Male													
1999	8,262	-	-	-	2	2	3	30	161	1,303	4,307	2,454	-
Female													
2001	7,139	-	-	-	-	1	4	12	87	753	3,288	2,994	-
2000	6,878	-	1	-	-	1	2	20	100	770	3,122	2,862	-
1999	6,331	1	-	-	-	-	2	13	81	717	2,803	2,714	-
Hispanic													
Both sexes													
2001	483	-	-	-	-	1	-	3	13	98	230	138	-
2000	425	-	-	-	-	-	-	5	15	84	191	130	-
1999	389	-	-	-	-	-	1	2	8	84	184	110	-
Male													
2001	274	-	-	-	-	1	-	2	7	56	141	67	-
2000	240	-	-	-	-	-	-	2	9	46	112	71	-
1999	222	-	-	-	-	-	1	2	5	52	113	49	-
Female													
2001	209	-	-	-	-	-	-	1	6	42	89	71	-
2000	185	-	-	-	-	-	-	3	6	38	79	59	-
1999	167	-	-	-	-	-	-	-	3	32	71	61	-
Non-Hispanic [1]													
Both sexes													
2001	16,019	-	-	-	-	1	15	41	288	2,046	7,873	5,755	-
2000	15,223	-	1	-	-	4	7	43	255	2,020	7,442	5,450	1
1999	14,177	1	-	-	2	2	4	41	234	1,930	6,917	5,046	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Parkinson's disease (G20-G21)													
Non-Hispanic [1]													
Male													
2001	9,104	-	-	-	-	-	11	30	207	1,335	4,681	2,840	-
2000	8,544	-	-	-	-	3	5	26	161	1,290	4,407	2,651	1
1999	8,029	-	-	-	2	2	2	28	156	1,248	4,189	2,402	-
Female													
2001	6,915	-	-	-	-	1	4	11	81	711	3,192	2,915	-
2000	6,679	-	1	-	-	1	2	17	94	730	3,035	2,799	-
1999	6,148	1	-	-	-	-	2	13	78	682	2,728	2,644	-
Non-Hispanic white													
Both sexes													
2001	15,239	-	-	-	-	1	11	36	257	1,911	7,521	5,502	-
2000	14,497	-	-	-	-	3	4	36	236	1,895	7,106	5,216	1
1999	13,475	1	-	-	2	1	4	37	207	1,807	6,594	4,822	-
Male													
2001	8,685	-	-	-	-	-	8	27	190	1,252	4,486	2,722	-
2000	8,116	-	-	-	-	3	3	22	149	1,214	4,191	2,533	1
1999	7,648	-	-	-	2	1	2	27	142	1,182	4,005	2,287	-
Female													
2001	6,554	-	-	-	-	1	3	9	67	659	3,035	2,780	-
2000	6,381	-	-	-	-	-	1	14	87	681	2,915	2,683	-
1999	5,827	1	-	-	-	-	2	10	65	625	2,589	2,535	-
Non-Hispanic black													
Both sexes													
2001	547	-	-	-	-	-	4	5	20	91	238	189	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Parkinson's disease (G20-G21)													
Non-Hispanic black													
Both sexes													
2000	528	-	-	-	-	1	3	6	14	93	246	165	-
1999	502	-	-	-	-	-	-	4	22	95	221	160	-
Male													
2001	291	-	-	-	-	-	3	3	13	59	127	86	-
2000	304	-	-	-	-	-	2	4	9	60	154	75	-
1999	269	-	-	-	-	-	-	1	13	58	123	74	-
Female													
2001	256	-	-	-	-	-	1	2	7	32	111	103	-
2000	224	-	-	-	-	1	1	2	5	33	92	90	-
1999	233	-	-	-	-	-	-	3	9	37	98	86	-
Origin not stated [2]													
Both sexes													
2001	42	-	-	-	-	-	-	1	-	3	22	16	-
2000	34	-	-	-	-	-	-	1	-	4	19	10	-
1999	27	-	-	-	-	-	-	-	-	6	9	12	-
Male													
2001	27	-	-	-	-	-	-	1	-	3	15	8	-
2000	20	-	-	-	-	-	-	1	-	2	11	6	-
1999	11	-	-	-	-	-	-	-	-	3	5	3	-
Female													
2001	15	-	-	-	-	-	-	-	-	-	7	8	-
2000	14	-	-	-	-	-	-	-	-	2	8	4	-
1999	16	-	-	-	-	-	-	-	-	3	4	9	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Alzheimer's disease (G30)													
All origins													
Both sexes													
2001	53,852	-	-	-	-	2	3	73	529	3,422	18,542	31,281	-
2000	49,558	-	-	-	-	2	6	64	491	3,431	17,253	28,309	2
1999	44,536	-	-	-	1	-	7	60	448	3,204	15,836	24,980	-
Male													
2001	15,762	-	-	-	-	2	3	38	228	1,472	6,795	7,224	-
2000	14,438	-	-	-	-	1	4	34	210	1,554	6,228	6,406	1
1999	13,391	-	-	-	1	-	3	28	222	1,437	5,904	5,796	-
Female													
2001	38,090	-	-	-	-	-	-	35	301	1,950	11,747	24,057	-
2000	35,120	-	-	-	-	1	2	30	281	1,877	11,025	21,903	1
1999	31,145	-	-	-	-	-	4	32	226	1,767	9,932	19,184	-
Hispanic													
Both sexes													
2001	1,239	-	-	-	-	-	1	3	19	112	465	639	-
2000	1,127	-	-	-	-	-	-	6	19	128	401	573	-
1999	981	-	-	-	1	-	-	3	14	100	362	501	-
Male													
2001	401	-	-	-	-	-	1	3	6	44	186	161	-
2000	383	-	-	-	-	-	-	4	9	63	166	141	-
1999	314	-	-	-	1	-	-	-	5	51	128	129	-
Female													
2001	838	-	-	-	-	-	-	-	13	68	279	478	-
2000	744	-	-	-	-	-	-	2	10	65	235	432	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Alzheimer's disease (G30)													
Hispanic													
Female													
1999	667	-	-	-	-	-	-	3	9	49	234	372	-
Non-Hispanic [1]													
Both sexes													
2001	52,516	-	-	-	-	2	2	70	509	3,305	18,051	30,577	-
2000	48,331	-	-	-	-	2	6	58	469	3,296	16,829	27,670	1
1999	43,471	-	-	-	-	-	7	57	434	3,099	15,447	24,427	-
Male													
2001	15,334	-	-	-	-	2	2	35	222	1,427	6,600	7,046	-
2000	14,034	-	-	-	-	1	4	30	200	1,489	6,058	6,252	-
1999	13,052	-	-	-	-	-	3	28	217	1,384	5,768	5,652	-
Female													
2001	37,182	-	-	-	-	-	-	35	287	1,878	11,451	23,531	-
2000	34,297	-	-	-	-	1	2	28	269	1,807	10,771	21,418	1
1999	30,419	-	-	-	-	-	4	29	217	1,715	9,679	18,775	-
Non-Hispanic white													
Both sexes													
2001	49,030	-	-	-	-	2	2	64	480	3,020	16,742	28,720	-
2000	45,246	-	-	-	-	2	6	50	428	3,004	15,704	26,052	-
1999	40,835	-	-	-	-	-	5	51	399	2,821	14,453	23,106	-
Male													
2001	14,295	-	-	-	-	2	2	31	205	1,305	6,137	6,613	-
2000	13,123	-	-	-	-	1	4	25	183	1,367	5,686	5,857	-
1999	12,253	-	-	-	-	-	2	26	200	1,263	5,432	5,330	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Alzheimer's disease (G30)													
Non-Hispanic white													
Female													
2001	34,735	-	-	-	-	-	-	33	275	1,715	10,605	22,107	-
2000	32,123	-	-	-	-	1	2	25	245	1,637	10,018	20,195	-
1999	28,582	-	-	-	-	-	3	25	199	1,558	9,021	17,776	-
Non-Hispanic black													
Both sexes													
2001	3,104	-	-	-	-	-	-	5	26	253	1,176	1,644	-
2000	2,719	-	-	-	-	-	-	5	30	261	1,004	1,418	1
1999	2,325	-	-	-	-	-	1	4	29	249	883	1,159	-
Male													
2001	906	-	-	-	-	-	-	3	17	110	412	364	-
2000	774	-	-	-	-	-	-	3	16	113	320	322	-
1999	689	-	-	-	-	-	-	2	15	112	290	270	-
Female													
2001	2,198	-	-	-	-	-	-	2	9	143	764	1,280	-
2000	1,945	-	-	-	-	-	-	2	14	148	684	1,096	1
1999	1,636	-	-	-	-	-	1	2	14	137	593	889	-
Origin not stated [2]													
Both sexes													
2001	97	-	-	-	-	-	-	-	1	5	26	65	-
2000	100	-	-	-	-	-	-	-	3	7	23	66	1
1999	84	-	-	-	-	-	-	-	-	5	27	52	-
Male													
2001	27	-	-	-	-	-	-	-	-	1	9	17	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Alzheimer's disease (G30)													
Origin not stated [2]													
Male													
2000	21	-	-	-	-	-	-	-	1	2	4	13	1
1999	25	-	-	-	-	-	-	-	-	2	8	15	-
Female													
2001	70	-	-	-	-	-	-	-	1	4	17	48	-
2000	79	-	-	-	-	-	-	-	2	5	19	53	-
1999	59	-	-	-	-	-	-	-	-	3	19	37	-
Major cardiovascular diseases (I00-I78)													
All origins													
Both sexes													
2001	922,334	600	285	365	1,280	4,038	16,686	44,450	76,224	148,409	291,995	337,930	72
2000	936,923	636	234	362	1,309	3,792	16,624	43,583	77,482	156,187	296,202	340,450	62
1999	950,314	644	234	354	1,315	3,873	17,010	42,512	77,833	163,428	302,552	340,510	49
Male													
2001	428,065	315	149	184	807	2,555	11,180	30,860	50,320	87,756	140,789	103,097	53
2000	435,912	341	128	206	783	2,378	11,110	30,535	51,423	92,510	142,797	103,658	43
1999	441,827	347	127	199	799	2,442	11,471	29,745	51,525	96,904	145,567	102,663	38
Female													
2001	494,269	285	136	181	473	1,483	5,506	13,590	25,904	60,653	151,206	234,833	19
2000	501,011	295	106	156	526	1,414	5,514	13,048	26,059	63,677	153,405	236,792	19
1999	508,487	297	107	155	516	1,431	5,539	12,767	26,308	66,524	156,985	237,847	11
Hispanic													
Both sexes													
2001	35,565	98	44	56	162	442	1,081	2,561	3,996	7,325	10,139	9,654	7

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Major cardiovascular diseases (I00-I78)													
Hispanic													
Both sexes													
2000	34,059	106	37	47	169	419	1,094	2,351	4,041	7,114	9,380	9,296	5
1999	33,665	115	41	47	160	408	1,057	2,386	3,973	7,150	9,291	9,031	6
Male													
2001	18,138	53	21	29	108	308	758	1,806	2,582	4,269	4,966	3,232	6
2000	17,442	54	21	30	110	283	739	1,671	2,657	4,123	4,622	3,128	4
1999	17,304	49	21	29	103	292	755	1,700	2,560	4,212	4,519	3,058	6
Female													
2001	17,427	45	23	27	54	134	323	755	1,414	3,056	5,173	6,422	1
2000	16,617	52	16	17	59	136	355	680	1,384	2,991	4,758	6,168	1
1999	16,361	66	20	18	57	116	302	686	1,413	2,938	4,772	5,973	-
Non-Hispanic [1]													
Both sexes													
2001	883,949	499	239	308	1,108	3,578	15,536	41,695	71,875	140,618	281,026	327,420	47
2000	899,801	526	196	313	1,137	3,357	15,433	41,021	73,064	148,517	285,960	330,234	43
1999	913,476	528	192	306	1,152	3,453	15,869	39,935	73,473	155,643	292,354	330,537	34
Male													
2001	408,437	260	128	155	692	2,231	10,366	28,916	47,469	83,178	135,395	99,611	36
2000	416,837	284	107	176	672	2,088	10,294	28,702	48,495	88,005	137,754	100,232	28
1999	422,824	298	106	169	694	2,142	10,649	27,907	48,675	92,273	140,588	99,298	25
Female													
2001	475,512	239	111	153	416	1,347	5,170	12,779	24,406	57,440	145,631	227,809	11
2000	482,964	242	89	137	465	1,269	5,139	12,319	24,569	60,512	148,206	230,002	15
1999	490,652	230	86	137	458	1,311	5,220	12,028	24,798	63,370	151,766	231,239	9

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Major cardiovascular diseases (I00-I78)													
Non-Hispanic white													
Both sexes													
2001	764,021	289	140	204	671	2,240	10,524	29,967	55,480	116,098	248,922	299,458	28
2000	780,445	330	126	195	690	2,084	10,556	29,780	56,666	123,708	254,089	302,192	29
1999	793,945	324	121	203	730	2,184	10,807	29,134	57,094	130,111	260,395	302,824	18
Male													
2001	352,013	160	75	108	425	1,441	7,379	21,723	37,726	70,213	121,332	91,409	22
2000	360,757	180	75	118	418	1,361	7,422	21,785	38,681	75,026	123,834	91,837	20
1999	366,310	189	70	110	430	1,400	7,606	21,200	38,851	78,853	126,770	90,818	13
Female													
2001	412,008	129	65	96	246	799	3,145	8,244	17,754	45,885	127,590	208,049	6
2000	419,688	150	51	77	272	723	3,134	7,995	17,985	48,682	130,255	210,355	9
1999	427,635	135	51	93	300	784	3,201	7,934	18,243	51,258	133,625	212,006	5
Non-Hispanic black													
Both sexes													
2001	103,213	184	86	85	395	1,209	4,506	10,539	14,483	21,084	27,025	23,600	17
2000	103,239	166	60	95	396	1,160	4,396	10,109	14,502	21,561	26,954	23,826	14
1999	103,629	172	60	93	388	1,131	4,610	9,721	14,487	22,242	27,129	23,581	15
Male													
2001	47,646	93	45	39	240	706	2,625	6,386	8,489	11,055	11,536	6,419	13
2000	47,408	88	27	47	218	652	2,544	6,143	8,568	11,074	11,434	6,605	8
1999	47,982	91	32	53	240	659	2,716	5,984	8,609	11,567	11,329	6,691	11
Female													
2001	55,567	91	41	46	155	503	1,881	4,153	5,994	10,029	15,489	17,181	4
2000	55,831	78	33	48	178	508	1,852	3,966	5,934	10,487	15,520	17,221	6

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Major cardiovascular diseases (I00-I78)													
Non-Hispanic black													
Female													
1999	55,647	81	28	40	148	472	1,894	3,737	5,878	10,675	15,800	16,890	4
Origin not stated [2]													
Both sexes													
2001	2,820	3	2	1	10	18	69	194	353	466	830	856	18
2000	3,063	4	1	2	3	16	97	211	377	556	862	920	14
1999	3,173	1	1	1	3	12	84	191	387	635	907	942	9
Male													
2001	1,490	2	-	-	7	16	56	138	269	309	428	254	11
2000	1,633	3	-	-	1	7	77	162	271	382	421	298	11
1999	1,699	-	-	1	2	8	67	138	290	419	460	307	7
Female													
2001	1,330	1	2	1	3	2	13	56	84	157	402	602	7
2000	1,430	1	1	2	2	9	20	49	106	174	441	622	3
1999	1,474	1	1	-	1	4	17	53	97	216	447	635	2
Diseases of heart (I00-I09,I11,I13,I20-I51)													
All origins													
Both sexes													
2001	700,142	479	225	272	999	3,160	13,326	36,399	62,486	116,299	216,992	249,439	66
2000	710,760	493	181	271	1,031	2,958	13,181	35,480	63,399	122,405	220,060	251,242	59
1999	725,192	524	183	277	1,069	3,066	13,600	34,994	64,167	129,253	226,152	251,860	47
Male													
2001	339,095	243	121	138	647	2,068	9,346	26,280	42,519	70,637	108,576	78,473	47
2000	344,807	260	107	163	638	1,922	9,286	25,927	43,285	74,589	109,693	78,896	41

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Diseases of heart (I00-I09,I11,I13,I20-I51)													
All origins													
Male													
1999	351,617	270	104	157	664	1,988	9,640	25,489	43,662	78,836	112,470	78,300	37
Female													
2001	361,047	236	104	134	352	1,092	3,980	10,119	19,967	45,662	108,416	170,966	19
2000	365,953	233	74	108	393	1,036	3,895	9,553	20,114	47,816	110,367	172,346	18
1999	373,575	254	79	120	405	1,078	3,960	9,505	20,505	50,417	113,682	173,560	10
Hispanic													
Both sexes													
2001	27,090	79	33	38	112	314	750	1,875	3,152	5,730	7,632	7,369	6
2000	25,819	86	29	38	125	280	760	1,754	3,136	5,453	7,067	7,086	5
1999	25,866	96	33	36	116	286	761	1,806	3,155	5,557	7,046	6,968	6
Male													
2001	14,195	43	16	19	76	226	565	1,397	2,104	3,424	3,808	2,512	5
2000	13,566	44	21	26	87	199	550	1,307	2,128	3,245	3,525	2,430	4
1999	13,554	40	20	23	73	219	572	1,348	2,091	3,355	3,483	2,324	6
Female													
2001	12,895	36	17	19	36	88	185	478	1,048	2,306	3,824	4,857	1
2000	12,253	42	8	12	38	81	210	447	1,008	2,208	3,542	4,656	1
1999	12,312	56	13	13	43	67	189	458	1,064	2,202	3,563	4,644	-
Non-Hispanic [1]													
Both sexes													
2001	670,787	397	191	234	877	2,834	12,515	34,359	59,038	110,181	208,704	241,415	42
2000	682,470	403	151	232	904	2,664	12,341	33,551	59,943	116,476	212,328	243,436	41
1999	696,719	427	149	240	950	2,770	12,767	33,029	60,684	123,161	218,359	244,151	32

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Diseases of heart (I00-I09,I11,I13,I20-I51)													
Non-Hispanic [1]													
Male													
2001	323,673	198	105	119	564	1,832	8,733	24,759	40,188	66,949	104,429	75,766	31
2000	329,888	213	86	137	550	1,716	8,673	24,487	40,924	71,010	105,834	76,231	27
1999	336,615	230	84	133	589	1,762	9,009	24,019	41,321	75,118	108,605	75,721	24
Female													
2001	347,114	199	86	115	313	1,002	3,782	9,600	18,850	43,232	104,275	165,649	11
2000	352,582	190	65	95	354	948	3,668	9,064	19,019	45,466	106,494	167,205	14
1999	360,104	197	65	107	361	1,008	3,758	9,010	19,363	48,043	109,754	168,430	8
Non-Hispanic white													
Both sexes													
2001	582,349	234	108	152	522	1,775	8,734	25,544	46,433	91,855	185,536	221,430	26
2000	594,465	258	98	140	531	1,653	8,743	25,153	47,292	97,748	189,366	223,454	29
1999	607,704	259	92	156	585	1,755	8,936	24,823	47,950	103,750	195,187	224,193	18
Male													
2001	280,713	127	59	80	336	1,184	6,395	19,178	32,504	57,059	93,996	69,775	20
2000	287,210	139	59	87	326	1,115	6,446	19,101	33,179	61,024	95,576	70,138	20
1999	293,206	145	55	84	357	1,152	6,590	18,747	33,466	64,699	98,422	69,476	13
Female													
2001	301,636	107	49	72	186	591	2,339	6,366	13,929	34,796	91,540	151,655	6
2000	307,255	119	39	53	205	538	2,297	6,052	14,113	36,724	93,790	153,316	9
1999	314,498	114	37	72	228	603	2,346	6,076	14,484	39,051	96,765	154,717	5
Non-Hispanic black													
Both sexes													
2001	76,794	140	73	70	322	961	3,424	7,956	11,209	15,940	19,681	17,004	14

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Diseases of heart (I00-I09,I11,I13,I20-I51)													
Non-Hispanic black													
Both sexes													
2000	76,706	118	47	75	331	929	3,259	7,606	11,257	16,413	19,600	17,059	12
1999	77,713	141	49	77	336	913	3,506	7,448	11,346	17,019	19,808	17,056	14
Male													
2001	36,541	65	40	34	205	578	2,071	4,955	6,713	8,491	8,621	4,758	10
2000	36,282	59	24	41	192	543	1,972	4,787	6,786	8,560	8,476	4,835	7
1999	37,077	71	27	45	212	546	2,165	4,716	6,907	9,016	8,416	4,945	11
Female													
2001	40,253	75	33	36	117	383	1,353	3,001	4,496	7,449	11,060	12,246	4
2000	40,424	59	23	34	139	386	1,287	2,819	4,471	7,853	11,124	12,224	5
1999	40,636	70	22	32	124	367	1,341	2,732	4,439	8,003	11,392	12,111	3
Origin not stated [2]													
Both sexes													
2001	2,265	3	1	-	10	12	61	165	296	388	656	655	18
2000	2,471	4	1	1	2	14	80	175	320	476	665	720	13
1999	2,607	1	1	1	3	10	72	159	328	535	747	741	9
Male													
2001	1,227	2	-	-	7	10	48	124	227	264	339	195	11
2000	1,353	3	-	-	1	7	63	133	233	334	334	235	10
1999	1,448	-	-	1	2	7	59	122	250	363	382	255	7
Female													
2001	1,038	1	1	-	3	2	13	41	69	124	317	460	7
2000	1,118	1	1	1	1	7	17	42	87	142	331	485	3
1999	1,159	1	1	-	1	3	13	37	78	172	365	486	2

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)													
All origins													
Both sexes													
2001	3,489	3	1	5	11	34	130	232	361	744	1,188	780	-
2000	3,582	6	2	4	17	33	141	252	420	819	1,176	712	-
1999	3,676	6	1	5	13	42	124	245	411	854	1,227	748	-
Male													
2001	1,031	3	1	2	7	15	54	87	135	231	330	166	-
2000	1,072	5	2	3	11	12	61	104	166	237	321	150	-
1999	1,042	2	1	2	10	17	49	90	133	290	325	123	-
Female													
2001	2,458	-	-	3	4	19	76	145	226	513	858	614	-
2000	2,510	1	-	1	6	21	80	148	254	582	855	562	-
1999	2,634	4	-	3	3	25	75	155	278	564	902	625	-
Hispanic													
Both sexes													
2001	156	-	1	1	2	2	12	16	27	48	30	17	-
2000	154	1	-	-	1	6	13	19	30	47	27	10	-
1999	166	2	-	-	3	4	16	23	26	44	35	13	-
Male													
2001	51	-	1	1	1	2	5	5	7	15	11	3	-
2000	49	1	-	-	-	1	4	9	12	13	6	3	-
1999	38	-	-	-	3	3	3	6	8	10	5	-	-
Female													
2001	105	-	-	-	1	-	7	11	20	33	19	14	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)													
Hispanic													
Female													
2000	105	-	-	-	1	5	9	10	18	34	21	7	-
1999	128	2	-	-	-	1	13	17	18	34	30	13	-
Non-Hispanic [1]													
Both sexes													
2001	3,327	3	-	4	8	32	118	214	333	696	1,157	762	-
2000	3,424	5	2	4	16	27	128	233	388	772	1,149	700	-
1999	3,498	4	1	5	10	38	106	221	383	806	1,190	734	-
Male													
2001	978	3	-	1	5	13	49	81	128	216	319	163	-
2000	1,022	4	2	3	11	11	57	95	153	224	315	147	-
1999	999	2	1	2	7	14	45	83	123	279	320	123	-
Female													
2001	2,349	-	-	3	3	19	69	133	205	480	838	599	-
2000	2,402	1	-	1	5	16	71	138	235	548	834	553	-
1999	2,499	2	-	3	3	24	61	138	260	527	870	611	-
Non-Hispanic white													
Both sexes													
2001	2,924	1	-	2	7	13	73	139	265	601	1,094	729	-
2000	3,073	4	2	4	9	19	83	178	320	687	1,102	665	-
1999	3,130	2	1	3	5	21	68	163	314	730	1,125	698	-
Male													
2001	857	1	-	-	5	7	35	58	108	187	302	154	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)													
Non-Hispanic white													
Male													
2000	917	3	2	3	6	9	42	78	136	197	303	138	-
1999	892	1	1	-	2	9	31	65	102	260	301	120	-
Female													
2001	2,067	-	-	2	2	6	38	81	157	414	792	575	-
2000	2,156	1	-	1	3	10	41	100	184	490	799	527	-
1999	2,238	1	-	3	3	12	37	98	212	470	824	578	-
Non-Hispanic black													
Both sexes													
2001	302	2	-	1	1	13	41	60	53	69	40	22	-
2000	263	1	-	-	7	8	35	43	48	58	36	27	-
1999	269	1	-	2	4	13	30	49	51	54	39	26	-
Male													
2001	86	2	-	-	-	3	13	20	17	17	8	6	-
2000	77	1	-	-	5	2	12	13	12	16	8	8	-
1999	86	1	-	2	4	3	12	18	16	15	14	1	-
Female													
2001	216	-	-	1	1	10	28	40	36	52	32	16	-
2000	186	-	-	-	2	6	23	30	36	42	28	19	-
1999	183	-	-	-	-	10	18	31	35	39	25	25	-
Origin not stated [2]													
Both sexes													
2001	6	-	-	-	1	-	-	2	1	-	1	1	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)													
Origin not stated [2]													
Both sexes													
2000	4	-	-	-	-	-	-	-	2	-	-	2	-
1999	12	-	-	-	-	-	2	1	2	4	2	1	-
Male													
2001	2	-	-	-	1	-	-	1	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	1	-	-	-	-
1999	5	-	-	-	-	-	1	1	2	1	-	-	-
Female													
2001	4	-	-	-	-	-	-	1	1	-	1	1	-
2000	3	-	-	-	-	-	-	-	1	-	-	2	-
1999	7	-	-	-	-	-	1	-	-	3	2	1	-
Hypertensive heart disease (I11)													
All origins													
Both sexes													
2001	24,689	-	-	1	24	246	1,091	2,464	2,769	3,619	6,119	8,345	11
2000	23,761	-	1	2	32	213	981	2,104	2,672	3,605	5,962	8,186	3
1999	22,702	1	-	-	33	210	964	1,930	2,537	3,517	5,830	7,679	1
Male													
2001	10,451	-	-	-	17	169	740	1,646	1,749	1,961	2,414	1,746	9
2000	9,876	-	-	1	24	139	643	1,436	1,684	1,971	2,269	1,707	2
1999	9,316	-	-	-	25	149	660	1,259	1,560	1,816	2,252	1,594	1
Female													
2001	14,238	-	-	1	7	77	351	818	1,020	1,658	3,705	6,599	2

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Hypertensive heart disease (I11)													
All origins													
Female													
2000	13,885	-	1	1	8	74	338	668	988	1,634	3,693	6,479	1
1999	13,386	1	-	-	8	61	304	671	977	1,701	3,578	6,085	-
Hispanic													
Both sexes													
2001	1,211	-	-	-	5	22	93	165	184	239	260	243	-
2000	991	-	-	-	1	20	64	118	141	198	233	216	-
1999	933	-	-	-	3	18	61	99	146	174	215	217	-
Male													
2001	648	-	-	-	4	15	75	118	120	132	108	76	-
2000	495	-	-	-	1	16	43	84	86	106	103	56	-
1999	455	-	-	-	2	16	45	69	96	95	82	50	-
Female													
2001	563	-	-	-	1	7	18	47	64	107	152	167	-
2000	496	-	-	-	-	4	21	34	55	92	130	160	-
1999	478	-	-	-	1	2	16	30	50	79	133	167	-
Non-Hispanic [1]													
Both sexes													
2001	23,343	-	-	1	18	222	991	2,270	2,554	3,364	5,837	8,077	9
2000	22,646	-	1	2	31	193	905	1,968	2,518	3,374	5,706	7,946	2
1999	21,639	1	-	-	30	192	898	1,815	2,363	3,315	5,591	7,433	1
Male													
2001	9,725	-	-	-	12	152	660	1,509	1,608	1,815	2,299	1,663	7
2000	9,298	-	-	1	23	123	591	1,338	1,586	1,839	2,152	1,644	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Hypertensive heart disease (I11)													
Non-Hispanic [1]													
Male													
1999	8,790	-	-	-	23	133	610	1,180	1,445	1,702	2,157	1,539	1
Female													
2001	13,618	-	-	1	6	70	331	761	946	1,549	3,538	6,414	2
2000	13,348	-	1	1	8	70	314	630	932	1,535	3,554	6,302	1
1999	12,849	1	-	-	7	59	288	635	918	1,613	3,434	5,894	-
Non-Hispanic white													
Both sexes													
2001	16,991	-	-	-	5	102	493	1,213	1,520	2,198	4,540	6,915	5
2000	16,373	-	-	2	14	75	476	1,042	1,441	2,193	4,386	6,744	-
1999	15,654	1	-	-	9	91	428	954	1,371	2,147	4,294	6,359	-
Male													
2001	6,728	-	-	-	3	80	367	866	1,016	1,237	1,781	1,374	4
2000	6,390	-	-	1	11	53	341	771	956	1,242	1,640	1,375	-
1999	6,013	-	-	-	7	75	323	676	882	1,122	1,647	1,281	-
Female													
2001	10,263	-	-	-	2	22	126	347	504	961	2,759	5,541	1
2000	9,983	-	-	1	3	22	135	271	485	951	2,746	5,369	-
1999	9,641	1	-	-	2	16	105	278	489	1,025	2,647	5,078	-
Non-Hispanic black													
Both sexes													
2001	5,880	-	-	1	13	114	474	1,007	972	1,078	1,164	1,053	4
2000	5,828	-	1	-	17	114	414	886	1,025	1,093	1,194	1,082	2
1999	5,556	-	-	-	20	98	447	838	932	1,084	1,182	954	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Hypertensive heart disease (I11)													
Non-Hispanic black													
Male													
2001	2,760	-	-	-	9	67	275	609	551	538	458	250	3
2000	2,685	-	-	-	12	66	240	538	594	547	452	235	1
1999	2,571	-	-	-	15	55	267	487	528	536	458	224	1
Female													
2001	3,120	-	-	1	4	47	199	398	421	540	706	803	1
2000	3,143	-	1	-	5	48	174	348	431	546	742	847	1
1999	2,985	-	-	-	5	43	180	351	404	548	724	730	-
Origin not stated [2]													
Both sexes													
2001	135	-	-	-	1	2	7	29	31	16	22	25	2
2000	124	-	-	-	-	-	12	18	13	33	23	24	1
1999	130	-	-	-	-	-	5	16	28	28	24	29	-
Male													
2001	78	-	-	-	1	2	5	19	21	14	7	7	2
2000	83	-	-	-	-	-	9	14	12	26	14	7	1
1999	71	-	-	-	-	-	5	10	19	19	13	5	-
Female													
2001	57	-	-	-	-	-	2	10	10	2	15	18	-
2000	41	-	-	-	-	-	3	4	1	7	9	17	-
1999	59	-	-	-	-	-	-	6	9	9	11	24	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Hypertensive heart and renal disease (I13)													
All origins													
Both sexes													
2001	2,826	1	-	-	7	28	99	146	247	417	855	1,026	-
2000	2,785	-	-	1	6	28	83	147	208	417	863	1,032	-
1999	3,327	-	-	2	6	21	114	156	262	470	1,087	1,209	-
Male													
2001	1,169	1	-	-	4	17	63	99	126	206	339	314	-
2000	1,163	-	-	-	4	18	49	87	108	206	350	341	-
1999	1,390	-	-	1	3	16	67	100	146	223	460	374	-
Female													
2001	1,657	-	-	-	3	11	36	47	121	211	516	712	-
2000	1,622	-	-	1	2	10	34	60	100	211	513	691	-
1999	1,937	-	-	1	3	5	47	56	116	247	627	835	-
Hispanic													
Both sexes													
2001	160	-	-	-	1	2	9	10	21	34	41	42	-
2000	138	-	-	-	1	3	8	14	15	27	32	38	-
1999	144	-	-	-	-	1	9	11	19	32	39	33	-
Male													
2001	76	-	-	-	1	-	5	6	14	17	17	16	-
2000	67	-	-	-	1	2	4	7	5	19	18	11	-
1999	65	-	-	-	-	-	6	6	12	14	14	13	-
Female													
2001	84	-	-	-	-	2	4	4	7	17	24	26	-
2000	71	-	-	-	-	1	4	7	10	8	14	27	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Hypertensive heart and renal disease (I13)													
Hispanic													
Female													
1999	79	-	-	-	-	1	3	5	7	18	25	20	-
Non-Hispanic [1]													
Both sexes													
2001	2,650	1	-	-	6	26	87	135	225	380	810	980	-
2000	2,637	-	-	1	5	25	75	133	192	389	826	991	-
1999	3,171	-	-	2	6	20	102	142	242	436	1,047	1,174	-
Male													
2001	1,086	1	-	-	3	17	56	92	111	188	321	297	-
2000	1,089	-	-	-	3	16	45	80	102	186	329	328	-
1999	1,318	-	-	1	3	16	59	91	134	208	446	360	-
Female													
2001	1,564	-	-	-	3	9	31	43	114	192	489	683	-
2000	1,548	-	-	1	2	9	30	53	90	203	497	663	-
1999	1,853	-	-	1	3	4	43	51	108	228	601	814	-
Non-Hispanic white													
Both sexes													
2001	1,759	-	-	-	2	5	16	43	95	204	601	793	-
2000	1,769	-	-	-	-	5	30	40	75	220	593	806	-
1999	2,091	-	-	1	-	6	19	43	91	215	768	948	-
Male													
2001	687	-	-	-	2	3	12	27	42	101	250	250	-
2000	710	-	-	-	-	4	18	24	41	109	256	258	-
1999	843	-	-	1	-	6	11	26	48	108	346	297	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Hypertensive heart and renal disease (I13)													
Non-Hispanic white													
Female													
2001	1,072	-	-	-	-	2	4	16	53	103	351	543	-
2000	1,059	-	-	-	-	1	12	16	34	111	337	548	-
1999	1,248	-	-	-	-	-	8	17	43	107	422	651	-
Non-Hispanic black													
Both sexes													
2001	821	1	-	-	4	21	65	92	125	164	187	162	-
2000	797	-	-	1	5	19	45	91	112	161	203	160	-
1999	989	-	-	1	6	13	82	92	143	205	243	204	-
Male													
2001	370	1	-	-	1	14	41	65	66	81	62	39	-
2000	348	-	-	-	3	12	27	55	57	74	65	55	-
1999	426	-	-	-	3	9	47	60	81	92	83	51	-
Female													
2001	451	-	-	-	3	7	24	27	59	83	125	123	-
2000	449	-	-	1	2	7	18	36	55	87	138	105	-
1999	563	-	-	1	3	4	35	32	62	113	160	153	-
Origin not stated [2]													
Both sexes													
2001	16	-	-	-	-	-	3	1	1	3	4	4	-
2000	10	-	-	-	-	-	-	-	1	1	5	3	-
1999	12	-	-	-	-	-	3	3	1	2	1	2	-
Male													
2001	7	-	-	-	-	-	2	1	1	1	1	1	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Hypertensive heart and renal disease (I13)													
Origin not stated [2]													
Male													
2000	7	-	-	-	-	-	-	-	1	1	3	2	-
1999	7	-	-	-	-	-	2	3	-	1	-	1	-
Female													
2001	9	-	-	-	-	-	1	-	-	2	3	3	-
2000	3	-	-	-	-	-	-	-	-	-	2	1	-
1999	5	-	-	-	-	-	1	-	1	1	1	1	-
Ischemic heart diseases (I20-I25)													
All origins													
Both sexes													
2001	502,189	26	9	20	163	1,116	7,534	25,073	46,574	87,463	159,854	174,309	48
2000	515,204	30	10	16	123	935	7,519	24,890	47,737	92,885	163,692	177,319	48
1999	529,659	27	5	22	137	1,014	7,751	24,775	48,377	98,840	169,296	179,376	39
Male													
2001	254,005	17	3	13	119	830	5,712	19,141	33,006	55,072	82,953	57,106	33
2000	260,574	17	7	9	81	704	5,787	19,149	33,795	58,554	84,747	57,690	34
1999	267,268	13	3	17	93	739	5,970	19,061	34,123	62,187	87,305	57,726	31
Female													
2001	248,184	9	6	7	44	286	1,822	5,932	13,568	32,391	76,901	117,203	15
2000	254,630	13	3	7	42	231	1,732	5,741	13,942	34,331	78,945	119,629	14
1999	262,391	14	2	5	44	275	1,781	5,714	14,254	36,653	81,991	121,650	8
Hispanic													
Both sexes													
2001	20,664	4	1	4	18	109	368	1,314	2,404	4,553	6,095	5,790	4

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Ischemic heart diseases (I20-I25)													
Hispanic													
Both sexes													
2000	19,744	6	1	1	12	69	392	1,240	2,457	4,364	5,626	5,572	4
1999	19,936	2	-	2	11	87	366	1,328	2,463	4,492	5,655	5,525	5
Male													
2001	11,019	3	1	1	13	90	295	1,016	1,667	2,804	3,116	2,010	3
2000	10,562	4	1	1	9	56	322	973	1,726	2,661	2,888	1,918	3
1999	10,604	-	-	2	8	72	310	1,042	1,677	2,770	2,869	1,849	5
Female													
2001	9,645	1	-	3	5	19	73	298	737	1,749	2,979	3,780	1
2000	9,182	2	-	-	3	13	70	267	731	1,703	2,738	3,654	1
1999	9,332	2	-	-	3	15	56	286	786	1,722	2,786	3,676	-
Non-Hispanic [1]													
Both sexes													
2001	479,842	21	8	16	144	1,004	7,135	23,663	43,946	82,604	153,239	168,032	30
2000	493,541	23	9	15	111	861	7,082	23,523	45,022	88,137	157,523	171,202	33
1999	507,700	25	5	20	125	926	7,344	23,331	45,666	93,919	163,035	173,278	26
Male													
2001	242,054	13	2	12	105	737	5,392	18,047	31,160	52,059	79,549	54,956	22
2000	248,932	12	6	8	72	645	5,426	18,074	31,880	55,623	81,578	55,585	23
1999	255,524	13	3	15	85	666	5,623	17,928	32,249	59,117	84,127	55,678	20
Female													
2001	237,788	8	6	4	39	267	1,743	5,616	12,786	30,545	73,690	113,076	8
2000	244,609	11	3	7	39	216	1,656	5,449	13,142	32,514	75,945	115,617	10
1999	252,176	12	2	5	40	260	1,721	5,403	13,417	34,802	78,908	117,600	6

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Ischemic heart diseases (I20-I25)													
Non-Hispanic white													
Both sexes													
2001	420,959	14	5	10	86	718	5,456	18,518	35,421	69,681	136,815	154,217	18
2000	434,505	16	9	12	80	606	5,437	18,531	36,367	74,867	141,111	157,445	24
1999	447,960	16	4	12	76	665	5,607	18,425	37,066	80,129	146,594	159,351	15
Male													
2001	212,928	10	1	8	65	539	4,250	14,590	25,776	44,881	72,036	50,758	14
2000	219,732	9	6	6	52	473	4,319	14,723	26,398	48,374	74,042	51,313	17
1999	225,909	8	3	10	57	482	4,416	14,585	26,754	51,553	76,763	51,267	11
Female													
2001	208,031	4	4	2	21	179	1,206	3,928	9,645	24,800	64,779	103,459	4
2000	214,773	7	3	6	28	133	1,118	3,808	9,969	26,493	67,069	106,132	7
1999	222,051	8	1	2	19	183	1,191	3,840	10,312	28,576	69,831	108,084	4
Non-Hispanic black													
Both sexes													
2001	50,295	7	3	5	52	246	1,474	4,533	7,451	11,107	13,782	11,625	10
2000	50,659	4	-	2	24	229	1,451	4,418	7,587	11,485	13,809	11,641	9
1999	51,368	8	1	8	44	213	1,552	4,362	7,542	11,929	13,890	11,808	11
Male													
2001	24,255	3	1	3	35	167	974	2,982	4,596	6,089	6,115	3,283	7
2000	24,289	2	-	2	14	150	946	2,886	4,713	6,121	6,118	3,331	6
1999	24,736	4	-	5	24	148	1,042	2,904	4,737	6,437	5,991	3,435	9
Female													
2001	26,040	4	2	2	17	79	500	1,551	2,855	5,018	7,667	8,342	3
2000	26,370	2	-	-	10	79	505	1,532	2,874	5,364	7,691	8,310	3

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Ischemic heart diseases (I20-I25)													
Non-Hispanic black													
Female													
1999	26,632	4	1	3	20	65	510	1,458	2,805	5,492	7,899	8,373	2
Origin not stated [2]													
Both sexes													
2001	1,683	1	-	-	1	3	31	96	224	306	520	487	14
2000	1,919	1	-	-	-	5	45	127	258	384	543	545	11
1999	2,023	-	-	-	1	1	41	116	248	429	606	573	8
Male													
2001	932	1	-	-	1	3	25	78	179	209	288	140	8
2000	1,080	1	-	-	-	3	39	102	189	270	281	187	8
1999	1,140	-	-	-	-	1	37	91	197	300	309	199	6
Female													
2001	751	-	-	-	-	-	6	18	45	97	232	347	6
2000	839	-	-	-	-	2	6	25	69	114	262	358	3
1999	883	-	-	-	1	-	4	25	51	129	297	374	2
Acute myocardial infarction (I21-I22)													
All origins													
Both sexes													
2001	184,757	15	2	7	64	475	3,176	10,817	19,830	35,380	59,254	55,727	10
2000	192,898	20	1	6	52	375	3,132	10,877	21,013	38,459	62,210	56,745	8
1999	199,454	13	2	11	60	434	3,249	10,823	21,406	41,441	64,711	57,297	7
Male													
2001	95,875	10	2	4	48	333	2,384	8,158	13,858	21,963	30,125	18,984	6
2000	100,306	10	1	4	30	269	2,387	8,265	14,638	23,823	31,631	19,243	5

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Acute myocardial infarction (I21-I22)													
All origins													
Male													
1999	104,169	6	1	7	40	306	2,470	8,300	15,029	25,708	32,943	19,357	2
Female													
2001	88,882	5	-	3	16	142	792	2,659	5,972	13,417	29,129	36,743	4
2000	92,592	10	-	2	22	106	745	2,612	6,375	14,636	30,579	37,502	3
1999	95,285	7	1	4	20	128	779	2,523	6,377	15,733	31,768	37,940	5
Hispanic													
Both sexes													
2001	7,403	4	1	1	6	46	139	503	914	1,701	2,190	1,896	2
2000	7,106	5	-	-	5	31	154	486	941	1,661	2,049	1,774	-
1999	7,158	2	-	-	4	32	142	528	919	1,699	2,082	1,750	-
Male													
2001	3,972	3	1	-	6	37	110	383	603	1,036	1,093	699	1
2000	3,825	3	-	-	4	24	125	363	643	984	1,047	632	-
1999	3,836	-	-	-	3	27	116	391	607	1,028	1,072	592	-
Female													
2001	3,431	1	-	1	-	9	29	120	311	665	1,097	1,197	1
2000	3,281	2	-	-	1	7	29	123	298	677	1,002	1,142	-
1999	3,322	2	-	-	1	5	26	137	312	671	1,010	1,158	-
Non-Hispanic [1]													
Both sexes													
2001	176,895	11	1	6	58	428	3,023	10,294	18,844	33,604	56,923	53,696	7
2000	185,283	14	1	6	47	344	2,963	10,363	20,000	36,695	60,020	54,823	7
1999	191,729	11	2	11	56	402	3,090	10,264	20,409	39,614	62,466	55,400	4

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Acute myocardial infarction (I21-I22)													
Non-Hispanic [1]													
Male													
2001	91,655	7	1	4	42	295	2,261	7,758	13,202	20,875	28,956	18,249	5
2000	96,218	6	1	4	26	245	2,250	7,886	13,950	22,772	30,516	18,558	4
1999	100,014	6	1	7	37	279	2,338	7,887	14,363	24,598	31,785	18,712	1
Female													
2001	85,240	4	-	2	16	133	762	2,536	5,642	12,729	27,967	35,447	2
2000	89,065	8	-	2	21	99	713	2,477	6,050	13,923	29,504	36,265	3
1999	91,715	5	1	4	19	123	752	2,377	6,046	15,016	30,681	36,688	3
Non-Hispanic white													
Both sexes													
2001	155,312	8	1	1	36	313	2,367	8,302	15,677	28,749	50,908	48,946	4
2000	163,203	10	1	4	34	243	2,336	8,410	16,678	31,568	53,840	50,073	6
1999	169,545	7	2	6	35	293	2,446	8,404	17,116	34,339	56,289	50,606	2
Male													
2001	81,175	6	1	1	28	219	1,834	6,491	11,303	18,302	26,282	16,704	4
2000	85,412	5	1	2	20	183	1,839	6,635	11,919	20,062	27,749	16,994	3
1999	89,142	3	1	4	26	202	1,907	6,671	12,319	21,791	29,104	17,114	-
Female													
2001	74,137	2	-	-	8	94	533	1,811	4,374	10,447	24,626	32,242	-
2000	77,791	5	-	2	14	60	497	1,775	4,759	11,506	26,091	33,079	3
1999	80,403	4	1	2	9	91	539	1,733	4,797	12,548	27,185	33,492	2
Non-Hispanic black													
Both sexes													
2001	18,399	3	-	4	21	97	572	1,740	2,752	4,173	5,039	3,995	3

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Acute myocardial infarction (I21-I22)													
Non-Hispanic black													
Both sexes													
2000	18,934	2	-	2	11	93	560	1,720	2,910	4,415	5,189	4,031	1
1999	19,120	4	-	5	19	88	580	1,653	2,862	4,542	5,256	4,109	2
Male													
2001	8,628	1	-	2	13	62	354	1,068	1,606	2,163	2,155	1,203	1
2000	8,959	1	-	2	5	55	355	1,068	1,731	2,279	2,247	1,215	1
1999	9,106	3	-	3	10	61	374	1,055	1,743	2,375	2,211	1,270	1
Female													
2001	9,771	2	-	2	8	35	218	672	1,146	2,010	2,884	2,792	2
2000	9,975	1	-	-	6	38	205	652	1,179	2,136	2,942	2,816	-
1999	10,014	1	-	2	9	27	206	598	1,119	2,167	3,045	2,839	1
Origin not stated [2]													
Both sexes													
2001	459	-	-	-	-	1	14	20	72	75	141	135	1
2000	509	1	-	-	-	-	15	28	72	103	141	148	1
1999	567	-	-	-	-	-	17	31	78	128	163	147	3
Male													
2001	248	-	-	-	-	1	13	17	53	52	76	36	-
2000	263	1	-	-	-	-	12	16	45	67	68	53	1
1999	319	-	-	-	-	-	16	22	59	82	86	53	1
Female													
2001	211	-	-	-	-	-	1	3	19	23	65	99	1
2000	246	-	-	-	-	-	3	12	27	36	73	95	-
1999	248	-	-	-	-	-	1	9	19	46	77	94	2

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other acute ischemic heart diseases (I24)													
All origins													
Both sexes													
2001	3,351	3	-	2	7	14	94	307	463	645	871	945	-
2000	3,363	1	1	-	8	16	92	282	484	644	906	928	1
1999	3,232	3	1	1	3	6	81	252	448	674	855	908	-
Male													
2001	1,720	2	-	1	6	10	65	230	320	393	424	269	-
2000	1,749	1	1	-	6	13	71	217	341	398	446	254	1
1999	1,665	1	1	1	3	4	63	178	315	410	423	266	-
Female													
2001	1,631	1	-	1	1	4	29	77	143	252	447	676	-
2000	1,614	-	-	-	2	3	21	65	143	246	460	674	-
1999	1,567	2	-	-	-	2	18	74	133	264	432	642	-
Hispanic													
Both sexes													
2001	63	-	-	-	-	-	1	4	12	19	14	13	-
2000	69	-	1	-	1	1	1	6	10	15	17	17	-
1999	53	-	-	1	-	-	1	7	13	8	12	11	-
Male													
2001	42	-	-	-	-	-	1	2	9	14	7	9	-
2000	39	-	1	-	1	1	1	3	5	11	9	7	-
1999	26	-	-	1	-	-	1	4	6	4	8	2	-
Female													
2001	21	-	-	-	-	-	-	2	3	5	7	4	-
2000	30	-	-	-	-	-	-	3	5	4	8	10	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other acute ischemic heart diseases (I24)													
Hispanic													
Female													
1999	27	-	-	-	-	-	-	3	7	4	4	9	-
Non-Hispanic [1]													
Both sexes													
2001	3,284	3	-	2	7	14	93	303	450	625	855	932	-
2000	3,282	1	-	-	7	15	90	273	473	626	887	909	1
1999	3,167	3	1	-	3	6	80	245	433	664	841	891	-
Male													
2001	1,675	2	-	1	6	10	64	228	310	378	416	260	-
2000	1,705	1	-	-	5	12	69	212	336	386	436	247	1
1999	1,635	1	1	-	3	4	62	174	307	406	414	263	-
Female													
2001	1,609	1	-	1	1	4	29	75	140	247	439	672	-
2000	1,577	-	-	-	2	3	21	61	137	240	451	662	-
1999	1,532	2	-	-	-	2	18	71	126	258	427	628	-
Non-Hispanic white													
Both sexes													
2001	2,756	3	-	2	4	11	62	222	342	511	757	842	-
2000	2,767	1	-	-	5	11	66	207	357	514	774	832	-
1999	2,689	1	1	-	3	5	52	169	344	553	736	825	-
Male													
2001	1,386	2	-	1	3	9	45	170	240	307	374	235	-
2000	1,412	1	-	-	3	8	52	165	261	314	382	226	-
1999	1,361	1	1	-	3	4	40	123	240	339	366	244	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other acute ischemic heart diseases (I24)													
Non-Hispanic white													
Female													
2001	1,370	1	-	1	1	2	17	52	102	204	383	607	-
2000	1,355	-	-	-	2	3	14	42	96	200	392	606	-
1999	1,328	-	-	-	-	1	12	46	104	214	370	581	-
Non-Hispanic black													
Both sexes													
2001	453	-	-	-	3	2	29	69	98	95	79	78	-
2000	449	-	-	-	1	4	22	61	103	91	97	69	1
1999	429	2	-	-	-	1	26	73	83	93	97	54	-
Male													
2001	249	-	-	-	3	-	19	50	62	59	35	21	-
2000	255	-	-	-	1	4	16	44	67	58	46	18	1
1999	246	-	-	-	-	-	20	49	63	56	41	17	-
Female													
2001	204	-	-	-	-	2	10	19	36	36	44	57	-
2000	194	-	-	-	-	-	6	17	36	33	51	51	-
1999	183	2	-	-	-	1	6	24	20	37	56	37	-
Origin not stated [2]													
Both sexes													
2001	4	-	-	-	-	-	-	-	1	1	2	-	-
2000	12	-	-	-	-	-	1	3	1	3	2	2	-
1999	12	-	-	-	-	-	-	-	2	2	2	6	-
Male													
2001	3	-	-	-	-	-	-	-	1	1	1	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other acute ischemic heart diseases (I24)													
Origin not stated [2]													
Male													
2000	5	-	-	-	-	-	1	2	-	1	1	-	-
1999	4	-	-	-	-	-	-	-	2	-	1	1	-
Female													
2001	1	-	-	-	-	-	-	-	-	-	1	-	-
2000	7	-	-	-	-	-	-	1	1	2	1	2	-
1999	8	-	-	-	-	-	-	-	-	2	1	5	-
Other forms of chronic ischemic heart disease (I20,I25)													
All origins													
Both sexes													
2001	314,081	8	7	11	92	627	4,264	13,949	26,281	51,438	99,729	117,637	38
2000	318,943	9	8	10	63	544	4,295	13,731	26,240	53,782	100,576	119,646	39
1999	326,973	11	2	10	74	574	4,421	13,700	26,523	56,725	103,730	121,171	32
Male													
2001	156,410	5	1	8	65	487	3,263	10,753	18,828	32,716	52,404	37,853	27
2000	158,519	6	5	5	45	422	3,329	10,667	18,816	34,333	52,670	38,193	28
1999	161,434	6	1	9	50	429	3,437	10,583	18,779	36,069	53,939	38,103	29
Female													
2001	157,671	3	6	3	27	140	1,001	3,196	7,453	18,722	47,325	79,784	11
2000	160,424	3	3	5	18	122	966	3,064	7,424	19,449	47,906	81,453	11
1999	165,539	5	1	1	24	145	984	3,117	7,744	20,656	49,791	83,068	3

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other forms of chronic ischemic heart disease (I20,I25)													
Hispanic													
Both sexes													
2001	13,198	-	-	3	12	63	228	807	1,478	2,833	3,891	3,881	2
2000	12,569	1	-	1	6	37	237	748	1,506	2,688	3,560	3,781	4
1999	12,725	-	-	1	7	55	223	793	1,531	2,785	3,561	3,764	5
Male													
2001	7,005	-	-	1	7	53	184	631	1,055	1,754	2,016	1,302	2
2000	6,698	1	-	1	4	31	196	607	1,078	1,666	1,832	1,279	3
1999	6,742	-	-	1	5	45	193	647	1,064	1,738	1,789	1,255	5
Female													
2001	6,193	-	-	2	5	10	44	176	423	1,079	1,875	2,579	-
2000	5,871	-	-	-	2	6	41	141	428	1,022	1,728	2,502	1
1999	5,983	-	-	-	2	10	30	146	467	1,047	1,772	2,509	-
Non-Hispanic [1]													
Both sexes													
2001	299,663	7	7	8	79	562	4,019	13,066	24,652	48,375	95,461	113,404	23
2000	304,976	8	8	9	57	502	4,029	12,887	24,549	50,816	96,616	115,470	25
1999	312,804	11	2	9	66	518	4,174	12,822	24,824	53,641	99,728	116,987	22
Male													
2001	148,724	4	1	7	57	432	3,067	10,061	17,648	30,806	50,177	36,447	17
2000	151,009	5	5	4	41	388	3,107	9,976	17,594	32,465	50,626	36,780	18
1999	153,875	6	1	8	45	383	3,223	9,867	17,579	34,113	51,928	36,703	19
Female													
2001	150,939	3	6	1	22	130	952	3,005	7,004	17,569	45,284	76,957	6

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other forms of chronic ischemic heart disease (I20,I25)													
Non-Hispanic [1]													
Female													
2000	153,967	3	3	5	16	114	922	2,911	6,955	18,351	45,990	78,690	7
1999	158,929	5	1	1	21	135	951	2,955	7,245	19,528	47,800	80,284	3
Non-Hispanic white													
Both sexes													
2001	262,891	3	4	7	46	394	3,027	9,994	19,402	40,421	85,150	104,429	14
2000	268,535	5	8	8	41	352	3,035	9,914	19,332	42,785	86,497	106,540	18
1999	275,726	8	1	6	38	367	3,109	9,852	19,606	45,237	89,569	107,920	13
Male													
2001	130,367	2	-	6	34	311	2,371	7,929	14,233	26,272	45,380	33,819	10
2000	132,908	3	5	4	29	282	2,428	7,923	14,218	27,998	45,911	34,093	14
1999	135,406	4	1	6	28	276	2,469	7,791	14,195	29,423	47,293	33,909	11
Female													
2001	132,524	1	4	1	12	83	656	2,065	5,169	14,149	39,770	70,610	4
2000	135,627	2	3	4	12	70	607	1,991	5,114	14,787	40,586	72,447	4
1999	140,320	4	-	-	10	91	640	2,061	5,411	15,814	42,276	74,011	2
Non-Hispanic black													
Both sexes													
2001	31,443	4	3	1	28	147	873	2,724	4,601	6,839	8,664	7,552	7
2000	31,276	2	-	-	12	132	869	2,637	4,574	6,979	8,523	7,541	7
1999	31,819	2	1	3	25	124	946	2,636	4,597	7,294	8,537	7,645	9
Male													
2001	15,378	2	1	1	19	105	601	1,864	2,928	3,867	3,925	2,059	6

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other forms of chronic ischemic heart disease (I20,I25)													
Non-Hispanic black													
Male													
2000	15,075	1	-	-	8	91	575	1,774	2,915	3,784	3,825	2,098	4
1999	15,384	1	-	2	14	87	648	1,800	2,931	4,006	3,739	2,148	8
Female													
2001	16,065	2	2	-	9	42	272	860	1,673	2,972	4,739	5,493	1
2000	16,201	1	-	-	4	41	294	863	1,659	3,195	4,698	5,443	3
1999	16,435	1	1	1	11	37	298	836	1,666	3,288	4,798	5,497	1
Origin not stated [2]													
Both sexes													
2001	1,220	1	-	-	1	2	17	76	151	230	377	352	13
2000	1,398	-	-	-	-	5	29	96	185	278	400	395	10
1999	1,444	-	-	-	1	1	24	85	168	299	441	420	5
Male													
2001	681	1	-	-	1	2	12	61	125	156	211	104	8
2000	812	-	-	-	-	3	26	84	144	202	212	134	7
1999	817	-	-	-	-	1	21	69	136	218	222	145	5
Female													
2001	539	-	-	-	-	-	5	15	26	74	166	248	5
2000	586	-	-	-	-	2	3	12	41	76	188	261	3
1999	627	-	-	-	1	-	3	16	32	81	219	275	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Atherosclerotic cardiovascular disease, so described (I25.0)													
All origins													
Both sexes													
2001	68,103	-	-	-	20	179	1,457	5,047	8,438	12,005	19,196	21,736	25
2000	69,554	1	-	1	15	144	1,376	4,980	8,242	12,567	19,545	22,664	19
1999	71,926	-	1	1	13	160	1,488	4,737	8,383	13,395	20,321	23,403	24
Male													
2001	35,440	-	-	-	16	133	1,130	4,037	6,245	7,668	9,671	6,521	19
2000	35,896	-	-	1	12	115	1,099	3,977	6,125	8,069	9,725	6,758	15
1999	36,633	-	-	1	10	119	1,192	3,752	6,123	8,546	9,963	6,905	22
Female													
2001	32,663	-	-	-	4	46	327	1,010	2,193	4,337	9,525	15,215	6
2000	33,658	1	-	-	3	29	277	1,003	2,117	4,498	9,820	15,906	4
1999	35,293	-	1	-	3	41	296	985	2,260	4,849	10,358	16,498	2
Hispanic													
Both sexes													
2001	3,150	-	-	-	1	25	81	343	548	748	735	667	2
2000	2,953	-	-	-	2	7	83	299	522	707	649	682	2
1999	3,063	-	-	-	1	14	73	308	540	747	708	669	3
Male													
2001	1,928	-	-	-	-	21	63	288	416	502	405	231	2
2000	1,794	-	-	-	2	6	71	257	406	438	368	245	1
1999	1,847	-	-	-	1	12	63	260	384	527	372	225	3
Female													
2001	1,222	-	-	-	1	4	18	55	132	246	330	436	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Atherosclerotic cardiovascular disease, so described (I25.0)													
Hispanic													
Female													
2000	1,159	-	-	-	-	1	12	42	116	269	281	437	1
1999	1,216	-	-	-	-	2	10	48	156	220	336	444	-
Non-Hispanic [1]													
Both sexes													
2001	64,534	-	-	-	19	154	1,366	4,664	7,816	11,159	18,346	20,999	11
2000	66,141	1	-	1	13	137	1,283	4,631	7,627	11,756	18,769	21,912	11
1999	68,373	-	1	1	12	146	1,404	4,387	7,754	12,524	19,465	22,661	18
Male													
2001	33,247	-	-	-	16	112	1,059	3,717	5,770	7,093	9,200	6,270	10
2000	33,786	-	-	1	10	109	1,019	3,674	5,646	7,549	9,281	6,487	10
1999	34,451	-	-	1	9	107	1,118	3,457	5,663	7,922	9,504	6,654	16
Female													
2001	31,287	-	-	-	3	42	307	947	2,046	4,066	9,146	14,729	1
2000	32,355	1	-	-	3	28	264	957	1,981	4,207	9,488	15,425	1
1999	33,922	-	1	-	3	39	286	930	2,091	4,602	9,961	16,007	2
Non-Hispanic white													
Both sexes													
2001	53,031	-	-	-	11	108	999	3,421	5,649	8,504	15,521	18,811	7
2000	54,544	-	-	1	11	95	944	3,407	5,537	9,079	15,871	19,591	8
1999	56,375	-	-	1	5	96	1,017	3,155	5,626	9,665	16,517	20,283	10
Male													
2001	26,960	-	-	-	8	82	794	2,809	4,291	5,496	7,849	5,625	6

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Atherosclerotic cardiovascular disease, so described (I25.0)													
Non-Hispanic white													
Male													
2000	27,461	-	-	1	8	77	775	2,786	4,210	5,949	7,864	5,783	8
1999	27,943	-	-	1	5	73	838	2,558	4,208	6,227	8,097	5,928	8
Female													
2001	26,071	-	-	-	3	26	205	612	1,358	3,008	7,672	13,186	1
2000	27,083	-	-	-	3	18	169	621	1,327	3,130	8,007	13,808	-
1999	28,432	-	-	-	-	23	179	597	1,418	3,438	8,420	14,355	2
Non-Hispanic black													
Both sexes													
2001	10,156	-	-	-	7	38	324	1,115	1,927	2,361	2,472	1,909	3
2000	10,283	1	-	-	1	36	292	1,120	1,877	2,380	2,522	2,051	3
1999	10,643	-	1	-	7	41	358	1,105	1,904	2,553	2,593	2,073	8
Male													
2001	5,476	-	-	-	7	24	230	799	1,299	1,412	1,174	528	3
2000	5,528	-	-	-	1	26	207	802	1,282	1,398	1,204	606	2
1999	5,692	-	-	-	4	27	254	792	1,278	1,518	1,210	601	8
Female													
2001	4,680	-	-	-	-	14	94	316	628	949	1,298	1,381	-
2000	4,755	1	-	-	-	10	85	318	595	982	1,318	1,445	1
1999	4,951	-	1	-	3	14	104	313	626	1,035	1,383	1,472	-
Origin not stated [2]													
Both sexes													
2001	419	-	-	-	-	-	10	40	74	98	115	70	12

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Atherosclerotic cardiovascular disease, so described (I25.0)													
Origin not stated [2]													
Both sexes													
2000	460	-	-	-	-	-	10	50	93	104	127	70	6
1999	490	-	-	-	-	-	11	42	89	124	148	73	3
Male													
2001	265	-	-	-	-	-	8	32	59	73	66	20	7
2000	316	-	-	-	-	-	9	46	73	82	76	26	4
1999	335	-	-	-	-	-	11	35	76	97	87	26	3
Female													
2001	154	-	-	-	-	-	2	8	15	25	49	50	5
2000	144	-	-	-	-	-	1	4	20	22	51	44	2
1999	155	-	-	-	-	-	-	7	13	27	61	47	-
All other forms of chronic ischemic heart disease (I20, I25.1-I25.9)													
All origins													
Both sexes													
2001	245,978	8	7	11	72	448	2,807	8,902	17,843	39,433	80,533	95,901	13
2000	249,389	8	8	9	48	400	2,919	8,751	17,998	41,215	81,031	96,982	20
1999	255,047	11	1	9	61	414	2,933	8,963	18,140	43,330	83,409	97,768	8
Male													
2001	120,970	5	1	8	49	354	2,133	6,716	12,583	25,048	42,733	31,332	8
2000	122,623	6	5	4	33	307	2,230	6,690	12,691	26,264	42,945	31,435	13
1999	124,801	6	1	8	40	310	2,245	6,831	12,656	27,523	43,976	31,198	7
Female													
2001	125,008	3	6	3	23	94	674	2,186	5,260	14,385	37,800	64,569	5

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
All other forms of chronic ischemic heart disease (I20, I25.1-I25.9)													
All origins													
Female													
2000	126,766	2	3	5	15	93	689	2,061	5,307	14,951	38,086	65,547	7
1999	130,246	5	-	1	21	104	688	2,132	5,484	15,807	39,433	66,570	1
Hispanic													
Both sexes													
2001	10,048	-	-	3	11	38	147	464	930	2,085	3,156	3,214	-
2000	9,616	1	-	1	4	30	154	449	984	1,981	2,911	3,099	2
1999	9,662	-	-	1	6	41	150	485	991	2,038	2,853	3,095	2
Male													
2001	5,077	-	-	1	7	32	121	343	639	1,252	1,611	1,071	-
2000	4,904	1	-	1	2	25	125	350	672	1,228	1,464	1,034	2
1999	4,895	-	-	1	4	33	130	387	680	1,211	1,417	1,030	2
Female													
2001	4,971	-	-	2	4	6	26	121	291	833	1,545	2,143	-
2000	4,712	-	-	-	2	5	29	99	312	753	1,447	2,065	-
1999	4,767	-	-	-	2	8	20	98	311	827	1,436	2,065	-
Non-Hispanic [1]													
Both sexes													
2001	235,129	7	7	8	60	408	2,653	8,402	16,836	37,216	77,115	92,405	12
2000	238,835	7	8	8	44	365	2,746	8,256	16,922	39,060	77,847	93,558	14
1999	244,431	11	1	8	54	372	2,770	8,435	17,070	41,117	80,263	94,326	4
Male													
2001	115,477	4	1	7	41	320	2,008	6,344	11,878	23,713	40,977	30,177	7

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
All other forms of chronic ischemic heart disease (I20, I25.1-I25.9)													
Non-Hispanic [1]													
Male													
2000	117,223	5	5	3	31	279	2,088	6,302	11,948	24,916	41,345	30,293	8
1999	119,424	6	1	7	36	276	2,105	6,410	11,916	26,191	42,424	30,049	3
Female													
2001	119,652	3	6	1	19	88	645	2,058	4,958	13,503	36,138	62,228	5
2000	121,612	2	3	5	13	86	658	1,954	4,974	14,144	36,502	63,265	6
1999	125,007	5	-	1	18	96	665	2,025	5,154	14,926	37,839	64,277	1
Non-Hispanic white													
Both sexes													
2001	209,860	3	4	7	35	286	2,028	6,573	13,753	31,917	69,629	85,618	7
2000	213,991	5	8	7	30	257	2,091	6,507	13,795	33,706	70,626	86,949	10
1999	219,351	8	1	5	33	271	2,092	6,697	13,980	35,572	73,052	87,637	3
Male													
2001	103,407	2	-	6	26	229	1,577	5,120	9,942	20,776	37,531	28,194	4
2000	105,447	3	5	3	21	205	1,653	5,137	10,008	22,049	38,047	28,310	6
1999	107,463	4	1	5	23	203	1,631	5,233	9,987	23,196	39,196	27,981	3
Female													
2001	106,453	1	4	1	9	57	451	1,453	3,811	11,141	32,098	57,424	3
2000	108,544	2	3	4	9	52	438	1,370	3,787	11,657	32,579	58,639	4
1999	111,888	4	-	-	10	68	461	1,464	3,993	12,376	33,856	59,656	-
Non-Hispanic black													
Both sexes													
2001	21,287	4	3	1	21	109	549	1,609	2,674	4,478	6,192	5,643	4

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
All other forms of chronic ischemic heart disease (I20, I25.1-I25.9)													
Non-Hispanic black													
Both sexes													
2000	20,993	1	-	-	11	96	577	1,517	2,697	4,599	6,001	5,490	4
1999	21,176	2	-	3	18	83	588	1,531	2,693	4,741	5,944	5,572	1
Male													
2001	9,902	2	1	1	12	81	371	1,065	1,629	2,455	2,751	1,531	3
2000	9,547	1	-	-	7	65	368	972	1,633	2,386	2,621	1,492	2
1999	9,692	1	-	2	10	60	394	1,008	1,653	2,488	2,529	1,547	-
Female													
2001	11,385	2	2	-	9	28	178	544	1,045	2,023	3,441	4,112	1
2000	11,446	-	-	-	4	31	209	545	1,064	2,213	3,380	3,998	2
1999	11,484	1	-	1	8	23	194	523	1,040	2,253	3,415	4,025	1
Origin not stated [2]													
Both sexes													
2001	801	1	-	-	1	2	7	36	77	132	262	282	1
2000	938	-	-	-	-	5	19	46	92	174	273	325	4
1999	954	-	-	-	1	1	13	43	79	175	293	347	2
Male													
2001	416	1	-	-	1	2	4	29	66	83	145	84	1
2000	496	-	-	-	-	3	17	38	71	120	136	108	3
1999	482	-	-	-	-	1	10	34	60	121	135	119	2
Female													
2001	385	-	-	-	-	-	3	7	11	49	117	198	-
2000	442	-	-	-	-	2	2	8	21	54	137	217	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
All other forms of chronic ischemic heart disease (I20, I25.1-I25.9)													
Origin not stated [2]													
Female													
1999	472	-	-	-	1	-	3	9	19	54	158	228	-
Other heart diseases (I26-I51)													
All origins													
Both sexes													
2001	166,949	449	215	246	794	1,736	4,472	8,484	12,535	24,056	48,976	64,979	7
2000	165,428	457	168	248	853	1,749	4,457	8,087	12,362	24,679	48,367	63,993	8
1999	165,828	490	177	248	880	1,779	4,647	7,888	12,580	25,572	48,712	62,848	7
Male													
2001	72,439	222	117	123	500	1,037	2,777	5,307	7,503	13,167	22,540	19,141	5
2000	72,122	238	98	150	518	1,049	2,746	5,151	7,532	13,621	22,006	19,008	5
1999	72,601	255	100	137	533	1,067	2,894	4,979	7,700	14,320	22,128	18,483	5
Female													
2001	94,510	227	98	123	294	699	1,695	3,177	5,032	10,889	26,436	45,838	2
2000	93,306	219	70	98	335	700	1,711	2,936	4,830	11,058	26,361	44,985	3
1999	93,227	235	77	111	347	712	1,753	2,909	4,880	11,252	26,584	44,365	2
Hispanic													
Both sexes													
2001	4,899	75	31	33	86	179	268	370	516	856	1,206	1,277	2
2000	4,792	79	28	37	110	182	283	363	493	817	1,149	1,250	1
1999	4,687	92	33	34	99	176	309	345	501	815	1,102	1,180	1
Male													
2001	2,401	40	14	17	57	119	185	252	296	456	556	407	2

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other heart diseases (I26-I51)													
Hispanic													
Male													
2000	2,393	39	20	25	76	124	177	234	299	446	510	442	1
1999	2,392	40	20	21	60	128	208	225	298	466	513	412	1
Female													
2001	2,498	35	17	16	29	60	83	118	220	400	650	870	-
2000	2,399	40	8	12	34	58	106	129	194	371	639	808	-
1999	2,295	52	13	13	39	48	101	120	203	349	589	768	-
Non-Hispanic [1]													
Both sexes													
2001	161,625	372	183	213	701	1,550	4,184	8,077	11,980	23,137	47,661	63,564	3
2000	160,222	375	139	210	741	1,558	4,151	7,694	11,823	23,804	47,124	62,597	6
1999	160,711	397	143	213	779	1,594	4,317	7,520	12,030	24,685	47,496	61,532	5
Male													
2001	69,830	181	103	106	439	913	2,576	5,030	7,181	12,671	21,941	18,687	2
2000	69,547	197	78	125	441	921	2,554	4,900	7,203	13,138	21,460	18,527	3
1999	69,984	215	80	115	471	933	2,672	4,737	7,370	13,812	21,555	18,021	3
Female													
2001	91,795	191	80	107	262	637	1,608	3,047	4,799	10,466	25,720	44,877	1
2000	90,675	178	61	85	300	637	1,597	2,794	4,620	10,666	25,664	44,070	3
1999	90,727	182	63	98	308	661	1,645	2,783	4,660	10,873	25,941	43,511	2
Non-Hispanic white													
Both sexes													
2001	139,716	219	103	140	422	937	2,696	5,631	9,132	19,171	42,486	58,776	3
2000	138,745	238	87	122	428	948	2,717	5,362	9,089	19,781	42,174	57,794	5

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other heart diseases (I26-I51)													
Non-Hispanic white													
Both sexes													
1999	138,869	240	87	140	495	972	2,814	5,238	9,108	20,529	42,406	56,837	3
Male													
2001	59,513	116	58	72	261	555	1,731	3,637	5,562	10,653	19,627	17,239	2
2000	59,461	127	51	77	257	576	1,726	3,505	5,648	11,102	19,335	17,054	3
1999	59,549	136	51	73	291	580	1,809	3,395	5,680	11,656	19,365	16,511	2
Female													
2001	80,203	103	45	68	161	382	965	1,994	3,570	8,518	22,859	41,537	1
2000	79,284	111	36	45	171	372	991	1,857	3,441	8,679	22,839	40,740	2
1999	79,320	104	36	67	204	392	1,005	1,843	3,428	8,873	23,041	40,326	1
Non-Hispanic black													
Both sexes													
2001	19,496	130	70	63	252	567	1,370	2,264	2,608	3,522	4,508	4,142	-
2000	19,159	113	46	72	278	559	1,314	2,168	2,485	3,616	4,358	4,149	1
1999	19,531	132	48	66	262	576	1,395	2,107	2,678	3,747	4,454	4,064	2
Male													
2001	9,070	59	39	31	160	327	768	1,279	1,483	1,766	1,978	1,180	-
2000	8,883	56	24	39	158	313	747	1,295	1,410	1,802	1,833	1,206	-
1999	9,258	66	27	38	166	331	797	1,247	1,545	1,936	1,870	1,234	1
Female													
2001	10,426	71	31	32	92	240	602	985	1,125	1,756	2,530	2,962	-
2000	10,276	57	22	33	120	246	567	873	1,075	1,814	2,525	2,943	1
1999	10,273	66	21	28	96	245	598	860	1,133	1,811	2,584	2,830	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other heart diseases (I26-I51)													
Origin not stated [2]													
Both sexes													
2001	425	2	1	-	7	7	20	37	39	63	109	138	2
2000	414	3	1	1	2	9	23	30	46	58	94	146	1
1999	430	1	1	1	2	9	21	23	49	72	114	136	1
Male													
2001	208	1	-	-	4	5	16	25	26	40	43	47	1
2000	182	2	-	-	1	4	15	17	30	37	36	39	1
1999	225	-	-	1	2	6	14	17	32	42	60	50	1
Female													
2001	217	1	1	-	3	2	4	12	13	23	66	91	1
2000	232	1	1	1	1	5	8	13	16	21	58	107	-
1999	205	1	1	-	-	3	7	6	17	30	54	86	-
Acute and subacute endocarditis (I33)													
All origins													
Both sexes													
2001	1,178	1	-	3	8	29	117	186	160	247	316	111	-
2000	1,172	3	1	3	13	40	116	186	165	224	302	119	-
1999	1,116	2	1	-	7	30	114	143	171	240	289	119	-
Male													
2001	668	1	-	1	5	17	62	129	95	132	177	49	-
2000	665	2	-	1	7	24	73	120	96	122	169	51	-
1999	639	2	1	-	3	24	75	103	103	118	154	56	-
Female													
2001	510	-	-	2	3	12	55	57	65	115	139	62	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Acute and subacute endocarditis (I33)													
All origins													
Female													
2000	507	1	1	2	6	16	43	66	69	102	133	68	-
1999	477	-	-	-	4	6	39	40	68	122	135	63	-
Hispanic													
Both sexes													
2001	56	-	-	1	1	6	8	10	9	9	10	2	-
2000	70	-	1	1	-	6	11	16	8	18	7	2	-
1999	63	-	1	-	2	6	10	7	10	13	12	2	-
Male													
2001	34	-	-	-	1	5	6	8	4	5	4	1	-
2000	39	-	-	-	-	4	7	10	4	11	2	1	-
1999	42	-	1	-	-	5	6	6	5	9	8	2	-
Female													
2001	22	-	-	1	-	1	2	2	5	4	6	1	-
2000	31	-	1	1	-	2	4	6	4	7	5	1	-
1999	21	-	-	-	2	1	4	1	5	4	4	-	-
Non-Hispanic [1]													
Both sexes													
2001	1,117	1	-	2	7	23	108	174	150	238	305	109	-
2000	1,096	3	-	2	13	34	102	170	157	205	293	117	-
1999	1,049	2	-	-	5	23	104	135	160	226	277	117	-
Male													
2001	631	1	-	1	4	12	55	120	90	127	173	48	-
2000	625	2	-	1	7	20	65	110	92	111	167	50	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Acute and subacute endocarditis (I33)													
Non-Hispanic [1]													
Male													
1999	594	2	-	-	3	18	69	96	97	109	146	54	-
Female													
2001	486	-	-	1	3	11	53	54	60	111	132	61	-
2000	471	1	-	1	6	14	37	60	65	94	126	67	-
1999	455	-	-	-	2	5	35	39	63	117	131	63	-
Non-Hispanic white													
Both sexes													
2001	859	1	-	2	5	15	65	114	107	182	265	103	-
2000	852	1	-	1	6	21	66	106	114	173	261	103	-
1999	814	1	-	-	4	17	59	82	107	182	252	110	-
Male													
2001	482	1	-	1	3	7	34	79	65	93	154	45	-
2000	496	1	-	-	3	15	42	73	72	93	151	46	-
1999	466	1	-	-	2	12	39	59	68	93	139	53	-
Female													
2001	377	-	-	1	2	8	31	35	42	89	111	58	-
2000	356	-	-	1	3	6	24	33	42	80	110	57	-
1999	348	-	-	-	2	5	20	23	39	89	113	57	-
Non-Hispanic black													
Both sexes													
2001	233	-	-	-	1	8	41	57	41	43	37	5	-
2000	213	2	-	-	6	11	31	57	39	30	26	11	-
1999	208	-	-	-	1	6	41	48	47	42	17	6	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Acute and subacute endocarditis (I33)													
Non-Hispanic black													
Male													
2001	134	-	-	-	-	5	20	39	24	26	18	2	-
2000	111	1	-	-	4	3	21	35	18	16	11	2	-
1999	115	-	-	-	1	6	27	34	26	15	5	1	-
Female													
2001	99	-	-	-	1	3	21	18	17	17	19	3	-
2000	102	1	-	-	2	8	10	22	21	14	15	9	-
1999	93	-	-	-	-	-	14	14	21	27	12	5	-
Origin not stated [2]													
Both sexes													
2001	5	-	-	-	-	-	1	2	1	-	1	-	-
2000	6	-	-	-	-	-	3	-	-	1	2	-	-
1999	4	-	-	-	-	1	-	1	1	1	-	-	-
Male													
2001	3	-	-	-	-	-	1	1	1	-	-	-	-
2000	1	-	-	-	-	-	1	-	-	-	-	-	-
1999	3	-	-	-	-	1	-	1	1	-	-	-	-
Female													
2001	2	-	-	-	-	-	-	1	-	-	1	-	-
2000	5	-	-	-	-	-	2	-	-	1	2	-	-
1999	1	-	-	-	-	-	-	-	-	1	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)													
All origins													
Both sexes													
2001	839	32	26	21	40	50	93	111	106	120	154	85	1
2000	767	20	24	20	38	53	86	85	89	131	145	76	-
1999	890	27	25	21	32	50	114	105	125	146	160	85	-
Male													
2001	401	18	15	7	22	33	41	59	61	59	61	24	1
2000	404	10	13	9	21	29	58	46	52	67	73	26	-
1999	442	13	14	7	20	34	64	56	70	69	65	30	-
Female													
2001	438	14	11	14	18	17	52	52	45	61	93	61	-
2000	363	10	11	11	17	24	28	39	37	64	72	50	-
1999	448	14	11	14	12	16	50	49	55	77	95	55	-
Hispanic													
Both sexes													
2001	58	2	4	2	4	11	5	6	6	6	7	5	-
2000	61	5	4	1	5	9	10	4	9	5	7	2	-
1999	64	2	3	4	10	8	9	9	2	7	8	2	-
Male													
2001	25	1	2	1	2	6	3	1	3	3	2	1	-
2000	29	2	4	-	1	4	6	2	3	-	6	1	-
1999	30	1	1	4	5	6	7	2	1	1	2	-	-
Female													
2001	33	1	2	1	2	5	2	5	3	3	5	4	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)													
Hispanic													
Female													
2000	32	3	-	1	4	5	4	2	6	5	1	1	-
1999	34	1	2	-	5	2	2	7	1	6	6	2	-
Non-Hispanic [1]													
Both sexes													
2001	779	30	22	19	36	38	87	105	100	114	147	80	1
2000	700	15	19	19	32	44	76	81	78	126	136	74	-
1999	824	25	22	17	22	41	104	96	123	139	152	83	-
Male													
2001	375	17	13	6	20	26	38	58	58	56	59	23	1
2000	374	8	9	9	20	25	52	44	48	67	67	25	-
1999	410	12	13	3	15	27	56	54	69	68	63	30	-
Female													
2001	404	13	9	13	16	12	49	47	42	58	88	57	-
2000	326	7	10	10	12	19	24	37	30	59	69	49	-
1999	414	13	9	14	7	14	48	42	54	71	89	53	-
Non-Hispanic white													
Both sexes													
2001	600	13	10	14	23	26	55	75	79	98	134	72	1
2000	528	6	11	9	20	25	49	55	62	103	122	66	-
1999	639	10	10	9	14	26	70	73	104	115	134	74	-
Male													
2001	286	10	6	5	10	18	27	41	45	48	53	22	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)													
Non-Hispanic white													
Male													
2000	280	4	7	5	12	14	34	30	41	52	59	22	-
1999	320	6	4	2	10	17	40	40	60	56	58	27	-
Female													
2001	314	3	4	9	13	8	28	34	34	50	81	50	-
2000	248	2	4	4	8	11	15	25	21	51	63	44	-
1999	319	4	6	7	4	9	30	33	44	59	76	47	-
Non-Hispanic black													
Both sexes													
2001	157	14	12	4	13	10	24	30	19	13	11	7	-
2000	140	7	7	8	7	16	24	21	15	18	10	7	-
1999	158	10	11	8	6	14	30	20	17	20	16	6	-
Male													
2001	78	5	7	1	10	7	7	17	11	8	5	-	-
2000	72	4	2	3	3	10	16	10	6	11	5	2	-
1999	76	4	9	1	3	10	13	12	9	11	4	-	-
Female													
2001	79	9	5	3	3	3	17	13	8	5	6	7	-
2000	68	3	5	5	4	6	8	11	9	7	5	5	-
1999	82	6	2	7	3	4	17	8	8	9	12	6	-
Origin not stated [2]													
Both sexes													
2001	2	-	-	-	-	1	1	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)													
Origin not stated [2]													
Both sexes													
2000	6	-	1	-	1	-	-	-	2	-	2	-	-
1999	2	-	-	-	-	1	1	-	-	-	-	-	-
Male													
2001	1	-	-	-	-	1	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	1	-	-	-	-
1999	2	-	-	-	-	1	1	-	-	-	-	-	-
Female													
2001	1	-	-	-	-	-	1	-	-	-	-	-	-
2000	5	-	1	-	1	-	-	-	1	-	2	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Heart failure (I50)													
All origins													
Both sexes													
2001	56,934	24	17	15	34	92	307	954	2,374	6,274	17,134	29,708	1
2000	55,704	31	9	21	34	63	262	811	2,211	6,403	16,820	29,035	4
1999	54,913	11	10	12	31	65	272	782	2,107	6,524	16,771	28,327	1
Male													
2001	21,632	16	5	8	16	56	182	588	1,347	3,344	7,672	8,398	-
2000	21,175	22	7	9	20	36	159	496	1,264	3,414	7,498	8,248	2
1999	20,755	5	7	6	23	32	164	486	1,228	3,526	7,318	7,959	1
Female													
2001	35,302	8	12	7	18	36	125	366	1,027	2,930	9,462	21,310	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Heart failure (I50)													
All origins													
Female													
2000	34,529	9	2	12	14	27	103	315	947	2,989	9,322	20,787	2
1999	34,158	6	3	6	8	33	108	296	879	2,998	9,453	20,368	-
Hispanic													
Both sexes													
2001	1,364	8	3	-	1	9	21	43	103	213	404	559	-
2000	1,270	7	1	2	-	3	10	22	100	203	365	557	-
1999	1,242	3	4	1	3	4	18	29	87	202	353	538	-
Male													
2001	565	6	1	-	1	6	15	31	52	109	169	175	-
2000	535	5	1	1	-	2	6	9	62	109	151	189	-
1999	522	1	3	-	3	3	12	16	47	112	159	166	-
Female													
2001	799	2	2	-	-	3	6	12	51	104	235	384	-
2000	735	2	-	1	-	1	4	13	38	94	214	368	-
1999	720	2	1	1	-	1	6	13	40	90	194	372	-
Non-Hispanic [1]													
Both sexes													
2001	55,437	16	14	15	33	82	286	906	2,258	6,044	16,688	29,094	1
2000	54,307	24	8	19	34	60	252	789	2,105	6,185	16,417	28,411	3
1999	53,528	8	6	11	28	61	253	750	2,007	6,308	16,377	27,718	1
Male													
2001	21,012	10	4	8	15	49	167	555	1,287	3,225	7,489	8,203	-
2000	20,599	17	6	8	20	34	153	487	1,197	3,296	7,335	8,045	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Heart failure (I50)													
Non-Hispanic [1]													
Male													
1999	20,170	4	4	6	20	29	152	469	1,173	3,406	7,140	7,766	1
Female													
2001	34,425	6	10	7	18	33	119	351	971	2,819	9,199	20,891	1
2000	33,708	7	2	11	14	26	99	302	908	2,889	9,082	20,366	2
1999	33,358	4	2	5	8	32	101	281	834	2,902	9,237	19,952	-
Non-Hispanic white													
Both sexes													
2001	49,788	11	12	11	17	49	167	608	1,730	5,052	15,033	27,097	1
2000	48,782	13	4	14	17	35	154	536	1,611	5,176	14,827	26,392	3
1999	47,977	5	4	8	15	30	156	480	1,537	5,302	14,702	25,738	-
Male													
2001	18,730	6	4	5	9	25	103	392	1,019	2,734	6,785	7,648	-
2000	18,386	10	4	6	10	20	103	342	942	2,828	6,671	7,449	1
1999	17,903	2	2	5	11	14	100	306	922	2,920	6,472	7,149	-
Female													
2001	31,058	5	8	6	8	24	64	216	711	2,318	8,248	19,449	1
2000	30,396	3	-	8	7	15	51	194	669	2,348	8,156	18,943	2
1999	30,074	3	2	3	4	16	56	174	615	2,382	8,230	18,589	-
Non-Hispanic black													
Both sexes													
2001	5,094	5	2	3	16	32	112	284	494	905	1,475	1,766	-
2000	4,936	11	4	4	13	24	94	241	453	909	1,407	1,776	-
1999	4,961	3	2	3	12	30	93	257	446	922	1,468	1,724	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Heart failure (I50)													
Non-Hispanic black													
Male													
2001	2,036	4	-	2	6	23	61	153	248	447	618	474	-
2000	1,942	7	2	2	8	13	47	138	231	420	572	502	-
1999	2,005	2	2	1	8	15	50	156	238	442	577	513	1
Female													
2001	3,058	1	2	1	10	9	51	131	246	458	857	1,292	-
2000	2,994	4	2	2	5	11	47	103	222	489	835	1,274	-
1999	2,956	1	-	2	4	15	43	101	208	480	891	1,211	-
Origin not stated [2]													
Both sexes													
2001	133	-	-	-	-	1	-	5	13	17	42	55	-
2000	127	-	-	-	-	-	-	-	6	15	38	67	1
1999	143	-	-	-	-	-	1	3	13	14	41	71	-
Male													
2001	55	-	-	-	-	1	-	2	8	10	14	20	-
2000	41	-	-	-	-	-	-	-	5	9	12	14	1
1999	63	-	-	-	-	-	-	1	8	8	19	27	-
Female													
2001	78	-	-	-	-	-	-	3	5	7	28	35	-
2000	86	-	-	-	-	-	-	-	1	6	26	53	-
1999	80	-	-	-	-	-	1	2	5	6	22	44	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)													
All origins													
Both sexes													
2001	107,998	392	172	207	712	1,565	3,955	7,233	9,895	17,415	31,372	35,075	5
2000	107,785	403	134	204	768	1,593	3,993	7,005	9,897	17,921	31,100	34,763	4
1999	108,909	450	141	215	810	1,634	4,147	6,858	10,177	18,662	31,492	34,317	6
Male													
2001	49,738	187	97	107	457	931	2,492	4,531	6,000	9,632	14,630	10,670	4
2000	49,878	204	78	131	470	960	2,456	4,489	6,120	10,018	14,266	10,683	3
1999	50,765	235	78	124	487	977	2,591	4,334	6,299	10,607	14,591	10,438	4
Female													
2001	58,260	205	75	100	255	634	1,463	2,702	3,895	7,783	16,742	24,405	1
2000	57,907	199	56	73	298	633	1,537	2,516	3,777	7,903	16,834	24,080	1
1999	58,144	215	63	91	323	657	1,556	2,524	3,878	8,055	16,901	23,879	2
Hispanic													
Both sexes													
2001	3,421	65	24	30	80	153	234	311	398	628	785	711	2
2000	3,391	67	22	33	105	164	252	321	376	591	770	689	1
1999	3,318	87	25	29	84	158	272	300	402	593	729	638	1
Male													
2001	1,777	33	11	16	53	102	161	212	237	339	381	230	2
2000	1,790	32	15	24	75	114	158	213	230	326	351	251	1
1999	1,798	38	15	17	52	114	183	201	245	344	344	244	1
Female													
2001	1,644	32	13	14	27	51	73	99	161	289	404	481	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)													
Hispanic													
Female													
2000	1,601	35	7	9	30	50	94	108	146	265	419	438	-
1999	1,520	49	10	12	32	44	89	99	157	249	385	394	-
Non-Hispanic [1]													
Both sexes													
2001	104,292	325	147	177	625	1,407	3,703	6,892	9,472	16,741	30,521	34,281	1
2000	104,119	333	112	170	662	1,420	3,721	6,654	9,483	17,288	30,278	33,995	3
1999	105,310	362	115	185	724	1,469	3,856	6,539	9,740	18,012	30,690	33,614	4
Male													
2001	47,812	153	86	91	400	826	2,316	4,297	5,746	9,263	14,220	10,413	1
2000	47,949	170	63	107	394	842	2,284	4,259	5,866	9,664	13,891	10,407	2
1999	48,810	197	63	106	433	859	2,395	4,118	6,031	10,229	14,206	10,171	2
Female													
2001	56,480	172	61	86	225	581	1,387	2,595	3,726	7,478	16,301	23,868	-
2000	56,170	163	49	63	268	578	1,437	2,395	3,617	7,624	16,387	23,588	1
1999	56,500	165	52	79	291	610	1,461	2,421	3,709	7,783	16,484	23,443	2
Non-Hispanic white													
Both sexes													
2001	88,469	194	81	113	377	847	2,409	4,834	7,216	13,839	27,054	31,504	1
2000	88,583	218	72	98	385	867	2,448	4,665	7,302	14,329	26,964	31,233	2
1999	89,439	224	73	123	462	899	2,529	4,603	7,360	14,930	27,318	30,915	3
Male													
2001	40,015	99	48	61	239	505	1,567	3,125	4,433	7,778	12,635	9,524	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)													
Non-Hispanic white													
Male													
2000	40,299	112	40	66	232	527	1,547	3,060	4,593	8,129	12,454	9,537	2
1999	40,860	127	45	66	268	537	1,630	2,990	4,630	8,587	12,696	9,282	2
Female													
2001	48,454	95	33	52	138	342	842	1,709	2,783	6,061	14,419	21,980	-
2000	48,284	106	32	32	153	340	901	1,605	2,709	6,200	14,510	21,696	-
1999	48,579	97	28	57	194	362	899	1,613	2,730	6,343	14,622	21,633	1
Non-Hispanic black													
Both sexes													
2001	14,012	111	56	56	222	517	1,193	1,893	2,054	2,561	2,985	2,364	-
2000	13,870	93	35	60	252	508	1,165	1,849	1,978	2,659	2,915	2,355	1
1999	14,204	119	35	55	243	526	1,231	1,782	2,168	2,763	2,953	2,328	1
Male													
2001	6,822	50	32	28	144	292	680	1,070	1,200	1,285	1,337	704	-
2000	6,758	44	20	34	143	287	663	1,112	1,155	1,355	1,245	700	-
1999	7,062	60	16	36	154	300	707	1,045	1,272	1,468	1,284	720	-
Female													
2001	7,190	61	24	28	78	225	513	823	854	1,276	1,648	1,660	-
2000	7,112	49	15	26	109	221	502	737	823	1,304	1,670	1,655	1
1999	7,142	59	19	19	89	226	524	737	896	1,295	1,669	1,608	1
Origin not stated [2]													
Both sexes													
2001	285	2	1	-	7	5	18	30	25	46	66	83	2

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)													
Origin not stated [2]													
Both sexes													
2000	275	3	-	1	1	9	20	30	38	42	52	79	-
1999	281	1	1	1	2	7	19	19	35	57	73	65	1
Male													
2001	149	1	-	-	4	3	15	22	17	30	29	27	1
2000	139	2	-	-	1	4	14	17	24	28	24	25	-
1999	157	-	-	1	2	4	13	15	23	34	41	23	1
Female													
2001	136	1	1	-	3	2	3	8	8	16	37	56	1
2000	136	1	-	1	-	5	6	13	14	14	28	54	-
1999	124	1	1	-	-	3	6	4	12	23	32	42	-
Essential (primary) hypertension and hypertensive renal disease (I10,I12)													
All origins													
Both sexes													
2001	19,250	-	1	1	23	104	331	931	1,462	2,832	5,993	7,572	-
2000	18,073	1	1	4	17	70	348	878	1,440	2,784	5,623	6,907	-
1999	16,968	3	3	1	13	78	313	799	1,318	2,799	5,324	6,317	-
Male													
2001	7,254	-	-	-	16	59	206	569	784	1,379	2,282	1,959	-
2000	6,895	1	-	3	4	39	216	555	818	1,308	2,204	1,747	-
1999	6,488	2	1	1	8	48	195	482	732	1,372	2,067	1,580	-
Female													
2001	11,996	-	1	1	7	45	125	362	678	1,453	3,711	5,613	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Essential (primary) hypertension and hypertensive renal disease (I10,I12)													
All origins													
Female													
2000	11,178	-	1	1	13	31	132	323	622	1,476	3,419	5,160	-
1999	10,480	1	2	-	5	30	118	317	586	1,427	3,257	4,737	-
Hispanic													
Both sexes													
2001	889	-	-	-	3	10	19	60	83	187	271	256	-
2000	821	1	1	-	4	7	27	55	109	162	232	223	-
1999	753	1	1	-	3	8	19	64	82	158	232	185	-
Male													
2001	365	-	-	-	2	8	16	39	34	84	114	68	-
2000	386	1	-	-	2	5	20	44	65	72	107	70	-
1999	340	-	-	-	2	5	14	35	40	84	99	61	-
Female													
2001	524	-	-	-	1	2	3	21	49	103	157	188	-
2000	435	-	1	-	2	2	7	11	44	90	125	153	-
1999	413	1	1	-	1	3	5	29	42	74	133	124	-
Non-Hispanic [1]													
Both sexes													
2001	18,306	-	1	1	20	94	310	868	1,370	2,636	5,706	7,300	-
2000	17,182	-	-	4	13	63	318	815	1,320	2,616	5,373	6,660	-
1999	16,158	2	2	1	10	69	292	734	1,232	2,633	5,073	6,110	-
Male													
2001	6,856	-	-	-	14	51	188	528	743	1,288	2,160	1,884	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Essential (primary) hypertension and hypertensive renal disease (I10,I12)													
Non-Hispanic [1]													
Male													
2000	6,475	-	-	3	2	34	194	504	746	1,232	2,088	1,672	-
1999	6,123	2	1	1	6	42	179	447	689	1,285	1,958	1,513	-
Female													
2001	11,450	-	1	1	6	43	122	340	627	1,348	3,546	5,416	-
2000	10,707	-	-	1	11	29	124	311	574	1,384	3,285	4,988	-
1999	10,035	-	1	-	4	27	113	287	543	1,348	3,115	4,597	-
Non-Hispanic white													
Both sexes													
2001	13,837	-	1	1	11	35	121	404	776	1,747	4,499	6,242	-
2000	12,917	-	-	3	3	25	132	381	736	1,744	4,252	5,641	-
1999	12,048	1	-	-	5	28	112	343	686	1,788	3,940	5,145	-
Male													
2001	4,970	-	-	-	7	20	77	253	439	873	1,701	1,600	-
2000	4,721	-	-	3	1	17	89	255	434	852	1,671	1,399	-
1999	4,395	1	-	-	2	22	77	220	410	897	1,526	1,240	-
Female													
2001	8,867	-	1	1	4	15	44	151	337	874	2,798	4,642	-
2000	8,196	-	-	-	2	8	43	126	302	892	2,581	4,242	-
1999	7,653	-	-	-	3	6	35	123	276	891	2,414	3,905	-
Non-Hispanic black													
Both sexes													
2001	4,012	-	-	-	9	55	177	439	553	789	1,056	934	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Essential (primary) hypertension and hypertensive renal disease (I10,I12)													
Non-Hispanic black													
Both sexes													
2000	3,862	-	-	1	10	35	176	405	555	801	980	899	-
1999	3,708	1	1	1	5	37	175	370	506	764	1,006	842	-
Male													
2001	1,698	-	-	-	7	29	102	263	284	374	399	240	-
2000	1,576	-	-	-	1	16	98	229	301	344	358	229	-
1999	1,550	1	1	1	4	17	97	218	256	348	381	226	-
Female													
2001	2,314	-	-	-	2	26	75	176	269	415	657	694	-
2000	2,286	-	-	1	9	19	78	176	254	457	622	670	-
1999	2,158	-	-	-	1	20	78	152	250	416	625	616	-
Origin not stated [2]													
Both sexes													
2001	55	-	-	-	-	-	2	3	9	9	16	16	-
2000	70	-	-	-	-	-	3	8	11	6	18	24	-
1999	57	-	-	-	-	1	2	1	4	8	19	22	-
Male													
2001	33	-	-	-	-	-	2	2	7	7	8	7	-
2000	34	-	-	-	-	-	2	7	7	4	9	5	-
1999	25	-	-	-	-	1	2	-	3	3	10	6	-
Female													
2001	22	-	-	-	-	-	-	1	2	2	8	9	-
2000	36	-	-	-	-	-	1	1	4	2	9	19	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Essential (primary) hypertension and hypertensive renal disease (I10,I12)													
Origin not stated [2]													
Female													
1999	32	-	-	-	-	-	-	1	1	5	9	16	-
Cerebrovascular diseases (I60-I69)													
All origins													
Both sexes													
2001	163,538	108	54	80	196	601	2,491	5,910	9,608	22,598	55,822	66,066	4
2000	167,661	124	47	76	199	602	2,599	6,011	9,956	23,649	57,020	67,376	2
1999	167,366	103	46	65	182	580	2,574	5,563	9,652	24,092	57,427	67,080	2
Male													
2001	63,177	63	25	42	103	311	1,267	3,205	5,277	11,649	23,301	17,930	4
2000	64,769	74	19	33	106	308	1,297	3,235	5,496	12,042	23,946	18,212	1
1999	64,485	65	22	34	95	299	1,308	2,996	5,305	12,220	24,088	18,052	1
Female													
2001	100,361	45	29	38	93	290	1,224	2,705	4,331	10,949	32,521	48,136	-
2000	102,892	50	28	43	93	294	1,302	2,776	4,460	11,607	33,074	49,164	1
1999	102,881	38	24	31	87	281	1,266	2,567	4,347	11,872	33,339	49,028	1
Hispanic													
Both sexes													
2001	6,416	17	10	16	40	99	275	539	654	1,206	1,892	1,668	-
2000	6,187	16	6	8	31	108	267	477	681	1,219	1,742	1,632	-
1999	5,907	16	7	9	32	102	241	464	617	1,211	1,672	1,536	-
Male													
2001	2,982	9	4	10	26	60	151	315	378	637	864	528	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Cerebrovascular diseases (I60-I69)													
Hispanic													
Male													
2000	2,865	8	-	3	17	65	145	280	392	645	799	511	-
1999	2,809	8	1	6	21	62	146	285	354	641	758	527	-
Female													
2001	3,434	8	6	6	14	39	124	224	276	569	1,028	1,140	-
2000	3,322	8	6	5	14	43	122	197	289	574	943	1,121	-
1999	3,098	8	6	3	11	40	95	179	263	570	914	1,009	-
Non-Hispanic [1]													
Both sexes													
2001	156,741	91	43	63	156	498	2,211	5,350	8,916	21,346	53,810	64,253	4
2000	161,057	108	41	67	167	493	2,323	5,509	9,238	22,373	55,129	65,608	1
1999	161,057	87	39	56	150	477	2,323	5,071	8,991	22,814	55,643	65,404	2
Male													
2001	60,030	54	21	32	77	247	1,111	2,882	4,872	10,989	22,377	17,364	4
2000	61,715	66	19	30	89	243	1,145	2,936	5,079	11,364	23,084	17,660	-
1999	61,499	57	21	28	74	237	1,156	2,697	4,922	11,542	23,275	17,489	1
Female													
2001	96,711	37	22	31	79	251	1,100	2,468	4,044	10,357	31,433	46,889	-
2000	99,342	42	22	37	78	250	1,178	2,573	4,159	11,009	32,045	47,948	1
1999	99,558	30	18	28	76	240	1,167	2,374	4,069	11,272	32,368	47,915	1
Non-Hispanic white													
Both sexes													
2001	133,879	46	28	42	98	320	1,333	3,200	6,208	16,838	47,304	58,461	1
2000	138,210	59	25	46	118	313	1,378	3,433	6,555	17,989	48,556	59,738	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Cerebrovascular diseases (I60-I69)													
Non-Hispanic white													
Both sexes													
1999	138,716	55	27	40	105	301	1,406	3,143	6,311	18,332	49,178	59,818	-
Male													
2001	50,382	27	14	24	55	161	684	1,745	3,400	8,753	19,762	15,756	1
2000	52,000	38	14	23	67	161	694	1,856	3,605	9,176	20,379	15,987	-
1999	51,968	36	15	21	51	151	712	1,662	3,479	9,337	20,618	15,886	-
Female													
2001	83,497	19	14	18	43	159	649	1,455	2,808	8,085	27,542	42,705	-
2000	86,210	21	11	23	51	152	684	1,577	2,950	8,813	28,177	43,751	-
1999	86,748	19	12	19	54	150	694	1,481	2,832	8,995	28,560	43,932	-
Non-Hispanic black													
Both sexes													
2001	18,844	42	12	14	51	153	758	1,886	2,289	3,684	5,257	4,695	3
2000	19,053	46	13	15	42	158	829	1,819	2,265	3,660	5,344	4,861	1
1999	18,731	27	10	15	40	149	808	1,662	2,269	3,792	5,314	4,644	1
Male													
2001	7,828	26	5	5	20	75	355	998	1,252	1,844	2,068	1,177	3
2000	7,945	27	3	5	19	70	394	955	1,240	1,817	2,155	1,260	-
1999	7,823	17	4	7	19	72	387	900	1,235	1,857	2,100	1,225	-
Female													
2001	11,016	16	7	9	31	78	403	888	1,037	1,840	3,189	3,518	-
2000	11,108	19	10	10	23	88	435	864	1,025	1,843	3,189	3,601	1
1999	10,908	10	6	8	21	77	421	762	1,034	1,935	3,214	3,419	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Cerebrovascular diseases (I60-I69)													
Origin not stated [2]													
Both sexes													
2001	381	-	1	1	-	4	5	21	38	46	120	145	-
2000	417	-	-	1	1	1	9	25	37	57	149	136	1
1999	402	-	-	-	-	1	10	28	44	67	112	140	-
Male													
2001	165	-	-	-	-	4	5	8	27	23	60	38	-
2000	189	-	-	-	-	-	7	19	25	33	63	41	1
1999	177	-	-	-	-	-	6	14	29	37	55	36	-
Female													
2001	216	-	1	1	-	-	-	13	11	23	60	107	-
2000	228	-	-	1	1	1	2	6	12	24	86	95	-
1999	225	-	-	-	-	1	4	14	15	30	57	104	-
Atherosclerosis (I70)													
All origins													
Both sexes													
2001	14,086	1	1	1	1	3	40	186	473	1,335	4,156	7,888	1
2000	14,393	2	-	-	3	8	48	176	451	1,423	4,277	8,005	-
1999	14,979	2	-	-	2	8	45	185	486	1,476	4,385	8,390	-
Male													
2001	5,143	1	1	1	1	2	26	118	286	743	1,791	2,172	1
2000	5,357	-	-	-	2	6	35	119	294	800	1,935	2,166	-
1999	5,443	1	-	-	-	5	32	121	307	795	1,927	2,255	-
Female													
2001	8,943	-	-	-	-	1	14	68	187	592	2,365	5,716	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Atherosclerosis (I70)													
All origins													
Female													
2000	9,036	2	-	-	1	2	13	57	157	623	2,342	5,839	-
1999	9,536	1	-	-	2	3	13	64	179	681	2,458	6,135	-
Hispanic													
Both sexes													
2001	416	-	-	-	-	-	-	14	19	46	128	209	-
2000	414	-	-	-	-	-	2	9	16	63	124	200	-
1999	410	-	-	-	-	-	3	14	25	54	130	184	-
Male													
2001	172	-	-	-	-	-	-	9	11	24	61	67	-
2000	173	-	-	-	-	-	2	7	8	39	65	52	-
1999	185	-	-	-	-	-	1	7	16	31	60	70	-
Female													
2001	244	-	-	-	-	-	-	5	8	22	67	142	-
2000	241	-	-	-	-	-	-	2	8	24	59	148	-
1999	225	-	-	-	-	-	2	7	9	23	70	114	-
Non-Hispanic [1]													
Both sexes													
2001	13,626	1	1	1	1	3	40	172	452	1,285	4,011	7,658	1
2000	13,945	2	-	-	3	8	46	167	434	1,357	4,145	7,783	-
1999	14,519	2	-	-	2	8	42	169	457	1,416	4,242	8,181	-
Male													
2001	4,952	1	1	1	1	2	26	109	274	718	1,721	2,097	1
2000	5,166	-	-	-	2	6	33	112	285	759	1,865	2,104	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Atherosclerosis (I70)													
Non-Hispanic [1]													
Male													
1999	5,236	1	-	-	-	5	31	112	287	761	1,861	2,178	-
Female													
2001	8,674	-	-	-	-	1	14	63	178	567	2,290	5,561	-
2000	8,779	2	-	-	1	2	13	55	149	598	2,280	5,679	-
1999	9,283	1	-	-	2	3	11	57	170	655	2,381	6,003	-
Non-Hispanic white													
Both sexes													
2001	12,517	1	1	1	-	2	33	137	361	1,119	3,675	7,186	1
2000	12,740	2	-	-	3	5	34	114	349	1,181	3,782	7,270	-
1999	13,286	1	-	-	2	6	30	135	374	1,241	3,880	7,617	-
Male													
2001	4,511	1	1	1	-	2	21	85	225	631	1,570	1,973	1
2000	4,688	-	-	-	2	5	26	79	238	673	1,716	1,949	-
1999	4,727	1	-	-	-	4	23	93	235	667	1,690	2,014	-
Female													
2001	8,006	-	-	-	-	-	12	52	136	488	2,105	5,213	-
2000	8,052	2	-	-	1	-	8	35	111	508	2,066	5,321	-
1999	8,559	-	-	-	2	2	7	42	139	574	2,190	5,603	-
Non-Hispanic black													
Both sexes													
2001	957	-	-	-	1	1	6	31	84	145	285	404	-
2000	1,049	-	-	-	-	3	12	47	79	147	320	441	-
1999	1,061	1	-	-	-	2	9	29	76	150	312	482	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Atherosclerosis (I70)													
Non-Hispanic black													
Male													
2001	365	-	-	-	1	-	4	21	43	74	124	98	-
2000	412	-	-	-	-	1	7	28	43	74	131	128	-
1999	430	-	-	-	-	1	6	15	46	80	142	140	-
Female													
2001	592	-	-	-	-	1	2	10	41	71	161	306	-
2000	637	-	-	-	-	2	5	19	36	73	189	313	-
1999	631	1	-	-	-	1	3	14	30	70	170	342	-
Origin not stated [2]													
Both sexes													
2001	44	-	-	-	-	-	-	-	2	4	17	21	-
2000	34	-	-	-	-	-	-	-	1	3	8	22	-
1999	50	-	-	-	-	-	-	2	4	6	13	25	-
Male													
2001	19	-	-	-	-	-	-	-	1	1	9	8	-
2000	18	-	-	-	-	-	-	-	1	2	5	10	-
1999	22	-	-	-	-	-	-	2	4	3	6	7	-
Female													
2001	25	-	-	-	-	-	-	-	1	3	8	13	-
2000	16	-	-	-	-	-	-	-	-	1	3	12	-
1999	28	-	-	-	-	-	-	-	-	3	7	18	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other diseases of circulatory system (I71-I78)													
All origins													
Both sexes													
2001	25,318	12	4	11	61	170	498	1,024	2,195	5,345	9,032	6,965	1
2000	26,036	16	5	11	59	154	448	1,038	2,236	5,926	9,222	6,920	1
1999	25,809	12	2	11	49	141	478	971	2,210	5,808	9,264	6,863	-
Male													
2001	13,396	8	2	3	40	115	335	688	1,454	3,348	4,839	2,563	1
2000	14,084	6	2	7	33	103	276	699	1,530	3,771	5,019	2,637	1
1999	13,794	9	-	7	32	102	296	657	1,519	3,681	5,015	2,476	-
Female													
2001	11,922	4	2	8	21	55	163	336	741	1,997	4,193	4,402	-
2000	11,952	10	3	4	26	51	172	339	706	2,155	4,203	4,283	-
1999	12,015	3	2	4	17	39	182	314	691	2,127	4,249	4,387	-
Hispanic													
Both sexes													
2001	754	2	1	2	7	19	37	73	88	156	216	152	1
2000	818	3	1	1	9	24	38	56	99	217	215	155	-
1999	729	2	-	2	9	12	33	38	94	170	211	158	-
Male													
2001	424	1	1	-	4	14	26	46	55	100	119	57	1
2000	452	1	-	1	4	14	22	33	64	122	126	65	-
1999	416	1	-	-	7	6	22	25	59	101	119	76	-
Female													
2001	330	1	-	2	3	5	11	27	33	56	97	95	-
2000	366	2	1	-	5	10	16	23	35	95	89	90	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other diseases of circulatory system (I71-I78)													
Hispanic													
Female													
1999	313	1	-	2	2	6	11	13	35	69	92	82	-
Non-Hispanic [1]													
Both sexes													
2001	24,489	10	3	9	54	149	460	946	2,099	5,170	8,795	6,794	-
2000	25,147	13	4	10	50	129	405	979	2,129	5,695	8,985	6,747	1
1999	25,023	10	2	9	40	129	445	932	2,109	5,619	9,037	6,691	-
Male													
2001	12,926	7	1	3	36	99	308	638	1,392	3,234	4,708	2,500	-
2000	13,593	5	2	6	29	89	249	663	1,461	3,640	4,883	2,565	1
1999	13,351	8	-	7	25	96	274	632	1,456	3,567	4,889	2,397	-
Female													
2001	11,563	3	2	6	18	50	152	308	707	1,936	4,087	4,294	-
2000	11,554	8	2	4	21	40	156	316	668	2,055	4,102	4,182	-
1999	11,672	2	2	2	15	33	171	300	653	2,052	4,148	4,294	-
Non-Hispanic white													
Both sexes													
2001	21,439	8	2	8	40	108	303	682	1,702	4,539	7,908	6,139	-
2000	22,113	11	3	6	35	88	269	699	1,734	5,046	8,133	6,089	-
1999	22,191	8	2	7	33	94	323	690	1,773	5,000	8,210	6,051	-
Male													
2001	11,437	5	1	3	27	74	202	462	1,158	2,897	4,303	2,305	-
2000	12,138	3	2	5	22	63	167	494	1,225	3,301	4,492	2,364	-
1999	12,014	6	-	5	20	71	204	478	1,261	3,253	4,514	2,202	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other diseases of circulatory system (I71-I78)													
Non-Hispanic white													
Female													
2001	10,002	3	1	5	13	34	101	220	544	1,642	3,605	3,834	-
2000	9,975	8	1	1	13	25	102	205	509	1,745	3,641	3,725	-
1999	10,177	2	2	2	13	23	119	212	512	1,747	3,696	3,849	-
Non-Hispanic black													
Both sexes													
2001	2,606	2	1	1	12	39	141	227	348	526	746	563	-
2000	2,569	2	-	4	13	35	120	232	346	540	710	566	1
1999	2,416	2	-	-	7	30	112	212	290	517	689	557	-
Male													
2001	1,214	2	-	-	7	24	93	149	197	272	324	146	-
2000	1,193	2	-	1	6	22	73	144	198	279	314	153	1
1999	1,102	2	-	-	5	23	61	135	165	266	290	155	-
Female													
2001	1,392	-	1	1	5	15	48	78	151	254	422	417	-
2000	1,376	-	-	3	7	13	47	88	148	261	396	413	-
1999	1,314	-	-	-	2	7	51	77	125	251	399	402	-
Origin not stated [2]													
Both sexes													
2001	75	-	-	-	-	2	1	5	8	19	21	19	-
2000	71	-	-	-	-	1	5	3	8	14	22	18	-
1999	57	-	-	-	-	-	-	1	7	19	16	14	-
Male													
2001	46	-	-	-	-	2	1	4	7	14	12	6	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other diseases of circulatory system (I71-I78)													
Origin not stated [2]													
Male													
2000	39	-	-	-	-	-	5	3	5	9	10	7	-
1999	27	-	-	-	-	-	-	-	4	13	7	3	-
Female													
2001	29	-	-	-	-	-	-	1	1	5	9	13	-
2000	32	-	-	-	-	1	-	-	3	5	12	11	-
1999	30	-	-	-	-	-	-	1	3	6	9	11	-
Aortic aneurysm and dissection (I71)													
All origins													
Both sexes													
2001	15,234	-	2	4	43	117	332	661	1,394	3,517	5,751	3,412	1
2000	15,810	-	-	5	36	107	303	662	1,493	4,012	5,835	3,357	-
1999	15,807	1	1	3	31	93	329	638	1,437	4,001	5,936	3,337	-
Male													
2001	9,114	-	2	3	31	90	264	506	1,031	2,382	3,313	1,491	1
2000	9,661	-	-	3	23	78	219	492	1,111	2,730	3,443	1,562	-
1999	9,615	1	-	2	20	78	232	486	1,088	2,712	3,501	1,495	-
Female													
2001	6,120	-	-	1	12	27	68	155	363	1,135	2,438	1,921	-
2000	6,149	-	-	2	13	29	84	170	382	1,282	2,392	1,795	-
1999	6,192	-	1	1	11	15	97	152	349	1,289	2,435	1,842	-
Hispanic													
Both sexes													
2001	403	-	1	-	4	14	24	41	47	93	127	51	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Aortic aneurysm and dissection (I71)													
Hispanic													
Both sexes													
2000	465	-	-	1	8	15	21	41	58	121	137	63	-
1999	384	-	-	1	4	6	23	27	55	95	115	58	-
Male													
2001	279	-	1	-	2	14	20	31	33	71	78	28	1
2000	294	-	-	1	4	11	14	28	42	76	86	32	-
1999	256	-	-	-	3	4	17	22	41	67	65	37	-
Female													
2001	124	-	-	-	2	-	4	10	14	22	49	23	-
2000	171	-	-	-	4	4	7	13	16	45	51	31	-
1999	128	-	-	1	1	2	6	5	14	28	50	21	-
Non-Hispanic [1]													
Both sexes													
2001	14,782	-	1	4	39	102	307	617	1,341	3,408	5,610	3,353	-
2000	15,296	-	-	4	28	91	277	618	1,431	3,881	5,682	3,284	-
1999	15,393	1	1	2	27	87	306	611	1,378	3,893	5,811	3,276	-
Male													
2001	8,798	-	1	3	29	75	243	472	992	2,299	3,225	1,459	-
2000	9,338	-	-	2	19	67	200	461	1,066	2,648	3,349	1,526	-
1999	9,342	1	-	2	17	74	215	464	1,045	2,635	3,432	1,457	-
Female													
2001	5,984	-	-	1	10	27	64	145	349	1,109	2,385	1,894	-
2000	5,958	-	-	2	9	24	77	157	365	1,233	2,333	1,758	-
1999	6,051	-	1	-	10	13	91	147	333	1,258	2,379	1,819	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Aortic aneurysm and dissection (I71)													
Non-Hispanic white													
Both sexes													
2001	13,217	-	1	4	28	77	203	435	1,117	3,079	5,155	3,118	-
2000	13,722	-	-	1	18	61	182	454	1,201	3,526	5,250	3,029	-
1999	13,934	1	1	1	23	64	217	440	1,195	3,548	5,407	3,037	-
Male													
2001	7,924	-	1	3	21	58	163	342	841	2,109	3,002	1,384	-
2000	8,488	-	-	1	14	45	134	355	914	2,453	3,136	1,436	-
1999	8,551	1	-	1	14	54	155	340	928	2,451	3,243	1,364	-
Female													
2001	5,293	-	-	1	7	19	40	93	276	970	2,153	1,734	-
2000	5,234	-	-	-	4	16	48	99	287	1,073	2,114	1,593	-
1999	5,383	-	1	-	9	10	62	100	267	1,097	2,164	1,673	-
Non-Hispanic black													
Both sexes													
2001	1,238	-	-	-	10	23	91	154	189	240	352	179	-
2000	1,262	-	-	3	9	25	86	135	196	279	331	198	-
1999	1,172	-	-	-	4	21	80	152	152	271	303	189	-
Male													
2001	659	-	-	-	7	16	69	108	124	134	158	43	-
2000	661	-	-	1	5	19	60	89	123	151	153	60	-
1999	615	-	-	-	3	19	52	109	96	147	124	65	-
Female													
2001	579	-	-	-	3	7	22	46	65	106	194	136	-
2000	601	-	-	2	4	6	26	46	73	128	178	138	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Aortic aneurysm and dissection (I71)													
Non-Hispanic black													
Female													
1999	557	-	-	-	1	2	28	43	56	124	179	124	-
Origin not stated [2]													
Both sexes													
2001	49	-	-	-	-	1	1	3	6	16	14	8	-
2000	49	-	-	-	-	1	5	3	4	10	16	10	-
1999	30	-	-	-	-	-	-	-	4	13	10	3	-
Male													
2001	37	-	-	-	-	1	1	3	6	12	10	4	-
2000	29	-	-	-	-	-	5	3	3	6	8	4	-
1999	17	-	-	-	-	-	-	-	2	10	4	1	-
Female													
2001	12	-	-	-	-	-	-	-	-	4	4	4	-
2000	20	-	-	-	-	1	-	-	1	4	8	6	-
1999	13	-	-	-	-	-	-	-	2	3	6	2	-
Other diseases of arteries, arterioles and capillaries (I72-I78)													
All origins													
Both sexes													
2001	10,084	12	2	7	18	53	166	363	801	1,828	3,281	3,553	-
2000	10,226	16	5	6	23	47	145	376	743	1,914	3,387	3,563	1
1999	10,002	11	1	8	18	48	149	333	773	1,807	3,328	3,526	-
Male													
2001	4,282	8	-	-	9	25	71	182	423	966	1,526	1,072	-
2000	4,423	6	2	4	10	25	57	207	419	1,041	1,576	1,075	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)													
All origins													
Male													
1999	4,179	8	-	5	12	24	64	171	431	969	1,514	981	-
Female													
2001	5,802	4	2	7	9	28	95	181	378	862	1,755	2,481	-
2000	5,803	10	3	2	13	22	88	169	324	873	1,811	2,488	-
1999	5,823	3	1	3	6	24	85	162	342	838	1,814	2,545	-
Hispanic													
Both sexes													
2001	351	2	-	2	3	5	13	32	41	63	89	101	-
2000	353	3	1	-	1	9	17	15	41	96	78	92	-
1999	345	2	-	1	5	6	10	11	39	75	96	100	-
Male													
2001	145	1	-	-	2	-	6	15	22	29	41	29	-
2000	158	1	-	-	-	3	8	5	22	46	40	33	-
1999	160	1	-	-	4	2	5	3	18	34	54	39	-
Female													
2001	206	1	-	2	1	5	7	17	19	34	48	72	-
2000	195	2	1	-	1	6	9	10	19	50	38	59	-
1999	185	1	-	1	1	4	5	8	21	41	42	61	-
Non-Hispanic [1]													
Both sexes													
2001	9,707	10	2	5	15	47	153	329	758	1,762	3,185	3,441	-
2000	9,851	13	4	6	22	38	128	361	698	1,814	3,303	3,463	1
1999	9,630	9	1	7	13	42	139	321	731	1,726	3,226	3,415	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)													
Non-Hispanic [1]													
Male													
2001	4,128	7	-	-	7	24	65	166	400	935	1,483	1,041	-
2000	4,255	5	2	4	10	22	49	202	395	992	1,534	1,039	1
1999	4,009	7	-	5	8	22	59	168	411	932	1,457	940	-
Female													
2001	5,579	3	2	5	8	23	88	163	358	827	1,702	2,400	-
2000	5,596	8	2	2	12	16	79	159	303	822	1,769	2,424	-
1999	5,621	2	1	2	5	20	80	153	320	794	1,769	2,475	-
Non-Hispanic white													
Both sexes													
2001	8,222	8	1	4	12	31	100	247	585	1,460	2,753	3,021	-
2000	8,391	11	3	5	17	27	87	245	533	1,520	2,883	3,060	-
1999	8,257	7	1	6	10	30	106	250	578	1,452	2,803	3,014	-
Male													
2001	3,513	5	-	-	6	16	39	120	317	788	1,301	921	-
2000	3,650	3	2	4	8	18	33	139	311	848	1,356	928	-
1999	3,463	5	-	4	6	17	49	138	333	802	1,271	838	-
Female													
2001	4,709	3	1	4	6	15	61	127	268	672	1,452	2,100	-
2000	4,741	8	1	1	9	9	54	106	222	672	1,527	2,132	-
1999	4,794	2	1	2	4	13	57	112	245	650	1,532	2,176	-
Non-Hispanic black													
Both sexes													
2001	1,368	2	1	1	2	16	50	73	159	286	394	384	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)													
Non-Hispanic black													
Both sexes													
2000	1,307	2	-	1	4	10	34	97	150	261	379	368	1
1999	1,244	2	-	-	3	9	32	60	138	246	386	368	-
Male													
2001	555	2	-	-	-	8	24	41	73	138	166	103	-
2000	532	2	-	-	1	3	13	55	75	128	161	93	1
1999	487	2	-	-	2	4	9	26	69	119	166	90	-
Female													
2001	813	-	1	1	2	8	26	32	86	148	228	281	-
2000	775	-	-	1	3	7	21	42	75	133	218	275	-
1999	757	-	-	-	1	5	23	34	69	127	220	278	-
Origin not stated [2]													
Both sexes													
2001	26	-	-	-	-	1	-	2	2	3	7	11	-
2000	22	-	-	-	-	-	-	-	4	4	6	8	-
1999	27	-	-	-	-	-	-	1	3	6	6	11	-
Male													
2001	9	-	-	-	-	1	-	1	1	2	2	2	-
2000	10	-	-	-	-	-	-	-	2	3	2	3	-
1999	10	-	-	-	-	-	-	-	2	3	3	2	-
Female													
2001	17	-	-	-	-	-	-	1	1	1	5	9	-
2000	12	-	-	-	-	-	-	-	2	1	4	5	-
1999	17	-	-	-	-	-	-	1	1	3	3	9	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other disorders of circulatory system (I80-I99)													
All origins													
Both sexes													
2001	4,665	22	5	-	38	124	285	454	563	841	1,183	1,149	1
2000	4,603	27	4	12	37	133	319	515	516	795	1,161	1,084	-
1999	4,025	23	1	2	30	93	271	378	445	692	1,117	972	1
Male													
2001	1,981	11	5	-	17	53	149	262	304	405	488	286	1
2000	1,981	16	2	6	16	57	190	295	264	390	472	273	-
1999	1,714	13	1	1	13	44	159	198	224	350	463	248	-
Female													
2001	2,684	11	-	-	21	71	136	192	259	436	695	863	-
2000	2,622	11	2	6	21	76	129	220	252	405	689	811	-
1999	2,311	10	-	1	17	49	112	180	221	342	654	724	1
Hispanic													
Both sexes													
2001	219	5	-	-	3	7	15	32	16	48	41	52	-
2000	229	6	-	4	3	11	32	34	23	35	44	37	-
1999	186	2	-	1	4	6	12	34	24	42	26	35	-
Male													
2001	104	4	-	-	2	4	7	22	9	20	20	16	-
2000	116	3	-	2	1	7	20	19	13	16	22	13	-
1999	83	-	-	1	2	4	8	15	11	17	13	12	-
Female													
2001	115	1	-	-	1	3	8	10	7	28	21	36	-
2000	113	3	-	2	2	4	12	15	10	19	22	24	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other disorders of circulatory system (I80-I99)													
Hispanic													
Female													
1999	103	2	-	-	2	2	4	19	13	25	13	23	-
Non-Hispanic [1]													
Both sexes													
2001	4,425	17	5	-	35	117	269	418	544	788	1,136	1,095	1
2000	4,359	20	4	8	34	120	283	479	491	758	1,117	1,045	-
1999	3,825	21	1	1	26	86	256	343	420	648	1,088	934	1
Male													
2001	1,869	7	5	-	15	49	141	239	293	384	466	269	1
2000	1,856	12	2	4	15	49	167	274	250	373	450	260	-
1999	1,623	13	1	-	11	40	148	182	212	331	450	235	-
Female													
2001	2,556	10	-	-	20	68	128	179	251	404	670	826	-
2000	2,503	8	2	4	19	71	116	205	241	385	667	785	-
1999	2,202	8	-	1	15	46	108	161	208	317	638	699	1
Non-Hispanic white													
Both sexes													
2001	3,617	12	3	-	18	73	175	306	414	642	987	987	-
2000	3,554	13	3	5	19	70	195	358	373	602	975	941	-
1999	3,149	17	-	1	12	57	168	243	342	524	949	835	1
Male													
2001	1,496	4	3	-	8	30	92	182	220	313	405	239	-
2000	1,496	9	1	3	8	26	118	207	188	307	398	231	-
1999	1,310	9	-	-	7	26	95	127	180	268	388	210	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other disorders of circulatory system (I80-I99)													
Non-Hispanic white													
Female													
2001	2,121	8	-	-	10	43	83	124	194	329	582	748	-
2000	2,058	4	2	2	11	44	77	151	185	295	577	710	-
1999	1,839	8	-	1	5	31	73	116	162	256	561	625	1
Non-Hispanic black													
Both sexes													
2001	729	5	2	-	14	43	87	98	122	129	133	95	1
2000	751	5	1	3	14	48	86	116	109	145	130	94	-
1999	613	3	1	-	14	28	81	95	69	109	125	88	-
Male													
2001	335	3	2	-	5	18	46	48	72	62	53	25	1
2000	333	2	1	1	7	22	48	66	56	61	43	26	-
1999	281	3	1	-	4	14	49	52	28	56	55	19	-
Female													
2001	394	2	-	-	9	25	41	50	50	67	80	70	-
2000	418	3	-	2	7	26	38	50	53	84	87	68	-
1999	332	-	-	-	10	14	32	43	41	53	70	69	-
Origin not stated [2]													
Both sexes													
2001	21	-	-	-	-	-	1	4	3	5	6	2	-
2000	15	1	-	-	-	2	4	2	2	2	-	2	-
1999	14	-	-	-	-	1	3	1	1	2	3	3	-
Male													
2001	8	-	-	-	-	-	1	1	2	1	2	1	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other disorders of circulatory system (I80-I99)													
Origin not stated [2]													
Male													
2000	9	1	-	-	-	1	3	2	1	1	-	-	-
1999	8	-	-	-	-	-	3	1	1	2	-	1	-
Female													
2001	13	-	-	-	-	-	-	3	1	4	4	1	-
2000	6	-	-	-	-	1	1	-	1	1	-	2	-
1999	6	-	-	-	-	1	-	-	-	-	3	2	-
Influenza and pneumonia (J10-J18)													
All origins													
Both sexes													
2001	62,034	299	112	92	181	339	983	1,801	2,704	6,650	18,677	30,191	5
2000	65,313	289	103	87	189	364	1,068	1,774	2,879	7,189	19,821	31,547	3
1999	63,730	320	130	93	179	339	1,063	1,697	2,625	6,861	19,192	31,229	2
Male													
2001	27,342	167	57	47	106	191	556	1,099	1,587	3,732	9,294	10,502	4
2000	28,658	163	54	48	102	204	630	1,066	1,680	4,013	9,751	10,945	2
1999	27,718	171	66	53	90	197	592	1,059	1,462	3,893	9,458	10,676	1
Female													
2001	34,692	132	55	45	75	148	427	702	1,117	2,918	9,383	19,689	1
2000	36,655	126	49	39	87	160	438	708	1,199	3,176	10,070	20,602	1
1999	36,012	149	64	40	89	142	471	638	1,163	2,968	9,734	20,553	1
Hispanic													
Both sexes													
2001	2,722	57	32	10	18	23	73	121	168	402	761	1,056	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Influenza and pneumonia (J10-J18)													
Hispanic													
Both sexes													
2000	2,625	60	17	21	21	48	82	120	191	405	690	970	-
1999	2,246	64	17	10	25	24	74	104	155	328	569	876	-
Male													
2001	1,309	31	16	6	15	14	48	85	99	204	388	402	1
2000	1,303	30	5	10	10	27	56	82	119	218	367	379	-
1999	1,092	40	10	6	11	15	54	71	80	187	284	334	-
Female													
2001	1,413	26	16	4	3	9	25	36	69	198	373	654	-
2000	1,322	30	12	11	11	21	26	38	72	187	323	591	-
1999	1,154	24	7	4	14	9	20	33	75	141	285	542	-
Non-Hispanic [1]													
Both sexes													
2001	59,103	240	79	82	163	312	906	1,669	2,516	6,209	17,863	29,062	2
2000	62,440	228	84	66	167	314	974	1,639	2,674	6,750	19,051	30,492	1
1999	61,218	255	110	83	153	315	979	1,577	2,446	6,499	18,553	30,247	1
Male													
2001	25,926	135	41	41	91	175	504	1,007	1,475	3,500	8,881	10,075	1
2000	27,218	132	47	38	91	175	566	972	1,549	3,772	9,339	10,536	1
1999	26,489	131	55	47	79	182	531	974	1,367	3,687	9,138	10,298	-
Female													
2001	33,177	105	38	41	72	137	402	662	1,041	2,709	8,982	18,987	1
2000	35,222	96	37	28	76	139	408	667	1,125	2,978	9,712	19,956	-
1999	34,729	124	55	36	74	133	448	603	1,079	2,812	9,415	19,949	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Influenza and pneumonia (J10-J18)													
Non-Hispanic white													
Both sexes													
2001	51,952	117	48	57	108	207	605	1,193	1,963	5,214	15,837	26,602	1
2000	55,135	114	51	49	116	213	678	1,165	2,061	5,546	17,068	28,073	1
1999	54,280	121	58	62	95	199	637	1,109	1,894	5,481	16,785	27,838	1
Male													
2001	22,389	71	26	28	64	112	333	707	1,132	2,919	7,830	9,166	1
2000	23,609	68	24	29	68	111	394	667	1,176	3,084	8,346	9,641	1
1999	23,097	64	22	36	51	119	334	670	1,049	3,087	8,255	9,410	-
Female													
2001	29,563	46	22	29	44	95	272	486	831	2,295	8,007	17,436	-
2000	31,526	46	27	20	48	102	284	498	885	2,462	8,722	18,432	-
1999	31,183	57	36	26	44	80	303	439	845	2,394	8,530	18,428	1
Non-Hispanic black													
Both sexes													
2001	5,686	103	24	17	45	87	274	430	471	796	1,571	1,868	-
2000	5,920	101	29	11	44	92	254	423	531	984	1,550	1,901	-
1999	5,794	114	40	15	49	107	305	425	486	880	1,410	1,963	-
Male													
2001	2,769	52	13	9	23	52	155	270	293	472	808	622	-
2000	2,880	55	20	7	21	58	151	271	322	562	763	650	-
1999	2,786	58	25	8	23	58	173	277	282	518	688	676	-
Female													
2001	2,917	51	11	8	22	35	119	160	178	324	763	1,246	-
2000	3,040	46	9	4	23	34	103	152	209	422	787	1,251	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Influenza and pneumonia (J10-J18)													
Non-Hispanic black													
Female													
1999	3,008	56	15	7	26	49	132	148	204	362	722	1,287	-
Origin not stated [2]													
Both sexes													
2001	209	2	1	-	-	4	4	11	20	39	53	73	2
2000	248	1	2	-	1	2	12	15	14	34	80	85	2
1999	266	1	3	-	1	-	10	16	24	34	70	106	1
Male													
2001	107	1	-	-	-	2	4	7	13	28	25	25	2
2000	137	1	2	-	1	2	8	12	12	23	45	30	1
1999	137	-	1	-	-	-	7	14	15	19	36	44	1
Female													
2001	102	1	1	-	-	2	-	4	7	11	28	48	-
2000	111	-	-	-	-	-	4	3	2	11	35	55	1
1999	129	1	2	-	1	-	3	2	9	15	34	62	-
Influenza (J10-J11)													
All origins													
Both sexes													
2001	257	7	6	12	7	10	6	15	21	21	56	96	-
2000	1,765	9	10	11	13	13	22	48	93	199	492	855	-
1999	1,665	13	12	11	11	9	16	26	71	164	451	881	-
Male													
2001	101	6	2	4	6	6	2	9	9	8	20	29	-
2000	664	5	4	5	6	5	14	25	44	106	196	254	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Influenza (J10-J11)													
All origins													
Male													
1999	602	6	9	5	2	5	9	15	37	74	202	238	-
Female													
2001	156	1	4	8	1	4	4	6	12	13	36	67	-
2000	1,101	4	6	6	7	8	8	23	49	93	296	601	-
1999	1,063	7	3	6	9	4	7	11	34	90	249	643	-
Hispanic													
Both sexes													
2001	13	-	3	1	1	-	1	1	-	1	5	-	-
2000	47	2	1	3	1	4	2	2	2	4	10	16	-
1999	45	4	3	-	2	-	2	2	5	8	9	10	-
Male													
2001	7	-	2	1	1	-	-	1	-	-	2	-	-
2000	21	-	1	1	1	1	1	1	2	2	5	6	-
1999	21	-	3	-	-	-	2	2	4	7	2	1	-
Female													
2001	6	-	1	-	-	-	1	-	-	1	3	-	-
2000	26	2	-	2	-	3	1	1	-	2	5	10	-
1999	24	4	-	-	2	-	-	-	1	1	7	9	-
Non-Hispanic [1]													
Both sexes													
2001	244	7	3	11	6	10	5	14	21	20	51	96	-
2000	1,716	7	9	8	12	9	20	46	91	195	481	838	-
1999	1,612	9	9	11	9	9	14	24	66	154	439	868	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Influenza (J10-J11)													
Non-Hispanic [1]													
Male													
2001	94	6	-	3	5	6	2	8	9	8	18	29	-
2000	642	5	3	4	5	4	13	24	42	104	190	248	-
1999	580	6	6	5	2	5	7	13	33	66	200	237	-
Female													
2001	150	1	3	8	1	4	3	6	12	12	33	67	-
2000	1,074	2	6	4	7	5	7	22	49	91	291	590	-
1999	1,032	3	3	6	7	4	7	11	33	88	239	631	-
Non-Hispanic white													
Both sexes													
2001	216	7	2	6	4	7	3	14	17	18	47	91	-
2000	1,613	5	8	6	11	7	16	38	85	176	458	803	-
1999	1,522	3	5	10	8	8	8	20	60	139	417	844	-
Male													
2001	81	6	-	2	3	3	2	8	6	6	18	27	-
2000	595	5	3	3	4	2	9	18	39	97	181	234	-
1999	547	1	3	5	2	5	4	11	32	59	192	233	-
Female													
2001	135	1	2	4	1	4	1	6	11	12	29	64	-
2000	1,018	-	5	3	7	5	7	20	46	79	277	569	-
1999	975	2	2	5	6	3	4	9	28	80	225	611	-
Non-Hispanic black													
Both sexes													
2001	20	-	-	2	1	2	1	-	3	2	4	5	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Influenza (J10-J11)													
Non-Hispanic black													
Both sexes													
2000	81	2	1	1	1	1	3	7	6	16	13	30	-
1999	65	4	1	-	1	1	5	3	5	12	15	18	-
Male													
2001	10	-	-	-	1	2	-	-	3	2	-	2	-
2000	35	-	-	1	1	1	3	5	3	5	5	11	-
1999	24	3	1	-	-	-	2	2	1	6	6	3	-
Female													
2001	10	-	-	2	-	-	1	-	-	-	4	3	-
2000	46	2	1	-	-	-	-	2	3	11	8	19	-
1999	41	1	-	-	1	1	3	1	4	6	9	15	-
Origin not stated [2]													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	1	1	-
1999	8	-	-	-	-	-	-	-	-	2	3	3	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	1	-	-	-	-	-	-	-	-	1	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	1	-
1999	7	-	-	-	-	-	-	-	-	1	3	3	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Pneumonia (J12-J18)													
All origins													
Both sexes													
2001	61,777	292	106	80	174	329	977	1,786	2,683	6,629	18,621	30,095	5
2000	63,548	280	93	76	176	351	1,046	1,726	2,786	6,990	19,329	30,692	3
1999	62,065	307	118	82	168	330	1,047	1,671	2,554	6,697	18,741	30,348	2
Male													
2001	27,241	161	55	43	100	185	554	1,090	1,578	3,724	9,274	10,473	4
2000	27,994	158	50	43	96	199	616	1,041	1,636	3,907	9,555	10,691	2
1999	27,116	165	57	48	88	192	583	1,044	1,425	3,819	9,256	10,438	1
Female													
2001	34,536	131	51	37	74	144	423	696	1,105	2,905	9,347	19,622	1
2000	35,554	122	43	33	80	152	430	685	1,150	3,083	9,774	20,001	1
1999	34,949	142	61	34	80	138	464	627	1,129	2,878	9,485	19,910	1
Hispanic													
Both sexes													
2001	2,709	57	29	9	17	23	72	120	168	401	756	1,056	1
2000	2,578	58	16	18	20	44	80	118	189	401	680	954	-
1999	2,201	60	14	10	23	24	72	102	150	320	560	866	-
Male													
2001	1,302	31	14	5	14	14	48	84	99	204	386	402	1
2000	1,282	30	4	9	9	26	55	81	117	216	362	373	-
1999	1,071	40	7	6	11	15	52	69	76	180	282	333	-
Female													
2001	1,407	26	15	4	3	9	24	36	69	197	370	654	-
2000	1,296	28	12	9	11	18	25	37	72	185	318	581	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Pneumonia (J12-J18)													
Hispanic													
Female													
1999	1,130	20	7	4	12	9	20	33	74	140	278	533	-
Non-Hispanic [1]													
Both sexes													
2001	58,859	233	76	71	157	302	901	1,655	2,495	6,189	17,812	28,966	2
2000	60,724	221	75	58	155	305	954	1,593	2,583	6,555	18,570	29,654	1
1999	59,606	246	101	72	144	306	965	1,553	2,380	6,345	18,114	29,379	1
Male													
2001	25,832	129	41	38	86	169	502	999	1,466	3,492	8,863	10,046	1
2000	26,576	127	44	34	86	171	553	948	1,507	3,668	9,149	10,288	1
1999	25,909	125	49	42	77	177	524	961	1,334	3,621	8,938	10,061	-
Female													
2001	33,027	104	35	33	71	133	399	656	1,029	2,697	8,949	18,920	1
2000	34,148	94	31	24	69	134	401	645	1,076	2,887	9,421	19,366	-
1999	33,697	121	52	30	67	129	441	592	1,046	2,724	9,176	19,318	1
Non-Hispanic white													
Both sexes													
2001	51,736	110	46	51	104	200	602	1,179	1,946	5,196	15,790	26,511	1
2000	53,522	109	43	43	105	206	662	1,127	1,976	5,370	16,610	27,270	1
1999	52,758	118	53	52	87	191	629	1,089	1,834	5,342	16,368	26,994	1
Male													
2001	22,308	65	26	26	61	109	331	699	1,126	2,913	7,812	9,139	1
2000	23,014	63	21	26	64	109	385	649	1,137	2,987	8,165	9,407	1
1999	22,550	63	19	31	49	114	330	659	1,017	3,028	8,063	9,177	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Pneumonia (J12-J18)													
Non-Hispanic white													
Female													
2001	29,428	45	20	25	43	91	271	480	820	2,283	7,978	17,372	-
2000	30,508	46	22	17	41	97	277	478	839	2,383	8,445	17,863	-
1999	30,208	55	34	21	38	77	299	430	817	2,314	8,305	17,817	1
Non-Hispanic black													
Both sexes													
2001	5,666	103	24	15	44	85	273	430	468	794	1,567	1,863	-
2000	5,839	99	28	10	43	91	251	416	525	968	1,537	1,871	-
1999	5,729	110	39	15	48	106	300	422	481	868	1,395	1,945	-
Male													
2001	2,759	52	13	9	22	50	155	270	290	470	808	620	-
2000	2,845	55	20	6	20	57	148	266	319	557	758	639	-
1999	2,762	55	24	8	23	58	171	275	281	512	682	673	-
Female													
2001	2,907	51	11	6	22	35	118	160	178	324	759	1,243	-
2000	2,994	44	8	4	23	34	103	150	206	411	779	1,232	-
1999	2,967	55	15	7	25	48	129	147	200	356	713	1,272	-
Origin not stated [2]													
Both sexes													
2001	209	2	1	-	-	4	4	11	20	39	53	73	2
2000	246	1	2	-	1	2	12	15	14	34	79	84	2
1999	258	1	3	-	1	-	10	16	24	32	67	103	1
Male													
2001	107	1	-	-	-	2	4	7	13	28	25	25	2

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Pneumonia (J12-J18)													
Origin not stated [2]													
Male													
2000	136	1	2	-	1	2	8	12	12	23	44	30	1
1999	136	-	1	-	-	-	7	14	15	18	36	44	1
Female													
2001	102	1	1	-	-	2	-	4	7	11	28	48	-
2000	110	-	-	-	-	-	4	3	2	11	35	54	1
1999	122	1	2	-	1	-	3	2	9	14	31	59	-
Other acute lower respiratory infections (J20-J22)													
All origins													
Both sexes													
2001	347	50	18	3	1	2	12	14	16	28	58	145	-
2000	425	43	11	7	5	8	12	12	10	43	82	192	-
1999	540	63	25	5	5	4	11	15	21	35	107	249	-
Male													
2001	117	26	8	2	1	-	7	7	6	10	20	30	-
2000	160	25	4	3	4	6	7	4	3	22	35	47	-
1999	196	36	15	2	1	3	5	8	10	17	51	48	-
Female													
2001	230	24	10	1	-	2	5	7	10	18	38	115	-
2000	265	18	7	4	1	2	5	8	7	21	47	145	-
1999	344	27	10	3	4	1	6	7	11	18	56	201	-
Hispanic													
Both sexes													
2001	22	8	1	-	-	1	-	2	-	2	4	4	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other acute lower respiratory infections (J20-J22)													
Hispanic													
Both sexes													
2000	24	9	2	-	-	1	-	1	-	1	2	8	-
1999	21	10	3	1	-	-	-	1	-	1	-	5	-
Male													
2001	9	6	-	-	-	-	-	1	-	1	1	-	-
2000	15	6	1	-	-	1	-	1	-	1	2	3	-
1999	10	5	2	1	-	-	-	-	-	1	-	1	-
Female													
2001	13	2	1	-	-	1	-	1	-	1	3	4	-
2000	9	3	1	-	-	-	-	-	-	-	-	5	-
1999	11	5	1	-	-	-	-	1	-	-	-	4	-
Non-Hispanic [1]													
Both sexes													
2001	325	42	17	3	1	1	12	12	16	26	54	141	-
2000	398	33	9	7	5	7	12	11	10	42	79	183	-
1999	518	53	22	4	5	4	10	14	21	34	107	244	-
Male													
2001	108	20	8	2	1	-	7	6	6	9	19	30	-
2000	144	19	3	3	4	5	7	3	3	21	32	44	-
1999	185	31	13	1	1	3	4	8	10	16	51	47	-
Female													
2001	217	22	9	1	-	1	5	6	10	17	35	111	-
2000	254	14	6	4	1	2	5	8	7	21	47	139	-
1999	333	22	9	3	4	1	6	6	11	18	56	197	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other acute lower respiratory infections (J20-J22)													
Non-Hispanic white													
Both sexes													
2001	278	24	12	3	1	-	12	8	14	20	50	134	-
2000	345	15	6	3	5	4	8	8	10	40	74	172	-
1999	440	26	12	3	4	3	8	10	17	26	99	232	-
Male													
2001	86	11	5	2	1	-	7	3	5	7	17	28	-
2000	118	6	3	2	4	3	6	3	3	19	30	39	-
1999	145	14	6	1	1	2	3	8	9	11	48	42	-
Female													
2001	192	13	7	1	-	-	5	5	9	13	33	106	-
2000	227	9	3	1	1	1	2	5	7	21	44	133	-
1999	295	12	6	2	3	1	5	2	8	15	51	190	-
Non-Hispanic black													
Both sexes													
2001	33	14	2	-	-	1	-	4	1	4	-	7	-
2000	46	14	3	3	-	2	4	3	-	2	5	10	-
1999	64	24	6	-	1	1	2	4	2	8	6	10	-
Male													
2001	15	8	1	-	-	-	-	3	-	1	-	2	-
2000	22	10	-	1	-	2	1	-	-	2	2	4	-
1999	30	15	3	-	-	1	1	-	1	5	1	3	-
Female													
2001	18	6	1	-	-	1	-	1	1	3	-	5	-
2000	24	4	3	2	-	-	3	3	-	-	3	6	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other acute lower respiratory infections (J20-J22)													
Non-Hispanic black													
Female													
1999	34	9	3	-	1	-	1	4	1	3	5	7	-
Origin not stated [2]													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	3	1	-	-	-	-	-	-	-	-	1	1	-
1999	1	-	-	-	-	-	1	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	1	-	-	-	-	-	1	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	1	-	-	-	-	-	-	-	-	-	1	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Acute bronchitis and bronchiolitis (J20-J21)													
All origins													
Both sexes													
2001	263	49	18	3	1	2	11	12	14	23	39	91	-
2000	290	42	11	6	4	6	10	9	7	31	53	111	-
1999	388	62	24	4	5	3	10	14	18	29	69	150	-
Male													
2001	89	25	8	2	1	-	6	6	6	8	10	17	-
2000	119	24	4	2	3	4	6	3	3	14	27	29	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Acute bronchitis and bronchiolitis (J20-J21)													
All origins													
Male													
1999	154	36	14	2	1	2	4	7	8	15	34	31	-
Female													
2001	174	24	10	1	-	2	5	6	8	15	29	74	-
2000	171	18	7	4	1	2	4	6	4	17	26	82	-
1999	234	26	10	2	4	1	6	7	10	14	35	119	-
Hispanic													
Both sexes													
2001	17	8	1	-	-	1	-	2	-	1	2	2	-
2000	22	9	2	-	-	1	-	1	-	1	2	6	-
1999	20	10	3	1	-	-	-	1	-	1	-	4	-
Male													
2001	7	6	-	-	-	-	-	1	-	-	-	-	-
2000	15	6	1	-	-	1	-	1	-	1	2	3	-
1999	10	5	2	1	-	-	-	-	-	1	-	1	-
Female													
2001	10	2	1	-	-	1	-	1	-	1	2	2	-
2000	7	3	1	-	-	-	-	-	-	-	-	3	-
1999	10	5	1	-	-	-	-	1	-	-	-	3	-
Non-Hispanic [1]													
Both sexes													
2001	246	41	17	3	1	1	11	10	14	22	37	89	-
2000	267	32	9	6	4	5	10	8	7	30	51	105	-
1999	367	52	21	3	5	3	9	13	18	28	69	146	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Acute bronchitis and bronchiolitis (J20-J21)													
Non-Hispanic [1]													
Male													
2001	82	19	8	2	1	-	6	5	6	8	10	17	-
2000	104	18	3	2	3	3	6	2	3	13	25	26	-
1999	143	31	12	1	1	2	3	7	8	14	34	30	-
Female													
2001	164	22	9	1	-	1	5	5	8	14	27	72	-
2000	163	14	6	4	1	2	4	6	4	17	26	79	-
1999	224	21	9	2	4	1	6	6	10	14	35	116	-
Non-Hispanic white													
Both sexes													
2001	202	23	12	3	1	-	11	6	12	17	34	83	-
2000	221	15	6	2	4	2	6	5	7	28	47	99	-
1999	299	25	12	2	4	3	7	9	14	20	63	140	-
Male													
2001	61	10	5	2	1	-	6	2	5	6	8	16	-
2000	83	6	3	1	3	1	5	2	3	11	24	24	-
1999	108	14	6	1	1	2	2	7	7	9	32	27	-
Female													
2001	141	13	7	1	-	-	5	4	7	11	26	67	-
2000	138	9	3	1	1	1	1	3	4	17	23	75	-
1999	191	11	6	1	3	1	5	2	7	11	31	113	-
Non-Hispanic black													
Both sexes													
2001	31	14	2	-	-	1	-	4	1	3	-	6	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Acute bronchitis and bronchiolitis (J20-J21)													
Non-Hispanic black													
Both sexes													
2000	40	13	3	3	-	2	4	3	-	2	4	6	-
1999	55	24	5	-	1	-	2	4	2	8	4	5	-
Male													
2001	14	8	1	-	-	-	-	3	-	1	-	1	-
2000	18	9	-	1	-	2	1	-	-	2	1	2	-
1999	26	15	2	-	-	-	1	-	1	5	-	2	-
Female													
2001	17	6	1	-	-	1	-	1	1	2	-	5	-
2000	22	4	3	2	-	-	3	3	-	-	3	4	-
1999	29	9	3	-	1	-	1	4	1	3	4	3	-
Origin not stated [2]													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	1	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	1	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Unspecified acute lower respiratory infection (J22)													
All origins													
Both sexes													
2001	84	1	-	-	-	-	1	2	2	5	19	54	-
2000	135	1	-	1	1	2	2	3	3	12	29	81	-
1999	152	1	1	1	-	1	1	1	3	6	38	99	-
Male													
2001	28	1	-	-	-	-	1	1	-	2	10	13	-
2000	41	1	-	1	1	2	1	1	-	8	8	18	-
1999	42	-	1	-	-	1	1	1	2	2	17	17	-
Female													
2001	56	-	-	-	-	-	-	1	2	3	9	41	-
2000	94	-	-	-	-	-	1	2	3	4	21	63	-
1999	110	1	-	1	-	-	-	-	1	4	21	82	-
Hispanic													
Both sexes													
2001	5	-	-	-	-	-	-	-	-	1	2	2	-
2000	2	-	-	-	-	-	-	-	-	-	-	2	-
1999	1	-	-	-	-	-	-	-	-	-	-	1	-
Male													
2001	2	-	-	-	-	-	-	-	-	1	1	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	3	-	-	-	-	-	-	-	-	-	1	2	-
2000	2	-	-	-	-	-	-	-	-	-	-	2	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Unspecified acute lower respiratory infection (J22)													
Hispanic													
Female													
1999	1	-	-	-	-	-	-	-	-	-	-	1	-
Non-Hispanic [1]													
Both sexes													
2001	79	1	-	-	-	-	1	2	2	4	17	52	-
2000	131	1	-	1	1	2	2	3	3	12	28	78	-
1999	151	1	1	1	-	1	1	1	3	6	38	98	-
Male													
2001	26	1	-	-	-	-	1	1	-	1	9	13	-
2000	40	1	-	1	1	2	1	1	-	8	7	18	-
1999	42	-	1	-	-	1	1	1	2	2	17	17	-
Female													
2001	53	-	-	-	-	-	-	1	2	3	8	39	-
2000	91	-	-	-	-	-	1	2	3	4	21	60	-
1999	109	1	-	1	-	-	-	-	1	4	21	81	-
Non-Hispanic white													
Both sexes													
2001	76	1	-	-	-	-	1	2	2	3	16	51	-
2000	124	-	-	1	1	2	2	3	3	12	27	73	-
1999	141	1	-	1	-	-	1	1	3	6	36	92	-
Male													
2001	25	1	-	-	-	-	1	1	-	1	9	12	-
2000	35	-	-	1	1	2	1	1	-	8	6	15	-
1999	37	-	-	-	-	-	1	1	2	2	16	15	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Unspecified acute lower respiratory infection (J22)													
Non-Hispanic white													
Female													
2001	51	-	-	-	-	-	-	1	2	2	7	39	-
2000	89	-	-	-	-	-	1	2	3	4	21	58	-
1999	104	1	-	1	-	-	-	-	1	4	20	77	-
Non-Hispanic black													
Both sexes													
2001	2	-	-	-	-	-	-	-	-	1	-	1	-
2000	6	1	-	-	-	-	-	-	-	-	1	4	-
1999	9	-	1	-	-	1	-	-	-	-	2	5	-
Male													
2001	1	-	-	-	-	-	-	-	-	-	-	1	-
2000	4	1	-	-	-	-	-	-	-	-	1	2	-
1999	4	-	1	-	-	1	-	-	-	-	1	1	-
Female													
2001	1	-	-	-	-	-	-	-	-	1	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	-	2	-
1999	5	-	-	-	-	-	-	-	-	-	1	4	-
Origin not stated [2]													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	1	1	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Unspecified acute lower respiratory infection (J22)													
Origin not stated [2]													
Male													
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	1	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Chronic lower respiratory diseases (J40-J47)													
All origins													
Both sexes													
2001	123,013	41	43	104	171	290	969	3,324	11,166	30,751	47,762	28,391	1
2000	122,009	36	51	139	190	296	930	3,251	10,739	31,157	47,722	27,496	2
1999	124,181	36	54	139	209	330	890	3,110	11,297	32,644	48,635	26,833	4
Male													
2001	59,697	22	29	61	99	146	455	1,702	5,702	15,877	23,691	11,912	1
2000	60,004	23	28	80	120	165	437	1,661	5,564	16,210	23,838	11,876	2
1999	62,415	19	30	93	123	152	410	1,551	5,958	17,477	24,705	11,894	3
Female													
2001	63,316	19	14	43	72	144	514	1,622	5,464	14,874	24,071	16,479	-
2000	62,005	13	23	59	70	131	493	1,590	5,175	14,947	23,884	15,620	-
1999	61,766	17	24	46	86	178	480	1,559	5,339	15,167	23,930	14,939	1
Hispanic													
Both sexes													
2001	2,832	7	8	9	10	22	52	124	258	634	940	768	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Chronic lower respiratory diseases (J40-J47)													
Hispanic													
Both sexes													
2000	2,689	7	9	15	16	20	55	105	244	572	921	725	-
1999	2,859	6	8	18	15	31	79	115	239	649	935	764	-
Male													
2001	1,482	4	4	6	6	13	21	54	140	378	501	355	-
2000	1,451	4	5	9	13	12	24	53	146	343	509	333	-
1999	1,547	4	3	11	9	13	36	52	135	373	523	388	-
Female													
2001	1,350	3	4	3	4	9	31	70	118	256	439	413	-
2000	1,238	3	4	6	3	8	31	52	98	229	412	392	-
1999	1,312	2	5	7	6	18	43	63	104	276	412	376	-
Non-Hispanic [1]													
Both sexes													
2001	119,839	34	35	94	161	267	909	3,180	10,860	30,028	46,713	27,558	-
2000	118,963	29	42	123	173	272	872	3,122	10,447	30,486	46,687	26,709	1
1999	120,968	30	46	121	194	296	807	2,978	11,018	31,902	47,578	25,995	3
Male													
2001	58,028	18	25	55	93	133	428	1,635	5,530	15,449	23,134	11,528	-
2000	58,349	19	23	71	107	150	411	1,593	5,388	15,805	23,266	11,515	1
1999	60,669	15	27	82	114	138	372	1,486	5,797	17,040	24,124	11,472	2
Female													
2001	61,811	16	10	39	68	134	481	1,545	5,330	14,579	23,579	16,030	-
2000	60,614	10	19	52	66	122	461	1,529	5,059	14,681	23,421	15,194	-
1999	60,299	15	19	39	80	158	435	1,492	5,221	14,862	23,454	14,523	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Chronic lower respiratory diseases (J40-J47)													
Non-Hispanic white													
Both sexes													
2001	110,753	16	13	39	75	134	613	2,570	9,678	27,693	43,972	25,950	-
2000	109,901	16	19	47	87	144	570	2,463	9,253	28,122	44,024	25,155	1
1999	111,620	10	23	55	110	146	533	2,372	9,786	29,400	44,662	24,521	2
Male													
2001	52,952	10	9	22	39	62	293	1,331	4,885	14,089	21,496	10,716	-
2000	53,231	12	8	24	52	71	267	1,259	4,728	14,398	21,703	10,708	1
1999	55,314	6	11	36	59	66	239	1,174	5,091	15,559	22,377	10,695	1
Female													
2001	57,801	6	4	17	36	72	320	1,239	4,793	13,604	22,476	15,234	-
2000	56,670	4	11	23	35	73	303	1,204	4,525	13,724	22,321	14,447	-
1999	56,306	4	12	19	51	80	294	1,198	4,695	13,841	22,285	13,826	1
Non-Hispanic black													
Both sexes													
2001	7,504	16	22	53	77	128	273	549	1,019	1,962	2,190	1,215	-
2000	7,528	11	21	70	80	114	272	593	1,063	1,993	2,128	1,183	-
1999	7,840	18	18	66	78	140	258	556	1,085	2,090	2,364	1,166	1
Male													
2001	4,143	6	16	32	48	69	124	271	554	1,147	1,285	591	-
2000	4,184	6	14	43	51	70	129	289	587	1,195	1,207	593	-
1999	4,452	9	13	46	52	67	121	284	624	1,238	1,402	595	1
Female													
2001	3,361	10	6	21	29	59	149	278	465	815	905	624	-
2000	3,344	5	7	27	29	44	143	304	476	798	921	590	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Chronic lower respiratory diseases (J40-J47)													
Non-Hispanic black													
Female													
1999	3,388	9	5	20	26	73	137	272	461	852	962	571	-
Origin not stated [2]													
Both sexes													
2001	342	-	-	1	-	1	8	20	48	89	109	65	1
2000	357	-	-	1	1	4	3	24	48	99	114	62	1
1999	354	-	-	-	-	3	4	17	40	93	122	74	1
Male													
2001	187	-	-	-	-	-	6	13	32	50	56	29	1
2000	204	-	-	-	-	3	2	15	30	62	63	28	1
1999	199	-	-	-	-	1	2	13	26	64	58	34	1
Female													
2001	155	-	-	1	-	1	2	7	16	39	53	36	-
2000	153	-	-	1	1	1	1	9	18	37	51	34	-
1999	155	-	-	-	-	2	2	4	14	29	64	40	-
Bronchitis, chronic and unspecified (J40-J42)													
All origins													
Both sexes													
2001	959	21	8	-	5	6	21	42	61	133	261	401	-
2000	1,167	18	14	5	5	8	20	38	67	158	333	501	-
1999	1,172	20	19	4	7	11	18	38	97	173	339	446	-
Male													
2001	396	11	5	-	3	2	8	21	36	76	115	119	-
2000	503	9	7	3	4	7	14	16	32	79	174	158	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Bronchitis, chronic and unspecified (J40-J42)													
All origins													
Male													
1999	498	9	11	3	4	3	5	19	51	94	157	142	-
Female													
2001	563	10	3	-	2	4	13	21	25	57	146	282	-
2000	664	9	7	2	1	1	6	22	35	79	159	343	-
1999	674	11	8	1	3	8	13	19	46	79	182	304	-
Hispanic													
Both sexes													
2001	37	4	1	-	1	1	2	1	2	3	9	13	-
2000	52	4	3	-	-	2	3	1	1	9	13	16	-
1999	52	3	3	1	-	2	7	-	2	4	13	17	-
Male													
2001	15	2	-	-	-	1	1	-	1	1	6	3	-
2000	28	2	2	-	-	2	2	-	1	7	6	6	-
1999	17	1	2	1	-	-	2	-	-	3	3	5	-
Female													
2001	22	2	1	-	1	-	1	1	1	2	3	10	-
2000	24	2	1	-	-	-	1	1	-	2	7	10	-
1999	35	2	1	-	-	2	5	-	2	1	10	12	-
Non-Hispanic [1]													
Both sexes													
2001	919	17	7	-	4	5	19	40	58	130	251	388	-
2000	1,112	14	11	5	5	6	17	37	66	149	320	482	-
1999	1,117	17	16	3	7	9	11	38	94	168	326	428	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Bronchitis, chronic and unspecified (J40-J42)													
Non-Hispanic [1]													
Male													
2001	380	9	5	-	3	1	7	21	34	75	109	116	-
2000	474	7	5	3	4	5	12	16	31	72	168	151	-
1999	478	8	9	2	4	3	3	19	50	90	154	136	-
Female													
2001	539	8	2	-	1	4	12	19	24	55	142	272	-
2000	638	7	6	2	1	1	5	21	35	77	152	331	-
1999	639	9	7	1	3	6	8	19	44	78	172	292	-
Non-Hispanic white													
Both sexes													
2001	839	7	5	-	3	4	16	35	54	113	232	370	-
2000	1,018	5	4	4	4	4	11	31	60	131	297	467	-
1999	1,002	8	12	3	6	6	9	24	86	150	293	405	-
Male													
2001	338	5	5	-	2	-	6	17	32	66	97	108	-
2000	429	3	1	2	3	3	6	15	27	64	159	146	-
1999	421	5	6	2	3	2	3	11	45	81	135	128	-
Female													
2001	501	2	-	-	1	4	10	18	22	47	135	262	-
2000	589	2	3	2	1	1	5	16	33	67	138	321	-
1999	581	3	6	1	3	4	6	13	41	69	158	277	-
Non-Hispanic black													
Both sexes													
2001	62	10	2	-	1	1	3	4	4	15	11	11	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Bronchitis, chronic and unspecified (J40-J42)													
Non-Hispanic black													
Both sexes													
2000	80	7	6	1	1	1	6	6	4	16	20	12	-
1999	88	7	1	-	1	3	2	12	7	15	26	14	-
Male													
2001	32	4	-	-	1	1	1	4	2	8	6	5	-
2000	35	3	3	1	1	1	6	1	3	6	7	3	-
1999	43	3	1	-	1	1	-	7	4	7	16	3	-
Female													
2001	30	6	2	-	-	-	2	-	2	7	5	6	-
2000	45	4	3	-	-	-	-	5	1	10	13	9	-
1999	45	4	-	-	-	2	2	5	3	8	10	11	-
Origin not stated [2]													
Both sexes													
2001	3	-	-	-	-	-	-	1	1	-	1	-	-
2000	3	-	-	-	-	-	-	-	-	-	-	3	-
1999	3	-	-	-	-	-	-	-	1	1	-	1	-
Male													
2001	1	-	-	-	-	-	-	-	1	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	1	-
1999	3	-	-	-	-	-	-	-	1	1	-	1	-
Female													
2001	2	-	-	-	-	-	-	1	-	-	1	-	-
2000	2	-	-	-	-	-	-	-	-	-	-	2	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Emphysema (J43)													
All origins													
Both sexes													
2001	16,242	3	-	1	2	7	75	448	1,829	4,804	6,337	2,736	-
2000	16,700	3	2	-	5	11	82	474	1,807	5,120	6,511	2,685	-
1999	17,787	1	-	1	1	9	97	468	2,019	5,656	6,955	2,580	-
Male													
2001	8,415	1	-	1	1	4	52	266	966	2,554	3,255	1,315	-
2000	8,671	3	-	-	5	5	46	280	943	2,738	3,373	1,278	-
1999	9,480	-	-	-	1	6	49	268	1,085	3,136	3,677	1,258	-
Female													
2001	7,827	2	-	-	1	3	23	182	863	2,250	3,082	1,421	-
2000	8,029	-	2	-	-	6	36	194	864	2,382	3,138	1,407	-
1999	8,307	1	-	1	-	3	48	200	934	2,520	3,278	1,322	-
Hispanic													
Both sexes													
2001	286	1	-	-	-	-	-	8	26	75	113	63	-
2000	266	-	-	-	1	-	3	11	26	65	113	47	-
1999	292	-	-	1	-	2	6	12	21	74	113	63	-
Male													
2001	175	-	-	-	-	-	-	5	16	46	76	32	-
2000	167	-	-	-	1	-	3	8	17	43	66	29	-
1999	183	-	-	-	-	1	2	8	16	51	69	36	-
Female													
2001	111	1	-	-	-	-	-	3	10	29	37	31	-
2000	99	-	-	-	-	-	-	3	9	22	47	18	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Emphysema (J43)													
Hispanic													
Female													
1999	109	-	-	1	-	1	4	4	5	23	44	27	-
Non-Hispanic [1]													
Both sexes													
2001	15,914	2	-	1	2	7	74	439	1,794	4,718	6,210	2,667	-
2000	16,382	3	2	-	4	11	79	457	1,777	5,036	6,382	2,631	-
1999	17,441	1	-	-	1	7	88	450	1,992	5,571	6,825	2,506	-
Male													
2001	8,213	1	-	1	1	4	51	260	943	2,500	3,172	1,280	-
2000	8,483	3	-	-	4	5	43	268	925	2,685	3,303	1,247	-
1999	9,260	-	-	-	1	5	45	255	1,063	3,076	3,598	1,217	-
Female													
2001	7,701	1	-	-	1	3	23	179	851	2,218	3,038	1,387	-
2000	7,899	-	2	-	-	6	36	189	852	2,351	3,079	1,384	-
1999	8,181	1	-	-	-	2	43	195	929	2,495	3,227	1,289	-
Non-Hispanic white													
Both sexes													
2001	14,930	1	-	-	1	4	64	374	1,633	4,414	5,913	2,526	-
2000	15,301	3	-	-	1	10	60	367	1,616	4,704	6,068	2,472	-
1999	16,338	-	-	-	-	5	64	372	1,825	5,238	6,469	2,365	-
Male													
2001	7,607	-	-	-	1	2	44	222	841	2,310	2,986	1,201	-
2000	7,805	3	-	-	1	4	31	206	830	2,470	3,121	1,139	-
1999	8,540	-	-	-	-	3	33	199	953	2,843	3,371	1,138	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Emphysema (J43)													
Non-Hispanic white													
Female													
2001	7,323	1	-	-	-	2	20	152	792	2,104	2,927	1,325	-
2000	7,496	-	-	-	-	6	29	161	786	2,234	2,947	1,333	-
1999	7,798	-	-	-	-	2	31	173	872	2,395	3,098	1,227	-
Non-Hispanic black													
Both sexes													
2001	800	1	-	1	1	3	9	59	138	254	219	115	-
2000	889	-	1	-	3	1	17	84	146	276	243	118	-
1999	923	1	-	-	-	-	22	75	146	279	283	117	-
Male													
2001	480	1	-	1	-	2	7	33	87	155	134	60	-
2000	551	-	-	-	3	1	11	56	87	183	134	76	-
1999	594	-	-	-	-	-	11	54	94	195	178	62	-
Female													
2001	320	-	-	-	1	1	2	26	51	99	85	55	-
2000	338	-	1	-	-	-	6	28	59	93	109	42	-
1999	329	1	-	-	-	-	11	21	52	84	105	55	-
Origin not stated [2]													
Both sexes													
2001	42	-	-	-	-	-	1	1	9	11	14	6	-
2000	52	-	-	-	-	-	-	6	4	19	16	7	-
1999	54	-	-	-	-	-	3	6	6	11	17	11	-
Male													
2001	27	-	-	-	-	-	1	1	7	8	7	3	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Emphysema (J43)													
Origin not stated [2]													
Male													
2000	21	-	-	-	-	-	-	4	1	10	4	2	-
1999	37	-	-	-	-	-	2	5	6	9	10	5	-
Female													
2001	15	-	-	-	-	-	-	-	2	3	7	3	-
2000	31	-	-	-	-	-	-	2	3	9	12	5	-
1999	17	-	-	-	-	-	1	1	-	2	7	6	-
Asthma (J45-J46)													
All origins													
Both sexes													
2001	4,269	10	31	99	140	233	455	603	553	634	802	709	-
2000	4,487	8	32	129	167	249	458	614	540	734	849	707	-
1999	4,657	5	27	126	182	262	447	607	583	773	925	720	-
Male													
2001	1,479	6	22	57	80	116	195	219	182	220	225	157	-
2000	1,632	6	20	74	102	142	196	238	200	253	243	158	-
1999	1,620	3	16	85	108	128	184	197	207	267	273	152	-
Female													
2001	2,790	4	9	42	60	117	260	384	371	414	577	552	-
2000	2,855	2	12	55	65	107	262	376	340	481	606	549	-
1999	3,037	2	11	41	74	134	263	410	376	506	652	568	-
Hispanic													
Both sexes													
2001	274	2	5	8	8	17	31	49	45	43	44	22	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Asthma (J45-J46)													
Hispanic													
Both sexes													
2000	292	1	5	13	14	16	29	41	48	60	40	25	-
1999	320	-	3	14	14	21	46	48	39	60	50	25	-
Male													
2001	103	2	3	5	6	10	13	14	17	18	13	2	-
2000	129	1	2	8	11	9	14	20	19	25	12	8	-
1999	119	-	-	9	8	11	22	20	12	19	12	6	-
Female													
2001	171	-	2	3	2	7	18	35	28	25	31	20	-
2000	163	-	3	5	3	7	15	21	29	35	28	17	-
1999	201	-	3	5	6	10	24	28	27	41	38	19	-
Non-Hispanic [1]													
Both sexes													
2001	3,976	8	26	90	132	215	421	550	507	587	756	684	-
2000	4,164	7	27	115	152	229	428	568	486	668	805	679	-
1999	4,324	5	24	112	168	238	400	557	543	711	874	692	-
Male													
2001	1,365	4	19	52	74	106	180	202	165	200	211	152	-
2000	1,486	5	18	66	91	130	181	215	176	226	229	149	-
1999	1,495	3	16	76	100	116	162	176	195	246	260	145	-
Female													
2001	2,611	4	7	38	58	109	241	348	342	387	545	532	-
2000	2,678	2	9	49	61	99	247	353	310	442	576	530	-
1999	2,829	2	8	36	68	122	238	381	348	465	614	547	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Asthma (J45-J46)													
Non-Hispanic white													
Both sexes													
2001	2,717	4	7	37	59	97	219	326	306	432	628	602	-
2000	2,836	4	13	40	75	114	227	317	296	484	683	583	-
1999	3,011	1	8	47	92	107	222	327	351	511	747	598	-
Male													
2001	834	3	4	21	29	42	91	113	95	141	167	128	-
2000	920	2	7	20	44	60	90	130	97	159	193	118	-
1999	930	1	5	31	52	48	80	97	124	164	218	110	-
Female													
2001	1,883	1	3	16	30	55	128	213	211	291	461	474	-
2000	1,916	2	6	20	31	54	137	187	199	325	490	465	-
1999	2,081	-	3	16	40	59	142	230	227	347	529	488	-
Non-Hispanic black													
Both sexes													
2001	1,092	3	19	51	67	115	186	203	170	126	99	53	-
2000	1,145	3	14	69	73	102	178	225	165	152	94	70	-
1999	1,134	4	14	65	71	126	166	211	162	157	94	64	-
Male													
2001	450	-	15	30	41	62	79	83	52	46	29	13	-
2000	472	3	11	42	44	62	79	71	65	53	21	21	-
1999	474	2	10	45	46	66	72	72	57	61	27	16	-
Female													
2001	642	3	4	21	26	53	107	120	118	80	70	40	-
2000	673	-	3	27	29	40	99	154	100	99	73	49	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Asthma (J45-J46)													
Non-Hispanic black													
Female													
1999	660	2	4	20	25	60	94	139	105	96	67	48	-
Origin not stated [2]													
Both sexes													
2001	19	-	-	1	-	1	3	4	1	4	2	3	-
2000	31	-	-	1	1	4	1	5	6	6	4	3	-
1999	13	-	-	-	-	3	1	2	1	2	1	3	-
Male													
2001	11	-	-	-	-	-	2	3	-	2	1	3	-
2000	17	-	-	-	-	3	1	3	5	2	2	1	-
1999	6	-	-	-	-	1	-	1	-	2	1	1	-
Female													
2001	8	-	-	1	-	1	1	1	1	2	1	-	-
2000	14	-	-	1	1	1	-	2	1	4	2	2	-
1999	7	-	-	-	-	2	1	1	1	-	-	2	-
Other chronic lower respiratory diseases (J44,J47)													
All origins													
Both sexes													
2001	101,543	7	4	4	24	44	418	2,231	8,723	25,180	40,362	24,545	1
2000	99,655	7	3	5	13	28	370	2,125	8,325	25,145	40,029	23,603	2
1999	100,565	10	8	8	19	48	328	1,997	8,598	26,042	40,416	23,087	4
Male													
2001	49,407	4	2	3	15	24	200	1,196	4,518	13,027	20,096	10,321	1
2000	49,198	5	1	3	9	11	181	1,127	4,389	13,140	20,048	10,282	2

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other chronic lower respiratory diseases (J44,J47)													
All origins													
Male													
1999	50,817	7	3	5	10	15	172	1,067	4,615	13,980	20,598	10,342	3
Female													
2001	52,136	3	2	1	9	20	218	1,035	4,205	12,153	20,266	14,224	-
2000	50,457	2	2	2	4	17	189	998	3,936	12,005	19,981	13,321	-
1999	49,748	3	5	3	9	33	156	930	3,983	12,062	19,818	12,745	1
Hispanic													
Both sexes													
2001	2,235	-	2	1	1	4	19	66	185	513	774	670	-
2000	2,079	2	1	2	1	2	20	52	169	438	755	637	-
1999	2,195	3	2	2	1	6	20	55	177	511	759	659	-
Male													
2001	1,189	-	1	1	-	2	7	35	106	313	406	318	-
2000	1,127	1	1	1	1	1	5	25	109	268	425	290	-
1999	1,228	3	1	1	1	1	10	24	107	300	439	341	-
Female													
2001	1,046	-	1	-	1	2	12	31	79	200	368	352	-
2000	952	1	-	1	-	1	15	27	60	170	330	347	-
1999	967	-	1	1	-	5	10	31	70	211	320	318	-
Non-Hispanic [1]													
Both sexes													
2001	99,030	7	2	3	23	40	395	2,151	8,501	24,593	39,496	23,819	-
2000	97,305	5	2	3	12	26	348	2,060	8,118	24,633	39,180	22,917	1
1999	98,086	7	6	6	18	42	308	1,933	8,389	25,452	39,553	22,369	3

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other chronic lower respiratory diseases (J44,J47)													
Non-Hispanic [1]													
Male													
2001	48,070	4	1	2	15	22	190	1,152	4,388	12,674	19,642	9,980	-
2000	47,906	4	-	2	8	10	175	1,094	4,256	12,822	19,566	9,968	1
1999	49,436	4	2	4	9	14	162	1,036	4,489	13,628	20,112	9,974	2
Female													
2001	50,960	3	1	1	8	18	205	999	4,113	11,919	19,854	13,839	-
2000	49,399	1	2	1	4	16	173	966	3,862	11,811	19,614	12,949	-
1999	48,650	3	4	2	9	28	146	897	3,900	11,824	19,441	12,395	1
Non-Hispanic white													
Both sexes													
2001	92,267	4	1	2	12	29	314	1,835	7,685	22,734	37,199	22,452	-
2000	90,746	4	2	3	7	16	272	1,748	7,281	22,803	36,976	21,633	1
1999	91,269	1	3	5	12	28	238	1,649	7,524	23,501	37,153	21,153	2
Male													
2001	44,173	2	-	1	7	18	152	979	3,917	11,572	18,246	9,279	-
2000	44,077	4	-	2	4	4	140	908	3,774	11,705	18,230	9,305	1
1999	45,423	-	-	3	4	13	123	867	3,969	12,471	18,653	9,319	1
Female													
2001	48,094	2	1	1	5	11	162	856	3,768	11,162	18,953	13,173	-
2000	46,669	-	2	1	3	12	132	840	3,507	11,098	18,746	12,328	-
1999	45,846	1	3	2	8	15	115	782	3,555	11,030	18,500	11,834	1
Non-Hispanic black													
Both sexes													
2001	5,550	2	1	1	8	9	75	283	707	1,567	1,861	1,036	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other chronic lower respiratory diseases (J44,J47)													
Non-Hispanic black													
Both sexes													
2000	5,414	1	-	-	3	10	71	278	748	1,549	1,771	983	-
1999	5,695	6	3	1	6	11	68	258	770	1,639	1,961	971	1
Male													
2001	3,181	1	1	1	6	4	37	151	413	938	1,116	513	-
2000	3,126	-	-	-	3	6	33	161	432	953	1,045	493	-
1999	3,341	4	2	1	5	-	38	151	469	975	1,181	514	1
Female													
2001	2,369	1	-	-	2	5	38	132	294	629	745	523	-
2000	2,288	1	-	-	-	4	38	117	316	596	726	490	-
1999	2,354	2	1	-	1	11	30	107	301	664	780	457	-
Origin not stated [2]													
Both sexes													
2001	278	-	-	-	-	-	4	14	37	74	92	56	1
2000	271	-	-	-	-	-	2	13	38	74	94	49	1
1999	284	-	-	-	-	-	-	9	32	79	104	59	1
Male													
2001	148	-	-	-	-	-	3	9	24	40	48	23	1
2000	165	-	-	-	-	-	1	8	24	50	57	24	1
1999	153	-	-	-	-	-	-	7	19	52	47	27	1
Female													
2001	130	-	-	-	-	-	1	5	13	34	44	33	-
2000	106	-	-	-	-	-	1	5	14	24	37	25	-
1999	131	-	-	-	-	-	-	2	13	27	57	32	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Pneumoconioses and chemical effects (J60-J66,J68)													
All origins													
Both sexes													
2001	1,136	-	1	-	-	1	5	11	64	266	516	272	-
2000	1,161	-	-	-	-	3	7	12	57	281	484	317	-
1999	1,113	-	3	-	1	-	7	15	57	274	503	253	-
Male													
2001	1,095	-	1	-	-	1	5	9	62	261	494	262	-
2000	1,113	-	-	-	-	3	6	12	54	265	467	306	-
1999	1,079	-	3	-	1	-	5	15	57	269	486	243	-
Female													
2001	41	-	-	-	-	-	-	2	2	5	22	10	-
2000	48	-	-	-	-	-	1	-	3	16	17	11	-
1999	34	-	-	-	-	-	2	-	-	5	17	10	-
Hispanic													
Both sexes													
2001	26	-	1	-	-	-	2	2	2	7	6	6	-
2000	27	-	-	-	-	1	1	1	1	5	13	5	-
1999	13	-	-	-	-	-	1	1	1	5	2	3	-
Male													
2001	25	-	1	-	-	-	2	2	2	7	6	5	-
2000	26	-	-	-	-	1	1	1	1	5	13	4	-
1999	11	-	-	-	-	-	1	1	1	5	2	1	-
Female													
2001	1	-	-	-	-	-	-	-	-	-	-	1	-
2000	1	-	-	-	-	-	-	-	-	-	-	1	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Pneumoconioses and chemical effects (J60-J66,J68)													
Hispanic													
Female													
1999	2	-	-	-	-	-	-	-	-	-	-	2	-
Non-Hispanic [1]													
Both sexes													
2001	1,108	-	-	-	-	1	3	9	62	259	509	265	-
2000	1,133	-	-	-	-	2	6	11	56	276	470	312	-
1999	1,100	-	3	-	1	-	6	14	56	269	501	250	-
Male													
2001	1,068	-	-	-	-	1	3	7	60	254	487	256	-
2000	1,086	-	-	-	-	2	5	11	53	260	453	302	-
1999	1,068	-	3	-	1	-	4	14	56	264	484	242	-
Female													
2001	40	-	-	-	-	-	-	2	2	5	22	9	-
2000	47	-	-	-	-	-	1	-	3	16	17	10	-
1999	32	-	-	-	-	-	2	-	-	5	17	8	-
Non-Hispanic white													
Both sexes													
2001	1,046	-	-	-	-	1	3	5	53	245	487	252	-
2000	1,072	-	-	-	-	2	3	11	54	261	442	299	-
1999	1,024	-	-	-	-	-	3	10	46	250	482	233	-
Male													
2001	1,008	-	-	-	-	1	3	5	51	240	465	243	-
2000	1,027	-	-	-	-	2	2	11	51	245	427	289	-
1999	995	-	-	-	-	-	2	10	46	246	465	226	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Pneumoconioses and chemical effects (J60-J66,J68)													
Non-Hispanic white													
Female													
2001	38	-	-	-	-	-	-	-	2	5	22	9	-
2000	45	-	-	-	-	-	1	-	3	16	15	10	-
1999	29	-	-	-	-	-	1	-	-	4	17	7	-
Non-Hispanic black													
Both sexes													
2001	56	-	-	-	-	-	-	4	8	12	20	12	-
2000	54	-	-	-	-	-	3	-	2	15	23	11	-
1999	67	-	3	-	1	-	3	4	8	16	16	16	-
Male													
2001	54	-	-	-	-	-	-	2	8	12	20	12	-
2000	52	-	-	-	-	-	3	-	2	15	21	11	-
1999	64	-	3	-	1	-	2	4	8	15	16	15	-
Female													
2001	2	-	-	-	-	-	-	2	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	2	-	-
1999	3	-	-	-	-	-	1	-	-	1	-	1	-
Origin not stated [2]													
Both sexes													
2001	2	-	-	-	-	-	-	-	-	-	1	1	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	2	-	-	-	-	-	-	-	-	-	1	1	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Pneumoconioses and chemical effects (J60-J66,J68)													
Origin not stated [2]													
Male													
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Pneumonitis due to solids and liquids (J69)													
All origins													
Both sexes													
2001	17,301	12	14	15	34	79	179	378	659	1,830	5,759	8,342	-
2000	16,636	10	11	13	37	62	181	366	598	1,899	5,505	7,954	-
1999	15,268	16	12	16	32	78	166	284	586	1,754	5,030	7,294	-
Male													
2001	8,576	6	6	7	22	40	117	232	384	1,106	3,145	3,511	-
2000	8,390	4	5	9	23	35	115	225	354	1,195	3,038	3,387	-
1999	7,755	9	3	6	22	47	97	165	365	1,083	2,875	3,083	-
Female													
2001	8,725	6	8	8	12	39	62	146	275	724	2,614	4,831	-
2000	8,246	6	6	4	14	27	66	141	244	704	2,467	4,567	-
1999	7,513	7	9	10	10	31	69	119	221	671	2,155	4,211	-
Hispanic													
Both sexes													
2001	555	-	3	5	3	4	10	18	33	69	152	258	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Pneumonitis due to solids and liquids (J69)													
Hispanic													
Both sexes													
2000	496	3	2	1	3	7	15	20	33	64	141	207	-
1999	413	2	2	5	2	4	3	8	26	57	118	186	-
Male													
2001	283	-	1	2	1	3	5	10	22	42	81	116	-
2000	253	-	-	-	2	2	8	15	20	35	87	84	-
1999	195	1	1	1	-	3	1	4	14	31	65	74	-
Female													
2001	272	-	2	3	2	1	5	8	11	27	71	142	-
2000	243	3	2	1	1	5	7	5	13	29	54	123	-
1999	218	1	1	4	2	1	2	4	12	26	53	112	-
Non-Hispanic [1]													
Both sexes													
2001	16,700	12	11	10	31	75	167	358	624	1,756	5,593	8,063	-
2000	16,094	7	9	12	34	55	166	343	562	1,833	5,346	7,727	-
1999	14,812	14	10	11	30	74	162	274	559	1,691	4,895	7,092	-
Male													
2001	8,270	6	5	5	21	37	110	221	361	1,061	3,057	3,386	-
2000	8,107	4	5	9	21	33	107	207	333	1,158	2,937	3,293	-
1999	7,534	8	2	5	22	44	95	159	350	1,046	2,803	3,000	-
Female													
2001	8,430	6	6	5	10	38	57	137	263	695	2,536	4,677	-
2000	7,987	3	4	3	13	22	59	136	229	675	2,409	4,434	-
1999	7,278	6	8	6	8	30	67	115	209	645	2,092	4,092	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Pneumonitis due to solids and liquids (J69)													
Non-Hispanic white													
Both sexes													
2001	14,843	9	9	6	22	58	131	280	518	1,469	5,012	7,329	-
2000	14,343	6	7	11	24	40	128	259	449	1,515	4,793	7,111	-
1999	13,139	11	6	7	20	53	118	209	446	1,400	4,361	6,508	-
Male													
2001	7,411	6	3	3	17	32	87	179	298	902	2,765	3,119	-
2000	7,214	4	4	8	16	21	85	154	270	945	2,648	3,059	-
1999	6,642	6	1	4	14	30	70	122	276	860	2,492	2,767	-
Female													
2001	7,432	3	6	3	5	26	44	101	220	567	2,247	4,210	-
2000	7,129	2	3	3	8	19	43	105	179	570	2,145	4,052	-
1999	6,497	5	5	3	6	23	48	87	170	540	1,869	3,741	-
Non-Hispanic black													
Both sexes													
2001	1,584	3	2	4	7	16	33	71	100	251	497	600	-
2000	1,507	1	2	1	9	14	37	79	102	289	468	505	-
1999	1,495	2	4	3	10	18	40	59	105	264	469	521	-
Male													
2001	731	-	2	2	2	4	20	38	60	140	252	211	-
2000	757	-	1	1	4	11	21	52	55	190	243	179	-
1999	768	1	1	1	8	12	22	33	68	168	261	193	-
Female													
2001	853	3	-	2	5	12	13	33	40	111	245	389	-
2000	750	1	1	-	5	3	16	27	47	99	225	326	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Pneumonitis due to solids and liquids (J69)													
Non-Hispanic black													
Female													
1999	727	1	3	2	2	6	18	26	37	96	208	328	-
Origin not stated [2]													
Both sexes													
2001	46	-	-	-	-	-	2	2	2	5	14	21	-
2000	46	-	-	-	-	-	-	3	3	2	18	20	-
1999	43	-	-	-	-	-	1	2	1	6	17	16	-
Male													
2001	23	-	-	-	-	-	2	1	1	3	7	9	-
2000	30	-	-	-	-	-	-	3	1	2	14	10	-
1999	26	-	-	-	-	-	1	2	1	6	7	9	-
Female													
2001	23	-	-	-	-	-	-	1	1	2	7	12	-
2000	16	-	-	-	-	-	-	-	2	-	4	10	-
1999	17	-	-	-	-	-	-	-	-	-	10	7	-
Other diseases of respiratory system (J00-J06,J30-J39, J67,J70-J98)													
All origins													
Both sexes													
2001	26,178	294	93	78	141	216	570	1,356	2,432	5,510	9,202	6,283	3
2000	25,535	273	87	74	131	253	615	1,332	2,411	5,448	8,900	6,008	3
1999	25,031	252	91	76	115	257	576	1,219	2,260	5,409	8,646	6,130	-
Male													
2001	12,495	186	47	43	76	104	304	737	1,322	3,025	4,453	2,197	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other diseases of respiratory system (J00-J06,J30-J39, J67,J70-J98)													
All origins													
Male													
2000	12,391	154	45	38	73	127	335	712	1,324	2,961	4,415	2,205	2
1999	12,110	145	55	39	69	127	317	637	1,213	3,005	4,328	2,175	-
Female													
2001	13,683	108	46	35	65	112	266	619	1,110	2,485	4,749	4,086	2
2000	13,144	119	42	36	58	126	280	620	1,087	2,487	4,485	3,803	1
1999	12,921	107	36	37	46	130	259	582	1,047	2,404	4,318	3,955	-
Hispanic													
Both sexes													
2001	1,275	41	17	7	18	17	39	106	165	287	363	215	-
2000	1,225	43	17	10	15	34	56	81	167	273	325	204	-
1999	1,157	38	10	8	15	23	45	94	157	269	307	191	-
Male													
2001	643	21	8	3	9	12	26	58	92	151	175	88	-
2000	608	23	8	4	8	19	34	46	96	139	165	66	-
1999	583	20	6	3	7	12	29	45	89	139	158	75	-
Female													
2001	632	20	9	4	9	5	13	48	73	136	188	127	-
2000	617	20	9	6	7	15	22	35	71	134	160	138	-
1999	574	18	4	5	8	11	16	49	68	130	149	116	-
Non-Hispanic [1]													
Both sexes													
2001	24,835	251	76	71	123	198	528	1,245	2,265	5,203	8,815	6,057	3

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other diseases of respiratory system (J00-J06,J30-J39, J67,J70-J98)													
Non-Hispanic [1]													
Both sexes													
2000	24,236	228	69	64	115	218	556	1,246	2,238	5,158	8,552	5,790	2
1999	23,802	211	81	68	100	233	529	1,120	2,092	5,127	8,315	5,926	-
Male													
2001	11,818	165	39	40	67	92	276	677	1,228	2,860	4,265	2,108	1
2000	11,752	130	37	34	65	108	298	663	1,227	2,815	4,239	2,135	1
1999	11,483	123	49	36	62	115	286	589	1,117	2,857	4,153	2,096	-
Female													
2001	13,017	86	37	31	56	106	252	568	1,037	2,343	4,550	3,949	2
2000	12,484	98	32	30	50	110	258	583	1,011	2,343	4,313	3,655	1
1999	12,319	88	32	32	38	118	243	531	975	2,270	4,162	3,830	-
Non-Hispanic white													
Both sexes													
2001	21,835	116	44	52	82	122	350	883	1,860	4,597	8,133	5,595	1
2000	21,397	122	46	43	71	148	385	928	1,847	4,557	7,866	5,384	-
1999	20,959	111	52	47	75	155	351	826	1,718	4,511	7,642	5,471	-
Male													
2001	10,433	76	20	31	46	52	202	498	1,018	2,574	3,963	1,953	-
2000	10,349	77	22	24	38	74	205	493	1,021	2,508	3,913	1,974	-
1999	10,154	66	28	26	51	86	200	443	919	2,561	3,848	1,926	-
Female													
2001	11,402	40	24	21	36	70	148	385	842	2,023	4,170	3,642	1
2000	11,048	45	24	19	33	74	180	435	826	2,049	3,953	3,410	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other diseases of respiratory system (J00-J06,J30-J39, J67,J70-J98)													
Non-Hispanic white													
Female													
1999	10,805	45	24	21	24	69	151	383	799	1,950	3,794	3,545	-
Non-Hispanic black													
Both sexes													
2001	2,462	116	26	17	38	67	163	326	343	482	527	355	2
2000	2,336	102	20	18	39	58	152	284	325	495	527	314	2
1999	2,339	90	26	19	21	73	160	258	317	503	523	349	-
Male													
2001	1,116	76	14	8	19	34	68	157	173	222	230	114	1
2000	1,126	50	14	10	23	30	84	151	166	242	240	115	1
1999	1,081	52	19	9	9	27	74	132	164	237	235	123	-
Female													
2001	1,346	40	12	9	19	33	95	169	170	260	297	241	1
2000	1,210	52	6	8	16	28	68	133	159	253	287	199	1
1999	1,258	38	7	10	12	46	86	126	153	266	288	226	-
Origin not stated [2]													
Both sexes													
2001	68	2	-	-	-	1	3	5	2	20	24	11	-
2000	74	2	1	-	1	1	3	5	6	17	23	14	1
1999	72	3	-	-	-	1	2	5	11	13	24	13	-
Male													
2001	34	-	-	-	-	-	2	2	2	14	13	1	-
2000	31	1	-	-	-	-	3	3	1	7	11	4	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other diseases of respiratory system (J00-J06,J30-J39, J67,J70-J98)													
Origin not stated [2]													
Male													
1999	44	2	-	-	-	-	2	3	7	9	17	4	-
Female													
2001	34	2	-	-	-	1	1	3	-	6	11	10	-
2000	43	1	1	-	1	1	-	2	5	10	12	10	-
1999	28	1	-	-	-	1	-	2	4	4	7	9	-
Peptic ulcer (K25-K28)													
All origins													
Both sexes													
2001	4,491	-	1	5	6	32	118	284	418	718	1,438	1,470	1
2000	4,558	-	-	-	9	14	113	307	443	748	1,415	1,509	-
1999	4,604	2	2	-	2	21	123	255	422	836	1,500	1,439	2
Male													
2001	2,074	-	1	2	3	21	87	186	273	397	649	454	1
2000	2,201	-	-	-	7	6	79	217	284	442	667	499	-
1999	2,189	-	-	-	2	12	76	169	265	504	712	448	1
Female													
2001	2,417	-	-	3	3	11	31	98	145	321	789	1,016	-
2000	2,357	-	-	-	2	8	34	90	159	306	748	1,010	-
1999	2,415	2	2	-	-	9	47	86	157	332	788	991	1
Hispanic													
Both sexes													
2001	166	-	-	-	1	-	1	16	22	33	54	38	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Peptic ulcer (K25-K28)													
Hispanic													
Both sexes													
2000	186	-	-	-	-	-	9	17	28	36	50	46	-
1999	172	-	1	-	-	8	4	15	25	37	44	38	-
Male													
2001	99	-	-	-	1	-	1	11	19	18	37	11	1
2000	97	-	-	-	-	-	6	14	23	21	17	16	-
1999	101	-	-	-	-	5	3	10	18	22	27	16	-
Female													
2001	67	-	-	-	-	-	-	5	3	15	17	27	-
2000	89	-	-	-	-	-	3	3	5	15	33	30	-
1999	71	-	1	-	-	3	1	5	7	15	17	22	-
Non-Hispanic [1]													
Both sexes													
2001	4,308	-	1	5	5	32	117	267	393	681	1,380	1,427	-
2000	4,357	-	-	-	9	14	102	290	414	709	1,361	1,458	-
1999	4,415	2	1	-	2	13	118	239	393	796	1,451	1,398	2
Male													
2001	1,966	-	1	2	2	21	86	175	252	375	609	443	-
2000	2,099	-	-	-	7	6	72	203	261	419	648	483	-
1999	2,079	-	-	-	2	7	72	158	244	481	682	432	1
Female													
2001	2,342	-	-	3	3	11	31	92	141	306	771	984	-
2000	2,258	-	-	-	2	8	30	87	153	290	713	975	-
1999	2,336	2	1	-	-	6	46	81	149	315	769	966	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Peptic ulcer (K25-K28)													
Non-Hispanic white													
Both sexes													
2001	3,766	-	1	4	2	16	77	197	308	583	1,248	1,330	-
2000	3,825	-	-	-	9	9	70	210	343	609	1,233	1,342	-
1999	3,859	1	1	-	2	7	73	171	322	677	1,313	1,291	1
Male													
2001	1,649	-	1	2	1	11	55	122	194	311	543	409	-
2000	1,799	-	-	-	7	4	49	150	217	357	580	435	-
1999	1,783	-	-	-	2	3	41	111	195	409	622	399	1
Female													
2001	2,117	-	-	2	1	5	22	75	114	272	705	921	-
2000	2,026	-	-	-	2	5	21	60	126	252	653	907	-
1999	2,076	1	1	-	-	4	32	60	127	268	691	892	-
Non-Hispanic black													
Both sexes													
2001	403	-	-	1	2	15	35	59	68	73	83	67	-
2000	420	-	-	-	-	4	29	70	63	79	91	84	-
1999	443	1	-	-	-	4	43	60	65	92	98	79	1
Male													
2001	238	-	-	-	1	10	27	46	47	47	39	21	-
2000	228	-	-	-	-	2	21	45	40	48	45	27	-
1999	240	-	-	-	-	3	30	43	46	55	39	24	-
Female													
2001	165	-	-	1	1	5	8	13	21	26	44	46	-
2000	192	-	-	-	-	2	8	25	23	31	46	57	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Peptic ulcer (K25-K28)													
Non-Hispanic black													
Female													
1999	203	1	-	-	-	1	13	17	19	37	59	55	1
Origin not stated [2]													
Both sexes													
2001	17	-	-	-	-	-	-	1	3	4	4	5	-
2000	15	-	-	-	-	-	2	-	1	3	4	5	-
1999	17	-	-	-	-	-	1	1	4	3	5	3	-
Male													
2001	9	-	-	-	-	-	-	-	2	4	3	-	-
2000	5	-	-	-	-	-	1	-	-	2	2	-	-
1999	9	-	-	-	-	-	1	1	3	1	3	-	-
Female													
2001	8	-	-	-	-	-	-	1	1	-	1	5	-
2000	10	-	-	-	-	-	1	-	1	1	2	5	-
1999	8	-	-	-	-	-	-	-	1	2	2	3	-
Diseases of appendix (K35-K38)													
All origins													
Both sexes													
2001	428	1	8	6	15	15	26	40	45	89	90	93	-
2000	435	3	5	15	11	11	27	43	60	71	121	68	-
1999	390	3	2	26	5	6	24	49	48	69	96	62	-
Male													
2001	252	-	7	5	8	11	17	28	30	57	52	37	-
2000	237	2	2	4	7	10	16	29	43	40	60	24	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Diseases of appendix (K35-K38)													
All origins													
Male													
1999	236	1	2	15	5	5	17	37	34	38	47	35	-
Female													
2001	176	1	1	1	7	4	9	12	15	32	38	56	-
2000	198	1	3	11	4	1	11	14	17	31	61	44	-
1999	154	2	-	11	-	1	7	12	14	31	49	27	-
Hispanic													
Both sexes													
2001	30	1	4	1	2	1	1	5	2	5	7	1	-
2000	25	1	1	4	5	2	1	1	5	2	3	-	-
1999	29	1	-	8	2	-	4	4	3	3	3	1	-
Male													
2001	16	-	3	1	-	-	-	3	1	5	3	-	-
2000	16	-	1	3	3	2	-	1	4	-	2	-	-
1999	20	1	-	4	2	-	4	3	2	2	1	1	-
Female													
2001	14	1	1	-	2	1	1	2	1	-	4	1	-
2000	9	1	-	1	2	-	1	-	1	2	1	-	-
1999	9	-	-	4	-	-	-	1	1	1	2	-	-
Non-Hispanic [1]													
Both sexes													
2001	398	-	4	5	13	14	25	35	43	84	83	92	-
2000	410	2	4	11	6	9	26	42	55	69	118	68	-
1999	358	2	2	17	3	6	20	44	45	65	93	61	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Diseases of appendix (K35-K38)													
Non-Hispanic [1]													
Male													
2001	236	-	4	4	8	11	17	25	29	52	49	37	-
2000	221	2	1	1	4	8	16	28	39	40	58	24	-
1999	213	-	2	10	3	5	13	33	32	35	46	34	-
Female													
2001	162	-	-	1	5	3	8	10	14	32	34	55	-
2000	189	-	3	10	2	1	10	14	16	29	60	44	-
1999	145	2	-	7	-	1	7	11	13	30	47	27	-
Non-Hispanic white													
Both sexes													
2001	313	-	1	3	8	7	16	24	27	75	70	82	-
2000	323	2	3	9	3	2	17	28	45	52	97	65	-
1999	284	2	-	10	1	4	15	33	27	54	81	57	-
Male													
2001	182	-	1	2	5	5	12	17	21	47	43	29	-
2000	167	2	1	1	2	2	9	18	30	31	49	22	-
1999	163	-	-	6	1	3	9	24	22	28	38	32	-
Female													
2001	131	-	-	1	3	2	4	7	6	28	27	53	-
2000	156	-	2	8	1	-	8	10	15	21	48	43	-
1999	121	2	-	4	-	1	6	9	5	26	43	25	-
Non-Hispanic black													
Both sexes													
2001	72	-	2	2	5	6	8	9	16	7	11	6	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Diseases of appendix (K35-K38)													
Non-Hispanic black													
Both sexes													
2000	77	-	-	2	2	7	9	12	10	16	16	3	-
1999	67	-	2	7	1	2	5	10	17	9	11	3	-
Male													
2001	47	-	2	2	3	6	5	7	8	4	5	5	-
2000	48	-	-	-	1	6	7	8	9	9	6	2	-
1999	47	-	2	4	1	2	4	9	9	6	8	2	-
Female													
2001	25	-	-	-	2	-	3	2	8	3	6	1	-
2000	29	-	-	2	1	1	2	4	1	7	10	1	-
1999	20	-	-	3	-	-	1	1	8	3	3	1	-
Origin not stated [2]													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	-	1	-	-	-	1	-	1	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	-	1	-	-	-	1	-	1	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Hernia (K40-K46)													
All origins													
Both sexes													
2001	1,522	43	2	3	3	15	34	82	131	213	443	553	-
2000	1,522	46	4	3	5	5	31	84	113	210	466	555	-
1999	1,464	44	3	2	3	16	39	55	107	215	451	529	-
Male													
2001	587	28	1	-	1	10	16	44	64	94	173	156	-
2000	586	28	3	3	3	1	13	45	49	92	195	154	-
1999	597	26	2	1	3	8	23	32	60	93	192	157	-
Female													
2001	935	15	1	3	2	5	18	38	67	119	270	397	-
2000	936	18	1	-	2	4	18	39	64	118	271	401	-
1999	867	18	1	1	-	8	16	23	47	122	259	372	-
Hispanic													
Both sexes													
2001	92	9	-	-	2	1	7	8	9	12	24	20	-
2000	85	9	-	1	1	1	1	7	9	14	23	19	-
1999	66	3	1	-	-	-	3	4	5	16	17	17	-
Male													
2001	32	7	-	-	1	-	3	3	3	4	6	5	-
2000	33	6	-	1	-	1	1	5	3	3	10	3	-
1999	27	2	1	-	-	-	2	3	4	6	6	3	-
Female													
2001	60	2	-	-	1	1	4	5	6	8	18	15	-
2000	52	3	-	-	1	-	-	2	6	11	13	16	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Hernia (K40-K46)													
Hispanic													
Female													
1999	39	1	-	-	-	-	1	1	1	10	11	14	-
Non-Hispanic [1]													
Both sexes													
2001	1,425	33	2	3	1	14	27	74	121	201	417	532	-
2000	1,431	37	4	2	4	4	30	76	103	194	442	535	-
1999	1,390	41	2	2	3	16	36	50	102	198	431	509	-
Male													
2001	552	20	1	-	-	10	13	41	60	90	166	151	-
2000	550	22	3	2	3	-	12	40	45	88	184	151	-
1999	566	24	1	1	3	8	21	28	56	86	185	153	-
Female													
2001	873	13	1	3	1	4	14	33	61	111	251	381	-
2000	881	15	1	-	1	4	18	36	58	106	258	384	-
1999	824	17	1	1	-	8	15	22	46	112	246	356	-
Non-Hispanic white													
Both sexes													
2001	1,248	22	2	1	-	12	18	59	97	173	374	490	-
2000	1,266	27	2	2	2	3	21	59	90	169	399	492	-
1999	1,227	31	2	2	2	10	27	36	89	167	390	471	-
Male													
2001	469	11	1	-	-	8	10	29	46	81	149	134	-
2000	486	17	2	2	1	-	9	29	40	78	170	138	-
1999	505	18	1	1	2	5	18	22	53	79	171	135	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Hernia (K40-K46)													
Non-Hispanic white													
Female													
2001	779	11	1	1	-	4	8	30	51	92	225	356	-
2000	780	10	-	-	1	3	12	30	50	91	229	354	-
1999	722	13	1	1	-	5	9	14	36	88	219	336	-
Non-Hispanic black													
Both sexes													
2001	155	10	-	1	1	2	9	15	23	25	33	36	-
2000	142	7	2	-	2	1	8	13	12	21	36	40	-
1999	139	9	-	-	1	4	9	12	12	24	37	31	-
Male													
2001	74	8	-	-	-	2	3	12	13	7	14	15	-
2000	52	3	1	-	2	-	2	9	4	9	11	11	-
1999	53	6	-	-	1	1	3	6	2	7	13	14	-
Female													
2001	81	2	-	1	1	-	6	3	10	18	19	21	-
2000	90	4	1	-	-	1	6	4	8	12	25	29	-
1999	86	3	-	-	-	3	6	6	10	17	24	17	-
Origin not stated [2]													
Both sexes													
2001	5	1	-	-	-	-	-	-	1	-	2	1	-
2000	6	-	-	-	-	-	-	1	1	2	1	1	-
1999	8	-	-	-	-	-	-	1	-	1	3	3	-
Male													
2001	3	1	-	-	-	-	-	-	1	-	1	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Hernia (K40-K46)													
Origin not stated [2]													
Male													
2000	3	-	-	-	-	-	-	-	1	1	1	-	-
1999	4	-	-	-	-	-	-	1	-	1	1	1	-
Female													
2001	2	-	-	-	-	-	-	-	-	-	1	1	-
2000	3	-	-	-	-	-	-	1	-	1	-	1	-
1999	4	-	-	-	-	-	-	-	-	-	2	2	-
Chronic liver disease and cirrhosis (K70,K73-K74)													
All origins													
Both sexes													
2001	27,035	3	-	2	26	387	3,336	7,259	5,750	5,486	3,801	979	6
2000	26,552	3	3	3	36	415	3,371	6,654	5,774	5,482	3,829	979	3
1999	26,259	5	2	2	26	407	3,302	6,368	5,637	5,642	3,901	965	2
Male													
2001	17,393	1	-	2	12	259	2,206	5,295	4,028	3,335	1,833	416	6
2000	17,214	3	1	2	19	280	2,314	4,975	4,005	3,317	1,874	421	3
1999	17,115	4	-	1	11	257	2,314	4,829	3,943	3,454	1,905	395	2
Female													
2001	9,642	2	-	-	14	128	1,130	1,964	1,722	2,151	1,968	563	-
2000	9,338	-	2	1	17	135	1,057	1,679	1,769	2,165	1,955	558	-
1999	9,144	1	2	1	15	150	988	1,539	1,694	2,188	1,996	570	-
Hispanic													
Both sexes													
2001	3,301	-	-	-	5	81	530	951	721	618	335	59	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Chronic liver disease and cirrhosis (K70,K73-K74)													
Hispanic													
Both sexes													
2000	3,187	-	1	-	7	92	501	854	708	628	323	73	-
1999	2,972	-	-	-	8	80	518	787	638	583	285	73	-
Male													
2001	2,410	-	-	-	2	67	425	782	555	386	170	22	1
2000	2,312	-	-	-	5	82	399	707	531	395	163	30	-
1999	2,143	-	-	-	5	62	421	647	481	360	136	31	-
Female													
2001	891	-	-	-	3	14	105	169	166	232	165	37	-
2000	875	-	1	-	2	10	102	147	177	233	160	43	-
1999	829	-	-	-	3	18	97	140	157	223	149	42	-
Non-Hispanic [1]													
Both sexes													
2001	23,629	3	-	1	21	305	2,796	6,280	5,002	4,846	3,455	917	3
2000	23,251	3	2	3	29	319	2,856	5,762	5,041	4,833	3,498	903	2
1999	23,185	5	2	2	18	326	2,767	5,547	4,978	5,044	3,606	888	2
Male													
2001	14,914	1	-	1	10	192	1,777	4,493	3,450	2,934	1,661	392	3
2000	14,814	3	1	2	14	195	1,903	4,237	3,452	2,908	1,708	389	2
1999	14,896	4	-	1	6	194	1,881	4,155	3,444	3,081	1,765	363	2
Female													
2001	8,715	2	-	-	11	113	1,019	1,787	1,552	1,912	1,794	525	-
2000	8,437	-	1	1	15	124	953	1,525	1,589	1,925	1,790	514	-
1999	8,289	1	2	1	12	132	886	1,392	1,534	1,963	1,841	525	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Chronic liver disease and cirrhosis (K70,K73-K74)													
Non-Hispanic white													
Both sexes													
2001	20,065	3	-	1	5	227	2,212	5,035	4,189	4,307	3,222	862	2
2000	19,739	-	2	2	14	217	2,240	4,637	4,235	4,285	3,264	842	1
1999	19,602	4	-	1	7	243	2,148	4,430	4,117	4,482	3,347	821	2
Male													
2001	12,609	1	-	1	4	145	1,403	3,645	2,875	2,606	1,557	370	2
2000	12,640	-	1	1	6	140	1,518	3,477	2,935	2,602	1,597	362	1
1999	12,613	4	-	-	3	151	1,478	3,365	2,885	2,746	1,641	338	2
Female													
2001	7,456	2	-	-	1	82	809	1,390	1,314	1,701	1,665	492	-
2000	7,099	-	1	1	8	77	722	1,160	1,300	1,683	1,667	480	-
1999	6,989	-	-	1	4	92	670	1,065	1,232	1,736	1,706	483	-
Non-Hispanic black													
Both sexes													
2001	2,733	-	-	-	10	45	420	993	649	412	169	34	1
2000	2,705	2	-	1	12	56	446	929	622	429	163	44	1
1999	2,793	-	2	1	10	54	453	905	691	432	193	52	-
Male													
2001	1,819	-	-	-	2	25	273	702	462	268	75	11	1
2000	1,709	2	-	1	6	26	279	633	408	255	81	17	1
1999	1,812	-	-	1	3	27	293	649	453	273	96	17	-
Female													
2001	914	-	-	-	8	20	147	291	187	144	94	23	-
2000	996	-	-	-	6	30	167	296	214	174	82	27	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Chronic liver disease and cirrhosis (K70,K73-K74)													
Non-Hispanic black													
Female													
1999	981	-	2	-	7	27	160	256	238	159	97	35	-
Origin not stated [2]													
Both sexes													
2001	105	-	-	1	-	1	10	28	27	22	11	3	2
2000	114	-	-	-	-	4	14	38	25	21	8	3	1
1999	102	-	-	-	-	1	17	34	21	15	10	4	-
Male													
2001	69	-	-	1	-	-	4	20	23	15	2	2	2
2000	88	-	-	-	-	3	12	31	22	14	3	2	1
1999	76	-	-	-	-	1	12	27	18	13	4	1	-
Female													
2001	36	-	-	-	-	1	6	8	4	7	9	1	-
2000	26	-	-	-	-	1	2	7	3	7	5	1	-
1999	26	-	-	-	-	-	5	7	3	2	6	3	-
Alcoholic liver disease (K70)													
All origins													
Both sexes													
2001	12,207	-	-	-	8	264	2,145	4,203	2,913	1,848	717	104	5
2000	12,109	-	-	-	8	279	2,226	3,888	2,969	1,891	751	95	2
1999	11,958	-	-	-	8	259	2,186	3,802	2,859	1,935	789	118	2
Male													
2001	8,932	-	-	-	5	179	1,446	3,099	2,242	1,379	505	72	5
2000	8,985	-	-	-	5	202	1,533	2,968	2,216	1,433	561	65	2

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Alcoholic liver disease (K70)													
All origins													
Male													
1999	8,922	-	-	-	6	165	1,520	2,927	2,194	1,497	538	73	2
Female													
2001	3,275	-	-	-	3	85	699	1,104	671	469	212	32	-
2000	3,124	-	-	-	3	77	693	920	753	458	190	30	-
1999	3,036	-	-	-	2	94	666	875	665	438	251	45	-
Hispanic													
Both sexes													
2001	1,729	-	-	-	-	65	341	582	396	240	94	10	1
2000	1,680	-	-	-	4	66	353	526	400	247	78	6	-
1999	1,547	-	-	-	2	50	342	499	356	213	69	16	-
Male													
2001	1,457	-	-	-	-	55	282	494	338	210	68	9	1
2000	1,442	-	-	-	4	62	282	455	344	221	69	5	-
1999	1,294	-	-	-	2	41	276	426	307	180	52	10	-
Female													
2001	272	-	-	-	-	10	59	88	58	30	26	1	-
2000	238	-	-	-	-	4	71	71	56	26	9	1	-
1999	253	-	-	-	-	9	66	73	49	33	17	6	-
Non-Hispanic [1]													
Both sexes													
2001	10,423	-	-	-	8	198	1,801	3,603	2,499	1,596	622	94	2
2000	10,370	-	-	-	4	211	1,865	3,337	2,554	1,639	671	88	1
1999	10,353	-	-	-	6	208	1,831	3,283	2,491	1,713	717	102	2

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Alcoholic liver disease (K70)													
Non-Hispanic [1]													
Male													
2001	7,431	-	-	-	5	124	1,162	2,592	1,888	1,158	437	63	2
2000	7,492	-	-	-	1	139	1,243	2,492	1,859	1,208	490	59	1
1999	7,580	-	-	-	4	123	1,235	2,485	1,877	1,308	483	63	2
Female													
2001	2,992	-	-	-	3	74	639	1,011	611	438	185	31	-
2000	2,878	-	-	-	3	72	622	845	695	431	181	29	-
1999	2,773	-	-	-	2	85	596	798	614	405	234	39	-
Non-Hispanic white													
Both sexes													
2001	8,635	-	-	-	2	146	1,431	2,935	2,092	1,381	566	80	2
2000	8,556	-	-	-	2	150	1,454	2,703	2,126	1,439	601	81	-
1999	8,470	-	-	-	3	155	1,414	2,649	2,045	1,480	636	86	2
Male													
2001	6,209	-	-	-	2	96	922	2,152	1,583	997	401	54	2
2000	6,296	-	-	-	-	105	979	2,068	1,579	1,065	446	54	-
1999	6,288	-	-	-	2	95	967	2,041	1,566	1,131	431	53	2
Female													
2001	2,426	-	-	-	-	50	509	783	509	384	165	26	-
2000	2,260	-	-	-	2	45	475	635	547	374	155	27	-
1999	2,182	-	-	-	1	60	447	608	479	349	205	33	-
Non-Hispanic black													
Both sexes													
2001	1,331	-	-	-	1	26	244	508	324	176	43	9	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Alcoholic liver disease (K70)													
Non-Hispanic black													
Both sexes													
2000	1,370	-	-	-	-	34	285	505	321	169	50	5	1
1999	1,445	-	-	-	2	28	281	507	358	191	64	14	-
Male													
2001	926	-	-	-	-	12	160	349	239	135	26	5	-
2000	910	-	-	-	-	17	185	339	208	123	33	4	1
1999	1,001	-	-	-	2	14	178	358	254	146	41	8	-
Female													
2001	405	-	-	-	1	14	84	159	85	41	17	4	-
2000	460	-	-	-	-	17	100	166	113	46	17	1	-
1999	444	-	-	-	-	14	103	149	104	45	23	6	-
Origin not stated [2]													
Both sexes													
2001	55	-	-	-	-	1	3	18	18	12	1	-	2
2000	59	-	-	-	-	2	8	25	15	5	2	1	1
1999	58	-	-	-	-	1	13	20	12	9	3	-	-
Male													
2001	44	-	-	-	-	-	2	13	16	11	-	-	2
2000	51	-	-	-	-	1	8	21	13	4	2	1	1
1999	48	-	-	-	-	1	9	16	10	9	3	-	-
Female													
2001	11	-	-	-	-	1	1	5	2	1	1	-	-
2000	8	-	-	-	-	1	-	4	2	1	-	-	-
1999	10	-	-	-	-	-	4	4	2	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other chronic liver disease and cirrhosis (K73-K74)													
All origins													
Both sexes													
2001	14,828	3	-	2	18	123	1,191	3,056	2,837	3,638	3,084	875	1
2000	14,443	3	3	3	28	136	1,145	2,766	2,805	3,591	3,078	884	1
1999	14,301	5	2	2	18	148	1,116	2,566	2,778	3,707	3,112	847	-
Male													
2001	8,461	1	-	2	7	80	760	2,196	1,786	1,956	1,328	344	1
2000	8,229	3	1	2	14	78	781	2,007	1,789	1,884	1,313	356	1
1999	8,193	4	-	1	5	92	794	1,902	1,749	1,957	1,367	322	-
Female													
2001	6,367	2	-	-	11	43	431	860	1,051	1,682	1,756	531	-
2000	6,214	-	2	1	14	58	364	759	1,016	1,707	1,765	528	-
1999	6,108	1	2	1	13	56	322	664	1,029	1,750	1,745	525	-
Hispanic													
Both sexes													
2001	1,572	-	-	-	5	16	189	369	325	378	241	49	-
2000	1,507	-	1	-	3	26	148	328	308	381	245	67	-
1999	1,425	-	-	-	6	30	176	288	282	370	216	57	-
Male													
2001	953	-	-	-	2	12	143	288	217	176	102	13	-
2000	870	-	-	-	1	20	117	252	187	174	94	25	-
1999	849	-	-	-	3	21	145	221	174	180	84	21	-
Female													
2001	619	-	-	-	3	4	46	81	108	202	139	36	-
2000	637	-	1	-	2	6	31	76	121	207	151	42	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other chronic liver disease and cirrhosis (K73-K74)													
Hispanic													
Female													
1999	576	-	-	-	3	9	31	67	108	190	132	36	-
Non-Hispanic [1]													
Both sexes													
2001	13,206	3	-	1	13	107	995	2,677	2,503	3,250	2,833	823	1
2000	12,881	3	2	3	25	108	991	2,425	2,487	3,194	2,827	815	1
1999	12,832	5	2	2	12	118	936	2,264	2,487	3,331	2,889	786	-
Male													
2001	7,483	1	-	1	5	68	615	1,901	1,562	1,776	1,224	329	1
2000	7,322	3	1	2	13	56	660	1,745	1,593	1,700	1,218	330	1
1999	7,316	4	-	1	2	71	646	1,670	1,567	1,773	1,282	300	-
Female													
2001	5,723	2	-	-	8	39	380	776	941	1,474	1,609	494	-
2000	5,559	-	1	1	12	52	331	680	894	1,494	1,609	485	-
1999	5,516	1	2	1	10	47	290	594	920	1,558	1,607	486	-
Non-Hispanic white													
Both sexes													
2001	11,430	3	-	1	3	81	781	2,100	2,097	2,926	2,656	782	-
2000	11,183	-	2	2	12	67	786	1,934	2,109	2,846	2,663	761	1
1999	11,132	4	-	1	4	88	734	1,781	2,072	3,002	2,711	735	-
Male													
2001	6,400	1	-	1	2	49	481	1,493	1,292	1,609	1,156	316	-
2000	6,344	-	1	1	6	35	539	1,409	1,356	1,537	1,151	308	1
1999	6,325	4	-	-	1	56	511	1,324	1,319	1,615	1,210	285	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other chronic liver disease and cirrhosis (K73-K74)													
Non-Hispanic white													
Female													
2001	5,030	2	-	-	1	32	300	607	805	1,317	1,500	466	-
2000	4,839	-	1	1	6	32	247	525	753	1,309	1,512	453	-
1999	4,807	-	-	1	3	32	223	457	753	1,387	1,501	450	-
Non-Hispanic black													
Both sexes													
2001	1,402	-	-	-	9	19	176	485	325	236	126	25	1
2000	1,335	2	-	1	12	22	161	424	301	260	113	39	-
1999	1,348	-	2	1	8	26	172	398	333	241	129	38	-
Male													
2001	893	-	-	-	2	13	113	353	223	133	49	6	1
2000	799	2	-	1	6	9	94	294	200	132	48	13	-
1999	811	-	-	1	1	13	115	291	199	127	55	9	-
Female													
2001	509	-	-	-	7	6	63	132	102	103	77	19	-
2000	536	-	-	-	6	13	67	130	101	128	65	26	-
1999	537	-	2	-	7	13	57	107	134	114	74	29	-
Origin not stated [2]													
Both sexes													
2001	50	-	-	1	-	-	7	10	9	10	10	3	-
2000	55	-	-	-	-	2	6	13	10	16	6	2	-
1999	44	-	-	-	-	-	4	14	9	6	7	4	-
Male													
2001	25	-	-	1	-	-	2	7	7	4	2	2	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other chronic liver disease and cirrhosis (K73-K74)													
Origin not stated [2]													
Male													
2000	37	-	-	-	-	2	4	10	9	10	1	1	-
1999	28	-	-	-	-	-	3	11	8	4	1	1	-
Female													
2001	25	-	-	-	-	-	5	3	2	6	8	1	-
2000	18	-	-	-	-	-	2	3	1	6	5	1	-
1999	16	-	-	-	-	-	1	3	1	2	6	3	-
Cholelithiasis and other disorders of gallbladder (K80-K82)													
All origins													
Both sexes													
2001	2,965	1	-	1	7	19	44	137	242	526	945	1,043	-
2000	2,810	-	-	2	9	19	53	109	189	450	926	1,053	-
1999	2,830	2	-	1	10	16	59	124	203	475	948	992	-
Male													
2001	1,341	1	-	-	3	9	25	79	142	269	464	349	-
2000	1,244	-	-	1	4	7	27	53	110	242	461	339	-
1999	1,248	1	-	1	5	4	30	58	120	252	447	330	-
Female													
2001	1,624	-	-	1	4	10	19	58	100	257	481	694	-
2000	1,566	-	-	1	5	12	26	56	79	208	465	714	-
1999	1,582	1	-	-	5	12	29	66	83	223	501	662	-
Hispanic													
Both sexes													
2001	197	-	-	-	2	3	8	13	23	51	47	50	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)													
Hispanic													
Both sexes													
2000	183	-	-	-	2	3	5	18	19	41	51	44	-
1999	185	-	-	-	-	3	11	14	20	34	57	46	-
Male													
2001	92	-	-	-	-	1	3	5	16	25	25	17	-
2000	99	-	-	-	-	1	5	9	15	25	32	12	-
1999	84	-	-	-	-	1	6	6	12	15	29	15	-
Female													
2001	105	-	-	-	2	2	5	8	7	26	22	33	-
2000	84	-	-	-	2	2	-	9	4	16	19	32	-
1999	101	-	-	-	-	2	5	8	8	19	28	31	-
Non-Hispanic [1]													
Both sexes													
2001	2,764	1	-	1	5	16	36	124	219	473	896	993	-
2000	2,619	-	-	2	7	16	48	91	170	405	873	1,007	-
1999	2,639	2	-	1	10	13	48	110	183	438	889	945	-
Male													
2001	1,248	1	-	-	3	8	22	74	126	243	439	332	-
2000	1,141	-	-	1	4	6	22	44	95	215	428	326	-
1999	1,163	1	-	1	5	3	24	52	108	236	418	315	-
Female													
2001	1,516	-	-	1	2	8	14	50	93	230	457	661	-
2000	1,478	-	-	1	3	10	26	47	75	190	445	681	-
1999	1,476	1	-	-	5	10	24	58	75	202	471	630	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)													
Non-Hispanic white													
Both sexes													
2001	2,368	1	-	1	3	12	25	86	168	381	787	904	-
2000	2,272	-	-	1	5	7	33	58	132	325	775	936	-
1999	2,299	-	-	1	6	10	26	85	134	361	805	871	-
Male													
2001	1,065	1	-	-	2	5	17	52	96	200	395	297	-
2000	1,012	-	-	1	3	4	17	31	77	181	391	307	-
1999	1,017	-	-	1	4	1	14	40	79	201	389	288	-
Female													
2001	1,303	-	-	1	1	7	8	34	72	181	392	607	-
2000	1,260	-	-	-	2	3	16	27	55	144	384	629	-
1999	1,282	-	-	-	2	9	12	45	55	160	416	583	-
Non-Hispanic black													
Both sexes													
2001	309	-	-	-	2	4	11	31	42	78	77	64	-
2000	274	-	-	1	2	8	14	27	29	61	75	57	-
1999	257	2	-	-	4	3	20	21	35	59	65	48	-
Male													
2001	130	-	-	-	1	3	5	18	24	34	21	24	-
2000	92	-	-	-	1	2	4	10	14	23	26	12	-
1999	102	1	-	-	1	2	9	9	19	25	23	13	-
Female													
2001	179	-	-	-	1	1	6	13	18	44	56	40	-
2000	182	-	-	1	1	6	10	17	15	38	49	45	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)													
Non-Hispanic black													
Female													
1999	155	1	-	-	3	1	11	12	16	34	42	35	-
Origin not stated [2]													
Both sexes													
2001	4	-	-	-	-	-	-	-	-	2	2	-	-
2000	8	-	-	-	-	-	-	-	-	4	2	2	-
1999	6	-	-	-	-	-	-	-	-	3	2	1	-
Male													
2001	1	-	-	-	-	-	-	-	-	1	-	-	-
2000	4	-	-	-	-	-	-	-	-	2	1	1	-
1999	1	-	-	-	-	-	-	-	-	1	-	-	-
Female													
2001	3	-	-	-	-	-	-	-	-	1	2	-	-
2000	4	-	-	-	-	-	-	-	-	2	1	1	-
1999	5	-	-	-	-	-	-	-	-	2	2	1	-
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)													
All origins													
Both sexes													
2001	39,480	133	16	18	92	225	768	1,822	3,284	7,356	13,097	12,668	1
2000	37,251	162	14	25	88	244	740	1,651	3,100	6,990	12,458	11,777	2
1999	35,525	166	15	31	72	252	707	1,480	2,864	6,841	11,927	11,170	-
Male													
2001	18,852	76	7	6	45	112	414	1,053	1,714	3,826	6,485	5,114	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)													
All origins													
Male													
2000	17,811	103	7	17	43	129	429	982	1,651	3,564	6,213	4,671	2
1999	17,016	90	9	18	33	125	402	846	1,530	3,504	6,031	4,428	-
Female													
2001	20,628	57	9	12	47	113	354	769	1,570	3,530	6,612	7,554	1
2000	19,440	59	7	8	45	115	311	669	1,449	3,426	6,245	7,106	-
1999	18,509	76	6	13	39	127	305	634	1,334	3,337	5,896	6,742	-
Hispanic													
Both sexes													
2001	1,857	18	2	4	16	25	74	132	237	470	524	355	-
2000	1,646	25	4	3	14	26	53	109	232	404	429	346	1
1999	1,584	20	3	9	6	21	62	107	230	369	426	331	-
Male													
2001	892	13	-	2	9	10	46	64	132	217	251	148	-
2000	805	16	3	2	7	11	36	69	115	194	206	145	1
1999	814	15	2	7	2	10	31	64	137	178	212	156	-
Female													
2001	965	5	2	2	7	15	28	68	105	253	273	207	-
2000	841	9	1	1	7	15	17	40	117	210	223	201	-
1999	770	5	1	2	4	11	31	43	93	191	214	175	-
Non-Hispanic [1]													
Both sexes													
2001	37,511	113	14	14	76	198	691	1,677	3,036	6,866	12,538	12,288	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)													
Non-Hispanic [1]													
Both sexes													
2000	35,509	134	10	22	73	217	686	1,536	2,854	6,567	12,000	11,410	-
1999	33,830	145	12	22	66	231	643	1,368	2,621	6,451	11,474	10,797	-
Male													
2001	17,908	61	7	4	36	100	366	982	1,577	3,598	6,220	4,957	-
2000	16,958	84	4	15	35	118	392	908	1,529	3,357	5,995	4,521	-
1999	16,143	74	7	11	31	115	370	777	1,384	3,312	5,808	4,254	-
Female													
2001	19,603	52	7	10	40	98	325	695	1,459	3,268	6,318	7,331	-
2000	18,551	50	6	7	38	99	294	628	1,325	3,210	6,005	6,889	-
1999	17,687	71	5	11	35	116	273	591	1,237	3,139	5,666	6,543	-
Non-Hispanic white													
Both sexes													
2001	29,449	58	11	6	45	100	359	904	1,861	4,984	10,398	10,723	-
2000	27,924	61	4	17	48	110	359	829	1,776	4,741	10,026	9,953	-
1999	26,418	81	10	13	36	116	319	717	1,636	4,718	9,504	9,268	-
Male													
2001	14,331	31	4	3	21	45	211	543	1,012	2,729	5,310	4,422	-
2000	13,567	41	2	12	25	62	206	491	990	2,574	5,141	4,023	-
1999	12,793	46	5	8	17	60	188	406	903	2,532	4,907	3,721	-
Female													
2001	15,118	27	7	3	24	55	148	361	849	2,255	5,088	6,301	-
2000	14,357	20	2	5	23	48	153	338	786	2,167	4,885	5,930	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)													
Non-Hispanic white													
Female													
1999	13,625	35	5	5	19	56	131	311	733	2,186	4,597	5,547	-
Non-Hispanic black													
Both sexes													
2001	7,214	52	3	7	27	93	306	702	1,048	1,698	1,891	1,387	-
2000	6,853	69	5	4	24	99	305	664	980	1,637	1,776	1,290	-
1999	6,659	58	1	8	30	103	310	615	879	1,554	1,750	1,351	-
Male													
2001	3,158	28	3	-	12	54	139	399	509	782	784	448	-
2000	3,045	40	2	2	10	50	175	394	484	706	759	423	-
1999	2,984	25	1	3	14	47	173	353	426	691	803	448	-
Female													
2001	4,056	24	-	7	15	39	167	303	539	916	1,107	939	-
2000	3,808	29	3	2	14	49	130	270	496	931	1,017	867	-
1999	3,675	33	-	5	16	56	137	262	453	863	947	903	-
Origin not stated [2]													
Both sexes													
2001	112	2	-	-	-	2	3	13	11	20	35	25	1
2000	96	3	-	-	1	1	1	6	14	19	29	21	1
1999	111	1	-	-	-	-	2	5	13	21	27	42	-
Male													
2001	52	2	-	-	-	2	2	7	5	11	14	9	-
2000	48	3	-	-	1	-	1	5	7	13	12	5	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)													
Origin not stated [2]													
Male													
1999	59	1	-	-	-	-	1	5	9	14	11	18	-
Female													
2001	60	-	-	-	-	-	1	6	6	9	21	16	1
2000	48	-	-	-	-	1	-	1	7	6	17	16	-
1999	52	-	-	-	-	-	1	-	4	7	16	24	-
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)													
All origins													
Both sexes													
2001	156	8	4	1	-	3	8	6	13	27	48	38	-
2000	177	8	3	2	1	4	3	16	10	29	52	49	-
1999	182	3	2	5	2	5	5	11	16	37	62	34	-
Male													
2001	73	4	3	-	-	2	6	5	9	10	19	15	-
2000	91	3	2	2	1	3	1	11	8	12	26	22	-
1999	90	3	2	2	2	4	2	6	6	17	32	14	-
Female													
2001	83	4	1	1	-	1	2	1	4	17	29	23	-
2000	86	5	1	-	-	1	2	5	2	17	26	27	-
1999	92	-	-	3	-	1	3	5	10	20	30	20	-
Hispanic													
Both sexes													
2001	13	1	-	-	-	1	1	2	2	3	1	2	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)													
Hispanic													
Both sexes													
2000	9	2	1	-	-	-	-	2	-	2	1	1	-
1999	12	1	-	2	-	-	-	1	2	4	1	1	-
Male													
2001	7	-	-	-	-	-	1	2	2	2	-	-	-
2000	4	-	1	-	-	-	-	1	-	-	1	1	-
1999	4	1	-	1	-	-	-	-	1	-	-	1	-
Female													
2001	6	1	-	-	-	1	-	-	-	1	1	2	-
2000	5	2	-	-	-	-	-	1	-	2	-	-	-
1999	8	-	-	1	-	-	-	1	1	4	1	-	-
Non-Hispanic [1]													
Both sexes													
2001	142	7	4	1	-	2	7	4	11	24	46	36	-
2000	168	6	2	2	1	4	3	14	10	27	51	48	-
1999	170	2	2	3	2	5	5	10	14	33	61	33	-
Male													
2001	65	4	3	-	-	2	5	3	7	8	18	15	-
2000	87	3	1	2	1	3	1	10	8	12	25	21	-
1999	86	2	2	1	2	4	2	6	5	17	32	13	-
Female													
2001	77	3	1	1	-	-	2	1	4	16	28	21	-
2000	81	3	1	-	-	1	2	4	2	15	26	27	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)													
Non-Hispanic [1]													
Female													
1999	84	-	-	2	-	1	3	4	9	16	29	20	-
Non-Hispanic white													
Both sexes													
2001	116	6	3	-	-	-	7	3	9	18	41	29	-
2000	132	4	-	1	1	3	1	4	10	20	45	43	-
1999	132	2	2	2	1	2	3	6	7	28	51	28	-
Male													
2001	52	3	2	-	-	-	5	3	5	4	18	12	-
2000	68	3	-	1	1	2	1	4	8	9	22	17	-
1999	63	2	2	1	1	1	1	4	2	14	26	9	-
Female													
2001	64	3	1	-	-	-	2	-	4	14	23	17	-
2000	64	1	-	-	-	1	-	-	2	11	23	26	-
1999	69	-	-	1	-	1	2	2	5	14	25	19	-
Non-Hispanic black													
Both sexes													
2001	21	-	1	1	-	2	-	1	1	6	5	4	-
2000	30	2	2	-	-	1	2	10	-	6	5	2	-
1999	28	-	-	1	1	2	2	4	5	3	6	4	-
Male													
2001	9	-	1	-	-	2	-	-	1	4	-	1	-
2000	13	-	1	-	-	1	-	6	-	2	2	1	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)													
Non-Hispanic black													
Male													
1999	16	-	-	-	1	2	1	2	2	2	3	3	-
Female													
2001	12	-	-	1	-	-	-	1	-	2	5	3	-
2000	17	2	1	-	-	-	2	4	-	4	3	1	-
1999	12	-	-	1	-	-	1	2	3	1	3	1	-
Origin not stated [2]													
Both sexes													
2001	1	-	-	-	-	-	-	-	-	-	1	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	1	-	-	-	-	-	-	-	-	-	1	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Chronic glomerulonephritis, nephritis and nephropathy, not specified as acute or chronic, and renal sclerosis, unspecified (N02-N03,N05-N07,N26)													
All origins													
Both sexes													
2001	511	-	2	1	8	12	27	41	40	111	147	122	-
2000	556	1	-	4	9	17	34	39	59	103	172	118	-
1999	589	-	1	1	7	15	28	44	60	101	200	132	-
Male													
2001	264	-	1	-	5	1	13	24	25	63	84	48	-
2000	284	-	-	1	2	11	20	27	31	54	83	55	-
1999	304	-	-	1	6	10	18	25	32	45	107	60	-
Female													
2001	247	-	1	1	3	11	14	17	15	48	63	74	-
2000	272	1	-	3	7	6	14	12	28	49	89	63	-
1999	285	-	1	-	1	5	10	19	28	56	93	72	-
Hispanic													
Both sexes													
2001	34	-	1	-	1	1	6	4	4	6	6	5	-
2000	28	-	-	-	-	2	1	4	4	5	7	5	-
1999	33	-	1	-	1	3	2	2	6	8	5	5	-
Male													
2001	17	-	-	-	1	-	5	-	3	2	4	2	-
2000	17	-	-	-	-	1	-	3	3	2	3	5	-
1999	18	-	-	-	1	3	2	1	4	4	1	2	-
Female													
2001	17	-	1	-	-	1	1	4	1	4	2	3	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Chronic glomerulonephritis, nephritis and nephropathy, not specified as acute or chronic, and renal sclerosis, unspecified (N02-N03,N05-N07,N26)													
Hispanic													
Female													
2000	11	-	-	-	-	1	1	1	1	3	4	-	-
1999	15	-	1	-	-	-	-	1	2	4	4	3	-
Non-Hispanic [1]													
Both sexes													
2001	477	-	1	1	7	11	21	37	36	105	141	117	-
2000	528	1	-	4	9	15	33	35	55	98	165	113	-
1999	555	-	-	1	6	12	26	42	53	93	195	127	-
Male													
2001	247	-	1	-	4	1	8	24	22	61	80	46	-
2000	267	-	-	1	2	10	20	24	28	52	80	50	-
1999	285	-	-	1	5	7	16	24	27	41	106	58	-
Female													
2001	230	-	-	1	3	10	13	13	14	44	61	71	-
2000	261	1	-	3	7	5	13	11	27	46	85	63	-
1999	270	-	-	-	1	5	10	18	26	52	89	69	-
Non-Hispanic white													
Both sexes													
2001	384	-	1	-	6	8	9	17	24	87	123	109	-
2000	411	-	-	3	7	9	19	22	39	80	143	89	-
1999	436	-	-	1	3	8	13	30	36	72	169	104	-
Male													
2001	200	-	1	-	3	1	3	10	14	52	71	45	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Chronic glomerulonephritis, nephritis and nephropathy, not specified as acute or chronic, and renal sclerosis, unspecified (N02-N03,N05-N07,N26)													
Non-Hispanic white													
Male													
2000	209	-	-	1	1	6	12	15	20	43	72	39	-
1999	227	-	-	1	2	5	8	18	19	33	93	48	-
Female													
2001	184	-	-	-	3	7	6	7	10	35	52	64	-
2000	202	-	-	2	6	3	7	7	19	37	71	50	-
1999	209	-	-	-	1	3	5	12	17	39	76	56	-
Non-Hispanic black													
Both sexes													
2001	68	-	-	1	-	2	8	15	8	14	13	7	-
2000	90	1	-	1	2	4	10	12	12	14	19	15	-
1999	92	-	-	-	3	2	11	10	11	17	19	19	-
Male													
2001	36	-	-	-	-	-	5	12	6	7	5	1	-
2000	44	-	-	-	1	2	5	8	7	8	6	7	-
1999	46	-	-	-	3	1	6	6	6	6	10	8	-
Female													
2001	32	-	-	1	-	2	3	3	2	7	8	6	-
2000	46	1	-	1	1	2	5	4	5	6	13	8	-
1999	46	-	-	-	-	1	5	4	5	11	9	11	-
Origin not stated [2]													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Chronic glomerulonephritis, nephritis and nephropathy, not specified as acute or chronic, and renal sclerosis, unspecified (N02-N03,N05-N07,N26)													
Origin not stated [2]													
Both sexes													
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	1	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	1	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Renal failure (N17-N19)													
All origins													
Both sexes													
2001	38,784	125	9	16	82	209	732	1,770	3,226	7,211	12,898	12,505	1
2000	36,471	152	11	19	78	221	701	1,590	3,023	6,848	12,223	11,603	2
1999	34,719	163	12	25	63	230	673	1,422	2,784	6,697	11,652	10,998	-
Male													
2001	18,509	72	3	6	40	109	394	1,022	1,680	3,752	6,381	5,050	-
2000	17,416	100	5	14	40	115	407	941	1,607	3,492	6,102	4,591	2
1999	16,611	87	7	15	25	110	382	814	1,492	3,440	5,888	4,351	-
Female													
2001	20,275	53	6	10	42	100	338	748	1,546	3,459	6,517	7,455	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Renal failure (N17-N19)													
All origins													
Female													
2000	19,055	52	6	5	38	106	294	649	1,416	3,356	6,121	7,012	-
1999	18,108	76	5	10	38	120	291	608	1,292	3,257	5,764	6,647	-
Hispanic													
Both sexes													
2001	1,806	17	1	4	15	23	66	125	230	460	517	348	-
2000	1,605	22	3	3	14	24	51	103	227	397	420	340	1
1999	1,535	19	2	7	5	18	60	104	222	356	417	325	-
Male													
2001	867	13	-	2	8	10	39	62	127	213	247	146	-
2000	782	16	2	2	7	10	35	65	111	192	202	139	1
1999	790	14	2	6	1	7	29	63	132	174	209	153	-
Female													
2001	939	4	1	2	7	13	27	63	103	247	270	202	-
2000	823	6	1	1	7	14	16	38	116	205	218	201	-
1999	745	5	-	1	4	11	31	41	90	182	208	172	-
Non-Hispanic [1]													
Both sexes													
2001	36,867	106	8	12	67	184	663	1,632	2,985	6,731	12,347	12,132	-
2000	34,771	127	8	16	63	196	649	1,481	2,783	6,432	11,774	11,242	-
1999	33,074	143	10	18	58	212	611	1,313	2,550	6,320	11,208	10,631	-
Male													
2001	17,591	57	3	4	32	97	353	953	1,548	3,528	6,121	4,895	-
2000	16,586	81	3	12	32	105	371	871	1,489	3,287	5,888	4,447	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Renal failure (N17-N19)													
Non-Hispanic [1]													
Male													
1999	15,763	72	5	9	24	103	352	746	1,352	3,252	5,668	4,180	-
Female													
2001	19,276	49	5	8	35	87	310	679	1,437	3,203	6,226	7,237	-
2000	18,185	46	5	4	31	91	278	610	1,294	3,145	5,886	6,795	-
1999	17,311	71	5	9	34	109	259	567	1,198	3,068	5,540	6,451	-
Non-Hispanic white													
Both sexes													
2001	28,932	52	6	6	38	91	343	881	1,825	4,873	10,233	10,584	-
2000	27,350	57	4	13	40	97	339	800	1,722	4,632	9,830	9,816	-
1999	25,831	79	8	10	32	104	302	679	1,592	4,614	9,279	9,132	-
Male													
2001	14,076	28	1	3	18	44	203	528	993	2,672	5,221	4,365	-
2000	13,275	38	2	10	23	54	193	471	959	2,516	5,045	3,964	-
1999	12,496	44	3	6	14	53	179	383	882	2,484	4,787	3,661	-
Female													
2001	14,856	24	5	3	20	47	140	353	832	2,201	5,012	6,219	-
2000	14,075	19	2	3	17	43	146	329	763	2,116	4,785	5,852	-
1999	13,335	35	5	4	18	51	123	296	710	2,130	4,492	5,471	-
Non-Hispanic black													
Both sexes													
2001	7,119	52	2	5	26	89	298	686	1,038	1,678	1,871	1,374	-
2000	6,724	66	3	3	22	93	292	639	968	1,616	1,751	1,271	-
1999	6,527	58	1	7	26	99	297	600	860	1,533	1,720	1,326	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Renal failure (N17-N19)													
Non-Hispanic black													
Male													
2001	3,111	28	2	-	12	52	134	387	502	771	778	445	-
2000	2,986	40	1	2	9	47	170	378	477	696	751	415	-
1999	2,920	25	1	3	10	44	166	345	418	682	789	437	-
Female													
2001	4,008	24	-	5	14	37	164	299	536	907	1,093	929	-
2000	3,738	26	2	1	13	46	122	261	491	920	1,000	856	-
1999	3,607	33	-	4	16	55	131	255	442	851	931	889	-
Origin not stated [2]													
Both sexes													
2001	111	2	-	-	-	2	3	13	11	20	34	25	1
2000	95	3	-	-	1	1	1	6	13	19	29	21	1
1999	110	1	-	-	-	-	2	5	12	21	27	42	-
Male													
2001	51	2	-	-	-	2	2	7	5	11	13	9	-
2000	48	3	-	-	1	-	1	5	7	13	12	5	1
1999	58	1	-	-	-	-	1	5	8	14	11	18	-
Female													
2001	60	-	-	-	-	-	1	6	6	9	21	16	1
2000	47	-	-	-	-	1	-	1	6	6	17	16	-
1999	52	-	-	-	-	-	1	-	4	7	16	24	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other disorders of kidney (N25,N27)													
All origins													
Both sexes													
2001	29	-	1	-	2	1	1	5	5	7	4	3	-
2000	47	1	-	-	-	2	2	6	8	10	11	7	-
1999	35	-	-	-	-	2	1	3	4	6	13	6	-
Male													
2001	6	-	-	-	-	-	1	2	-	1	1	1	-
2000	20	-	-	-	-	-	1	3	5	6	2	3	-
1999	11	-	-	-	-	1	-	1	-	2	4	3	-
Female													
2001	23	-	1	-	2	1	-	3	5	6	3	2	-
2000	27	1	-	-	-	2	1	3	3	4	9	4	-
1999	24	-	-	-	-	1	1	2	4	4	9	3	-
Hispanic													
Both sexes													
2001	4	-	-	-	-	-	1	1	1	1	-	-	-
2000	4	1	-	-	-	-	1	-	1	-	1	-	-
1999	4	-	-	-	-	-	-	-	-	1	3	-	-
Male													
2001	1	-	-	-	-	-	1	-	-	-	-	-	-
2000	2	-	-	-	-	-	1	-	1	-	-	-	-
1999	2	-	-	-	-	-	-	-	-	-	2	-	-
Female													
2001	3	-	-	-	-	-	-	1	1	1	-	-	-
2000	2	1	-	-	-	-	-	-	-	-	1	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other disorders of kidney (N25,N27)													
Hispanic													
Female													
1999	2	-	-	-	-	-	-	-	-	1	1	-	-
Non-Hispanic [1]													
Both sexes													
2001	25	-	1	-	2	1	-	4	4	6	4	3	-
2000	42	-	-	-	-	2	1	6	6	10	10	7	-
1999	31	-	-	-	-	2	1	3	4	5	10	6	-
Male													
2001	5	-	-	-	-	-	-	2	-	1	1	1	-
2000	18	-	-	-	-	-	-	3	4	6	2	3	-
1999	9	-	-	-	-	1	-	1	-	2	2	3	-
Female													
2001	20	-	1	-	2	1	-	2	4	5	3	2	-
2000	24	-	-	-	-	2	1	3	2	4	8	4	-
1999	22	-	-	-	-	1	1	2	4	3	8	3	-
Non-Hispanic white													
Both sexes													
2001	17	-	1	-	1	1	-	3	3	6	1	1	-
2000	31	-	-	-	-	1	-	3	5	9	8	5	-
1999	19	-	-	-	-	2	1	2	1	4	5	4	-
Male													
2001	3	-	-	-	-	-	-	2	-	1	-	-	-
2000	15	-	-	-	-	-	-	1	3	6	2	3	-
1999	7	-	-	-	-	1	-	1	-	1	1	3	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other disorders of kidney (N25,N27)													
Non-Hispanic white													
Female													
2001	14	-	1	-	1	1	-	1	3	5	1	1	-
2000	16	-	-	-	-	1	-	2	2	3	6	2	-
1999	12	-	-	-	-	1	1	1	1	3	4	1	-
Non-Hispanic black													
Both sexes													
2001	6	-	-	-	1	-	-	-	1	-	2	2	-
2000	9	-	-	-	-	1	1	3	-	1	1	2	-
1999	12	-	-	-	-	-	-	1	3	1	5	2	-
Male													
2001	2	-	-	-	-	-	-	-	-	-	1	1	-
2000	2	-	-	-	-	-	-	2	-	-	-	-	-
1999	2	-	-	-	-	-	-	-	-	1	1	-	-
Female													
2001	4	-	-	-	1	-	-	-	1	-	1	1	-
2000	7	-	-	-	-	1	1	1	-	1	1	2	-
1999	10	-	-	-	-	-	-	1	3	-	4	2	-
Origin not stated [2]													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	1	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other disorders of kidney (N25,N27)													
Origin not stated [2]													
Male													
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	1	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Infections of kidney (N10-N12,N13.6,N15.1)													
All origins													
Both sexes													
2001	748	2	4	-	5	11	33	46	46	87	249	265	-
2000	813	5	3	-	2	9	21	42	59	104	248	320	-
1999	846	3	1	1	4	9	28	55	59	133	242	311	-
Male													
2001	213	1	2	-	2	3	10	11	12	40	76	56	-
2000	214	3	-	-	1	3	8	15	15	26	74	69	-
1999	263	1	-	-	1	-	7	19	27	44	83	81	-
Female													
2001	535	1	2	-	3	8	23	35	34	47	173	209	-
2000	599	2	3	-	1	6	13	27	44	78	174	251	-
1999	583	2	1	1	3	9	21	36	32	89	159	230	-
Hispanic													
Both sexes													
2001	33	-	-	-	-	1	3	5	1	8	9	6	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Infections of kidney (N10-N12,N13.6,N15.1)													
Hispanic													
Both sexes													
2000	39	1	-	-	-	-	2	4	4	7	10	11	-
1999	55	-	-	-	-	1	7	6	9	8	15	9	-
Male													
2001	10	-	-	-	-	-	-	3	-	3	1	3	-
2000	11	1	-	-	-	-	-	3	1	3	2	1	-
1999	21	-	-	-	-	-	2	2	5	4	6	2	-
Female													
2001	23	-	-	-	-	1	3	2	1	5	8	3	-
2000	28	-	-	-	-	-	2	1	3	4	8	10	-
1999	34	-	-	-	-	1	5	4	4	4	9	7	-
Non-Hispanic [1]													
Both sexes													
2001	714	2	4	-	5	10	30	41	45	79	239	259	-
2000	771	4	3	-	2	9	19	38	55	96	236	309	-
1999	786	3	1	1	4	8	20	49	49	125	225	301	-
Male													
2001	203	1	2	-	2	3	10	8	12	37	75	53	-
2000	201	2	-	-	1	3	8	12	14	22	71	68	-
1999	240	1	-	-	1	-	4	17	22	40	76	79	-
Female													
2001	511	1	2	-	3	7	20	33	33	42	164	206	-
2000	570	2	3	-	1	6	11	26	41	74	165	241	-
1999	546	2	1	1	3	8	16	32	27	85	149	222	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Infections of kidney (N10-N12,N13.6,N15.1)													
Non-Hispanic white													
Both sexes													
2001	614	1	2	-	2	7	23	29	36	65	210	239	-
2000	647	4	3	-	1	7	10	26	44	74	198	280	-
1999	689	1	1	1	2	7	12	38	38	102	206	281	-
Male													
2001	166	-	1	-	1	2	9	6	9	27	64	47	-
2000	169	2	-	-	-	3	4	9	12	13	62	64	-
1999	209	1	-	-	1	-	2	12	17	30	70	76	-
Female													
2001	448	1	1	-	1	5	14	23	27	38	146	192	-
2000	478	2	3	-	1	4	6	17	32	61	136	216	-
1999	480	-	1	1	1	7	10	26	21	72	136	205	-
Non-Hispanic black													
Both sexes													
2001	80	1	-	-	1	3	7	10	7	13	22	16	-
2000	101	-	-	-	1	2	7	11	8	19	28	25	-
1999	85	1	-	-	2	1	6	10	11	20	16	18	-
Male													
2001	28	1	-	-	1	1	1	2	3	9	6	4	-
2000	27	-	-	-	1	-	2	3	2	8	7	4	-
1999	29	-	-	-	-	-	2	5	5	10	4	3	-
Female													
2001	52	-	-	-	-	2	6	8	4	4	16	12	-
2000	74	-	-	-	-	2	5	8	6	11	21	21	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Infections of kidney (N10-N12,N13.6,N15.1)													
Non-Hispanic black													
Female													
1999	56	1	-	-	2	1	4	5	6	10	12	15	-
Origin not stated [2]													
Both sexes													
2001	1	-	-	-	-	-	-	-	-	-	1	-	-
2000	3	-	-	-	-	-	-	-	-	1	2	-	-
1999	5	-	-	-	-	-	1	-	1	-	2	1	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	1	1	-	-
1999	2	-	-	-	-	-	1	-	-	-	1	-	-
Female													
2001	1	-	-	-	-	-	-	-	-	-	1	-	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-
1999	3	-	-	-	-	-	-	-	1	-	1	1	-
Hyperplasia of prostate (N40)													
All origins													
Both sexes													
2001	439	-	-	-	-	-	-	2	9	45	141	242	-
2000	433	-	-	-	-	-	1	1	11	38	159	222	1
1999	430	-	-	-	-	-	-	1	13	47	160	209	-
Male													
2001	439	-	-	-	-	-	-	2	9	45	141	242	-
2000	433	-	-	-	-	-	1	1	11	38	159	222	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Hyperplasia of prostate (N40)													
All origins													
Male													
1999	430	-	-	-	-	-	-	1	13	47	160	209	-
Female													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Hispanic													
Both sexes													
2001	11	-	-	-	-	-	-	-	1	4	4	2	-
2000	19	-	-	-	-	-	-	-	1	5	3	10	-
1999	11	-	-	-	-	-	-	-	1	3	4	3	-
Male													
2001	11	-	-	-	-	-	-	-	1	4	4	2	-
2000	19	-	-	-	-	-	-	-	1	5	3	10	-
1999	11	-	-	-	-	-	-	-	1	3	4	3	-
Female													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic [1]													
Both sexes													
2001	423	-	-	-	-	-	-	2	8	39	134	240	-
2000	414	-	-	-	-	-	1	1	10	33	156	212	1
1999	417	-	-	-	-	-	-	-	12	43	156	206	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Hyperplasia of prostate (N40)													
Non-Hispanic [1]													
Male													
2001	423	-	-	-	-	-	-	2	8	39	134	240	-
2000	414	-	-	-	-	-	1	1	10	33	156	212	1
1999	417	-	-	-	-	-	-	-	12	43	156	206	-
Female													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic white													
Both sexes													
2001	388	-	-	-	-	-	-	1	4	35	122	226	-
2000	370	-	-	-	-	-	-	1	6	30	142	191	-
1999	377	-	-	-	-	-	-	-	8	36	143	190	-
Male													
2001	388	-	-	-	-	-	-	1	4	35	122	226	-
2000	370	-	-	-	-	-	-	1	6	30	142	191	-
1999	377	-	-	-	-	-	-	-	8	36	143	190	-
Female													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic black													
Both sexes													
2001	31	-	-	-	-	-	-	1	4	3	12	11	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Hyperplasia of prostate (N40)													
Non-Hispanic black													
Both sexes													
2000	37	-	-	-	-	-	1	-	4	2	14	15	1
1999	38	-	-	-	-	-	-	-	4	6	13	15	-
Male													
2001	31	-	-	-	-	-	-	1	4	3	12	11	-
2000	37	-	-	-	-	-	1	-	4	2	14	15	1
1999	38	-	-	-	-	-	-	-	4	6	13	15	-
Female													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Origin not stated [2]													
Both sexes													
2001	5	-	-	-	-	-	-	-	-	2	3	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	-	-	-	-	-	-	1	-	1	-	-	-
Male													
2001	5	-	-	-	-	-	-	-	-	2	3	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	-	-	-	-	-	-	1	-	1	-	-	-
Female													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Inflammatory diseases of female pelvic organs (N70-N76)													
All origins													
Both sexes													
2001	127	1	-	-	2	3	10	12	12	18	31	38	-
2000	120	2	-	1	2	6	15	9	8	21	29	27	-
1999	108	2	1	-	3	6	12	6	13	15	32	18	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	127	1	-	-	2	3	10	12	12	18	31	38	-
2000	120	2	-	1	2	6	15	9	8	21	29	27	-
1999	108	2	1	-	3	6	12	6	13	15	32	18	-
Hispanic													
Both sexes													
2001	5	-	-	-	-	-	-	2	-	2	-	1	-
2000	7	-	-	-	-	1	1	-	2	2	-	1	-
1999	7	-	-	-	-	-	1	2	1	-	1	2	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	5	-	-	-	-	-	-	2	-	2	-	1	-
2000	7	-	-	-	-	1	1	-	2	2	-	1	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Inflammatory diseases of female pelvic organs (N70-N76)													
Hispanic													
Female													
1999	7	-	-	-	-	-	1	2	1	-	1	2	-
Non-Hispanic [1]													
Both sexes													
2001	122	1	-	-	2	3	10	10	12	16	31	37	-
2000	111	2	-	1	2	5	14	7	6	19	29	26	-
1999	101	2	1	-	3	6	11	4	12	15	31	16	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	122	1	-	-	2	3	10	10	12	16	31	37	-
2000	111	2	-	1	2	5	14	7	6	19	29	26	-
1999	101	2	1	-	3	6	11	4	12	15	31	16	-
Non-Hispanic white													
Both sexes													
2001	98	1	-	-	2	3	6	6	8	12	27	33	-
2000	87	1	-	-	2	3	6	6	5	15	26	23	-
1999	73	1	1	-	2	3	3	2	11	10	25	15	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Inflammatory diseases of female pelvic organs (N70-N76)													
Non-Hispanic white													
Female													
2001	98	1	-	-	2	3	6	6	8	12	27	33	-
2000	87	1	-	-	2	3	6	6	5	15	26	23	-
1999	73	1	1	-	2	3	3	2	11	10	25	15	-
Non-Hispanic black													
Both sexes													
2001	23	-	-	-	-	-	4	4	4	4	4	3	-
2000	21	1	-	1	-	1	7	1	1	4	3	2	-
1999	26	1	-	-	1	2	7	2	1	5	6	1	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	23	-	-	-	-	-	4	4	4	4	4	3	-
2000	21	1	-	1	-	1	7	1	1	4	3	2	-
1999	26	1	-	-	1	2	7	2	1	5	6	1	-
Origin not stated [2]													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	2	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Inflammatory diseases of female pelvic organs (N70-N76)													
Origin not stated [2]													
Male													
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	2	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Pregnancy, childbirth and the puerperium (O00-O99)													
All origins													
Both sexes													
2001	416	. . .	. . .	3	110	190	106	6	-	-	-	1	-
2000	404	. . .	. . .	-	95	182	121	6	-	-	-	-	-
1999	406	. . .	. . .	-	97	181	121	7	-	-	-	-	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	416	. . .	. . .	3	110	190	106	6	-	-	-	1	-
2000	404	. . .	. . .	-	95	182	121	6	-	-	-	-	-
1999	406	. . .	. . .	-	97	181	121	7	-	-	-	-	-
Hispanic													
Both sexes													
2001	83	. . .	. . .	1	26	38	17	1	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Pregnancy, childbirth and the puerperium (O00-O99)													
Hispanic													
Both sexes													
2000	84	. . .	. . .	-	24	36	23	1	-	-	-	-	-
1999	70	. . .	. . .	-	19	26	25	-	-	-	-	-	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	83	. . .	. . .	1	26	38	17	1	-	-	-	-	-
2000	84	. . .	. . .	-	24	36	23	1	-	-	-	-	-
1999	70	. . .	. . .	-	19	26	25	-	-	-	-	-	-
Non-Hispanic [1]													
Both sexes													
2001	332	. . .	. . .	2	84	151	89	5	-	-	-	1	-
2000	319	. . .	. . .	-	71	146	97	5	-	-	-	-	-
1999	333	. . .	. . .	-	76	154	96	7	-	-	-	-	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	332	. . .	. . .	2	84	151	89	5	-	-	-	1	-
2000	319	. . .	. . .	-	71	146	97	5	-	-	-	-	-
1999	333	. . .	. . .	-	76	154	96	7	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Pregnancy, childbirth and the puerperium (O00-O99)													
Non-Hispanic white													
Both sexes													
2001	156	. . .	. . .	1	38	77	37	2	-	-	-	1	-
2000	161	. . .	. . .	-	33	72	54	2	-	-	-	-	-
1999	152	. . .	. . .	-	26	74	48	4	-	-	-	-	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	156	. . .	. . .	1	38	77	37	2	-	-	-	1	-
2000	161	. . .	. . .	-	33	72	54	2	-	-	-	-	-
1999	152	. . .	. . .	-	26	74	48	4	-	-	-	-	-
Non-Hispanic black													
Both sexes													
2001	154	. . .	. . .	1	45	65	40	3	-	-	-	-	-
2000	139	. . .	. . .	-	34	65	37	3	-	-	-	-	-
1999	158	. . .	. . .	-	46	70	40	2	-	-	-	-	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	154	. . .	. . .	1	45	65	40	3	-	-	-	-	-
2000	139	. . .	. . .	-	34	65	37	3	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Pregnancy, childbirth and the puerperium (O00-O99)													
Non-Hispanic black													
Female													
1999	158	. . .	. . .	-	46	70	40	2	-	-	-	-	-
Origin not stated [2]													
Both sexes													
2001	1	. . .	. . .	-	-	1	-	-	-	-	-	-	-
2000	1	. . .	. . .	-	-	-	1	-	-	-	-	-	-
1999	3	. . .	. . .	-	2	1	-	-	-	-	-	-	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	1	. . .	. . .	-	-	1	-	-	-	-	-	-	-
2000	1	. . .	. . .	-	-	-	1	-	-	-	-	-	-
1999	3	. . .	. . .	-	2	1	-	-	-	-	-	-	-
Pregnancy with abortive outcome (O00-O07)													
All origins													
Both sexes													
2001	38	. . .	. . .	2	13	14	9	-	-	-	-	-	-
2000	37	. . .	. . .	-	6	17	14	-	-	-	-	-	-
1999	31	. . .	. . .	-	8	15	8	-	-	-	-	-	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Pregnancy with abortive outcome (O00-O07)													
All origins													
Male													
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	38	. . .	. . .	2	13	14	9	-	-	-	-	-	-
2000	37	. . .	. . .	-	6	17	14	-	-	-	-	-	-
1999	31	. . .	. . .	-	8	15	8	-	-	-	-	-	-
Hispanic													
Both sexes													
2001	1	. . .	. . .	1	-	-	-	-	-	-	-	-	-
2000	6	. . .	. . .	-	2	3	1	-	-	-	-	-	-
1999	4	. . .	. . .	-	2	-	2	-	-	-	-	-	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	1	. . .	. . .	1	-	-	-	-	-	-	-	-	-
2000	6	. . .	. . .	-	2	3	1	-	-	-	-	-	-
1999	4	. . .	. . .	-	2	-	2	-	-	-	-	-	-
Non-Hispanic [1]													
Both sexes													
2001	37	. . .	. . .	1	13	14	9	-	-	-	-	-	-
2000	30	. . .	. . .	-	4	14	12	-	-	-	-	-	-
1999	26	. . .	. . .	-	5	15	6	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Pregnancy with abortive outcome (O00-O07)													
Non-Hispanic [1]													
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	37	. . .	. . .	1	13	14	9	-	-	-	-	-	-
2000	30	. . .	. . .	-	4	14	12	-	-	-	-	-	-
1999	26	. . .	. . .	-	5	15	6	-	-	-	-	-	-
Non-Hispanic white													
Both sexes													
2001	19	. . .	. . .	-	7	8	4	-	-	-	-	-	-
2000	12	. . .	. . .	-	-	5	7	-	-	-	-	-	-
1999	11	. . .	. . .	-	2	6	3	-	-	-	-	-	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	19	. . .	. . .	-	7	8	4	-	-	-	-	-	-
2000	12	. . .	. . .	-	-	5	7	-	-	-	-	-	-
1999	11	. . .	. . .	-	2	6	3	-	-	-	-	-	-
Non-Hispanic black													
Both sexes													
2001	16	. . .	. . .	1	6	5	4	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Pregnancy with abortive outcome (O00-O07)													
Non-Hispanic black													
Both sexes													
2000	18	. . .	. . .	-	4	9	5	-	-	-	-	-	-
1999	14	. . .	. . .	-	3	8	3	-	-	-	-	-	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	16	. . .	. . .	1	6	5	4	-	-	-	-	-	-
2000	18	. . .	. . .	-	4	9	5	-	-	-	-	-	-
1999	14	. . .	. . .	-	3	8	3	-	-	-	-	-	-
Origin not stated [2]													
Both sexes													
2001	-	. . .	. . .	-	-	-	-	-	-	-	-	-	-
2000	1	. . .	. . .	-	-	-	1	-	-	-	-	-	-
1999	1	. . .	. . .	-	1	-	-	-	-	-	-	-	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	-	. . .	. . .	-	-	-	-	-	-	-	-	-	-
2000	1	. . .	. . .	-	-	-	1	-	-	-	-	-	-
1999	1	. . .	. . .	-	1	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other complications of pregnancy, childbirth and the puerperium (O10-O99)													
All origins													
Both sexes													
2001	378	. . .	. . .	1	97	176	97	6	-	-	-	1	-
2000	367	. . .	. . .	-	89	165	107	6	-	-	-	-	-
1999	375	. . .	. . .	-	89	166	113	7	-	-	-	-	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	378	. . .	. . .	1	97	176	97	6	-	-	-	1	-
2000	367	. . .	. . .	-	89	165	107	6	-	-	-	-	-
1999	375	. . .	. . .	-	89	166	113	7	-	-	-	-	-
Hispanic													
Both sexes													
2001	82	. . .	. . .	-	26	38	17	1	-	-	-	-	-
2000	78	. . .	. . .	-	22	33	22	1	-	-	-	-	-
1999	66	. . .	. . .	-	17	26	23	-	-	-	-	-	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	82	. . .	. . .	-	26	38	17	1	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other complications of pregnancy, childbirth and the puerperium (O10-O99)													
Hispanic													
Female													
2000	78	. . .	. . .	-	22	33	22	1	-	-	-	-	-
1999	66	. . .	. . .	-	17	26	23	-	-	-	-	-	-
Non-Hispanic [1]													
Both sexes													
2001	295	. . .	. . .	1	71	137	80	5	-	-	-	1	-
2000	289	. . .	. . .	-	67	132	85	5	-	-	-	-	-
1999	307	. . .	. . .	-	71	139	90	7	-	-	-	-	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	295	. . .	. . .	1	71	137	80	5	-	-	-	1	-
2000	289	. . .	. . .	-	67	132	85	5	-	-	-	-	-
1999	307	. . .	. . .	-	71	139	90	7	-	-	-	-	-
Non-Hispanic white													
Both sexes													
2001	137	. . .	. . .	1	31	69	33	2	-	-	-	1	-
2000	149	. . .	. . .	-	33	67	47	2	-	-	-	-	-
1999	141	. . .	. . .	-	24	68	45	4	-	-	-	-	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other complications of pregnancy, childbirth and the puerperium (O10-O99)													
Non-Hispanic white													
Male													
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	137	. . .	. . .	1	31	69	33	2	-	-	-	1	-
2000	149	. . .	. . .	-	33	67	47	2	-	-	-	-	-
1999	141	. . .	. . .	-	24	68	45	4	-	-	-	-	-
Non-Hispanic black													
Both sexes													
2001	138	. . .	. . .	-	39	60	36	3	-	-	-	-	-
2000	121	. . .	. . .	-	30	56	32	3	-	-	-	-	-
1999	144	. . .	. . .	-	43	62	37	2	-	-	-	-	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	138	. . .	. . .	-	39	60	36	3	-	-	-	-	-
2000	121	. . .	. . .	-	30	56	32	3	-	-	-	-	-
1999	144	. . .	. . .	-	43	62	37	2	-	-	-	-	-
Origin not stated [2]													
Both sexes													
2001	1	. . .	. . .	-	-	1	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other complications of pregnancy, childbirth and the puerperium (O10-O99)													
Origin not stated [2]													
Both sexes													
2000	-	. . .	. . .	-	-	-	-	-	-	-	-	-	-
1999	2	. . .	. . .	-	1	1	-	-	-	-	-	-	-
Male													
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female													
2001	1	. . .	. . .	-	-	1	-	-	-	-	-	-	-
2000	-	. . .	. . .	-	-	-	-	-	-	-	-	-	-
1999	2	. . .	. . .	-	1	1	-	-	-	-	-	-	-
Certain conditions originating in the perinatal period (P00-P96)													
All origins													
Both sexes													
2001	13,887	13,734	72	36	17	6	9	5	2	3	1	1	1
2000	14,069	13,918	79	41	15	7	4	2	2	-	-	-	1
1999	14,259	14,084	92	43	16	7	3	2	3	2	1	1	5
Male													
2001	7,868	7,775	44	24	7	2	6	5	1	2	1	-	1
2000	7,968	7,872	52	23	11	6	2	2	-	-	-	-	-
1999	8,044	7,941	56	23	10	2	2	2	1	1	1	1	4
Female													
2001	6,019	5,959	28	12	10	4	3	-	1	1	-	1	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Certain conditions originating in the perinatal period (P00-P96)													
All origins													
Female													
2000	6,101	6,046	27	18	4	1	2	-	2	-	-	-	1
1999	6,215	6,143	36	20	6	5	1	-	2	1	-	-	1
Hispanic													
Both sexes													
2001	2,227	2,203	14	3	2	4	-	1	-	-	-	-	-
2000	2,145	2,120	12	10	2	-	1	-	-	-	-	-	-
1999	2,153	2,127	18	2	4	1	-	-	-	-	-	-	1
Male													
2001	1,268	1,258	6	2	-	1	-	1	-	-	-	-	-
2000	1,194	1,180	7	6	1	-	-	-	-	-	-	-	-
1999	1,192	1,181	5	2	3	-	-	-	-	-	-	-	1
Female													
2001	959	945	8	1	2	3	-	-	-	-	-	-	-
2000	951	940	5	4	1	-	1	-	-	-	-	-	-
1999	961	946	13	-	1	1	-	-	-	-	-	-	-
Non-Hispanic [1]													
Both sexes													
2001	11,500	11,371	58	33	15	2	9	4	2	3	1	1	1
2000	11,672	11,546	67	31	13	7	3	2	2	-	-	-	1
1999	11,815	11,668	73	41	12	6	3	2	3	2	1	1	3
Male													
2001	6,511	6,428	38	22	7	1	6	4	1	2	1	-	1
2000	6,637	6,555	45	17	10	6	2	2	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Certain conditions originating in the perinatal period (P00-P96)													
Non-Hispanic [1]													
Male													
1999	6,680	6,589	50	21	7	2	2	2	1	1	1	1	3
Female													
2001	4,989	4,943	20	11	8	1	3	-	1	1	-	1	-
2000	5,035	4,991	22	14	3	1	1	-	2	-	-	-	1
1999	5,135	5,079	23	20	5	4	1	-	2	1	-	-	-
Non-Hispanic white													
Both sexes													
2001	6,323	6,239	34	18	11	2	8	4	1	3	1	1	1
2000	6,318	6,242	34	20	11	3	3	2	2	-	-	-	1
1999	6,384	6,286	37	31	11	6	3	2	3	2	1	1	1
Male													
2001	3,574	3,518	22	14	6	1	5	4	-	2	1	-	1
2000	3,605	3,553	23	14	9	2	2	2	-	-	-	-	-
1999	3,627	3,570	24	15	7	2	2	2	1	1	1	1	1
Female													
2001	2,749	2,721	12	4	5	1	3	-	1	1	-	1	-
2000	2,713	2,689	11	6	2	1	1	-	2	-	-	-	1
1999	2,757	2,716	13	16	4	4	1	-	2	1	-	-	-
Non-Hispanic black													
Both sexes													
2001	4,713	4,675	22	11	4	-	1	-	-	-	-	-	-
2000	4,878	4,836	27	10	1	4	-	-	-	-	-	-	-
1999	5,000	4,963	29	5	1	-	-	-	-	-	-	-	2

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Certain conditions originating in the perinatal period (P00-P96)													
Non-Hispanic black													
Male													
2001	2,674	2,653	14	5	1	-	1	-	-	-	-	-	-
2000	2,760	2,734	18	3	1	4	-	-	-	-	-	-	-
1999	2,818	2,794	19	3	-	-	-	-	-	-	-	-	2
Female													
2001	2,039	2,022	8	6	3	-	-	-	-	-	-	-	-
2000	2,118	2,102	9	7	-	-	-	-	-	-	-	-	-
1999	2,182	2,169	10	2	1	-	-	-	-	-	-	-	-
Origin not stated [2]													
Both sexes													
2001	160	160	-	-	-	-	-	-	-	-	-	-	-
2000	252	252	-	-	-	-	-	-	-	-	-	-	-
1999	291	289	1	-	-	-	-	-	-	-	-	-	1
Male													
2001	89	89	-	-	-	-	-	-	-	-	-	-	-
2000	137	137	-	-	-	-	-	-	-	-	-	-	-
1999	172	171	1	-	-	-	-	-	-	-	-	-	-
Female													
2001	71	71	-	-	-	-	-	-	-	-	-	-	-
2000	115	115	-	-	-	-	-	-	-	-	-	-	-
1999	119	118	-	-	-	-	-	-	-	-	-	-	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)													
All origins													
Both sexes													
2001	10,442	5,513	557	376	505	458	586	636	581	448	465	317	-
2000	10,578	5,743	495	399	441	477	569	612	569	464	488	320	1
1999	10,393	5,473	549	428	434	465	552	647	557	469	491	326	2
Male													
2001	5,448	2,954	282	210	306	249	317	323	276	228	203	100	-
2000	5,517	3,028	259	207	268	279	316	347	264	222	197	130	-
1999	5,346	2,875	310	215	262	242	296	347	283	209	204	102	1
Female													
2001	4,994	2,559	275	166	199	209	269	313	305	220	262	217	-
2000	5,061	2,715	236	192	173	198	253	265	305	242	291	190	1
1999	5,047	2,598	239	213	172	223	256	300	274	260	287	224	1
Hispanic													
Both sexes													
2001	1,632	1,195	103	58	73	45	42	36	24	22	24	10	-
2000	1,612	1,182	111	52	54	46	43	36	31	21	24	12	-
1999	1,498	1,072	101	57	56	54	34	42	27	30	16	9	-
Male													
2001	892	639	54	36	39	26	27	25	14	14	13	5	-
2000	857	605	60	28	39	26	27	22	17	11	15	7	-
1999	756	534	63	29	30	25	15	24	9	17	7	3	-
Female													
2001	740	556	49	22	34	19	15	11	10	8	11	5	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)													
Hispanic													
Female													
2000	755	577	51	24	15	20	16	14	14	10	9	5	-
1999	742	538	38	28	26	29	19	18	18	13	9	6	-
Non-Hispanic [1]													
Both sexes													
2001	8,748	4,271	452	317	432	412	542	596	557	422	441	306	-
2000	8,900	4,514	384	345	386	427	525	574	532	443	462	307	1
1999	8,825	4,349	447	370	378	410	516	601	529	435	473	315	2
Male													
2001	4,527	2,297	227	173	267	222	289	294	262	211	190	95	-
2000	4,623	2,400	199	177	229	250	288	323	243	211	181	122	-
1999	4,550	2,311	246	185	232	217	280	320	274	191	196	97	1
Female													
2001	4,221	1,974	225	144	165	190	253	302	295	211	251	211	-
2000	4,277	2,114	185	168	157	177	237	251	289	232	281	185	1
1999	4,275	2,038	201	185	146	193	236	281	255	244	277	218	1
Non-Hispanic white													
Both sexes													
2001	6,709	3,053	287	226	302	330	439	503	503	378	405	283	-
2000	6,792	3,199	240	240	279	344	419	501	460	390	431	288	1
1999	6,782	3,127	284	244	277	334	426	514	473	383	427	291	2
Male													
2001	3,487	1,664	144	135	183	186	236	251	235	196	174	83	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)													
Non-Hispanic white													
Male													
2000	3,519	1,720	124	119	163	203	233	276	217	177	172	115	-
1999	3,498	1,693	154	119	175	178	230	271	248	169	171	89	1
Female													
2001	3,222	1,389	143	91	119	144	203	252	268	182	231	200	-
2000	3,273	1,479	116	121	116	141	186	225	243	213	259	173	1
1999	3,284	1,434	130	125	102	156	196	243	225	214	256	202	1
Non-Hispanic black													
Both sexes													
2001	1,674	999	127	73	113	67	89	76	48	35	28	19	-
2000	1,739	1,062	123	89	91	74	91	67	58	43	25	16	-
1999	1,680	994	137	105	82	60	76	73	46	46	38	23	-
Male													
2001	861	525	65	33	72	30	46	34	23	12	12	9	-
2000	911	545	67	50	58	43	49	41	17	26	9	6	-
1999	880	518	79	53	50	32	41	42	22	16	19	8	-
Female													
2001	813	474	62	40	41	37	43	42	25	23	16	10	-
2000	828	517	56	39	33	31	42	26	41	17	16	10	-
1999	800	476	58	52	32	28	35	31	24	30	19	15	-
Origin not stated [2]													
Both sexes													
2001	62	47	2	1	-	1	2	4	-	4	-	1	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)													
Origin not stated [2]													
Both sexes													
2000	66	47	-	2	1	4	1	2	6	-	2	1	-
1999	70	52	1	1	-	1	2	4	1	4	2	2	-
Male													
2001	29	18	1	1	-	1	1	4	-	3	-	-	-
2000	37	23	-	2	-	3	1	2	4	-	1	1	-
1999	40	30	1	1	-	-	1	3	-	1	1	2	-
Female													
2001	33	29	1	-	-	-	1	-	-	1	-	1	-
2000	29	24	-	-	1	1	-	-	2	-	1	-	-
1999	30	22	-	-	-	1	1	1	1	3	1	-	-
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)													
All origins													
Both sexes													
2001	32,326	3,491	272	181	949	1,530	3,126	3,504	2,292	2,774	5,003	9,089	115
2000	31,876	3,639	248	177	868	1,554	2,987	3,142	2,253	3,040	5,029	8,824	115
1999	26,672	3,538	236	137	609	1,095	2,062	2,151	1,860	2,698	4,413	7,798	75
Male													
2001	15,779	2,041	162	99	654	1,040	2,007	2,286	1,457	1,497	2,184	2,274	78
2000	15,863	2,156	137	99	581	1,060	1,994	2,123	1,429	1,708	2,244	2,242	90
1999	12,877	2,072	136	77	422	721	1,339	1,446	1,181	1,513	1,930	1,984	56
Female													
2001	16,547	1,450	110	82	295	490	1,119	1,218	835	1,277	2,819	6,815	37

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)													
All origins													
Female													
2000	16,013	1,483	111	78	287	494	993	1,019	824	1,332	2,785	6,582	25
1999	13,795	1,466	100	60	187	374	723	705	679	1,185	2,483	5,814	19
Hispanic													
Both sexes													
2001	2,190	496	47	36	184	286	327	283	127	112	107	173	12
2000	2,071	506	51	31	140	260	342	251	114	107	104	154	11
1999	1,281	415	30	16	83	138	148	93	63	73	93	124	5
Male													
2001	1,410	280	28	15	139	221	239	212	92	71	56	46	11
2000	1,369	310	31	16	102	201	262	192	72	68	52	53	10
1999	768	237	14	10	64	99	110	68	42	38	46	36	4
Female													
2001	780	216	19	21	45	65	88	71	35	41	51	127	1
2000	702	196	20	15	38	59	80	59	42	39	52	101	1
1999	513	178	16	6	19	39	38	25	21	35	47	88	1
Non-Hispanic [1]													
Both sexes													
2001	29,931	2,978	225	144	757	1,227	2,782	3,193	2,154	2,653	4,886	8,896	36
2000	29,514	3,108	196	144	720	1,279	2,604	2,849	2,109	2,908	4,908	8,654	35
1999	25,181	3,097	204	121	521	951	1,885	2,023	1,780	2,601	4,310	7,658	30
Male													
2001	14,246	1,750	134	84	510	807	1,754	2,058	1,355	1,422	2,125	2,221	26

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)													
Non-Hispanic [1]													
Male													
2000	14,288	1,834	106	82	474	848	1,700	1,901	1,331	1,621	2,178	2,183	30
1999	11,962	1,817	121	67	355	619	1,209	1,347	1,125	1,463	1,876	1,944	19
Female													
2001	15,685	1,228	91	60	247	420	1,028	1,135	799	1,231	2,761	6,675	10
2000	15,226	1,274	90	62	246	431	904	948	778	1,287	2,730	6,471	5
1999	13,219	1,280	83	54	166	332	676	676	655	1,138	2,434	5,714	11
Non-Hispanic white													
Both sexes													
2001	24,596	1,752	139	89	563	907	2,175	2,510	1,713	2,191	4,350	8,178	29
2000	24,064	1,805	134	88	518	918	2,027	2,173	1,652	2,394	4,358	7,975	22
1999	20,200	1,812	129	81	348	660	1,341	1,459	1,374	2,166	3,775	7,033	22
Male													
2001	11,404	1,046	79	52	376	614	1,387	1,634	1,058	1,184	1,920	2,032	22
2000	11,351	1,067	79	52	356	625	1,366	1,465	1,036	1,328	1,945	2,013	19
1999	9,249	1,083	82	45	244	431	884	963	870	1,222	1,638	1,771	16
Female													
2001	13,192	706	60	37	187	293	788	876	655	1,007	2,430	6,146	7
2000	12,713	738	55	36	162	293	661	708	616	1,066	2,413	5,962	3
1999	10,951	729	47	36	104	229	457	496	504	944	2,137	5,262	6
Non-Hispanic black													
Both sexes													
2001	4,606	1,081	69	42	158	266	518	605	373	401	466	622	5

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)													
Non-Hispanic black													
Both sexes													
2000	4,892	1,186	54	46	179	308	514	609	411	472	505	595	13
1999	4,508	1,149	67	33	152	260	504	520	371	395	488	563	6
Male													
2001	2,414	614	44	26	109	150	300	364	262	207	174	161	3
2000	2,614	694	26	28	105	191	283	390	266	267	214	139	11
1999	2,460	658	35	20	100	172	299	352	238	218	214	152	2
Female													
2001	2,192	467	25	16	49	116	218	241	111	194	292	461	2
2000	2,278	492	28	18	74	117	231	219	145	205	291	456	2
1999	2,048	491	32	13	52	88	205	168	133	177	274	411	4
Origin not stated [2]													
Both sexes													
2001	205	17	-	1	8	17	17	28	11	9	10	20	67
2000	291	25	1	2	8	15	41	42	30	25	17	16	69
1999	210	26	2	-	5	6	29	35	17	24	10	16	40
Male													
2001	123	11	-	-	5	12	14	16	10	4	3	7	41
2000	206	12	-	1	5	11	32	30	26	19	14	6	50
1999	147	18	1	-	3	3	20	31	14	12	8	4	33
Female													
2001	82	6	-	1	3	5	3	12	1	5	7	13	26
2000	85	13	1	1	3	4	9	12	4	6	3	10	19

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)													
Origin not stated [2]													
Female													
1999	63	8	1	-	2	3	9	4	3	12	2	12	7
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25, G31-H93,K00-K22,K29-K31,K50-K66,K71-K72, K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0, N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)													
All origins													
Both sexes													
2001	186,406	1,131	646	860	1,799	3,072	8,084	14,203	16,048	25,803	49,711	65,032	17
2000	178,378	1,105	609	879	1,764	3,097	8,113	13,326	15,156	25,007	48,019	61,292	11
1999	171,683	1,148	647	921	1,771	3,062	7,989	12,383	14,390	25,165	45,598	58,599	10
Male													
2001	75,162	649	358	454	1,064	1,729	4,774	8,397	8,652	12,732	19,817	16,520	16
2000	72,227	629	338	474	1,057	1,811	4,801	7,911	8,236	12,321	19,102	15,536	11
1999	69,784	680	354	483	1,045	1,790	4,835	7,272	7,870	12,332	18,495	14,623	5
Female													
2001	111,244	482	288	406	735	1,343	3,310	5,806	7,396	13,071	29,894	48,512	1
2000	106,151	476	271	405	707	1,286	3,312	5,415	6,920	12,686	28,917	45,756	-
1999	101,899	468	293	438	726	1,272	3,154	5,111	6,520	12,833	27,103	43,976	5
Hispanic													
Both sexes													
2001	7,847	187	112	122	246	354	671	892	918	1,218	1,539	1,585	3
2000	7,325	196	109	113	214	381	706	862	813	1,113	1,438	1,376	4
1999	7,291	200	106	107	218	409	690	837	842	1,133	1,265	1,482	2

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25, G31-H93,K00-K22,K29-K31,K50-K66,K71-K72, K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0, N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)													
Hispanic													
Male													
2001	3,849	106	66	65	147	222	451	584	513	587	612	493	3
2000	3,663	100	54	63	140	246	483	552	464	568	557	432	4
1999	3,655	124	50	54	128	275	479	541	481	564	530	428	1
Female													
2001	3,998	81	46	57	99	132	220	308	405	631	927	1,092	-
2000	3,662	96	55	50	74	135	223	310	349	545	881	944	-
1999	3,636	76	56	53	90	134	211	296	361	569	735	1,054	1
Non-Hispanic [1]													
Both sexes													
2001	177,985	934	532	736	1,545	2,709	7,370	13,230	15,063	24,506	48,059	63,290	11
2000	170,462	898	497	762	1,541	2,697	7,366	12,376	14,283	23,799	46,458	59,780	5
1999	163,832	935	537	814	1,547	2,645	7,247	11,467	13,479	23,956	44,232	56,968	5
Male													
2001	71,014	539	291	388	912	1,499	4,293	7,751	8,087	12,104	19,155	15,985	10
2000	68,263	522	282	411	911	1,556	4,287	7,294	7,734	11,700	18,493	15,068	5
1999	65,835	546	302	429	913	1,509	4,316	6,676	7,341	11,723	17,924	14,153	3
Female													
2001	106,971	395	241	348	633	1,210	3,077	5,479	6,976	12,402	28,904	47,305	1
2000	102,199	376	215	351	630	1,141	3,079	5,082	6,549	12,099	27,965	44,712	-
1999	97,997	389	235	385	634	1,136	2,931	4,791	6,138	12,233	26,308	42,815	2

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25, G31-H93,K00-K22,K29-K31,K50-K66,K71-K72, K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0, N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)													
Non-Hispanic white													
Both sexes													
2001	153,156	556	362	541	1,089	1,808	5,201	9,814	11,982	20,807	42,990	58,000	6
2000	146,518	544	316	537	1,113	1,807	5,202	9,186	11,332	20,188	41,562	54,729	2
1999	141,002	554	375	569	1,100	1,845	5,036	8,437	10,749	20,267	39,663	52,406	1
Male													
2001	60,022	311	209	286	677	1,040	3,130	5,873	6,476	10,312	17,140	14,563	5
2000	57,565	310	183	307	683	1,105	3,128	5,569	6,149	9,933	16,519	13,677	2
1999	55,517	321	214	291	681	1,110	3,144	5,029	5,906	9,977	16,013	12,830	1
Female													
2001	93,134	245	153	255	412	768	2,071	3,941	5,506	10,495	25,850	43,437	1
2000	88,953	234	133	230	430	702	2,074	3,617	5,183	10,255	25,043	41,052	-
1999	85,485	233	161	278	419	735	1,892	3,408	4,843	10,290	23,650	39,576	-
Non-Hispanic black													
Both sexes													
2001	21,752	329	131	165	386	758	1,944	3,106	2,691	3,236	4,415	4,586	5
2000	21,081	296	144	189	373	810	1,948	2,897	2,607	3,153	4,248	4,413	3
1999	20,137	332	129	201	385	704	1,997	2,728	2,437	3,251	3,963	4,006	4
Male													
2001	9,561	192	62	83	196	381	1,030	1,713	1,411	1,570	1,767	1,151	5
2000	9,351	180	77	87	197	400	1,033	1,562	1,411	1,548	1,711	1,142	3
1999	8,967	192	67	118	202	344	1,040	1,487	1,270	1,518	1,630	1,097	2

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25, G31-H93,K00-K22,K29-K31,K50-K66,K71-K72, K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0, N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)													
Non-Hispanic black													
Female													
2001	12,191	137	69	82	190	377	914	1,393	1,280	1,666	2,648	3,435	-
2000	11,730	116	67	102	176	410	915	1,335	1,196	1,605	2,537	3,271	-
1999	11,170	140	62	83	183	360	957	1,241	1,167	1,733	2,333	2,909	2
Origin not stated [2]													
Both sexes													
2001	574	10	2	2	8	9	43	81	67	79	113	157	3
2000	591	11	3	4	9	19	41	88	60	95	123	136	2
1999	560	13	4	-	6	8	52	79	69	76	101	149	3
Male													
2001	299	4	1	1	5	8	30	62	52	41	50	42	3
2000	301	7	2	-	6	9	31	65	38	53	52	36	2
1999	294	10	2	-	4	6	40	55	48	45	41	42	1
Female													
2001	275	6	1	1	3	1	13	19	15	38	63	115	-
2000	290	4	1	4	3	10	10	23	22	42	71	100	-
1999	266	3	2	-	2	2	12	24	21	31	60	107	2
Accidents (unintentional injuries) (V01-X59,Y85-Y86)													
All origins													
Both sexes													
2001	101,537	976	1,714	2,836	14,411	11,839	15,945	13,344	7,658	7,835	12,688	12,171	120
2000	97,900	881	1,826	2,979	14,113	11,769	15,413	12,278	7,505	7,698	11,758	11,595	85

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Accidents (unintentional injuries) (V01-X59,Y85-Y86)													
All origins													
Both sexes													
1999	97,860	845	1,898	3,091	13,656	11,890	15,231	11,639	7,285	8,208	12,282	11,729	106
Male													
2001	66,060	541	1,045	1,803	10,832	9,092	11,612	9,743	5,228	4,751	6,652	4,657	104
2000	63,817	501	1,098	1,861	10,460	9,059	11,444	9,063	5,133	4,639	6,063	4,413	83
1999	63,535	494	1,127	1,974	10,040	9,097	11,248	8,602	5,015	4,992	6,372	4,480	94
Female													
2001	35,477	435	669	1,033	3,579	2,747	4,333	3,601	2,430	3,084	6,036	7,514	16
2000	34,083	380	728	1,118	3,653	2,710	3,969	3,215	2,372	3,059	5,695	7,182	2
1999	34,325	351	771	1,117	3,616	2,793	3,983	3,037	2,270	3,216	5,910	7,249	12
Hispanic													
Both sexes													
2001	9,523	154	310	379	2,182	1,938	1,760	1,078	564	454	409	266	29
2000	8,830	118	306	383	2,095	1,776	1,618	953	492	418	365	278	28
1999	8,650	134	341	362	1,946	1,790	1,607	926	466	440	334	260	44
Male													
2001	7,157	75	195	224	1,773	1,619	1,418	822	391	275	235	104	26
2000	6,696	67	174	210	1,717	1,499	1,317	750	347	271	191	125	28
1999	6,578	80	197	227	1,575	1,494	1,307	734	334	288	182	120	40
Female													
2001	2,366	79	115	155	409	319	342	256	173	179	174	162	3
2000	2,134	51	132	173	378	277	301	203	145	147	174	153	-
1999	2,072	54	144	135	371	296	300	192	132	152	152	140	4

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Accidents (unintentional injuries) (V01-X59,Y85-Y86)													
Non-Hispanic [1]													
Both sexes													
2001	91,522	817	1,399	2,445	12,187	9,857	14,089	12,184	7,042	7,347	12,245	11,874	36
2000	88,558	759	1,512	2,585	11,968	9,935	13,706	11,219	6,966	7,247	11,361	11,288	12
1999	88,669	710	1,550	2,721	11,658	10,034	13,529	10,622	6,753	7,738	11,896	11,432	26
Male													
2001	58,528	462	845	1,570	9,024	7,438	10,117	8,852	4,795	4,455	6,397	4,544	29
2000	56,736	431	921	1,648	8,707	7,512	10,054	8,231	4,747	4,344	5,855	4,275	11
1999	56,551	414	926	1,739	8,428	7,550	9,871	7,791	4,627	4,681	6,158	4,343	23
Female													
2001	32,994	355	554	875	3,163	2,419	3,972	3,332	2,247	2,892	5,848	7,330	7
2000	31,822	328	591	937	3,261	2,423	3,652	2,988	2,219	2,903	5,506	7,013	1
1999	32,118	296	624	982	3,230	2,484	3,658	2,831	2,126	3,057	5,738	7,089	3
Non-Hispanic white													
Both sexes													
2001	76,262	500	1,006	1,688	9,867	7,798	11,340	9,704	5,776	6,309	11,157	11,092	25
2000	73,536	455	1,033	1,843	9,735	7,730	11,008	8,882	5,753	6,181	10,356	10,552	8
1999	73,326	435	1,042	1,908	9,410	7,855	10,844	8,392	5,455	6,568	10,757	10,647	13
Male													
2001	48,180	281	624	1,066	7,297	5,906	8,188	7,034	3,905	3,837	5,800	4,220	22
2000	46,470	266	638	1,176	7,065	5,861	8,113	6,521	3,871	3,692	5,292	3,968	7
1999	46,113	258	617	1,205	6,749	5,932	7,949	6,142	3,730	3,962	5,551	4,006	12
Female													
2001	28,082	219	382	622	2,570	1,892	3,152	2,670	1,871	2,472	5,357	6,872	3
2000	27,066	189	395	667	2,670	1,869	2,895	2,361	1,882	2,489	5,064	6,584	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Accidents (unintentional injuries) (V01-X59,Y85-Y86)													
Non-Hispanic white													
Female													
1999	27,213	177	425	703	2,661	1,923	2,895	2,250	1,725	2,606	5,206	6,641	1
Non-Hispanic black													
Both sexes													
2001	12,239	272	331	630	1,713	1,600	2,282	2,103	1,045	836	830	589	8
2000	12,087	268	396	619	1,662	1,740	2,253	1,961	983	857	768	576	4
1999	12,533	251	428	701	1,694	1,738	2,245	1,903	1,047	943	925	646	12
Male													
2001	8,381	158	188	432	1,285	1,192	1,608	1,574	750	505	461	222	6
2000	8,380	145	232	406	1,233	1,316	1,649	1,454	736	537	432	236	4
1999	8,629	140	260	467	1,281	1,313	1,628	1,441	742	596	483	268	10
Female													
2001	3,858	114	143	198	428	408	674	529	295	331	369	367	2
2000	3,707	123	164	213	429	424	604	507	247	320	336	340	-
1999	3,904	111	168	234	413	425	617	462	305	347	442	378	2
Origin not stated [2]													
Both sexes													
2001	492	5	5	12	42	44	96	82	52	34	34	31	55
2000	512	4	8	11	50	58	89	106	47	33	32	29	45
1999	541	1	7	8	52	66	95	91	66	30	52	37	36
Male													
2001	375	4	5	9	35	35	77	69	42	21	20	9	49
2000	385	3	3	3	36	48	73	82	39	24	17	13	44
1999	406	-	4	8	37	53	70	77	54	23	32	17	31

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Accidents (unintentional injuries) (V01-X59,Y85-Y86)													
Origin not stated [2]													
Female													
2001	117	1	-	3	7	9	19	13	10	13	14	22	6
2000	127	1	5	8	14	10	16	24	8	9	15	16	1
1999	135	1	3	-	15	13	25	14	12	7	20	20	5
Transport accidents (V01-V99,Y85)													
All origins													
Both sexes													
2001	47,288	151	678	1,823	11,139	7,454	7,827	6,251	3,861	3,282	3,507	1,272	43
2000	46,749	174	691	1,922	11,028	7,400	7,661	5,965	3,883	3,315	3,342	1,335	33
1999	46,423	189	712	1,975	10,689	7,434	7,553	5,645	3,812	3,579	3,468	1,330	37
Male													
2001	32,794	68	392	1,115	8,111	5,661	5,613	4,484	2,631	2,015	1,983	684	37
2000	32,166	94	391	1,138	7,890	5,570	5,560	4,260	2,617	2,021	1,858	734	33
1999	31,642	101	400	1,206	7,558	5,511	5,376	4,016	2,561	2,162	1,959	758	34
Female													
2001	14,494	83	286	708	3,028	1,793	2,214	1,767	1,230	1,267	1,524	588	6
2000	14,583	80	300	784	3,138	1,830	2,101	1,705	1,266	1,294	1,484	601	-
1999	14,781	88	312	769	3,131	1,923	2,177	1,629	1,251	1,417	1,509	572	3
Hispanic													
Both sexes													
2001	5,774	43	159	265	1,740	1,293	942	557	331	229	167	38	10
2000	5,217	42	154	279	1,616	1,126	831	508	268	220	121	38	14
1999	4,880	40	157	250	1,457	1,125	758	445	253	228	120	36	11

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Transport accidents (V01-V99,Y85)													
Hispanic													
Male													
2001	4,334	16	92	155	1,383	1,067	731	390	214	141	114	23	8
2000	3,921	24	85	155	1,285	921	639	373	189	140	74	22	14
1999	3,630	22	85	151	1,136	904	590	330	161	144	77	19	11
Female													
2001	1,440	27	67	110	357	226	211	167	117	88	53	15	2
2000	1,296	18	69	124	331	205	192	135	79	80	47	16	-
1999	1,250	18	72	99	321	221	168	115	92	84	43	17	-
Non-Hispanic [1]													
Both sexes													
2001	41,339	108	517	1,554	9,371	6,141	6,848	5,668	3,513	3,044	3,330	1,230	15
2000	41,341	132	533	1,637	9,375	6,244	6,801	5,430	3,595	3,082	3,214	1,295	3
1999	41,330	149	553	1,719	9,192	6,272	6,766	5,175	3,535	3,338	3,329	1,291	11
Male													
2001	28,327	52	298	957	6,706	4,578	4,855	4,071	2,405	1,869	1,865	659	12
2000	28,096	70	304	982	6,580	4,624	4,896	3,865	2,411	1,871	1,779	711	3
1999	27,850	79	313	1,049	6,396	4,578	4,764	3,666	2,379	2,009	1,869	738	10
Female													
2001	13,012	56	219	597	2,665	1,563	1,993	1,597	1,108	1,175	1,465	571	3
2000	13,245	62	229	655	2,795	1,620	1,905	1,565	1,184	1,211	1,435	584	-
1999	13,480	70	240	670	2,796	1,694	2,002	1,509	1,156	1,329	1,460	553	1
Non-Hispanic white													
Both sexes													
2001	33,753	74	353	1,122	7,563	4,771	5,494	4,611	2,937	2,628	3,044	1,148	8

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Transport accidents (V01-V99,Y85)													
Non-Hispanic white													
Both sexes													
2000	33,808	79	357	1,246	7,669	4,759	5,471	4,402	3,039	2,649	2,937	1,197	3
1999	33,803	98	349	1,291	7,493	4,889	5,425	4,211	2,946	2,897	3,020	1,180	4
Male													
2001	23,047	36	208	686	5,401	3,561	3,887	3,326	2,009	1,617	1,698	611	7
2000	22,855	44	205	748	5,361	3,518	3,935	3,160	2,014	1,597	1,609	661	3
1999	22,682	58	192	782	5,181	3,561	3,827	2,992	1,986	1,745	1,686	668	4
Female													
2001	10,706	38	145	436	2,162	1,210	1,607	1,285	928	1,011	1,346	537	1
2000	10,953	35	152	498	2,308	1,241	1,536	1,242	1,025	1,052	1,328	536	-
1999	11,121	40	157	509	2,312	1,328	1,598	1,219	960	1,152	1,334	512	-
Non-Hispanic black													
Both sexes													
2001	5,803	18	127	354	1,322	1,058	1,071	829	455	307	198	57	7
2000	5,757	40	139	309	1,262	1,152	1,059	809	417	322	179	69	-
1999	5,810	44	168	360	1,262	1,075	1,074	758	430	314	232	86	7
Male													
2001	4,143	7	69	234	966	797	779	615	332	185	123	31	5
2000	4,127	21	77	191	907	875	796	571	314	218	118	39	-
1999	4,097	18	96	229	909	817	769	557	303	196	139	58	6
Female													
2001	1,660	11	58	120	356	261	292	214	123	122	75	26	2
2000	1,630	19	62	118	355	277	263	238	103	104	61	30	-
1999	1,713	26	72	131	353	258	305	201	127	118	93	28	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Transport accidents (V01-V99,Y85)													
Origin not stated [2]													
Both sexes													
2001	175	-	2	4	28	20	37	26	17	9	10	4	18
2000	191	-	4	6	37	30	29	27	20	13	7	2	16
1999	213	-	2	6	40	37	29	25	24	13	19	3	15
Male													
2001	133	-	2	3	22	16	27	23	12	5	4	2	17
2000	149	-	2	1	25	25	25	22	17	10	5	1	16
1999	162	-	2	6	26	29	22	20	21	9	13	1	13
Female													
2001	42	-	-	1	6	4	10	3	5	4	6	2	1
2000	42	-	2	5	12	5	4	5	3	3	2	1	-
1999	51	-	-	-	14	8	7	5	3	4	6	2	2
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1, V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)													
All origins													
Both sexes													
2001	43,788	144	626	1,686	10,725	6,937	7,083	5,564	3,465	2,990	3,312	1,223	33
2000	43,354	168	651	1,772	10,560	6,884	6,927	5,361	3,506	3,038	3,173	1,288	26
1999	42,401	184	650	1,771	10,128	6,778	6,738	4,972	3,370	3,276	3,243	1,261	30
Male													
2001	30,064	66	356	1,029	7,778	5,236	5,016	3,938	2,309	1,786	1,859	664	27
2000	29,451	90	368	1,043	7,516	5,135	4,940	3,780	2,306	1,802	1,739	706	26
1999	28,552	97	366	1,062	7,127	4,993	4,735	3,473	2,200	1,931	1,827	714	27

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1, V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)													
All origins													
Female													
2001	13,724	78	270	657	2,947	1,701	2,067	1,626	1,156	1,204	1,453	559	6
2000	13,903	78	283	729	3,044	1,749	1,987	1,581	1,200	1,236	1,434	582	-
1999	13,849	87	284	709	3,001	1,785	2,003	1,499	1,170	1,345	1,416	547	3
Hispanic													
Both sexes													
2001	5,272	40	142	246	1,677	1,197	834	470	279	201	146	32	8
2000	4,900	40	144	260	1,557	1,066	756	472	246	198	113	34	14
1999	4,525	40	148	231	1,384	1,035	684	405	238	204	113	33	10
Male													
2001	3,987	15	82	144	1,330	992	653	340	184	123	99	19	6
2000	3,658	24	78	142	1,236	869	574	341	170	123	68	19	14
1999	3,337	22	82	136	1,077	828	530	294	146	124	71	17	10
Female													
2001	1,285	25	60	102	347	205	181	130	95	78	47	13	2
2000	1,242	16	66	118	321	197	182	131	76	75	45	15	-
1999	1,188	18	66	95	307	207	154	111	92	80	42	16	-
Non-Hispanic [1]													
Both sexes													
2001	38,362	104	483	1,436	9,021	5,722	6,217	5,072	3,173	2,780	3,156	1,187	11
2000	38,295	128	505	1,507	8,969	5,792	6,147	4,866	3,243	2,829	3,054	1,252	3
1999	37,684	144	500	1,534	8,707	5,712	6,026	4,546	3,110	3,059	3,112	1,225	9

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1, V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)													
Non-Hispanic [1]													
Male													
2001	25,962	51	273	882	6,426	4,230	4,340	3,579	2,116	1,658	1,756	643	8
2000	25,670	66	288	900	6,257	4,243	4,346	3,421	2,122	1,671	1,667	686	3
1999	25,072	75	282	920	6,027	4,141	4,184	3,163	2,034	1,798	1,744	696	8
Female													
2001	12,400	53	210	554	2,595	1,492	1,877	1,493	1,057	1,122	1,400	544	3
2000	12,625	62	217	607	2,712	1,549	1,801	1,445	1,121	1,158	1,387	566	-
1999	12,612	69	218	614	2,680	1,571	1,842	1,383	1,076	1,261	1,368	529	1
Non-Hispanic white													
Both sexes													
2001	31,283	71	323	1,036	7,266	4,424	4,980	4,121	2,652	2,398	2,891	1,113	8
2000	31,219	75	340	1,136	7,330	4,379	4,923	3,925	2,728	2,431	2,791	1,158	3
1999	30,743	94	315	1,144	7,099	4,425	4,797	3,678	2,579	2,655	2,832	1,122	3
Male													
2001	21,073	35	185	631	5,164	3,270	3,468	2,915	1,764	1,436	1,600	598	7
2000	20,789	40	197	680	5,094	3,196	3,473	2,781	1,759	1,424	1,505	637	3
1999	20,362	55	175	683	4,884	3,201	3,338	2,565	1,689	1,561	1,575	633	3
Female													
2001	10,210	36	138	405	2,102	1,154	1,512	1,206	888	962	1,291	515	1
2000	10,430	35	143	456	2,236	1,183	1,450	1,144	969	1,007	1,286	521	-
1999	10,381	39	140	461	2,215	1,224	1,459	1,113	890	1,094	1,257	489	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1, V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)													
Non-Hispanic black													
Both sexes													
2001	5,423	18	123	324	1,285	1,007	985	747	410	283	184	54	3
2000	5,424	40	132	294	1,216	1,095	977	750	388	298	169	65	-
1999	5,388	43	152	331	1,209	1,006	985	685	387	291	214	79	6
Male													
2001	3,842	7	67	214	935	757	706	550	295	163	116	31	1
2000	3,857	21	72	179	868	828	727	524	289	199	112	38	-
1999	3,763	17	84	203	869	761	696	499	267	178	132	52	5
Female													
2001	1,581	11	56	110	350	250	279	197	115	120	68	23	2
2000	1,567	19	60	115	348	267	250	226	99	99	57	27	-
1999	1,625	26	68	128	340	245	289	186	120	113	82	27	1
Origin not stated [2]													
Both sexes													
2001	154	-	1	4	27	18	32	22	13	9	10	4	14
2000	159	-	2	5	34	26	24	23	17	11	6	2	9
1999	192	-	2	6	37	31	28	21	22	13	18	3	11
Male													
2001	115	-	1	3	22	14	23	19	9	5	4	2	13
2000	123	-	2	1	23	23	20	18	14	8	4	1	9
1999	143	-	2	6	23	24	21	16	20	9	12	1	9

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1, V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)													
Origin not stated [2]													
Female													
2001	39	-	-	1	5	4	9	3	4	4	6	2	1
2000	36	-	-	4	11	3	4	5	3	3	2	1	-
1999	49	-	-	-	14	7	7	5	2	4	6	2	2
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11, V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9, V88.9,V89.1,V89.3,V89.9)													
All origins													
Both sexes													
2001	1,405	3	28	80	198	177	285	247	133	117	98	29	10
2000	1,492	2	31	98	243	212	317	234	122	100	91	35	7
1999	2,118	4	48	128	362	317	408	304	182	145	159	54	7
Male													
2001	1,067	-	20	49	158	152	235	189	104	84	56	10	10
2000	1,125	2	17	61	194	171	258	166	98	75	57	19	7
1999	1,559	4	26	91	279	244	319	234	135	97	88	35	7
Female													
2001	338	3	8	31	40	25	50	58	29	33	42	19	-
2000	367	-	14	37	49	41	59	68	24	25	34	16	-
1999	559	-	22	37	83	73	89	70	47	48	71	19	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11, V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9, V88.9,V89.1,V89.3,V89.9)													
Hispanic													
Both sexes													
2001	200	-	8	7	39	38	42	29	14	5	13	3	2
2000	201	-	10	15	34	36	44	25	14	14	5	4	-
1999	264	-	8	14	55	60	57	31	8	21	7	2	1
Male													
2001	173	-	6	4	37	35	39	22	12	4	10	2	2
2000	166	-	7	11	31	32	36	22	11	10	3	3	-
1999	220	-	3	12	45	53	47	27	8	17	6	1	1
Female													
2001	27	-	2	3	2	3	3	7	2	1	3	1	-
2000	35	-	3	4	3	4	8	3	3	4	2	1	-
1999	44	-	5	2	10	7	10	4	-	4	1	1	-
Non-Hispanic [1]													
Both sexes													
2001	1,187	3	19	73	158	137	241	214	115	112	85	26	4
2000	1,262	2	19	82	206	172	268	207	105	85	85	31	-
1999	1,836	4	40	114	304	252	350	271	172	124	151	52	2
Male													
2001	878	-	13	45	121	115	194	163	89	80	46	8	4
2000	936	2	10	50	161	137	217	142	84	64	53	16	-
1999	1,323	4	23	79	231	187	271	205	126	80	81	34	2

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11, V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9, V88.9,V89.1,V89.3,V89.9)													
Non-Hispanic [1]													
Female													
2001	309	3	6	28	37	22	47	51	26	32	39	18	-
2000	326	-	9	32	45	35	51	65	21	21	32	15	-
1999	513	-	17	35	73	65	79	66	46	44	70	18	-
Non-Hispanic white													
Both sexes													
2001	901	2	16	49	126	103	165	165	89	92	73	21	-
2000	997	2	9	67	168	132	201	161	84	70	75	28	-
1999	1,480	3	25	86	239	195	282	219	151	102	131	46	1
Male													
2001	673	-	12	30	99	89	133	129	73	63	40	5	-
2000	741	2	3	41	130	106	165	110	68	51	49	16	-
1999	1,051	3	12	57	177	143	218	163	110	67	71	29	1
Female													
2001	228	2	4	19	27	14	32	36	16	29	33	16	-
2000	256	-	6	26	38	26	36	51	16	19	26	12	-
1999	429	-	13	29	62	52	64	56	41	35	60	17	-
Non-Hispanic black													
Both sexes													
2001	216	-	3	23	21	24	59	38	22	15	6	1	4
2000	203	-	7	11	30	32	54	34	15	10	7	3	-
1999	248	1	12	20	33	42	52	40	17	14	11	5	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11, V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9, V88.9,V89.1,V89.3,V89.9)													
Non-Hispanic black													
Male													
2001	159	-	1	15	15	18	48	28	14	13	3	-	4
2000	155	-	5	8	24	25	43	26	12	9	3	-	-
1999	192	1	9	19	27	34	41	32	12	9	3	4	1
Female													
2001	57	-	2	8	6	6	11	10	8	2	3	1	-
2000	48	-	2	3	6	7	11	8	3	1	4	3	-
1999	56	-	3	1	6	8	11	8	5	5	8	1	-
Origin not stated [2]													
Both sexes													
2001	18	-	1	-	1	2	2	4	4	-	-	-	4
2000	29	-	2	1	3	4	5	2	3	1	1	-	7
1999	18	-	-	-	3	5	1	2	2	-	1	-	4
Male													
2001	16	-	1	-	-	2	2	4	3	-	-	-	4
2000	23	-	-	-	2	2	5	2	3	1	1	-	7
1999	16	-	-	-	3	4	1	2	1	-	1	-	4
Female													
2001	2	-	-	-	1	-	-	-	1	-	-	-	-
2000	6	-	2	1	1	2	-	-	-	-	-	-	-
1999	2	-	-	-	-	1	-	-	1	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)													
All origins													
Both sexes													
2001	2,095	4	24	57	216	340	459	440	263	175	97	20	-
2000	1,903	4	9	52	225	304	417	370	255	177	78	12	-
1999	1,904	1	14	76	199	339	407	369	260	158	66	15	-
Male													
2001	1,663	2	16	37	175	273	362	357	218	145	68	10	-
2000	1,590	2	6	34	180	264	362	314	213	144	62	9	-
1999	1,531	-	8	53	152	274	322	309	226	134	44	9	-
Female													
2001	432	2	8	20	41	67	97	83	45	30	29	10	-
2000	313	2	3	18	45	40	55	56	42	33	16	3	-
1999	373	1	6	23	47	65	85	60	34	24	22	6	-
Hispanic													
Both sexes													
2001	302	3	9	12	24	58	66	58	38	23	8	3	-
2000	116	2	-	4	25	24	31	11	8	8	3	-	-
1999	91	-	1	5	18	30	17	9	7	3	-	1	-
Male													
2001	174	1	4	7	16	40	39	28	18	14	5	2	-
2000	97	-	-	2	18	20	29	10	8	7	3	-	-
1999	73	-	-	3	14	23	13	9	7	3	-	1	-
Female													
2001	128	2	5	5	8	18	27	30	20	9	3	1	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)													
Hispanic													
Female													
2000	19	2	-	2	7	4	2	1	-	1	-	-	-
1999	18	-	1	2	4	7	4	-	-	-	-	-	-
Non-Hispanic [1]													
Both sexes													
2001	1,790	1	15	45	192	282	390	382	225	152	89	17	-
2000	1,784	2	9	48	200	280	386	357	247	168	75	12	-
1999	1,810	1	13	71	181	308	390	358	253	155	66	14	-
Male													
2001	1,487	1	12	30	159	233	321	329	200	131	63	8	-
2000	1,490	2	6	32	162	244	333	302	205	136	59	9	-
1999	1,455	-	8	50	138	250	309	298	219	131	44	8	-
Female													
2001	303	-	3	15	33	49	69	53	25	21	26	9	-
2000	294	-	3	16	38	36	53	55	42	32	16	3	-
1999	355	1	5	21	43	58	81	60	34	24	22	6	-
Non-Hispanic white													
Both sexes													
2001	1,569	1	14	37	171	244	349	325	196	138	80	14	-
2000	1,592	2	8	43	171	248	347	316	227	148	71	11	-
1999	1,580	1	9	61	155	269	346	314	216	140	57	12	-
Male													
2001	1,301	1	11	25	138	202	286	282	172	118	58	8	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)													
Non-Hispanic white													
Male													
2000	1,325	2	5	27	137	216	297	269	187	122	55	8	-
1999	1,269	-	5	42	120	217	271	264	187	117	40	6	-
Female													
2001	268	-	3	12	33	42	63	43	24	20	22	6	-
2000	267	-	3	16	34	32	50	47	40	26	16	3	-
1999	311	1	4	19	35	52	75	50	29	23	17	6	-
Non-Hispanic black													
Both sexes													
2001	164	-	1	7	16	27	27	44	23	9	8	2	-
2000	130	-	-	4	16	25	28	25	14	14	3	1	-
1999	174	-	4	9	20	27	37	33	26	9	7	2	-
Male													
2001	142	-	1	5	16	22	25	37	23	9	4	-	-
2000	115	-	-	4	15	22	26	21	13	10	3	1	-
1999	142	-	3	7	13	22	32	26	24	9	4	2	-
Female													
2001	22	-	-	2	-	5	2	7	-	-	4	2	-
2000	15	-	-	-	1	3	2	4	1	4	-	-	-
1999	32	-	1	2	7	5	5	7	2	-	3	-	-
Origin not stated [2]													
Both sexes													
2001	3	-	-	-	-	-	3	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)													
Origin not stated [2]													
Both sexes													
2000	3	-	-	-	-	-	-	2	-	1	-	-	-
1999	3	-	-	-	-	1	-	2	-	-	-	-	-
Male													
2001	2	-	-	-	-	-	2	-	-	-	-	-	-
2000	3	-	-	-	-	-	-	2	-	1	-	-	-
1999	3	-	-	-	-	1	-	2	-	-	-	-	-
Female													
2001	1	-	-	-	-	-	1	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Nontransport accidents (W00-X59,Y86)													
All origins													
Both sexes													
2001	54,249	825	1,036	1,013	3,272	4,385	8,118	7,093	3,797	4,553	9,181	10,899	77
2000	51,151	707	1,135	1,057	3,085	4,369	7,752	6,313	3,622	4,383	8,416	10,260	52
1999	51,437	656	1,186	1,116	2,967	4,456	7,678	5,994	3,473	4,629	8,814	10,399	69
Male													
2001	33,266	473	653	688	2,721	3,431	5,999	5,259	2,597	2,736	4,669	3,973	67
2000	31,651	407	707	723	2,570	3,489	5,884	4,803	2,516	2,618	4,205	3,679	50
1999	31,893	393	727	768	2,482	3,586	5,872	4,586	2,454	2,830	4,413	3,722	60
Female													
2001	20,983	352	383	325	551	954	2,119	1,834	1,200	1,817	4,512	6,926	10

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Nontransport accidents (W00-X59,Y86)													
All origins													
Female													
2000	19,500	300	428	334	515	880	1,868	1,510	1,106	1,765	4,211	6,581	2
1999	19,544	263	459	348	485	870	1,806	1,408	1,019	1,799	4,401	6,677	9
Hispanic													
Both sexes													
2001	3,749	111	151	114	442	645	818	521	233	225	242	228	19
2000	3,613	76	152	104	479	650	787	445	224	198	244	240	14
1999	3,770	94	184	112	489	665	849	481	213	212	214	224	33
Male													
2001	2,823	59	103	69	390	552	687	432	177	134	121	81	18
2000	2,775	43	89	55	432	578	678	377	158	131	117	103	14
1999	2,948	58	112	76	439	590	717	404	173	144	105	101	29
Female													
2001	926	52	48	45	52	93	131	89	56	91	121	147	1
2000	838	33	63	49	47	72	109	68	66	67	127	137	-
1999	822	36	72	36	50	75	132	77	40	68	109	123	4
Non-Hispanic [1]													
Both sexes													
2001	50,183	709	882	891	2,816	3,716	7,241	6,516	3,529	4,303	8,915	10,644	21
2000	47,217	627	979	948	2,593	3,691	6,905	5,789	3,371	4,165	8,147	9,993	9
1999	47,339	561	997	1,002	2,466	3,762	6,763	5,447	3,218	4,400	8,567	10,141	15
Male													
2001	30,201	410	547	613	2,318	2,860	5,262	4,781	2,390	2,586	4,532	3,885	17
2000	28,640	361	617	666	2,127	2,888	5,158	4,366	2,336	2,473	4,076	3,564	8

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Nontransport accidents (W00-X59,Y86)													
Non-Hispanic [1]													
Male													
1999	28,701	335	613	690	2,032	2,972	5,107	4,125	2,248	2,672	4,289	3,605	13
Female													
2001	19,982	299	335	278	498	856	1,979	1,735	1,139	1,717	4,383	6,759	4
2000	18,577	266	362	282	466	803	1,747	1,423	1,035	1,692	4,071	6,429	1
1999	18,638	226	384	312	434	790	1,656	1,322	970	1,728	4,278	6,536	2
Non-Hispanic white													
Both sexes													
2001	42,509	426	653	566	2,304	3,027	5,846	5,093	2,839	3,681	8,113	9,944	17
2000	39,728	376	676	597	2,066	2,971	5,537	4,480	2,714	3,532	7,419	9,355	5
1999	39,523	337	693	617	1,917	2,966	5,419	4,181	2,509	3,671	7,737	9,467	9
Male													
2001	25,133	245	416	380	1,896	2,345	4,301	3,708	1,896	2,220	4,102	3,609	15
2000	23,615	222	433	428	1,704	2,343	4,178	3,361	1,857	2,095	3,683	3,307	4
1999	23,431	200	425	423	1,568	2,371	4,122	3,150	1,744	2,217	3,865	3,338	8
Female													
2001	17,376	181	237	186	408	682	1,545	1,385	943	1,461	4,011	6,335	2
2000	16,113	154	243	169	362	628	1,359	1,119	857	1,437	3,736	6,048	1
1999	16,092	137	268	194	349	595	1,297	1,031	765	1,454	3,872	6,129	1
Non-Hispanic black													
Both sexes													
2001	6,436	254	204	276	391	542	1,211	1,274	590	529	632	532	1
2000	6,330	228	257	310	400	588	1,194	1,152	566	535	589	507	4
1999	6,723	207	260	341	432	663	1,171	1,145	617	629	693	560	5

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Nontransport accidents (W00-X59,Y86)													
Non-Hispanic black													
Male													
2001	4,238	151	119	198	319	395	829	959	418	320	338	191	1
2000	4,253	124	155	215	326	441	853	883	422	319	314	197	4
1999	4,532	122	164	238	372	496	859	884	439	400	344	210	4
Female													
2001	2,198	103	85	78	72	147	382	315	172	209	294	341	-
2000	2,077	104	102	95	74	147	341	269	144	216	275	310	-
1999	2,191	85	96	103	60	167	312	261	178	229	349	350	1
Origin not stated [2]													
Both sexes													
2001	317	5	3	8	14	24	59	56	35	25	24	27	37
2000	321	4	4	5	13	28	60	79	27	20	25	27	29
1999	328	1	5	2	12	29	66	66	42	17	33	34	21
Male													
2001	242	4	3	6	13	19	50	46	30	16	16	7	32
2000	236	3	1	2	11	23	48	60	22	14	12	12	28
1999	244	-	2	2	11	24	48	57	33	14	19	16	18
Female													
2001	75	1	-	2	1	5	9	10	5	9	8	20	5
2000	85	1	3	3	2	5	12	19	5	6	13	15	1
1999	84	1	3	-	1	5	18	9	9	3	14	18	3

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Falls (W00-W19)													
All origins													
Both sexes													
2001	15,019	23	32	66	256	340	647	1,024	1,004	1,833	4,440	5,350	4
2000	13,322	8	36	37	237	303	608	871	949	1,660	3,841	4,772	-
1999	13,162	12	55	53	242	343	645	824	887	1,663	3,856	4,578	4
Male													
2001	8,089	15	25	46	232	295	544	821	718	1,090	2,277	2,023	3
2000	7,122	6	23	30	203	272	496	690	680	991	1,947	1,784	-
1999	7,109	8	37	40	209	294	528	661	650	1,027	1,950	1,701	4
Female													
2001	6,930	8	7	20	24	45	103	203	286	743	2,163	3,327	1
2000	6,200	2	13	7	34	31	112	181	269	669	1,894	2,988	-
1999	6,053	4	18	13	33	49	117	163	237	636	1,906	2,877	-
Hispanic													
Both sexes													
2001	757	7	6	14	44	73	100	88	78	103	131	112	1
2000	693	3	9	3	46	73	66	77	74	86	134	122	-
1999	652	4	9	3	45	68	70	65	63	84	116	123	2
Male													
2001	525	4	5	13	44	71	93	75	62	57	62	38	1
2000	485	2	8	3	44	70	62	69	53	60	62	52	-
1999	464	4	6	3	43	65	62	56	55	52	60	56	2
Female													
2001	232	3	1	1	-	2	7	13	16	46	69	74	-
2000	208	1	1	-	2	3	4	8	21	26	72	70	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Falls (W00-W19)													
Hispanic													
Female													
1999	188	-	3	-	2	3	8	9	8	32	56	67	-
Non-Hispanic [1]													
Both sexes													
2001	14,204	15	26	51	212	266	544	929	921	1,718	4,299	5,223	-
2000	12,585	5	27	34	191	229	537	784	869	1,569	3,700	4,640	-
1999	12,447	8	46	50	197	273	573	753	813	1,572	3,721	4,440	1
Male													
2001	7,527	10	20	33	188	223	448	739	651	1,024	2,209	1,982	-
2000	6,603	4	15	27	159	201	429	612	622	928	1,881	1,725	-
1999	6,602	4	31	37	166	227	464	599	587	969	1,878	1,639	1
Female													
2001	6,677	5	6	18	24	43	96	190	270	694	2,090	3,241	-
2000	5,982	1	12	7	32	28	108	172	247	641	1,819	2,915	-
1999	5,845	4	15	13	31	46	109	154	226	603	1,843	2,801	-
Non-Hispanic white													
Both sexes													
2001	13,037	8	23	42	193	226	435	768	780	1,540	4,044	4,978	-
2000	11,537	2	15	27	172	192	438	646	725	1,393	3,487	4,440	-
1999	11,295	4	34	39	168	216	477	593	653	1,379	3,500	4,231	1
Male													
2001	6,785	5	17	27	173	191	360	603	551	921	2,067	1,870	-
2000	5,949	1	6	22	144	169	354	505	519	829	1,764	1,636	-
1999	5,860	3	23	29	144	181	380	474	471	846	1,759	1,549	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Falls (W00-W19)													
Non-Hispanic white													
Female													
2001	6,252	3	6	15	20	35	75	165	229	619	1,977	3,108	-
2000	5,588	1	9	5	28	23	84	141	206	564	1,723	2,804	-
1999	5,435	1	11	10	24	35	97	119	182	533	1,741	2,682	-
Non-Hispanic black													
Both sexes													
2001	837	6	3	6	13	26	90	128	108	143	173	141	-
2000	745	2	12	7	12	22	71	111	114	134	137	123	-
1999	865	2	11	10	21	42	73	136	127	150	153	140	-
Male													
2001	537	4	3	4	9	19	72	109	76	85	97	59	-
2000	471	2	9	5	10	18	54	88	87	76	70	52	-
1999	554	1	8	7	16	34	65	107	90	100	77	49	-
Female													
2001	300	2	-	2	4	7	18	19	32	58	76	82	-
2000	274	-	3	2	2	4	17	23	27	58	67	71	-
1999	311	1	3	3	5	8	8	29	37	50	76	91	-
Origin not stated [2]													
Both sexes													
2001	58	1	-	1	-	1	3	7	5	12	10	15	3
2000	44	-	-	-	-	1	5	10	6	5	7	10	-
1999	63	-	-	-	-	2	2	6	11	7	19	15	1
Male													
2001	37	1	-	-	-	1	3	7	5	9	6	3	2

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Falls (W00-W19)													
Origin not stated [2]													
Male													
2000	34	-	-	-	-	1	5	9	5	3	4	7	-
1999	43	-	-	-	-	2	2	6	8	6	12	6	1
Female													
2001	21	-	-	1	-	-	-	-	-	3	4	12	1
2000	10	-	-	-	-	-	-	1	1	2	3	3	-
1999	20	-	-	-	-	-	-	-	3	1	7	9	-
Accidental discharge of firearms (W32-W34)													
All origins													
Both sexes													
2001	802	-	15	57	206	122	146	103	69	33	41	8	2
2000	776	1	18	67	202	131	153	78	50	41	24	11	-
1999	824	-	12	76	251	143	136	92	48	33	23	10	-
Male													
2001	690	-	12	49	193	103	121	87	55	26	36	7	1
2000	671	-	11	62	183	109	125	69	44	34	23	11	-
1999	707	-	7	60	229	122	112	78	41	31	20	7	-
Female													
2001	112	-	3	8	13	19	25	16	14	7	5	1	1
2000	105	1	7	5	19	22	28	9	6	7	1	-	-
1999	117	-	5	16	22	21	24	14	7	2	3	3	-
Hispanic													
Both sexes													
2001	86	-	1	4	36	21	12	6	1	1	2	1	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Accidental discharge of firearms (W32-W34)													
Hispanic													
Both sexes													
2000	59	-	2	6	25	12	11	2	1	-	-	-	-
1999	66	-	-	3	31	17	8	4	1	2	-	-	-
Male													
2001	76	-	1	3	33	16	12	5	1	1	2	1	1
2000	50	-	1	4	24	10	9	2	-	-	-	-	-
1999	61	-	-	3	29	16	7	3	1	2	-	-	-
Female													
2001	10	-	-	1	3	5	-	1	-	-	-	-	-
2000	9	-	1	2	1	2	2	-	1	-	-	-	-
1999	5	-	-	-	2	1	1	1	-	-	-	-	-
Non-Hispanic [1]													
Both sexes													
2001	715	-	14	53	169	101	134	97	68	32	39	7	1
2000	717	1	16	61	177	119	142	76	49	41	24	11	-
1999	758	-	12	73	220	126	128	88	47	31	23	10	-
Male													
2001	613	-	11	46	159	87	109	82	54	25	34	6	-
2000	621	-	10	58	159	99	116	67	44	34	23	11	-
1999	646	-	7	57	200	106	105	75	40	29	20	7	-
Female													
2001	102	-	3	7	10	14	25	15	14	7	5	1	1
2000	96	1	6	3	18	20	26	9	5	7	1	-	-
1999	112	-	5	16	20	20	23	13	7	2	3	3	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Accidental discharge of firearms (W32-W34)													
Non-Hispanic white													
Both sexes													
2001	591	-	9	41	117	82	116	92	61	29	36	7	1
2000	555	1	9	46	117	83	118	65	46	37	22	11	-
1999	585	-	5	54	145	98	108	77	40	28	20	10	-
Male													
2001	506	-	7	38	109	71	96	78	48	22	31	6	-
2000	483	-	5	44	106	66	99	59	41	31	21	11	-
1999	495	-	3	43	131	80	87	65	36	26	17	7	-
Female													
2001	85	-	2	3	8	11	20	14	13	7	5	1	1
2000	72	1	4	2	11	17	19	6	5	6	1	-	-
1999	90	-	2	11	14	18	21	12	4	2	3	3	-
Non-Hispanic black													
Both sexes													
2001	110	-	5	8	48	17	15	5	6	3	3	-	-
2000	137	-	5	12	52	32	19	10	3	3	1	-	-
1999	156	-	7	17	67	23	18	11	7	3	3	-	-
Male													
2001	97	-	4	6	46	14	12	4	5	3	3	-	-
2000	117	-	3	12	46	29	14	7	3	2	1	-	-
1999	135	-	4	12	61	22	16	10	4	3	3	-	-
Female													
2001	13	-	1	2	2	3	3	1	1	-	-	-	-
2000	20	-	2	-	6	3	5	3	-	1	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Accidental discharge of firearms (W32-W34)													
Non-Hispanic black													
Female													
1999	21	-	3	5	6	1	2	1	3	-	-	-	-
Origin not stated [2]													
Both sexes													
2001	1	-	-	-	1	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	1	-	-	-	1	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Accidental drowning and submersion (W65-W74)													
All origins													
Both sexes													
2001	3,281	68	458	333	596	374	462	359	206	159	157	75	34
2000	3,482	75	493	375	646	419	480	354	217	179	156	64	24
1999	3,529	68	490	369	647	446	475	370	219	185	158	68	34
Male													
2001	2,560	30	291	237	544	328	376	299	155	124	102	45	29
2000	2,735	43	316	267	583	364	406	299	179	133	88	34	23

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Accidental drowning and submersion (W65-W74)													
All origins													
Male													
1999	2,794	39	310	275	592	397	402	311	170	129	111	28	30
Female													
2001	721	38	167	96	52	46	86	60	51	35	55	30	5
2000	747	32	177	108	63	55	74	55	38	46	68	30	1
1999	735	29	180	94	55	49	73	59	49	56	47	40	4
Hispanic													
Both sexes													
2001	421	16	78	40	101	74	57	25	7	5	4	2	12
2000	433	9	75	43	147	62	41	20	11	12	4	1	8
1999	480	16	78	46	145	79	45	25	8	7	6	-	25
Male													
2001	348	9	53	28	99	65	48	21	6	3	3	2	11
2000	350	6	44	22	135	60	36	18	8	10	3	-	8
1999	405	12	43	35	140	75	38	23	7	7	4	-	21
Female													
2001	73	7	25	12	2	9	9	4	1	2	1	-	1
2000	83	3	31	21	12	2	5	2	3	2	1	1	-
1999	75	4	35	11	5	4	7	2	1	-	2	-	4
Non-Hispanic [1]													
Both sexes													
2001	2,832	51	380	291	493	300	403	329	198	153	153	73	8
2000	3,018	65	417	331	496	355	437	332	205	166	152	61	1
1999	3,029	52	409	323	500	363	429	343	210	178	152	67	3

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Accidental drowning and submersion (W65-W74)													
Non-Hispanic [1]													
Male													
2001	2,188	21	238	207	443	263	326	275	148	120	99	43	5
2000	2,358	37	272	244	445	302	368	279	170	122	85	33	1
1999	2,370	27	265	240	450	318	363	286	162	122	107	27	3
Female													
2001	644	30	142	84	50	37	77	54	50	33	54	30	3
2000	660	28	145	87	51	53	69	53	35	44	67	28	-
1999	659	25	144	83	50	45	66	57	48	56	45	40	-
Non-Hispanic white													
Both sexes													
2001	2,082	36	321	135	322	202	306	265	158	133	134	66	4
2000	2,235	38	342	176	316	258	344	255	175	144	134	53	-
1999	2,230	30	337	142	317	250	349	291	174	147	132	59	2
Male													
2001	1,577	16	205	87	284	171	245	220	114	105	88	39	3
2000	1,737	22	226	129	278	222	287	212	150	104	78	29	-
1999	1,708	14	217	99	279	216	290	241	133	99	93	25	2
Female													
2001	505	20	116	48	38	31	61	45	44	28	46	27	1
2000	498	16	116	47	38	36	57	43	25	40	56	24	-
1999	522	16	120	43	38	34	59	50	41	48	39	34	-
Non-Hispanic black													
Both sexes													
2001	570	11	47	133	133	62	70	51	32	13	13	4	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Accidental drowning and submersion (W65-W74)													
Non-Hispanic black													
Both sexes													
2000	580	25	54	134	128	58	77	52	17	15	12	7	1
1999	624	21	59	158	137	86	58	36	28	21	15	5	-
Male													
2001	471	3	28	102	126	61	58	42	26	12	9	3	1
2000	470	14	33	101	123	52	67	45	13	13	4	4	1
1999	521	12	40	124	128	78	53	34	22	17	11	2	-
Female													
2001	99	8	19	31	7	1	12	9	6	1	4	1	-
2000	110	11	21	33	5	6	10	7	4	2	8	3	-
1999	103	9	19	34	9	8	5	2	6	4	4	3	-
Origin not stated [2]													
Both sexes													
2001	28	1	-	2	2	-	2	5	1	1	-	-	14
2000	31	1	1	1	3	2	2	2	1	1	-	2	15
1999	20	-	3	-	2	4	1	2	1	-	-	1	6
Male													
2001	24	-	-	2	2	-	2	3	1	1	-	-	13
2000	27	-	-	1	3	2	2	2	1	1	-	1	14
1999	19	-	2	-	2	4	1	2	1	-	-	1	6
Female													
2001	4	1	-	-	-	-	-	2	-	-	-	-	1
2000	4	1	1	-	-	-	-	-	-	-	-	1	1
1999	1	-	1	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Accidental exposure to smoke, fire and flames (X00-X09)													
All origins													
Both sexes													
2001	3,309	50	225	252	209	247	441	419	378	411	423	251	3
2000	3,377	37	290	266	192	241	402	439	369	401	472	263	5
1999	3,348	41	302	262	201	282	421	368	319	416	487	245	4
Male													
2001	1,998	30	142	152	137	161	308	288	236	229	204	108	3
2000	2,025	20	174	145	121	157	268	303	241	232	244	115	5
1999	2,025	21	173	147	133	197	300	254	194	244	247	111	4
Female													
2001	1,311	20	83	100	72	86	133	131	142	182	219	143	-
2000	1,352	17	116	121	71	84	134	136	128	169	228	148	-
1999	1,323	20	129	115	68	85	121	114	125	172	240	134	-
Hispanic													
Both sexes													
2001	176	9	18	23	19	21	28	17	13	9	9	10	-
2000	178	4	21	19	21	19	25	11	17	18	17	6	-
1999	192	5	41	23	24	17	21	16	14	11	15	5	-
Male													
2001	109	5	13	11	10	16	19	11	9	6	5	4	-
2000	105	2	11	10	11	14	17	7	9	10	10	4	-
1999	127	3	23	12	16	12	16	12	11	8	9	5	-
Female													
2001	67	4	5	12	9	5	9	6	4	3	4	6	-
2000	73	2	10	9	10	5	8	4	8	8	7	2	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Accidental exposure to smoke, fire and flames (X00-X09)													
Hispanic													
Female													
1999	65	2	18	11	8	5	5	4	3	3	6	-	-
Non-Hispanic [1]													
Both sexes													
2001	3,106	41	205	226	190	223	410	398	361	398	412	240	2
2000	3,178	31	268	246	169	222	375	426	349	382	451	256	3
1999	3,133	36	260	239	173	264	395	347	302	404	470	240	3
Male													
2001	1,873	25	127	139	127	145	286	274	224	222	199	103	2
2000	1,907	16	163	135	108	143	250	295	230	222	232	110	3
1999	1,879	18	150	135	113	184	279	238	181	235	237	106	3
Female													
2001	1,233	16	78	87	63	78	124	124	137	176	213	137	-
2000	1,271	15	105	111	61	79	125	131	119	160	219	146	-
1999	1,254	18	110	104	60	80	116	109	121	169	233	134	-
Non-Hispanic white													
Both sexes													
2001	2,204	26	124	136	139	153	297	284	261	291	311	180	2
2000	2,194	17	139	139	120	151	267	320	242	278	328	190	3
1999	2,172	21	137	136	125	199	286	253	218	281	343	171	2
Male													
2001	1,322	14	82	82	96	105	207	190	160	157	149	78	2
2000	1,330	7	88	76	79	99	179	222	160	159	172	86	3
1999	1,325	11	80	76	83	145	204	173	136	160	174	81	2

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Accidental exposure to smoke, fire and flames (X00-X09)													
Non-Hispanic white													
Female													
2001	882	12	42	54	43	48	90	94	101	134	162	102	-
2000	864	10	51	63	41	52	88	98	82	119	156	104	-
1999	847	10	57	60	42	54	82	80	82	121	169	90	-
Non-Hispanic black													
Both sexes													
2001	829	14	80	85	42	63	103	105	89	99	92	57	-
2000	915	14	120	101	42	64	97	97	100	98	117	65	-
1999	887	15	110	102	40	52	100	86	80	113	120	68	1
Male													
2001	501	10	44	53	24	34	73	79	56	61	44	23	-
2000	539	9	70	57	25	39	63	67	67	60	58	24	-
1999	507	7	63	58	24	29	69	59	44	69	59	25	1
Female													
2001	328	4	36	32	18	29	30	26	33	38	48	34	-
2000	376	5	50	44	17	25	34	30	33	38	59	41	-
1999	380	8	47	44	16	23	31	27	36	44	61	43	-
Origin not stated [2]													
Both sexes													
2001	27	-	2	3	-	3	3	4	4	4	2	1	1
2000	21	2	1	1	2	-	2	2	3	1	4	1	2
1999	23	-	1	-	4	1	5	5	3	1	2	-	1
Male													
2001	16	-	2	2	-	-	3	3	3	1	-	1	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Accidental exposure to smoke, fire and flames (X00-X09)													
Origin not stated [2]													
Male													
2000	13	2	-	-	2	-	1	1	2	-	2	1	2
1999	19	-	-	-	4	1	5	4	2	1	1	-	1
Female													
2001	11	-	-	1	-	3	-	1	1	3	2	-	-
2000	8	-	1	1	-	-	1	1	1	1	2	-	-
1999	4	-	1	-	-	-	-	1	1	-	1	-	-
Accidental poisoning and exposure to noxious substances (X40-X49)													
All origins													
Both sexes													
2001	14,078	15	31	50	1,362	2,507	5,036	3,547	798	291	279	152	10
2000	12,757	14	32	45	1,160	2,380	4,663	3,061	688	278	245	184	7
1999	12,186	12	34	40	964	2,355	4,549	2,844	668	300	265	143	12
Male													
2001	9,885	8	21	23	1,080	1,862	3,522	2,503	500	170	137	49	10
2000	9,138	11	21	27	920	1,820	3,384	2,223	401	147	102	76	6
1999	8,887	11	17	25	754	1,828	3,349	2,106	448	172	116	53	8
Female													
2001	4,193	7	10	27	282	645	1,514	1,044	298	121	142	103	-
2000	3,619	3	11	18	240	560	1,279	838	287	131	143	108	1
1999	3,299	1	17	15	210	527	1,200	738	220	128	149	90	4

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Accidental poisoning and exposure to noxious substances (X40-X49)													
Hispanic													
Both sexes													
2001	1,324	1	5	10	144	334	475	263	64	19	7	1	1
2000	1,308	2	4	4	132	344	493	251	53	15	5	5	-
1999	1,502	4	3	6	146	383	585	289	58	18	5	2	3
Male													
2001	1,061	1	3	2	111	274	385	218	50	13	3	-	1
2000	1,090	2	2	2	113	295	415	211	35	10	2	3	-
1999	1,271	4	3	4	123	332	493	240	51	15	2	1	3
Female													
2001	263	-	2	8	33	60	90	45	14	6	4	1	-
2000	218	-	2	2	19	49	78	40	18	5	3	2	-
1999	231	-	-	2	23	51	92	49	7	3	3	1	-
Non-Hispanic [1]													
Both sexes													
2001	12,631	14	26	40	1,210	2,157	4,519	3,250	719	270	271	151	4
2000	11,311	12	27	38	1,022	2,018	4,131	2,758	624	259	240	179	3
1999	10,542	8	31	33	814	1,955	3,907	2,516	596	280	258	139	5
Male													
2001	8,720	7	18	21	962	1,574	3,103	2,257	437	155	133	49	4
2000	7,944	9	18	24	802	1,512	2,938	1,975	357	134	100	73	2
1999	7,510	7	14	20	627	1,482	2,817	1,833	387	155	114	51	3
Female													
2001	3,911	7	8	19	248	583	1,416	993	282	115	138	102	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Accidental poisoning and exposure to noxious substances (X40-X49)													
Non-Hispanic [1]													
Female													
2000	3,367	3	9	14	220	506	1,193	783	267	125	140	106	1
1999	3,032	1	17	13	187	473	1,090	683	209	125	144	88	2
Non-Hispanic white													
Both sexes													
2001	10,269	8	16	30	1,078	1,836	3,698	2,468	546	226	229	130	4
2000	9,124	8	15	24	916	1,683	3,331	2,071	482	216	224	152	2
1999	8,311	6	17	21	694	1,581	3,096	1,867	444	224	238	120	3
Male													
2001	7,111	4	12	17	864	1,358	2,579	1,688	304	129	109	43	4
2000	6,413	6	9	16	736	1,272	2,399	1,464	251	104	93	62	1
1999	5,921	5	6	13	536	1,217	2,265	1,337	263	124	105	48	2
Female													
2001	3,158	4	4	13	214	478	1,119	780	242	97	120	87	-
2000	2,711	2	6	8	180	411	932	607	231	112	131	90	1
1999	2,390	1	11	8	158	364	831	530	181	100	133	72	1
Non-Hispanic black													
Both sexes													
2001	2,122	6	8	6	93	273	749	736	157	39	34	21	-
2000	1,985	4	11	8	76	294	732	645	132	41	15	26	1
1999	2,008	1	12	7	93	328	729	613	139	49	18	17	2
Male													
2001	1,452	3	5	2	67	186	477	540	121	25	20	6	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Accidental poisoning and exposure to noxious substances (X40-X49)													
Non-Hispanic black													
Male													
2000	1,400	3	8	6	49	209	496	480	101	29	7	11	1
1999	1,450	1	7	5	76	231	503	471	114	29	9	3	1
Female													
2001	670	3	3	4	26	87	272	196	36	14	14	15	-
2000	585	1	3	2	27	85	236	165	31	12	8	15	-
1999	558	-	5	2	17	97	226	142	25	20	9	14	1
Origin not stated [2]													
Both sexes													
2001	123	-	-	-	8	16	42	34	15	2	1	-	5
2000	138	-	1	3	6	18	39	52	11	4	-	-	4
1999	142	-	-	1	4	17	57	39	14	2	2	2	4
Male													
2001	104	-	-	-	7	14	34	28	13	2	1	-	5
2000	104	-	1	1	5	13	31	37	9	3	-	-	4
1999	106	-	-	1	4	14	39	33	10	2	-	1	2
Female													
2001	19	-	-	-	1	2	8	6	2	-	-	-	-
2000	34	-	-	2	1	5	8	15	2	1	-	-	-
1999	36	-	-	-	-	3	18	6	4	-	2	1	2

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)													
All origins													
Both sexes													
2001	17,760	669	275	255	643	795	1,386	1,641	1,342	1,826	3,841	5,063	24
2000	17,437	572	266	267	648	895	1,446	1,510	1,349	1,824	3,678	4,966	16
1999	18,388	523	293	316	662	887	1,452	1,496	1,332	2,032	4,025	5,355	15
Male													
2001	10,044	390	162	181	535	682	1,128	1,261	933	1,097	1,913	1,741	21
2000	9,960	327	162	192	560	767	1,205	1,219	971	1,081	1,801	1,659	16
1999	10,371	314	183	221	565	748	1,181	1,176	951	1,227	1,969	1,822	14
Female													
2001	7,716	279	113	74	108	113	258	380	409	729	1,928	3,322	3
2000	7,477	245	104	75	88	128	241	291	378	743	1,877	3,307	-
1999	8,017	209	110	95	97	139	271	320	381	805	2,056	3,533	1
Hispanic													
Both sexes													
2001	985	78	43	23	98	122	146	122	70	88	89	102	4
2000	942	58	41	29	108	140	151	84	68	67	84	106	6
1999	878	65	53	31	98	101	120	82	69	90	72	94	3
Male													
2001	704	40	28	12	93	110	130	102	49	54	46	36	4
2000	695	31	23	14	105	129	139	70	53	41	40	44	6
1999	620	35	37	19	88	90	101	70	48	60	30	39	3
Female													
2001	281	38	15	11	5	12	16	20	21	34	43	66	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)													
Hispanic													
Female													
2000	247	27	18	15	3	11	12	14	15	26	44	62	-
1999	258	30	16	12	10	11	19	12	21	30	42	55	-
Non-Hispanic [1]													
Both sexes													
2001	16,695	588	231	230	542	669	1,231	1,513	1,262	1,732	3,741	4,950	6
2000	16,408	513	224	238	538	748	1,283	1,413	1,275	1,748	3,580	4,846	2
1999	17,430	457	239	284	562	781	1,331	1,400	1,250	1,935	3,943	5,245	3
Male													
2001	9,280	347	133	167	439	568	990	1,154	876	1,040	1,858	1,702	6
2000	9,207	295	139	178	454	631	1,057	1,138	913	1,033	1,755	1,612	2
1999	9,694	279	146	201	476	655	1,079	1,094	891	1,162	1,933	1,775	3
Female													
2001	7,415	241	98	63	103	101	241	359	386	692	1,883	3,248	-
2000	7,201	218	85	60	84	117	226	275	362	715	1,825	3,234	-
1999	7,736	178	93	83	86	126	252	306	359	773	2,010	3,470	-
Non-Hispanic white													
Both sexes													
2001	14,326	348	160	182	455	528	994	1,216	1,033	1,462	3,359	4,583	6
2000	14,083	310	156	185	425	604	1,039	1,123	1,044	1,464	3,224	4,509	-
1999	14,930	276	163	225	468	622	1,103	1,100	980	1,612	3,504	4,876	1
Male													
2001	7,832	206	93	129	370	449	814	929	719	886	1,658	1,573	6

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)													
Non-Hispanic white													
Male													
2000	7,703	186	99	141	361	515	860	899	736	868	1,555	1,483	-
1999	8,122	167	96	163	395	532	896	860	705	962	1,717	1,628	1
Female													
2001	6,494	142	67	53	85	79	180	287	314	576	1,701	3,010	-
2000	6,380	124	57	44	64	89	179	224	308	596	1,669	3,026	-
1999	6,808	109	67	62	73	90	207	240	275	650	1,787	3,248	-
Non-Hispanic black													
Both sexes													
2001	1,968	217	61	38	62	101	184	249	198	232	317	309	-
2000	1,968	183	55	48	90	118	198	237	200	244	307	286	2
1999	2,183	168	61	47	74	132	193	263	236	293	384	330	2
Male													
2001	1,180	131	35	31	47	81	137	185	134	134	165	100	-
2000	1,256	96	32	34	73	94	159	196	151	139	174	106	2
1999	1,365	101	42	32	67	102	153	203	165	182	185	131	2
Female													
2001	788	86	26	7	15	20	47	64	64	98	152	209	-
2000	712	87	23	14	17	24	39	41	49	105	133	180	-
1999	818	67	19	15	7	30	40	60	71	111	199	199	-
Origin not stated [2]													
Both sexes													
2001	80	3	1	2	3	4	9	6	10	6	11	11	14

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)													
Origin not stated [2]													
Both sexes													
2000	87	1	1	-	2	7	12	13	6	9	14	14	8
1999	80	1	1	1	2	5	1	14	13	7	10	16	9
Male													
2001	60	3	1	2	3	4	8	5	8	3	9	3	11
2000	58	1	-	-	1	7	9	11	5	7	6	3	8
1999	57	-	-	1	1	3	1	12	12	5	6	8	8
Female													
2001	20	-	-	-	-	-	1	1	2	3	2	8	3
2000	29	-	1	-	1	-	3	2	1	2	8	11	-
1999	23	1	1	-	1	2	-	2	1	2	4	8	1
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)													
All origins													
Both sexes													
2001 [3]	30,622	. . .	. . .	279	3,971	5,070	6,635	5,942	3,317	2,432	2,192	769	15
2000	29,350	. . .	. . .	307	3,994	4,792	6,562	5,437	2,945	2,292	2,181	833	7
1999	29,199	. . .	. . .	244	3,901	5,106	6,466	5,081	2,896	2,471	2,217	801	16
Male													
2001 [3]	24,672	. . .	. . .	214	3,409	4,199	5,180	4,504	2,563	2,040	1,885	664	14
2000	23,618	. . .	. . .	244	3,424	3,938	5,114	4,149	2,265	1,888	1,884	705	7
1999	23,458	. . .	. . .	194	3,326	4,194	5,019	3,854	2,255	2,051	1,868	681	16
Female													
2001 [3]	5,950	. . .	. . .	65	562	871	1,455	1,438	754	392	307	105	1
2000	5,732	. . .	. . .	63	570	854	1,448	1,288	680	404	297	128	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)													
All origins													
Female													
1999	5,741	. . .	. . .	50	575	912	1,447	1,227	641	420	349	120	-
Hispanic													
Both sexes													
2001 [3]	1,850	. . .	. . .	37	414	490	397	239	114	86	49	22	2
2000	1,787	. . .	. . .	29	447	420	385	244	102	84	59	16	1
1999	1,695	. . .	. . .	30	391	423	369	210	112	73	63	22	2
Male													
2001 [3]	1,576	. . .	. . .	26	343	419	350	192	100	77	45	22	2
2000	1,525	. . .	. . .	22	387	370	320	194	89	74	53	15	1
1999	1,430	. . .	. . .	21	338	364	307	169	94	63	54	18	2
Female													
2001 [3]	274	. . .	. . .	11	71	71	47	47	14	9	4	-	-
2000	262	. . .	. . .	7	60	50	65	50	13	10	6	1	-
1999	265	. . .	. . .	9	53	59	62	41	18	10	9	4	-
Non-Hispanic [1]													
Both sexes													
2001 [3]	28,647	. . .	. . .	242	3,545	4,560	6,213	5,682	3,184	2,334	2,138	744	5
2000	27,426	. . .	. . .	276	3,529	4,350	6,145	5,165	2,827	2,198	2,119	813	4
1999	27,379	. . .	. . .	212	3,500	4,665	6,068	4,842	2,773	2,388	2,146	776	9
Male													
2001 [3]	22,983	. . .	. . .	188	3,055	3,762	4,809	4,292	2,446	1,952	1,835	639	5
2000	21,981	. . .	. . .	221	3,023	3,553	4,767	3,931	2,161	1,805	1,828	688	4
1999	21,929	. . .	. . .	171	2,983	3,816	4,689	3,663	2,152	1,978	1,807	661	9
Female													
2001 [3]	5,664	. . .	. . .	54	490	798	1,404	1,390	738	382	303	105	-
2000	5,445	. . .	. . .	55	506	797	1,378	1,234	666	393	291	125	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)													
Non-Hispanic [1]													
Female													
1999	5,450	. . .	. . .	41	517	849	1,379	1,179	621	410	339	115	-
Non-Hispanic white													
Both sexes													
2001 [3]	25,813	. . .	. . .	183	2,926	3,871	5,604	5,292	2,979	2,187	2,049	718	4
2000	24,616	. . .	. . .	225	2,868	3,658	5,540	4,774	2,659	2,075	2,028	786	3
1999	24,534	. . .	. . .	174	2,830	3,927	5,469	4,487	2,580	2,249	2,058	754	6
Male													
2001 [3]	20,710	. . .	. . .	144	2,522	3,190	4,329	4,016	2,284	1,836	1,763	622	4
2000	19,711	. . .	. . .	179	2,467	2,989	4,287	3,627	2,031	1,701	1,761	666	3
1999	19,620	. . .	. . .	141	2,416	3,205	4,210	3,388	2,001	1,869	1,740	644	6
Female													
2001 [3]	5,103	. . .	. . .	39	404	681	1,275	1,276	695	351	286	96	-
2000	4,905	. . .	. . .	46	401	669	1,253	1,147	628	374	267	120	-
1999	4,914	. . .	. . .	33	414	722	1,259	1,099	579	380	318	110	-
Non-Hispanic black													
Both sexes													
2001 [3]	1,904	. . .	. . .	40	425	472	418	268	126	90	52	13	-
2000	1,923	. . .	. . .	39	471	469	427	266	101	85	53	12	-
1999	1,924	. . .	. . .	29	463	497	430	226	116	94	52	15	2
Male													
2001 [3]	1,578	. . .	. . .	31	388	406	329	196	102	73	44	9	-
2000	1,601	. . .	. . .	34	407	397	346	210	80	76	40	11	-
1999	1,630	. . .	. . .	22	408	432	353	183	95	78	45	12	2
Female													
2001 [3]	326	. . .	. . .	9	37	66	89	72	24	17	8	4	-
2000	322	. . .	. . .	5	64	72	81	56	21	9	13	1	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)													
Non-Hispanic black													
Female													
1999	294	. . .	. . .	7	55	65	77	43	21	16	7	3	-
Origin not stated [2]													
Both sexes													
2001 [3]	125	. . .	. . .	-	12	20	25	21	19	12	5	3	8
2000	137	. . .	. . .	2	18	22	32	28	16	10	3	4	2
1999	125	. . .	. . .	2	10	18	29	29	11	10	8	3	5
Male													
2001 [3]	113	. . .	. . .	-	11	18	21	20	17	11	5	3	7
2000	112	. . .	. . .	1	14	15	27	24	15	9	3	2	2
1999	99	. . .	. . .	2	5	14	23	22	9	10	7	2	5
Female													
2001 [3]	12	. . .	. . .	-	1	2	4	1	2	1	-	-	1
2000	25	. . .	. . .	1	4	7	5	4	1	1	-	2	-
1999	26	. . .	. . .	-	5	4	6	7	2	-	1	1	-
Intentional self-harm (suicide) by discharge of firearms (X72-X74)													
All origins													
Both sexes													
2001	16,869	. . .	. . .	90	2,130	2,564	3,030	3,023	2,083	1,758	1,648	537	6
2000	16,586	. . .	. . .	110	2,267	2,453	3,191	2,858	1,833	1,700	1,611	558	5
1999	16,599	. . .	. . .	103	2,315	2,621	3,097	2,677	1,860	1,791	1,606	524	5
Male													
2001	14,758	. . .	. . .	69	1,935	2,246	2,551	2,551	1,771	1,592	1,531	506	6
2000	14,454	. . .	. . .	90	2,043	2,123	2,657	2,409	1,568	1,528	1,505	526	5
1999	14,479	. . .	. . .	80	2,083	2,270	2,586	2,251	1,594	1,615	1,499	496	5

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Intentional self-harm (suicide) by discharge of firearms (X72-X74)													
All origins													
Female													
2001	2,111	. . .	. . .	21	195	318	479	472	312	166	117	31	-
2000	2,132	. . .	. . .	20	224	330	534	449	265	172	106	32	-
1999	2,120	. . .	. . .	23	232	351	511	426	266	176	107	28	-
Hispanic													
Both sexes													
2001	798	. . .	. . .	5	193	224	135	102	59	45	26	8	1
2000	813	. . .	. . .	7	224	181	144	124	47	49	27	9	1
1999	794	. . .	. . .	13	219	182	148	96	48	37	39	11	1
Male													
2001	731	. . .	. . .	4	172	203	126	90	57	44	26	8	1
2000	737	. . .	. . .	7	205	165	126	106	46	46	26	9	1
1999	705	. . .	. . .	7	196	164	127	81	45	35	38	11	1
Female													
2001	67	. . .	. . .	1	21	21	9	12	2	1	-	-	-
2000	76	. . .	. . .	-	19	16	18	18	1	3	1	-	-
1999	89	. . .	. . .	6	23	18	21	15	3	2	1	-	-
Non-Hispanic [1]													
Both sexes													
2001	16,020	. . .	. . .	85	1,933	2,335	2,887	2,916	2,011	1,706	1,617	526	4
2000	15,717	. . .	. . .	103	2,035	2,264	3,036	2,722	1,779	1,645	1,583	547	3
1999	15,755	. . .	. . .	90	2,092	2,431	2,937	2,573	1,806	1,749	1,563	511	3
Male													
2001	13,978	. . .	. . .	65	1,759	2,039	2,417	2,457	1,701	1,541	1,500	495	4

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Intentional self-harm (suicide) by discharge of firearms (X72-X74)													
Non-Hispanic [1]													
Male													
2000	13,664	. . .	. . .	83	1,831	1,951	2,520	2,291	1,515	1,476	1,478	516	3
1999	13,732	. . .	. . .	73	1,884	2,101	2,448	2,164	1,543	1,575	1,457	484	3
Female													
2001	2,042	. . .	. . .	20	174	296	470	459	310	165	117	31	-
2000	2,053	. . .	. . .	20	204	313	516	431	264	169	105	31	-
1999	2,023	. . .	. . .	17	208	330	489	409	263	174	106	27	-
Non-Hispanic white													
Both sexes													
2001	14,648	. . .	. . .	69	1,565	2,001	2,629	2,760	1,919	1,628	1,560	514	3
2000	14,333	. . .	. . .	96	1,655	1,916	2,780	2,552	1,679	1,572	1,545	536	2
1999	14,316	. . .	. . .	81	1,684	2,071	2,676	2,413	1,703	1,672	1,514	499	3
Male													
2001	12,771	. . .	. . .	55	1,414	1,739	2,199	2,333	1,619	1,474	1,451	484	3
2000	12,448	. . .	. . .	79	1,489	1,648	2,296	2,151	1,427	1,405	1,446	505	2
1999	12,441	. . .	. . .	65	1,514	1,776	2,213	2,029	1,451	1,505	1,412	473	3
Female													
2001	1,877	. . .	. . .	14	151	262	430	427	300	154	109	30	-
2000	1,885	. . .	. . .	17	166	268	484	401	252	167	99	31	-
1999	1,875	. . .	. . .	16	170	295	463	384	252	167	102	26	-
Non-Hispanic black													
Both sexes													
2001	1,069	. . .	. . .	12	293	258	198	122	67	62	47	10	-
2000	1,073	. . .	. . .	4	304	268	193	131	71	62	32	8	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Intentional self-harm (suicide) by discharge of firearms (X72-X74)													
Non-Hispanic black													
Both sexes													
1999	1,112	. . .	. . .	5	323	268	197	118	81	65	43	12	-
Male													
2001	949	. . .	. . .	9	277	233	169	98	61	53	40	9	-
2000	949	. . .	. . .	4	277	237	168	107	61	60	27	8	-
1999	1,007	. . .	. . .	4	294	247	178	102	73	59	39	11	-
Female													
2001	120	. . .	. . .	3	16	25	29	24	6	9	7	1	-
2000	124	. . .	. . .	-	27	31	25	24	10	2	5	-	-
1999	105	. . .	. . .	1	29	21	19	16	8	6	4	1	-
Origin not stated [2]													
Both sexes													
2001	51	. . .	. . .	-	4	5	8	5	13	7	5	3	1
2000	56	. . .	. . .	-	8	8	11	12	7	6	1	2	1
1999	50	. . .	. . .	-	4	8	12	8	6	5	4	2	1
Male													
2001	49	. . .	. . .	-	4	4	8	4	13	7	5	3	1
2000	53	. . .	. . .	-	7	7	11	12	7	6	1	1	1
1999	42	. . .	. . .	-	3	5	11	6	6	5	4	1	1
Female													
2001	2	. . .	. . .	-	-	1	-	1	-	-	-	-	-
2000	3	. . .	. . .	-	1	1	-	-	-	-	-	1	-
1999	8	. . .	. . .	-	1	3	1	2	-	-	-	1	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84, Y87.0)													
All origins													
Both sexes													
2001 [3]	13,753	. . .	. . .	189	1,841	2,506	3,605	2,919	1,234	674	544	232	9
2000	12,764	. . .	. . .	197	1,727	2,339	3,371	2,579	1,112	592	570	275	2
1999	12,600	. . .	. . .	141	1,586	2,485	3,369	2,404	1,036	680	611	277	11
Male													
2001 [3]	9,914	. . .	. . .	145	1,474	1,953	2,629	1,953	792	448	354	158	8
2000	9,164	. . .	. . .	154	1,381	1,815	2,457	1,740	697	360	379	179	2
1999	8,979	. . .	. . .	114	1,243	1,924	2,433	1,603	661	436	369	185	11
Female													
2001 [3]	3,839	. . .	. . .	44	367	553	976	966	442	226	190	74	1
2000	3,600	. . .	. . .	43	346	524	914	839	415	232	191	96	-
1999	3,621	. . .	. . .	27	343	561	936	801	375	244	242	92	-
Hispanic													
Both sexes													
2001 [3]	1,052	. . .	. . .	32	221	266	262	137	55	41	23	14	1
2000	974	. . .	. . .	22	223	239	241	120	55	35	32	7	-
1999	901	. . .	. . .	17	172	241	221	114	64	36	24	11	1
Male													
2001 [3]	845	. . .	. . .	22	171	216	224	102	43	33	19	14	1
2000	788	. . .	. . .	15	182	205	194	88	43	28	27	6	-
1999	725	. . .	. . .	14	142	200	180	88	49	28	16	7	1
Female													
2001 [3]	207	. . .	. . .	10	50	50	38	35	12	8	4	-	-
2000	186	. . .	. . .	7	41	34	47	32	12	7	5	1	-
1999	176	. . .	. . .	3	30	41	41	26	15	8	8	4	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84, Y87.0)													
Non-Hispanic [1]													
Both sexes													
2001 [3]	12,627	. . .	. . .	157	1,612	2,225	3,326	2,766	1,173	628	521	218	1
2000	11,709	. . .	. . .	173	1,494	2,086	3,109	2,443	1,048	553	536	266	1
1999	11,624	. . .	. . .	122	1,408	2,234	3,131	2,269	967	639	583	265	6
Male													
2001 [3]	9,005	. . .	. . .	123	1,296	1,723	2,392	1,835	745	411	335	144	1
2000	8,317	. . .	. . .	138	1,192	1,602	2,247	1,640	646	329	350	172	1
1999	8,197	. . .	. . .	98	1,099	1,715	2,241	1,499	609	403	350	177	6
Female													
2001 [3]	3,622	. . .	. . .	34	316	502	934	931	428	217	186	74	-
2000	3,392	. . .	. . .	35	302	484	862	803	402	224	186	94	-
1999	3,427	. . .	. . .	24	309	519	890	770	358	236	233	88	-
Non-Hispanic white													
Both sexes													
2001 [3]	11,165	. . .	. . .	114	1,361	1,870	2,975	2,532	1,060	559	489	204	1
2000	10,283	. . .	. . .	129	1,213	1,742	2,760	2,222	980	503	483	250	1
1999	10,218	. . .	. . .	93	1,146	1,856	2,793	2,074	877	577	544	255	3
Male													
2001 [3]	7,939	. . .	. . .	89	1,108	1,451	2,130	1,683	665	362	312	138	1
2000	7,263	. . .	. . .	100	978	1,341	1,991	1,476	604	296	315	161	1
1999	7,179	. . .	. . .	76	902	1,429	1,997	1,359	550	364	328	171	3
Female													
2001 [3]	3,226	. . .	. . .	25	253	419	845	849	395	197	177	66	-
2000	3,020	. . .	. . .	29	235	401	769	746	376	207	168	89	-
1999	3,039	. . .	. . .	17	244	427	796	715	327	213	216	84	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84, Y87.0)													
Non-Hispanic black													
Both sexes													
2001 [3]	835	. . .	. . .	28	132	214	220	146	59	28	5	3	-
2000	850	. . .	. . .	35	167	201	234	135	30	23	21	4	-
1999	812	. . .	. . .	24	140	229	233	108	35	29	9	3	2
Male													
2001 [3]	629	. . .	. . .	22	111	173	160	98	41	20	4	-	-
2000	652	. . .	. . .	30	130	160	178	103	19	16	13	3	-
1999	623	. . .	. . .	18	114	185	175	81	22	19	6	1	2
Female													
2001 [3]	206	. . .	. . .	6	21	41	60	48	18	8	1	3	-
2000	198	. . .	. . .	5	37	41	56	32	11	7	8	1	-
1999	189	. . .	. . .	6	26	44	58	27	13	10	3	2	-
Origin not stated [2]													
Both sexes													
2001 [3]	74	. . .	. . .	-	8	15	17	16	6	5	-	-	7
2000	81	. . .	. . .	2	10	14	21	16	9	4	2	2	1
1999	75	. . .	. . .	2	6	10	17	21	5	5	4	1	4
Male													
2001 [3]	64	. . .	. . .	-	7	14	13	16	4	4	-	-	6
2000	59	. . .	. . .	1	7	8	16	12	8	3	2	1	1
1999	57	. . .	. . .	2	2	9	12	16	3	5	3	1	4
Female													
2001 [3]	10	. . .	. . .	-	1	1	4	-	2	1	-	-	1
2000	22	. . .	. . .	1	3	6	5	4	1	1	-	1	-
1999	18	. . .	. . .	-	4	1	5	5	2	-	1	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)													
All origins													
Both sexes													
2001 [3]	20,308	332	415	326	5,297	5,204	4,268	2,467	1,018	532	312	105	32
2000	16,765	349	356	371	4,939	4,164	3,219	1,755	738	443	301	100	30
1999	16,889	331	376	432	4,998	4,231	3,206	1,668	722	473	304	99	49
Male													
2001 [3]	15,555	197	238	193	4,541	4,157	3,122	1,832	723	332	155	42	23
2000	12,820	203	198	227	4,203	3,361	2,319	1,284	537	279	154	40	15
1999	12,785	187	194	250	4,191	3,346	2,307	1,229	541	304	162	41	33
Female													
2001 [3]	4,753	135	177	133	756	1,047	1,146	635	295	200	157	63	9
2000	3,945	146	158	144	736	803	900	471	201	164	147	60	15
1999	4,104	144	182	182	807	885	899	439	181	169	142	58	16
Hispanic													
Both sexes													
2001 [3]	3,331	55	69	56	1,222	960	567	248	82	45	16	3	8
2000	2,917	54	57	76	1,126	810	447	211	77	36	16	4	3
1999	2,864	53	52	89	1,119	801	432	197	69	33	11	3	5
Male													
2001 [3]	2,756	33	38	38	1,099	800	451	190	59	33	8	2	5
2000	2,431	26	30	56	1,014	697	359	158	57	22	8	2	2
1999	2,399	26	25	59	990	698	354	150	56	26	8	2	5
Female													
2001 [3]	575	22	31	18	123	160	116	58	23	12	8	1	3
2000	486	28	27	20	112	113	88	53	20	14	8	2	1
1999	465	27	27	30	129	103	78	47	13	7	3	1	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)													
Non-Hispanic [1]													
Both sexes													
2001 [3]	16,726	270	343	269	4,028	4,178	3,652	2,182	916	480	296	101	11
2000	13,717	291	299	293	3,776	3,337	2,749	1,526	652	402	285	95	12
1999	13,873	270	323	341	3,860	3,402	2,743	1,452	642	437	292	95	16
Male													
2001 [3]	12,608	161	199	154	3,401	3,309	2,632	1,612	651	294	147	39	9
2000	10,290	174	168	170	3,159	2,650	1,939	1,115	473	254	146	37	5
1999	10,275	158	168	191	3,188	2,624	1,930	1,063	476	275	153	39	10
Female													
2001 [3]	4,118	109	144	115	627	869	1,020	570	265	186	149	62	2
2000	3,427	117	131	123	617	687	810	411	179	148	139	58	7
1999	3,598	112	155	150	672	778	813	389	166	162	139	56	6
Non-Hispanic white													
Both sexes													
2001 [3]	7,928	139	160	164	1,058	1,642	2,081	1,373	652	342	225	85	7
2000	5,429	141	144	164	874	1,023	1,273	834	415	264	211	78	8
1999	5,748	143	156	176	962	1,135	1,327	814	418	319	214	76	8
Male													
2001 [3]	5,453	84	95	83	729	1,177	1,498	990	454	197	110	31	5
2000	3,510	96	86	93	588	677	836	560	288	153	101	28	4
1999	3,747	94	83	93	644	753	881	579	293	188	109	27	3
Female													
2001 [3]	2,475	55	65	81	329	465	583	383	198	145	115	54	2
2000	1,919	45	58	71	286	346	437	274	127	111	110	50	4
1999	2,001	49	73	83	318	382	446	235	125	131	105	49	5

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)													
Non-Hispanic black													
Both sexes													
2001 [3]	8,077	119	166	92	2,817	2,335	1,427	704	219	117	63	14	4
2000	7,748	139	143	122	2,771	2,193	1,359	615	201	123	64	14	4
1999	7,536	114	151	145	2,752	2,122	1,305	557	194	103	67	18	8
Male													
2001 [3]	6,658	68	100	66	2,551	1,998	1,036	544	165	85	34	7	4
2000	6,382	73	73	70	2,459	1,883	1,027	498	159	93	39	7	1
1999	6,131	58	74	89	2,433	1,771	980	431	159	79	39	11	7
Female													
2001 [3]	1,419	51	66	26	266	337	391	160	54	32	29	7	-
2000	1,366	66	70	52	312	310	332	117	42	30	25	7	3
1999	1,405	56	77	56	319	351	325	126	35	24	28	7	1
Origin not stated [2]													
Both sexes													
2001 [3]	251	7	3	1	47	66	49	37	20	7	-	1	13
2000	131	4	-	2	37	17	23	18	9	5	-	1	15
1999	152	8	1	2	19	28	31	19	11	3	1	1	28
Male													
2001 [3]	191	3	1	1	41	48	39	30	13	5	-	1	9
2000	99	3	-	1	30	14	21	11	7	3	-	1	8
1999	111	3	1	-	13	24	23	16	9	3	1	-	18
Female													
2001 [3]	60	4	2	-	6	18	10	7	7	2	-	-	4
2000	32	1	-	1	7	3	2	7	2	2	-	-	7
1999	41	5	-	2	6	4	8	3	2	-	-	1	10

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)													
All origins													
Both sexes													
2001	11,348	11	55	180	4,200	3,308	1,978	934	364	192	99	16	11
2000	10,801	12	28	187	3,963	3,074	1,906	959	367	184	81	28	12
1999	10,828	8	50	224	4,038	3,048	1,868	893	373	193	95	23	15
Male													
2001	9,532	7	27	125	3,796	2,864	1,528	726	264	123	54	7	11
2000	9,006	9	17	138	3,548	2,650	1,487	730	266	108	37	9	7
1999	8,944	5	26	155	3,569	2,609	1,414	676	294	124	48	10	14
Female													
2001	1,816	4	28	55	404	444	450	208	100	69	45	9	-
2000	1,795	3	11	49	415	424	419	229	101	76	44	19	5
1999	1,884	3	24	69	469	439	454	217	79	69	47	13	1
Hispanic													
Both sexes													
2001	2,123	4	12	30	974	654	306	102	20	14	3	1	3
2000	1,958	1	5	41	904	581	249	126	34	14	1	-	2
1999	1,939	2	5	52	908	550	258	112	35	9	5	1	2
Male													
2001	1,892	3	5	23	898	596	257	81	15	10	-	1	3
2000	1,738	-	3	34	837	524	207	99	26	7	-	-	1
1999	1,709	2	3	39	824	499	218	84	27	6	4	1	2
Female													
2001	231	1	7	7	76	58	49	21	5	4	3	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)													
Hispanic													
Female													
2000	220	1	2	7	67	57	42	27	8	7	1	-	1
1999	230	-	2	13	84	51	40	28	8	3	1	-	-
Non-Hispanic [1]													
Both sexes													
2001	9,134	7	43	149	3,186	2,631	1,662	827	341	175	96	14	3
2000	8,767	11	23	144	3,032	2,480	1,640	828	330	168	80	28	3
1999	8,821	6	45	172	3,115	2,476	1,597	776	335	184	89	22	4
Male													
2001	7,559	4	22	101	2,863	2,246	1,264	641	246	110	54	5	3
2000	7,205	9	14	103	2,686	2,115	1,264	628	237	101	37	9	2
1999	7,177	3	23	116	2,734	2,091	1,184	588	264	118	43	9	4
Female													
2001	1,575	3	21	48	323	385	398	186	95	65	42	9	-
2000	1,562	2	9	41	346	365	376	200	93	67	43	19	1
1999	1,644	3	22	56	381	385	413	188	71	66	46	13	-
Non-Hispanic white													
Both sexes													
2001	3,085	4	21	90	618	681	759	460	235	126	75	14	2
2000	2,861	4	12	72	544	624	729	445	224	118	63	24	2
1999	2,995	2	21	80	591	694	749	416	215	140	67	20	-
Male													
2001	2,172	2	12	56	468	495	536	335	155	69	37	5	2

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)													
Non-Hispanic white													
Male													
2000	1,967	2	6	47	401	444	510	311	149	63	25	8	1
1999	2,119	2	11	50	443	515	514	308	157	84	27	8	-
Female													
2001	913	2	9	34	150	186	223	125	80	57	38	9	-
2000	894	2	6	25	143	180	219	134	75	55	38	16	1
1999	876	-	10	30	148	179	235	108	58	56	40	12	-
Non-Hispanic black													
Both sexes													
2001	5,790	2	18	56	2,490	1,874	864	332	92	43	18	-	1
2000	5,622	7	8	68	2,396	1,790	854	347	85	45	17	4	1
1999	5,508	2	18	81	2,426	1,695	796	320	104	39	21	2	4
Male													
2001	5,190	1	9	43	2,329	1,697	700	280	78	37	15	-	1
2000	5,011	7	5	52	2,206	1,616	715	290	69	37	12	1	1
1999	4,826	1	7	60	2,208	1,512	636	256	94	31	16	1	4
Female													
2001	600	1	9	13	161	177	164	52	14	6	3	-	-
2000	611	-	3	16	190	174	139	57	16	8	5	3	-
1999	682	1	11	21	218	183	160	64	10	8	5	1	-
Origin not stated [2]													
Both sexes													
2001	91	-	-	1	40	23	10	5	3	3	-	1	5

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)													
Origin not stated [2]													
Both sexes													
2000	76	-	-	2	27	13	17	5	3	2	-	-	7
1999	68	-	-	-	15	22	13	5	3	-	1	-	9
Male													
2001	81	-	-	1	35	22	7	4	3	3	-	1	5
2000	63	-	-	1	25	11	16	3	3	-	-	-	4
1999	58	-	-	-	11	19	12	4	3	-	1	-	8
Female													
2001	10	-	-	-	5	1	3	1	-	-	-	-	-
2000	13	-	-	1	2	2	1	2	-	2	-	-	3
1999	10	-	-	-	4	3	1	1	-	-	-	-	1
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)													
All origins													
Both sexes													
2001 [3]	8,960	321	360	146	1,097	1,896	2,290	1,533	654	340	213	89	21
2000	5,964	337	328	184	976	1,090	1,313	796	371	259	220	72	18
1999	6,061	323	326	208	960	1,183	1,338	775	349	280	209	76	34
Male													
2001 [3]	6,023	190	211	68	745	1,293	1,594	1,106	459	209	101	35	12
2000	3,814	194	181	89	655	711	832	554	271	171	117	31	8
1999	3,841	182	168	95	622	737	893	553	247	180	114	31	19
Female													
2001 [3]	2,937	131	149	78	352	603	696	427	195	131	112	54	9
2000	2,150	143	147	95	321	379	481	242	100	88	103	41	10

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)													
All origins													
Female													
1999	2,220	141	158	113	338	446	445	222	102	100	95	45	15
Hispanic													
Both sexes													
2001 [3]	1,208	51	57	26	248	306	261	146	62	31	13	2	5
2000	959	53	52	35	222	229	198	85	43	22	15	4	1
1999	925	51	47	37	211	251	174	85	34	24	6	2	3
Male													
2001 [3]	864	30	33	15	201	204	194	109	44	23	8	1	2
2000	693	26	27	22	177	173	152	59	31	15	8	2	1
1999	690	24	22	20	166	199	136	66	29	20	4	1	3
Female													
2001 [3]	344	21	24	11	47	102	67	37	18	8	5	1	3
2000	266	27	25	13	45	56	46	26	12	7	7	2	-
1999	235	27	25	17	45	52	38	19	5	4	2	1	-
Non-Hispanic [1]													
Both sexes													
2001 [3]	7,592	263	300	120	842	1,547	1,990	1,355	575	305	200	87	8
2000	4,950	280	276	149	744	857	1,109	698	322	234	205	67	9
1999	5,052	264	278	169	745	926	1,146	676	307	253	203	73	12
Male													
2001 [3]	5,049	157	177	53	538	1,063	1,368	971	405	184	93	34	6
2000	3,085	165	154	67	473	535	675	487	236	153	109	28	3
1999	3,098	155	145	75	454	533	746	475	212	157	110	30	6
Female													
2001 [3]	2,543	106	123	67	304	484	622	384	170	121	107	53	2

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)													
Non-Hispanic [1]													
Female													
2000	1,865	115	122	82	271	322	434	211	86	81	96	39	6
1999	1,954	109	133	94	291	393	400	201	95	96	93	43	6
Non-Hispanic white													
Both sexes													
2001 [3]	4,843	135	139	74	440	961	1,322	913	417	216	150	71	5
2000	2,568	137	132	92	330	399	544	389	191	146	148	54	6
1999	2,753	141	135	96	371	441	578	398	203	179	147	56	8
Male													
2001 [3]	3,281	82	83	27	261	682	962	655	299	128	73	26	3
2000	1,543	94	80	46	187	233	326	249	139	90	76	20	3
1999	1,628	92	72	43	201	238	367	271	136	104	82	19	3
Female													
2001 [3]	1,562	53	56	47	179	279	360	258	118	88	77	45	2
2000	1,025	43	52	46	143	166	218	140	52	56	72	34	3
1999	1,125	49	63	53	170	203	211	127	67	75	65	37	5
Non-Hispanic black													
Both sexes													
2001 [3]	2,287	117	148	36	327	461	563	372	127	74	45	14	3
2000	2,126	132	135	54	375	403	505	268	116	78	47	10	3
1999	2,028	112	133	64	326	427	509	237	90	64	46	16	4
Male													
2001 [3]	1,468	67	91	23	222	301	336	264	87	48	19	7	3
2000	1,371	66	68	18	253	267	312	208	90	56	27	6	-
1999	1,305	57	67	29	225	259	344	175	65	48	23	10	3

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)													
Non-Hispanic black													
Female													
2001 [3]	819	50	57	13	105	160	227	108	40	26	26	7	-
2000	755	66	67	36	122	136	193	60	26	22	20	4	3
1999	723	55	66	35	101	168	165	62	25	16	23	6	1
Origin not stated [2]													
Both sexes													
2001 [3]	160	7	3	-	7	43	39	32	17	4	-	-	8
2000	55	4	-	-	10	4	6	13	6	3	-	1	8
1999	84	8	1	2	4	6	18	14	8	3	-	1	19
Male													
2001 [3]	110	3	1	-	6	26	32	26	10	2	-	-	4
2000	36	3	-	-	5	3	5	8	4	3	-	1	4
1999	53	3	1	-	2	5	11	12	6	3	-	-	10
Female													
2001 [3]	50	4	2	-	1	17	7	6	7	2	-	-	4
2000	19	1	-	-	5	1	1	5	2	-	-	-	4
1999	31	5	-	2	2	1	7	2	2	-	-	1	9
Legal intervention (Y35,Y89.0)													
All origins													
Both sexes													
2001	396	-	-	-	88	107	115	51	24	6	4	1	-
2000	359	-	-	3	73	110	101	55	10	5	1	1	-
1999	398	-	-	1	67	136	121	50	20	2	1	-	-
Male													
2001	380	-	-	-	84	104	108	51	22	6	4	1	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Legal intervention (Y35,Y89.0)													
All origins													
Male													
2000	348	-	-	3	71	108	96	54	9	5	1	1	-
1999	392	-	-	1	67	135	117	49	20	2	1	-	-
Female													
2001	16	-	-	-	4	3	7	-	2	-	-	-	-
2000	11	-	-	-	2	2	5	1	1	-	-	-	-
1999	6	-	-	-	-	1	4	1	-	-	-	-	-
Hispanic													
Both sexes													
2001	65	-	-	-	23	24	13	4	1	-	-	-	-
2000	48	-	-	1	18	15	9	4	1	-	-	-	-
1999	58	-	-	-	13	23	14	5	2	1	-	-	-
Male													
2001	65	-	-	-	23	24	13	4	1	-	-	-	-
2000	47	-	-	1	17	15	9	4	1	-	-	-	-
1999	58	-	-	-	13	23	14	5	2	1	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	1	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic [1]													
Both sexes													
2001	330	-	-	-	65	83	102	46	23	6	4	1	-
2000	308	-	-	2	54	95	90	51	9	5	1	1	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Legal intervention (Y35,Y89.0)													
Non-Hispanic [1]													
Both sexes													
1999	338	-	-	1	54	112	106	45	18	1	1	-	-
Male													
2001	314	-	-	-	61	80	95	46	21	6	4	1	-
2000	298	-	-	2	53	93	85	50	8	5	1	1	-
1999	332	-	-	1	54	111	102	44	18	1	1	-	-
Female													
2001	16	-	-	-	4	3	7	-	2	-	-	-	-
2000	10	-	-	-	1	2	5	1	1	-	-	-	-
1999	6	-	-	-	-	1	4	1	-	-	-	-	-
Non-Hispanic white													
Both sexes													
2001	230	-	-	-	37	56	74	36	17	6	4	-	-
2000	187	-	-	-	26	57	52	39	8	3	1	1	-
1999	197	-	-	-	20	62	65	35	13	1	1	-	-
Male													
2001	221	-	-	-	35	55	69	36	16	6	4	-	-
2000	182	-	-	-	25	55	51	39	7	3	1	1	-
1999	192	-	-	-	20	62	61	34	13	1	1	-	-
Female													
2001	9	-	-	-	2	1	5	-	1	-	-	-	-
2000	5	-	-	-	1	2	1	-	1	-	-	-	-
1999	5	-	-	-	-	-	4	1	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Legal intervention (Y35,Y89.0)													
Non-Hispanic black													
Both sexes													
2001	82	-	-	-	26	21	22	9	3	-	-	1	-
2000	110	-	-	2	27	33	35	11	-	2	-	-	-
1999	125	-	-	1	31	42	36	10	5	-	-	-	-
Male													
2001	77	-	-	-	24	20	20	9	3	-	-	1	-
2000	107	-	-	2	27	33	33	10	-	2	-	-	-
1999	124	-	-	1	31	41	36	10	5	-	-	-	-
Female													
2001	5	-	-	-	2	1	2	-	-	-	-	-	-
2000	3	-	-	-	-	-	2	1	-	-	-	-	-
1999	1	-	-	-	-	1	-	-	-	-	-	-	-
Origin not stated [2]													
Both sexes													
2001	1	-	-	-	-	-	-	1	-	-	-	-	-
2000	3	-	-	-	1	-	2	-	-	-	-	-	-
1999	2	-	-	-	-	1	1	-	-	-	-	-	-
Male													
2001	1	-	-	-	-	-	-	1	-	-	-	-	-
2000	3	-	-	-	1	-	2	-	-	-	-	-	-
1999	2	-	-	-	-	1	1	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Legal intervention (Y35,Y89.0)													
Origin not stated [2]													
Female													
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)													
All origins													
Both sexes													
2001	4,198	83	44	51	412	728	1,396	990	225	99	108	44	18
2000	3,819	65	39	60	370	679	1,286	841	228	104	79	57	11
1999	3,917	55	43	49	430	743	1,295	793	222	114	96	58	19
Male													
2001	2,833	49	19	38	316	515	931	662	146	64	57	20	16
2000	2,635	27	26	50	285	489	885	570	152	70	49	21	11
1999	2,692	33	21	38	345	523	887	532	148	72	55	25	13
Female													
2001	1,365	34	25	13	96	213	465	328	79	35	51	24	2
2000	1,184	38	13	10	85	190	401	271	76	34	30	36	-
1999	1,225	22	22	11	85	220	408	261	74	42	41	33	6
Hispanic													
Both sexes													
2001	283	7	6	4	34	63	92	54	9	6	2	4	2
2000	215	5	5	1	34	61	64	25	12	5	2	-	1
1999	249	6	9	4	52	69	61	28	13	3	3	-	1
Male													
2001	216	4	3	4	26	52	73	36	7	5	1	3	2
2000	165	2	2	1	30	49	47	20	9	2	2	-	1

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)													
Hispanic													
Male													
1999	200	3	6	3	46	54	49	23	11	2	2	-	1
Female													
2001	67	3	3	-	8	11	19	18	2	1	1	1	-
2000	50	3	3	-	4	12	17	5	3	3	-	-	-
1999	49	3	3	1	6	15	12	5	2	1	1	-	-
Non-Hispanic [1]													
Both sexes													
2001	3,876	76	38	47	377	661	1,298	927	210	91	105	40	6
2000	3,571	58	34	59	336	614	1,214	812	211	96	77	56	4
1999	3,630	49	34	45	376	666	1,230	757	207	107	93	56	10
Male													
2001	2,586	45	16	34	289	459	853	622	134	57	55	17	5
2000	2,445	25	24	49	255	436	834	546	140	65	47	20	4
1999	2,463	30	15	35	297	462	834	503	135	67	53	24	8
Female													
2001	1,290	31	22	13	88	202	445	305	76	34	50	23	1
2000	1,126	33	10	10	81	178	380	266	71	31	30	36	-
1999	1,167	19	19	10	79	204	396	254	72	40	40	32	2
Non-Hispanic white													
Both sexes													
2001	3,155	40	25	35	318	549	1,054	749	176	75	93	37	4
2000	2,823	35	22	37	271	490	965	647	161	76	70	47	2
1999	2,848	30	15	38	290	531	960	578	172	92	86	50	6

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)													
Non-Hispanic white													
Male													
2001	2,054	25	10	25	238	377	673	485	106	45	51	16	3
2000	1,900	16	15	31	201	341	661	425	101	48	43	16	2
1999	1,887	19	8	30	232	357	646	363	101	57	47	22	5
Female													
2001	1,101	15	15	10	80	172	381	264	70	30	42	21	1
2000	923	19	7	6	70	149	304	222	60	28	27	31	-
1999	961	11	7	8	58	174	314	215	71	35	39	28	1
Non-Hispanic black													
Both sexes													
2001	629	36	13	11	43	90	222	163	30	12	5	3	1
2000	654	14	11	21	50	103	230	153	42	15	6	7	2
1999	686	18	15	7	70	119	239	166	28	12	5	3	4
Male													
2001	467	20	6	8	37	65	165	127	26	10	1	1	1
2000	478	3	8	17	45	78	160	113	32	13	4	3	2
1999	514	10	7	5	55	91	170	132	27	9	4	1	3
Female													
2001	162	16	7	3	6	25	57	36	4	2	4	2	-
2000	176	11	3	4	5	25	70	40	10	2	2	4	-
1999	172	8	8	2	15	28	69	34	1	3	1	2	1
Origin not stated [2]													
Both sexes													
2001	39	-	-	-	1	4	6	9	6	2	1	-	10

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)													
Origin not stated [2]													
Both sexes													
2000	33	2	-	-	-	4	8	4	5	3	-	1	6
1999	38	-	-	-	2	8	4	8	2	4	-	2	8
Male													
2001	31	-	-	-	1	4	5	4	5	2	1	-	9
2000	25	-	-	-	-	4	4	4	3	3	-	1	6
1999	29	-	-	-	2	7	4	6	2	3	-	1	4
Female													
2001	8	-	-	-	-	-	1	5	1	-	-	-	1
2000	8	2	-	-	-	-	4	-	2	-	-	-	-
1999	9	-	-	-	-	1	-	2	-	1	-	1	4
Discharge of firearms, undetermined intent (Y22-Y24)													
All origins													
Both sexes													
2001	231	-	-	6	64	48	50	29	12	10	11	1	-
2000	230	-	-	10	71	52	39	23	15	11	6	2	1
1999	324	-	3	12	127	51	57	28	14	15	13	4	-
Male													
2001	190	-	-	6	59	42	33	21	11	8	9	1	-
2000	190	-	-	9	62	43	30	18	11	9	5	2	1
1999	277	-	3	12	111	44	45	21	11	14	12	4	-
Female													
2001	41	-	-	-	5	6	17	8	1	2	2	-	-
2000	40	-	-	1	9	9	9	5	4	2	1	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Discharge of firearms, undetermined intent (Y22-Y24)													
All origins													
Female													
1999	47	-	-	-	16	7	12	7	3	1	1	-	-
Hispanic													
Both sexes													
2001	21	-	-	1	8	4	4	3	1	-	-	-	-
2000	20	-	-	-	13	4	2	-	-	1	-	-	-
1999	31	-	1	1	18	3	3	2	1	2	-	-	-
Male													
2001	16	-	-	1	7	3	2	2	1	-	-	-	-
2000	17	-	-	-	13	2	1	-	-	1	-	-	-
1999	26	-	1	1	16	2	2	1	1	2	-	-	-
Female													
2001	5	-	-	-	1	1	2	1	-	-	-	-	-
2000	3	-	-	-	-	2	1	-	-	-	-	-	-
1999	5	-	-	-	2	1	1	1	-	-	-	-	-
Non-Hispanic [1]													
Both sexes													
2001	208	-	-	5	56	43	46	26	11	9	11	1	-
2000	209	-	-	10	58	48	37	23	15	10	6	2	-
1999	293	-	2	11	109	48	54	26	13	13	13	4	-
Male													
2001	172	-	-	5	52	38	31	19	10	7	9	1	-
2000	172	-	-	9	49	41	29	18	11	8	5	2	-
1999	251	-	2	11	95	42	43	20	10	12	12	4	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Discharge of firearms, undetermined intent (Y22-Y24)													
Non-Hispanic [1]													
Female													
2001	36	-	-	-	4	5	15	7	1	2	2	-	-
2000	37	-	-	1	9	7	8	5	4	2	1	-	-
1999	42	-	-	-	14	6	11	6	3	1	1	-	-
Non-Hispanic white													
Both sexes													
2001	166	-	-	3	36	35	41	23	8	9	10	1	-
2000	148	-	-	5	36	30	30	20	11	8	6	2	-
1999	217	-	1	10	67	33	47	21	11	11	12	4	-
Male													
2001	133	-	-	3	33	31	27	16	7	7	8	1	-
2000	120	-	-	5	30	26	24	15	7	6	5	2	-
1999	187	-	1	10	60	29	39	15	8	10	11	4	-
Female													
2001	33	-	-	-	3	4	14	7	1	2	2	-	-
2000	28	-	-	-	6	4	6	5	4	2	1	-	-
1999	30	-	-	-	7	4	8	6	3	1	1	-	-
Non-Hispanic black													
Both sexes													
2001	31	-	-	2	15	5	2	3	3	-	1	-	-
2000	53	-	-	5	19	15	7	2	3	2	-	-	-
1999	65	-	1	1	35	14	6	4	1	2	1	-	-
Male													
2001	28	-	-	2	14	4	1	3	3	-	1	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Discharge of firearms, undetermined intent (Y22-Y24)													
Non-Hispanic black													
Male													
2000	46	-	-	4	18	12	5	2	3	2	-	-	-
1999	55	-	1	1	29	12	4	4	1	2	1	-	-
Female													
2001	3	-	-	-	1	1	1	-	-	-	-	-	-
2000	7	-	-	1	1	3	2	-	-	-	-	-	-
1999	10	-	-	-	6	2	2	-	-	-	-	-	-
Origin not stated [2]													
Both sexes													
2001	2	-	-	-	-	1	-	-	-	1	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	1
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	2	-	-	-	-	1	-	-	-	1	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	1
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2, Y89.9)													
All origins													
Both sexes													
2001	3,967	83	44	45	348	680	1,346	961	213	89	97	43	18
2000	3,589	65	39	50	299	627	1,247	818	213	93	73	55	10
1999	3,593	55	40	37	303	692	1,238	765	208	99	83	54	19
Male													
2001	2,643	49	19	32	257	473	898	641	135	56	48	19	16
2000	2,445	27	26	41	223	446	855	552	141	61	44	19	10
1999	2,415	33	18	26	234	479	842	511	137	58	43	21	13
Female													
2001	1,324	34	25	13	91	207	448	320	78	33	49	24	2
2000	1,144	38	13	9	76	181	392	266	72	32	29	36	-
1999	1,178	22	22	11	69	213	396	254	71	41	40	33	6
Hispanic													
Both sexes													
2001	262	7	6	3	26	59	88	51	8	6	2	4	2
2000	195	5	5	1	21	57	62	25	12	4	2	-	1
1999	218	6	8	3	34	66	58	26	12	1	3	-	1
Male													
2001	200	4	3	3	19	49	71	34	6	5	1	3	2
2000	148	2	2	1	17	47	46	20	9	1	2	-	1
1999	174	3	5	2	30	52	47	22	10	-	2	-	1
Female													
2001	62	3	3	-	7	10	17	17	2	1	1	1	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2, Y89.9)													
Hispanic													
Female													
2000	47	3	3	-	4	10	16	5	3	3	-	-	-
1999	44	3	3	1	4	14	11	4	2	1	1	-	-
Non-Hispanic [1]													
Both sexes													
2001	3,668	76	38	42	321	618	1,252	901	199	82	94	39	6
2000	3,362	58	34	49	278	566	1,177	789	196	86	71	54	4
1999	3,337	49	32	34	267	618	1,176	731	194	94	80	52	10
Male													
2001	2,414	45	16	29	237	421	822	603	124	50	46	16	5
2000	2,273	25	24	40	206	395	805	528	129	57	42	18	4
1999	2,212	30	13	24	202	420	791	483	125	55	41	20	8
Female													
2001	1,254	31	22	13	84	197	430	298	75	32	48	23	1
2000	1,089	33	10	9	72	171	372	261	67	29	29	36	-
1999	1,125	19	19	10	65	198	385	248	69	39	39	32	2
Non-Hispanic white													
Both sexes													
2001	2,989	40	25	32	282	514	1,013	726	168	66	83	36	4
2000	2,675	35	22	32	235	460	935	627	150	68	64	45	2
1999	2,631	30	14	28	223	498	913	557	161	81	74	46	6
Male													
2001	1,921	25	10	22	205	346	646	469	99	38	43	15	3

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2, Y89.9)													
Non-Hispanic white													
Male													
2000	1,780	16	15	26	171	315	637	410	94	42	38	14	2
1999	1,700	19	7	20	172	328	607	348	93	47	36	18	5
Female													
2001	1,068	15	15	10	77	168	367	257	69	28	40	21	1
2000	895	19	7	6	64	145	298	217	56	26	26	31	-
1999	931	11	7	8	51	170	306	209	68	34	38	28	1
Non-Hispanic black													
Both sexes													
2001	598	36	13	9	28	85	220	160	27	12	4	3	1
2000	601	14	11	16	31	88	223	151	39	13	6	7	2
1999	621	18	14	6	35	105	233	162	27	10	4	3	4
Male													
2001	439	20	6	6	23	61	164	124	23	10	-	1	1
2000	432	3	8	13	27	66	155	111	29	11	4	3	2
1999	459	10	6	4	26	79	166	128	26	7	3	1	3
Female													
2001	159	16	7	3	5	24	56	36	4	2	4	2	-
2000	169	11	3	3	4	22	68	40	10	2	2	4	-
1999	162	8	8	2	9	26	67	34	1	3	1	2	1
Origin not stated [2]													
Both sexes													
2001	37	-	-	-	1	3	6	9	6	1	1	-	10

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2, Y89.9)													
Origin not stated [2]													
Both sexes													
2000	32	2	-	-	-	4	8	4	5	3	-	1	5
1999	38	-	-	-	2	8	4	8	2	4	-	2	8
Male													
2001	29	-	-	-	1	3	5	4	5	1	1	-	9
2000	24	-	-	-	-	4	4	4	3	3	-	1	5
1999	29	-	-	-	2	7	4	6	2	3	-	1	4
Female													
2001	8	-	-	-	-	-	1	5	1	-	-	-	1
2000	8	2	-	-	-	-	4	-	2	-	-	-	-
1999	9	-	-	-	-	1	-	2	-	1	-	1	4
Operations of war and their sequelae (Y36,Y89.1)													
All origins													
Both sexes													
2001	17	-	-	-	1	-	1	2	-	4	8	1	-
2000	16	-	-	-	-	-	-	2	1	3	9	1	-
1999	23	-	-	-	-	-	1	2	1	8	8	3	-
Male													
2001	16	-	-	-	1	-	1	2	-	4	7	1	-
2000	16	-	-	-	-	-	-	2	1	3	9	1	-
1999	22	-	-	-	-	-	1	2	1	8	7	3	-
Female													
2001	1	-	-	-	-	-	-	-	-	-	1	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Operations of war and their sequelae (Y36,Y89.1)													
All origins													
Female													
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	1	-	-
Hispanic													
Both sexes													
2001	2	-	-	-	1	-	-	-	-	-	1	-	-
2000	1	-	-	-	-	-	-	-	-	1	-	-	-
1999	2	-	-	-	-	-	-	-	-	-	1	1	-
Male													
2001	2	-	-	-	1	-	-	-	-	-	1	-	-
2000	1	-	-	-	-	-	-	-	-	1	-	-	-
1999	2	-	-	-	-	-	-	-	-	-	1	1	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic [1]													
Both sexes													
2001	15	-	-	-	-	-	1	2	-	4	7	1	-
2000	15	-	-	-	-	-	-	2	1	2	9	1	-
1999	21	-	-	-	-	-	1	2	1	8	7	2	-
Male													
2001	14	-	-	-	-	-	1	2	-	4	6	1	-
2000	15	-	-	-	-	-	-	2	1	2	9	1	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Operations of war and their sequelae (Y36,Y89.1)													
Non-Hispanic [1]													
Male													
1999	20	-	-	-	-	-	1	2	1	8	6	2	-
Female													
2001	1	-	-	-	-	-	-	-	-	-	1	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	1	-	-
Non-Hispanic white													
Both sexes													
2001	14	-	-	-	-	-	1	2	-	3	7	1	-
2000	14	-	-	-	-	-	-	1	1	2	9	1	-
1999	20	-	-	-	-	-	1	2	1	7	7	2	-
Male													
2001	13	-	-	-	-	-	1	2	-	3	6	1	-
2000	14	-	-	-	-	-	-	1	1	2	9	1	-
1999	19	-	-	-	-	-	1	2	1	7	6	2	-
Female													
2001	1	-	-	-	-	-	-	-	-	-	1	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	1	-	-
Non-Hispanic black													
Both sexes													
2001	1	-	-	-	-	-	-	-	-	1	-	-	-
2000	1	-	-	-	-	-	-	1	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	1	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Operations of war and their sequelae (Y36,Y89.1)													
Non-Hispanic black													
Male													
2001	1	-	-	-	-	-	-	-	-	1	-	-	-
2000	1	-	-	-	-	-	-	1	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	1	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated [2]													
Both sexes													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Male													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-
Female													
2001	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Complications of medical and surgical care (Y40-Y84,Y88)													
All origins													
Both sexes													
2001	3,021	26	24	14	43	79	172	283	384	567	864	564	1
2000	3,059	19	22	31	32	85	149	295	376	647	825	578	-
1999	2,823	54	30	18	36	78	152	214	315	586	811	529	-
Male													
2001	1,357	15	14	8	19	42	83	143	196	265	383	189	-
2000	1,430	12	11	15	16	47	76	143	181	331	402	196	-
1999	1,308	31	12	9	21	43	78	98	165	279	375	197	-
Female													
2001	1,664	11	10	6	24	37	89	140	188	302	481	375	1
2000	1,629	7	11	16	16	38	73	152	195	316	423	382	-
1999	1,515	23	18	9	15	35	74	116	150	307	436	332	-
Hispanic													
Both sexes													
2001	145	3	6	4	8	7	12	22	13	25	29	15	1
2000	136	3	4	6	7	4	11	13	11	35	26	16	-
1999	100	5	6	3	4	8	8	11	16	15	12	12	-
Male													
2001	65	-	3	3	3	3	4	11	8	8	14	8	-
2000	70	2	2	4	6	2	5	4	8	16	15	6	-
1999	52	3	3	2	2	3	7	6	9	5	8	4	-
Female													
2001	80	3	3	1	5	4	8	11	5	17	15	7	1
2000	66	1	2	2	1	2	6	9	3	19	11	10	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Complications of medical and surgical care (Y40-Y84,Y88)													
Hispanic													
Female													
1999	48	2	3	1	2	5	1	5	7	10	4	8	-
Non-Hispanic [1]													
Both sexes													
2001	2,861	23	18	10	34	72	158	261	369	537	831	548	-
2000	2,914	16	18	25	25	81	137	281	363	610	797	561	-
1999	2,714	49	24	15	32	70	141	202	298	570	798	515	-
Male													
2001	1,286	15	11	5	15	39	78	132	188	254	368	181	-
2000	1,354	10	9	11	10	45	70	139	172	313	385	190	-
1999	1,253	28	9	7	19	40	70	91	156	274	366	193	-
Female													
2001	1,575	8	7	5	19	33	80	129	181	283	463	367	-
2000	1,560	6	9	14	15	36	67	142	191	297	412	371	-
1999	1,461	21	15	8	13	30	71	111	142	296	432	322	-
Non-Hispanic white													
Both sexes													
2001	2,294	14	10	6	25	51	105	185	279	417	735	467	-
2000	2,378	12	8	17	17	56	103	218	272	498	691	486	-
1999	2,212	30	16	11	26	47	103	148	224	478	688	441	-
Male													
2001	1,016	9	7	2	12	26	50	91	137	197	333	152	-
2000	1,135	9	6	7	7	31	58	111	135	255	347	169	-
1999	1,025	19	8	5	15	29	52	65	117	230	321	164	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Complications of medical and surgical care (Y40-Y84,Y88)													
Non-Hispanic white													
Female													
2001	1,278	5	3	4	13	25	55	94	142	220	402	315	-
2000	1,243	3	2	10	10	25	45	107	137	243	344	317	-
1999	1,187	11	8	6	11	18	51	83	107	248	367	277	-
Non-Hispanic black													
Both sexes													
2001	515	9	7	3	7	21	48	70	84	104	86	76	-
2000	484	4	8	5	7	23	31	57	86	101	95	67	-
1999	442	18	8	4	6	18	35	51	61	84	94	63	-
Male													
2001	247	6	4	2	2	13	26	38	47	53	30	26	-
2000	192	1	2	2	2	13	12	25	34	52	33	16	-
1999	195	8	1	2	4	10	16	26	30	40	38	20	-
Female													
2001	268	3	3	1	5	8	22	32	37	51	56	50	-
2000	292	3	6	3	5	10	19	32	52	49	62	51	-
1999	247	10	7	2	2	8	19	25	31	44	56	43	-
Origin not stated [2]													
Both sexes													
2001	15	-	-	-	1	-	2	-	2	5	4	1	-
2000	9	-	-	-	-	-	1	1	2	2	2	1	-
1999	9	-	-	-	-	-	3	1	1	1	1	2	-
Male													
2001	6	-	-	-	1	-	1	-	-	3	1	-	-

See footnotes at end of table.

Work table Orig290F. Deaths from 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Not stated
Complications of medical and surgical care (Y40-Y84,Y88)													
Origin not stated [2]													
Male													
2000	6	-	-	-	-	-	1	-	1	2	2	-	-
1999	3	-	-	-	-	-	1	1	-	-	1	-	-
Female													
2001	9	-	-	-	-	-	1	-	2	2	3	1	-
2000	3	-	-	-	-	-	-	1	1	-	-	1	-
1999	6	-	-	-	-	-	2	-	1	1	-	2	-

- Quantity zero.
... Category not applicable.
[1] Includes races other than white and black.
[2] Includes deaths for which Hispanic origin was not reported on the death certificate.
[3] Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

Source: CDC/NCHS, National Vital Statistics System, Mortality.