Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All causes												
All origins [3]												
Both sexes												
2001	848.5	683.4	33.3	17.3	80.7	105.2	203.6	428.9	964.6	2,353.3	5,582.4	15,112.8
2000 [4]	854.0	736.7	32.4	18.0	79.9	101.4	198.9	425.6	992.2	2,399.1	5,666.5	15,524.4
1999 [4]	857.0	736.0	34.2	18.6	79.3	102.2	198.0	418.2	1,005.0	2,457.3	5,714.5	15,554.6
Male												
2001	846.4	749.8	37.0	19.8	117.0	143.7	259.6	545.1	1,192.7	2,911.5	6,833.0	16,744.8
2000 [4]	853.0	806.5	35.9	20.9	114.9	138.6	255.2	542.8	1,230.7	2,979.6	6,972.6	17,501.4
1999 [4]	859.2	805.0	37.9	21.5	113.1	139.7	254.9	533.1	1,252.0	3,073.7	7,083.3	17,597.2
Female												
2001	850.4	613.9	29.5	14.6	42.6	66.0	148.2	316.8	754.0	1,890.8	4,760.5	14,429.9
2000 [4]	855.0	663.4	28.7	15.0	43.1	63.5	143.2	312.5	772.2	1,921.2	4,814.7	14,719.2
1999 [4]	854.9	663.6	30.3	15.6	43.7	64.1	141.8	307.6	777.6	1,952.3	4,825.4	14,731.3
Hispanic												
Both sexes												
2001	306.8	572.7	30.6	14.7	75.6	84.5	154.8	334.4	722.2	1,776.0	4,262.6	11,821.8
2000 [4]	303.8	596.3	29.6	15.7	72.8	84.6	157.6	330.2	744.4	1,803.8	4,335.9	11,822.9
1999 [4]	305.7	583.8	30.9	15.6	71.3	85.5	158.2	333.2	752.5	1,853.0	4,356.3	12,177.9
Male												
2001	332.9	624.4	33.8	16.6	111.5	118.0	208.5	443.9	923.9	2,242.6	5,258.0	12,888.3
2000 [4]	331.3	637.1	31.5	17.9	107.7	120.2	211.0	439.0	965.7	2,287.9	5,395.3	13,086.2
1999 [4]	332.6	623.4	32.9	17.8	104.1	120.6	215.1	444.4	974.8	2,368.9	5,379.2	13,485.9
Female												
2001	279.0	518.9	27.2	12.7	33.7	45.2	97.0	226.7	543.0	1,408.0	3,589.8	11,300.5

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All causes												
Hispanic												
Female												
2000 [4]	274.6	553.6	27.5	13.4	31.7	43.4	100.5	223.8	548.4	1,423.2	3,624.5	11,202.8
1999 [4]	277.2	542.3	28.7	13.3	32.9	44.9	97.4	224.9	555.8	1,448.8	3,675.7	11,547.3
Non-Hispanic [5]												
Both sexes												
2001	926.2	703.5	33.8	17.7	81.4	108.9	209.2	435.6	979.5	2,383.9	5,628.5	15,199.0
2000 [4]	929.6	760.4	32.9	18.4	80.9	104.1	202.9	431.8	1,006.5	2,427.9	5,707.6	15,619.8
1999 [4]	929.9	760.7	34.8	19.1	80.5	104.7	201.6	423.3	1,019.2	2,485.0	5,753.4	15,631.0
Male												
2001	923.6	773.2	37.6	20.4	117.5	148.5	265.2	551.6	1,207.0	2,944.1	6,888.0	16,866.3
2000 [4]	928.1	836.8	36.8	21.5	115.9	141.6	259.2	548.7	1,244.2	3,009.8	7,020.7	17,639.8
1999 [4]	932.2	834.9	38.8	22.1	114.6	142.7	258.1	537.3	1,265.6	3,102.8	7,133.0	17,710.4
Female												
2001	928.6	630.4	29.9	14.9	44.1	69.6	154.3	324.0	768.7	1,918.4	4,801.8	14,505.6
2000 [4]	931.0	680.1	28.8	15.2	45.0	66.8	147.7	319.2	786.8	1,947.5	4,852.4	14,802.9
1999 [4]	927.8	682.9	30.6	16.0	45.4	67.1	146.3	313.7	791.7	1,977.5	4,858.2	14,798.4
Non-Hispanic white												
Both sexes												
2001	991.1	555.4	30.1	16.3	74.1	96.4	188.3	396.0	924.0	2,325.5	5,604.4	15,371.2
2000 [4]	993.2	596.5	28.5	17.1	73.6	90.1	181.0	390.5	950.2	2,366.2	5,675.4	15,799.7
1999 [4]	990.7	593.1	30.1	17.3	72.4	91.3	177.8	381.5	957.8	2,420.0	5,716.0	15,795.3
Male												
2001	975.6	611.6	33.7	18.6	105.1	131.5	241.6	502.6	1,136.3	2,869.4	6,851.5	17,055.3

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All causes												
Non-Hispanic white												
Male												
2000 [4]	978.5	658.7	32.4	20.0	103.5	123.0	233.9	497.7	1,170.9	2,930.5	6,977.8	17,853.2
1999 [4]	979.6	658.1	33.4	19.9	100.8	124.5	230.0	483.7	1,187.4	3,023.2	7,088.0	17,871.2
Female												
2001	1,006.1	496.4	26.3	13.9	41.9	60.9	134.9	291.0	723.5	1,864.1	4,777.3	14,670.6
2000 [4]	1,007.3	530.9	24.4	13.9	42.6	56.8	128.1	285.0	742.1	1,891.0	4,819.3	14,971.7
1999 [4]	1,001.3	524.6	26.7	14.7	42.9	57.8	125.6	281.0	741.6	1,915.1	4,817.7	14,967.5
Non-Hispanic black												
Both sexes												
2001	798.1	1,354.4	49.5	24.0	121.4	191.5	365.8	786.9	1,602.5	3,300.7	6,645.3	14,660.8
2000 [4]	805.5	1,477.8	51.7	25.0	123.2	194.6	365.1	802.2	1,633.9	3,378.8	6,764.3	14,837.5
1999 [4]	812.1	1,510.2	54.6	27.5	125.5	191.1	376.9	796.7	1,681.7	3,454.5	6,805.4	14,839.2
Male												
2001	849.7	1,484.2	55.1	28.6	186.2	267.9	456.3	1,014.6	2,055.4	4,218.9	8,426.8	16,576.2
2000 [4]	859.5	1,623.2	56.1	29.1	187.2	270.2	462.7	1,036.2	2,111.4	4,303.9	8,551.0	16,944.8
1999 [4]	872.8	1,650.8	61.5	33.1	191.4	267.9	479.5	1,048.1	2,176.7	4,397.8	8,615.3	17,385.3
Female												
2001	751.2	1,217.8	43.6	19.3	56.8	122.4	285.7	591.5	1,238.2	2,653.0	5,645.1	13,951.4
2000 [4]	756.7	1,327.7	47.1	20.7	60.1	126.5	278.8	601.2	1,249.4	2,728.0	5,758.6	14,054.9
1999 [4]	757.3	1,365.4	47.4	21.8	60.5	122.1	286.4	581.7	1,285.1	2,789.2	5,792.0	13,887.7

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Salmonella infections (A01-A02)												
All origins [3]												
Both sexes												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Salmonella infections (A01-A02)												
Hispanic												
Female												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Salmonella infections (A01-A02)												
Non-Hispanic white												
Male												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Shigellosis and amebiasis (A03,A06)												
All origins [3]												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Shigellosis and amebiasis (A03,A06)												
Hispanic												
Female												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Shigellosis and amebiasis (A03,A06)												
Non-Hispanic white												
Male												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain other intestinal infections (A04,A07-A09)												
All origins [3]												
Both sexes												
2001	0.5	*	*	*	*	*	*	0.1	0.3	1.3	4.5	12.8
2000 [4]	0.5	*	*	*	*	*	*	0.1	0.3	1.2	3.7	11.8
1999 [4]	0.4	*	*	*	*	*	*	0.1	0.2	1.0	3.2	10.3
Male												
2001	0.4	*	*	*	*	*	*	*	0.3	1.3	4.6	12.1
2000 [4]	0.4	*	*	*	*	*	*	*	0.3	1.3	4.0	11.4
1999 [4]	0.3	*	*	*	*	*	*	*	0.2	0.9	3.5	9.5
Female												
2001	0.7	*	*	*	*	*	*	0.1	0.3	1.3	4.5	13.1
2000 [4]	0.6	*	*	*	*	*	*	*	0.2	1.2	3.6	11.9
1999 [4]	0.5	*	*	*	*	*	*	*	0.2	1.0	3.0	10.7
Hispanic												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain other intestinal infections (A04,A07-A09)												
Hispanic												
Female												
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.6	*	*	*	*	*	*	0.1	0.3	1.3	4.5	12.8
2000 [4]	0.5	*	*	*	*	*	*	0.1	0.3	1.2	3.8	11.9
1999 [4]	0.4	*	*	*	*	*	*	*	0.2	1.0	3.3	10.4
Male												
2001	0.5	*	*	*	*	*	*	*	0.3	1.3	4.7	12.0
2000 [4]	0.4	*	*	*	*	*	*	*	0.3	1.3	4.0	11.5
1999 [4]	0.3	*	*	*	*	*	*	*	0.2	0.9	3.6	9.6
Female												
2001	0.7	*	*	*	*	*	*	0.1	0.4	1.3	4.5	13.2
2000 [4]	0.6	*	*	*	*	*	*	*	0.2	1.2	3.6	12.1
1999 [4]	0.5	*	*	*	*	*	*	*	0.2	1.0	3.1	10.8
Non-Hispanic white												
Both sexes												
2001	0.7	*	*	*	*	*	*	0.1	0.3	1.4	4.7	13.6
2000 [4]	0.6	*	*	*	*	*	*	0.1	0.3	1.3	4.0	12.3
1999 [4]	0.5	*	*	*	*	*	*	*	0.2	1.0	3.4	11.0
Male												
2001	0.5	*	*	*	*	*	*	*	0.3	1.4	4.7	12.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain other intestinal infections (A04,A07-A09)												
Non-Hispanic white												
Male												
2000 [4]	0.5	*	*	*	*	*	*	*	0.3	1.3	4.3	12.0
1999 [4]	0.4	*	*	*	*	*	*	*	0.2	1.0	3.8	10.0
Female												
2001	0.9	*	*	*	*	*	*	0.2	0.3	1.3	4.7	14.0
2000 [4]	0.7	*	*	*	*	*	*	*	0.3	1.2	3.9	12.5
1999 [4]	0.6	*	*	*	*	*	*	*	*	1.0	3.2	11.4
Non-Hispanic black												
Both sexes												
2001	0.3	*	*	*	*	*	*	*	*	1.3	3.0	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	8.3
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	2.8	*
Male												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Tuberculosis (A16-A19)												
All origins [3]												
Both sexes												
2001	0.3	*	*	*	*	0.1	0.1	0.2	0.5	0.8	1.6	2.8
2000 [4]	0.3	*	*	*	*	0.1	0.1	0.2	0.3	0.8	2.0	2.7
1999 [4]	0.3	*	*	*	*	*	0.1	0.3	0.5	0.9	2.1	4.2
Male												
2001	0.3	*	*	*	*	*	0.1	0.3	0.7	1.2	2.0	4.4
2000 [4]	0.3	*	*	*	*	*	0.1	0.4	0.5	1.0	2.8	4.6
1999 [4]	0.4	*	*	*	*	*	0.2	0.5	0.8	1.4	2.8	7.1
Female												
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	2.2
2000 [4]	0.2	*	*	*	*	*	*	*	0.2	0.7	1.5	2.0
1999 [4]	0.2	*	*	*	*	*	*	*	0.3	0.6	1.6	3.0
Hispanic												
Both sexes												
2001	0.3	*	*	*	*	*	*	*	*	*	3.6	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	4.5	*
1999 [4]	0.4	*	*	*	*	*	*	*	1.2	2.3	5.6	*
Male												
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.4	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Tuberculosis (A16-A19)												
Hispanic												
Female												
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	1.5	2.7
2000 [4]	0.3	*	*	*	*	*	0.1	0.2	0.3	0.8	1.9	2.5
1999 [4]	0.3	*	*	*	*	*	0.1	0.2	0.5	0.9	1.9	3.9
Male												
2001	0.3	*	*	*	*	*	0.1	0.3	0.6	1.1	1.9	4.3
2000 [4]	0.3	*	*	*	*	*	0.1	0.4	0.5	1.0	2.6	4.3
1999 [4]	0.4	*	*	*	*	*	0.2	0.4	0.8	1.3	2.6	6.8
Female												
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.3	2.1
2000 [4]	0.2	*	*	*	*	*	*	*	*	0.7	1.4	1.8
1999 [4]	0.2	*	*	*	*	*	*	*	0.2	0.5	1.4	2.8
Non-Hispanic white												
Both sexes												
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.1	2.0
2000 [4]	0.2	*	*	*	*	*	*	0.1	0.1	0.5	1.2	2.0
1999 [4]	0.2	*	*	*	*	*	*	0.1	0.3	0.6	1.3	3.0
Male												
2001	0.2	*	*	*	*	*	*	0.1	0.4	0.6	1.4	2.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Tuberculosis (A16-A19)												
Non-Hispanic white												
Male												
2000 [4]	0.2	*	*	*	*	*	*	0.2	*	0.6	1.5	3.6
1999 [4]	0.3	*	*	*	*	*	*	0.2	0.4	0.8	1.8	4.9
Female												
2001	0.2	*	*	*	*	*	*	*	*	0.4	0.9	1.7
2000 [4]	0.1	*	*	*	*	*	*	*	*	0.3	1.1	1.4
1999 [4]	0.2	*	*	*	*	*	*	*	*	0.3	1.0	2.2
Non-Hispanic black												
Both sexes												
2001	0.6	*	*	*	*	*	*	0.8	1.5	2.7	3.8	6.9
2000 [4]	0.7	*	*	*	*	*	0.5	1.0	1.4	3.2	6.4	*
1999 [4]	0.7	*	*	*	*	*	0.4	1.0	1.7	2.6	5.6	8.4
Male												
2001	0.7	*	*	*	*	*	*	1.4	2.4	4.4	*	*
2000 [4]	0.9	*	*	*	*	*	*	1.9	2.5	4.2	11.7	*
1999 [4]	0.9	*	*	*	*	*	*	1.8	3.0	4.2	7.8	*
Female												
2001	0.4	*	*	*	*	*	*	*	*	*	3.5	*
2000 [4]	0.4	*	*	*	*	*	*	*	*	2.5	3.5	*
1999 [4]	0.5	*	*	*	*	*	*	*	*	*	4.4	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Respiratory tuberculosis (A16)												
All origins [3]												
Both sexes												
2001	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	1.3	2.4
2000 [4]	0.2	*	*	*	*	*	0.1	0.2	0.2	0.6	1.7	2.3
1999 [4]	0.3	*	*	*	*	*	0.1	0.2	0.5	0.8	1.8	3.7
Male												
2001	0.3	*	*	*	*	*	*	0.3	0.6	0.9	1.7	3.8
2000 [4]	0.3	*	*	*	*	*	0.1	0.3	0.4	0.9	2.4	4.1
1999 [4]	0.4	*	*	*	*	*	0.1	0.4	0.7	1.2	2.5	6.4
Female												
2001	0.2	*	*	*	*	*	*	*	*	0.4	1.1	1.8
2000 [4]	0.2	*	*	*	*	*	*	*	*	0.5	1.2	1.6
1999 [4]	0.2	*	*	*	*	*	*	*	0.2	0.4	1.3	2.6
Hispanic												
Both sexes												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	5.0	*
Male												
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Respiratory tuberculosis (A16)												
Hispanic												
Female												
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.2	*	*	*	*	*	0.1	0.1	0.3	0.6	1.2	2.4
2000 [4]	0.2	*	*	*	*	*	0.1	0.2	0.2	0.6	1.6	2.2
1999 [4]	0.3	*	*	*	*	*	0.1	0.2	0.4	0.7	1.6	3.5
Male												
2001	0.3	*	*	*	*	*	*	0.2	0.6	0.8	1.6	3.8
2000 [4]	0.3	*	*	*	*	*	0.1	0.3	0.4	0.8	2.2	3.8
1999 [4]	0.4	*	*	*	*	*	0.1	0.3	0.7	1.1	2.3	6.1
Female												
2001	0.2	*	*	*	*	*	*	*	*	0.4	1.0	1.8
2000 [4]	0.2	*	*	*	*	*	*	*	*	0.4	1.2	1.5
1999 [4]	0.2	*	*	*	*	*	*	*	*	0.4	1.2	2.4
Non-Hispanic white												
Both sexes												
2001	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.9	1.7
2000 [4]	0.1	*	*	*	*	*	*	0.1	*	0.3	1.1	1.8
1999 [4]	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.1	2.8
Male												
2001	0.2	*	*	*	*	*	*	*	0.4	0.4	1.1	2.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Respiratory tuberculosis (A16)												
Non-Hispanic white												
Male												
2000 [4]	0.2	*	*	*	*	*	*	0.2	*	0.5	1.3	3.2
1999 [4]	0.2	*	*	*	*	*	*	0.2	0.4	0.7	1.6	4.8
Female												
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.4
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.9	1.2
1999 [4]	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.9
Non-Hispanic black												
Both sexes												
2001	0.4	*	*	*	*	*	*	0.6	1.2	2.0	3.0	6.2
2000 [4]	0.5	*	*	*	*	*	0.4	0.8	0.9	2.7	5.1	*
1999 [4]	0.6	*	*	*	*	*	*	0.7	1.5	2.0	4.8	7.1
Male												
2001	0.6	*	*	*	*	*	*	1.2	2.1	3.4	*	*
2000 [4]	0.8	*	*	*	*	*	*	1.5	2.1	3.9	9.3	*
1999 [4]	0.7	*	*	*	*	*	*	1.3	2.7	3.2	6.9	*
Female												
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.4	*	*	*	*	*	*	*	*	*	3.7	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other tuberculosis (A17-A19)												
All origins [3]												
Both sexes												
2001	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.3	*
2000 [4]	0.1	*	*	*	*	*	*	*	0.1	0.2	0.3	*
1999 [4]	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.3	0.5
Male												
2001	0.1	*	*	*	*	*	*	*	*	0.3	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.5	*
1999 [4]	0.1	*	*	*	*	*	*	0.1	*	*	*	*
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	0.3	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	0.2	0.3	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	0.3	*
Hispanic												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other tuberculosis (A17-A19)												
Hispanic												
Female												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.1	*	*	*	*	*	*	0.1	*	0.2	0.3	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	0.2	0.3	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	0.2	0.3	*
Male												
2001	0.1	*	*	*	*	*	*	*	*	0.3	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.4	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	0.3	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	0.2	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2001	0.0	*	*	*	*	*	*	*	*	0.1	0.2	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	0.1	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	0.2	*
Male												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other tuberculosis (A17-A19)												
Non-Hispanic white												
Male												
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Whooping cough (A37)												
All origins [3]												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Whooping cough (A37)												
Hispanic												
Female												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Whooping cough (A37)												
Non-Hispanic white												
Male												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Scarlet fever and erysipelas (A38,A46)												
All origins [3]												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Scarlet fever and erysipelas (A38,A46)												
Hispanic												
Female												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Scarlet fever and erysipelas (A38,A46)												
Non-Hispanic white												
Male												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningococcal infection (A39)												
All origins [3]												
Both sexes												
2001	0.1	*	*	0.1	0.1	*	*	*	*	*	*	*
2000 [4]	0.1	0.6	*	*	0.1	*	*	0.1	*	*	*	*
1999 [4]	0.1	0.8	0.2	0.1	0.1	0.1	0.1	*	*	*	*	*
Male												
2001	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	0.2	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	0.1	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	0.1	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningococcal infection (A39)												
Hispanic												
Female												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.1	*	*	0.1	0.2	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	0.1	*	*	0.1	*	*	*	*
1999 [4]	0.1	0.8	0.2	0.1	0.1	*	0.1	*	*	*	*	*
Male												
2001	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	0.2	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.1	*	*	*	0.2	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2001	0.1	*	*	*	0.2	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	0.1	*	*	*	*	*	*	*
1999 [4]	0.1	1.0	0.3	*	0.1	*	*	*	*	*	*	*
Male												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningococcal infection (A39)												
Non-Hispanic white												
Male												
2000 [4]	0.1	*	*	*	0.2	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.1	*	*	*	0.2	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Septicemia (A40-A41)												
All origins [3]												
Both sexes												
2001	11.3	7.7	0.7	0.2	0.3	0.7	1.8	5.0	12.3	32.8	82.3	205.9
2000 [4]	11.1	7.2	0.6	0.2	0.3	0.7	1.9	4.9	11.9	31.0	80.4	215.7
1999 [4]	11.0	7.5	0.6	0.2	0.3	0.7	1.8	4.6	11.4	31.2	79.4	220.7
Male												
2001	10.2	8.4	0.8	0.2	0.4	0.7	2.0	5.4	13.2	36.7	95.5	224.3
2000 [4]	9.8	7.1	0.7	0.2	0.3	0.8	2.0	5.6	13.0	34.5	91.9	223.0
1999 [4]	9.8	9.0	0.6	0.2	0.3	0.7	2.1	5.1	12.4	35.2	91.9	233.5
Female												
2001	12.4	7.0	0.6	0.1	0.3	0.7	1.7	4.7	11.4	29.5	73.7	198.3
2000 [4]	12.3	7.3	0.6	0.1	0.2	0.6	1.9	4.2	11.0	28.1	72.9	212.7
1999 [4]	12.2	5.8	0.5	0.2	0.3	0.6	1.6	4.1	10.5	28.0	71.3	215.5
Hispanic												
Both sexes												
2001	3.7	6.1	*	*	*	0.3	1.3	3.9	8.2	25.4	62.4	171.3
2000 [4]	3.5	8.0	*	*	*	0.5	1.5	3.9	8.5	21.8	59.3	144.7
1999 [4]	3.6	5.7	*	*	*	0.4	1.5	3.1	8.0	25.6	61.7	180.3
Male												
2001	3.6	7.0	*	*	*	*	1.5	4.7	9.9	29.8	76.0	160.0
2000 [4]	3.4	8.4	*	*	*	0.6	1.7	5.0	10.0	26.6	70.3	157.2
1999 [4]	3.5	7.0	*	*	*	*	2.0	3.8	10.2	29.4	75.1	181.2
Female												
2001	3.9	5.2	*	*	*	*	1.1	3.1	6.6	21.9	53.2	176.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Septicemia (A40-A41)												
Hispanic												
Female												
2000 [4]	3.5	7.7	*	*	*	*	1.3	2.8	7.2	18.1	51.8	138.5
1999 [4]	3.7	*	*	*	*	*	1.0	2.4	6.1	22.5	52.7	179.9
Non-Hispanic [5]												
Both sexes												
2001	12.4	8.1	0.7	0.1	0.4	0.8	1.9	5.1	12.5	33.1	83.0	206.8
2000 [4]	12.1	6.9	0.7	0.2	0.3	0.7	2.0	5.0	12.1	31.4	81.0	217.5
1999 [4]	12.0	7.9	0.6	0.2	0.3	0.7	1.8	4.7	11.6	31.4	79.8	221.2
Male												
2001	11.2	8.8	0.9	0.1	0.4	0.8	2.1	5.5	13.4	37.0	96.1	226.8
2000 [4]	10.7	6.7	0.7	0.2	0.3	0.8	2.0	5.6	13.1	34.8	92.3	224.8
1999 [4]	10.7	9.4	0.6	0.2	0.4	0.8	2.1	5.2	12.5	35.4	92.2	234.5
Female												
2001	13.5	7.4	0.6	0.2	0.3	0.8	1.7	4.8	11.8	29.9	74.4	198.5
2000 [4]	13.5	7.0	0.6	0.1	0.2	0.7	1.9	4.3	11.2	28.7	73.6	214.5
1999 [4]	13.2	6.2	0.5	0.2	0.3	0.7	1.6	4.2	10.8	28.2	71.8	215.9
Non-Hispanic white												
Both sexes												
2001	12.3	5.1	0.7	0.1	0.3	0.6	1.4	4.0	10.4	29.4	77.6	195.6
2000 [4]	11.9	4.8	0.5	0.2	0.2	0.5	1.5	3.8	10.0	27.7	74.2	204.3
1999 [4]	11.7	5.3	0.5	0.2	0.3	0.6	1.4	3.6	9.5	27.3	73.0	206.0
Male												
2001	11.0	5.8	0.9	*	0.3	0.5	1.7	4.2	11.1	32.5	89.7	213.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Septicemia (A40-A41)												
Non-Hispanic white												
Male												
2000 [4]	10.4	4.7	0.5	0.2	0.3	0.6	1.6	4.3	10.9	30.9	84.7	211.2
1999 [4]	10.2	6.7	0.5	0.2	0.3	0.7	1.5	4.0	9.9	30.5	84.6	218.1
Female												
2001	13.6	4.4	0.5	*	0.3	0.6	1.2	3.8	9.9	26.8	69.6	188.1
2000 [4]	13.4	4.8	0.6	*	*	0.5	1.5	3.3	9.2	24.9	67.2	201.6
1999 [4]	13.1	3.8	0.5	0.2	0.2	0.5	1.2	3.3	9.1	24.5	65.4	201.2
Non-Hispanic black												
Both sexes												
2001	16.4	20.3	0.9	*	0.7	1.9	4.8	13.7	32.6	72.8	161.9	368.6
2000 [4]	16.7	16.6	1.1	*	0.6	2.1	4.8	13.9	31.8	72.5	175.1	395.9
1999 [4]	16.6	18.6	*	*	0.6	1.6	5.1	13.2	30.8	75.3	169.8	414.2
Male												
2001	15.4	21.4	*	*	0.9	1.9	4.9	15.2	37.4	86.9	196.8	430.8
2000 [4]	15.3	16.4	*	*	*	2.4	4.8	16.8	35.9	81.6	208.9	432.1
1999 [4]	15.7	20.8	*	*	*	1.7	6.0	15.4	37.9	91.5	202.6	451.6
Female												
2001	17.2	19.1	*	*	*	1.9	4.7	12.5	28.8	62.8	142.4	345.6
2000 [4]	17.9	16.9	*	*	*	1.8	4.9	11.5	28.5	66.1	156.1	382.5
1999 [4]	17.4	16.4	*	*	*	1.6	4.3	11.3	25.1	63.9	151.4	400.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Syphilis (A50-A53)												
All origins [3]												
Both sexes												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Syphilis (A50-A53)												
Hispanic												
Female												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Syphilis (A50-A53)												
Non-Hispanic white												
Male												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute poliomyelitis (A80)												
All origins [3]												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute poliomyelitis (A80)												
Hispanic												
Female												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute poliomyelitis (A80)												
Non-Hispanic white												
Male												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Arthropod-borne viral encephalitis (A83-A84,A85.2)												
All origins [3]												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Arthropod-borne viral encephalitis (A83-A84,A85.2)												
Hispanic												
Female												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Arthropod-borne viral encephalitis (A83-A84,A85.2)												
Non-Hispanic white												
Male												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Measles (B05)												
All origins [3]												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Measles (B05)												
Hispanic												
Female												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Measles (B05)												
Non-Hispanic white												
Male												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Viral hepatitis (B15-B19)												
All origins [3]												
Both sexes												
2001	2.0	*	*	*	0.1	0.2	1.8	5.3	4.0	4.6	4.9	3.8
2000 [4]	1.9	*	*	*	*	0.2	1.7	5.0	3.7	4.6	5.3	4.0
1999 [4]	1.7	*	*	*	*	0.2	1.6	4.1	3.3	4.8	5.3	4.7
Male												
2001	2.5	*	*	*	*	0.2	2.5	8.2	5.3	4.8	5.3	4.4
2000 [4]	2.4	*	*	*	*	0.2	2.4	7.7	5.0	4.8	6.4	5.7
1999 [4]	2.2	*	*	*	*	0.3	2.3	6.3	4.3	5.1	6.6	6.9
Female												
2001	1.4	*	*	*	*	0.2	1.1	2.5	2.8	4.4	4.6	3.5
2000 [4]	1.4	*	*	*	*	0.2	1.1	2.4	2.6	4.5	4.7	3.3
1999 [4]	1.3	*	*	*	*	0.2	0.9	2.0	2.4	4.5	4.5	3.9
Hispanic												
Both sexes												
2001	2.1	*	*	*	*	*	2.3	7.8	10.4	9.9	11.0	*
2000 [4]	2.0	*	*	*	*	*	2.5	7.0	8.6	11.1	13.0	*
1999 [4]	1.9	*	*	*	*	*	2.1	6.8	7.8	12.4	13.5	*
Male												
2001	2.6	*	*	*	*	*	3.1	12.2	14.2	7.0	10.3	*
2000 [4]	2.5	*	*	*	*	*	3.8	10.6	11.7	11.0	12.3	*
1999 [4]	2.3	*	*	*	*	*	3.3	10.5	9.8	13.0	*	*
Female												
2001	1.7	*	*	*	*	*	1.4	3.5	7.1	12.3	11.5	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Viral hepatitis (B15-B19)												
Hispanic												
Female												
2000 [4]	1.5	*	*	*	*	*	1.0	3.3	5.8	11.3	13.5	*
1999 [4]	1.5	*	*	*	*	*	0.8	3.3	6.0	12.0	16.3	*
Non-Hispanic [5]												
Both sexes												
2001	1.9	*	*	*	*	0.2	1.7	5.0	3.5	4.2	4.6	3.6
2000 [4]	1.9	*	*	*	*	0.2	1.6	4.8	3.3	4.2	5.0	3.8
1999 [4]	1.7	*	*	*	*	0.2	1.5	3.8	3.0	4.3	5.0	4.7
Male												
2001	2.5	*	*	*	*	0.2	2.4	7.7	4.6	4.6	5.0	4.3
2000 [4]	2.4	*	*	*	*	0.2	2.2	7.4	4.4	4.4	6.1	5.4
1999 [4]	2.2	*	*	*	*	0.2	2.2	5.9	3.9	4.7	6.4	6.8
Female												
2001	1.4	*	*	*	*	0.2	1.0	2.3	2.5	3.8	4.3	3.3
2000 [4]	1.4	*	*	*	*	0.2	1.0	2.4	2.3	4.0	4.3	3.2
1999 [4]	1.2	*	*	*	*	0.2	0.9	1.8	2.1	4.0	4.0	3.9
Non-Hispanic white												
Both sexes												
2001	1.8	*	*	*	*	0.2	1.7	4.6	2.9	3.7	4.3	3.3
2000 [4]	1.8	*	*	*	*	0.1	1.6	4.3	2.7	3.7	4.7	3.7
1999 [4]	1.6	*	*	*	*	0.2	1.4	3.4	2.5	3.7	4.7	4.5
Male												
2001	2.5	*	*	*	*	0.2	2.3	7.2	4.0	4.3	5.1	4.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Viral hepatitis (B15-B19)												
Non-Hispanic white												
Male												
2000 [4]	2.3	*	*	*	*	*	2.1	6.7	3.6	3.8	5.9	5.4
1999 [4]	2.1	*	*	*	*	0.2	2.1	5.2	3.4	4.3	6.4	6.5
Female												
2001	1.2	*	*	*	*	*	1.0	2.0	2.0	3.1	3.8	2.9
2000 [4]	1.3	*	*	*	*	*	1.0	2.0	1.8	3.6	4.0	3.1
1999 [4]	1.1	*	*	*	*	0.2	0.8	1.7	1.6	3.3	3.6	3.7
Non-Hispanic black												
Both sexes												
2001	2.3	*	*	*	*	*	2.2	8.1	6.8	5.6	5.7	*
2000 [4]	2.2	*	*	*	*	0.4	2.1	8.0	5.8	5.8	6.3	*
1999 [4]	2.1	*	*	*	*	*	2.2	7.0	5.1	6.8	5.9	*
Male												
2001	2.8	*	*	*	*	*	3.0	12.0	9.0	5.0	*	*
2000 [4]	2.9	*	*	*	*	*	3.0	12.3	8.1	7.2	7.7	*
1999 [4]	2.6	*	*	*	*	*	3.1	11.8	6.3	5.7	*	*
Female												
2001	1.8	*	*	*	*	*	1.4	4.7	5.0	6.0	6.7	*
2000 [4]	1.6	*	*	*	*	*	1.4	4.3	3.9	4.7	5.6	*
1999 [4]	1.6	*	*	*	*	*	1.4	2.9	4.1	7.5	6.7	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)												
All origins [3]												
Both sexes												
2001	5.0	*	*	0.1	0.6	5.3	13.0	10.5	5.2	2.1	0.7	*
2000 [4]	5.1	*	*	0.1	0.5	6.1	13.1	11.0	5.1	2.2	0.7	*
1999 [4]	5.3	*	0.2	0.2	0.5	6.8	13.8	10.7	4.8	2.2	0.6	*
Male												
2001	7.6	*	*	0.1	0.5	7.1	19.5	16.8	8.6	3.5	1.5	*
2000 [4]	7.9	*	*	0.1	0.5	8.0	19.8	17.8	8.7	3.8	1.3	*
1999 [4]	8.2	*	*	0.2	0.5	9.5	21.0	17.5	8.3	3.8	1.0	*
Female												
2001	2.5	*	*	*	0.6	3.5	6.7	4.4	2.0	0.9	*	*
2000 [4]	2.5	*	*	0.1	0.4	4.2	6.5	4.4	1.8	0.8	0.3	*
1999 [4]	2.5	*	*	0.2	0.5	4.1	6.7	4.1	1.6	0.8	0.3	*
Hispanic												
Both sexes												
2001	5.0	*	*	*	0.5	4.4	13.9	15.5	9.2	4.5	*	*
2000 [4]	5.4	*	*	*	0.3	5.6	14.9	16.9	9.5	3.8	*	*
1999 [4]	5.8	*	*	*	0.5	6.4	17.3	15.4	8.4	4.7	*	*
Male												
2001	7.6	*	*	*	0.6	6.0	20.8	24.9	14.8	8.6	*	*
2000 [4]	8.2	*	*	*	*	7.8	22.7	27.5	15.3	7.2	*	*
1999 [4]	8.7	*	*	*	0.6	9.3	26.2	24.2	15.4	9.0	*	*
Female												
2001	2.3	*	*	*	*	2.5	6.5	6.2	4.1	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)												
Hispanic												
Female												
2000 [4]	2.4	*	*	*	*	3.0	6.4	6.6	4.4	*	*	*
1999 [4]	2.6	*	*	*	*	3.2	7.9	6.7	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	4.9	*	*	0.1	0.6	5.4	12.7	9.9	4.8	1.9	0.6	*
2000 [4]	5.0	*	*	0.1	0.5	6.2	12.7	10.3	4.7	2.0	0.6	*
1999 [4]	5.2	*	*	0.2	0.5	6.8	13.2	10.0	4.5	2.0	0.6	*
Male												
2001	7.5	*	*	0.1	0.5	7.3	19.0	15.8	8.0	3.1	1.3	*
2000 [4]	7.7	*	*	*	0.5	7.9	19.1	16.7	8.0	3.4	1.2	*
1999 [4]	8.0	*	*	0.2	0.5	9.4	20.1	16.5	7.6	3.4	1.0	*
Female												
2001	2.5	*	*	*	0.7	3.6	6.6	4.2	1.8	0.9	*	*
2000 [4]	2.5	*	*	0.1	0.5	4.4	6.4	4.2	1.6	0.8	*	*
1999 [4]	2.5	*	*	0.1	0.6	4.2	6.5	3.9	1.6	0.7	0.3	*
Non-Hispanic white												
Both sexes												
2001	2.2	*	*	*	0.1	2.1	6.0	4.2	2.1	0.7	0.3	*
2000 [4]	2.3	*	*	*	0.1	2.4	6.2	4.6	2.2	0.8	0.3	*
1999 [4]	2.4	*	*	*	0.1	3.0	6.4	4.6	2.0	0.8	0.3	*
Male												
2001	3.8	*	*	*	*	3.2	10.4	7.5	3.8	1.4	0.6	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)												
Non-Hispanic white												
Male												
2000 [4]	4.0	*	*	*	0.2	3.7	10.4	8.2	4.0	1.5	0.6	*
1999 [4]	4.2	*	*	*	0.2	4.9	10.9	8.2	3.6	1.5	0.6	*
Female												
2001	0.6	*	*	*	*	0.9	1.7	1.0	0.5	*	*	*
2000 [4]	0.7	*	*	*	*	1.1	2.0	1.1	0.5	*	*	*
1999 [4]	0.7	*	*	*	*	1.2	1.9	1.1	0.4	*	*	*
Non-Hispanic black												
Both sexes												
2001	21.6	*	*	0.4	2.8	24.0	55.0	51.9	28.0	12.8	5.3	*
2000 [4]	22.0	*	*	0.4	2.2	27.2	54.5	53.9	26.3	13.2	4.6	*
1999 [4]	22.3	*	*	0.6	2.3	28.3	57.2	52.2	26.4	13.2	3.7	*
Male												
2001	30.8	*	*	*	2.4	30.8	77.8	81.4	47.7	21.4	11.4	*
2000 [4]	31.7	*	*	*	2.2	33.4	79.0	86.6	46.4	23.8	8.9	*
1999 [4]	32.6	*	*	0.7	1.9	37.1	83.5	85.5	45.6	24.0	6.6	*
Female												
2001	13.3	*	*	*	3.2	17.8	34.8	26.6	12.1	6.7	*	*
2000 [4]	13.1	*	*	*	2.3	21.7	32.8	25.8	10.1	5.8	*	*
1999 [4]	13.0	*	*	*	2.8	20.3	34.1	23.8	11.0	5.6	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malaria (B50-B54)												
All origins [3]												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malaria (B50-B54)												
Hispanic												
Female												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malaria (B50-B54)												
Non-Hispanic white												
Male												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44, A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8, A86-B04,B06-B09,B25-B49,B55-B99)												
All origins [3]												
Both sexes												
2001	2.0	4.7	0.4	0.2	0.2	0.4	0.8	1.7	3.0	6.2	10.6	20.1
2000 [4]	2.0	4.4	0.5	0.2	0.2	0.4	0.8	1.6	2.9	5.8	11.1	21.2
1999 [4]	2.6	5.3	0.5	0.2	0.2	0.6	1.2	2.3	3.8	7.8	14.2	27.0
Male												
2001	2.1	4.9	0.5	0.1	0.2	0.4	1.0	2.2	3.7	6.8	13.2	23.6
2000 [4]	2.1	4.8	0.5	0.2	0.3	0.4	1.0	1.9	3.3	7.2	13.6	27.6
1999 [4]	2.7	6.1	0.6	0.2	0.2	0.6	1.5	2.7	4.2	9.2	17.2	31.9
Female												
2001	1.9	4.6	0.4	0.2	0.2	0.3	0.6	1.2	2.4	5.8	8.9	18.7
2000 [4]	1.9	3.9	0.4	0.2	0.2	0.3	0.6	1.2	2.6	4.7	9.5	18.6
1999 [4]	2.5	4.5	0.4	0.2	0.2	0.6	0.9	1.9	3.4	6.7	12.3	25.1
Hispanic												
Both sexes												
2001	1.1	5.0	*	*	*	0.5	0.8	2.0	3.0	5.9	10.3	15.0
2000 [4]	1.2	3.0	*	*	*	0.4	1.0	1.9	3.2	5.8	11.3	19.2
1999 [4]	1.5	4.8	*	*	*	0.6	1.4	2.3	3.8	7.7	14.2	26.0
Male												
2001	1.3	5.0	*	*	*	0.6	1.1	2.8	4.4	7.6	12.1	*
2000 [4]	1.3	*	*	*	*	0.6	1.4	2.1	4.4	5.9	13.3	*
1999 [4]	1.5	5.7	*	*	*	0.6	1.9	2.7	3.8	7.9	15.1	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44, A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8, A86-B04,B06-B09,B25-B49,B55-B99)												
Hispanic												
Female												
2001	1.0	5.0	*	*	*	*	*	1.3	*	4.6	9.1	*
2000 [4]	1.0	*	*	*	*	*	*	1.8	*	5.6	9.9	*
1999 [4]	1.4	*	*	*	*	*	*	1.9	3.8	7.6	13.5	28.1
Non-Hispanic [5]												
Both sexes												
2001	2.1	4.6	0.5	0.2	0.2	0.3	0.8	1.6	3.0	6.2	10.6	20.3
2000 [4]	2.1	4.7	0.4	0.2	0.2	0.3	0.7	1.5	2.9	5.8	11.1	21.2
1999 [4]	2.8	5.5	0.5	0.2	0.2	0.6	1.2	2.3	3.8	7.8	14.2	27.0
Male												
2001	2.2	4.8	0.5	0.1	0.2	0.4	1.0	2.1	3.6	6.7	13.2	24.0
2000 [4]	2.2	5.1	0.5	0.2	0.3	0.3	0.9	1.9	3.2	7.3	13.5	27.8
1999 [4]	2.8	6.2	0.6	0.2	0.3	0.6	1.4	2.6	4.2	9.2	17.2	32.3
Female												
2001	2.1	4.5	0.4	0.2	0.2	0.3	0.6	1.2	2.5	5.8	8.9	18.8
2000 [4]	2.0	4.3	0.4	0.2	0.2	0.4	0.6	1.1	2.6	4.7	9.5	18.6
1999 [4]	2.7	4.7	0.4	0.2	0.2	0.6	1.0	1.9	3.3	6.7	12.2	24.9
Non-Hispanic white												
Both sexes												
2001	2.1	3.3	0.4	0.1	0.2	0.3	0.7	1.5	2.8	6.0	10.4	20.2
2000 [4]	2.1	3.5	0.4	0.2	0.2	0.2	0.6	1.3	2.6	5.6	10.9	21.0
1999 [4]	2.7	4.4	0.5	0.2	0.2	0.4	0.9	1.8	3.3	7.4	13.6	26.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44, A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8, A86-B04,B06-B09,B25-B49,B55-B99)												
Non-Hispanic white												
Male												
2001	2.2	3.3	0.4	*	0.2	0.3	0.8	1.9	3.3	6.5	13.1	24.2
2000 [4]	2.2	3.9	*	*	0.2	0.2	0.7	1.7	3.0	7.0	13.2	27.0
1999 [4]	2.7	5.5	0.6	0.2	0.2	0.4	1.2	2.2	3.8	8.8	16.3	30.8
Female												
2001	2.1	3.4	*	0.2	0.2	0.2	0.5	1.0	2.3	5.7	8.7	18.6
2000 [4]	2.0	3.1	*	0.2	0.2	0.2	0.5	1.0	2.3	4.5	9.5	18.6
1999 [4]	2.6	3.3	*	*	0.2	0.4	0.6	1.5	2.9	6.3	11.8	24.7
Non-Hispanic black												
Both sexes												
2001	2.3	10.0	*	*	0.4	0.9	1.6	3.0	5.2	8.3	11.3	20.9
2000 [4]	2.3	10.4	*	*	0.4	1.0	1.7	3.0	4.9	8.0	13.3	24.2
1999 [4]	3.6	10.0	1.0	*	0.4	1.7	3.1	5.6	7.3	11.9	20.9	34.3
Male												
2001	2.5	11.5	*	*	*	*	2.1	4.0	6.4	9.8	12.6	*
2000 [4]	2.6	10.6	*	*	*	0.9	2.2	4.0	5.8	10.5	17.9	31.7
1999 [4]	3.7	10.0	*	*	*	1.9	3.1	6.3	7.9	14.1	29.4	46.4
Female												
2001	2.1	8.4	*	*	*	1.0	1.2	2.1	4.3	7.2	10.6	20.5
2000 [4]	2.1	10.2	*	*	*	1.0	1.3	2.2	4.1	6.2	10.8	21.4
1999 [4]	3.5	10.0	*	*	*	1.5	3.0	4.9	6.7	10.4	16.1	29.8

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms (C00-C97)												
All origins [3]												
Both sexes												
2001	194.4	1.6	2.7	2.5	4.3	10.1	36.8	126.5	356.5	802.8	1,315.8	1,765.6
2000 [4]	196.5	2.4	2.7	2.5	4.4	9.8	36.6	127.5	366.7	816.3	1,335.6	1,819.4
1999 [4]	197.0	1.8	2.7	2.5	4.5	10.0	37.1	127.6	374.6	827.1	1,331.5	1,805.8
Male												
2001	205.3	1.5	2.9	2.5	5.0	9.3	32.6	130.3	405.2	984.6	1,727.1	2,613.6
2000 [4]	207.2	2.6	3.0	2.7	5.1	9.2	32.7	130.9	415.8	1,001.9	1,760.6	2,710.7
1999 [4]	208.9	1.9	2.9	2.7	5.2	9.7	33.3	131.7	428.4	1,019.4	1,767.0	2,722.2
Female												
2001	183.9	1.8	2.5	2.4	3.5	10.9	41.0	122.7	311.5	652.2	1,045.4	1,410.7
2000 [4]	186.2	2.3	2.5	2.2	3.6	10.4	40.4	124.2	321.3	663.6	1,058.5	1,456.4
1999 [4]	185.6	1.7	2.5	2.3	3.7	10.3	40.9	123.7	325.0	669.5	1,048.5	1,436.5
Hispanic												
Both sexes												
2001	60.5	*	2.8	2.5	4.3	7.2	25.6	82.8	223.4	504.0	913.9	1,363.7
2000 [4]	59.9	*	2.5	2.9	4.2	7.3	25.2	82.1	221.0	516.2	953.5	1,383.5
1999 [4]	59.6	*	2.5	2.7	4.1	7.4	24.4	84.9	225.1	520.5	926.8	1,399.6
Male												
2001	62.2	*	2.9	2.5	5.5	6.3	21.7	81.5	253.5	642.8	1,258.3	1,967.4
2000 [4]	61.3	*	2.9	3.5	4.5	6.9	20.1	79.4	253.1	651.2	1,306.4	2,049.7
1999 [4]	61.2	*	2.3	2.8	4.6	6.6	19.8	84.6	268.1	668.9	1,252.9	1,945.3
Female												
2001	58.7	*	2.7	2.5	2.9	8.3	29.8	84.2	196.7	394.5	681.2	1,068.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms (C00-C97)												
Hispanic												
Female												
2000 [4]	58.5	*	2.2	2.3	3.7	7.8	30.7	84.7	192.5	410.0	716.5	1,056.5
1999 [4]	58.0	*	2.7	2.6	3.5	8.3	29.4	85.1	187.2	404.3	709.8	1,136.5
Non-Hispanic [5]												
Both sexes												
2001	213.9	1.6	2.7	2.4	4.2	10.6	38.2	130.3	365.9	820.5	1,331.4	1,776.6
2000 [4]	215.6	2.5	2.7	2.4	4.4	10.3	37.9	131.2	376.6	832.8	1,348.9	1,831.1
1999 [4]	215.5	1.7	2.7	2.4	4.5	10.4	38.6	131.0	384.6	843.2	1,344.5	1,816.1
Male												
2001	227.2	1.6	2.9	2.5	4.9	9.9	34.0	134.6	415.3	1,003.8	1,745.1	2,634.8
2000 [4]	228.7	2.6	2.9	2.5	5.2	9.7	34.3	135.1	426.6	1,020.2	1,776.0	2,732.4
1999 [4]	229.9	1.6	3.0	2.6	5.3	10.3	35.0	135.4	438.8	1,036.8	1,783.7	2,747.5
Female												
2001	201.2	1.6	2.5	2.3	3.6	11.3	42.4	126.0	320.0	668.1	1,059.8	1,419.6
2000 [4]	203.1	2.4	2.6	2.2	3.6	10.9	41.4	127.3	330.3	678.1	1,070.6	1,466.6
1999 [4]	201.9	1.8	2.5	2.2	3.7	10.5	42.1	126.8	334.5	684.2	1,059.4	1,443.2
Non-Hispanic white												
Both sexes												
2001	230.6	1.8	2.9	2.4	4.3	10.3	36.5	124.6	358.3	815.9	1,326.5	1,775.2
2000 [4]	232.2	2.6	2.8	2.4	4.3	10.1	35.6	125.1	369.3	827.7	1,342.9	1,826.6
1999 [4]	231.5	1.6	3.0	2.3	4.3	10.2	36.6	124.8	375.0	837.1	1,336.4	1,808.3
Male												
2001	243.4	1.7	3.1	2.4	4.9	10.0	32.6	127.3	401.7	988.2	1,721.8	2,616.8

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms (C00-C97)												
Non-Hispanic white												
Male												
2000 [4]	244.4	2.9	3.1	2.6	5.0	9.7	32.3	127.2	412.0	1,002.1	1,750.2	2,714.1
1999 [4]	245.1	1.7	3.3	2.6	5.0	10.1	33.5	126.7	422.3	1,019.3	1,757.7	2,721.3
Female												
2001	218.4	1.8	2.8	2.4	3.6	10.7	40.4	121.9	317.3	669.7	1,064.4	1,425.1
2000 [4]	220.6	2.4	2.4	2.3	3.5	10.5	38.9	123.0	328.9	681.0	1,075.3	1,468.7
1999 [4]	218.4	*	2.7	2.1	3.6	10.3	39.7	123.0	330.5	684.6	1,060.5	1,444.2
Non-Hispanic black												
Both sexes												
2001	172.8	*	2.0	2.7	4.4	13.4	52.8	188.7	494.5	1,005.6	1,559.5	1,947.5
2000 [4]	174.9	*	2.8	2.3	5.3	12.3	55.3	196.7	505.2	1,019.3	1,582.8	2,015.1
1999 [4]	176.9	*	2.0	2.8	5.6	12.6	55.3	196.8	530.5	1,033.6	1,592.9	2,024.0
Male												
2001	190.8	*	2.4	3.0	5.1	10.9	45.8	209.6	614.1	1,354.7	2,310.5	3,238.5
2000 [4]	194.9	*	2.6	2.6	6.4	10.5	49.6	219.0	637.4	1,382.7	2,372.3	3,290.1
1999 [4]	197.4	*	2.0	2.9	6.6	11.9	48.6	225.8	670.9	1,387.0	2,377.4	3,361.1
Female												
2001	156.5	*	*	2.3	3.8	15.7	59.0	170.7	398.3	759.4	1,137.8	1,469.3
2000 [4]	156.9	*	3.0	1.9	4.2	14.0	60.4	177.5	398.8	763.6	1,138.4	1,541.6
1999 [4]	158.3	*	2.0	2.7	4.8	13.3	61.3	172.0	417.9	784.2	1,153.6	1,524.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
All origins [3]												
Both sexes												
2001	2.7	*	*	*	0.1	0.1	0.6	2.7	6.9	10.4	14.2	18.5
2000 [4]	2.7	*	*	*	0.1	0.1	0.6	2.8	6.8	10.7	13.2	19.4
1999 [4]	2.7	*	*	*	0.1	0.1	0.6	2.9	6.8	10.3	14.3	19.3
Male												
2001	3.7	*	*	*	*	0.2	0.9	4.4	11.3	16.0	20.6	25.8
2000 [4]	3.6	*	*	*	*	0.1	1.0	4.5	10.9	16.0	19.2	27.1
1999 [4]	3.6	*	*	*	0.1	0.1	0.9	4.6	10.8	15.8	20.9	27.1
Female												
2001	1.8	*	*	*	*	0.1	0.4	1.2	2.8	5.8	10.0	15.4
2000 [4]	1.8	*	*	*	*	0.1	0.3	1.3	2.9	6.3	9.2	16.3
1999 [4]	1.8	*	*	*	*	*	0.4	1.3	3.1	5.8	9.9	16.1
Hispanic												
Both sexes												
2001	0.8	*	*	*	*	*	*	1.6	3.7	6.9	10.5	19.4
2000 [4]	0.8	*	*	*	*	*	*	1.3	4.4	6.8	9.5	18.6
1999 [4]	0.8	*	*	*	*	*	*	1.4	4.0	6.9	13.1	14.0
Male												
2001	1.2	*	*	*	*	*	*	2.4	6.3	11.6	17.4	*
2000 [4]	1.2	*	*	*	*	*	*	2.2	7.7	12.2	13.8	40.3
1999 [4]	1.2	*	*	*	*	*	*	2.3	7.2	13.6	18.2	*
Female												
2001	0.5	*	*	*	*	*	*	*	*	3.1	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
Hispanic												
Female												
2000 [4]	0.4	*	*	*	*	*	*	*	*	*	6.6	*
1999 [4]	0.5	*	*	*	*	*	*	*	*	*	9.7	*
Non-Hispanic [5]												
Both sexes												
2001	3.0	*	*	*	0.1	0.2	0.7	2.8	7.1	10.6	14.3	18.4
2000 [4]	2.9	*	*	*	0.1	0.1	0.7	3.0	6.9	10.8	13.3	19.5
1999 [4]	2.9	*	*	*	0.1	0.1	0.7	3.0	7.0	10.5	14.3	19.4
Male												
2001	4.1	*	*	*	*	0.2	1.0	4.6	11.6	16.2	20.6	25.8
2000 [4]	3.9	*	*	*	*	0.1	1.0	4.6	11.1	16.1	19.4	26.5
1999 [4]	4.0	*	*	*	0.1	0.2	1.0	4.8	11.0	15.8	21.0	27.3
Female												
2001	1.9	*	*	*	*	0.1	0.4	1.2	2.9	6.0	10.2	15.3
2000 [4]	2.0	*	*	*	*	0.1	0.4	1.3	3.0	6.5	9.3	16.6
1999 [4]	2.0	*	*	*	*	*	0.4	1.4	3.2	6.0	9.9	16.2
Non-Hispanic white												
Both sexes												
2001	3.1	*	*	*	*	0.1	0.5	2.5	6.5	10.2	14.5	18.9
2000 [4]	3.0	*	*	*	*	0.1	0.6	2.5	6.1	10.4	13.3	19.8
1999 [4]	3.0	*	*	*	*	0.1	0.5	2.4	6.3	10.1	14.2	19.9
Male												
2001	4.1	*	*	*	*	0.2	0.8	4.0	10.4	15.3	20.5	26.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
Non-Hispanic white												
Male												
2000 [4]	3.9	*	*	*	*	*	0.8	3.8	9.7	15.0	18.9	27.4
1999 [4]	3.9	*	*	*	*	*	0.8	3.8	9.7	15.1	20.3	27.5
Female												
2001	2.1	*	*	*	*	*	0.3	1.0	2.8	5.9	10.5	15.7
2000 [4]	2.1	*	*	*	*	*	0.4	1.2	2.8	6.5	9.6	16.8
1999 [4]	2.1	*	*	*	*	*	0.3	1.1	3.0	5.9	10.1	16.8
Non-Hispanic black												
Both sexes												
2001	3.0	*	*	*	*	*	1.3	5.3	12.9	15.7	13.9	13.1
2000 [4]	3.1	*	*	*	*	*	1.1	6.1	13.7	15.7	15.1	14.9
1999 [4]	3.3	*	*	*	*	*	1.2	7.3	13.6	15.4	16.8	14.5
Male												
2001	4.6	*	*	*	*	*	1.7	8.9	23.9	27.3	23.4	*
2000 [4]	4.9	*	*	*	*	*	1.8	10.7	24.7	28.6	29.0	*
1999 [4]	5.0	*	*	*	*	*	1.8	12.3	23.5	25.4	31.9	25.0
Female												
2001	1.5	*	*	*	*	*	0.9	2.1	4.0	7.5	8.6	10.7
2000 [4]	1.5	*	*	*	*	*	*	2.1	4.9	6.6	7.3	14.4
1999 [4]	1.7	*	*	*	*	*	0.7	3.0	5.7	8.4	8.4	10.7

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of esophagus (C15)												
All origins [3]												
Both sexes												
2001	4.4	*	*	*	*	0.1	0.7	3.6	10.1	20.6	26.3	25.8
2000 [4]	4.3	*	*	*	*	0.1	0.7	3.6	10.8	20.0	25.6	25.9
1999 [4]	4.3	*	*	*	*	0.1	0.7	3.5	10.6	20.1	24.4	26.8
Male												
2001	6.9	*	*	*	*	0.1	1.1	6.3	17.6	36.0	47.4	48.1
2000 [4]	6.7	*	*	*	*	0.2	1.2	6.3	18.6	34.1	45.8	47.8
1999 [4]	6.6	*	*	*	*	0.1	1.1	6.1	18.3	35.3	44.4	48.0
Female												
2001	2.0	*	*	*	*	*	0.2	1.0	3.2	7.9	12.4	16.5
2000 [4]	2.1	*	*	*	*	*	0.1	1.0	3.5	8.3	12.4	17.1
1999 [4]	2.0	*	*	*	*	*	0.2	0.9	3.6	7.6	11.5	18.3
Hispanic												
Both sexes												
2001	1.1	*	*	*	*	*	*	1.9	5.2	8.4	18.2	20.0
2000 [4]	1.1	*	*	*	*	*	*	1.2	5.0	11.1	19.4	19.9
1999 [4]	1.1	*	*	*	*	*	*	1.5	5.2	9.2	18.1	26.7
Male												
2001	1.7	*	*	*	*	*	*	3.6	8.5	16.0	30.4	41.9
2000 [4]	1.6	*	*	*	*	*	*	2.2	8.0	19.4	35.4	40.3
1999 [4]	1.7	*	*	*	*	*	*	2.7	10.2	16.3	32.8	49.6
Female												
2001	0.5	*	*	*	*	*	*	*	2.2	*	10.0	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of esophagus (C15)												
Hispanic												
Female												
2000 [4]	0.5	*	*	*	*	*	*	*	2.4	4.6	8.6	*
1999 [4]	0.4	*	*	*	*	*	*	*	*	3.6	8.3	*
Non-Hispanic [5]												
Both sexes												
2001	4.9	*	*	*	*	0.1	0.7	3.7	10.4	21.3	26.6	26.0
2000 [4]	4.8	*	*	*	*	0.1	0.7	3.8	11.2	20.4	25.8	26.1
1999 [4]	4.7	*	*	*	*	0.1	0.7	3.6	11.0	20.6	24.6	26.8
Male												
2001	7.7	*	*	*	*	0.1	1.3	6.5	18.2	37.1	48.1	48.4
2000 [4]	7.5	*	*	*	*	0.2	1.3	6.6	19.4	34.8	46.1	47.8
1999 [4]	7.4	*	*	*	*	0.2	1.2	6.4	18.8	36.2	44.7	48.0
Female												
2001	2.2	*	*	*	*	*	0.3	1.1	3.2	8.2	12.5	16.7
2000 [4]	2.3	*	*	*	*	*	0.1	1.0	3.6	8.5	12.6	17.3
1999 [4]	2.2	*	*	*	*	*	0.2	0.9	3.8	7.8	11.6	18.3
Non-Hispanic white												
Both sexes												
2001	5.1	*	*	*	*	0.1	0.8	3.5	9.7	20.5	26.8	26.5
2000 [4]	5.1	*	*	*	*	0.1	0.7	3.5	10.4	20.0	25.5	26.7
1999 [4]	4.9	*	*	*	*	0.1	0.7	3.3	10.2	20.0	24.4	27.2
Male												
2001	8.1	*	*	*	*	*	1.3	6.1	17.0	36.0	48.6	49.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of esophagus (C15)												
Non-Hispanic white												
Male												
2000 [4]	8.0	*	*	*	*	0.2	1.4	6.3	18.3	34.3	45.9	49.4
1999 [4]	7.7	*	*	*	*	0.2	1.2	6.0	17.5	35.4	44.4	48.0
Female												
2001	2.3	*	*	*	*	*	0.2	1.0	2.7	7.5	12.4	17.1
2000 [4]	2.3	*	*	*	*	*	*	0.8	3.0	8.0	12.0	17.5
1999 [4]	2.2	*	*	*	*	*	0.2	0.8	3.3	7.1	11.3	18.9
Non-Hispanic black												
Both sexes												
2001	4.7	*	*	*	*	*	1.0	6.1	18.9	32.8	30.0	24.0
2000 [4]	4.7	*	*	*	*	*	0.8	6.8	20.2	28.6	34.1	22.6
1999 [4]	4.7	*	*	*	*	*	1.0	6.6	20.3	30.2	31.4	23.6
Male												
2001	7.2	*	*	*	*	*	1.6	10.5	33.0	56.2	55.5	47.4
2000 [4]	7.0	*	*	*	*	*	1.2	11.5	34.8	47.6	58.9	38.7
1999 [4]	7.2	*	*	*	*	*	1.7	11.5	35.3	51.1	57.2	49.9
Female												
2001	2.4	*	*	*	*	*	*	2.3	7.5	16.2	15.7	15.4
2000 [4]	2.6	*	*	*	*	*	*	2.7	8.5	15.2	20.1	16.6
1999 [4]	2.4	*	*	*	*	*	*	2.4	8.3	15.5	17.0	13.8

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of stomach (C16)												
All origins [3]												
Both sexes												
2001	4.3	*	*	*	0.1	0.3	1.0	2.9	7.0	16.5	30.2	45.4
2000 [4]	4.5	*	*	*	*	0.3	1.1	2.9	7.4	16.6	31.0	52.7
1999 [4]	4.6	*	*	*	*	0.3	1.1	3.0	7.8	17.1	31.6	50.1
Male												
2001	5.1	*	*	*	*	0.4	1.1	3.8	10.2	23.2	43.1	62.6
2000 [4]	5.3	*	*	*	*	0.3	1.3	3.8	10.5	23.7	43.9	75.6
1999 [4]	5.5	*	*	*	*	0.4	1.3	4.2	11.0	24.2	46.5	71.3
Female												
2001	3.5	*	*	*	*	0.3	0.9	2.0	4.1	10.9	21.7	38.3
2000 [4]	3.7	*	*	*	*	0.3	0.9	2.1	4.6	10.8	22.6	43.4
1999 [4]	3.7	*	*	*	*	0.3	1.0	1.9	4.9	11.3	21.9	41.6
Hispanic												
Both sexes												
2001	3.2	*	*	*	*	0.6	1.8	5.4	11.3	26.4	45.6	63.8
2000 [4]	3.1	*	*	*	*	0.4	2.0	4.8	9.8	28.4	50.2	65.7
1999 [4]	3.1	*	*	*	*	0.6	1.7	4.1	11.7	26.2	49.8	73.7
Male												
2001	3.6	*	*	*	*	*	1.9	6.2	15.0	37.1	60.4	95.2
2000 [4]	3.6	*	*	*	*	*	2.0	5.0	13.7	43.3	70.8	98.8
1999 [4]	3.5	*	*	*	*	0.7	1.6	5.2	16.4	35.6	73.5	75.5
Female												
2001	2.8	*	*	*	*	0.7	1.7	4.6	8.1	18.0	35.7	48.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of stomach (C16)												
Hispanic												
Female												
2000 [4]	2.7	*	*	*	*	*	2.1	4.6	6.3	16.8	36.3	49.5
1999 [4]	2.7	*	*	*	*	*	1.8	3.0	7.6	18.8	34.0	72.8
Non-Hispanic [5]												
Both sexes												
2001	4.5	*	*	*	*	0.2	0.9	2.6	6.6	15.8	29.4	44.7
2000 [4]	4.7	*	*	*	*	0.3	0.9	2.7	7.2	15.8	30.1	52.0
1999 [4]	4.7	*	*	*	*	0.3	1.0	2.9	7.5	16.5	30.8	49.2
Male												
2001	5.4	*	*	*	*	0.3	1.0	3.6	9.8	22.3	42.1	61.0
2000 [4]	5.5	*	*	*	*	0.3	1.2	3.7	10.3	22.4	42.6	74.5
1999 [4]	5.7	*	*	*	*	0.3	1.2	4.0	10.6	23.4	45.3	71.0
Female												
2001	3.6	*	*	*	*	0.2	0.8	1.7	3.7	10.4	21.0	37.9
2000 [4]	3.8	*	*	*	*	0.3	0.7	1.8	4.4	10.4	21.9	42.9
1999 [4]	3.8	*	*	*	*	0.3	0.9	1.8	4.7	10.8	21.3	40.5
Non-Hispanic white												
Both sexes												
2001	4.2	*	*	*	*	0.2	0.7	2.1	5.6	13.5	25.8	40.0
2000 [4]	4.4	*	*	*	*	0.2	0.7	2.2	6.1	13.5	26.1	46.3
1999 [4]	4.4	*	*	*	*	0.2	0.7	2.3	6.1	14.4	27.0	43.5
Male												
2001	5.0	*	*	*	*	0.2	0.8	2.8	8.2	18.8	37.4	55.7

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of stomach (C16)												
Non-Hispanic white												
Male												
2000 [4]	5.2	*	*	*	*	0.2	0.8	3.0	8.9	19.3	37.4	67.3
1999 [4]	5.4	*	*	*	*	0.2	0.9	3.3	8.7	20.6	39.9	62.6
Female												
2001	3.4	*	*	*	*	*	0.6	1.3	3.1	9.0	18.1	33.5
2000 [4]	3.5	*	*	*	*	0.2	0.5	1.4	3.6	8.6	18.7	37.9
1999 [4]	3.5	*	*	*	*	0.2	0.6	1.4	3.7	9.1	18.6	35.9
Non-Hispanic black												
Both sexes												
2001	5.8	*	*	*	*	0.5	2.2	5.5	13.1	31.0	59.1	89.3
2000 [4]	6.1	*	*	*	*	0.7	2.1	5.8	14.1	32.3	64.3	102.7
1999 [4]	6.3	*	*	*	*	0.7	2.3	6.0	16.1	31.5	65.1	103.1
Male												
2001	6.9	*	*	*	*	*	2.4	8.0	20.0	49.2	87.9	106.3
2000 [4]	7.2	*	*	*	*	0.8	3.0	8.1	20.9	47.0	91.0	138.6
1999 [4]	7.6	*	*	*	*	*	2.8	8.7	23.3	46.4	101.6	149.8
Female												
2001	4.7	*	*	*	*	*	2.0	3.3	7.6	18.1	43.0	83.0
2000 [4]	5.2	*	*	*	*	*	1.4	3.9	8.6	22.0	49.3	89.4
1999 [4]	5.1	*	*	*	*	*	1.9	3.7	10.2	21.0	44.6	85.7

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of colon, rectum and anus (C18-C21)												
All origins [3]												
Both sexes												
2001	20.0	*	*	*	0.1	0.8	3.2	11.6	31.5	75.3	139.4	255.7
2000 [4]	20.4	*	*	*	0.1	0.7	3.0	11.2	32.7	76.7	146.0	270.4
1999 [4]	20.5	*	*	*	0.1	0.7	3.0	11.1	33.5	77.4	146.7	271.7
Male												
2001	20.2	*	*	*	0.1	0.8	3.5	13.1	38.3	93.7	169.5	298.3
2000 [4]	20.6	*	*	*	0.1	0.8	3.4	12.8	39.6	95.4	179.5	315.6
1999 [4]	20.6	*	*	*	0.1	0.8	3.0	12.5	40.5	95.1	181.9	324.0
Female												
2001	19.8	*	*	*	0.1	0.7	2.9	10.2	25.3	60.1	119.6	237.8
2000 [4]	20.2	*	*	*	*	0.6	2.7	9.8	26.4	61.2	124.2	251.9
1999 [4]	20.3	*	*	*	0.1	0.6	2.9	9.7	27.0	62.9	123.8	250.6
Hispanic												
Both sexes												
2001	6.1	*	*	*	*	0.3	2.3	7.2	22.5	52.1	96.5	193.8
2000 [4]	6.0	*	*	*	*	0.5	2.3	8.4	23.1	49.2	98.2	181.1
1999 [4]	6.0	*	*	*	*	0.5	1.7	8.3	21.0	56.1	101.0	176.8
Male												
2001	6.2	*	*	*	*	*	2.7	7.4	28.2	66.5	116.7	257.1
2000 [4]	6.2	*	*	*	*	*	2.0	9.0	28.2	63.5	131.3	235.8
1999 [4]	6.1	*	*	*	*	*	1.5	9.1	24.1	72.7	134.0	211.4
Female												
2001	5.9	*	*	*	*	*	1.9	7.1	17.4	40.8	82.8	162.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of colon, rectum and anus (C18-C21)												
Hispanic												
Female												
2000 [4]	5.8	*	*	*	*	*	2.7	7.8	18.5	38.0	75.9	154.3
1999 [4]	5.9	*	*	*	*	*	2.0	7.5	18.3	43.0	79.1	160.1
Non-Hispanic [5]												
Both sexes												
2001	22.0	*	*	*	0.1	0.9	3.3	12.0	32.2	76.7	141.1	257.4
2000 [4]	22.4	*	*	*	0.1	0.7	3.1	11.5	33.3	78.2	147.8	273.1
1999 [4]	22.4	*	*	*	0.1	0.7	3.1	11.3	34.3	78.5	148.1	274.3
Male												
2001	22.3	*	*	*	0.1	0.9	3.6	13.6	38.9	95.2	171.8	299.6
2000 [4]	22.8	*	*	*	0.1	0.8	3.6	13.1	40.3	97.2	181.3	318.6
1999 [4]	22.7	*	*	*	0.1	0.9	3.2	12.8	41.6	96.2	183.2	327.5
Female												
2001	21.7	*	*	*	*	0.8	3.0	10.5	25.9	61.3	121.0	239.9
2000 [4]	22.1	*	*	*	*	0.6	2.7	9.9	26.9	62.6	126.1	254.7
1999 [4]	22.2	*	*	*	*	0.6	3.0	9.9	27.6	63.9	125.3	253.0
Non-Hispanic white												
Both sexes												
2001	23.5	*	*	*	0.1	0.8	3.1	11.1	30.6	74.7	138.5	256.9
2000 [4]	24.0	*	*	*	0.1	0.6	2.8	10.6	31.7	76.5	145.7	271.9
1999 [4]	24.0	*	*	*	0.1	0.7	2.9	10.6	32.8	76.7	145.9	273.4
Male												
2001	23.9	*	*	*	*	0.9	3.3	12.8	37.4	93.2	169.3	298.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of colon, rectum and anus (C18-C21)												
Non-Hispanic white												
Male												
2000 [4]	24.2	*	*	*	*	0.8	3.2	12.2	38.2	95.1	178.4	315.8
1999 [4]	24.3	*	*	*	*	0.8	3.1	12.1	39.7	94.8	180.6	328.2
Female												
2001	23.1	*	*	*	*	0.7	2.8	9.5	24.2	59.0	118.1	239.8
2000 [4]	23.8	*	*	*	*	0.5	2.4	9.0	25.5	60.9	124.1	254.1
1999 [4]	23.7	*	*	*	*	0.6	2.7	9.0	26.2	61.5	123.2	251.5
Non-Hispanic black												
Both sexes												
2001	19.2	*	*	*	*	1.1	5.0	19.5	50.6	105.9	189.7	289.0
2000 [4]	19.4	*	*	*	*	1.0	5.2	20.0	51.1	106.3	192.5	312.9
1999 [4]	19.3	*	*	*	*	1.1	4.9	18.1	53.7	107.2	194.5	308.8
Male												
2001	19.0	*	*	*	*	1.2	5.8	21.4	59.3	130.7	229.9	352.3
2000 [4]	19.8	*	*	*	*	1.1	6.1	22.7	63.7	133.7	244.4	381.6
1999 [4]	19.2	*	*	*	*	1.2	4.8	20.6	67.0	127.8	245.8	351.8
Female												
2001	19.4	*	*	*	*	1.0	4.3	17.9	43.6	88.4	167.1	265.6
2000 [4]	19.1	*	*	*	*	1.0	4.4	17.6	40.9	87.1	163.2	287.4
1999 [4]	19.5	*	*	*	*	0.9	5.0	15.9	43.1	92.7	165.8	292.7

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)												
All origins [3]												
Both sexes												
2001	4.7	*	*	*	0.1	0.2	0.9	4.8	9.3	18.8	29.0	34.1
2000 [4]	4.6	*	0.1	0.0	0.1	0.2	1.0	4.5	9.3	18.7	28.8	32.7
1999 [4]	4.4	*	0.2	*	0.1	0.2	1.0	4.1	8.8	19.0	28.0	30.6
Male												
2001	6.1	*	*	*	*	0.2	1.3	7.7	13.9	26.8	41.0	50.3
2000 [4]	5.9	*	*	*	0.1	0.3	1.3	7.2	13.8	27.0	42.1	44.7
1999 [4]	5.6	*	0.3	*	0.1	0.2	1.4	6.5	12.9	27.5	39.0	40.1
Female												
2001	3.4	*	*	*	*	0.2	0.4	1.9	5.0	12.1	21.1	27.2
2000 [4]	3.3	*	*	*	*	0.1	0.6	1.9	5.1	11.8	20.1	27.8
1999 [4]	3.3	*	*	*	*	0.2	0.6	1.8	5.1	11.9	20.9	26.8
Hispanic												
Both sexes												
2001	3.4	*	*	*	*	*	0.8	6.7	14.1	28.8	52.1	52.5
2000 [4]	3.3	*	*	*	*	*	1.2	5.6	13.7	31.1	51.8	58.4
1999 [4]	3.2	*	*	*	*	*	0.9	5.7	13.1	31.3	49.8	53.3
Male												
2001	4.3	*	*	*	*	*	1.3	11.1	21.5	41.1	68.0	64.8
2000 [4]	4.2	*	*	*	*	*	1.7	9.1	20.5	43.9	74.7	72.6
1999 [4]	3.9	*	*	*	*	*	1.1	9.6	20.7	42.8	63.1	66.9
Female												
2001	2.5	*	*	*	*	*	*	2.3	7.6	19.1	41.4	46.5

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)												
Hispanic												
Female												
2000 [4]	2.4	*	*	*	*	*	*	2.2	7.6	21.1	36.3	51.4
1999 [4]	2.4	*	*	*	*	*	*	1.9	6.3	22.2	40.9	46.8
Non-Hispanic [5]												
Both sexes												
2001	4.9	*	*	*	0.1	0.2	0.9	4.6	8.9	18.1	27.9	33.2
2000 [4]	4.8	*	0.2	*	0.1	0.2	0.9	4.3	8.9	17.9	27.7	31.7
1999 [4]	4.6	*	0.2	*	0.1	0.2	1.0	3.9	8.4	18.2	27.1	29.7
Male												
2001	6.3	*	*	*	*	0.2	1.3	7.4	13.2	25.8	39.7	49.7
2000 [4]	6.2	*	*	*	*	0.3	1.3	6.9	13.3	25.9	40.5	43.4
1999 [4]	5.8	*	*	*	0.1	0.3	1.5	6.2	12.2	26.5	37.9	38.7
Female												
2001	3.5	*	*	*	*	0.2	0.4	1.9	4.8	11.7	20.2	26.4
2000 [4]	3.4	*	*	*	*	0.1	0.6	1.9	4.9	11.2	19.4	27.0
1999 [4]	3.4	*	*	*	*	0.2	0.5	1.8	4.9	11.3	20.0	26.1
Non-Hispanic white												
Both sexes												
2001	4.8	*	*	*	0.1	0.1	0.6	3.7	7.6	16.6	26.5	32.4
2000 [4]	4.6	*	*	*	*	0.1	0.6	3.4	7.5	16.2	26.5	30.8
1999 [4]	4.5	*	0.2	*	0.1	0.1	0.7	3.1	7.1	16.8	26.1	28.9
Male												
2001	6.1	*	*	*	*	*	0.9	5.9	11.2	24.2	38.0	48.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)												
Non-Hispanic white												
Male												
2000 [4]	5.9	*	*	*	*	0.2	0.8	5.3	11.0	23.6	39.2	41.6
1999 [4]	5.7	*	*	*	*	*	1.1	4.8	10.2	24.6	36.7	39.0
Female												
2001	3.4	*	*	*	*	*	0.3	1.5	4.1	10.2	18.9	25.5
2000 [4]	3.4	*	*	*	*	*	0.4	1.6	4.3	10.1	18.2	26.5
1999 [4]	3.4	*	*	*	*	*	0.4	1.6	4.2	10.2	19.1	24.8
Non-Hispanic black												
Both sexes												
2001	4.6	*	*	*	*	*	1.7	8.9	15.1	21.9	32.3	35.3
2000 [4]	4.5	*	*	*	*	0.4	1.8	8.8	14.5	23.7	29.1	31.8
1999 [4]	4.3	*	*	*	*	0.4	1.7	7.4	13.4	24.6	29.7	34.9
Male												
2001	6.1	*	*	*	*	*	2.7	15.1	24.2	30.3	42.6	42.7
2000 [4]	6.3	*	*	*	*	*	2.5	15.0	24.1	37.3	40.4	48.1
1999 [4]	5.7	*	*	*	*	*	2.5	13.1	20.3	36.8	41.6	30.9
Female												
2001	3.3	*	*	*	*	*	0.8	3.6	7.8	16.0	26.4	32.5
2000 [4]	2.9	*	*	*	*	*	1.2	3.4	6.7	14.1	22.8	25.7
1999 [4]	3.1	*	*	*	*	*	0.9	2.5	7.9	15.9	23.1	36.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of pancreas (C25)												
All origins [3]												
Both sexes												
2001	10.5	*	*	*	*	0.1	1.3	6.1	19.2	44.5	74.1	100.1
2000 [4]	10.4	*	*	*	*	0.1	1.2	6.3	19.4	44.1	76.5	96.4
1999 [4]	10.4	*	*	*	*	0.1	1.3	6.4	19.8	45.0	73.1	99.7
Male												
2001	10.3	*	*	*	*	0.1	1.4	7.6	23.6	50.8	84.0	105.1
2000 [4]	10.3	*	*	*	*	0.2	1.4	8.0	23.6	50.4	85.9	102.9
1999 [4]	10.4	*	*	*	*	0.2	1.6	8.3	23.9	52.4	81.8	107.5
Female												
2001	10.6	*	*	*	*	0.1	1.1	4.7	15.2	39.4	67.6	98.1
2000 [4]	10.5	*	*	*	*	0.1	1.0	4.6	15.5	38.8	70.4	93.8
1999 [4]	10.5	*	*	*	*	0.1	1.0	4.7	16.0	39.0	67.4	96.6
Hispanic												
Both sexes												
2001	3.6	*	*	*	*	*	0.8	3.7	14.2	36.9	61.3	83.8
2000 [4]	3.5	*	*	*	*	*	0.7	4.1	13.7	36.9	61.8	82.9
1999 [4]	3.3	*	*	*	*	*	0.9	4.6	13.3	32.6	56.7	84.2
Male												
2001	3.4	*	*	*	*	*	0.9	4.3	16.2	42.3	68.9	62.9
2000 [4]	3.4	*	*	*	*	*	0.8	5.2	16.6	45.2	62.9	80.6
1999 [4]	3.2	*	*	*	*	*	1.2	5.7	15.2	40.0	56.3	86.3
Female												
2001	3.9	*	*	*	*	*	0.8	3.2	12.5	32.6	56.2	94.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of pancreas (C25)												
Hispanic												
Female												
2000 [4]	3.6	*	*	*	*	*	*	3.0	11.1	30.4	61.1	84.1
1999 [4]	3.5	*	*	*	*	*	*	3.5	11.6	26.8	56.9	83.2
Non-Hispanic [5]												
Both sexes												
2001	11.5	*	*	*	*	0.1	1.3	6.4	19.6	44.9	74.6	100.6
2000 [4]	11.4	*	*	*	*	0.1	1.3	6.5	19.8	44.4	77.0	96.6
1999 [4]	11.4	*	*	*	*	0.1	1.4	6.6	20.2	45.6	73.5	100.1
Male												
2001	11.4	*	*	*	*	0.2	1.5	7.9	24.2	51.2	84.6	106.4
2000 [4]	11.3	*	*	*	*	0.2	1.5	8.3	24.1	50.7	86.7	103.5
1999 [4]	11.4	*	*	*	*	0.2	1.7	8.4	24.4	52.9	82.7	108.2
Female												
2001	11.5	*	*	*	*	0.1	1.2	4.9	15.4	39.8	68.0	98.2
2000 [4]	11.4	*	*	*	*	*	1.0	4.7	15.8	39.2	70.6	93.8
1999 [4]	11.4	*	*	*	*	*	1.0	4.8	16.4	39.6	67.6	96.8
Non-Hispanic white												
Both sexes												
2001	12.3	*	*	*	*	0.1	1.3	6.1	19.3	44.1	73.7	99.4
2000 [4]	12.2	*	*	*	*	0.2	1.2	6.3	19.2	43.4	75.7	94.7
1999 [4]	12.1	*	*	*	*	0.1	1.3	6.3	19.5	44.7	72.1	98.9
Male												
2001	12.4	*	*	*	*	0.2	1.5	7.7	24.0	50.3	84.5	108.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of pancreas (C25)												
Non-Hispanic white												
Male												
2000 [4]	12.2	*	*	*	*	0.2	1.5	8.0	23.4	50.0	86.1	103.0
1999 [4]	12.1	*	*	*	*	0.2	1.6	8.0	23.4	52.1	81.6	108.1
Female												
2001	12.3	*	*	*	*	*	1.2	4.6	14.9	38.8	66.6	95.8
2000 [4]	12.2	*	*	*	*	*	1.0	4.5	15.3	37.9	68.9	91.4
1999 [4]	12.1	*	*	*	*	*	1.0	4.7	15.7	38.5	65.9	95.2
Non-Hispanic black												
Both sexes												
2001	9.4	*	*	*	*	*	1.6	9.2	25.9	59.0	93.3	122.0
2000 [4]	9.7	*	*	*	*	*	1.7	9.2	27.9	60.0	98.9	121.2
1999 [4]	9.7	*	*	*	*	*	2.0	9.9	28.8	60.4	95.0	119.0
Male												
2001	8.9	*	*	*	*	*	1.8	11.4	30.8	68.2	100.5	100.5
2000 [4]	9.3	*	*	*	*	*	2.1	12.1	34.6	66.9	108.0	110.4
1999 [4]	9.7	*	*	*	*	*	2.4	13.6	37.4	68.7	107.6	115.3
Female												
2001	9.9	*	*	*	*	*	1.4	7.3	21.9	52.4	89.3	130.0
2000 [4]	10.0	*	*	*	*	*	1.5	6.8	22.5	55.1	93.8	125.2
1999 [4]	9.7	*	*	*	*	*	1.7	6.7	21.9	54.6	87.9	120.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of larynx (C32)												
All origins [3]												
Both sexes												
2001	1.3	*	*	*	*	*	0.2	1.1	3.6	6.4	7.2	6.7
2000 [4]	1.4	*	*	*	*	*	0.1	1.1	3.6	6.6	8.1	6.5
1999 [4]	1.4	*	*	*	*	*	0.2	1.1	3.8	6.7	7.3	7.0
Male												
2001	2.2	*	*	*	*	*	0.3	1.8	6.1	11.1	14.2	18.3
2000 [4]	2.2	*	*	*	*	*	0.2	1.8	6.2	11.4	15.7	17.4
1999 [4]	2.2	*	*	*	*	*	0.3	1.9	6.4	11.4	14.1	17.8
Female												
2001	0.5	*	*	*	*	*	0.1	0.4	1.3	2.5	2.6	1.9
2000 [4]	0.6	*	*	*	*	*	*	0.4	1.2	2.7	3.1	2.1
1999 [4]	0.6	*	*	*	*	*	*	0.3	1.4	2.9	2.8	2.6
Hispanic												
Both sexes												
2001	0.4	*	*	*	*	*	*	0.6	2.1	5.3	5.2	*
2000 [4]	0.4	*	*	*	*	*	*	*	1.7	4.2	6.9	*
1999 [4]	0.5	*	*	*	*	*	*	0.8	1.8	5.3	8.3	*
Male												
2001	0.7	*	*	*	*	*	*	*	3.9	10.6	10.7	*
2000 [4]	0.7	*	*	*	*	*	*	*	3.5	7.8	14.8	*
1999 [4]	0.9	*	*	*	*	*	*	1.6	3.5	10.5	16.2	*
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of larynx (C32)												
Hispanic												
Female												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	1.5	*	*	*	*	*	0.2	1.1	3.7	6.4	7.3	6.7
2000 [4]	1.5	*	*	*	*	*	0.1	1.2	3.7	6.7	8.1	6.5
1999 [4]	1.5	*	*	*	*	*	0.2	1.1	3.9	6.8	7.2	6.8
Male												
2001	2.4	*	*	*	*	*	0.3	1.9	6.1	11.1	14.3	18.4
2000 [4]	2.4	*	*	*	*	*	0.2	1.9	6.4	11.5	15.7	17.3
1999 [4]	2.4	*	*	*	*	*	0.3	1.9	6.6	11.4	14.0	17.3
Female												
2001	0.6	*	*	*	*	*	0.1	0.4	1.4	2.6	2.7	1.8
2000 [4]	0.6	*	*	*	*	*	*	0.5	1.2	2.8	3.1	2.1
1999 [4]	0.6	*	*	*	*	*	*	0.3	1.4	3.0	2.8	2.6
Non-Hispanic white												
Both sexes												
2001	1.5	*	*	*	*	*	0.2	0.9	3.3	6.0	6.9	6.5
2000 [4]	1.5	*	*	*	*	*	0.1	1.0	3.2	6.2	7.7	6.5
1999 [4]	1.5	*	*	*	*	*	0.1	0.9	3.5	6.4	7.1	6.5
Male												
2001	2.3	*	*	*	*	*	0.3	1.5	5.5	10.2	13.3	18.3

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of larynx (C32)												
Non-Hispanic white												
Male												
2000 [4]	2.4	*	*	*	*	*	0.2	1.5	5.4	10.5	15.0	17.4
1999 [4]	2.3	*	*	*	*	*	0.2	1.6	5.8	10.7	13.6	16.0
Female												
2001	0.6	*	*	*	*	*	*	0.3	1.2	2.4	2.7	1.6
2000 [4]	0.6	*	*	*	*	*	*	0.4	1.1	2.6	3.0	2.1
1999 [4]	0.7	*	*	*	*	*	*	0.3	1.3	2.9	2.9	2.7
Non-Hispanic black												
Both sexes												
2001	1.9	*	*	*	*	*	0.4	2.9	7.9	11.5	12.0	8.4
2000 [4]	2.0	*	*	*	*	*	*	2.9	9.2	12.9	14.3	7.0
1999 [4]	1.9	*	*	*	*	*	0.6	3.0	8.7	11.7	10.0	8.4
Male												
2001	3.1	*	*	*	*	*	*	5.2	14.2	21.4	28.2	*
2000 [4]	3.4	*	*	*	*	*	*	5.2	17.5	24.6	29.0	*
1999 [4]	3.2	*	*	*	*	*	1.0	5.3	16.3	22.1	22.5	25.0
Female												
2001	0.7	*	*	*	*	*	*	*	2.9	4.6	*	*
2000 [4]	0.8	*	*	*	*	*	*	1.0	2.4	4.7	6.1	*
1999 [4]	0.7	*	*	*	*	*	*	0.9	2.7	4.3	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)												
All origins [3]												
Both sexes												
2001	54.8	*	*	*	0.1	0.4	6.2	30.7	117.7	279.7	371.4	302.7
2000 [4]	55.3	*	*	*	0.1	0.5	6.1	31.6	122.4	284.2	370.8	302.1
1999 [4]	54.5	*	*	*	0.1	0.4	6.0	31.3	123.4	281.5	362.1	297.0
Male												
2001	64.7	*	*	*	*	0.4	6.7	37.4	147.1	370.0	529.5	512.4
2000 [4]	65.5	*	*	*	*	0.5	6.9	38.5	154.0	377.9	532.2	521.2
1999 [4]	65.4	*	*	*	*	0.5	6.6	38.2	158.0	377.2	527.6	529.2
Female												
2001	45.3	*	*	*	*	0.4	5.7	24.3	90.5	204.9	267.5	215.0
2000 [4]	45.4	*	*	*	*	0.5	5.3	25.0	93.3	206.9	265.6	212.8
1999 [4]	44.1	*	*	*	*	0.4	5.5	24.6	91.5	203.1	254.5	203.5
Hispanic												
Both sexes												
2001	10.3	*	*	*	*	*	1.9	10.7	41.0	108.3	182.3	200.7
2000 [4]	10.3	*	*	*	*	*	1.8	10.9	39.3	110.6	198.7	213.7
1999 [4]	10.4	*	*	*	*	*	2.0	10.4	44.2	112.9	187.7	227.3
Male												
2001	13.1	*	*	*	*	*	2.4	14.2	59.5	163.7	293.8	335.2
2000 [4]	13.3	*	*	*	*	*	2.1	14.8	58.6	167.3	327.5	368.8
1999 [4]	13.6	*	*	*	*	*	2.3	14.2	68.1	175.3	316.0	362.3
Female												
2001	7.4	*	*	*	*	*	1.4	7.3	24.6	64.6	107.0	135.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)												
Hispanic												
Female												
2000 [4]	7.2	*	*	*	*	*	1.4	7.1	22.2	66.0	112.3	137.5
1999 [4]	7.1	*	*	*	*	*	1.7	6.7	23.1	64.0	102.3	162.2
Non-Hispanic [5]												
Both sexes												
2001	61.3	*	*	*	*	0.4	6.8	32.5	123.3	290.4	379.4	305.8
2000 [4]	61.5	*	*	*	*	0.5	6.6	33.4	128.4	294.2	377.4	304.5
1999 [4]	60.5	*	*	*	*	0.5	6.5	33.0	128.9	290.7	368.2	298.8
Male												
2001	72.6	*	*	*	*	0.4	7.3	39.5	153.2	382.4	539.5	518.4
2000 [4]	73.2	*	*	*	*	0.6	7.5	40.5	160.6	389.7	539.9	526.2
1999 [4]	72.7	*	*	*	*	0.5	7.1	40.1	164.0	387.8	534.9	534.6
Female												
2001	50.5	*	*	*	*	0.5	6.2	25.8	95.6	213.9	274.3	217.4
2000 [4]	50.4	*	*	*	*	0.5	5.8	26.5	98.6	215.4	271.6	214.8
1999 [4]	48.8	*	*	*	*	0.4	6.0	26.1	96.5	211.0	260.0	204.4
Non-Hispanic white												
Both sexes												
2001	67.3	*	*	*	*	0.5	6.6	31.5	124.0	295.2	383.0	308.6
2000 [4]	67.4	*	*	*	*	0.5	6.2	32.0	129.1	298.0	381.7	306.8
1999 [4]	65.9	*	*	*	*	0.5	6.2	31.5	128.9	293.5	370.3	300.5
Male												
2001	78.4	*	*	*	*	0.4	6.9	37.0	150.3	381.1	536.5	520.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)												
Non-Hispanic white												
Male												
2000 [4]	78.9	*	*	*	*	0.6	6.7	37.7	157.7	387.3	537.7	527.3
1999 [4]	78.1	*	*	*	*	0.5	6.6	37.1	160.1	384.4	531.1	536.9
Female												
2001	56.6	*	*	*	*	0.5	6.4	26.0	99.2	222.3	281.3	220.6
2000 [4]	56.4	*	*	*	*	0.5	5.7	26.4	102.2	222.9	279.2	218.0
1999 [4]	54.3	*	*	*	*	0.5	5.8	26.1	99.5	217.5	265.0	206.2
Non-Hispanic black												
Both sexes												
2001	44.6	*	*	*	*	*	9.2	47.5	149.4	311.4	390.9	281.6
2000 [4]	45.4	*	*	*	*	0.6	10.4	51.5	155.0	320.2	379.4	291.0
1999 [4]	45.6	*	*	*	*	0.4	9.6	51.5	161.1	317.7	390.9	293.3
Male												
2001	59.2	*	*	*	*	*	11.4	67.5	218.3	484.4	668.0	550.9
2000 [4]	60.3	*	*	*	*	*	13.8	72.4	227.1	495.4	649.5	567.1
1999 [4]	60.9	*	*	*	*	*	11.7	73.7	241.9	491.0	665.1	580.0
Female												
2001	31.3	*	*	*	*	*	7.3	30.4	94.0	189.4	235.3	181.8
2000 [4]	31.9	*	*	*	*	*	7.4	33.6	97.0	196.9	227.3	188.4
1999 [4]	31.8	*	*	*	*	*	7.9	32.5	96.4	195.5	237.4	186.1

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant melanoma of skin (C43)												
All origins [3]												
Both sexes												
2001	2.6	*	*	*	0.1	0.5	1.4	3.1	5.3	9.2	13.3	17.8
2000 [4]	2.6	*	*	*	0.1	0.5	1.4	3.2	5.4	8.9	13.1	18.9
1999 [4]	2.6	*	*	*	0.1	0.5	1.4	3.1	5.3	9.0	13.4	16.4
Male												
2001	3.4	*	*	*	0.1	0.5	1.7	3.9	7.3	13.9	21.3	29.9
2000 [4]	3.3	*	*	*	0.1	0.5	1.6	4.2	7.4	12.9	20.8	31.8
1999 [4]	3.3	*	*	*	0.1	0.5	1.8	4.0	7.5	13.2	20.9	27.5
Female												
2001	1.9	*	*	*	*	0.4	1.1	2.3	3.3	5.3	8.1	12.8
2000 [4]	2.0	*	*	*	0.1	0.4	1.2	2.2	3.5	5.7	8.2	13.7
1999 [4]	1.9	*	*	*	*	0.4	1.0	2.2	3.3	5.5	8.5	11.9
Hispanic												
Both sexes												
2001	0.4	*	*	*	*	*	0.4	0.7	1.4	2.5	5.0	*
2000 [4]	0.4	*	*	*	*	*	0.5	0.9	1.6	2.4	*	*
1999 [4]	0.4	*	*	*	*	*	*	0.7	1.8	2.6	4.2	*
Male												
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.5	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.5	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.4	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant melanoma of skin (C43)												
Hispanic												
Female												
2000 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	3.0	*	*	*	0.1	0.5	1.5	3.3	5.6	9.7	13.7	18.3
2000 [4]	3.0	*	*	*	0.1	0.5	1.5	3.4	5.7	9.3	13.5	19.3
1999 [4]	2.9	*	*	*	0.1	0.6	1.5	3.3	5.6	9.3	13.7	16.7
Male												
2001	3.9	*	*	*	0.1	0.6	1.9	4.2	7.8	14.6	22.0	31.0
2000 [4]	3.7	*	*	*	0.1	0.6	1.7	4.4	7.8	13.4	21.3	32.6
1999 [4]	3.7	*	*	*	0.1	0.6	2.0	4.2	7.8	13.7	21.5	28.2
Female												
2001	2.1	*	*	*	*	0.5	1.2	2.4	3.5	5.6	8.3	12.9
2000 [4]	2.2	*	*	*	*	0.4	1.3	2.4	3.7	5.9	8.4	13.9
1999 [4]	2.1	*	*	*	*	0.5	1.1	2.3	3.5	5.7	8.7	12.0
Non-Hispanic white												
Both sexes												
2001	3.7	*	*	*	0.2	0.7	1.9	4.0	6.5	10.8	15.0	19.9
2000 [4]	3.6	*	*	*	0.2	0.7	1.8	4.0	6.5	10.4	14.7	20.8
1999 [4]	3.5	*	*	*	0.1	0.7	1.9	3.9	6.4	10.5	14.9	17.9
Male												
2001	4.8	*	*	*	0.2	0.8	2.3	5.0	9.0	16.3	24.1	34.1

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant melanoma of skin (C43)												
Non-Hispanic white												
Male												
2000 [4]	4.6	*	*	*	0.2	0.8	2.0	5.2	8.9	15.0	23.2	35.7
1999 [4]	4.5	*	*	*	*	0.8	2.4	5.0	8.9	15.4	23.3	30.5
Female												
2001	2.6	*	*	*	*	0.6	1.5	2.9	4.1	6.2	9.0	13.9
2000 [4]	2.7	*	*	*	*	0.6	1.6	2.8	4.2	6.6	9.1	14.7
1999 [4]	2.5	*	*	*	*	0.6	1.4	2.8	4.1	6.4	9.4	12.9
Non-Hispanic black												
Both sexes												
2001	0.3	*	*	*	*	*	*	*	*	2.2	2.6	*
2000 [4]	0.4	*	*	*	*	*	*	*	*	1.9	3.6	6.7
1999 [4]	0.3	*	*	*	*	*	*	*	*	1.4	2.8	*
Male												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.4	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.4	*	*	*	*	*	*	*	*	2.1	*	*
2000 [4]	0.4	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of breast (C50)												
All origins [3]												
Both sexes												
2001	14.7	*	*	*	*	1.2	6.3	16.8	30.2	47.6	76.8	134.2
2000 [4]	15.0	*	*	*	*	1.2	6.3	16.9	31.2	49.1	78.9	147.4
1999 [4]	14.9	*	*	*	*	1.1	6.1	16.9	31.1	49.5	78.9	144.4
Male												
2001	0.3	*	*	*	*	*	*	0.2	0.6	1.4	2.3	3.5
2000 [4]	0.3	*	*	*	*	*	*	0.2	0.7	1.4	2.2	4.2
1999 [4]	0.3	*	*	*	*	*	*	0.2	0.6	1.2	2.0	4.5
Female												
2001	28.6	*	*	*	*	2.4	12.4	32.8	57.5	85.8	125.8	188.9
2000 [4]	29.2	*	*	*	*	2.3	12.4	33.0	59.3	88.3	128.9	205.7
1999 [4]	28.9	*	*	*	*	2.2	12.0	32.9	59.2	88.9	128.9	200.8
Hispanic												
Both sexes												
2001	4.8	*	*	*	*	0.8	4.5	12.0	22.0	28.0	39.5	68.8
2000 [4]	4.8	*	*	*	*	0.8	4.3	12.1	21.0	31.3	45.6	71.7
1999 [4]	4.6	*	*	*	*	0.7	4.0	12.2	19.6	28.0	43.5	86.3
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	9.7	*	*	*	*	1.7	9.2	23.8	41.5	49.3	64.7	102.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of breast (C50)												
Hispanic												
Female												
2000 [4]	9.7	*	*	*	*	1.8	8.7	23.9	39.1	54.9	74.9	105.8
1999 [4]	9.3	*	*	*	*	1.6	8.3	23.9	36.6	49.1	71.8	127.9
Non-Hispanic [5]												
Both sexes												
2001	16.1	*	*	*	*	1.2	6.5	17.2	30.7	48.8	78.4	136.4
2000 [4]	16.5	*	*	*	*	1.2	6.5	17.3	31.9	50.1	80.1	149.7
1999 [4]	16.3	*	*	*	*	1.2	6.3	17.3	31.9	50.6	80.2	146.2
Male												
2001	0.3	*	*	*	*	*	*	0.3	0.6	1.5	2.3	3.7
2000 [4]	0.3	*	*	*	*	*	*	0.3	0.7	1.4	2.2	4.2
1999 [4]	0.3	*	*	*	*	*	*	0.2	0.6	1.2	2.1	4.7
Female												
2001	31.1	*	*	*	*	2.5	12.8	33.6	58.6	88.1	128.3	191.6
2000 [4]	31.8	*	*	*	*	2.4	12.8	33.8	60.7	90.3	130.8	208.6
1999 [4]	31.4	*	*	*	*	2.3	12.4	33.6	60.7	91.2	130.9	202.8
Non-Hispanic white												
Both sexes												
2001	17.0	*	*	*	*	1.0	5.8	16.0	29.6	48.2	78.7	137.1
2000 [4]	17.4	*	*	*	*	1.1	5.8	16.1	30.8	50.3	80.6	149.6
1999 [4]	17.1	*	*	*	*	1.0	5.6	16.0	30.6	50.2	80.3	147.8
Male												
2001	0.3	*	*	*	*	*	*	0.2	0.6	1.4	2.2	3.5

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of breast (C50)												
Non-Hispanic white												
Male												
2000 [4]	0.4	*	*	*	*	*	*	0.2	0.7	1.4	2.2	4.0
1999 [4]	0.3	*	*	*	*	*	*	0.2	0.6	1.1	2.0	4.5
Female												
2001	32.9	*	*	*	*	2.0	11.5	31.6	57.0	88.0	129.4	192.6
2000 [4]	33.8	*	*	*	*	2.1	11.6	31.7	59.2	91.4	132.2	208.3
1999 [4]	33.3	*	*	*	*	2.0	11.1	31.5	58.9	91.2	131.5	204.9
Non-Hispanic black												
Both sexes												
2001	15.5	*	*	*	*	2.7	11.5	27.9	45.6	64.6	93.0	153.9
2000 [4]	15.3	*	*	*	*	2.3	11.5	28.4	46.4	59.4	91.7	178.7
1999 [4]	15.5	*	*	*	*	2.4	12.0	29.2	47.1	64.5	95.6	149.1
Male												
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.4	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.4	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	29.2	*	*	*	*	5.1	21.5	51.3	81.6	108.4	143.0	208.3
2000 [4]	28.9	*	*	*	*	4.3	21.5	52.5	82.5	100.1	141.5	241.6
1999 [4]	29.2	*	*	*	*	4.5	22.3	53.7	83.7	108.2	147.2	201.7

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of cervix uteri (C53)												
All origins [3]												
Both sexes												
2001	1.4	*	*	*	*	0.5	1.6	2.4	2.9	3.2	4.6	7.2
2000 [4]	1.5	*	*	*	*	0.6	1.5	2.4	3.1	3.7	4.8	7.3
1999 [4]	1.5	*	*	*	*	0.5	1.6	2.5	3.2	3.8	4.6	7.9
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	2.8	*	*	*	*	1.1	3.1	4.8	5.5	5.8	7.7	10.1
2000 [4]	2.9	*	*	*	*	1.2	3.1	4.8	5.9	6.8	8.0	10.3
1999 [4]	3.0	*	*	*	*	1.1	3.1	4.9	6.2	6.9	7.7	11.1
Hispanic												
Both sexes												
2001	1.1	*	*	*	*	0.5	1.9	3.0	4.1	3.4	7.0	*
2000 [4]	1.2	*	*	*	*	0.5	2.1	2.8	4.6	5.6	8.5	13.9
1999 [4]	1.1	*	*	*	*	0.5	1.8	3.3	3.9	4.9	6.5	*
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	2.3	*	*	*	*	1.1	3.9	6.0	7.7	6.1	11.8	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of cervix uteri (C53)												
Hispanic												
Female												
2000 [4]	2.5	*	*	*	*	1.1	4.4	5.5	8.6	10.0	14.2	20.8
1999 [4]	2.3	*	*	*	*	1.1	3.6	6.6	7.4	8.8	10.8	*
Non-Hispanic [5]												
Both sexes												
2001	1.5	*	*	*	*	0.5	1.5	2.4	2.8	3.2	4.5	7.0
2000 [4]	1.5	*	*	*	*	0.6	1.5	2.4	3.0	3.6	4.7	7.1
1999 [4]	1.6	*	*	*	*	0.5	1.5	2.4	3.2	3.7	4.6	7.9
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	2.9	*	*	*	*	1.0	3.0	4.6	5.3	5.8	7.5	10.0
2000 [4]	3.0	*	*	*	*	1.2	2.9	4.7	5.7	6.6	7.7	9.9
1999 [4]	3.0	*	*	*	*	1.1	3.0	4.7	6.1	6.7	7.5	11.0
Non-Hispanic white												
Both sexes												
2001	1.4	*	*	*	*	0.5	1.4	2.1	2.4	2.6	3.8	6.1
2000 [4]	1.4	*	*	*	*	0.5	1.3	2.1	2.5	2.9	4.1	6.0
1999 [4]	1.4	*	*	*	*	0.6	1.4	2.3	2.7	3.0	3.9	6.7
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of cervix uteri (C53)												
Non-Hispanic white												
Male												
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	2.7	*	*	*	*	1.0	2.8	4.2	4.7	4.9	6.3	8.6
2000 [4]	2.7	*	*	*	*	1.1	2.5	4.2	4.8	5.3	6.8	8.4
1999 [4]	2.8	*	*	*	*	1.1	2.8	4.5	5.2	5.5	6.5	9.3
Non-Hispanic black												
Both sexes												
2001	2.2	*	*	*	*	0.8	2.5	3.8	5.3	7.9	12.0	19.0
2000 [4]	2.5	*	*	*	*	1.0	2.6	4.3	6.4	9.8	12.0	19.7
1999 [4]	2.4	*	*	*	*	0.7	2.5	3.6	7.2	10.8	11.3	21.7
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	4.2	*	*	*	*	1.6	4.6	7.1	9.6	13.5	18.7	26.1
2000 [4]	4.7	*	*	*	*	1.9	5.0	8.0	11.5	16.6	18.8	27.0
1999 [4]	4.5	*	*	*	*	1.2	4.8	6.6	13.0	18.3	17.7	29.8

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)												
All origins [3]												
Both sexes												
2001	2.4	*	*	*	*	0.1	0.3	1.4	4.4	9.7	16.4	24.7
2000 [4]	2.3	*	*	*	*	0.1	0.3	1.3	4.6	9.7	15.8	24.7
1999 [4]	2.3	*	*	*	*	0.1	0.3	1.3	4.3	9.8	16.4	24.0
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	4.7	*	*	*	*	0.2	0.6	2.8	8.4	17.8	27.2	35.0
2000 [4]	4.6	*	*	*	*	0.1	0.6	2.6	8.9	17.7	26.2	34.8
1999 [4]	4.5	*	*	*	*	0.1	0.6	2.5	8.2	17.8	27.1	33.7
Hispanic												
Both sexes												
2001	0.9	*	*	*	*	*	*	1.4	3.8	7.5	11.7	18.8
2000 [4]	0.8	*	*	*	*	*	*	1.1	3.6	7.0	10.7	19.9
1999 [4]	0.8	*	*	*	*	*	*	1.1	3.7	8.3	10.6	15.4
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	1.8	*	*	*	*	*	*	2.8	7.3	13.4	19.6	27.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)												
Hispanic												
Female												
2000 [4]	1.6	*	*	*	*	*	*	2.2	6.8	12.4	17.8	29.7
1999 [4]	1.6	*	*	*	*	*	*	2.2	7.0	14.8	17.7	22.9
Non-Hispanic [5]												
Both sexes												
2001	2.6	*	*	*	*	0.1	0.3	1.4	4.4	9.9	16.6	24.8
2000 [4]	2.6	*	*	*	*	0.1	0.3	1.4	4.7	9.8	16.0	24.9
1999 [4]	2.5	*	*	*	*	0.1	0.3	1.3	4.3	9.9	16.6	24.3
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	5.1	*	*	*	*	0.2	0.6	2.8	8.5	18.1	27.5	35.1
2000 [4]	5.0	*	*	*	*	0.1	0.6	2.7	9.0	17.9	26.5	34.9
1999 [4]	4.9	*	*	*	*	0.1	0.6	2.5	8.3	18.0	27.4	34.1
Non-Hispanic white												
Both sexes												
2001	2.7	*	*	*	*	*	0.3	1.3	4.0	8.9	15.9	24.5
2000 [4]	2.6	*	*	*	*	*	0.3	1.3	4.3	8.6	15.0	24.5
1999 [4]	2.6	*	*	*	*	*	0.3	1.3	4.0	8.9	16.0	23.6
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)												
Non-Hispanic white												
Male												
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	5.2	*	*	*	*	*	0.6	2.7	7.8	16.4	26.5	34.7
2000 [4]	5.1	*	*	*	*	*	0.5	2.7	8.4	15.9	24.8	34.3
1999 [4]	5.1	*	*	*	*	*	0.6	2.5	7.7	16.3	26.4	33.1
Non-Hispanic black												
Both sexes												
2001	2.9	*	*	*	*	*	0.4	2.1	7.7	21.6	28.6	32.8
2000 [4]	3.0	*	*	*	*	*	0.4	1.7	8.4	22.9	31.1	33.7
1999 [4]	2.8	*	*	*	*	*	0.5	1.4	8.1	21.5	27.5	35.9
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	5.5	*	*	*	*	*	0.7	4.0	13.9	36.8	44.6	44.9
2000 [4]	5.7	*	*	*	*	*	0.7	3.1	15.3	39.1	48.6	46.2
1999 [4]	5.4	*	*	*	*	*	0.9	2.7	14.7	36.7	42.9	49.3

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of ovary (C56)												
All origins [3]												
Both sexes												
2001	5.1	*	*	*	0.1	0.3	1.0	4.1	10.3	20.0	33.2	39.6
2000 [4]	5.0	*	*	*	0.1	0.3	1.0	4.2	10.1	19.6	33.2	40.9
1999 [4]	4.9	*	*	*	0.1	0.2	1.1	4.2	10.2	19.6	31.7	37.6
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	9.9	*	*	*	0.1	0.5	2.1	8.1	19.9	36.6	55.0	56.1
2000 [4]	9.8	*	*	*	0.1	0.5	2.1	8.3	19.4	35.7	54.8	57.6
1999 [4]	9.6	*	*	*	0.1	0.4	2.2	8.3	19.6	35.7	52.3	52.8
Hispanic												
Both sexes												
2001	1.6	*	*	*	*	*	0.8	2.7	7.1	14.4	20.0	30.6
2000 [4]	1.6	*	*	*	*	*	0.6	3.0	6.5	14.3	24.9	32.5
1999 [4]	1.4	*	*	*	*	*	0.6	2.7	6.1	12.2	20.8	23.9
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	3.4	*	*	*	*	*	1.6	5.3	13.4	25.8	33.5	45.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of ovary (C56)												
Hispanic												
Female												
2000 [4]	3.4	*	*	*	*	*	1.2	5.9	12.2	25.5	41.6	48.5
1999 [4]	2.9	*	*	*	*	*	1.2	5.3	11.5	21.7	34.7	35.4
Non-Hispanic [5]												
Both sexes												
2001	5.6	*	*	*	0.1	0.3	1.1	4.3	10.5	20.3	33.7	39.8
2000 [4]	5.5	*	*	*	*	0.3	1.1	4.3	10.3	19.9	33.4	41.2
1999 [4]	5.3	*	*	*	0.1	0.2	1.2	4.4	10.5	20.0	32.0	38.1
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	10.9	*	*	*	0.1	0.5	2.1	8.4	20.3	37.3	55.8	56.4
2000 [4]	10.7	*	*	*	*	0.5	2.2	8.4	19.9	36.3	55.2	57.8
1999 [4]	10.4	*	*	*	0.2	0.4	2.3	8.6	20.1	36.4	52.7	53.3
Non-Hispanic white												
Both sexes												
2001	6.2	*	*	*	*	0.3	1.1	4.5	10.8	21.0	34.8	40.6
2000 [4]	6.2	*	*	*	*	0.3	1.1	4.5	10.8	20.5	34.5	42.2
1999 [4]	6.0	*	*	*	0.1	0.2	1.2	4.5	10.9	20.5	32.7	39.2
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of ovary (C56)												
Non-Hispanic white												
Male												
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	12.2	*	*	*	*	0.5	2.2	9.0	21.0	38.8	57.8	57.5
2000 [4]	12.0	*	*	*	*	0.6	2.3	8.9	21.0	37.7	57.2	59.2
1999 [4]	11.7	*	*	*	0.2	0.4	2.4	9.0	21.2	37.7	54.1	54.9
Non-Hispanic black												
Both sexes												
2001	3.2	*	*	*	*	*	1.2	3.1	10.2	18.5	27.2	36.5
2000 [4]	3.0	*	*	*	*	*	1.1	3.4	8.6	18.2	27.1	34.7
1999 [4]	3.1	*	*	*	*	*	0.9	3.6	9.1	18.7	28.7	29.1
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	6.1	*	*	*	*	*	2.2	5.7	18.4	31.5	42.5	50.0
2000 [4]	5.8	*	*	*	*	*	2.0	6.3	15.6	31.1	42.4	47.5
1999 [4]	5.8	*	*	*	*	*	1.6	6.7	16.4	31.9	44.8	40.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of prostate (C61)												
All origins [3]												
Both sexes												
2001	10.8	*	*	*	*	*	0.1	1.1	7.5	35.7	103.2	200.9
2000 [4]	11.0	*	*	*	*	*	0.1	1.0	7.9	37.0	105.7	209.3
1999 [4]	11.4	*	*	*	*	*	0.1	1.0	8.2	39.7	109.9	207.5
Male												
2001	22.0	*	*	*	*	*	0.1	2.1	15.7	78.9	260.2	681.2
2000 [4]	22.5	*	*	*	*	*	0.1	2.0	16.6	82.0	267.9	723.1
1999 [4]	23.2	*	*	*	*	*	0.1	2.1	17.1	88.2	279.1	722.4
Female												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Hispanic												
Both sexes												
2001	3.3	*	*	*	*	*	*	0.8	5.8	29.2	87.8	170.1
2000 [4]	2.9	*	*	*	*	*	*	0.8	4.6	25.7	79.2	165.9
1999 [4]	3.1	*	*	*	*	*	*	0.7	5.6	28.5	80.4	176.1
Male												
2001	6.4	*	*	*	*	*	*	1.6	12.4	66.3	217.8	518.0
2000 [4]	5.7	*	*	*	*	*	*	1.7	9.7	58.5	197.2	503.9
1999 [4]	6.0	*	*	*	*	*	*	1.5	12.0	64.8	201.3	541.3
Female												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of prostate (C61)												
Hispanic												
Female												
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic [5]												
Both sexes												
2001	11.9	*	*	*	*	*	0.1	1.1	7.6	36.1	103.6	201.6
2000 [4]	12.2	*	*	*	*	*	0.1	1.0	8.2	37.6	106.6	210.4
1999 [4]	12.5	*	*	*	*	*	0.1	1.0	8.3	40.3	110.9	208.2
Male												
2001	24.3	*	*	*	*	*	0.1	2.2	15.9	79.5	261.4	686.3
2000 [4]	25.0	*	*	*	*	*	0.1	2.0	17.0	83.2	270.2	730.7
1999 [4]	25.6	*	*	*	*	*	0.2	2.1	17.4	89.4	281.8	728.2
Female												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic white												
Both sexes												
2001	12.0	*	*	*	*	*	*	0.9	6.4	31.9	95.1	189.5
2000 [4]	12.3	*	*	*	*	*	*	0.8	6.9	32.8	97.6	199.3
1999 [4]	12.6	*	*	*	*	*	0.1	0.9	6.9	35.6	102.6	195.4
Male												
2001	24.6	*	*	*	*	*	*	1.8	13.3	69.6	238.4	645.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of prostate (C61)												
Non-Hispanic white												
Male												
2000 [4]	25.2	*	*	*	*	*	*	1.7	14.2	71.7	246.1	693.6
1999 [4]	25.9	*	*	*	*	*	0.1	1.8	14.3	78.2	259.4	685.4
Female												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic black												
Both sexes												
2001	14.6	*	*	*	*	*	*	2.9	19.5	84.2	221.6	365.8
2000 [4]	15.1	*	*	*	*	*	*	2.4	20.8	91.6	230.8	359.6
1999 [4]	15.3	*	*	*	*	*	*	2.7	22.0	94.7	227.5	368.6
Male												
2001	30.8	*	*	*	*	*	*	6.2	43.7	203.5	616.3	1,353.6
2000 [4]	31.7	*	*	*	*	*	*	5.3	46.6	221.7	640.8	1,328.0
1999 [4]	32.2	*	*	*	*	*	*	5.8	49.5	229.1	633.8	1,354.9
Female												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of kidney and renal pelvis (C64-C65)												
All origins [3]												
Both sexes												
2001	4.2	*	0.1	0.1	0.1	0.1	0.8	3.3	9.2	17.6	26.3	32.3
2000 [4]	4.2	*	*	0.1	0.1	0.1	0.7	3.3	9.0	17.1	26.6	33.8
1999 [4]	4.0	*	*	0.1	*	0.1	0.8	3.1	8.9	16.7	24.6	31.4
Male												
2001	5.4	*	*	*	*	0.1	1.1	4.9	13.4	25.2	37.4	52.9
2000 [4]	5.3	*	*	*	*	0.1	1.0	4.8	13.1	24.2	38.2	55.1
1999 [4]	5.0	*	*	*	*	0.2	1.1	4.5	12.9	23.6	35.7	49.4
Female												
2001	3.1	*	*	0.1	*	0.1	0.5	1.9	5.3	11.3	19.0	23.7
2000 [4]	3.1	*	*	*	*	*	0.5	1.9	5.3	11.3	18.9	25.1
1999 [4]	3.0	*	*	*	*	*	0.5	1.8	5.1	11.0	17.4	24.1
Hispanic												
Both sexes												
2001	1.7	*	*	*	*	*	0.7	3.3	6.9	14.4	23.3	31.9
2000 [4]	1.6	*	*	*	*	*	0.6	2.8	7.9	12.3	24.7	25.9
1999 [4]	1.6	*	*	*	*	*	0.6	2.8	7.9	13.9	22.5	28.8
Male												
2001	2.2	*	*	*	*	*	1.2	4.7	9.9	21.4	35.3	57.1
2000 [4]	2.0	*	*	*	*	*	1.0	3.9	11.4	17.9	38.4	40.3
1999 [4]	2.0	*	*	*	*	*	0.9	4.3	11.8	21.1	29.7	*
Female												
2001	1.2	*	*	*	*	*	*	1.9	4.2	8.8	15.1	19.5

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of kidney and renal pelvis (C64-C65)												
Hispanic												
Female												
2000 [4]	1.1	*	*	*	*	*	*	1.8	4.7	7.8	15.5	*
1999 [4]	1.2	*	*	*	*	*	*	1.4	4.4	8.3	17.7	23.9
Non-Hispanic [5]												
Both sexes												
2001	4.6	*	*	0.1	0.1	0.1	0.8	3.3	9.4	17.8	26.4	32.3
2000 [4]	4.5	*	*	0.1	*	0.1	0.8	3.3	9.1	17.3	26.6	33.9
1999 [4]	4.3	*	*	0.1	*	0.1	0.8	3.2	8.9	16.8	24.6	31.4
Male												
2001	5.9	*	*	*	*	0.1	1.0	4.9	13.7	25.4	37.5	52.6
2000 [4]	5.7	*	*	*	*	0.1	1.0	4.8	13.2	24.4	38.1	55.4
1999 [4]	5.4	*	*	*	*	0.2	1.1	4.6	13.0	23.7	35.9	49.8
Female												
2001	3.4	*	*	0.1	*	0.1	0.6	1.8	5.4	11.4	19.1	23.8
2000 [4]	3.4	*	*	*	*	*	0.5	1.9	5.3	11.4	19.1	25.2
1999 [4]	3.2	*	*	*	*	*	0.5	1.8	5.2	11.2	17.3	24.1
Non-Hispanic white												
Both sexes												
2001	5.1	*	*	*	*	0.1	0.7	3.4	9.6	18.4	26.8	32.8
2000 [4]	5.0	*	*	*	*	0.1	0.7	3.4	9.3	17.7	26.9	34.6
1999 [4]	4.7	*	*	0.1	*	0.1	0.7	3.1	9.2	17.2	24.9	31.5
Male												
2001	6.5	*	*	*	*	*	1.0	4.9	13.8	26.2	37.7	54.3

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of kidney and renal pelvis (C64-C65)												
Non-Hispanic white												
Male												
2000 [4]	6.3	*	*	*	*	*	1.0	4.9	13.5	24.6	38.5	56.9
1999 [4]	6.0	*	*	*	*	*	1.0	4.5	13.3	24.0	36.5	50.2
Female												
2001	3.8	*	*	*	*	*	0.5	1.8	5.5	11.8	19.5	23.8
2000 [4]	3.8	*	*	*	*	*	0.5	1.9	5.4	11.8	19.3	25.7
1999 [4]	3.5	*	*	*	*	*	0.5	1.8	5.3	11.5	17.3	24.1
Non-Hispanic black												
Both sexes												
2001	3.1	*	*	*	*	*	1.3	3.8	9.7	15.3	26.0	30.0
2000 [4]	2.9	*	*	*	*	*	0.9	3.5	9.0	16.8	25.9	29.6
1999 [4]	3.0	*	*	*	*	0.4	1.1	3.9	8.7	16.4	23.9	31.7
Male												
2001	4.0	*	*	*	*	*	1.5	5.5	15.3	22.7	40.8	46.2
2000 [4]	3.7	*	*	*	*	*	1.0	5.0	13.4	26.8	38.3	49.3
1999 [4]	3.7	*	*	*	*	*	1.6	6.0	13.3	23.9	32.5	49.9
Female												
2001	2.3	*	*	*	*	*	1.1	2.2	5.3	10.0	17.7	24.0
2000 [4]	2.2	*	*	*	*	*	0.8	2.2	5.4	9.7	18.9	22.2
1999 [4]	2.2	*	*	*	*	*	*	2.1	5.0	11.0	19.1	24.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of bladder (C67)												
All origins [3]												
Both sexes												
2001	4.3	*	*	*	*	*	0.3	1.2	5.0	15.4	35.7	68.9
2000 [4]	4.3	*	*	*	*	*	0.2	1.2	5.0	15.5	35.7	69.6
1999 [4]	4.3	*	*	*	*	*	0.2	1.3	5.2	15.9	35.3	68.5
Male												
2001	6.0	*	*	*	*	*	0.3	1.8	7.9	25.2	62.5	134.8
2000 [4]	5.9	*	*	*	*	*	0.3	1.7	7.9	25.2	62.6	139.4
1999 [4]	5.9	*	*	*	*	*	0.3	2.0	8.2	25.7	62.5	134.8
Female												
2001	2.7	*	*	*	*	*	0.3	0.6	2.3	7.2	18.1	41.4
2000 [4]	2.7	*	*	*	*	*	0.2	0.6	2.4	7.5	18.1	41.2
1999 [4]	2.7	*	*	*	*	*	0.2	0.6	2.5	7.8	17.7	41.8
Hispanic												
Both sexes												
2001	0.9	*	*	*	*	*	*	0.6	2.2	7.9	18.8	46.3
2000 [4]	0.9	*	*	*	*	*	*	*	2.9	6.6	17.6	47.1
1999 [4]	0.9	*	*	*	*	*	*	0.7	3.1	8.5	16.5	38.6
Male												
2001	1.2	*	*	*	*	*	*	*	4.3	11.8	30.0	87.6
2000 [4]	1.2	*	*	*	*	*	*	*	4.6	10.6	33.9	84.6
1999 [4]	1.3	*	*	*	*	*	*	*	5.6	15.2	27.1	77.6
Female												
2001	0.6	*	*	*	*	*	*	*	*	4.9	11.2	26.1

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of bladder (C67)												
Hispanic												
Female												
2000 [4]	0.5	*	*	*	*	*	*	*	*	3.5	6.6	28.7
1999 [4]	0.5	*	*	*	*	*	*	*	*	*	9.4	*
Non-Hispanic [5]												
Both sexes												
2001	4.8	*	*	*	*	*	0.3	1.3	5.2	15.8	36.4	69.6
2000 [4]	4.7	*	*	*	*	*	0.3	1.2	5.2	16.1	36.3	70.3
1999 [4]	4.7	*	*	*	*	*	0.2	1.3	5.4	16.3	36.0	69.3
Male												
2001	6.7	*	*	*	*	*	0.4	1.9	8.1	26.0	63.8	136.4
2000 [4]	6.6	*	*	*	*	*	0.3	1.8	8.1	26.1	63.6	141.3
1999 [4]	6.6	*	*	*	*	*	0.3	2.0	8.4	26.3	63.8	136.6
Female												
2001	3.0	*	*	*	*	*	0.3	0.6	2.5	7.4	18.3	41.9
2000 [4]	3.0	*	*	*	*	*	0.2	0.6	2.5	7.8	18.5	41.6
1999 [4]	3.0	*	*	*	*	*	0.2	0.6	2.6	8.1	17.9	42.3
Non-Hispanic white												
Both sexes												
2001	5.5	*	*	*	*	*	0.3	1.3	5.4	16.6	37.4	71.5
2000 [4]	5.4	*	*	*	*	*	0.3	1.3	5.5	16.8	37.1	72.8
1999 [4]	5.4	*	*	*	*	*	0.2	1.3	5.5	16.8	37.1	70.7
Male												
2001	7.8	*	*	*	*	*	0.4	2.0	8.6	27.6	66.6	143.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of bladder (C67)												
Non-Hispanic white												
Male												
2000 [4]	7.7	*	*	*	*	*	0.3	2.0	8.6	27.5	65.9	148.0
1999 [4]	7.6	*	*	*	*	*	0.3	2.0	8.7	27.6	66.2	143.0
Female												
2001	3.2	*	*	*	*	*	0.3	0.6	2.4	7.4	18.1	41.7
2000 [4]	3.3	*	*	*	*	*	0.2	0.7	2.6	7.8	18.2	42.5
1999 [4]	3.2	*	*	*	*	*	0.2	0.7	2.5	7.8	18.0	41.9
Non-Hispanic black												
Both sexes												
2001	2.4	*	*	*	*	*	0.4	1.3	5.1	12.1	31.4	55.6
2000 [4]	2.4	*	*	*	*	*	*	1.1	4.2	13.5	34.9	49.6
1999 [4]	2.6	*	*	*	*	*	*	1.5	5.6	15.2	29.4	60.8
Male												
2001	2.6	*	*	*	*	*	*	1.8	6.5	16.5	43.2	78.5
2000 [4]	2.8	*	*	*	*	*	*	1.4	5.9	17.8	52.8	88.1
1999 [4]	3.0	*	*	*	*	*	*	2.6	7.4	19.2	46.6	86.8
Female												
2001	2.3	*	*	*	*	*	*	*	3.9	9.0	24.8	47.1
2000 [4]	2.1	*	*	*	*	*	*	*	2.7	10.4	24.8	35.3
1999 [4]	2.3	*	*	*	*	*	*	*	4.2	12.3	19.8	51.1

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
All origins [3]												
Both sexes												
2001	4.4	*	0.7	0.7	0.6	1.1	2.3	5.0	9.7	15.4	20.4	16.4
2000 [4]	4.5	0.5	0.7	0.8	0.6	1.0	2.4	5.1	9.6	16.5	20.0	16.9
1999 [4]	4.6	*	0.7	0.8	0.6	1.1	2.4	5.1	10.3	16.9	20.4	16.0
Male												
2001	5.0	*	0.7	0.7	0.6	1.3	2.8	6.1	12.1	19.2	25.8	20.1
2000 [4]	5.1	*	0.7	0.8	0.7	1.3	3.1	6.2	11.8	20.2	24.3	22.4
1999 [4]	5.1	*	0.7	0.7	0.7	1.4	2.8	6.2	12.4	20.8	25.5	20.7
Female												
2001	3.9	*	0.6	0.8	0.5	0.8	1.7	3.9	7.6	12.3	16.8	14.8
2000 [4]	3.9	*	0.6	0.7	0.5	0.7	1.7	4.0	7.6	13.4	17.3	14.7
1999 [4]	4.1	*	0.8	0.8	0.5	0.8	1.9	4.1	8.4	13.7	17.1	14.1
Hispanic												
Both sexes												
2001	1.7	*	0.7	0.5	0.4	0.6	1.4	2.6	5.9	10.3	14.8	*
2000 [4]	1.8	*	*	0.9	0.4	0.7	1.5	2.9	6.2	9.8	14.2	15.9
1999 [4]	1.7	*	0.9	0.7	*	0.5	1.3	3.0	6.0	10.0	14.0	*
Male												
2001	1.8	*	*	*	*	0.6	1.4	3.0	6.8	14.4	17.9	*
2000 [4]	1.9	*	*	1.2	*	0.8	1.5	2.6	7.7	12.5	18.7	*
1999 [4]	1.7	*	*	0.7	*	*	1.8	2.9	7.8	10.5	16.2	*
Female												
2001	1.6	*	*	0.6	*	*	1.4	2.2	5.1	7.1	12.7	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
Hispanic												
Female												
2000 [4]	1.7	*	*	0.7	*	*	1.5	3.2	4.9	7.6	11.2	*
1999 [4]	1.7	*	*	0.8	*	*	0.8	3.1	4.5	9.6	12.5	*
Non-Hispanic [5]												
Both sexes												
2001	4.8	*	0.6	0.8	0.6	1.2	2.4	5.2	10.0	15.7	20.6	16.5
2000 [4]	4.9	*	0.7	0.7	0.6	1.1	2.5	5.3	9.8	16.9	20.2	16.9
1999 [4]	5.0	*	0.7	0.8	0.6	1.2	2.5	5.3	10.6	17.3	20.6	16.1
Male												
2001	5.5	*	0.6	0.7	0.7	1.5	3.0	6.4	12.5	19.4	26.1	20.5
2000 [4]	5.5	*	0.7	0.8	0.7	1.4	3.3	6.5	12.1	20.6	24.5	22.3
1999 [4]	5.6	*	0.7	0.8	0.7	1.5	3.0	6.5	12.7	21.4	25.8	21.3
Female												
2001	4.2	*	0.6	0.8	0.5	0.8	1.8	4.1	7.7	12.7	17.0	14.9
2000 [4]	4.2	*	0.7	0.7	0.5	0.7	1.8	4.1	7.8	13.8	17.5	14.7
1999 [4]	4.4	*	0.7	0.8	0.5	0.9	2.0	4.2	8.6	14.0	17.2	14.0
Non-Hispanic white												
Both sexes												
2001	5.6	*	0.7	0.8	0.6	1.3	2.7	5.7	10.8	16.8	21.9	17.2
2000 [4]	5.6	*	0.7	0.8	0.6	1.2	2.8	5.9	10.7	18.3	21.2	17.5
1999 [4]	5.7	*	0.7	0.8	0.6	1.4	2.8	5.8	11.5	18.7	21.7	16.7
Male												
2001	6.3	*	0.7	0.7	0.7	1.7	3.5	7.0	13.4	20.9	27.5	20.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
Non-Hispanic white												
Male												
2000 [4]	6.3	*	0.7	0.8	0.7	1.5	3.7	7.1	13.1	22.2	25.7	23.5
1999 [4]	6.4	*	0.8	0.8	0.7	1.8	3.3	7.1	13.7	23.0	27.2	22.4
Female												
2001	4.8	*	0.6	0.9	0.5	0.9	2.0	4.5	8.4	13.4	18.2	15.7
2000 [4]	4.9	*	0.7	0.7	0.5	0.9	1.9	4.6	8.4	15.0	18.3	15.1
1999 [4]	5.0	*	0.7	0.7	0.5	1.1	2.3	4.6	9.5	15.1	18.1	14.4
Non-Hispanic black												
Both sexes												
2001	2.1	*	*	0.7	0.5	0.5	1.1	3.1	5.9	9.4	9.9	11.9
2000 [4]	2.2	*	*	0.8	0.6	0.7	1.6	2.9	5.8	8.2	12.1	10.5
1999 [4]	2.2	*	*	0.8	0.7	0.7	1.4	3.3	6.1	8.2	11.7	10.0
Male												
2001	2.2	*	*	0.8	*	*	1.3	3.7	7.9	10.4	12.0	*
2000 [4]	2.4	*	*	0.9	0.8	1.1	2.0	3.7	7.2	9.9	13.3	*
1999 [4]	2.4	*	*	0.7	0.9	0.9	1.8	4.5	7.4	10.2	12.5	*
Female												
2001	1.9	*	*	*	*	*	1.0	2.5	4.2	8.7	8.8	9.8
2000 [4]	1.9	*	*	0.7	*	*	1.2	2.2	4.7	7.1	11.5	10.5
1999 [4]	1.9	*	*	0.9	*	*	1.0	2.2	5.1	6.8	11.2	9.3

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
All origins [3]												
Both sexes												
2001	19.8	0.7	1.0	0.9	1.8	2.5	4.1	10.4	29.5	74.6	145.2	199.9
2000 [4]	20.1	1.0	0.9	0.9	1.8	2.3	4.1	10.7	30.0	76.3	149.2	204.6
1999 [4]	20.2	0.8	0.9	0.9	1.9	2.4	4.2	10.8	31.8	77.8	148.0	199.4
Male												
2001	21.5	*	1.0	1.0	2.1	2.9	5.1	12.7	36.3	94.7	187.0	275.3
2000 [4]	21.6	1.1	1.0	1.1	2.3	2.7	4.8	13.0	36.1	96.8	192.6	281.8
1999 [4]	21.7	*	1.0	1.0	2.2	2.8	5.1	13.2	38.8	97.3	189.0	279.0
Female												
2001	18.1	*	1.0	0.7	1.4	2.1	3.2	8.1	23.1	57.9	117.7	168.4
2000 [4]	18.6	*	0.9	0.7	1.4	1.9	3.4	8.5	24.4	59.5	120.9	173.1
1999 [4]	18.7	*	0.8	0.7	1.5	2.0	3.4	8.5	25.3	61.9	121.3	167.3
Hispanic												
Both sexes												
2001	7.4	*	1.3	1.2	2.2	2.2	3.9	8.2	23.9	52.6	103.3	134.4
2000 [4]	7.5	*	1.0	1.1	2.0	1.9	3.9	8.9	25.0	58.8	103.3	143.3
1999 [4]	7.5	*	1.0	1.2	2.0	2.1	3.9	10.1	24.2	58.5	101.0	131.9
Male												
2001	8.3	*	1.4	1.3	2.7	2.7	5.0	9.7	28.9	67.5	142.2	180.9
2000 [4]	8.0	*	*	1.4	2.3	2.4	4.3	11.1	30.7	71.3	126.9	197.5
1999 [4]	7.6	*	*	1.3	2.1	2.2	4.2	12.0	29.4	68.5	117.8	135.9
Female												
2001	6.5	*	*	1.1	1.5	1.7	2.6	6.7	19.5	40.8	77.1	111.7

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
Hispanic												
Female												
2000 [4]	6.9	*	*	0.8	1.5	1.4	3.4	6.8	19.8	48.9	87.5	116.7
1999 [4]	7.4	*	*	1.0	1.9	2.1	3.4	8.3	19.6	50.6	89.9	130.0
Non-Hispanic [5]												
Both sexes												
2001	21.6	0.6	0.9	0.8	1.7	2.5	4.1	10.5	29.8	75.8	146.8	202.1
2000 [4]	21.8	1.0	0.9	0.9	1.8	2.4	4.1	10.8	30.3	77.3	150.9	206.5
1999 [4]	21.9	0.7	0.9	0.8	1.8	2.4	4.3	10.8	32.3	78.8	149.5	201.4
Male												
2001	23.5	*	1.0	1.0	1.9	2.9	5.0	12.9	36.8	96.3	188.6	278.8
2000 [4]	23.7	*	1.0	1.0	2.2	2.7	4.8	13.1	36.4	98.1	195.2	285.0
1999 [4]	23.7	*	0.9	0.9	2.2	2.9	5.2	13.3	39.5	98.7	191.5	284.3
Female												
2001	19.7	*	0.9	0.7	1.4	2.2	3.3	8.2	23.3	58.9	119.3	170.2
2000 [4]	20.1	*	0.9	0.7	1.3	2.0	3.4	8.6	24.7	60.1	122.1	174.8
1999 [4]	20.1	*	0.8	0.7	1.4	2.0	3.4	8.5	25.6	62.5	122.3	168.2
Non-Hispanic white												
Both sexes												
2001	23.8	*	1.1	0.8	1.7	2.4	4.0	10.3	29.5	77.2	150.5	208.2
2000 [4]	24.2	1.3	1.0	0.9	1.7	2.3	4.0	10.5	30.3	79.0	155.1	210.7
1999 [4]	24.1	*	1.0	0.8	1.8	2.4	4.2	10.6	32.0	80.4	153.0	206.4
Male												
2001	25.9	*	1.0	0.8	2.0	2.8	4.7	12.8	36.4	97.9	193.6	288.7

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
Non-Hispanic white												
Male												
2000 [4]	26.2	*	1.1	1.0	2.1	2.6	4.8	12.7	36.3	100.1	201.0	293.8
1999 [4]	26.1	*	1.0	0.9	2.2	2.7	5.0	13.1	39.2	100.7	196.1	291.8
Female												
2001	21.8	*	1.1	0.7	1.4	2.0	3.2	7.9	23.0	59.5	122.0	174.6
2000 [4]	22.3	*	0.9	0.8	1.2	1.9	3.2	8.4	24.6	61.2	125.0	177.2
1999 [4]	22.2	*	0.9	0.6	1.3	2.0	3.3	8.3	25.2	63.4	124.7	172.4
Non-Hispanic black												
Both sexes												
2001	14.5	*	*	1.0	1.8	3.5	5.8	13.5	37.1	79.0	127.6	157.3
2000 [4]	14.3	*	*	0.7	2.3	3.0	5.4	15.0	36.5	76.2	122.3	177.1
1999 [4]	14.5	*	*	0.9	2.4	3.0	5.6	14.1	39.5	77.5	128.2	162.3
Male												
2001	15.6	*	*	1.3	2.0	4.0	7.6	16.3	46.6	100.7	161.4	206.7
2000 [4]	15.2	*	*	0.8	2.8	3.7	5.7	18.2	45.3	97.8	157.4	226.6
1999 [4]	15.6	*	*	1.2	2.6	3.9	7.0	17.5	48.4	96.9	162.9	221.1
Female												
2001	13.5	*	*	*	1.7	3.1	4.2	11.1	29.5	63.7	108.7	139.0
2000 [4]	13.5	*	*	*	1.8	2.5	5.2	12.2	29.4	61.0	102.6	158.7
1999 [4]	13.6	*	*	0.7	2.1	2.2	4.3	11.1	32.3	63.8	108.7	140.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hodgkin's disease (C81)												
All origins [3]												
Both sexes												
2001	0.5	*	*	*	0.2	0.4	0.4	0.5	0.6	1.2	2.2	2.2
2000 [4]	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7	1.3	2.1	2.5
1999 [4]	0.5	*	*	*	0.2	0.4	0.4	0.5	0.7	1.5	2.1	2.4
Male												
2001	0.5	*	*	*	0.2	0.4	0.5	0.6	0.7	1.6	2.6	2.6
2000 [4]	0.5	*	*	*	0.2	0.4	0.4	0.5	0.7	1.6	2.5	3.7
1999 [4]	0.6	*	*	*	0.3	0.4	0.5	0.5	1.0	1.9	2.5	3.8
Female												
2001	0.4	*	*	*	0.2	0.4	0.3	0.4	0.4	0.8	2.0	2.0
2000 [4]	0.4	*	*	*	0.2	0.3	0.3	0.3	0.6	1.0	1.8	2.0
1999 [4]	0.4	*	*	*	0.2	0.3	0.3	0.4	0.5	1.2	1.9	1.8
Hispanic												
Both sexes												
2001	0.3	*	*	*	*	*	*	*	*	1.8	*	*
2000 [4]	0.3	*	*	*	*	*	*	*	*	*	4.5	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	2.5	*	*
Male												
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.4	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hodgkin's disease (C81)												
Hispanic												
Female												
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.5	*	*	*	0.2	0.5	0.4	0.5	0.5	1.1	2.2	2.2
2000 [4]	0.5	*	*	*	0.2	0.4	0.4	0.4	0.6	1.3	1.9	2.4
1999 [4]	0.5	*	*	*	0.3	0.4	0.4	0.4	0.7	1.5	2.1	2.3
Male												
2001	0.5	*	*	*	0.2	0.5	0.5	0.6	0.7	1.5	2.6	2.5
2000 [4]	0.5	*	*	*	0.2	0.4	0.4	0.5	0.7	1.6	2.3	3.7
1999 [4]	0.6	*	*	*	0.3	0.5	0.5	0.5	1.0	1.9	2.3	3.7
Female												
2001	0.4	*	*	*	0.2	0.4	0.3	0.4	0.5	0.8	2.0	2.0
2000 [4]	0.4	*	*	*	0.2	0.4	0.3	0.3	0.6	1.0	1.7	2.0
1999 [4]	0.5	*	*	*	0.2	0.3	0.3	0.4	0.5	1.2	1.9	1.7
Non-Hispanic white												
Both sexes												
2001	0.5	*	*	*	0.2	0.5	0.4	0.5	0.6	1.2	2.3	2.3
2000 [4]	0.5	*	*	*	0.2	0.5	0.4	0.4	0.6	1.3	2.1	2.6
1999 [4]	0.6	*	*	*	0.3	0.4	0.4	0.5	0.8	1.5	2.2	2.3
Male												
2001	0.6	*	*	*	0.3	0.5	0.5	0.6	0.7	1.6	2.7	2.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hodgkin's disease (C81)												
Non-Hispanic white												
Male												
2000 [4]	0.6	*	*	*	0.2	0.5	0.4	0.5	0.7	1.6	2.5	3.9
1999 [4]	0.7	*	*	*	0.3	0.5	0.5	0.5	1.1	2.0	2.5	3.8
Female												
2001	0.5	*	*	*	0.2	0.4	0.3	0.4	0.5	0.9	2.0	2.3
2000 [4]	0.5	*	*	*	0.2	0.4	0.3	0.4	0.6	1.1	1.9	2.1
1999 [4]	0.5	*	*	*	0.2	0.4	0.3	0.4	0.5	1.2	2.0	1.7
Non-Hispanic black												
Both sexes												
2001	0.4	*	*	*	*	0.8	0.4	*	*	*	*	*
2000 [4]	0.3	*	*	*	*	*	0.4	*	*	*	*	*
1999 [4]	0.4	*	*	*	*	0.5	*	*	*	*	*	*
Male												
2001	0.4	*	*	*	*	0.8	*	*	*	*	*	*
2000 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.4	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)												
All origins [3]												
Both sexes												
2001	7.8	*	*	0.1	0.4	0.8	1.7	4.4	12.2	29.3	58.6	79.6
2000 [4]	8.1	*	*	0.1	0.5	0.7	1.7	4.6	13.1	30.4	61.0	80.2
1999 [4]	8.2	*	*	0.1	0.4	0.7	1.8	4.8	13.3	31.7	61.8	77.7
Male												
2001	8.4	*	*	0.1	0.5	0.9	2.2	5.8	15.0	36.5	71.3	106.6
2000 [4]	8.6	*	*	0.1	0.6	0.9	2.1	6.0	15.9	37.4	74.5	105.9
1999 [4]	8.6	*	*	0.1	0.5	0.9	2.3	6.2	16.2	38.3	74.7	104.4
Female												
2001	7.3	*	*	*	0.3	0.6	1.1	3.1	9.7	23.3	50.2	68.3
2000 [4]	7.6	*	*	*	0.3	0.6	1.2	3.3	10.5	24.7	52.1	69.8
1999 [4]	7.7	*	*	*	0.3	0.6	1.2	3.4	10.6	26.4	53.4	66.9
Hispanic												
Both sexes												
2001	2.7	*	*	*	0.3	0.5	1.4	3.6	10.9	18.3	43.3	59.4
2000 [4]	2.9	*	*	*	0.3	0.6	1.6	3.6	11.1	25.7	41.5	59.7
1999 [4]	2.8	*	*	*	*	0.6	1.4	4.4	9.3	25.2	42.7	56.1
Male												
2001	3.0	*	*	*	*	0.6	2.0	4.4	13.2	24.4	58.6	68.6
2000 [4]	3.1	*	*	*	*	0.7	1.9	4.6	13.9	30.2	48.2	82.6
1999 [4]	2.8	*	*	*	*	*	1.7	5.7	11.8	27.5	48.0	60.4
Female												
2001	2.4	*	*	*	*	*	0.8	2.8	8.8	13.5	32.9	54.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)												
Hispanic												
Female												
2000 [4]	2.7	*	*	*	*	*	1.3	2.6	8.6	22.2	37.0	48.5
1999 [4]	2.8	*	*	*	*	*	1.0	3.1	7.1	23.4	39.2	54.1
Non-Hispanic [5]												
Both sexes												
2001	8.6	*	*	0.1	0.4	0.8	1.7	4.5	12.3	29.9	59.2	80.3
2000 [4]	8.8	*	*	0.1	0.5	0.8	1.7	4.7	13.2	30.7	61.7	80.9
1999 [4]	8.9	*	*	0.1	0.4	0.7	1.8	4.8	13.5	32.1	62.4	78.3
Male												
2001	9.2	*	*	0.1	0.5	0.9	2.3	5.9	15.1	37.2	71.7	108.1
2000 [4]	9.4	*	*	0.2	0.7	0.9	2.2	6.1	15.9	37.7	75.6	106.7
1999 [4]	9.4	*	*	0.2	0.5	0.9	2.3	6.2	16.5	38.8	75.5	106.0
Female												
2001	8.0	*	*	*	0.3	0.6	1.2	3.1	9.7	23.8	50.9	68.7
2000 [4]	8.3	*	*	*	0.3	0.6	1.2	3.4	10.6	24.8	52.6	70.4
1999 [4]	8.4	*	*	*	0.3	0.5	1.3	3.4	10.8	26.5	53.9	67.2
Non-Hispanic white												
Both sexes												
2001	9.7	*	*	0.1	0.3	0.7	1.7	4.5	12.7	31.4	62.3	84.1
2000 [4]	10.1	*	*	0.1	0.4	0.7	1.7	4.8	13.8	32.3	65.1	85.1
1999 [4]	10.1	*	*	0.1	0.4	0.7	1.7	4.9	14.1	33.9	65.6	82.0
Male												
2001	10.4	*	*	*	0.4	0.9	2.1	6.0	15.5	38.8	75.2	113.1

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)												
Non-Hispanic white												
Male												
2000 [4]	10.6	*	*	0.2	0.6	0.8	2.2	6.1	16.5	39.5	79.6	113.0
1999 [4]	10.6	*	*	0.2	0.5	0.8	2.2	6.3	17.0	40.8	78.9	110.0
Female												
2001	9.2	*	*	*	0.3	0.5	1.2	3.1	10.1	25.2	53.7	72.0
2000 [4]	9.5	*	*	*	0.3	0.6	1.1	3.4	11.3	26.3	55.6	73.8
1999 [4]	9.6	*	*	*	0.2	0.5	1.2	3.5	11.4	28.1	56.9	70.8
Non-Hispanic black												
Both sexes												
2001	4.1	*	*	*	0.6	1.2	2.3	5.1	10.9	21.3	30.6	40.0
2000 [4]	3.9	*	*	*	0.6	1.1	2.1	5.0	10.8	19.7	27.1	36.6
1999 [4]	4.1	*	*	*	0.6	1.1	2.4	4.8	11.2	20.5	32.1	40.4
Male												
2001	4.7	*	*	*	0.8	1.3	3.4	6.7	13.7	28.2	37.2	55.4
2000 [4]	4.3	*	*	*	*	1.2	2.6	7.0	14.8	25.3	31.8	43.4
1999 [4]	4.7	*	*	*	*	1.6	3.3	6.6	14.8	26.2	41.0	59.4
Female												
2001	3.7	*	*	*	*	1.1	1.4	3.7	8.6	16.5	27.0	34.2
2000 [4]	3.5	*	*	*	*	0.9	1.7	3.3	7.5	15.7	24.5	34.0
1999 [4]	3.6	*	*	*	*	*	1.7	3.2	8.2	16.5	27.1	33.3

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leukemia (C91-C95)												
All origins [3]												
Both sexes												
2001	7.5	0.6	1.0	0.8	1.2	1.3	1.7	3.8	10.0	26.7	53.3	81.4
2000 [4]	7.6	1.0	0.9	0.8	1.2	1.2	1.8	3.7	9.9	27.2	54.5	83.6
1999 [4]	7.5	0.7	0.8	0.7	1.2	1.2	1.8	3.7	10.6	27.2	52.8	82.4
Male												
2001	8.5	*	1.0	0.9	1.4	1.5	1.9	4.2	12.8	35.7	74.2	117.5
2000 [4]	8.5	1.1	0.9	0.9	1.5	1.4	1.9	4.3	12.5	37.2	75.7	119.8
1999 [4]	8.4	*	0.8	0.8	1.5	1.4	1.9	4.3	13.1	36.8	73.2	119.2
Female												
2001	6.6	*	1.0	0.7	1.0	1.1	1.5	3.3	7.5	19.2	39.5	66.3
2000 [4]	6.7	*	0.9	0.6	0.9	1.0	1.7	3.2	7.6	19.0	40.6	68.8
1999 [4]	6.7	*	0.7	0.7	1.0	1.1	1.6	3.2	8.3	19.3	39.6	67.6
Hispanic												
Both sexes												
2001	3.0	*	1.3	1.1	1.7	1.5	1.9	2.9	6.5	17.3	33.4	42.5
2000 [4]	2.9	*	0.9	1.1	1.5	1.1	1.9	3.5	7.8	16.9	32.6	49.1
1999 [4]	2.9	*	0.9	1.1	1.6	1.4	2.0	3.4	7.0	17.3	31.0	44.2
Male												
2001	3.4	*	1.4	1.1	2.2	1.8	2.3	3.3	8.4	19.6	45.6	68.6
2000 [4]	3.3	*	*	1.3	1.7	1.5	2.0	4.1	10.2	20.7	42.8	76.6
1999 [4]	3.1	*	*	1.1	1.7	1.4	1.9	3.6	8.6	23.5	39.1	49.6
Female												
2001	2.6	*	*	1.1	1.1	1.1	1.4	2.6	4.8	15.6	25.1	29.8

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leukemia (C91-C95)												
Hispanic												
Female												
2000 [4]	2.6	*	*	0.8	1.3	*	1.8	2.9	5.6	13.9	25.8	35.6
1999 [4]	2.8	*	*	1.0	1.5	1.3	2.1	3.2	5.6	12.4	25.7	41.6
Non-Hispanic [5]												
Both sexes												
2001	8.2	*	0.9	0.7	1.1	1.2	1.7	3.8	10.3	27.2	54.1	82.7
2000 [4]	8.2	1.0	0.9	0.7	1.1	1.2	1.7	3.8	10.1	27.8	55.3	84.7
1999 [4]	8.2	*	0.7	0.7	1.2	1.2	1.7	3.8	10.8	27.8	53.6	83.7
Male												
2001	9.3	*	0.9	0.8	1.2	1.4	1.9	4.3	13.1	36.6	75.4	119.3
2000 [4]	9.3	*	0.9	0.8	1.4	1.4	1.9	4.3	12.6	38.2	77.0	121.5
1999 [4]	9.2	*	0.8	0.8	1.4	1.3	1.9	4.4	13.4	37.5	74.6	122.1
Female												
2001	7.2	*	0.9	0.6	0.9	1.1	1.5	3.4	7.7	19.4	40.1	67.5
2000 [4]	7.2	*	0.9	0.6	0.8	1.0	1.6	3.3	7.7	19.3	41.1	69.8
1999 [4]	7.2	*	0.7	0.6	0.9	1.1	1.5	3.2	8.5	19.8	40.0	68.3
Non-Hispanic white												
Both sexes												
2001	9.1	*	1.0	0.7	1.1	1.2	1.7	3.8	10.5	28.1	56.0	86.1
2000 [4]	9.2	1.3	0.9	0.7	1.1	1.1	1.7	3.7	10.2	29.0	57.5	87.3
1999 [4]	9.1	*	0.8	0.7	1.1	1.2	1.7	3.7	11.0	28.6	55.3	86.8
Male												
2001	10.3	*	1.0	0.7	1.3	1.4	1.8	4.3	13.3	37.7	78.0	125.3

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leukemia (C91-C95)												
Non-Hispanic white												
Male												
2000 [4]	10.4	*	1.0	0.8	1.3	1.3	1.9	4.2	12.7	39.5	79.9	125.5
1999 [4]	10.2	*	0.8	0.7	1.4	1.3	1.9	4.3	13.6	38.6	77.0	127.2
Female												
2001	8.0	*	1.1	0.6	0.9	1.0	1.6	3.3	7.8	19.9	41.4	69.8
2000 [4]	8.1	*	0.9	0.6	0.8	0.9	1.5	3.2	7.8	20.1	42.7	72.0
1999 [4]	8.0	*	0.9	0.6	0.8	1.1	1.5	3.2	8.5	20.3	41.1	70.8
Non-Hispanic black												
Both sexes												
2001	4.9	*	*	0.9	1.0	1.5	2.1	4.3	10.1	25.0	42.1	57.1
2000 [4]	4.9	*	*	0.6	1.4	1.7	2.1	5.0	10.8	23.1	38.8	66.1
1999 [4]	4.9	*	*	0.8	1.5	1.3	1.9	4.4	10.8	25.2	41.6	55.9
Male												
2001	5.4	*	*	1.2	1.1	1.8	2.6	4.6	13.4	34.4	56.1	75.1
2000 [4]	5.5	*	*	0.7	1.9	2.0	2.0	5.6	14.0	32.3	55.8	98.6
1999 [4]	5.4	*	*	1.0	1.6	1.5	2.1	5.7	13.7	33.4	55.7	78.4
Female												
2001	4.4	*	*	*	1.0	1.1	1.7	4.0	7.4	18.4	34.2	50.5
2000 [4]	4.4	*	*	*	0.9	1.4	2.2	4.4	8.3	16.5	29.2	54.1
1999 [4]	4.4	*	*	0.6	1.3	1.1	1.7	3.3	8.5	19.4	33.8	47.5

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Multiple myeloma and immunoproliferative neoplasms (C88,C90)												
All origins [3]												
Both sexes												
2001	3.9	*	*	*	*	*	0.3	1.7	6.6	17.4	30.9	36.3
2000 [4]	3.9	*	*	*	*	*	0.3	1.9	6.3	17.3	31.5	37.8
1999 [4]	3.9	*	*	*	*	0.1	0.3	1.8	7.1	17.2	31.0	36.6
Male												
2001	4.1	*	*	*	*	*	0.4	2.1	7.8	20.8	38.6	48.0
2000 [4]	4.0	*	*	*	*	*	0.4	2.1	6.9	20.4	39.6	51.8
1999 [4]	4.1	*	*	*	*	*	0.4	2.1	8.5	20.1	38.3	51.3
Female												
2001	3.8	*	*	*	*	*	0.3	1.3	5.5	14.5	25.8	31.3
2000 [4]	3.8	*	*	*	*	*	0.3	1.6	5.7	14.7	26.3	32.2
1999 [4]	3.8	*	*	*	*	*	0.3	1.5	5.9	14.8	26.2	30.6
Hispanic												
Both sexes												
2001	1.4	*	*	*	*	*	*	1.3	5.8	14.9	24.7	29.4
2000 [4]	1.4	*	*	*	*	*	*	1.4	5.3	15.0	24.5	31.2
1999 [4]	1.4	*	*	*	*	*	*	1.7	6.8	13.4	23.1	28.1
Male												
2001	1.5	*	*	*	*	*	*	1.4	6.0	20.4	35.8	38.1
2000 [4]	1.3	*	*	*	*	*	*	1.5	5.5	18.4	30.0	*
1999 [4]	1.3	*	*	*	*	*	*	1.9	7.8	14.7	24.5	*
Female												
2001	1.3	*	*	*	*	*	*	1.2	5.6	10.6	17.2	25.1

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Multiple myeloma and immunoproliferative neoplasms (C88,C90)												
Hispanic												
Female												
2000 [4]	1.4	*	*	*	*	*	*	*	5.1	12.3	20.8	30.7
1999 [4]	1.5	*	*	*	*	*	*	1.5	5.9	12.4	22.2	31.2
Non-Hispanic [5]												
Both sexes												
2001	4.3	*	*	*	*	*	0.3	1.7	6.6	17.5	31.1	36.5
2000 [4]	4.3	*	*	*	*	*	0.3	1.9	6.4	17.4	31.8	38.0
1999 [4]	4.3	*	*	*	*	*	0.4	1.8	7.1	17.4	31.2	36.8
Male												
2001	4.5	*	*	*	*	*	0.4	2.2	7.9	20.8	38.6	48.4
2000 [4]	4.4	*	*	*	*	*	0.4	2.2	7.0	20.5	39.9	52.5
1999 [4]	4.5	*	*	*	*	*	0.4	2.1	8.5	20.4	38.9	52.2
Female												
2001	4.1	*	*	*	*	*	0.3	1.4	5.5	14.8	26.1	31.5
2000 [4]	4.2	*	*	*	*	*	0.3	1.6	5.7	14.8	26.5	32.1
1999 [4]	4.1	*	*	*	*	*	0.3	1.5	5.8	15.0	26.3	30.6
Non-Hispanic white												
Both sexes												
2001	4.4	*	*	*	*	*	0.3	1.5	5.7	16.3	29.7	35.1
2000 [4]	4.3	*	*	*	*	*	0.3	1.6	5.6	16.2	30.2	35.2
1999 [4]	4.3	*	*	*	*	*	0.3	1.5	6.1	16.2	29.6	35.0
Male												
2001	4.6	*	*	*	*	*	0.3	1.9	6.9	19.5	37.4	47.3

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Multiple myeloma and immunoproliferative neoplasms (C88,C90)												
Non-Hispanic white												
Male												
2000 [4]	4.6	*	*	*	*	*	0.4	1.8	6.3	19.2	38.5	50.9
1999 [4]	4.6	*	*	*	*	*	0.3	1.9	7.5	19.2	37.4	50.6
Female												
2001	4.1	*	*	*	*	*	0.2	1.1	4.7	13.6	24.6	30.0
2000 [4]	4.1	*	*	*	*	*	0.2	1.4	4.9	13.6	24.8	28.9
1999 [4]	4.1	*	*	*	*	*	0.2	1.2	4.7	13.7	24.6	28.7
Non-Hispanic black												
Both sexes												
2001	5.0	*	*	*	*	*	0.9	3.7	15.3	31.8	53.2	59.3
2000 [4]	5.2	*	*	*	*	*	0.8	4.5	14.1	32.5	55.6	72.8
1999 [4]	5.1	*	*	*	*	*	0.9	4.3	16.8	30.7	53.6	63.4
Male												
2001	5.0	*	*	*	*	*	1.1	4.3	18.5	37.2	66.0	72.8
2000 [4]	5.0	*	*	*	*	*	*	5.0	15.4	39.1	68.5	81.0
1999 [4]	5.0	*	*	*	*	*	1.1	4.4	19.0	35.6	65.0	80.8
Female												
2001	5.0	*	*	*	*	*	0.8	3.1	12.7	28.0	46.0	54.3
2000 [4]	5.4	*	*	*	*	*	0.9	4.1	13.1	27.9	48.3	69.8
1999 [4]	5.3	*	*	*	*	*	0.7	4.3	15.1	27.2	47.3	56.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)												
All origins [3]												
Both sexes												
2001	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.5
2000 [4]	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.5
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	0.2	*
Male												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)												
Hispanic												
Female												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.5
2000 [4]	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.5
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	0.2	*
Male												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2001	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	0.2	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	0.2	*
Male												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)												
Non-Hispanic white												
Male												
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49, C51-C52,C57-C60,C62-C63,C66,C68-C69, C73-C80,C97)												
All origins [3]												
Both sexes												
2001	22.0	*	0.8	0.7	1.3	1.8	4.6	14.1	37.3	82.0	148.7	234.5
2000 [4]	22.4	0.6	0.8	0.6	1.3	1.8	4.6	14.1	38.3	85.5	152.6	239.8
1999 [4]	23.7	*	0.8	0.7	1.3	1.9	5.1	15.0	41.5	91.4	160.8	250.4
Male												
2001	22.6	*	0.9	0.7	1.7	2.2	4.9	16.4	43.8	98.5	181.1	295.2
2000 [4]	23.0	*	1.0	0.7	1.6	2.1	5.1	16.0	44.9	103.2	187.8	300.7
1999 [4]	24.5	*	0.8	0.8	1.7	2.4	5.8	17.3	49.1	110.4	196.1	318.9
Female												
2001	21.4	*	0.6	0.6	0.9	1.5	4.2	11.7	31.3	68.4	127.4	209.1
2000 [4]	21.9	*	0.7	0.6	1.0	1.4	4.2	12.2	32.2	70.8	129.6	215.0
1999 [4]	23.1	*	0.7	0.7	1.0	1.4	4.5	12.8	34.5	75.8	137.9	222.8
Hispanic												
Both sexes												
2001	7.7	*	*	0.6	1.3	1.2	3.4	9.8	26.0	60.7	110.9	193.8
2000 [4]	7.8	*	0.8	0.7	1.2	1.5	2.7	9.2	26.6	64.1	124.6	192.4
1999 [4]	8.1	*	*	0.7	1.3	1.6	3.5	10.6	28.8	64.8	122.3	202.1
Male												
2001	7.7	*	*	0.6	1.8	1.5	3.4	10.7	30.2	67.7	140.9	213.3
2000 [4]	7.7	*	*	0.7	1.4	1.9	2.9	9.5	29.2	73.6	151.9	231.8
1999 [4]	8.0	*	*	0.7	1.8	1.9	3.5	11.1	33.3	77.1	144.9	235.1

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49, C51-C52,C57-C60,C62-C63,C66,C68-C69, C73-C80,C97)												
Hispanic												
Female												
2001	7.7	*	*	0.6	0.7	0.8	3.5	8.9	22.4	55.3	90.7	184.3
2000 [4]	7.9	*	*	0.6	1.1	1.1	2.5	9.0	24.3	56.6	106.3	173.1
1999 [4]	8.2	*	*	0.7	0.8	1.2	3.6	10.1	24.7	55.1	107.2	186.1
Non-Hispanic [5]												
Both sexes												
2001	24.1	*	0.8	0.7	1.3	2.0	4.7	14.4	38.1	83.1	150.1	235.3
2000 [4]	24.4	0.7	0.8	0.6	1.3	1.8	4.8	14.5	39.1	86.5	153.3	241.0
1999 [4]	25.8	*	0.8	0.7	1.3	1.9	5.3	15.3	42.3	92.7	161.9	251.6
Male												
2001	24.9	*	1.0	0.8	1.7	2.4	5.1	16.9	44.7	100.0	182.7	297.9
2000 [4]	25.2	*	0.9	0.6	1.6	2.2	5.3	16.5	45.9	104.7	188.7	303.0
1999 [4]	26.8	*	1.0	0.8	1.6	2.5	6.1	17.8	50.1	112.0	197.6	321.9
Female												
2001	23.3	*	0.6	0.6	0.9	1.6	4.3	12.0	31.9	69.1	128.7	209.3
2000 [4]	23.7	*	0.8	0.6	1.0	1.4	4.4	12.5	32.8	71.5	130.2	215.9
1999 [4]	25.0	*	0.7	0.6	1.1	1.4	4.6	13.0	35.1	76.9	138.8	223.5
Non-Hispanic white												
Both sexes												
2001	26.0	*	0.9	0.7	1.3	2.0	4.5	13.6	36.8	82.5	150.8	238.9
2000 [4]	26.5	*	0.8	0.7	1.3	1.9	4.5	13.6	38.2	86.3	153.9	244.9
1999 [4]	27.9	*	0.9	0.7	1.3	2.0	5.0	14.6	40.9	92.7	162.3	253.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49, C51-C52,C57-C60,C62-C63,C66,C68-C69, C73-C80,C97)												
Non-Hispanic white												
Male												
2001	26.8	*	1.1	0.7	1.7	2.3	5.0	15.9	42.8	99.3	183.7	302.9
2000 [4]	27.2	*	1.0	0.7	1.6	2.3	5.0	15.6	44.1	104.6	189.1	309.5
1999 [4]	28.8	*	1.1	0.8	1.5	2.6	5.8	16.6	48.3	111.5	198.9	326.9
Female												
2001	25.3	*	0.7	0.6	1.0	1.6	4.0	11.4	31.3	68.3	129.1	212.3
2000 [4]	25.8	*	0.6	0.6	1.0	1.4	4.1	11.7	32.6	71.0	130.8	218.9
1999 [4]	27.0	*	0.8	0.6	1.1	1.3	4.2	12.6	33.9	77.0	138.3	224.3
Non-Hispanic black												
Both sexes												
2001	18.9	*	*	0.7	1.3	2.4	6.1	22.2	54.0	101.8	158.4	218.5
2000 [4]	18.9	*	0.9	0.6	1.5	1.7	7.7	22.6	52.8	101.2	163.6	211.1
1999 [4]	20.2	*	*	0.8	1.5	2.0	7.6	23.6	60.6	105.9	172.7	244.8
Male												
2001	20.0	*	*	0.8	1.8	2.8	6.3	27.1	69.0	128.4	195.0	287.6
2000 [4]	20.0	*	*	0.7	2.0	1.9	8.7	26.9	69.1	123.8	212.0	253.6
1999 [4]	21.3	*	*	0.7	2.0	2.4	8.5	29.7	77.5	134.5	208.6	293.6
Female												
2001	17.8	*	*	0.6	0.9	2.0	5.8	18.0	42.0	83.0	137.8	192.9
2000 [4]	17.8	*	*	*	1.0	1.5	6.8	18.9	39.6	85.2	136.3	195.4
1999 [4]	19.2	*	*	0.9	1.0	1.7	6.9	18.3	47.0	85.8	152.7	226.5

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)												
All origins [3]												
Both sexes												
2001	4.8	1.3	0.4	0.3	0.3	0.4	0.8	1.8	5.0	14.1	37.0	82.7
2000 [4]	4.8	1.6	0.3	0.2	0.2	0.4	0.8	1.8	5.0	13.8	37.6	84.9
1999 [4]	4.7	1.5	0.4	0.2	0.3	0.4	0.8	1.8	4.8	14.6	36.6	82.9
Male												
2001	4.8	1.3	0.4	0.2	0.3	0.4	0.9	2.0	5.5	17.3	48.2	107.6
2000 [4]	4.8	1.6	0.4	0.2	0.3	0.4	0.8	2.1	5.6	17.0	47.5	108.9
1999 [4]	4.8	1.5	0.4	0.2	0.3	0.5	0.9	2.0	5.8	18.4	46.1	109.2
Female												
2001	4.8	1.3	0.4	0.3	0.3	0.4	0.8	1.6	4.4	11.3	29.7	72.3
2000 [4]	4.9	1.5	0.3	0.2	0.2	0.4	0.8	1.6	4.5	11.2	31.1	75.2
1999 [4]	4.7	1.5	0.4	0.3	0.2	0.4	0.6	1.6	3.9	11.5	30.4	72.3
Hispanic												
Both sexes												
2001	1.5	*	*	0.3	*	0.3	0.7	1.5	3.6	8.8	23.1	57.5
2000 [4]	1.5	*	*	*	*	0.4	0.7	1.4	3.1	9.1	24.3	70.3
1999 [4]	1.5	*	*	*	*	0.4	0.6	1.3	3.7	10.5	23.1	51.9
Male												
2001	1.5	*	*	*	*	*	0.8	1.8	4.0	10.0	30.0	70.5
2000 [4]	1.3	*	*	*	*	*	*	*	3.1	9.7	26.6	86.7
1999 [4]	1.3	*	*	*	*	*	0.8	1.8	3.9	10.3	24.5	49.6
Female												
2001	1.5	*	*	*	*	*	*	1.2	3.3	7.9	18.4	51.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)												
Hispanic												
Female												
2000 [4]	1.7	*	*	*	*	*	0.9	1.6	3.1	8.6	22.8	62.3
1999 [4]	1.6	*	*	*	*	*	*	*	3.5	10.7	22.2	53.0
Non-Hispanic [5]												
Both sexes												
2001	5.3	1.4	0.4	0.2	0.3	0.4	0.8	1.9	5.0	14.4	37.5	83.5
2000 [4]	5.3	1.7	0.4	0.2	0.3	0.4	0.8	1.9	5.2	14.1	38.1	85.2
1999 [4]	5.2	1.6	0.4	0.2	0.3	0.4	0.8	1.8	4.8	14.8	37.0	83.8
Male												
2001	5.3	1.5	0.4	0.2	0.3	0.4	0.9	2.1	5.6	17.8	48.9	109.0
2000 [4]	5.3	1.8	0.5	0.2	0.3	0.4	0.9	2.2	5.8	17.5	48.3	109.4
1999 [4]	5.3	1.6	0.5	0.2	0.3	0.5	0.9	2.0	5.9	18.8	46.9	111.4
Female												
2001	5.2	1.4	0.4	0.3	0.3	0.4	0.8	1.7	4.5	11.5	30.1	72.9
2000 [4]	5.3	1.7	*	0.3	0.2	0.4	0.8	1.6	4.6	11.4	31.4	75.4
1999 [4]	5.1	1.7	0.4	0.3	0.2	0.4	0.7	1.7	3.9	11.5	30.6	72.8
Non-Hispanic white												
Both sexes												
2001	5.9	1.3	0.4	0.2	0.3	0.4	0.8	1.9	5.0	14.7	38.6	84.8
2000 [4]	5.9	1.8	0.4	0.2	0.3	0.4	0.8	1.8	5.2	14.2	39.1	87.6
1999 [4]	5.7	1.5	0.4	0.3	0.2	0.4	0.7	1.8	4.8	15.0	37.8	86.3
Male												
2001	6.0	*	*	0.2	0.2	0.5	0.9	2.1	5.7	18.3	50.1	111.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)												
Non-Hispanic white												
Male												
2000 [4]	5.9	1.8	0.5	0.2	0.3	0.4	0.8	2.1	5.8	17.4	49.9	113.1
1999 [4]	5.9	*	*	0.2	0.2	0.4	0.9	2.0	5.9	19.2	48.1	116.8
Female												
2001	5.8	*	*	0.3	0.3	0.4	0.7	1.7	4.5	11.6	30.9	73.7
2000 [4]	5.9	1.8	*	0.3	0.2	0.4	0.8	1.5	4.7	11.5	32.1	77.3
1999 [4]	5.6	*	*	0.3	0.2	0.4	0.6	1.5	3.7	11.5	31.1	74.1
Non-Hispanic black												
Both sexes												
2001	3.2	*	*	0.4	0.4	*	1.1	2.1	6.0	14.5	31.9	72.7
2000 [4]	3.2	*	*	*	*	0.6	1.1	2.7	5.6	15.1	31.3	64.2
1999 [4]	3.3	*	*	*	0.4	0.6	1.2	2.6	6.4	15.9	33.1	62.7
Male												
2001	3.1	*	*	*	*	*	1.1	1.9	6.6	17.6	42.3	87.8
2000 [4]	3.1	*	*	*	*	*	1.2	2.8	7.0	19.5	36.4	76.3
1999 [4]	3.1	*	*	*	*	0.8	1.2	2.4	7.3	19.2	38.8	64.2
Female												
2001	3.3	*	*	*	*	*	1.0	2.2	5.4	12.2	26.1	67.2
2000 [4]	3.3	*	*	*	*	0.8	1.0	2.6	4.5	12.1	28.5	59.7
1999 [4]	3.5	*	*	*	*	*	1.2	2.8	5.7	13.6	29.9	62.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Anemias (D50-D64)												
All origins [3]												
Both sexes												
2001	1.6	*	0.1	0.1	0.3	0.4	0.5	0.6	1.0	2.7	9.8	42.0
2000 [4]	1.6	*	0.2	0.1	0.2	0.4	0.4	0.6	1.1	2.8	9.5	42.4
1999 [4]	1.6	*	0.2	0.1	0.3	0.4	0.4	0.6	1.0	2.6	9.5	43.8
Male												
2001	1.3	*	*	0.1	0.3	0.5	0.4	0.6	0.9	2.9	9.9	40.0
2000 [4]	1.2	*	*	0.1	0.2	0.3	0.3	0.6	1.1	2.8	9.5	40.1
1999 [4]	1.3	*	*	0.1	0.3	0.4	0.5	0.6	0.9	2.7	10.1	43.4
Female												
2001	2.0	*	*	0.1	0.3	0.3	0.5	0.6	1.0	2.5	9.6	42.9
2000 [4]	2.0	*	*	0.1	0.2	0.4	0.5	0.6	1.0	2.8	9.4	43.3
1999 [4]	1.9	*	*	0.2	0.3	0.4	0.4	0.6	1.0	2.6	9.2	44.0
Hispanic												
Both sexes												
2001	0.5	*	*	*	*	*	*	*	*	2.2	7.8	28.8
2000 [4]	0.4	*	*	*	*	*	*	*	*	2.2	7.1	17.3
1999 [4]	0.5	*	*	*	*	*	*	*	*	2.6	7.3	35.1
Male												
2001	0.4	*	*	*	*	*	*	*	*	*	9.4	*
2000 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.5	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.6	*	*	*	*	*	*	*	*	*	6.6	32.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Anemias (D50-D64)												
Hispanic												
Female												
2000 [4]	0.5	*	*	*	*	*	*	*	*	*	6.9	*
1999 [4]	0.5	*	*	*	*	*	*	*	*	*	*	32.2
Non-Hispanic [5]												
Both sexes												
2001	1.8	*	*	0.1	0.3	0.4	0.5	0.6	1.0	2.7	9.8	42.5
2000 [4]	1.8	*	0.3	0.1	0.2	0.4	0.5	0.6	1.1	2.8	9.5	43.2
1999 [4]	1.8	*	0.2	0.1	0.4	0.4	0.5	0.7	1.0	2.6	9.6	44.0
Male												
2001	1.4	*	*	0.1	0.3	0.5	0.4	0.6	0.9	2.9	10.0	40.8
2000 [4]	1.3	*	*	0.1	0.2	0.4	0.4	0.6	1.2	2.8	9.6	41.0
1999 [4]	1.4	*	*	0.1	0.4	0.5	0.5	0.7	1.0	2.6	10.1	43.3
Female												
2001	2.2	*	*	0.2	0.3	0.4	0.6	0.6	1.0	2.6	9.7	43.1
2000 [4]	2.2	*	*	*	0.3	0.4	0.5	0.6	1.0	2.8	9.5	44.1
1999 [4]	2.1	*	*	0.2	0.3	0.4	0.4	0.6	1.0	2.6	9.3	44.2
Non-Hispanic white												
Both sexes												
2001	1.7	*	*	0.1	0.1	0.1	0.1	0.3	0.8	2.5	9.4	42.1
2000 [4]	1.7	*	*	*	*	*	0.1	0.3	0.8	2.5	9.3	42.9
1999 [4]	1.6	*	*	*	0.1	*	0.1	0.3	0.6	2.4	9.4	43.2
Male												
2001	1.3	*	*	*	*	*	*	0.3	0.7	2.6	9.8	41.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Anemias (D50-D64)												
Non-Hispanic white												
Male												
2000 [4]	1.2	*	*	*	*	*	0.1	0.3	0.9	2.7	9.3	39.9
1999 [4]	1.2	*	*	*	*	*	*	0.3	0.5	2.4	9.9	43.4
Female												
2001	2.1	*	*	*	*	*	0.2	0.3	0.8	2.4	9.2	42.4
2000 [4]	2.1	*	*	*	*	*	*	0.3	0.7	2.4	9.4	44.1
1999 [4]	2.0	*	*	*	*	*	*	0.3	0.7	2.4	9.1	43.1
Non-Hispanic black												
Both sexes												
2001	2.8	*	*	0.4	1.2	2.3	2.8	3.1	3.0	5.5	16.1	49.3
2000 [4]	2.7	*	*	0.5	1.1	2.4	2.5	3.1	3.5	5.5	13.4	51.8
1999 [4]	2.9	*	0.9	0.5	1.5	2.7	2.6	3.2	4.3	5.0	13.4	53.0
Male												
2001	2.6	*	*	*	1.4	3.0	2.5	3.6	3.2	6.8	15.3	42.7
2000 [4]	2.4	*	*	*	1.0	2.2	2.1	3.4	3.5	4.6	13.9	56.4
1999 [4]	2.7	*	*	*	1.8	2.9	2.8	3.4	5.1	5.6	12.8	44.0
Female												
2001	2.9	*	*	*	1.1	1.6	3.0	2.7	2.9	4.6	16.5	51.8
2000 [4]	3.1	*	*	*	1.1	2.5	2.8	2.9	3.5	6.1	13.2	50.2
1999 [4]	3.1	*	*	*	1.2	2.4	2.4	3.0	3.6	4.7	13.7	56.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diabetes mellitus (E10-E14)												
All origins [3]												
Both sexes												
2001	25.1	*	*	0.1	0.4	1.5	4.3	13.6	37.8	91.4	181.4	321.8
2000 [4]	24.6	*	*	0.1	0.4	1.6	4.3	13.1	37.8	90.7	179.5	319.7
1999 [4]	24.5	*	*	0.1	0.4	1.4	4.3	12.9	38.3	91.8	178.0	317.2
Male												
2001	23.5	*	*	0.1	0.4	1.7	5.3	16.4	43.1	101.7	198.9	343.1
2000 [4]	22.9	*	*	*	0.5	1.9	5.1	15.3	42.4	99.1	199.6	345.1
1999 [4]	22.8	*	*	*	0.4	1.6	5.1	15.6	42.6	99.8	198.2	345.7
Female												
2001	26.6	*	*	*	0.3	1.3	3.4	11.0	32.9	82.8	169.9	312.9
2000 [4]	26.3	*	*	*	0.4	1.2	3.4	11.1	33.6	83.7	166.3	309.4
1999 [4]	26.2	*	*	*	0.4	1.3	3.5	10.4	34.3	85.2	164.9	305.8
Hispanic												
Both sexes												
2001	15.3	*	*	*	*	0.7	3.1	14.9	53.2	141.2	279.2	489.6
2000 [4]	15.1	*	*	*	*	1.0	3.0	14.0	55.5	147.9	278.5	465.1
1999 [4]	15.3	*	*	*	*	0.7	3.2	13.9	56.6	149.7	285.2	493.9
Male												
2001	13.6	*	*	*	*	0.8	3.9	18.5	58.7	151.4	285.3	422.8
2000 [4]	13.8	*	*	*	*	1.4	3.6	15.5	64.7	160.8	299.4	449.4
1999 [4]	13.4	*	*	*	*	0.6	3.3	17.2	62.5	155.8	293.0	468.0
Female												
2001	17.1	*	*	*	*	0.7	2.2	11.4	48.3	133.2	275.0	522.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diabetes mellitus (E10-E14)												
Hispanic												
Female												
2000 [4]	16.5	*	*	*	*	*	2.4	12.6	47.4	137.7	264.5	472.8
1999 [4]	17.3	*	*	*	*	0.8	3.1	10.7	51.4	144.9	280.0	506.4
Non-Hispanic [5]												
Both sexes												
2001	26.4	*	*	0.1	0.4	1.7	4.5	13.4	36.5	87.7	176.4	314.8
2000 [4]	25.9	*	*	0.1	0.5	1.7	4.4	13.0	36.4	86.8	174.8	313.4
1999 [4]	25.7	*	*	0.1	0.4	1.6	4.4	12.8	36.8	88.1	173.1	310.0
Male												
2001	25.0	*	*	0.1	0.4	1.9	5.5	16.1	41.8	98.0	194.3	338.9
2000 [4]	24.2	*	*	*	0.5	2.0	5.3	15.2	40.7	95.0	194.6	339.7
1999 [4]	24.1	*	*	*	0.4	1.8	5.3	15.3	41.0	96.3	193.8	339.5
Female												
2001	27.8	*	*	*	0.3	1.4	3.6	10.9	31.6	79.1	164.6	304.7
2000 [4]	27.6	*	*	*	0.4	1.4	3.5	11.0	32.4	80.1	161.8	302.8
1999 [4]	27.3	*	*	*	0.4	1.4	3.6	10.4	32.8	81.4	159.6	298.2
Non-Hispanic white												
Both sexes												
2001	26.0	*	*	*	0.3	1.3	3.9	11.3	30.7	77.0	162.7	297.4
2000 [4]	25.4	*	*	*	0.4	1.4	3.9	10.7	30.9	76.1	159.9	295.5
1999 [4]	25.0	*	*	*	0.4	1.4	3.8	10.5	30.1	76.8	159.1	290.2
Male												
2001	25.1	*	*	*	0.3	1.5	4.8	13.7	36.4	89.8	183.5	332.1

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diabetes mellitus (E10-E14)												
Non-Hispanic white												
Male												
2000 [4]	24.3	*	*	*	0.5	1.7	4.8	13.0	35.4	86.6	184.5	330.4
1999 [4]	24.0	*	*	*	0.4	1.5	4.6	13.0	35.0	87.7	184.6	326.0
Female												
2001	26.9	*	*	*	0.2	1.2	2.9	8.8	25.3	66.0	148.9	283.0
2000 [4]	26.5	*	*	*	0.3	1.1	3.1	8.6	26.6	67.3	143.8	281.4
1999 [4]	26.0	*	*	*	0.3	1.2	3.0	8.1	25.4	67.7	142.4	275.9
Non-Hispanic black												
Both sexes												
2001	34.1	*	*	*	1.0	3.8	9.1	30.1	86.5	190.8	337.9	533.8
2000 [4]	34.0	*	*	*	1.0	3.6	8.2	31.0	83.9	190.6	354.8	530.1
1999 [4]	34.1	*	*	*	0.8	3.1	9.1	31.0	92.0	194.0	336.1	530.9
Male												
2001	29.4	*	*	*	0.9	5.0	10.5	35.0	93.8	189.5	345.1	452.7
2000 [4]	28.3	*	*	*	1.0	4.0	9.7	34.2	89.7	186.9	339.4	456.8
1999 [4]	28.6	*	*	*	*	3.4	10.7	35.3	95.4	189.7	317.7	506.3
Female												
2001	38.4	*	*	*	1.0	2.8	7.8	25.9	80.6	191.7	333.9	563.8
2000 [4]	39.1	*	*	*	1.0	3.3	7.0	28.2	79.1	193.3	363.5	557.3
1999 [4]	39.0	*	*	*	0.9	2.7	7.7	27.2	89.2	197.0	346.5	540.1

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nutritional deficiencies (E40-E64)												
All origins [3]												
Both sexes												
2001	1.3	*	*	*	*	*	0.1	0.3	0.6	2.0	8.7	42.9
2000 [4]	1.5	*	*	*	*	*	0.1	0.3	0.7	2.5	10.4	50.0
1999 [4]	1.5	*	*	*	*	*	0.1	0.3	0.7	2.3	10.2	53.9
Male												
2001	0.9	*	*	*	*	*	0.1	0.3	0.7	2.1	8.5	35.2
2000 [4]	1.0	*	*	*	*	*	*	0.2	0.8	2.7	9.7	43.7
1999 [4]	1.1	*	*	*	*	*	0.1	0.3	0.9	2.7	10.3	48.4
Female												
2001	1.7	*	*	*	*	*	0.1	0.3	0.5	2.0	8.9	46.1
2000 [4]	2.0	*	*	*	*	*	0.2	0.3	0.6	2.4	10.9	52.6
1999 [4]	2.0	*	*	*	*	*	0.2	0.3	0.6	2.1	10.2	56.2
Hispanic												
Both sexes												
2001	0.4	*	*	*	*	*	*	*	*	*	8.1	38.8
2000 [4]	0.4	*	*	*	*	*	*	*	*	2.2	8.5	39.8
1999 [4]	0.4	*	*	*	*	*	*	*	*	*	6.5	51.2
Male												
2001	0.3	*	*	*	*	*	*	*	*	*	8.9	43.8
2000 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	*	51.8
Female												
2001	0.5	*	*	*	*	*	*	*	*	*	7.6	36.3

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nutritional deficiencies (E40-E64)												
Hispanic												
Female												
2000 [4]	0.5	*	*	*	*	*	*	*	*	*	8.6	41.5
1999 [4]	0.5	*	*	*	*	*	*	*	*	*	*	51.0
Non-Hispanic [5]												
Both sexes												
2001	1.4	*	*	*	*	*	0.1	0.3	0.6	2.1	8.8	43.0
2000 [4]	1.7	*	*	*	*	*	0.1	0.3	0.7	2.5	10.4	50.3
1999 [4]	1.7	*	*	*	*	*	0.1	0.3	0.7	2.4	10.3	53.9
Male												
2001	1.0	*	*	*	*	*	*	0.3	0.7	2.2	8.5	34.8
2000 [4]	1.1	*	*	*	*	*	*	0.3	0.9	2.7	9.6	43.9
1999 [4]	1.2	*	*	*	*	*	0.1	0.3	0.9	2.7	10.4	48.1
Female												
2001	1.9	*	*	*	*	*	*	0.3	0.6	2.0	8.9	46.4
2000 [4]	2.2	*	*	*	*	*	0.2	0.3	0.6	2.4	10.9	52.8
1999 [4]	2.2	*	*	*	*	*	0.2	0.3	0.6	2.1	10.3	56.2
Non-Hispanic white												
Both sexes												
2001	1.5	*	*	*	*	*	0.1	0.2	0.6	2.0	8.2	42.2
2000 [4]	1.8	*	*	*	*	*	0.1	0.2	0.6	2.3	10.0	49.0
1999 [4]	1.8	*	*	*	*	*	0.1	0.3	0.6	2.0	9.6	52.8
Male												
2001	1.0	*	*	*	*	*	*	0.2	0.5	1.9	7.9	33.3

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nutritional deficiencies (E40-E64)												
Non-Hispanic white												
Male												
2000 [4]	1.2	*	*	*	*	*	*	0.2	0.7	2.3	9.0	43.0
1999 [4]	1.2	*	*	*	*	*	*	0.2	0.7	2.2	9.4	46.2
Female												
2001	2.0	*	*	*	*	*	*	0.3	0.6	2.0	8.3	45.9
2000 [4]	2.3	*	*	*	*	*	0.2	0.3	0.6	2.3	10.6	51.5
1999 [4]	2.3	*	*	*	*	*	0.2	0.3	0.5	1.9	9.8	55.4
Non-Hispanic black												
Both sexes												
2001	1.3	*	*	*	*	*	*	0.8	1.2	3.7	16.1	55.2
2000 [4]	1.5	*	*	*	*	*	*	0.6	1.7	4.8	16.3	67.4
1999 [4]	1.7	*	*	*	*	*	*	0.7	2.0	5.6	19.6	70.8
Male												
2001	1.2	*	*	*	*	*	*	1.2	1.9	5.6	16.5	58.9
2000 [4]	1.3	*	*	*	*	*	*	*	2.8	6.6	17.9	64.6
1999 [4]	1.5	*	*	*	*	*	*	*	2.9	8.3	23.1	76.1
Female												
2001	1.4	*	*	*	*	*	*	*	*	2.4	15.8	53.9
2000 [4]	1.8	*	*	*	*	*	*	*	*	3.6	15.5	68.5
1999 [4]	1.8	*	*	*	*	*	*	*	*	3.7	17.7	68.8

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malnutrition (E40-E46)												
All origins [3]												
Both sexes												
2001	1.2	*	*	*	*	*	0.1	0.3	0.6	1.9	8.2	40.2
2000 [4]	1.4	*	*	*	*	*	0.1	0.2	0.7	2.3	9.7	46.3
1999 [4]	1.4	*	*	*	*	*	0.1	0.3	0.6	2.2	9.5	50.0
Male												
2001	0.8	*	*	*	*	*	*	0.2	0.6	1.9	8.0	33.6
2000 [4]	0.9	*	*	*	*	*	*	0.2	0.7	2.5	8.9	40.7
1999 [4]	1.0	*	*	*	*	*	0.1	0.3	0.8	2.5	9.4	44.4
Female												
2001	1.6	*	*	*	*	*	0.1	0.3	0.5	1.9	8.2	43.0
2000 [4]	1.8	*	*	*	*	*	0.2	0.3	0.6	2.1	10.2	48.6
1999 [4]	1.8	*	*	*	*	*	0.1	0.3	0.5	2.0	9.5	52.3
Hispanic												
Both sexes												
2001	0.4	*	*	*	*	*	*	*	*	*	7.9	38.1
2000 [4]	0.4	*	*	*	*	*	*	*	*	2.0	7.7	38.5
1999 [4]	0.4	*	*	*	*	*	*	*	*	*	6.2	47.7
Male												
2001	0.3	*	*	*	*	*	*	*	*	*	8.9	41.9
2000 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	*	45.3
Female												
2001	0.4	*	*	*	*	*	*	*	*	*	7.3	36.3

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malnutrition (E40-E46)												
Hispanic												
Female												
2000 [4]	0.5	*	*	*	*	*	*	*	*	*	7.6	41.5
1999 [4]	0.5	*	*	*	*	*	*	*	*	*	*	48.9
Non-Hispanic [5]												
Both sexes												
2001	1.3	*	*	*	*	*	0.1	0.3	0.6	2.0	8.2	40.2
2000 [4]	1.5	*	*	*	*	*	0.1	0.2	0.7	2.3	9.7	46.5
1999 [4]	1.6	*	*	*	*	*	0.1	0.3	0.7	2.2	9.6	50.0
Male												
2001	0.9	*	*	*	*	*	*	0.3	0.6	2.0	8.0	33.2
2000 [4]	1.0	*	*	*	*	*	*	0.2	0.8	2.5	8.8	40.9
1999 [4]	1.1	*	*	*	*	*	0.1	0.3	0.8	2.5	9.5	44.3
Female												
2001	1.7	*	*	*	*	*	*	0.3	0.6	1.9	8.3	43.2
2000 [4]	2.0	*	*	*	*	*	0.2	0.3	0.6	2.2	10.3	48.7
1999 [4]	2.0	*	*	*	*	*	0.2	0.3	0.5	2.0	9.7	52.2
Non-Hispanic white												
Both sexes												
2001	1.4	*	*	*	*	*	*	0.2	0.5	1.8	7.6	39.4
2000 [4]	1.6	*	*	*	*	*	0.1	0.2	0.6	2.1	9.3	45.3
1999 [4]	1.6	*	*	*	*	*	0.1	0.2	0.5	1.9	8.9	48.8
Male												
2001	0.9	*	*	*	*	*	*	0.2	0.5	1.7	7.4	31.7

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malnutrition (E40-E46)												
Non-Hispanic white												
Male												
2000 [4]	1.1	*	*	*	*	*	*	0.2	0.6	2.1	8.3	39.8
1999 [4]	1.1	*	*	*	*	*	*	0.2	0.6	2.0	8.6	42.0
Female												
2001	1.9	*	*	*	*	*	*	0.3	0.6	1.9	7.7	42.7
2000 [4]	2.2	*	*	*	*	*	0.1	0.2	0.6	2.0	10.0	47.4
1999 [4]	2.2	*	*	*	*	*	0.2	0.3	0.4	1.8	9.2	51.6
Non-Hispanic black												
Both sexes												
2001	1.2	*	*	*	*	*	*	0.6	1.1	3.4	15.0	52.1
2000 [4]	1.4	*	*	*	*	*	*	0.6	1.6	4.8	15.1	64.2
1999 [4]	1.6	*	*	*	*	*	*	0.6	1.8	5.5	18.1	66.0
Male												
2001	1.1	*	*	*	*	*	*	1.0	*	5.0	15.9	56.6
2000 [4]	1.2	*	*	*	*	*	*	*	2.5	6.4	17.3	62.2
1999 [4]	1.4	*	*	*	*	*	*	*	2.6	8.0	20.6	74.9
Female												
2001	1.3	*	*	*	*	*	*	*	*	2.3	14.5	50.5
2000 [4]	1.6	*	*	*	*	*	*	*	*	3.6	13.9	65.0
1999 [4]	1.7	*	*	*	*	*	*	*	*	3.7	16.6	62.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other nutritional deficiencies (E50-E64)												
All origins [3]												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	0.1	0.6	2.7
2000 [4]	0.1	*	*	*	*	*	*	*	*	0.2	0.7	3.7
1999 [4]	0.1	*	*	*	*	*	*	*	*	0.1	0.7	3.9
Male												
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	1.6
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.8	3.0
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	0.9	3.9
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	0.6	3.1
2000 [4]	0.1	*	*	*	*	*	*	*	*	0.2	0.7	4.0
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	0.6	3.9
Hispanic												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other nutritional deficiencies (E50-E64)												
Hispanic												
Female												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	0.2	0.6	2.7
2000 [4]	0.1	*	*	*	*	*	*	*	*	0.2	0.7	3.8
1999 [4]	0.1	*	*	*	*	*	*	*	*	0.1	0.7	3.9
Male												
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	1.6
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.8	3.0
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	0.9	3.8
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	0.7	3.2
2000 [4]	0.2	*	*	*	*	*	*	*	*	0.2	0.6	4.1
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	0.7	4.0
Non-Hispanic white												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	0.1	0.6	2.8
2000 [4]	0.1	*	*	*	*	*	*	*	*	0.2	0.6	3.8
1999 [4]	0.1	*	*	*	*	*	*	*	*	0.1	0.7	3.9
Male												
2001	0.1	*	*	*	*	*	*	*	*	*	0.6	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other nutritional deficiencies (E50-E64)												
Non-Hispanic white												
Male												
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.8	3.1
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	0.8	4.2
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	0.6	3.3
2000 [4]	0.2	*	*	*	*	*	*	*	*	0.3	0.6	4.0
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	0.6	3.9
Non-Hispanic black												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningitis (G00,G03)												
All origins [3]												
Both sexes												
2001	0.3	1.9	0.2	0.1	0.1	0.1	0.2	0.3	0.4	0.5	0.8	0.9
2000 [4]	0.3	2.4	0.2	0.1	*	0.1	0.2	0.3	0.4	0.6	1.0	1.2
1999 [4]	0.3	3.1	0.3	0.1	0.1	0.1	0.2	0.3	0.4	0.6	1.0	1.5
Male												
2001	0.3	1.9	*	*	0.1	0.1	0.2	0.3	0.5	0.6	0.9	*
2000 [4]	0.3	2.5	0.3	*	*	0.1	0.2	0.3	0.5	0.7	1.3	*
1999 [4]	0.3	3.1	0.3	*	0.1	0.1	0.2	0.4	0.4	0.7	1.0	1.7
Female												
2001	0.2	2.0	*	*	*	*	0.1	0.2	0.4	0.4	0.6	0.8
2000 [4]	0.2	2.3	*	*	*	0.1	0.1	0.2	0.3	0.5	0.8	1.0
1999 [4]	0.3	3.0	*	*	*	0.1	0.1	0.2	0.4	0.5	1.0	1.4
Hispanic												
Both sexes												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningitis (G00,G03)												
Hispanic												
Female												
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.3	2.2	0.2	0.1	0.1	0.1	0.2	0.3	0.5	0.5	0.7	0.9
2000 [4]	0.3	2.7	0.3	0.1	*	0.1	0.2	0.3	0.4	0.6	1.0	1.2
1999 [4]	0.3	3.2	0.3	0.1	0.1	0.1	0.2	0.3	0.4	0.6	1.0	1.4
Male												
2001	0.3	2.1	*	*	0.1	0.1	0.2	0.3	0.5	0.6	0.9	*
2000 [4]	0.3	2.7	*	*	*	*	0.2	0.3	0.5	0.8	1.2	*
1999 [4]	0.3	3.1	0.3	*	0.1	0.1	0.2	0.4	0.4	0.7	1.1	*
Female												
2001	0.2	2.3	*	*	*	*	0.1	0.2	0.4	0.4	0.6	0.7
2000 [4]	0.3	2.7	*	*	*	0.1	0.1	0.2	0.3	0.5	0.8	1.1
1999 [4]	0.3	3.3	*	*	*	0.1	0.1	0.2	0.4	0.5	1.0	1.4
Non-Hispanic white												
Both sexes												
2001	0.2	1.6	*	*	0.1	0.1	0.1	0.2	0.4	0.5	0.7	0.7
2000 [4]	0.2	2.1	*	*	*	0.1	0.1	0.2	0.3	0.5	0.9	1.1
1999 [4]	0.3	2.2	0.3	*	*	0.1	0.1	0.2	0.3	0.5	0.9	1.3
Male												
2001	0.3	*	*	*	*	*	*	0.3	0.4	0.5	0.8	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningitis (G00,G03)												
Non-Hispanic white												
Male												
2000 [4]	0.3	2.3	*	*	*	*	0.1	0.3	0.4	0.7	1.1	*
1999 [4]	0.3	2.8	*	*	*	*	0.1	0.3	0.4	0.6	0.9	*
Female												
2001	0.2	1.9	*	*	*	*	*	0.2	0.3	0.4	0.6	*
2000 [4]	0.2	1.8	*	*	*	*	*	0.2	0.2	0.3	0.8	1.0
1999 [4]	0.3	*	*	*	*	*	0.1	0.2	0.3	0.5	0.8	1.2
Non-Hispanic black												
Both sexes												
2001	0.5	4.7	*	*	*	*	0.5	0.7	1.0	*	*	*
2000 [4]	0.6	5.9	*	*	*	*	0.5	0.6	0.9	1.7	*	*
1999 [4]	0.6	7.0	*	*	*	*	0.4	0.9	*	1.3	2.7	*
Male												
2001	0.6	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.6	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.7	*	*	*	*	*	*	1.5	*	*	*	*
Female												
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.6	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.5	9.6	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Parkinson's disease (G20-G21)												
All origins [3]												
Both sexes												
2001	5.8	*	*	*	*	*	*	0.1	1.2	11.7	64.6	134.2
2000 [4]	5.6	*	*	*	*	*	*	0.1	1.1	11.5	61.9	131.9
1999 [4]	5.2	*	*	*	*	*	*	0.1	1.0	11.0	58.2	124.4
Male												
2001	6.7	*	*	*	*	*	*	0.2	1.8	16.8	97.0	224.4
2000 [4]	6.4	*	*	*	*	*	*	0.2	1.5	16.1	92.8	222.3
1999 [4]	6.0	*	*	*	*	*	*	0.2	1.4	15.7	89.5	205.6
Female												
2001	4.9	*	*	*	*	*	*	*	0.7	7.5	43.3	96.4
2000 [4]	4.8	*	*	*	*	*	*	0.1	0.8	7.6	41.7	95.0
1999 [4]	4.5	*	*	*	*	*	*	*	0.7	7.1	37.8	91.7
Hispanic												
Both sexes												
2001	1.3	*	*	*	*	*	*	*	*	8.6	41.5	86.3
2000 [4]	1.2	*	*	*	*	*	*	*	*	7.8	37.7	86.3
1999 [4]	1.1	*	*	*	*	*	*	*	*	8.1	38.3	77.2
Male												
2001	1.4	*	*	*	*	*	*	*	*	11.2	63.0	127.6
2000 [4]	1.3	*	*	*	*	*	*	*	*	9.7	55.1	143.1
1999 [4]	1.3	*	*	*	*	*	*	*	*	11.4	58.9	105.7
Female												
2001	1.2	*	*	*	*	*	*	*	*	6.6	26.9	66.1

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Parkinson's disease (G20-G21)												
Hispanic												
Female												
2000 [4]	1.1	*	*	*	*	*	*	*	*	6.3	26.1	58.4
1999 [4]	1.0	*	*	*	*	*	*	*	*	5.5	24.6	63.4
Non-Hispanic [5]												
Both sexes												
2001	6.5	*	*	*	*	*	*	0.1	1.2	11.9	65.5	135.6
2000 [4]	6.2	*	*	*	*	*	*	0.1	1.1	11.7	62.8	133.3
1999 [4]	5.8	*	*	*	*	*	*	0.1	1.1	11.1	58.9	125.8
Male												
2001	7.5	*	*	*	*	*	*	0.2	1.8	17.1	98.3	227.8
2000 [4]	7.1	*	*	*	*	*	*	0.2	1.5	16.5	94.2	225.2
1999 [4]	6.7	*	*	*	*	*	*	0.2	1.5	15.9	90.6	209.4
Female												
2001	5.4	*	*	*	*	*	*	*	0.7	7.6	44.0	97.3
2000 [4]	5.3	*	*	*	*	*	*	*	0.8	7.7	42.3	96.1
1999 [4]	4.9	*	*	*	*	*	*	*	0.7	7.1	38.3	92.3
Non-Hispanic white												
Both sexes												
2001	7.7	*	*	*	*	*	*	0.1	1.3	12.8	69.9	143.5
2000 [4]	7.3	*	*	*	*	*	*	0.1	1.2	12.6	66.7	141.2
1999 [4]	6.8	*	*	*	*	*	*	0.1	1.1	11.9	62.4	132.8
Male												
2001	9.0	*	*	*	*	*	*	0.2	2.0	18.3	104.5	241.7

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Parkinson's disease (G20-G21)												
Non-Hispanic white												
Male												
2000 [4]	8.4	*	*	*	*	*	*	0.2	1.6	17.6	99.2	238.6
1999 [4]	7.9	*	*	*	*	*	*	0.2	1.6	17.1	95.7	221.0
Female												
2001	6.5	*	*	*	*	*	*	*	0.7	8.2	46.9	102.7
2000 [4]	6.3	*	*	*	*	*	*	*	0.9	8.3	45.3	101.9
1999 [4]	5.8	*	*	*	*	*	*	*	0.7	7.6	40.5	97.7
Non-Hispanic black												
Both sexes												
2001	1.5	*	*	*	*	*	*	*	0.8	5.6	25.7	59.0
2000 [4]	1.5	*	*	*	*	*	*	*	*	5.8	27.3	52.5
1999 [4]	1.4	*	*	*	*	*	*	*	0.9	5.9	24.8	51.7
Male												
2001	1.7	*	*	*	*	*	*	*	*	8.8	38.1	99.3
2000 [4]	1.8	*	*	*	*	*	*	*	*	9.0	47.5	88.1
1999 [4]	1.6	*	*	*	*	*	*	*	*	8.7	38.5	87.9
Female												
2001	1.4	*	*	*	*	*	*	*	*	3.3	18.7	44.1
2000 [4]	1.2	*	*	*	*	*	*	*	*	3.5	16.0	39.2
1999 [4]	1.3	*	*	*	*	*	*	*	*	3.9	17.2	38.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alzheimer's disease (G30)												
All origins [3]												
Both sexes												
2001	18.9	*	*	*	*	*	*	0.2	2.1	18.7	147.5	710.3
2000 [4]	17.6	*	*	*	*	*	*	0.2	2.0	18.7	139.6	667.7
1999 [4]	16.0	*	*	*	*	*	*	0.2	1.9	17.4	129.5	601.3
Male												
2001	11.3	*	*	*	*	*	*	0.2	1.9	17.7	136.3	556.1
2000 [4]	10.5	*	*	*	*	*	*	0.2	1.8	18.7	127.6	522.1
1999 [4]	9.8	*	*	*	*	*	*	0.2	1.9	17.3	122.6	485.7
Female												
2001	26.3	*	*	*	*	*	*	0.2	2.3	19.5	154.8	774.9
2000 [4]	24.5	*	*	*	*	*	*	0.2	2.2	18.6	147.4	727.0
1999 [4]	21.9	*	*	*	*	*	*	0.2	1.8	17.5	134.0	648.0
Hispanic												
Both sexes												
2001	3.4	*	*	*	*	*	*	*	*	9.9	83.9	399.5
2000 [4]	3.2	*	*	*	*	*	*	*	*	11.9	79.2	380.2
1999 [4]	2.9	*	*	*	*	*	*	*	*	9.7	75.4	351.5
Male												
2001	2.1	*	*	*	*	*	*	*	*	8.8	83.2	306.6
2000 [4]	2.1	*	*	*	*	*	*	*	*	13.3	81.6	284.2
1999 [4]	1.8	*	*	*	*	*	*	*	*	11.2	66.7	278.2
Female												
2001	4.7	*	*	*	*	*	*	*	*	10.7	84.3	444.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alzheimer's disease (G30)												
Hispanic												
Female												
2000 [4]	4.3	*	*	*	*	*	*	*	*	10.8	77.6	427.3
1999 [4]	4.0	*	*	*	*	*	*	*	*	8.4	81.2	386.8
Non-Hispanic [5]												
Both sexes												
2001	21.2	*	*	*	*	*	*	0.2	2.2	19.2	150.2	720.5
2000 [4]	19.6	*	*	*	*	*	*	0.2	2.1	19.0	142.0	676.7
1999 [4]	17.7	*	*	*	*	*	*	0.2	2.0	17.8	131.5	608.9
Male												
2001	12.7	*	*	*	*	*	*	0.2	2.0	18.3	138.6	565.2
2000 [4]	11.7	*	*	*	*	*	*	0.2	1.8	19.0	129.6	531.0
1999 [4]	10.9	*	*	*	*	*	*	0.2	2.0	17.7	124.8	492.8
Female												
2001	29.3	*	*	*	*	*	*	0.2	2.4	20.0	157.8	785.1
2000 [4]	27.2	*	*	*	*	*	*	0.2	2.3	19.1	150.0	735.6
1999 [4]	24.2	*	*	*	*	*	*	0.2	1.9	18.0	135.9	655.4
Non-Hispanic white												
Both sexes												
2001	24.8	*	*	*	*	*	*	0.2	2.4	20.3	155.6	749.2
2000 [4]	22.9	*	*	*	*	*	*	0.2	2.2	19.9	147.4	705.1
1999 [4]	20.7	*	*	*	*	*	*	0.2	2.1	18.6	136.7	636.6
Male												
2001	14.7	*	*	*	*	*	*	0.2	2.1	19.1	143.0	587.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alzheimer's disease (G30)												
Non-Hispanic white												
Male												
2000 [4]	13.6	*	*	*	*	*	*	0.2	2.0	19.8	134.6	551.7
1999 [4]	12.7	*	*	*	*	*	*	0.2	2.2	18.3	129.8	515.1
Female												
2001	34.4	*	*	*	*	*	*	0.2	2.7	21.3	163.9	816.6
2000 [4]	31.9	*	*	*	*	*	*	0.2	2.5	20.0	155.8	767.0
1999 [4]	28.4	*	*	*	*	*	*	0.2	2.0	18.8	141.2	685.0
Non-Hispanic black												
Both sexes												
2001	8.7	*	*	*	*	*	*	*	1.1	15.5	126.9	513.2
2000 [4]	7.7	*	*	*	*	*	*	*	1.3	16.1	111.6	450.9
1999 [4]	6.7	*	*	*	*	*	*	*	1.2	15.5	99.1	374.7
Male												
2001	5.3	*	*	*	*	*	*	*	*	16.3	123.6	420.4
2000 [4]	4.6	*	*	*	*	*	*	*	*	16.9	98.7	378.1
1999 [4]	4.2	*	*	*	*	*	*	*	*	16.8	90.7	320.9
Female												
2001	11.8	*	*	*	*	*	*	*	*	15.0	128.7	547.5
2000 [4]	10.6	*	*	*	*	*	*	*	*	15.6	118.8	478.0
1999 [4]	9.0	*	*	*	*	*	*	*	*	14.5	103.8	394.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Major cardiovascular diseases (I00-I78)												
All origins [3]												
Both sexes												
2001	323.9	14.9	1.9	0.9	3.2	10.2	37.1	113.4	301.2	810.4	2,322.2	7,673.7
2000 [4]	332.9	16.7	1.5	0.9	3.3	9.5	36.8	115.7	319.2	849.3	2,396.2	8,030.3
1999 [4]	340.6	17.0	1.5	0.9	3.4	9.6	37.7	116.2	327.3	887.3	2,474.9	8,197.1
Male												
2001	306.2	15.3	1.9	0.9	3.9	12.8	49.9	160.4	414.0	1,057.6	2,823.3	7,936.0
2000 [4]	315.8	17.5	1.6	1.0	3.9	11.8	49.5	165.1	441.6	1,114.1	2,926.6	8,448.1
1999 [4]	323.0	17.9	1.6	1.0	4.0	12.1	51.2	165.8	452.0	1,168.4	3,024.0	8,603.2
Female												
2001	340.9	14.5	1.8	0.9	2.4	7.6	24.3	68.1	196.9	605.6	1,992.9	7,563.9
2000 [4]	349.5	15.9	1.4	0.8	2.8	7.2	24.3	68.0	206.3	631.2	2,050.4	7,860.1
1999 [4]	357.5	16.0	1.4	0.8	2.7	7.2	24.4	68.5	212.5	657.0	2,118.2	8,033.5
Hispanic												
Both sexes												
2001	96.2	12.0	1.5	0.8	2.4	6.4	19.8	75.4	216.4	644.9	1,828.4	6,036.2
2000 [4]	96.5	13.7	1.3	0.7	2.6	6.4	21.3	75.0	236.3	660.8	1,852.8	6,168.2
1999 [4]	99.2	15.2	1.4	0.7	2.5	6.5	21.4	80.5	242.7	690.0	1,935.5	6,335.9
Male												
2001	95.4	12.7	1.4	0.8	3.0	8.3	26.8	107.2	297.2	852.4	2,220.5	6,155.6
2000 [4]	96.0	13.7	1.4	0.9	3.1	8.1	27.9	107.7	330.6	870.1	2,272.6	6,304.3
1999 [4]	99.2	12.7	1.4	0.9	3.0	8.6	29.7	116.3	333.2	925.3	2,356.2	6,595.2
Female												
2001	97.1	11.2	1.6	0.8	1.8	4.2	12.3	44.1	144.6	481.2	1,563.4	5,977.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Major cardiovascular diseases (I00-I78)												
Hispanic												
Female												
2000 [4]	96.9	13.8	*	*	2.0	4.5	14.3	42.9	152.6	496.2	1,570.9	6,101.4
1999 [4]	99.1	17.9	1.4	*	2.0	4.0	12.7	45.7	162.6	505.6	1,655.5	6,210.9
Non-Hispanic [5]												
Both sexes												
2001	356.7	15.5	1.9	0.9	3.3	10.9	39.3	116.5	306.3	818.6	2,338.1	7,715.2
2000 [4]	365.6	17.3	1.6	0.9	3.5	10.1	38.6	118.8	323.8	857.8	2,412.2	8,076.4
1999 [4]	372.7	17.4	1.5	0.9	3.6	10.2	39.5	118.8	331.8	895.4	2,489.2	8,239.8
Male												
2001	338.1	15.8	2.0	0.9	4.1	13.7	53.0	164.7	420.6	1,066.8	2,842.6	7,990.6
2000 [4]	347.7	18.3	1.7	1.0	4.1	12.6	52.0	169.4	447.3	1,124.0	2,946.0	8,513.1
1999 [4]	354.2	19.1	1.7	1.0	4.2	12.7	53.6	169.3	457.9	1,177.2	3,041.8	8,657.6
Female												
2001	374.3	15.2	1.8	0.9	2.5	8.2	25.9	70.1	200.4	612.3	2,006.9	7,600.7
2000 [4]	382.6	16.3	1.5	0.8	2.9	7.6	25.4	70.0	209.6	638.0	2,064.5	7,899.8
1999 [4]	390.2	15.5	1.4	0.8	2.9	7.7	25.7	70.2	215.4	664.0	2,130.7	8,072.5
Non-Hispanic white												
Both sexes												
2001	385.8	12.1	1.5	0.8	2.7	8.9	33.3	101.3	278.5	779.4	2,313.0	7,811.4
2000 [4]	395.5	14.6	1.4	0.8	2.8	8.1	32.8	103.9	295.3	820.2	2,384.9	8,179.0
1999 [4]	402.7	14.2	1.3	0.8	3.0	8.3	33.3	104.1	302.5	856.8	2,462.9	8,342.9
Male												
2001	363.0	13.0	1.6	0.8	3.3	11.4	46.7	148.1	389.7	1,027.1	2,827.4	8,116.3

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Major cardiovascular diseases (I00-I78)												
Non-Hispanic white												
Male												
2000 [4]	373.6	15.5	1.6	0.9	3.3	10.5	46.1	153.2	415.4	1,088.2	2,930.7	8,650.5
1999 [4]	379.8	16.1	1.5	0.8	3.4	10.6	46.9	152.7	424.4	1,139.8	3,030.2	8,776.1
Female												
2001	407.6	11.0	1.5	0.8	2.0	6.4	19.9	55.3	173.3	569.4	1,971.8	7,684.6
2000 [4]	416.5	13.6	1.1	0.6	2.2	5.7	19.5	55.3	182.0	594.6	2,026.2	7,988.9
1999 [4]	424.7	12.1	1.1	0.7	2.5	6.0	19.8	56.2	187.7	620.0	2,091.4	8,170.1
Non-Hispanic black												
Both sexes												
2001	289.7	30.1	3.7	1.3	6.8	23.5	80.7	243.2	587.2	1,293.5	2,915.8	7,366.6
2000 [4]	294.2	28.8	2.5	1.5	7.0	22.3	79.1	246.4	609.6	1,333.1	2,995.1	7,576.5
1999 [4]	298.5	30.2	2.6	1.4	7.0	21.6	83.7	246.8	623.7	1,382.8	3,044.9	7,624.4
Male												
2001	281.2	29.7	3.8	1.2	8.3	28.9	100.1	319.1	772.1	1,639.6	3,461.6	7,413.8
2000 [4]	284.3	30.0	2.3	1.4	7.8	26.5	97.6	324.0	807.5	1,658.1	3,527.8	7,755.5
1999 [4]	291.1	31.5	2.7	1.6	8.7	26.6	105.2	329.7	833.1	1,738.6	3,542.5	7,952.2
Female												
2001	297.4	30.6	3.6	1.4	5.3	18.6	63.5	178.1	438.5	1,049.3	2,609.4	7,349.1
2000 [4]	303.1	27.5	2.9	1.5	6.3	18.6	62.8	179.8	450.3	1,104.6	2,695.3	7,510.0
1999 [4]	305.1	28.9	2.4	1.3	5.3	17.2	64.7	176.0	455.9	1,131.8	2,766.3	7,501.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13,I20-I51)												
All origins [3]												
Both sexes												
2001	245.8	11.9	1.5	0.7	2.5	8.0	29.6	92.9	246.9	635.1	1,725.7	5,664.2
2000 [4]	252.6	13.0	1.2	0.7	2.6	7.4	29.2	94.2	261.2	665.6	1,780.3	5,926.1
1999 [4]	259.9	13.8	1.2	0.7	2.8	7.6	30.2	95.7	269.9	701.7	1,849.9	6,063.0
Male												
2001	242.5	11.8	1.5	0.7	3.2	10.3	41.7	136.6	349.8	851.3	2,177.3	6,040.5
2000 [4]	249.8	13.3	1.4	0.8	3.2	9.6	41.4	140.2	371.7	898.3	2,248.1	6,430.0
1999 [4]	257.0	13.9	1.3	0.8	3.4	9.8	43.0	142.0	383.0	950.6	2,336.4	6,561.6
Female												
2001	249.0	12.0	1.4	0.7	1.8	5.6	17.6	50.7	151.8	455.9	1,428.9	5,506.8
2000 [4]	255.3	12.5	1.0	0.5	2.1	5.2	17.2	49.8	159.3	474.0	1,475.1	5,720.9
1999 [4]	262.6	13.7	1.1	0.6	2.1	5.4	17.5	51.0	165.6	497.9	1,533.9	5,862.1
Hispanic												
Both sexes												
2001	73.3	9.7	1.1	0.5	1.7	4.6	13.8	55.2	170.7	504.5	1,376.3	4,607.5
2000 [4]	73.1	11.2	1.0	0.6	1.9	4.3	14.8	55.9	183.3	506.5	1,395.9	4,701.8
1999 [4]	76.2	12.7	1.2	0.6	1.8	4.5	15.4	60.9	192.7	536.3	1,467.8	4,888.6
Male												
2001	74.6	10.3	*	*	2.1	6.1	20.0	82.9	242.2	683.7	1,702.7	4,784.3
2000 [4]	74.7	11.2	1.4	0.7	2.4	5.7	20.7	84.3	264.8	684.8	1,733.2	4,897.5
1999 [4]	77.7	10.3	1.4	0.7	2.1	6.5	22.5	92.2	272.1	737.0	1,816.0	5,012.2
Female												
2001	71.8	9.0	*	*	1.2	2.8	7.1	27.9	107.2	363.1	1,155.7	4,521.1

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13,I20-I51)												
Hispanic												
Female												
2000 [4]	71.5	11.2	*	*	1.3	2.7	8.5	28.2	111.2	366.3	1,169.4	4,605.8
1999 [4]	74.6	15.2	*	*	1.5	2.3	7.9	30.5	122.5	379.0	1,236.1	4,828.9
Non-Hispanic [5]												
Both sexes												
2001	270.7	12.3	1.5	0.7	2.6	8.7	31.6	96.0	251.6	641.4	1,736.4	5,688.6
2000 [4]	277.3	13.3	1.2	0.7	2.8	8.0	30.8	97.1	265.7	672.7	1,791.1	5,953.6
1999 [4]	284.3	14.0	1.2	0.7	2.9	8.2	31.8	98.3	274.1	708.5	1,859.2	6,086.3
Male												
2001	268.0	12.0	1.7	0.7	3.3	11.2	44.6	141.1	356.1	858.7	2,192.5	6,077.8
2000 [4]	275.2	13.7	1.4	0.8	3.3	10.3	43.8	144.5	377.5	907.0	2,263.4	6,474.6
1999 [4]	282.0	14.8	1.3	0.8	3.6	10.5	45.4	145.7	388.7	958.3	2,349.8	6,602.0
Female												
2001	273.3	12.7	1.4	0.7	1.9	6.1	18.9	52.6	154.8	460.9	1,437.0	5,526.8
2000 [4]	279.3	12.8	1.1	0.6	2.2	5.7	18.1	51.5	162.2	479.3	1,483.4	5,742.9
1999 [4]	286.4	13.3	1.1	0.6	2.3	5.9	18.5	52.6	168.2	503.4	1,540.8	5,879.8
Non-Hispanic white												
Both sexes												
2001	294.1	9.8	1.2	0.6	2.1	7.1	27.6	86.4	233.0	616.7	1,724.0	5,776.0
2000 [4]	301.3	11.4	1.1	0.5	2.1	6.4	27.2	87.7	246.4	648.1	1,777.4	6,047.9
1999 [4]	308.2	11.3	1.0	0.6	2.4	6.7	27.6	88.7	254.0	683.2	1,846.1	6,176.6
Male												
2001	289.5	10.3	1.3	0.6	2.6	9.4	40.4	130.7	335.8	834.7	2,190.4	6,195.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13,I20-I51)												
Non-Hispanic white												
Male												
2000 [4]	297.5	12.0	1.2	0.7	2.6	8.6	40.1	134.3	356.3	885.1	2,261.9	6,606.6
1999 [4]	304.0	12.4	1.1	0.6	2.8	8.7	40.6	135.0	365.6	935.2	2,352.6	6,713.7
Female												
2001	298.4	9.2	1.1	0.6	1.5	4.7	14.8	42.7	136.0	431.8	1,414.7	5,601.6
2000 [4]	304.9	10.8	0.9	0.4	1.7	4.2	14.3	41.9	142.9	448.5	1,458.9	5,822.7
1999 [4]	312.3	10.2	0.8	0.6	1.9	4.6	14.5	43.1	149.0	472.3	1,514.5	5,962.4
Non-Hispanic black												
Both sexes												
2001	215.5	22.9	3.1	1.1	5.6	18.7	61.3	183.6	454.5	977.9	2,123.4	5,307.7
2000 [4]	218.6	20.5	2.0	1.2	5.9	17.9	58.7	185.4	473.2	1,014.8	2,177.9	5,424.6
1999 [4]	223.8	24.8	2.1	1.2	6.0	17.5	63.6	189.1	488.5	1,058.0	2,223.2	5,514.7
Male												
2001	215.7	20.7	3.4	1.0	7.1	23.6	78.9	247.6	610.5	1,259.3	2,586.9	5,495.4
2000 [4]	217.6	20.1	2.0	1.2	6.9	22.1	75.6	252.5	639.6	1,281.7	2,615.1	5,677.2
1999 [4]	224.9	24.6	2.3	1.4	7.7	22.1	83.8	259.8	668.4	1,355.1	2,631.6	5,877.1
Female												
2001	215.4	25.2	2.9	1.1	4.0	14.2	45.7	128.7	328.9	779.4	1,863.2	5,238.2
2000 [4]	219.5	20.8	2.0	1.1	4.9	14.1	43.7	127.8	339.3	827.1	1,931.9	5,330.8
1999 [4]	222.8	25.0	1.9	1.0	4.4	13.3	45.8	128.7	344.3	848.5	1,994.5	5,379.3

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
All origins [3]												
Both sexes												
2001	1.2	*	*	*	*	0.1	0.3	0.6	1.4	4.1	9.4	17.7
2000 [4]	1.3	*	*	*	*	0.1	0.3	0.7	1.7	4.5	9.5	16.8
1999 [4]	1.3	*	*	*	*	0.1	0.3	0.7	1.7	4.6	10.0	18.0
Male												
2001	0.7	*	*	*	*	*	0.2	0.5	1.1	2.8	6.6	12.8
2000 [4]	0.8	*	*	*	*	*	0.3	0.6	1.4	2.9	6.6	12.2
1999 [4]	0.8	*	*	*	*	*	0.2	0.5	1.2	3.5	6.8	10.3
Female												
2001	1.7	*	*	*	*	*	0.3	0.7	1.7	5.1	11.3	19.8
2000 [4]	1.8	*	*	*	*	0.1	0.4	0.8	2.0	5.8	11.4	18.7
1999 [4]	1.9	*	*	*	*	0.1	0.3	0.8	2.2	5.6	12.2	21.1
Hispanic												
Both sexes												
2001	0.4	*	*	*	*	*	*	*	1.5	4.2	5.4	*
2000 [4]	0.4	*	*	*	*	*	*	*	1.8	4.4	5.3	*
1999 [4]	0.5	*	*	*	*	*	*	0.8	1.6	4.2	7.3	*
Male												
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.6	*	*	*	*	*	*	*	2.0	5.2	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Hispanic												
Female												
2000 [4]	0.6	*	*	*	*	*	*	*	*	5.6	6.9	*
1999 [4]	0.8	*	*	*	*	*	*	*	*	5.9	10.4	*
Non-Hispanic [5]												
Both sexes												
2001	1.3	*	*	*	*	0.1	0.3	0.6	1.4	4.1	9.6	18.0
2000 [4]	1.4	*	*	*	*	0.1	0.3	0.7	1.7	4.5	9.7	17.1
1999 [4]	1.4	*	*	*	*	0.1	0.3	0.7	1.7	4.6	10.1	18.3
Male												
2001	0.8	*	*	*	*	*	0.3	0.5	1.1	2.8	6.7	13.1
2000 [4]	0.9	*	*	*	*	*	0.3	0.6	1.4	2.9	6.7	12.5
1999 [4]	0.8	*	*	*	*	*	0.2	0.5	1.2	3.6	6.9	10.7
Female												
2001	1.8	*	*	*	*	*	0.3	0.7	1.7	5.1	11.5	20.0
2000 [4]	1.9	*	*	*	*	*	0.4	0.8	2.0	5.8	11.6	19.0
1999 [4]	2.0	*	*	*	*	0.1	0.3	0.8	2.3	5.5	12.2	21.3
Non-Hispanic white												
Both sexes												
2001	1.5	*	*	*	*	*	0.2	0.5	1.3	4.0	10.2	19.0
2000 [4]	1.6	*	*	*	*	*	0.3	0.6	1.7	4.6	10.3	18.0
1999 [4]	1.6	*	*	*	*	0.1	0.2	0.6	1.7	4.8	10.6	19.2
Male												
2001	0.9	*	*	*	*	*	0.2	0.4	1.1	2.7	7.0	13.7

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Non-Hispanic white												
Male												
2000 [4]	0.9	*	*	*	*	*	0.3	0.5	1.5	2.9	7.2	13.0
1999 [4]	0.9	*	*	*	*	*	0.2	0.5	1.1	3.8	7.2	11.6
Female												
2001	2.0	*	*	*	*	*	0.2	0.5	1.5	5.1	12.2	21.2
2000 [4]	2.1	*	*	*	*	*	0.3	0.7	1.9	6.0	12.4	20.0
1999 [4]	2.2	*	*	*	*	*	0.2	0.7	2.2	5.7	12.9	22.3
Non-Hispanic black												
Both sexes												
2001	0.8	*	*	*	*	*	0.7	1.4	2.1	4.2	4.3	6.9
2000 [4]	0.7	*	*	*	*	*	0.6	1.0	2.0	3.6	4.0	8.6
1999 [4]	0.8	*	*	*	*	*	0.5	1.2	2.2	3.4	4.4	8.4
Male												
2001	0.5	*	*	*	*	*	*	1.0	*	*	*	*
2000 [4]	0.5	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.5	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	1.2	*	*	*	*	*	0.9	1.7	2.6	5.4	5.4	*
2000 [4]	1.0	*	*	*	*	*	0.8	1.4	2.7	4.4	4.9	*
1999 [4]	1.0	*	*	*	*	*	*	1.5	2.7	4.1	4.4	11.1

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart disease (I11)												
All origins [3]												
Both sexes												
2001	8.7	*	*	*	0.1	0.6	2.4	6.3	10.9	19.8	48.7	189.5
2000 [4]	8.4	*	*	*	0.1	0.5	2.2	5.6	11.0	19.6	48.2	193.1
1999 [4]	8.1	*	*	*	0.1	0.5	2.1	5.3	10.7	19.1	47.7	184.9
Male												
2001	7.5	*	*	*	*	0.8	3.3	8.6	14.4	23.6	48.4	134.4
2000 [4]	7.2	*	*	*	0.1	0.7	2.9	7.8	14.5	23.7	46.5	139.1
1999 [4]	6.8	*	*	*	0.1	0.7	2.9	7.0	13.7	21.9	46.8	133.6
Female												
2001	9.8	*	*	*	*	0.4	1.6	4.1	7.8	16.6	48.8	212.6
2000 [4]	9.7	*	*	*	*	0.4	1.5	3.5	7.8	16.2	49.4	215.1
1999 [4]	9.4	*	*	*	*	0.3	1.3	3.6	7.9	16.8	48.3	205.5
Hispanic												
Both sexes												
2001	3.3	*	*	*	*	0.3	1.7	4.9	10.0	21.0	46.9	151.9
2000 [4]	2.8	*	*	*	*	0.3	1.2	3.8	8.2	18.4	46.0	143.3
1999 [4]	2.7	*	*	*	*	*	1.2	3.3	8.9	16.8	44.8	152.2
Male												
2001	3.4	*	*	*	*	*	2.7	7.0	13.8	26.4	48.3	144.7
2000 [4]	2.7	*	*	*	*	*	1.6	5.4	10.7	22.4	50.6	112.9
1999 [4]	2.6	*	*	*	*	*	1.8	4.7	12.5	20.9	42.8	107.8
Female												
2001	3.1	*	*	*	*	*	*	2.7	6.5	16.8	45.9	155.5

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart disease (I11)												
Hispanic												
Female												
2000 [4]	2.9	*	*	*	*	*	0.8	2.1	6.1	15.3	42.9	158.3
1999 [4]	2.9	*	*	*	*	*	*	2.0	5.8	13.6	46.1	173.7
Non-Hispanic [5]												
Both sexes												
2001	9.4	*	*	*	*	0.7	2.5	6.3	10.9	19.6	48.6	190.3
2000 [4]	9.2	*	*	*	0.1	0.6	2.3	5.7	11.2	19.5	48.1	194.3
1999 [4]	8.8	*	*	*	0.1	0.6	2.2	5.4	10.7	19.1	47.6	185.3
Male												
2001	8.1	*	*	*	*	0.9	3.4	8.6	14.2	23.3	48.3	133.4
2000 [4]	7.8	*	*	*	0.1	0.7	3.0	7.9	14.6	23.5	46.0	139.6
1999 [4]	7.4	*	*	*	0.1	0.8	3.1	7.2	13.6	21.7	46.7	134.2
Female												
2001	10.7	*	*	*	*	0.4	1.7	4.2	7.8	16.5	48.8	214.0
2000 [4]	10.6	*	*	*	*	0.4	1.6	3.6	8.0	16.2	49.5	216.5
1999 [4]	10.2	*	*	*	*	0.3	1.4	3.7	8.0	16.9	48.2	205.8
Non-Hispanic white												
Both sexes												
2001	8.6	*	*	*	*	0.4	1.6	4.1	7.6	14.8	42.2	180.4
2000 [4]	8.3	*	*	*	*	0.3	1.5	3.6	7.5	14.5	41.2	182.5
1999 [4]	7.9	*	*	*	*	0.3	1.3	3.4	7.3	14.1	40.6	175.2
Male												
2001	6.9	*	*	*	*	0.6	2.3	5.9	10.5	18.1	41.5	122.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart disease (I11)												
Non-Hispanic white												
Male												
2000 [4]	6.6	*	*	*	*	0.4	2.1	5.4	10.3	18.0	38.8	129.5
1999 [4]	6.2	*	*	*	*	0.6	2.0	4.9	9.6	16.2	39.4	123.8
Female												
2001	10.2	*	*	*	*	0.2	0.8	2.3	4.9	11.9	42.6	204.7
2000 [4]	9.9	*	*	*	*	0.2	0.8	1.9	4.9	11.6	42.7	203.9
1999 [4]	9.6	*	*	*	*	*	0.6	2.0	5.0	12.4	41.4	195.7
Non-Hispanic black												
Both sexes												
2001	16.5	*	*	*	*	2.2	8.5	23.2	39.4	66.1	125.6	328.7
2000 [4]	16.6	*	*	*	*	2.2	7.5	21.6	43.1	67.6	132.7	344.1
1999 [4]	16.0	*	*	*	0.4	1.9	8.1	21.3	40.1	67.4	132.7	308.5
Male												
2001	16.3	*	*	*	*	2.7	10.5	30.4	50.1	79.8	137.4	288.7
2000 [4]	16.1	*	*	*	*	2.7	9.2	28.4	56.0	81.9	139.5	275.9
1999 [4]	15.6	*	*	*	*	2.2	10.3	26.8	51.1	80.6	143.2	266.2
Female												
2001	16.7	*	*	*	*	1.7	6.7	17.1	30.8	56.5	118.9	343.5
2000 [4]	17.1	*	*	*	*	1.8	5.9	15.8	32.7	57.5	128.9	369.4
1999 [4]	16.4	*	*	*	*	1.6	6.1	16.5	31.3	58.1	126.8	324.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart and renal disease (I13)												
All origins [3]												
Both sexes												
2001	1.0	*	*	*	*	0.1	0.2	0.4	1.0	2.3	6.8	23.3
2000 [4]	1.0	*	*	*	*	0.1	0.2	0.4	0.9	2.3	7.0	24.3
1999 [4]	1.2	*	*	*	*	0.1	0.3	0.4	1.1	2.6	8.9	29.1
Male												
2001	0.8	*	*	*	*	*	0.3	0.5	1.0	2.5	6.8	24.2
2000 [4]	0.8	*	*	*	*	*	0.2	0.5	0.9	2.5	7.2	27.8
1999 [4]	1.0	*	*	*	*	*	0.3	0.6	1.3	2.7	9.6	31.3
Female												
2001	1.1	*	*	*	*	*	0.2	0.2	0.9	2.1	6.8	22.9
2000 [4]	1.1	*	*	*	*	*	0.1	0.3	0.8	2.1	6.9	22.9
1999 [4]	1.4	*	*	*	*	*	0.2	0.3	0.9	2.4	8.5	28.2
Hispanic												
Both sexes												
2001	0.4	*	*	*	*	*	*	*	1.1	3.0	7.4	26.3
2000 [4]	0.4	*	*	*	*	*	*	*	*	2.5	6.3	25.2
1999 [4]	0.4	*	*	*	*	*	*	*	*	3.1	8.1	23.2
Male												
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.4	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.4	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.5	*	*	*	*	*	*	*	*	*	7.3	24.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart and renal disease (I13)												
Hispanic												
Female												
2000 [4]	0.4	*	*	*	*	*	*	*	*	*	*	26.7
1999 [4]	0.5	*	*	*	*	*	*	*	*	*	8.7	20.8
Non-Hispanic [5]												
Both sexes												
2001	1.1	*	*	*	*	0.1	0.2	0.4	1.0	2.2	6.7	23.1
2000 [4]	1.1	*	*	*	*	0.1	0.2	0.4	0.9	2.2	7.0	24.2
1999 [4]	1.3	*	*	*	*	0.1	0.3	0.4	1.1	2.5	8.9	29.3
Male												
2001	0.9	*	*	*	*	*	0.3	0.5	1.0	2.4	6.7	23.8
2000 [4]	0.9	*	*	*	*	*	0.2	0.5	0.9	2.4	7.0	27.9
1999 [4]	1.1	*	*	*	*	*	0.3	0.6	1.3	2.7	9.6	31.4
Female												
2001	1.2	*	*	*	*	*	0.2	0.2	0.9	2.0	6.7	22.8
2000 [4]	1.2	*	*	*	*	*	0.1	0.3	0.8	2.1	6.9	22.8
1999 [4]	1.5	*	*	*	*	*	0.2	0.3	0.9	2.4	8.4	28.4
Non-Hispanic white												
Both sexes												
2001	0.9	*	*	*	*	*	*	0.1	0.5	1.4	5.6	20.7
2000 [4]	0.9	*	*	*	*	*	0.1	0.1	0.4	1.5	5.6	21.8
1999 [4]	1.1	*	*	*	*	*	*	0.2	0.5	1.4	7.3	26.1
Male												
2001	0.7	*	*	*	*	*	*	0.2	0.4	1.5	5.8	22.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart and renal disease (I13)												
Non-Hispanic white												
Male												
2000 [4]	0.7	*	*	*	*	*	*	0.2	0.4	1.6	6.1	24.3
1999 [4]	0.9	*	*	*	*	*	*	0.2	0.5	1.6	8.3	28.7
Female												
2001	1.1	*	*	*	*	*	*	*	0.5	1.3	5.4	20.1
2000 [4]	1.1	*	*	*	*	*	*	*	0.3	1.4	5.2	20.8
1999 [4]	1.2	*	*	*	*	*	*	*	0.4	1.3	6.6	25.1
Non-Hispanic black												
Both sexes												
2001	2.3	*	*	*	*	0.4	1.2	2.1	5.1	10.1	20.2	50.6
2000 [4]	2.3	*	*	*	*	*	0.8	2.2	4.7	10.0	22.6	50.9
1999 [4]	2.8	*	*	*	*	*	1.5	2.3	6.2	12.7	27.3	66.0
Male												
2001	2.2	*	*	*	*	*	1.6	3.2	6.0	12.0	18.6	45.0
2000 [4]	2.1	*	*	*	*	*	1.0	2.9	5.4	11.1	20.1	64.6
1999 [4]	2.6	*	*	*	*	*	1.8	3.3	7.8	13.8	26.0	60.6
Female												
2001	2.4	*	*	*	*	*	0.8	1.2	4.3	8.7	21.1	52.6
2000 [4]	2.4	*	*	*	*	*	*	1.6	4.2	9.2	24.0	45.8
1999 [4]	3.1	*	*	*	*	*	1.2	1.5	4.8	12.0	28.0	68.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ischemic heart diseases (I20-I25)												
All origins [3]												
Both sexes												
2001	176.3	0.6	*	0.0	0.4	2.8	16.7	64.0	184.0	477.6	1,271.3	3,958.2
2000 [4]	183.1	0.8	*	*	0.3	2.3	16.7	66.1	196.7	505.1	1,324.2	4,182.5
1999 [4]	189.8	0.7	*	0.1	0.4	2.5	17.2	67.7	203.5	536.6	1,384.8	4,318.1
Male												
2001	181.7	*	*	*	0.6	4.1	25.5	99.5	271.6	663.7	1,663.5	4,395.8
2000 [4]	188.7	*	*	*	0.4	3.5	25.8	103.5	290.2	705.2	1,736.8	4,701.7
1999 [4]	195.4	*	*	*	0.5	3.7	26.6	106.2	299.3	749.8	1,813.7	4,837.5
Female												
2001	171.2	*	*	*	0.2	1.5	8.1	29.7	103.1	323.4	1,013.6	3,775.1
2000 [4]	177.6	*	*	*	0.2	1.2	7.6	29.9	110.4	340.3	1,055.2	3,971.0
1999 [4]	184.5	*	*	*	0.2	1.4	7.9	30.7	115.1	362.0	1,106.3	4,108.8
Hispanic												
Both sexes												
2001	55.9	*	*	*	*	1.6	6.8	38.7	130.2	400.9	1,099.1	3,620.2
2000 [4]	55.9	*	*	*	*	1.1	7.6	39.5	143.6	405.3	1,111.3	3,697.2
1999 [4]	58.7	*	*	*	*	1.4	7.4	44.8	150.4	433.5	1,178.0	3,876.2
Male												
2001	57.9	*	*	*	*	2.4	10.4	60.3	191.9	559.9	1,393.3	3,828.2
2000 [4]	58.2	*	*	*	*	1.6	12.1	62.7	214.8	561.5	1,420.0	3,865.6
1999 [4]	60.8	*	*	*	*	2.1	12.2	71.3	218.3	608.5	1,495.9	3,987.7
Female												
2001	53.7	*	*	*	*	*	2.8	17.4	75.4	275.4	900.3	3,518.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ischemic heart diseases (I20-I25)												
Hispanic												
Female												
2000 [4]	53.6	*	*	*	*	*	2.8	16.8	80.6	282.5	904.0	3,614.6
1999 [4]	56.6	*	*	*	*	*	2.3	19.0	90.5	296.4	966.5	3,822.4
Non-Hispanic [5]												
Both sexes												
2001	193.6	0.7	*	*	0.4	3.1	18.0	66.1	187.3	480.9	1,274.9	3,959.4
2000 [4]	200.5	0.8	*	*	0.3	2.6	17.7	68.1	199.5	509.0	1,328.8	4,187.0
1999 [4]	207.1	0.8	*	0.1	0.4	2.7	18.3	69.4	206.3	540.3	1,388.1	4,319.6
Male												
2001	200.4	*	*	*	0.6	4.5	27.5	102.8	276.1	667.7	1,670.1	4,408.5
2000 [4]	207.6	*	*	*	0.4	3.9	27.4	106.7	294.0	710.4	1,744.6	4,721.1
1999 [4]	214.1	*	*	*	0.5	4.0	28.3	108.8	303.4	754.2	1,820.2	4,854.5
Female												
2001	187.2	*	*	*	0.2	1.6	8.7	30.8	105.0	325.6	1,015.5	3,772.7
2000 [4]	193.8	*	*	*	0.2	1.3	8.2	31.0	112.1	342.8	1,057.9	3,971.0
1999 [4]	200.6	*	*	*	0.3	1.5	8.5	31.5	116.6	364.6	1,107.8	4,105.4
Non-Hispanic white												
Both sexes												
2001	212.6	*	*	*	0.3	2.9	17.3	62.6	177.8	467.8	1,271.3	4,022.8
2000 [4]	220.2	*	*	*	0.3	2.4	16.9	64.6	189.5	496.4	1,324.5	4,261.3
1999 [4]	227.2	*	*	*	0.3	2.5	17.3	65.8	196.4	527.7	1,386.5	4,390.2
Male												
2001	219.6	*	*	*	0.5	4.3	26.9	99.5	266.3	656.5	1,678.7	4,506.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ischemic heart diseases (I20-I25)												
Non-Hispanic white												
Male												
2000 [4]	227.6	*	*	*	0.4	3.7	26.8	103.5	283.5	701.7	1,752.3	4,833.4
1999 [4]	234.2	*	*	*	0.5	3.6	27.2	105.1	292.3	745.2	1,834.9	4,954.1
Female												
2001	205.8	*	*	*	0.2	1.4	7.6	26.4	94.2	307.7	1,001.1	3,821.4
2000 [4]	213.1	*	*	*	0.2	1.0	6.9	26.4	100.9	323.6	1,043.3	4,030.7
1999 [4]	220.5	*	*	*	*	1.4	7.4	27.2	106.1	345.6	1,092.9	4,165.3
Non-Hispanic black												
Both sexes												
2001	141.2	*	*	*	0.9	4.8	26.4	104.6	302.1	681.4	1,487.0	3,628.7
2000 [4]	144.4	*	*	*	0.4	4.4	26.1	107.7	318.9	710.1	1,534.5	3,701.7
1999 [4]	147.9	*	*	*	0.8	4.1	28.2	110.8	324.7	741.6	1,559.0	3,817.9
Male												
2001	143.2	*	*	*	1.2	6.8	37.1	149.0	418.0	903.1	1,834.9	3,791.8
2000 [4]	145.7	*	*	*	*	6.1	36.3	152.2	444.2	916.5	1,887.6	3,911.2
1999 [4]	150.0	*	*	*	0.9	6.0	40.3	160.0	458.4	967.5	1,873.3	4,082.5
Female												
2001	139.4	*	*	*	*	2.9	16.9	66.5	208.9	525.0	1,291.6	3,568.3
2000 [4]	143.2	*	*	*	*	2.9	17.1	69.4	218.1	565.0	1,335.7	3,623.9
1999 [4]	146.0	*	*	*	0.7	2.4	17.4	68.7	217.5	582.3	1,383.0	3,719.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute myocardial infarction (I21-I22)												
All origins [3]												
Both sexes												
2001	64.9	*	*	*	0.2	1.2	7.1	27.6	78.4	193.2	471.2	1,265.4
2000 [4]	68.5	0.5	*	*	0.1	0.9	6.9	28.9	86.6	209.1	503.3	1,338.5
1999 [4]	71.5	*	*	*	0.2	1.1	7.2	29.6	90.0	225.0	529.3	1,379.3
Male												
2001	68.6	*	*	*	0.2	1.7	10.6	42.4	114.0	264.7	604.1	1,461.3
2000 [4]	72.7	*	*	*	0.1	1.3	10.6	44.7	125.7	286.9	648.3	1,568.3
1999 [4]	76.1	*	*	*	0.2	1.5	11.0	46.3	131.8	310.0	684.4	1,622.1
Female												
2001	61.3	*	*	*	*	0.7	3.5	13.3	45.4	134.0	383.9	1,183.5
2000 [4]	64.6	*	*	*	0.1	0.5	3.3	13.6	50.5	145.1	408.7	1,244.8
1999 [4]	67.0	*	*	*	0.1	0.6	3.4	13.5	51.5	155.4	428.6	1,281.5
Hispanic												
Both sexes												
2001	20.0	*	*	*	*	0.7	2.5	14.8	49.5	149.8	394.9	1,185.5
2000 [4]	20.1	*	*	*	*	0.5	3.0	15.5	55.0	154.3	404.7	1,177.1
1999 [4]	21.1	*	*	*	*	0.5	2.9	17.8	56.1	164.0	433.7	1,227.8
Male												
2001	20.9	*	*	*	*	1.0	3.9	22.7	69.4	206.9	488.7	1,331.3
2000 [4]	21.1	*	*	*	*	0.7	4.7	23.4	80.0	207.6	514.8	1,273.8
1999 [4]	22.0	*	*	*	*	0.8	4.6	26.7	79.0	225.8	558.9	1,276.8
Female												
2001	19.1	*	*	*	*	*	1.1	7.0	31.8	104.7	331.5	1,114.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute myocardial infarction (I21-I22)												
Hispanic												
Female												
2000 [4]	19.1	*	*	*	*	*	1.2	7.8	32.9	112.3	330.8	1,129.7
1999 [4]	20.1	*	*	*	*	*	1.1	9.1	35.9	115.5	350.4	1,204.1
Non-Hispanic [5]												
Both sexes												
2001	71.4	*	*	*	0.2	1.3	7.6	28.8	80.3	195.6	473.6	1,265.3
2000 [4]	75.3	*	*	*	0.1	1.0	7.4	30.0	88.6	211.9	506.3	1,340.8
1999 [4]	78.2	*	*	*	0.2	1.2	7.7	30.5	92.2	227.9	531.9	1,381.0
Male												
2001	75.9	*	*	*	0.2	1.8	11.6	44.2	117.0	267.7	607.9	1,463.9
2000 [4]	80.3	*	*	*	0.2	1.5	11.4	46.5	128.7	290.9	652.6	1,576.2
1999 [4]	83.8	*	*	*	0.2	1.7	11.8	47.8	135.1	313.8	687.7	1,631.5
Female												
2001	67.1	*	*	*	*	0.8	3.8	13.9	46.3	135.7	385.4	1,182.7
2000 [4]	70.6	*	*	*	0.1	0.6	3.5	14.1	51.6	146.8	411.0	1,245.6
1999 [4]	72.9	*	*	*	*	0.7	3.7	13.9	52.5	157.3	430.7	1,280.8
Non-Hispanic white												
Both sexes												
2001	78.4	*	*	*	0.1	1.2	7.5	28.1	78.7	193.0	473.0	1,276.8
2000 [4]	82.7	*	*	*	0.1	0.9	7.3	29.3	86.9	209.3	505.3	1,355.3
1999 [4]	86.0	*	*	*	0.1	1.1	7.5	30.0	90.7	226.1	532.4	1,394.2
Male												
2001	83.7	*	*	*	0.2	1.7	11.6	44.3	116.8	267.7	612.5	1,483.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute myocardial infarction (I21-I22)												
Non-Hispanic white												
Male												
2000 [4]	88.5	*	*	*	0.2	1.4	11.4	46.7	128.0	291.0	656.7	1,600.7
1999 [4]	92.4	*	*	*	0.2	1.5	11.8	48.1	134.6	315.0	695.7	1,653.8
Female												
2001	73.4	*	*	*	*	0.8	3.4	12.2	42.7	129.6	380.6	1,190.9
2000 [4]	77.2	*	*	*	*	0.5	3.1	12.3	48.2	140.5	405.9	1,256.3
1999 [4]	79.8	*	*	*	*	0.7	3.3	12.3	49.3	151.8	425.5	1,290.7
Non-Hispanic black												
Both sexes												
2001	51.6	*	*	*	0.4	1.9	10.2	40.2	111.6	256.0	543.7	1,247.0
2000 [4]	54.0	*	*	*	*	1.8	10.1	41.9	122.3	273.0	576.6	1,281.8
1999 [4]	55.1	*	*	*	*	1.7	10.5	42.0	123.2	282.4	589.9	1,328.6
Male												
2001	50.9	*	*	*	*	2.5	13.5	53.4	146.1	320.8	646.6	1,389.4
2000 [4]	53.7	*	*	*	*	2.2	13.6	56.3	163.1	341.2	693.3	1,426.6
1999 [4]	55.2	*	*	*	*	2.5	14.5	58.1	168.7	357.0	691.4	1,509.4
Female												
2001	52.3	*	*	*	*	1.3	7.4	28.8	83.8	210.3	485.9	1,194.3
2000 [4]	54.2	*	*	*	*	1.4	7.0	29.6	89.5	225.0	510.9	1,228.0
1999 [4]	54.9	*	*	*	*	1.0	7.0	28.2	86.8	229.7	533.1	1,261.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute ischemic heart diseases (I24)												
All origins [3]												
Both sexes												
2001	1.2	*	*	*	*	*	0.2	0.8	1.8	3.5	6.9	21.5
2000 [4]	1.2	*	*	*	*	*	0.2	0.7	2.0	3.5	7.3	21.9
1999 [4]	1.2	*	*	*	*	*	0.2	0.7	1.9	3.7	7.0	21.9
Male												
2001	1.2	*	*	*	*	*	0.3	1.2	2.6	4.7	8.5	20.7
2000 [4]	1.3	*	*	*	*	*	0.3	1.2	2.9	4.8	9.1	20.7
1999 [4]	1.2	*	*	*	*	*	0.3	1.0	2.8	4.9	8.8	22.3
Female												
2001	1.1	*	*	*	*	*	0.1	0.4	1.1	2.5	5.9	21.8
2000 [4]	1.1	*	*	*	*	*	0.1	0.3	1.1	2.4	6.1	22.4
1999 [4]	1.1	*	*	*	*	*	*	0.4	1.1	2.6	5.8	21.7
Hispanic												
Both sexes												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute ischemic heart diseases (I24)												
Hispanic												
Female												
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	1.3	*	*	*	*	*	0.2	0.8	1.9	3.6	7.1	22.0
2000 [4]	1.3	*	*	*	*	*	0.2	0.8	2.1	3.6	7.5	22.2
1999 [4]	1.3	*	*	*	*	*	0.2	0.7	2.0	3.8	7.2	22.2
Male												
2001	1.4	*	*	*	*	*	0.3	1.3	2.7	4.8	8.7	20.9
2000 [4]	1.4	*	*	*	*	*	0.3	1.3	3.1	4.9	9.3	21.0
1999 [4]	1.4	*	*	*	*	*	0.3	1.1	2.9	5.2	9.0	22.9
Female												
2001	1.3	*	*	*	*	*	0.1	0.4	1.1	2.6	6.0	22.4
2000 [4]	1.2	*	*	*	*	*	0.1	0.3	1.2	2.5	6.3	22.7
1999 [4]	1.2	*	*	*	*	*	*	0.4	1.1	2.7	6.0	21.9
Non-Hispanic white												
Both sexes												
2001	1.4	*	*	*	*	*	0.2	0.8	1.7	3.4	7.0	22.0
2000 [4]	1.4	*	*	*	*	*	0.2	0.7	1.9	3.4	7.3	22.5
1999 [4]	1.4	*	*	*	*	*	0.2	0.6	1.8	3.6	7.0	22.7
Male												
2001	1.4	*	*	*	*	*	0.3	1.2	2.5	4.5	8.7	20.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute ischemic heart diseases (I24)												
Non-Hispanic white												
Male												
2000 [4]	1.5	*	*	*	*	*	0.3	1.2	2.8	4.6	9.0	21.3
1999 [4]	1.4	*	*	*	*	*	0.2	0.9	2.6	4.9	8.7	23.6
Female												
2001	1.4	*	*	*	*	*	*	0.3	1.0	2.5	5.9	22.4
2000 [4]	1.3	*	*	*	*	*	*	0.3	1.0	2.4	6.1	23.0
1999 [4]	1.3	*	*	*	*	*	*	0.3	1.1	2.6	5.8	22.4
Non-Hispanic black												
Both sexes												
2001	1.3	*	*	*	*	*	0.5	1.6	4.0	5.8	8.5	24.3
2000 [4]	1.3	*	*	*	*	*	0.4	1.5	4.3	5.6	10.8	21.9
1999 [4]	1.2	*	*	*	*	*	0.5	1.9	3.6	5.8	10.9	17.5
Male												
2001	1.5	*	*	*	*	*	*	2.5	5.6	8.8	10.5	24.3
2000 [4]	1.5	*	*	*	*	*	*	2.3	6.3	8.7	14.2	*
1999 [4]	1.5	*	*	*	*	*	0.8	2.7	6.1	8.4	12.8	*
Female												
2001	1.1	*	*	*	*	*	*	*	2.6	3.8	7.4	24.4
2000 [4]	1.1	*	*	*	*	*	*	*	2.7	3.5	8.9	22.2
1999 [4]	1.0	*	*	*	*	*	*	1.1	1.6	3.9	9.8	16.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)												
All origins [3]												
Both sexes												
2001	110.3	*	*	*	0.2	1.6	9.5	35.6	103.8	280.9	793.1	2,671.3
2000 [4]	113.3	*	*	*	0.2	1.4	9.5	36.4	108.1	292.4	813.6	2,822.1
1999 [4]	117.2	*	*	*	0.2	1.4	9.8	37.5	111.5	308.0	848.5	2,917.0
Male												
2001	111.9	*	*	*	0.3	2.4	14.6	55.9	154.9	394.3	1,050.9	2,913.8
2000 [4]	114.8	*	*	*	0.2	2.1	14.8	57.7	161.6	413.5	1,079.4	3,112.7
1999 [4]	118.0	*	*	*	0.3	2.1	15.3	59.0	164.7	434.9	1,120.5	3,193.1
Female												
2001	108.8	*	*	*	0.1	0.7	4.4	16.0	56.7	186.9	623.7	2,569.8
2000 [4]	111.9	*	*	*	*	0.6	4.3	16.0	58.8	192.8	640.3	2,703.8
1999 [4]	116.4	*	*	*	0.1	0.7	4.3	16.7	62.6	204.0	671.8	2,805.7
Hispanic												
Both sexes												
2001	35.7	*	*	*	*	0.9	4.2	23.8	80.0	249.4	701.7	2,426.6
2000 [4]	35.6	*	*	*	*	0.6	4.6	23.9	88.0	249.7	703.2	2,508.8
1999 [4]	37.5	*	*	*	*	0.9	4.5	26.8	93.5	268.8	741.8	2,640.7
Male												
2001	36.8	*	*	*	*	1.4	6.5	37.5	121.4	350.2	901.5	2,479.8
2000 [4]	36.9	*	*	*	*	0.9	7.4	39.1	134.2	351.6	900.8	2,577.7
1999 [4]	38.7	*	*	*	*	1.3	7.6	44.3	138.5	381.8	932.8	2,706.7
Female												
2001	34.5	*	*	*	*	*	1.7	10.3	43.3	169.9	566.7	2,400.7

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)												
Hispanic												
Female												
2000 [4]	34.2	*	*	*	*	*	1.7	8.9	47.2	169.6	570.5	2,475.0
1999 [4]	36.3	*	*	*	*	*	1.3	9.7	53.8	180.2	614.8	2,608.9
Non-Hispanic [5]												
Both sexes												
2001	120.9	*	*	*	0.2	1.7	10.2	36.5	105.1	281.6	794.2	2,672.2
2000 [4]	123.9	*	*	*	0.2	1.5	10.1	37.3	108.8	293.5	815.0	2,824.0
1999 [4]	127.6	*	*	*	0.2	1.5	10.4	38.1	112.1	308.6	849.1	2,916.3
Male												
2001	123.1	*	*	*	0.3	2.6	15.7	57.3	156.4	395.1	1,053.5	2,923.7
2000 [4]	126.0	*	*	*	0.2	2.3	15.7	58.9	162.3	414.7	1,082.7	3,123.9
1999 [4]	128.9	*	*	*	0.3	2.3	16.2	59.9	165.4	435.2	1,123.5	3,200.1
Female												
2001	118.8	*	*	*	0.1	0.8	4.8	16.5	57.5	187.3	624.1	2,567.6
2000 [4]	122.0	*	*	*	*	0.7	4.6	16.5	59.3	193.5	640.6	2,702.7
1999 [4]	126.4	*	*	*	0.1	0.8	4.7	17.3	62.9	204.6	671.1	2,802.7
Non-Hispanic white												
Both sexes												
2001	132.7	*	*	*	0.2	1.6	9.6	33.8	97.4	271.4	791.2	2,724.1
2000 [4]	136.1	*	*	*	0.2	1.4	9.4	34.6	100.7	283.7	811.9	2,883.6
1999 [4]	139.9	*	*	*	0.2	1.4	9.6	35.2	103.9	297.9	847.2	2,973.2
Male												
2001	134.4	*	*	*	0.3	2.5	15.0	54.1	147.0	384.3	1,057.5	3,002.8

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)												
Non-Hispanic white												
Male												
2000 [4]	137.7	*	*	*	0.2	2.2	15.1	55.7	152.7	406.1	1,086.5	3,211.3
1999 [4]	140.4	*	*	*	0.2	2.1	15.2	56.1	155.1	425.3	1,130.5	3,276.7
Female												
2001	131.1	*	*	*	*	0.7	4.1	13.9	50.5	175.6	614.6	2,608.1
2000 [4]	134.6	*	*	*	*	0.5	3.8	13.8	51.8	180.6	631.3	2,751.4
1999 [4]	139.3	*	*	*	*	0.7	4.0	14.6	55.7	191.3	661.7	2,852.2
Non-Hispanic black												
Both sexes												
2001	88.2	*	*	*	0.5	2.9	15.6	62.9	186.6	419.6	934.8	2,357.3
2000 [4]	89.1	*	*	*	*	2.5	15.6	64.3	192.3	431.5	947.1	2,398.0
1999 [4]	91.6	*	*	*	0.4	2.4	17.2	66.9	197.9	453.5	958.2	2,471.9
Male												
2001	90.8	*	*	*	*	4.3	22.9	93.1	266.3	573.5	1,177.8	2,378.1
2000 [4]	90.4	*	*	*	*	3.7	22.1	93.6	274.7	566.6	1,180.1	2,463.5
1999 [4]	93.3	*	*	*	*	3.5	25.1	99.2	283.6	602.1	1,169.2	2,552.9
Female												
2001	86.0	*	*	*	*	1.6	9.2	36.9	122.4	310.9	798.4	2,349.6
2000 [4]	88.0	*	*	*	*	1.5	10.0	39.1	125.9	336.5	815.9	2,373.7
1999 [4]	90.1	*	*	*	*	1.3	10.2	39.4	129.2	348.6	840.0	2,441.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)												
All origins [3]												
Both sexes												
2001	23.9	*	*	*	0.1	0.5	3.2	12.9	33.3	65.6	152.7	493.6
2000 [4]	24.7	*	*	*	*	0.4	3.0	13.2	34.0	68.3	158.1	534.6
1999 [4]	25.8	*	*	*	*	0.4	3.3	13.0	35.3	72.7	166.2	563.4
Male												
2001	25.3	*	*	*	*	0.7	5.0	21.0	51.4	92.4	193.9	502.0
2000 [4]	26.0	*	*	*	*	0.6	4.9	21.5	52.6	97.2	199.3	550.8
1999 [4]	26.8	*	*	*	*	0.6	5.3	20.9	53.7	103.0	207.0	578.6
Female												
2001	22.5	*	*	*	*	0.2	1.4	5.1	16.7	43.3	125.5	490.1
2000 [4]	23.5	*	*	*	*	0.1	1.2	5.2	16.8	44.6	131.3	528.0
1999 [4]	24.8	*	*	*	*	0.2	1.3	5.3	18.3	47.9	139.8	557.2
Hispanic												
Both sexes												
2001	8.5	*	*	*	*	0.4	1.5	10.1	29.7	65.9	132.5	417.0
2000 [4]	8.4	*	*	*	*	*	1.6	9.5	30.5	65.7	128.2	452.5
1999 [4]	9.0	*	*	*	*	*	1.5	10.4	33.0	72.1	147.5	469.4
Male												
2001	10.1	*	*	*	*	0.6	2.2	17.1	47.9	100.2	181.1	440.0
2000 [4]	9.9	*	*	*	*	*	2.7	16.6	50.5	92.4	180.9	493.8
1999 [4]	10.6	*	*	*	*	*	2.5	17.8	50.0	115.8	194.0	485.3
Female												
2001	6.8	*	*	*	*	*	*	3.2	13.5	38.7	99.7	405.8

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)												
Hispanic												
Female												
2000 [4]	6.8	*	*	*	*	*	*	2.6	12.8	44.6	92.8	432.3
1999 [4]	7.4	*	*	*	*	*	*	3.2	18.0	37.9	116.6	461.7
Non-Hispanic [5]												
Both sexes												
2001	26.0	*	*	*	*	0.5	3.5	13.0	33.3	65.0	152.6	494.8
2000 [4]	26.9	*	*	*	*	0.4	3.2	13.4	33.8	67.9	158.3	535.9
1999 [4]	27.9	*	*	*	*	0.4	3.5	13.1	35.0	72.0	165.7	564.9
Male												
2001	27.5	*	*	*	*	0.7	5.4	21.2	51.1	91.0	193.2	503.0
2000 [4]	28.2	*	*	*	*	0.7	5.1	21.7	52.1	96.4	198.5	551.0
1999 [4]	28.9	*	*	*	*	0.6	5.6	21.0	53.3	101.1	205.6	580.2
Female												
2001	24.6	*	*	*	*	0.3	1.5	5.2	16.8	43.3	126.0	491.4
2000 [4]	25.6	*	*	*	*	0.2	1.3	5.4	16.9	44.4	132.2	529.8
1999 [4]	27.0	*	*	*	*	0.2	1.4	5.4	18.2	48.2	139.8	558.8
Non-Hispanic white												
Both sexes												
2001	26.8	*	*	*	*	0.4	3.2	11.6	28.4	57.1	144.2	490.7
2000 [4]	27.6	*	*	*	*	0.4	2.9	11.9	28.9	60.2	149.0	530.2
1999 [4]	28.6	*	*	*	*	0.4	3.1	11.3	29.8	63.6	156.2	558.8
Male												
2001	27.8	*	*	*	*	0.6	5.0	19.1	44.3	80.4	182.9	499.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)												
Non-Hispanic white												
Male												
2000 [4]	28.4	*	*	*	*	0.6	4.8	19.6	45.2	86.3	186.1	544.7
1999 [4]	29.0	*	*	*	*	0.6	5.2	18.4	46.0	90.0	193.5	572.8
Female												
2001	25.8	*	*	*	*	0.2	1.3	4.1	13.3	37.3	118.6	487.0
2000 [4]	26.9	*	*	*	*	*	1.1	4.3	13.4	38.2	124.6	524.4
1999 [4]	28.2	*	*	*	*	0.2	1.1	4.2	14.6	41.6	131.8	553.2
Non-Hispanic black												
Both sexes												
2001	28.5	*	*	*	*	0.7	5.8	25.7	78.1	144.8	266.7	595.9
2000 [4]	29.3	*	*	*	*	0.7	5.3	27.3	78.9	147.2	280.2	652.2
1999 [4]	30.7	*	*	*	*	0.8	6.5	28.1	82.0	158.7	291.0	670.3
Male												
2001	32.3	*	*	*	*	1.0	8.8	39.9	118.1	209.4	352.3	609.8
2000 [4]	33.2	*	*	*	*	1.1	7.9	42.3	120.8	209.3	371.5	711.6
1999 [4]	34.5	*	*	*	*	1.1	9.8	43.6	123.7	228.2	378.4	714.3
Female												
2001	25.0	*	*	*	*	*	3.2	13.6	45.9	99.3	218.7	590.7
2000 [4]	25.8	*	*	*	*	*	2.9	14.4	45.1	103.4	228.9	630.2
1999 [4]	27.1	*	*	*	*	*	3.6	14.7	48.5	109.7	242.1	653.8

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20, I25.1-I25.9)												
All origins [3]												
Both sexes												
2001	86.4	*	*	*	0.2	1.1	6.2	22.7	70.5	215.3	640.5	2,177.7
2000 [4]	88.6	*	*	*	0.1	1.0	6.5	23.2	74.1	224.1	655.5	2,287.5
1999 [4]	91.4	*	*	*	0.2	1.0	6.5	24.5	76.3	235.2	682.3	2,353.6
Male												
2001	86.5	*	*	*	0.2	1.8	9.5	34.9	103.5	301.9	856.9	2,411.8
2000 [4]	88.8	*	*	*	0.2	1.5	9.9	36.2	109.0	316.3	880.1	2,561.9
1999 [4]	91.2	*	*	*	0.2	1.5	10.0	38.1	111.0	331.9	913.6	2,614.4
Female												
2001	86.2	*	*	*	0.1	0.5	3.0	11.0	40.0	143.6	498.2	2,079.8
2000 [4]	88.4	*	*	*	*	0.5	3.0	10.7	42.0	148.2	509.0	2,175.8
1999 [4]	91.6	*	*	*	0.1	0.5	3.0	11.4	44.3	156.1	532.1	2,248.4
Hispanic												
Both sexes												
2001	27.2	*	*	*	*	0.6	2.7	13.7	50.4	183.6	569.1	2,009.6
2000 [4]	27.2	*	*	*	*	0.5	3.0	14.3	57.5	184.0	575.0	2,056.3
1999 [4]	28.5	*	*	*	*	0.6	3.0	16.4	60.5	196.7	594.3	2,171.4
Male												
2001	26.7	*	*	*	*	0.9	4.3	20.4	73.6	250.0	720.4	2,039.8
2000 [4]	27.0	*	*	*	*	0.7	4.7	22.6	83.6	259.1	719.8	2,084.0
1999 [4]	28.1	*	*	*	*	1.0	5.1	26.5	88.5	266.0	738.8	2,221.4
Female												
2001	27.7	*	*	*	*	*	1.0	7.1	29.8	131.2	466.9	1,994.8

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20, I25.1-I25.9)												
Hispanic												
Female												
2000 [4]	27.5	*	*	*	*	*	1.2	6.2	34.4	124.9	477.7	2,042.7
1999 [4]	28.9	*	*	*	*	*	0.8	6.5	35.8	142.3	498.2	2,147.2
Non-Hispanic [5]												
Both sexes												
2001	94.9	*	*	*	0.2	1.2	6.7	23.5	71.8	216.7	641.6	2,177.4
2000 [4]	97.0	*	*	*	0.1	1.1	6.9	23.9	75.0	225.6	656.7	2,288.1
1999 [4]	99.7	*	*	*	0.2	1.1	6.9	25.1	77.1	236.5	683.4	2,351.4
Male												
2001	95.6	*	*	*	0.2	2.0	10.3	36.1	105.3	304.1	860.3	2,420.7
2000 [4]	97.8	*	*	*	0.2	1.7	10.5	37.2	110.2	318.2	884.2	2,572.9
1999 [4]	100.0	*	*	*	0.2	1.6	10.6	38.9	112.1	334.1	917.9	2,619.9
Female												
2001	94.2	*	*	*	*	0.5	3.2	11.3	40.7	143.9	498.0	2,076.2
2000 [4]	96.3	*	*	*	*	0.5	3.3	11.1	42.4	149.1	508.5	2,172.9
1999 [4]	99.4	*	*	*	*	0.6	3.3	11.8	44.8	156.4	531.2	2,243.9
Non-Hispanic white												
Both sexes												
2001	106.0	*	*	*	0.1	1.1	6.4	22.2	69.0	214.3	647.0	2,233.4
2000 [4]	108.4	*	*	*	0.1	1.0	6.5	22.7	71.9	223.5	662.9	2,353.3
1999 [4]	111.3	*	*	*	0.1	1.0	6.5	23.9	74.1	234.2	690.9	2,414.4
Male												
2001	106.6	*	*	*	0.2	1.8	10.0	34.9	102.7	303.9	874.6	2,503.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20, I25.1-I25.9)												
Non-Hispanic white												
Male												
2000 [4]	109.2	*	*	*	0.2	1.6	10.3	36.1	107.5	319.8	900.4	2,666.6
1999 [4]	111.4	*	*	*	0.2	1.5	10.1	37.7	109.1	335.3	936.9	2,703.9
Female												
2001	105.3	*	*	*	*	0.5	2.9	9.8	37.2	138.2	496.0	2,121.0
2000 [4]	107.7	*	*	*	*	0.4	2.7	9.5	38.3	142.4	506.8	2,227.0
1999 [4]	111.1	*	*	*	*	0.5	2.8	10.4	41.1	149.7	529.9	2,299.0
Non-Hispanic black												
Both sexes												
2001	59.7	*	*	*	0.4	2.1	9.8	37.1	108.4	274.7	668.1	1,761.4
2000 [4]	59.8	*	*	*	*	1.8	10.4	37.0	113.4	284.4	666.8	1,745.8
1999 [4]	61.0	*	*	*	*	1.6	10.7	38.9	115.9	294.7	667.1	1,801.6
Male												
2001	58.4	*	*	*	*	3.3	14.1	53.2	148.2	364.1	825.5	1,768.3
2000 [4]	57.3	*	*	*	*	2.6	14.1	51.3	153.9	357.2	808.7	1,751.9
1999 [4]	58.8	*	*	*	*	2.4	15.3	55.5	160.0	374.0	790.8	1,838.6
Female												
2001	60.9	*	*	*	*	1.0	6.0	23.3	76.5	211.7	579.7	1,758.9
2000 [4]	62.1	*	*	*	*	1.1	7.1	24.7	80.7	233.1	587.0	1,743.5
1999 [4]	63.0	*	*	*	*	0.8	6.6	24.6	80.7	238.9	597.9	1,787.8

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other heart diseases (I26-I51)												
All origins [3]												
Both sexes												
2001	58.6	11.1	1.4	0.6	2.0	4.4	9.9	21.6	49.5	131.4	389.5	1,475.5
2000 [4]	58.8	12.0	1.1	0.6	2.2	4.4	9.9	21.5	50.9	134.2	391.3	1,509.4
1999 [4]	59.4	12.9	1.2	0.6	2.3	4.4	10.3	21.6	52.9	138.8	398.5	1,512.9
Male												
2001	51.8	10.8	1.5	0.6	2.4	5.2	12.4	27.6	61.7	158.7	452.0	1,473.4
2000 [4]	52.2	12.2	1.2	0.7	2.6	5.2	12.2	27.8	64.7	164.0	451.0	1,549.1
1999 [4]	53.1	13.1	1.3	0.7	2.7	5.3	12.9	27.7	67.5	172.7	459.7	1,548.9
Female												
2001	65.2	11.5	1.3	0.6	1.5	3.6	7.5	15.9	38.3	108.7	348.4	1,476.4
2000 [4]	65.1	11.8	0.9	0.5	1.8	3.5	7.5	15.3	38.2	109.6	352.3	1,493.2
1999 [4]	65.5	12.7	1.0	0.6	1.8	3.6	7.7	15.6	39.4	111.1	358.7	1,498.5
Hispanic												
Both sexes												
2001	13.3	9.2	1.0	0.5	1.3	2.6	4.9	10.9	27.9	75.4	217.5	798.5
2000 [4]	13.6	10.2	1.0	0.5	1.7	2.8	5.5	11.6	28.8	75.9	227.0	829.4
1999 [4]	13.8	12.2	1.2	0.5	1.6	2.8	6.3	11.6	30.6	78.7	229.6	827.9
Male												
2001	12.6	9.6	*	*	1.6	3.2	6.5	15.0	34.1	91.1	248.6	775.2
2000 [4]	13.2	9.9	1.3	0.7	2.1	3.5	6.7	15.1	37.2	94.1	250.8	890.8
1999 [4]	13.7	10.3	1.4	0.6	1.8	3.8	8.2	15.4	38.8	102.4	267.5	888.6
Female												
2001	13.9	8.7	*	*	0.9	1.9	3.2	6.9	22.5	63.0	196.4	809.8

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other heart diseases (I26-I51)												
Hispanic												
Female												
2000 [4]	14.0	10.6	*	*	1.1	1.9	4.3	8.1	21.4	61.6	211.0	799.3
1999 [4]	13.9	14.1	*	*	1.3	1.6	4.2	8.0	23.4	60.1	204.3	798.6
Non-Hispanic [5]												
Both sexes												
2001	65.2	11.6	1.5	0.6	2.1	4.7	10.6	22.6	51.1	134.7	396.5	1,497.8
2000 [4]	65.1	12.4	1.1	0.6	2.3	4.7	10.4	22.3	52.4	137.5	397.5	1,530.9
1999 [4]	65.6	13.1	1.1	0.6	2.4	4.7	10.8	22.4	54.3	142.0	404.4	1,533.9
Male												
2001	57.8	11.0	1.6	0.6	2.6	5.6	13.2	28.7	63.6	162.5	460.6	1,499.0
2000 [4]	58.0	12.7	1.2	0.7	2.7	5.5	12.9	28.9	66.4	167.8	458.9	1,573.6
1999 [4]	58.6	13.8	1.3	0.7	2.9	5.5	13.5	28.7	69.3	176.2	466.4	1,571.2
Female												
2001	72.3	12.2	1.3	0.6	1.6	3.9	8.0	16.7	39.4	111.6	354.4	1,497.3
2000 [4]	71.8	12.0	1.0	0.5	1.9	3.8	7.9	15.9	39.4	112.5	357.5	1,513.7
1999 [4]	72.2	12.3	1.0	0.6	1.9	3.9	8.1	16.2	40.5	113.9	364.2	1,519.0
Non-Hispanic white												
Both sexes												
2001	70.6	9.1	1.1	0.5	1.7	3.7	8.5	19.0	45.8	128.7	394.8	1,533.2
2000 [4]	70.3	10.5	0.9	0.5	1.7	3.7	8.4	18.7	47.4	131.2	395.8	1,564.2
1999 [4]	70.4	10.5	0.9	0.5	2.0	3.7	8.7	18.7	48.3	135.2	401.1	1,565.9
Male												
2001	61.4	9.4	1.2	0.6	2.0	4.4	10.9	24.8	57.5	155.8	457.4	1,530.7

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other heart diseases (I26-I51)												
Non-Hispanic white												
Male												
2000 [4]	61.6	10.9	1.1	0.6	2.0	4.4	10.7	24.6	60.7	161.0	457.6	1,606.4
1999 [4]	61.7	11.6	1.1	0.5	2.3	4.4	11.2	24.5	62.1	168.5	462.9	1,595.5
Female												
2001	79.4	8.8	1.0	0.5	1.3	3.1	6.1	13.4	34.8	105.7	353.3	1,534.2
2000 [4]	78.7	10.1	0.8	0.4	1.4	2.9	6.2	12.9	34.8	106.0	355.3	1,547.2
1999 [4]	78.8	9.4	0.8	0.5	1.7	3.0	6.2	13.1	35.3	107.3	360.6	1,554.1
Non-Hispanic black												
Both sexes												
2001	54.7	21.3	3.0	1.0	4.4	11.0	24.5	52.2	105.7	216.1	486.4	1,292.9
2000 [4]	54.6	19.6	2.0	1.1	4.9	10.8	23.7	52.8	104.5	223.6	484.3	1,319.4
1999 [4]	56.2	23.2	2.0	1.0	4.7	11.0	25.3	53.5	115.3	232.9	499.9	1,314.0
Male												
2001	53.5	18.8	3.3	0.9	5.5	13.4	29.3	63.9	134.9	261.9	593.5	1,362.9
2000 [4]	53.3	19.1	2.0	1.2	5.7	12.7	28.7	68.3	132.9	269.8	565.5	1,416.1
1999 [4]	56.2	22.9	2.3	1.2	6.0	13.4	30.9	68.7	149.5	291.0	584.7	1,466.6
Female												
2001	55.8	23.8	2.7	1.0	3.2	8.9	20.3	42.2	82.3	183.7	426.2	1,267.0
2000 [4]	55.8	20.1	1.9	1.0	4.2	9.0	19.2	39.6	81.6	191.1	438.5	1,283.4
1999 [4]	56.3	23.6	1.8	0.9	3.4	8.9	20.4	40.5	87.9	192.0	452.4	1,257.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and subacute endocarditis (I33)												
All origins [3]												
Both sexes												
2001	0.4	*	*	*	*	0.1	0.3	0.5	0.6	1.3	2.5	2.5
2000 [4]	0.4	*	*	*	*	0.1	0.3	0.5	0.7	1.2	2.4	2.8
1999 [4]	0.4	*	*	*	*	0.1	0.3	0.4	0.7	1.3	2.4	2.9
Male												
2001	0.5	*	*	*	*	*	0.3	0.7	0.8	1.6	3.5	3.8
2000 [4]	0.5	*	*	*	*	0.1	0.3	0.6	0.8	1.5	3.5	4.2
1999 [4]	0.5	*	*	*	*	0.1	0.3	0.6	0.9	1.4	3.2	4.7
Female												
2001	0.4	*	*	*	*	*	0.2	0.3	0.5	1.1	1.8	2.0
2000 [4]	0.4	*	*	*	*	*	0.2	0.3	0.5	1.0	1.8	2.3
1999 [4]	0.3	*	*	*	*	*	0.2	0.2	0.5	1.2	1.8	2.1
Hispanic												
Both sexes												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and subacute endocarditis (I33)												
Hispanic												
Female												
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.5	*	*	*	*	0.1	0.3	0.5	0.6	1.4	2.5	2.6
2000 [4]	0.4	*	*	*	*	0.1	0.3	0.5	0.7	1.2	2.5	2.9
1999 [4]	0.4	*	*	*	*	0.1	0.3	0.4	0.7	1.3	2.4	2.9
Male												
2001	0.5	*	*	*	*	*	0.3	0.7	0.8	1.6	3.6	3.9
2000 [4]	0.5	*	*	*	*	0.1	0.3	0.6	0.8	1.4	3.6	4.2
1999 [4]	0.5	*	*	*	*	*	0.3	0.6	0.9	1.4	3.2	4.7
Female												
2001	0.4	*	*	*	*	*	0.3	0.3	0.5	1.2	1.8	2.0
2000 [4]	0.4	*	*	*	*	*	0.2	0.3	0.6	1.0	1.8	2.3
1999 [4]	0.4	*	*	*	*	*	0.2	0.2	0.5	1.2	1.8	2.2
Non-Hispanic white												
Both sexes												
2001	0.4	*	*	*	*	*	0.2	0.4	0.5	1.2	2.5	2.7
2000 [4]	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.1	2.4	2.8
1999 [4]	0.4	*	*	*	*	*	0.2	0.3	0.6	1.2	2.4	3.0
Male												
2001	0.5	*	*	*	*	*	0.2	0.5	0.7	1.4	3.6	4.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and subacute endocarditis (I33)												
Non-Hispanic white												
Male												
2000 [4]	0.5	*	*	*	*	*	0.3	0.5	0.8	1.3	3.6	4.3
1999 [4]	0.5	*	*	*	*	*	0.2	0.4	0.7	1.3	3.3	5.1
Female												
2001	0.4	*	*	*	*	*	0.2	0.2	0.4	1.1	1.7	2.1
2000 [4]	0.4	*	*	*	*	*	0.1	0.2	0.4	1.0	1.7	2.2
1999 [4]	0.3	*	*	*	*	*	0.1	0.2	0.4	1.1	1.8	2.2
Non-Hispanic black												
Both sexes												
2001	0.7	*	*	*	*	*	0.7	1.3	1.7	2.6	4.0	*
2000 [4]	0.6	*	*	*	*	*	0.6	1.4	1.6	1.9	2.9	*
1999 [4]	0.6	*	*	*	*	*	0.7	1.2	2.0	2.6	*	*
Male												
2001	0.8	*	*	*	*	*	0.8	1.9	2.2	3.9	*	*
2000 [4]	0.7	*	*	*	*	*	0.8	1.8	*	*	*	*
1999 [4]	0.7	*	*	*	*	*	1.0	1.9	2.5	*	*	*
Female												
2001	0.5	*	*	*	*	*	0.7	*	*	*	*	*
2000 [4]	0.6	*	*	*	*	*	*	1.0	1.6	*	*	*
1999 [4]	0.5	*	*	*	*	*	*	*	1.6	2.9	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
All origins [3]												
Both sexes												
2001	0.3	0.8	0.2	0.1	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.9
2000 [4]	0.3	0.5	0.2	0.0	0.1	0.1	0.2	0.2	0.4	0.7	1.2	1.8
1999 [4]	0.3	0.7	0.2	0.1	0.1	0.1	0.3	0.3	0.5	0.8	1.3	2.0
Male												
2001	0.3	*	*	*	0.1	0.2	0.2	0.3	0.5	0.7	1.2	1.8
2000 [4]	0.3	*	*	*	0.1	0.1	0.3	0.2	0.4	0.8	1.5	2.1
1999 [4]	0.3	*	*	*	0.1	0.2	0.3	0.3	0.6	0.8	1.4	2.5
Female												
2001	0.3	*	*	*	*	*	0.2	0.3	0.3	0.6	1.2	2.0
2000 [4]	0.3	*	*	*	*	0.1	0.1	0.2	0.3	0.6	1.0	1.7
1999 [4]	0.3	*	*	*	*	*	0.2	0.3	0.4	0.8	1.3	1.9
Hispanic												
Both sexes												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Hispanic												
Female												
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.3	0.9	0.2	*	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.9
2000 [4]	0.3	*	*	*	0.1	0.1	0.2	0.2	0.3	0.7	1.1	1.8
1999 [4]	0.3	0.8	0.2	*	0.1	0.1	0.3	0.3	0.6	0.8	1.3	2.1
Male												
2001	0.3	*	*	*	0.1	0.2	0.2	0.3	0.5	0.7	1.2	1.8
2000 [4]	0.3	*	*	*	0.1	0.2	0.3	0.3	0.4	0.9	1.4	2.1
1999 [4]	0.3	*	*	*	*	0.2	0.3	0.3	0.6	0.9	1.4	2.6
Female												
2001	0.3	*	*	*	*	*	0.2	0.3	0.3	0.6	1.2	1.9
2000 [4]	0.3	*	*	*	*	*	0.1	0.2	0.3	0.6	1.0	1.7
1999 [4]	0.3	*	*	*	*	*	0.2	0.2	0.5	0.7	1.2	1.9
Non-Hispanic white												
Both sexes												
2001	0.3	*	*	*	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.9
2000 [4]	0.3	*	*	*	0.1	0.1	0.2	0.2	0.3	0.7	1.1	1.8
1999 [4]	0.3	*	*	*	*	0.1	0.2	0.3	0.6	0.8	1.3	2.0
Male												
2001	0.3	*	*	*	*	*	0.2	0.3	0.5	0.7	1.2	2.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Non-Hispanic white												
Male												
2000 [4]	0.3	*	*	*	*	*	0.2	0.2	0.4	0.8	1.4	2.1
1999 [4]	0.3	*	*	*	*	*	0.2	0.3	0.7	0.8	1.4	2.6
Female												
2001	0.3	*	*	*	*	*	0.2	0.2	0.3	0.6	1.3	1.8
2000 [4]	0.2	*	*	*	*	*	*	0.2	0.2	0.6	1.0	1.7
1999 [4]	0.3	*	*	*	*	*	0.2	0.2	0.5	0.7	1.2	1.8
Non-Hispanic black												
Both sexes												
2001	0.4	*	*	*	*	*	0.4	0.7	*	*	*	*
2000 [4]	0.4	*	*	*	*	*	0.4	0.5	*	*	*	*
1999 [4]	0.5	*	*	*	*	*	0.5	0.5	*	1.2	*	*
Male												
2001	0.5	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.4	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.5	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.4	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.4	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Heart failure (I50)												
All origins [3]												
Both sexes												
2001	20.0	0.6	*	*	0.1	0.2	0.7	2.4	9.4	34.3	136.3	674.6
2000 [4]	19.8	0.8	*	0.1	0.1	0.2	0.6	2.2	9.1	34.8	136.1	684.9
1999 [4]	19.7	*	*	*	0.1	0.2	0.6	2.1	8.9	35.4	137.2	681.9
Male												
2001	15.5	*	*	*	*	0.3	0.8	3.1	11.1	40.3	153.8	646.4
2000 [4]	15.3	1.1	*	*	0.1	0.2	0.7	2.7	10.9	41.1	153.7	672.2
1999 [4]	15.2	*	*	*	0.1	0.2	0.7	2.7	10.8	42.5	152.0	667.0
Female												
2001	24.3	*	*	*	*	0.2	0.6	1.8	7.8	29.3	124.7	686.4
2000 [4]	24.1	*	*	*	*	0.1	0.5	1.6	7.5	29.6	124.6	690.0
1999 [4]	24.0	*	*	*	*	0.2	0.5	1.6	7.1	29.6	127.6	687.9
Hispanic												
Both sexes												
2001	3.7	*	*	*	*	*	0.4	1.3	5.6	18.8	72.9	349.5
2000 [4]	3.6	*	*	*	*	*	*	0.7	5.8	18.9	72.1	369.6
1999 [4]	3.7	*	*	*	*	*	*	1.0	5.3	19.5	73.5	377.4
Male												
2001	3.0	*	*	*	*	*	*	1.8	6.0	21.8	75.6	333.3
2000 [4]	2.9	*	*	*	*	*	*	*	7.7	23.0	74.2	380.9
1999 [4]	3.0	*	*	*	*	*	*	*	6.1	24.6	82.9	358.0
Female												
2001	4.5	*	*	*	*	*	*	*	5.2	16.4	71.0	357.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Heart failure (I50)												
Hispanic												
Female												
2000 [4]	4.3	*	*	*	*	*	*	*	4.2	15.6	70.7	364.0
1999 [4]	4.4	*	*	*	*	*	*	*	4.6	15.5	67.3	386.8
Non-Hispanic [5]												
Both sexes												
2001	22.4	*	*	*	0.1	0.3	0.7	2.5	9.6	35.2	138.8	685.6
2000 [4]	22.1	0.8	*	*	0.1	0.2	0.6	2.3	9.3	35.7	138.5	694.8
1999 [4]	21.8	*	*	*	0.1	0.2	0.6	2.2	9.1	36.3	139.4	691.0
Male												
2001	17.4	*	*	*	*	0.3	0.9	3.2	11.4	41.4	157.2	658.0
2000 [4]	17.2	*	*	*	0.1	0.2	0.8	2.9	11.0	42.1	156.9	683.3
1999 [4]	16.9	*	*	*	0.1	0.2	0.8	2.8	11.0	43.5	154.5	677.1
Female												
2001	27.1	*	*	*	*	0.2	0.6	1.9	8.0	30.1	126.8	697.0
2000 [4]	26.7	*	*	*	*	0.2	0.5	1.7	7.7	30.5	126.5	699.5
1999 [4]	26.5	*	*	*	*	0.2	0.5	1.6	7.2	30.4	129.7	696.5
Non-Hispanic white												
Both sexes												
2001	25.1	*	*	*	*	0.2	0.5	2.1	8.7	33.9	139.7	706.8
2000 [4]	24.7	*	*	*	*	0.1	0.5	1.9	8.4	34.3	139.2	714.3
1999 [4]	24.3	*	*	*	*	0.1	0.5	1.7	8.1	34.9	139.1	709.1
Male												
2001	19.3	*	*	*	*	0.2	0.7	2.7	10.5	40.0	158.1	679.1

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Heart failure (I50)												
Non-Hispanic white												
Male												
2000 [4]	19.0	*	*	*	*	0.2	0.6	2.4	10.1	41.0	157.9	701.6
1999 [4]	18.6	*	*	*	*	*	0.6	2.2	10.1	42.2	154.7	690.8
Female												
2001	30.7	*	*	*	*	0.2	0.4	1.4	6.9	28.8	127.5	718.4
2000 [4]	30.2	*	*	*	*	*	0.3	1.3	6.8	28.7	126.9	719.4
1999 [4]	29.9	*	*	*	*	*	0.3	1.2	6.3	28.8	128.8	716.4
Non-Hispanic black												
Both sexes												
2001	14.3	*	*	*	*	0.6	2.0	6.6	20.0	55.5	159.1	551.2
2000 [4]	14.1	*	*	*	*	0.5	1.7	5.9	19.0	56.2	156.3	564.8
1999 [4]	14.3	*	*	*	*	0.6	1.7	6.5	19.2	57.3	164.8	557.4
Male												
2001	12.0	*	*	*	*	0.9	2.3	7.6	22.6	66.3	185.4	547.5
2000 [4]	11.6	*	*	*	*	*	1.8	7.3	21.8	62.9	176.5	589.4
1999 [4]	12.2	*	*	*	*	*	1.9	8.6	23.0	66.4	180.4	609.7
Female												
2001	16.4	*	*	*	*	*	1.7	5.6	18.0	47.9	144.4	552.6
2000 [4]	16.3	*	*	*	*	*	1.6	4.7	16.8	51.5	145.0	555.6
1999 [4]	16.2	*	*	*	*	*	1.5	4.8	16.1	50.9	156.0	537.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
All origins [3]												
Both sexes												
2001	37.9	9.7	1.1	0.5	1.8	4.0	8.8	18.5	39.1	95.1	249.5	796.5
2000 [4]	38.3	10.6	0.9	0.5	2.0	4.0	8.8	18.6	40.8	97.4	251.6	820.0
1999 [4]	39.0	11.9	0.9	0.5	2.1	4.1	9.2	18.7	42.8	101.3	257.6	826.1
Male												
2001	35.6	9.1	1.2	0.5	2.2	4.7	11.1	23.6	49.4	116.1	293.4	821.3
2000 [4]	36.1	10.5	1.0	0.6	2.3	4.8	10.9	24.3	52.6	120.7	292.4	870.7
1999 [4]	37.1	12.1	1.0	0.6	2.5	4.8	11.6	24.2	55.3	127.9	303.1	874.7
Female												
2001	40.2	10.4	1.0	0.5	1.3	3.2	6.5	13.5	29.6	77.7	220.7	786.1
2000 [4]	40.4	10.7	0.7	0.4	1.6	3.2	6.8	13.1	29.9	78.3	225.0	799.3
1999 [4]	40.9	11.6	0.8	0.5	1.7	3.3	6.9	13.5	31.3	79.6	228.0	806.5
Hispanic												
Both sexes												
2001	9.3	8.0	0.8	0.4	1.2	2.2	4.3	9.2	21.6	55.3	141.6	444.6
2000 [4]	9.6	8.7	0.7	0.5	1.6	2.5	4.9	10.2	22.0	54.9	152.1	457.2
1999 [4]	9.8	11.5	0.9	0.4	1.3	2.5	5.5	10.1	24.6	57.2	151.9	447.6
Male												
2001	9.3	7.9	*	*	1.5	2.8	5.7	12.6	27.3	67.7	170.4	438.1
2000 [4]	9.9	8.1	*	0.7	2.1	3.3	6.0	13.7	28.6	68.8	172.6	505.9
1999 [4]	10.3	9.8	*	*	1.5	3.4	7.2	13.7	31.9	75.6	179.4	526.2
Female												
2001	9.2	8.0	*	*	0.9	1.6	2.8	5.8	16.5	45.5	122.1	447.7

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
Hispanic												
Female												
2000 [4]	9.3	9.3	*	*	1.0	1.7	3.8	6.8	16.1	44.0	138.3	433.3
1999 [4]	9.2	13.3	*	*	1.1	1.5	3.7	6.6	18.1	42.9	133.6	409.7
Non-Hispanic [5]												
Both sexes												
2001	42.1	10.1	1.2	0.5	1.9	4.3	9.4	19.3	40.4	97.5	253.9	807.8
2000 [4]	42.3	11.0	0.9	0.5	2.0	4.3	9.3	19.3	42.0	99.8	255.4	831.4
1999 [4]	43.0	11.9	0.9	0.5	2.2	4.3	9.6	19.5	44.0	103.6	261.3	837.9
Male												
2001	39.6	9.3	1.4	0.5	2.4	5.1	11.8	24.5	50.9	118.8	298.5	835.3
2000 [4]	40.0	10.9	1.0	0.6	2.4	5.1	11.5	25.1	54.1	123.4	297.1	883.9
1999 [4]	40.9	12.7	1.0	0.6	2.6	5.1	12.1	25.0	56.7	130.5	307.4	886.8
Female												
2001	44.5	11.0	1.0	0.5	1.4	3.5	6.9	14.2	30.6	79.7	224.6	796.3
2000 [4]	44.5	11.0	0.8	0.4	1.7	3.4	7.1	13.6	30.9	80.4	228.3	810.2
1999 [4]	44.9	11.1	0.9	0.5	1.8	3.6	7.2	14.1	32.2	81.5	231.4	818.4
Non-Hispanic white												
Both sexes												
2001	44.7	8.1	0.9	0.4	1.5	3.4	7.6	16.3	36.2	92.9	251.4	821.8
2000 [4]	44.9	9.6	0.8	0.4	1.6	3.4	7.6	16.3	38.0	95.0	253.1	845.3
1999 [4]	45.4	9.8	0.8	0.5	1.9	3.4	7.8	16.4	39.0	98.3	258.4	851.7
Male												
2001	41.3	8.1	1.0	0.5	1.9	4.0	9.9	21.3	45.8	113.8	294.4	845.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
Non-Hispanic white												
Male												
2000 [4]	41.7	9.6	0.8	0.5	1.8	4.1	9.6	21.5	49.3	117.9	294.7	898.3
1999 [4]	42.4	10.8	0.9	0.5	2.1	4.1	10.1	21.5	50.6	124.1	303.5	897.0
Female												
2001	47.9	8.1	0.7	0.4	1.1	2.7	5.3	11.5	27.2	75.2	222.8	811.9
2000 [4]	47.9	9.6	0.7	0.3	1.3	2.7	5.6	11.1	27.4	75.7	225.7	824.0
1999 [4]	48.2	8.7	0.6	0.5	1.6	2.8	5.5	11.4	28.1	76.7	228.9	833.7
Non-Hispanic black												
Both sexes												
2001	39.3	18.2	2.4	0.9	3.8	10.0	21.4	43.7	83.3	157.1	322.1	737.9
2000 [4]	39.5	16.1	1.5	0.9	4.5	9.8	21.0	45.1	83.1	164.4	323.9	748.9
1999 [4]	40.9	20.9	1.5	0.9	4.4	10.1	22.3	45.2	93.3	171.8	331.4	752.7
Male												
2001	40.3	16.0	2.7	0.9	5.0	11.9	25.9	53.5	109.1	190.6	401.2	813.1
2000 [4]	40.5	15.0	1.7	1.0	5.1	11.7	25.4	58.6	108.9	202.9	384.1	821.9
1999 [4]	42.8	20.8	*	1.1	5.6	12.1	27.4	57.6	123.1	220.6	401.5	855.7
Female												
2001	38.5	20.5	2.1	0.9	2.7	8.3	17.3	35.3	62.5	133.5	277.6	710.1
2000 [4]	38.6	17.3	*	0.8	3.8	8.1	17.0	33.4	62.4	137.3	290.0	721.7
1999 [4]	39.2	21.1	*	*	3.2	8.2	17.9	34.7	69.5	137.3	292.2	714.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Essential (primary) hypertension and hypertensive renal disease (I10,I12)												
All origins [3]												
Both sexes												
2001	6.8	*	*	*	0.1	0.3	0.7	2.4	5.8	15.5	47.7	171.9
2000 [4]	6.4	*	*	*	*	0.2	0.8	2.3	5.9	15.1	45.5	162.9
1999 [4]	6.1	*	*	*	*	0.2	0.7	2.2	5.5	15.2	43.6	152.1
Male												
2001	5.2	*	*	*	*	0.3	0.9	3.0	6.5	16.6	45.8	150.8
2000 [4]	5.0	*	*	*	*	0.2	1.0	3.0	7.0	15.8	45.2	142.4
1999 [4]	4.7	*	*	*	*	0.2	0.9	2.7	6.4	16.5	42.9	132.4
Female												
2001	8.3	*	*	*	*	0.2	0.6	1.8	5.2	14.5	48.9	180.8
2000 [4]	7.8	*	*	*	*	0.2	0.6	1.7	4.9	14.6	45.7	171.3
1999 [4]	7.4	*	*	*	*	0.2	0.5	1.7	4.7	14.1	43.9	160.0
Hispanic												
Both sexes												
2001	2.4	*	*	*	*	*	*	1.8	4.5	16.5	48.9	160.1
2000 [4]	2.3	*	*	*	*	*	0.5	1.8	6.4	15.0	45.8	148.0
1999 [4]	2.2	*	*	*	*	*	*	2.2	5.0	15.2	48.3	129.8
Male												
2001	1.9	*	*	*	*	*	*	2.3	3.9	16.8	51.0	129.5
2000 [4]	2.1	*	*	*	*	*	0.8	2.8	8.1	15.2	52.6	141.1
1999 [4]	1.9	*	*	*	*	*	*	2.4	5.2	18.5	51.6	131.6
Female												
2001	2.9	*	*	*	*	*	*	1.2	5.0	16.2	47.4	175.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Essential (primary) hypertension and hypertensive renal disease (I10,I12)												
Hispanic												
Female												
2000 [4]	2.5	*	*	*	*	*	*	*	4.9	14.9	41.3	151.3
1999 [4]	2.5	*	*	*	*	*	*	1.9	4.8	12.7	46.1	128.9
Non-Hispanic [5]												
Both sexes												
2001	7.4	*	*	*	0.1	0.3	0.8	2.4	5.8	15.3	47.5	172.0
2000 [4]	7.0	*	*	*	*	0.2	0.8	2.4	5.8	15.1	45.3	162.9
1999 [4]	6.6	*	*	*	*	0.2	0.7	2.2	5.6	15.1	43.2	152.3
Male												
2001	5.7	*	*	*	*	0.3	1.0	3.0	6.6	16.5	45.3	151.1
2000 [4]	5.4	*	*	*	*	0.2	1.0	3.0	6.9	15.7	44.7	142.0
1999 [4]	5.1	*	*	*	*	0.2	0.9	2.7	6.5	16.4	42.4	131.9
Female												
2001	9.0	*	*	*	*	0.3	0.6	1.9	5.1	14.4	48.9	180.7
2000 [4]	8.5	*	*	*	*	0.2	0.6	1.8	4.9	14.6	45.8	171.3
1999 [4]	8.0	*	*	*	*	0.2	0.6	1.7	4.7	14.1	43.7	160.5
Non-Hispanic white												
Both sexes												
2001	7.0	*	*	*	*	0.1	0.4	1.4	3.9	11.7	41.8	162.8
2000 [4]	6.5	*	*	*	*	0.1	0.4	1.3	3.8	11.6	39.9	152.7
1999 [4]	6.1	*	*	*	*	0.1	0.3	1.2	3.6	11.8	37.3	141.7
Male												
2001	5.1	*	*	*	*	0.2	0.5	1.7	4.5	12.8	39.6	142.1

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Essential (primary) hypertension and hypertensive renal disease (I10,I12)												
Non-Hispanic white												
Male												
2000 [4]	4.9	*	*	*	*	*	0.6	1.8	4.7	12.4	39.5	131.8
1999 [4]	4.6	*	*	*	*	0.2	0.5	1.6	4.5	13.0	36.5	119.8
Female												
2001	8.8	*	*	*	*	*	0.3	1.0	3.3	10.8	43.2	171.5
2000 [4]	8.1	*	*	*	*	*	0.3	0.9	3.1	10.9	40.1	161.1
1999 [4]	7.6	*	*	*	*	*	0.2	0.9	2.8	10.8	37.8	150.5
Non-Hispanic black												
Both sexes												
2001	11.3	*	*	*	*	1.1	3.2	10.1	22.4	48.4	113.9	291.5
2000 [4]	11.0	*	*	*	*	0.7	3.2	9.9	23.3	49.5	108.9	285.9
1999 [4]	10.7	*	*	*	*	0.7	3.2	9.4	21.8	47.5	112.9	272.2
Male												
2001	10.0	*	*	*	*	1.2	3.9	13.1	25.8	55.5	119.7	277.2
2000 [4]	9.5	*	*	*	*	*	3.8	12.1	28.4	51.5	110.5	268.9
1999 [4]	9.4	*	*	*	*	*	3.8	12.0	24.8	52.3	119.1	268.6
Female												
2001	12.4	*	*	*	*	1.0	2.5	7.5	19.7	43.4	110.7	296.9
2000 [4]	12.4	*	*	*	*	*	2.6	8.0	19.3	48.1	108.0	292.2
1999 [4]	11.8	*	*	*	*	0.7	2.7	7.2	19.4	44.1	109.4	273.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebrovascular diseases (I60-I69)												
All origins [3]												
Both sexes												
2001	57.4	2.7	0.4	0.2	0.5	1.5	5.5	15.1	38.0	123.4	443.9	1,500.2
2000 [4]	59.6	3.3	0.3	0.2	0.5	1.5	5.8	16.0	41.0	128.6	461.3	1,589.2
1999 [4]	60.0	2.7	0.3	0.2	0.5	1.4	5.7	15.2	40.6	130.8	469.8	1,614.8
Male												
2001	45.2	3.1	0.3	0.2	0.5	1.6	5.7	16.7	43.4	140.4	467.3	1,380.2
2000 [4]	46.9	3.8	*	0.2	0.5	1.5	5.8	17.5	47.2	145.0	490.8	1,484.3
1999 [4]	47.1	3.3	0.3	0.2	0.5	1.5	5.8	16.7	46.5	147.3	500.4	1,512.8
Female												
2001	69.2	2.3	0.4	0.2	0.5	1.5	5.4	13.6	32.9	109.3	428.6	1,550.4
2000 [4]	71.8	2.7	0.4	0.2	0.5	1.5	5.7	14.5	35.3	115.1	442.1	1,632.0
1999 [4]	72.3	2.1	0.3	0.2	0.5	1.4	5.6	13.8	35.1	117.2	449.8	1,656.0
Hispanic												
Both sexes												
2001	17.4	*	*	*	0.6	1.4	5.0	15.9	35.4	106.2	341.2	1,042.9
2000 [4]	17.5	*	*	*	0.5	1.7	5.2	15.2	39.8	113.2	344.1	1,082.9
1999 [4]	17.4	*	*	*	0.5	1.6	4.9	15.7	37.7	116.9	348.3	1,077.6
Male												
2001	15.7	*	*	*	0.7	1.6	5.3	18.7	43.5	127.2	386.3	1,005.6
2000 [4]	15.8	*	*	*	*	1.9	5.5	18.1	48.8	136.1	392.9	1,029.9
1999 [4]	16.1	*	*	*	0.6	1.8	5.7	19.5	46.1	140.8	395.2	1,136.6
Female												
2001	19.1	*	*	*	*	1.2	4.7	13.1	28.2	89.6	310.7	1,061.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebrovascular diseases (I60-I69)												
Hispanic												
Female												
2000 [4]	19.4	*	*	*	*	1.4	4.9	12.4	31.9	95.2	311.3	1,108.9
1999 [4]	18.8	*	*	*	*	1.4	4.0	11.9	30.3	98.1	317.1	1,049.2
Non-Hispanic [5]												
Both sexes												
2001	63.2	2.8	0.3	0.2	0.5	1.5	5.6	14.9	38.0	124.3	447.7	1,514.0
2000 [4]	65.4	3.6	0.3	0.2	0.5	1.5	5.8	15.9	40.9	129.2	465.0	1,604.5
1999 [4]	65.7	2.9	0.3	0.2	0.5	1.4	5.8	15.1	40.6	131.2	473.8	1,630.4
Male												
2001	49.7	3.3	0.3	0.2	0.5	1.5	5.7	16.4	43.2	140.9	469.8	1,392.9
2000 [4]	51.5	4.2	*	0.2	0.5	1.5	5.8	17.3	46.8	145.1	493.7	1,499.9
1999 [4]	51.5	3.7	0.3	0.2	0.5	1.4	5.8	16.4	46.3	147.3	503.6	1,524.8
Female												
2001	76.1	2.4	0.4	0.2	0.5	1.5	5.5	13.5	33.2	110.4	433.2	1,564.4
2000 [4]	78.7	2.8	0.4	0.2	0.5	1.5	5.8	14.6	35.5	116.1	446.4	1,646.8
1999 [4]	79.2	2.0	*	0.2	0.5	1.4	5.8	13.9	35.4	118.1	454.4	1,672.7
Non-Hispanic white												
Both sexes												
2001	67.6	1.9	0.3	0.2	0.4	1.3	4.2	10.8	31.2	113.0	439.5	1,525.0
2000 [4]	70.0	2.6	0.3	0.2	0.5	1.2	4.3	12.0	34.2	119.3	455.7	1,616.8
1999 [4]	70.4	2.4	0.3	0.2	0.4	1.1	4.3	11.2	33.4	120.7	465.1	1,648.0
Male												
2001	52.0	2.2	*	0.2	0.4	1.3	4.3	11.9	35.1	128.0	460.5	1,399.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebrovascular diseases (I60-I69)												
Non-Hispanic white												
Male												
2000 [4]	53.9	3.3	*	0.2	0.5	1.2	4.3	13.0	38.7	133.1	482.3	1,505.9
1999 [4]	53.9	3.1	*	0.2	0.4	1.1	4.4	12.0	38.0	135.0	492.8	1,535.1
Female												
2001	82.6	*	*	*	0.3	1.3	4.1	9.8	27.4	100.3	425.6	1,577.4
2000 [4]	85.5	1.9	*	0.2	0.4	1.2	4.3	10.9	29.9	107.6	438.3	1,661.6
1999 [4]	86.1	*	*	*	0.4	1.1	4.3	10.5	29.1	108.8	447.0	1,693.0
Non-Hispanic black												
Both sexes												
2001	52.9	6.9	*	*	0.9	3.0	13.6	43.5	92.8	226.0	567.2	1,465.5
2000 [4]	54.3	8.0	*	*	0.7	3.0	14.9	44.3	95.2	226.3	593.8	1,545.8
1999 [4]	53.9	4.7	*	*	0.7	2.9	14.7	42.2	97.7	235.7	596.4	1,501.5
Male												
2001	46.2	8.3	*	*	0.7	3.1	13.5	49.9	113.9	273.5	620.5	1,359.4
2000 [4]	47.7	9.2	*	*	*	2.8	15.1	50.4	116.9	272.1	664.9	1,479.5
1999 [4]	47.5	*	*	*	*	2.9	15.0	49.6	119.5	279.1	656.7	1,455.9
Female												
2001	59.0	*	*	*	1.1	2.9	13.6	38.1	75.9	192.5	537.2	1,504.8
2000 [4]	60.3	*	*	*	0.8	3.2	14.8	39.2	77.8	194.1	553.8	1,570.4
1999 [4]	59.8	*	*	*	0.8	2.8	14.4	35.9	80.2	205.1	562.7	1,518.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerosis (I70)												
All origins [3]												
Both sexes												
2001	4.9	*	*	*	*	*	0.1	0.5	1.9	7.3	33.1	179.1
2000 [4]	5.1	*	*	*	*	*	0.1	0.5	1.9	7.7	34.6	188.8
1999 [4]	5.4	*	*	*	*	*	0.1	0.5	2.0	8.0	35.9	202.0
Male												
2001	3.7	*	*	*	*	*	0.1	0.6	2.4	9.0	35.9	167.2
2000 [4]	3.9	*	*	*	*	*	0.2	0.6	2.5	9.6	39.7	176.5
1999 [4]	4.0	*	*	*	*	*	0.1	0.7	2.7	9.6	40.0	189.0
Female												
2001	6.2	*	*	*	*	*	*	0.3	1.4	5.9	31.2	184.1
2000 [4]	6.3	*	*	*	*	*	*	0.3	1.2	6.2	31.3	193.8
1999 [4]	6.7	*	*	*	*	*	*	0.3	1.4	6.7	33.2	207.2
Hispanic												
Both sexes												
2001	1.1	*	*	*	*	*	*	*	*	4.0	23.1	130.7
2000 [4]	1.2	*	*	*	*	*	*	*	*	5.9	24.5	132.7
1999 [4]	1.2	*	*	*	*	*	*	*	1.5	5.2	27.1	129.1
Male												
2001	0.9	*	*	*	*	*	*	*	*	4.8	27.3	127.6
2000 [4]	1.0	*	*	*	*	*	*	*	*	8.2	32.0	104.8
1999 [4]	1.1	*	*	*	*	*	*	*	*	6.8	31.3	151.0
Female												
2001	1.4	*	*	*	*	*	*	*	*	3.5	20.2	132.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerosis (I70)												
Hispanic												
Female												
2000 [4]	1.4	*	*	*	*	*	*	*	*	4.0	19.5	146.4
1999 [4]	1.4	*	*	*	*	*	*	*	*	4.0	24.3	118.5
Non-Hispanic [5]												
Both sexes												
2001	5.5	*	*	*	*	*	0.1	0.5	1.9	7.5	33.4	180.5
2000 [4]	5.7	*	*	*	*	*	0.1	0.5	1.9	7.8	35.0	190.3
1999 [4]	5.9	*	*	*	*	*	0.1	0.5	2.1	8.1	36.1	203.9
Male												
2001	4.1	*	*	*	*	*	0.1	0.6	2.4	9.2	36.1	168.2
2000 [4]	4.3	*	*	*	*	*	0.2	0.7	2.6	9.7	39.9	178.7
1999 [4]	4.4	*	*	*	*	*	0.2	0.7	2.7	9.7	40.3	189.9
Female												
2001	6.8	*	*	*	*	*	*	0.3	1.5	6.0	31.6	185.5
2000 [4]	7.0	*	*	*	*	*	*	0.3	1.3	6.3	31.8	195.1
1999 [4]	7.4	*	*	*	*	*	*	0.3	1.5	6.9	33.4	209.6
Non-Hispanic white												
Both sexes												
2001	6.3	*	*	*	*	*	0.1	0.5	1.8	7.5	34.1	187.4
2000 [4]	6.5	*	*	*	*	*	0.1	0.4	1.8	7.8	35.5	196.8
1999 [4]	6.7	*	*	*	*	*	0.1	0.5	2.0	8.2	36.7	209.9
Male												
2001	4.7	*	*	*	*	*	0.1	0.6	2.3	9.2	36.6	175.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerosis (I70)												
Non-Hispanic white												
Male												
2000 [4]	4.9	*	*	*	*	*	0.2	0.6	2.6	9.8	40.6	183.6
1999 [4]	4.9	*	*	*	*	*	0.1	0.7	2.6	9.6	40.4	194.6
Female												
2001	7.9	*	*	*	*	*	*	0.3	1.3	6.1	32.5	192.6
2000 [4]	8.0	*	*	*	*	*	*	0.2	1.1	6.2	32.1	202.1
1999 [4]	8.5	*	*	*	*	*	*	0.3	1.4	6.9	34.3	215.9
Non-Hispanic black												
Both sexes												
2001	2.7	*	*	*	*	*	*	0.7	3.4	8.9	30.7	126.1
2000 [4]	3.0	*	*	*	*	*	*	1.1	3.3	9.1	35.6	140.2
1999 [4]	3.1	*	*	*	*	*	*	0.7	3.3	9.3	35.0	155.8
Male												
2001	2.2	*	*	*	*	*	*	1.0	3.9	11.0	37.2	113.2
2000 [4]	2.5	*	*	*	*	*	*	1.5	4.1	11.1	40.4	150.3
1999 [4]	2.6	*	*	*	*	*	*	*	4.5	12.0	44.4	166.4
Female												
2001	3.2	*	*	*	*	*	*	*	3.0	7.4	27.1	130.9
2000 [4]	3.5	*	*	*	*	*	*	*	2.7	7.7	32.8	136.5
1999 [4]	3.5	*	*	*	*	*	*	*	2.3	7.4	29.8	151.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of circulatory system (I71-I78)												
All origins [3]												
Both sexes												
2001	8.9	*	*	*	0.2	0.4	1.1	2.6	8.7	29.2	71.8	158.2
2000 [4]	9.3	*	*	*	0.2	0.4	1.0	2.8	9.2	32.2	74.6	163.2
1999 [4]	9.2	*	*	*	0.1	0.4	1.1	2.7	9.3	31.5	75.8	165.2
Male												
2001	9.6	*	*	*	0.2	0.6	1.5	3.6	12.0	40.3	97.0	197.3
2000 [4]	10.2	*	*	*	0.2	0.5	1.2	3.8	13.1	45.4	102.9	214.9
1999 [4]	10.1	*	*	*	0.2	0.5	1.3	3.7	13.3	44.4	104.2	207.5
Female												
2001	8.2	*	*	*	0.1	0.3	0.7	1.7	5.6	19.9	55.3	141.8
2000 [4]	8.3	*	*	*	0.1	0.3	0.8	1.8	5.6	21.4	56.2	142.2
1999 [4]	8.4	*	*	*	*	0.2	0.8	1.7	5.6	21.0	57.3	148.2
Hispanic												
Both sexes												
2001	2.0	*	*	*	*	*	0.7	2.1	4.8	13.7	39.0	95.0
2000 [4]	2.3	*	*	*	*	0.4	0.7	1.8	5.8	20.2	42.5	102.8
1999 [4]	2.1	*	*	*	*	*	0.7	1.3	5.7	16.4	44.0	110.8
Male												
2001	2.2	*	*	*	*	*	0.9	2.7	6.3	20.0	53.2	108.6
2000 [4]	2.5	*	*	*	*	*	0.8	2.1	8.0	25.7	62.0	131.0
1999 [4]	2.4	*	*	*	*	*	0.9	1.7	7.7	22.2	62.0	163.9
Female												
2001	1.8	*	*	*	*	*	*	1.6	3.4	8.8	29.3	88.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of circulatory system (I71-I78)												
Hispanic												
Female												
2000 [4]	2.1	*	*	*	*	*	*	1.5	3.9	15.8	29.4	89.0
1999 [4]	1.9	*	*	*	*	*	*	*	4.0	11.9	31.9	85.3
Non-Hispanic [5]												
Both sexes												
2001	9.9	*	*	*	0.2	0.5	1.2	2.6	8.9	30.1	73.2	160.1
2000 [4]	10.2	*	*	*	0.2	0.4	1.0	2.8	9.4	32.9	75.8	165.0
1999 [4]	10.2	*	*	*	0.1	0.4	1.1	2.8	9.5	32.3	76.9	166.8
Male												
2001	10.7	*	*	*	0.2	0.6	1.6	3.6	12.3	41.5	98.8	200.5
2000 [4]	11.3	*	*	*	0.2	0.5	1.3	3.9	13.5	46.5	104.4	217.9
1999 [4]	11.2	*	*	*	0.2	0.6	1.4	3.8	13.7	45.5	105.8	209.0
Female												
2001	9.1	*	*	*	*	0.3	0.8	1.7	5.8	20.6	56.3	143.3
2000 [4]	9.2	*	*	*	0.1	0.2	0.8	1.8	5.7	21.7	57.1	143.6
1999 [4]	9.3	*	*	*	*	0.2	0.8	1.8	5.7	21.5	58.2	149.9
Non-Hispanic white												
Both sexes												
2001	10.8	*	*	*	0.2	0.4	1.0	2.3	8.5	30.5	73.5	160.1
2000 [4]	11.2	*	*	*	0.1	0.3	0.8	2.4	9.0	33.5	76.3	164.8
1999 [4]	11.3	*	*	*	0.1	0.4	1.0	2.5	9.4	32.9	77.7	166.7
Male												
2001	11.8	*	*	*	0.2	0.6	1.3	3.1	12.0	42.4	100.3	204.7

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of circulatory system (I71-I78)												
Non-Hispanic white												
Male												
2000 [4]	12.6	*	*	*	0.2	0.5	1.0	3.5	13.2	47.9	106.3	222.7
1999 [4]	12.5	*	*	*	0.2	0.5	1.3	3.4	13.8	47.0	107.9	212.8
Female												
2001	9.9	*	*	*	*	0.3	0.6	1.5	5.3	20.4	55.7	141.6
2000 [4]	9.9	*	*	*	*	0.2	0.6	1.4	5.2	21.3	56.6	141.5
1999 [4]	10.1	*	*	*	*	0.2	0.7	1.5	5.3	21.1	57.8	148.3
Non-Hispanic black												
Both sexes												
2001	7.3	*	*	*	*	0.8	2.5	5.2	14.1	32.3	80.5	175.7
2000 [4]	7.3	*	*	*	*	0.7	2.2	5.7	14.5	33.4	78.9	180.0
1999 [4]	7.0	*	*	*	*	0.6	2.0	5.4	12.5	32.1	77.3	180.1
Male												
2001	7.2	*	*	*	*	1.0	3.5	7.4	17.9	40.3	97.2	168.6
2000 [4]	7.2	*	*	*	*	0.9	2.8	7.6	18.7	41.8	96.9	179.7
1999 [4]	6.7	*	*	*	*	0.9	2.4	7.4	16.0	40.0	90.7	184.2
Female												
2001	7.4	*	*	*	*	*	1.6	3.3	11.0	26.6	71.1	178.4
2000 [4]	7.5	*	*	*	*	*	1.6	4.0	11.2	27.5	68.8	180.1
1999 [4]	7.2	*	*	*	*	*	1.7	3.6	9.7	26.6	69.9	178.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Aortic aneurysm and dissection (I71)												
All origins [3]												
Both sexes												
2001	5.3	*	*	*	0.1	0.3	0.7	1.7	5.5	19.2	45.7	77.5
2000 [4]	5.6	*	*	*	0.1	0.3	0.7	1.8	6.2	21.8	47.2	79.2
1999 [4]	5.7	*	*	*	0.1	0.2	0.7	1.7	6.0	21.7	48.6	80.3
Male												
2001	6.5	*	*	*	0.2	0.4	1.2	2.6	8.5	28.7	66.4	114.8
2000 [4]	7.0	*	*	*	0.1	0.4	1.0	2.7	9.5	32.9	70.6	127.3
1999 [4]	7.0	*	*	*	0.1	0.4	1.0	2.7	9.5	32.7	72.7	125.3
Female												
2001	4.2	*	*	*	*	0.1	0.3	0.8	2.8	11.3	32.1	61.9
2000 [4]	4.3	*	*	*	*	0.1	0.4	0.9	3.0	12.7	32.0	59.6
1999 [4]	4.4	*	*	*	*	*	0.4	0.8	2.8	12.7	32.9	62.2
Hispanic												
Both sexes												
2001	1.1	*	*	*	*	*	0.4	1.2	2.5	8.2	22.9	31.9
2000 [4]	1.3	*	*	*	*	*	0.4	1.3	3.4	11.2	27.1	41.8
1999 [4]	1.1	*	*	*	*	*	0.5	0.9	3.4	9.2	24.0	40.7
Male												
2001	1.5	*	*	*	*	*	0.7	1.8	3.8	14.2	34.9	53.3
2000 [4]	1.6	*	*	*	*	*	*	1.8	5.2	16.0	42.3	64.5
1999 [4]	1.5	*	*	*	*	*	*	1.5	5.3	14.7	33.9	79.8
Female												
2001	0.7	*	*	*	*	*	*	*	*	3.5	14.8	21.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Aortic aneurysm and dissection (I71)												
Hispanic												
Female												
2000 [4]	1.0	*	*	*	*	*	*	*	*	7.5	16.8	30.7
1999 [4]	0.8	*	*	*	*	*	*	*	*	4.8	17.3	21.8
Non-Hispanic [5]												
Both sexes												
2001	6.0	*	*	*	0.1	0.3	0.8	1.7	5.7	19.8	46.7	79.0
2000 [4]	6.2	*	*	*	0.1	0.3	0.7	1.8	6.3	22.4	47.9	80.3
1999 [4]	6.3	*	*	*	0.1	0.3	0.8	1.8	6.2	22.4	49.5	81.7
Male												
2001	7.3	*	*	*	0.2	0.5	1.2	2.7	8.8	29.5	67.7	117.0
2000 [4]	7.8	*	*	*	*	0.4	1.0	2.7	9.8	33.8	71.6	129.6
1999 [4]	7.8	*	*	*	*	0.4	1.1	2.8	9.8	33.6	74.3	127.0
Female												
2001	4.7	*	*	*	*	0.2	0.3	0.8	2.9	11.8	32.9	63.2
2000 [4]	4.7	*	*	*	*	0.1	0.4	0.9	3.1	13.0	32.5	60.4
1999 [4]	4.8	*	*	*	*	*	0.4	0.9	2.9	13.2	33.4	63.5
Non-Hispanic white												
Both sexes												
2001	6.7	*	*	*	0.1	0.3	0.6	1.5	5.6	20.7	47.9	81.3
2000 [4]	7.0	*	*	*	*	0.2	0.6	1.6	6.3	23.4	49.3	82.0
1999 [4]	7.1	*	*	*	0.1	0.2	0.7	1.6	6.3	23.4	51.1	83.7
Male												
2001	8.2	*	*	*	0.2	0.5	1.0	2.3	8.7	30.9	70.0	122.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Aortic aneurysm and dissection (I71)												
Non-Hispanic white												
Male												
2000 [4]	8.8	*	*	*	*	0.3	0.8	2.5	9.8	35.6	74.2	135.3
1999 [4]	8.9	*	*	*	*	0.4	1.0	2.4	10.1	35.4	77.5	131.8
Female												
2001	5.2	*	*	*	*	*	0.3	0.6	2.7	12.0	33.3	64.0
2000 [4]	5.2	*	*	*	*	*	0.3	0.7	2.9	13.1	32.9	60.5
1999 [4]	5.3	*	*	*	*	*	0.4	0.7	2.7	13.3	33.9	64.5
Non-Hispanic black												
Both sexes												
2001	3.5	*	*	*	*	0.4	1.6	3.6	7.7	14.7	38.0	55.9
2000 [4]	3.6	*	*	*	*	0.5	1.5	3.3	8.2	17.3	36.8	63.0
1999 [4]	3.4	*	*	*	*	0.4	1.5	3.9	6.5	16.8	34.0	61.1
Male												
2001	3.9	*	*	*	*	*	2.6	5.4	11.3	19.9	47.4	49.7
2000 [4]	4.0	*	*	*	*	*	2.3	4.7	11.6	22.6	47.2	70.5
1999 [4]	3.7	*	*	*	*	*	2.0	6.0	9.3	22.1	38.8	77.3
Female												
2001	3.1	*	*	*	*	*	0.7	2.0	4.8	11.1	32.7	58.2
2000 [4]	3.3	*	*	*	*	*	0.9	2.1	5.5	13.5	30.9	60.2
1999 [4]	3.1	*	*	*	*	*	1.0	2.0	4.3	13.1	31.3	55.1

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)												
All origins [3]												
Both sexes												
2001	3.5	*	*	*	*	0.1	0.4	0.9	3.2	10.0	26.1	80.7
2000 [4]	3.6	*	*	*	0.1	0.1	0.3	1.0	3.1	10.4	27.4	84.0
1999 [4]	3.6	*	*	*	*	0.1	0.3	0.9	3.3	9.8	27.2	84.9
Male												
2001	3.1	*	*	*	*	0.1	0.3	0.9	3.5	11.6	30.6	82.5
2000 [4]	3.2	*	*	*	*	0.1	0.3	1.1	3.6	12.5	32.3	87.6
1999 [4]	3.1	*	*	*	*	0.1	0.3	1.0	3.8	11.7	31.5	82.2
Female												
2001	4.0	*	*	*	*	0.1	0.4	0.9	2.9	8.6	23.1	79.9
2000 [4]	4.0	*	*	*	*	0.1	0.4	0.9	2.6	8.7	24.2	82.6
1999 [4]	4.1	*	*	*	*	0.1	0.4	0.9	2.8	8.3	24.5	86.0
Hispanic												
Both sexes												
2001	0.9	*	*	*	*	*	*	0.9	2.2	5.5	16.0	63.2
2000 [4]	1.0	*	*	*	*	*	*	*	2.4	8.9	15.4	61.0
1999 [4]	1.0	*	*	*	*	*	*	*	2.4	7.2	20.0	70.2
Male												
2001	0.8	*	*	*	*	*	*	*	2.5	5.8	18.3	55.2
2000 [4]	0.9	*	*	*	*	*	*	*	2.7	9.7	19.7	66.5
1999 [4]	0.9	*	*	*	*	*	*	*	*	7.5	28.2	84.1
Female												
2001	1.1	*	*	*	*	*	*	*	*	5.4	14.5	67.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Hispanic												
Female												
2000 [4]	1.1	*	*	*	*	*	*	*	*	8.3	12.5	58.4
1999 [4]	1.1	*	*	*	*	*	*	*	2.4	7.1	14.6	63.4
Non-Hispanic [5]												
Both sexes												
2001	3.9	*	*	*	*	0.1	0.4	0.9	3.2	10.3	26.5	81.1
2000 [4]	4.0	*	*	*	0.1	0.1	0.3	1.0	3.1	10.5	27.9	84.7
1999 [4]	3.9	*	*	*	*	0.1	0.3	1.0	3.3	9.9	27.5	85.1
Male												
2001	3.4	*	*	*	*	0.1	0.3	0.9	3.5	12.0	31.1	83.5
2000 [4]	3.5	*	*	*	*	0.1	0.2	1.2	3.6	12.7	32.8	88.2
1999 [4]	3.4	*	*	*	*	0.1	0.3	1.0	3.9	11.9	31.5	82.0
Female												
2001	4.4	*	*	*	*	0.1	0.4	0.9	2.9	8.8	23.5	80.1
2000 [4]	4.4	*	*	*	*	*	0.4	0.9	2.6	8.7	24.6	83.3
1999 [4]	4.5	*	*	*	*	0.1	0.4	0.9	2.8	8.3	24.8	86.4
Non-Hispanic white												
Both sexes												
2001	4.2	*	*	*	*	0.1	0.3	0.8	2.9	9.8	25.6	78.8
2000 [4]	4.3	*	*	*	*	0.1	0.3	0.9	2.8	10.1	27.1	82.8
1999 [4]	4.2	*	*	*	*	0.1	0.3	0.9	3.1	9.6	26.5	83.0
Male												
2001	3.6	*	*	*	*	*	0.2	0.8	3.3	11.5	30.3	81.8

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Non-Hispanic white												
Male												
2000 [4]	3.8	*	*	*	*	*	0.2	1.0	3.3	12.3	32.1	87.4
1999 [4]	3.6	*	*	*	*	*	0.3	1.0	3.6	11.6	30.4	81.0
Female												
2001	4.7	*	*	*	*	*	0.4	0.9	2.6	8.3	22.4	77.6
2000 [4]	4.7	*	*	*	*	*	0.3	0.7	2.2	8.2	23.8	81.0
1999 [4]	4.8	*	*	*	*	*	0.4	0.8	2.5	7.9	24.0	83.9
Non-Hispanic black												
Both sexes												
2001	3.8	*	*	*	*	*	0.9	1.7	6.4	17.5	42.5	119.9
2000 [4]	3.7	*	*	*	*	*	0.6	2.4	6.3	16.1	42.1	117.0
1999 [4]	3.6	*	*	*	*	*	0.6	1.5	5.9	15.3	43.3	119.0
Male												
2001	3.3	*	*	*	*	*	0.9	2.0	6.6	20.5	49.8	119.0
2000 [4]	3.2	*	*	*	*	*	*	2.9	7.1	19.2	49.7	109.2
1999 [4]	3.0	*	*	*	*	*	*	1.4	6.7	17.9	51.9	107.0
Female												
2001	4.4	*	*	*	*	*	0.9	1.4	6.3	15.5	38.4	120.2
2000 [4]	4.2	*	*	*	*	*	0.7	1.9	5.7	14.0	37.9	119.9
1999 [4]	4.2	*	*	*	*	*	0.8	1.6	5.4	13.5	38.5	123.5

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of circulatory system (I80-I99)												
All origins [3]												
Both sexes												
2001	1.6	0.5	*	*	0.1	0.3	0.6	1.2	2.2	4.6	9.4	26.1
2000 [4]	1.6	0.7	*	*	0.1	0.3	0.7	1.4	2.1	4.3	9.4	25.6
1999 [4]	1.4	0.6	*	*	0.1	0.2	0.6	1.0	1.9	3.8	9.1	23.4
Male												
2001	1.4	*	*	*	*	0.3	0.7	1.4	2.5	4.9	9.8	22.0
2000 [4]	1.4	*	*	*	*	0.3	0.8	1.6	2.3	4.7	9.7	22.2
1999 [4]	1.3	*	*	*	*	0.2	0.7	1.1	2.0	4.2	9.6	20.8
Female												
2001	1.9	*	*	*	0.1	0.4	0.6	1.0	2.0	4.4	9.2	27.8
2000 [4]	1.8	*	*	*	0.1	0.4	0.6	1.1	2.0	4.0	9.2	26.9
1999 [4]	1.6	*	*	*	*	0.2	0.5	1.0	1.8	3.4	8.8	24.5
Hispanic												
Both sexes												
2001	0.6	*	*	*	*	*	*	0.9	*	4.2	7.4	32.5
2000 [4]	0.6	*	*	*	*	*	0.6	1.1	1.3	3.3	8.7	24.6
1999 [4]	0.5	*	*	*	*	*	*	1.1	1.5	4.1	5.4	24.6
Male												
2001	0.5	*	*	*	*	*	*	1.3	*	4.0	8.9	*
2000 [4]	0.6	*	*	*	*	*	0.8	*	*	*	10.8	*
1999 [4]	0.5	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.6	*	*	*	*	*	*	*	*	4.4	6.3	33.5

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of circulatory system (I80-I99)												
Hispanic												
Female												
2000 [4]	0.7	*	*	*	*	*	*	*	*	*	7.3	23.7
1999 [4]	0.6	*	*	*	*	*	*	*	*	4.3	*	23.9
Non-Hispanic [5]												
Both sexes												
2001	1.8	*	*	*	0.1	0.4	0.7	1.2	2.3	4.6	9.5	25.8
2000 [4]	1.8	0.7	*	*	0.1	0.4	0.7	1.4	2.2	4.4	9.4	25.6
1999 [4]	1.6	0.7	*	*	0.1	0.3	0.6	1.0	1.9	3.7	9.3	23.3
Male												
2001	1.5	*	*	*	*	0.3	0.7	1.4	2.6	4.9	9.8	21.6
2000 [4]	1.5	*	*	*	*	0.3	0.8	1.6	2.3	4.8	9.6	22.1
1999 [4]	1.4	*	*	*	*	0.2	0.7	1.1	2.0	4.2	9.7	20.5
Female												
2001	2.0	*	*	*	0.1	0.4	0.6	1.0	2.1	4.3	9.2	27.6
2000 [4]	2.0	*	*	*	*	0.4	0.6	1.2	2.1	4.1	9.3	27.0
1999 [4]	1.8	*	*	*	*	0.3	0.5	0.9	1.8	3.3	9.0	24.4
Non-Hispanic white												
Both sexes												
2001	1.8	*	*	*	*	0.3	0.6	1.0	2.1	4.3	9.2	25.7
2000 [4]	1.8	*	*	*	*	0.3	0.6	1.2	1.9	4.0	9.2	25.5
1999 [4]	1.6	*	*	*	*	0.2	0.5	0.9	1.8	3.5	9.0	23.0
Male												
2001	1.5	*	*	*	*	0.2	0.6	1.2	2.3	4.6	9.4	21.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of circulatory system (I80-I99)												
Non-Hispanic white												
Male												
2000 [4]	1.5	*	*	*	*	0.2	0.7	1.5	2.0	4.5	9.4	21.8
1999 [4]	1.4	*	*	*	*	0.2	0.6	0.9	2.0	3.9	9.3	20.3
Female												
2001	2.1	*	*	*	*	0.3	0.5	0.8	1.9	4.1	9.0	27.6
2000 [4]	2.0	*	*	*	*	0.3	0.5	1.0	1.9	3.6	9.0	27.0
1999 [4]	1.8	*	*	*	*	0.2	0.5	0.8	1.7	3.1	8.8	24.1
Non-Hispanic black												
Both sexes												
2001	2.0	*	*	*	*	0.8	1.6	2.3	4.9	7.9	14.3	29.7
2000 [4]	2.1	*	*	*	*	0.9	1.5	2.8	4.6	9.0	14.4	29.9
1999 [4]	1.8	*	*	*	*	0.5	1.5	2.4	3.0	6.8	14.0	28.5
Male												
2001	2.0	*	*	*	*	*	1.8	2.4	6.5	9.2	15.9	28.9
2000 [4]	2.0	*	*	*	*	0.9	1.8	3.5	5.3	9.1	13.3	30.5
1999 [4]	1.7	*	*	*	*	*	1.9	2.9	2.7	8.4	17.2	*
Female												
2001	2.1	*	*	*	*	0.9	1.4	2.1	3.7	7.0	13.5	29.9
2000 [4]	2.3	*	*	*	*	1.0	1.3	2.3	4.0	8.8	15.1	29.7
1999 [4]	1.8	*	*	*	*	*	1.1	2.0	3.2	5.6	12.3	30.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza and pneumonia (J10-J18)												
All origins [3]												
Both sexes												
2001	21.8	7.4	0.7	0.2	0.5	0.9	2.2	4.6	10.7	36.3	148.5	685.6
2000 [4]	23.2	7.6	0.7	0.2	0.5	0.9	2.4	4.7	11.9	39.1	160.3	744.1
1999 [4]	22.8	8.4	0.8	0.2	0.5	0.8	2.4	4.6	11.0	37.2	157.0	751.8
Male												
2001	19.6	8.1	0.7	0.2	0.5	1.0	2.5	5.7	13.1	45.0	186.4	808.4
2000 [4]	20.8	8.4	0.7	0.2	0.5	1.0	2.8	5.8	14.4	48.3	199.8	892.0
1999 [4]	20.3	8.8	0.8	0.3	0.5	1.0	2.6	5.9	12.8	46.9	196.5	894.7
Female												
2001	23.9	6.7	0.7	0.2	0.4	0.8	1.9	3.5	8.5	29.1	123.7	634.2
2000 [4]	25.6	6.8	0.7	0.2	0.5	0.8	1.9	3.7	9.5	31.5	134.6	683.9
1999 [4]	25.3	8.0	0.9	0.2	0.5	0.7	2.1	3.4	9.4	29.3	131.3	694.2
Hispanic												
Both sexes												
2001	7.4	7.0	1.1	*	*	0.3	1.3	3.6	9.1	35.4	137.2	660.3
2000 [4]	7.4	7.8	*	0.3	0.3	0.7	1.6	3.8	11.2	37.6	136.3	643.6
1999 [4]	6.6	8.5	*	*	0.4	0.4	1.5	3.5	9.5	31.7	118.5	614.6
Male												
2001	6.9	7.4	*	*	*	*	1.7	5.0	11.4	40.7	173.5	765.6
2000 [4]	7.2	7.6	*	*	*	0.8	2.1	5.3	14.8	46.0	180.4	763.9
1999 [4]	6.3	10.3	*	*	*	*	2.1	4.9	10.4	41.1	148.1	720.3
Female												
2001	7.9	6.5	*	*	*	*	1.0	2.1	7.1	31.2	112.7	608.8

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza and pneumonia (J10-J18)												
Hispanic												
Female												
2000 [4]	7.7	8.0	*	*	*	0.7	1.0	2.4	7.9	31.0	106.6	584.6
1999 [4]	7.0	6.5	*	*	*	*	0.8	2.2	8.6	24.3	98.9	563.6
Non-Hispanic [5]												
Both sexes												
2001	23.8	7.5	0.6	0.2	0.5	1.0	2.3	4.7	10.7	36.1	148.6	684.8
2000 [4]	25.4	7.5	0.7	0.2	0.5	0.9	2.4	4.7	11.9	39.0	160.7	745.7
1999 [4]	25.0	8.4	0.9	0.2	0.5	0.9	2.4	4.7	11.0	37.4	158.0	754.0
Male												
2001	21.5	8.2	0.6	0.2	0.5	1.1	2.6	5.7	13.1	44.9	186.5	808.2
2000 [4]	22.7	8.5	0.7	0.2	0.6	1.1	2.9	5.7	14.3	48.2	199.7	894.9
1999 [4]	22.2	8.4	0.9	0.3	0.5	1.1	2.7	5.9	12.9	47.0	197.7	897.9
Female												
2001	26.1	6.7	0.6	0.2	0.4	0.8	2.0	3.6	8.5	28.9	123.8	633.5
2000 [4]	27.9	6.5	0.6	0.2	0.5	0.8	2.0	3.8	9.6	31.4	135.3	685.4
1999 [4]	27.6	8.4	0.9	0.2	0.5	0.8	2.2	3.5	9.4	29.5	132.2	696.4
Non-Hispanic white												
Both sexes												
2001	26.2	4.9	0.5	0.2	0.4	0.8	1.9	4.0	9.9	35.0	147.2	693.9
2000 [4]	27.9	5.0	0.5	0.2	0.5	0.8	2.1	4.1	10.7	36.8	160.2	759.8
1999 [4]	27.5	5.3	0.6	0.2	0.4	0.8	2.0	4.0	10.0	36.1	158.8	766.9
Male												
2001	23.1	5.8	0.6	0.2	0.5	0.9	2.1	4.8	11.7	42.7	182.5	813.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza and pneumonia (J10-J18)												
Non-Hispanic white												
Male												
2000 [4]	24.5	5.8	0.5	0.2	0.5	0.9	2.4	4.7	12.6	44.7	197.5	908.1
1999 [4]	23.9	5.5	0.5	0.3	0.4	0.9	2.1	4.8	11.5	44.6	197.3	909.3
Female												
2001	29.2	3.9	0.5	0.2	0.4	0.8	1.7	3.3	8.1	28.5	123.7	644.0
2000 [4]	31.3	4.2	0.6	0.2	0.4	0.8	1.8	3.4	9.0	30.1	135.7	700.0
1999 [4]	31.0	5.1	0.8	0.2	0.4	0.6	1.9	3.1	8.7	29.0	133.5	710.2
Non-Hispanic black												
Both sexes												
2001	16.0	16.9	1.0	*	0.8	1.7	4.9	9.9	19.1	48.8	169.5	583.1
2000 [4]	16.9	17.5	1.2	*	0.8	1.8	4.6	10.3	22.3	60.8	172.2	604.5
1999 [4]	16.7	20.0	1.7	*	0.9	2.0	5.5	10.8	20.9	54.7	158.3	634.7
Male												
2001	16.3	16.6	*	*	0.8	2.1	5.9	13.5	26.6	70.0	242.5	718.4
2000 [4]	17.3	18.8	1.7	*	0.8	2.4	5.8	14.3	30.3	84.1	235.4	763.2
1999 [4]	16.9	20.1	2.1	*	0.8	2.3	6.7	15.3	27.3	77.9	215.1	803.4
Female												
2001	15.6	17.1	*	*	0.8	1.3	4.0	6.9	13.0	33.9	128.5	533.0
2000 [4]	16.5	16.2	*	*	0.8	1.2	3.5	6.9	15.9	44.4	136.7	545.6
1999 [4]	16.5	20.0	*	*	0.9	1.8	4.5	7.0	15.8	38.4	126.4	571.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza (J10-J11)												
All origins [3]												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	0.1	0.1	0.4	2.2
2000 [4]	0.6	*	*	*	*	*	0.0	0.1	0.4	1.1	4.0	20.2
1999 [4]	0.6	*	*	*	*	*	*	0.1	0.3	0.9	3.7	21.2
Male												
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	2.2
2000 [4]	0.5	*	*	*	*	*	*	0.1	0.4	1.3	4.0	20.7
1999 [4]	0.4	*	*	*	*	*	*	*	0.3	0.9	4.2	19.9
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	2.2
2000 [4]	0.8	*	*	*	*	*	*	0.1	0.4	0.9	4.0	19.9
1999 [4]	0.7	*	*	*	*	*	*	*	0.3	0.9	3.4	21.7
Hispanic												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza (J10-J11)												
Hispanic												
Female												
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	0.1	0.1	0.4	2.3
2000 [4]	0.7	*	*	*	*	*	0.0	0.1	0.4	1.1	4.1	20.5
1999 [4]	0.7	*	*	*	*	*	*	0.1	0.3	0.9	3.7	21.6
Male												
2001	0.1	*	*	*	*	*	*	*	*	*	*	2.3
2000 [4]	0.5	*	*	*	*	*	*	0.1	0.4	1.3	4.1	21.1
1999 [4]	0.5	*	*	*	*	*	*	*	0.3	0.8	4.3	20.7
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	2.2
2000 [4]	0.9	*	*	*	*	*	*	0.1	0.4	1.0	4.1	20.3
1999 [4]	0.8	*	*	*	*	*	*	*	0.3	0.9	3.4	22.0
Non-Hispanic white												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	2.4
2000 [4]	0.8	*	*	*	*	*	*	0.1	0.4	1.2	4.3	21.7
1999 [4]	0.8	*	*	*	*	*	*	0.1	0.3	0.9	3.9	23.3
Male												
2001	0.1	*	*	*	*	*	*	*	*	*	*	2.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza (J10-J11)												
Non-Hispanic white												
Male												
2000 [4]	0.6	*	*	*	*	*	*	*	0.4	1.4	4.3	22.0
1999 [4]	0.6	*	*	*	*	*	*	*	0.3	0.9	4.6	22.5
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	2.4
2000 [4]	1.0	*	*	*	*	*	*	0.1	0.5	1.0	4.3	21.6
1999 [4]	1.0	*	*	*	*	*	*	*	0.3	1.0	3.5	23.5
Non-Hispanic black												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	9.5
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia (J12-J18)												
All origins [3]												
Both sexes												
2001	21.7	7.2	0.7	0.2	0.4	0.8	2.2	4.6	10.6	36.2	148.1	683.4
2000 [4]	22.6	7.4	0.6	0.2	0.4	0.9	2.3	4.6	11.5	38.0	156.4	723.9
1999 [4]	22.2	8.1	0.8	0.2	0.4	0.8	2.3	4.6	10.7	36.4	153.3	730.6
Male												
2001	19.5	7.8	0.7	0.2	0.5	0.9	2.5	5.7	13.0	44.9	186.0	806.2
2000 [4]	20.3	8.1	0.6	0.2	0.5	1.0	2.7	5.6	14.0	47.1	195.8	871.3
1999 [4]	19.8	8.5	0.7	0.2	0.4	0.9	2.6	5.8	12.5	46.0	192.3	874.7
Female												
2001	23.8	6.7	0.7	0.2	0.4	0.7	1.9	3.5	8.4	29.0	123.2	632.0
2000 [4]	24.8	6.6	0.6	0.2	0.4	0.8	1.9	3.6	9.1	30.6	130.6	663.9
1999 [4]	24.6	7.7	0.8	0.2	0.4	0.7	2.0	3.4	9.1	28.4	128.0	672.5
Hispanic												
Both sexes												
2001	7.3	7.0	1.0	*	*	0.3	1.3	3.5	9.1	35.3	136.3	660.3
2000 [4]	7.3	7.5	*	*	0.3	0.7	1.6	3.8	11.0	37.2	134.3	633.0
1999 [4]	6.5	7.9	*	*	0.4	0.4	1.5	3.4	9.2	30.9	116.7	607.6
Male												
2001	6.8	7.4	*	*	*	*	1.7	5.0	11.4	40.7	172.6	765.6
2000 [4]	7.1	7.6	*	*	*	0.7	2.1	5.2	14.6	45.6	178.0	751.8
1999 [4]	6.1	10.3	*	*	*	*	2.0	4.7	9.9	39.5	147.0	718.2
Female												
2001	7.8	6.5	*	*	*	*	0.9	2.1	7.1	31.0	111.8	608.8

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia (J12-J18)												
Hispanic												
Female												
2000 [4]	7.6	7.4	*	*	*	*	1.0	2.3	7.9	30.7	105.0	574.7
1999 [4]	6.8	5.4	*	*	*	*	0.8	2.2	8.5	24.1	96.4	554.2
Non-Hispanic [5]												
Both sexes												
2001	23.8	7.2	0.6	0.2	0.5	0.9	2.3	4.6	10.6	36.0	148.2	682.5
2000 [4]	24.7	7.3	0.6	0.2	0.5	0.9	2.4	4.6	11.4	37.9	156.6	725.2
1999 [4]	24.3	8.1	0.8	0.2	0.4	0.9	2.4	4.6	10.7	36.5	154.2	732.4
Male												
2001	21.4	7.8	0.6	0.2	0.5	1.0	2.6	5.7	13.0	44.8	186.1	805.9
2000 [4]	22.2	8.2	0.7	0.2	0.5	1.0	2.8	5.6	13.9	46.8	195.7	873.8
1999 [4]	21.7	8.0	0.8	0.2	0.5	1.1	2.6	5.8	12.5	46.2	193.4	877.2
Female												
2001	26.0	6.6	0.6	0.2	0.4	0.8	2.0	3.6	8.5	28.8	123.3	631.3
2000 [4]	27.1	6.4	0.5	0.1	0.4	0.8	2.0	3.7	9.2	30.4	131.2	665.2
1999 [4]	26.8	8.2	0.9	0.2	0.4	0.8	2.2	3.5	9.1	28.5	128.8	674.4
Non-Hispanic white												
Both sexes												
2001	26.1	4.6	0.5	0.2	0.4	0.8	1.9	4.0	9.8	34.9	146.7	691.5
2000 [4]	27.1	4.8	0.5	0.2	0.4	0.8	2.1	3.9	10.3	35.6	155.9	738.1
1999 [4]	26.8	5.2	0.6	0.2	0.4	0.7	1.9	3.9	9.7	35.2	154.8	743.7
Male												
2001	23.0	5.3	0.6	0.2	0.5	0.9	2.1	4.8	11.6	42.6	182.0	811.5

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia (J12-J18)												
Non-Hispanic white												
Male												
2000 [4]	23.8	5.4	0.4	0.2	0.5	0.8	2.4	4.6	12.2	43.3	193.2	886.1
1999 [4]	23.4	5.4	*	0.2	0.4	0.9	2.0	4.7	11.1	43.8	192.7	886.8
Female												
2001	29.1	3.8	0.4	0.2	0.3	0.7	1.7	3.2	8.0	28.3	123.3	641.7
2000 [4]	30.3	4.2	0.5	*	0.3	0.8	1.7	3.3	8.5	29.1	131.4	678.4
1999 [4]	30.0	4.9	0.7	0.2	0.3	0.6	1.8	3.0	8.4	28.0	130.0	686.6
Non-Hispanic black												
Both sexes												
2001	15.9	16.9	1.0	*	0.8	1.7	4.9	9.9	19.0	48.7	169.1	581.5
2000 [4]	16.6	17.2	1.2	*	0.8	1.8	4.5	10.1	22.1	59.9	170.8	595.0
1999 [4]	16.5	19.3	1.7	*	0.9	2.0	5.4	10.7	20.7	54.0	156.6	628.9
Male												
2001	16.3	16.6	*	*	0.8	2.0	5.9	13.5	26.4	69.7	242.5	716.1
2000 [4]	17.1	18.8	1.7	*	0.7	2.3	5.7	14.0	30.1	83.4	233.9	750.3
1999 [4]	16.8	19.1	2.0	*	0.8	2.3	6.6	15.1	27.2	77.0	213.3	799.9
Female												
2001	15.6	17.1	*	*	0.8	1.3	4.0	6.9	13.0	33.9	127.9	531.7
2000 [4]	16.3	15.5	*	*	0.8	1.2	3.5	6.8	15.6	43.3	135.3	537.3
1999 [4]	16.3	19.6	*	*	0.9	1.7	4.4	6.9	15.5	37.7	124.8	565.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute lower respiratory infections (J20-J22)												
All origins [3]												
Both sexes												
2001	0.1	1.2	*	*	*	*	*	*	*	0.2	0.5	3.3
2000 [4]	0.2	1.1	*	*	*	*	*	*	*	0.2	0.7	4.5
1999 [4]	0.2	1.7	0.2	*	*	*	*	*	0.1	0.2	0.9	6.0
Male												
2001	0.1	1.3	*	*	*	*	*	*	*	*	0.4	2.3
2000 [4]	0.1	1.3	*	*	*	*	*	*	*	0.3	0.7	3.8
1999 [4]	0.1	1.9	*	*	*	*	*	*	*	*	1.1	4.0
Female												
2001	0.2	1.2	*	*	*	*	*	*	*	*	0.5	3.7
2000 [4]	0.2	*	*	*	*	*	*	*	*	0.2	0.6	4.8
1999 [4]	0.2	1.5	*	*	*	*	*	*	*	*	0.8	6.8
Hispanic												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute lower respiratory infections (J20-J22)												
Hispanic												
Female												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.1	1.3	*	*	*	*	*	*	*	0.2	0.4	3.3
2000 [4]	0.2	1.1	*	*	*	*	*	*	*	0.2	0.7	4.5
1999 [4]	0.2	1.7	0.2	*	*	*	*	*	0.1	0.2	0.9	6.1
Male												
2001	0.1	1.2	*	*	*	*	*	*	*	*	*	2.4
2000 [4]	0.1	*	*	*	*	*	*	*	*	0.3	0.7	3.7
1999 [4]	0.2	2.0	*	*	*	*	*	*	*	*	1.1	4.1
Female												
2001	0.2	1.4	*	*	*	*	*	*	*	*	0.5	3.7
2000 [4]	0.2	*	*	*	*	*	*	*	*	0.2	0.7	4.8
1999 [4]	0.3	1.5	*	*	*	*	*	*	*	*	0.8	6.9
Non-Hispanic white												
Both sexes												
2001	0.1	1.0	*	*	*	*	*	*	*	0.1	0.5	3.5
2000 [4]	0.2	*	*	*	*	*	*	*	*	0.3	0.7	4.7
1999 [4]	0.2	1.1	*	*	*	*	*	*	*	0.2	0.9	6.4
Male												
2001	0.1	*	*	*	*	*	*	*	*	*	*	2.5

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute lower respiratory infections (J20-J22)												
Non-Hispanic white												
Male												
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.7	3.7
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	1.1	4.1
Female												
2001	0.2	*	*	*	*	*	*	*	*	*	0.5	3.9
2000 [4]	0.2	*	*	*	*	*	*	*	*	0.3	0.7	5.1
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	0.8	7.3
Non-Hispanic black												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	4.2	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute bronchitis and bronchiolitis (J20-J21)												
All origins [3]												
Both sexes												
2001	0.1	1.2	*	*	*	*	*	*	*	0.1	0.3	2.1
2000 [4]	0.1	1.1	*	*	*	*	*	*	*	0.2	0.4	2.6
1999 [4]	0.1	1.6	0.2	*	*	*	*	*	*	0.2	0.6	3.6
Male												
2001	0.1	1.2	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	1.2	*	*	*	*	*	*	*	*	0.6	2.4
1999 [4]	0.1	1.9	*	*	*	*	*	*	*	*	0.7	2.6
Female												
2001	0.1	1.2	*	*	*	*	*	*	*	*	0.4	2.4
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.3	2.7
1999 [4]	0.2	1.4	*	*	*	*	*	*	*	*	0.5	4.0
Hispanic												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute bronchitis and bronchiolitis (J20-J21)												
Hispanic												
Female												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.1	1.3	*	*	*	*	*	*	*	0.1	0.3	2.1
2000 [4]	0.1	1.1	*	*	*	*	*	*	*	0.2	0.4	2.6
1999 [4]	0.1	1.7	0.2	*	*	*	*	*	*	0.2	0.6	3.6
Male												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.5	2.2
1999 [4]	0.1	2.0	*	*	*	*	*	*	*	*	0.7	2.6
Female												
2001	0.1	1.4	*	*	*	*	*	*	*	*	0.4	2.4
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.4	2.7
1999 [4]	0.2	1.4	*	*	*	*	*	*	*	*	0.5	4.0
Non-Hispanic white												
Both sexes												
2001	0.1	1.0	*	*	*	*	*	*	*	*	0.3	2.2
2000 [4]	0.1	*	*	*	*	*	*	*	*	0.2	0.4	2.7
1999 [4]	0.2	1.1	*	*	*	*	*	*	*	0.1	0.6	3.9
Male												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute bronchitis and bronchiolitis (J20-J21)												
Non-Hispanic white												
Male												
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.6	2.3
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	0.8	2.6
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	2.5
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.4	2.8
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	0.5	4.4
Non-Hispanic black												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	4.2	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Unspecified acute lower respiratory infection (J22)												
All origins [3]												
Both sexes												
2001	0.0	*	*	*	*	*	*	*	*	*	*	1.2
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	0.2	1.9
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	0.3	2.4
Male												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.0	*	*	*	*	*	*	*	*	*	*	1.3
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.3	2.1
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	0.3	2.8
Hispanic												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Unspecified acute lower respiratory infection (J22)												
Hispanic												
Female												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.0	*	*	*	*	*	*	*	*	*	*	1.2
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.2	1.9
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	0.3	2.4
Male												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.0	*	*	*	*	*	*	*	*	*	*	1.3
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.3	2.1
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	0.3	2.8
Non-Hispanic white												
Both sexes												
2001	0.0	*	*	*	*	*	*	*	*	*	*	1.3
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.3	2.0
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	0.3	2.5
Male												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Unspecified acute lower respiratory infection (J22)												
Non-Hispanic white												
Male												
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	*	1.4
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	0.3	3.0
Non-Hispanic black												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)												
All origins [3]												
Both sexes												
2001	43.2	1.0	0.3	0.3	0.4	0.7	2.2	8.5	44.1	167.9	379.8	644.7
2000 [4]	43.4	0.9	0.3	0.3	0.5	0.7	2.1	8.6	44.2	169.4	386.1	648.6
1999 [4]	44.5	0.9	0.4	0.3	0.5	0.8	2.0	8.5	47.5	177.2	397.8	646.0
Male												
2001	42.7	1.1	0.4	0.3	0.5	0.7	2.0	8.8	46.9	191.3	475.1	916.9
2000 [4]	43.5	1.2	0.4	0.4	0.6	0.8	1.9	9.0	47.8	195.2	488.5	967.9
1999 [4]	45.6	*	0.4	0.4	0.6	0.8	1.8	8.6	52.3	210.7	513.2	996.7
Female												
2001	43.7	*	*	0.2	0.4	0.7	2.3	8.1	41.5	148.5	317.3	530.8
2000 [4]	43.2	*	0.3	0.3	0.4	0.7	2.2	8.3	41.0	148.2	319.2	518.5
1999 [4]	43.4	*	0.3	0.2	0.5	0.9	2.1	8.4	43.1	149.8	322.9	504.6
Hispanic												
Both sexes												
2001	7.7	*	*	*	*	0.3	1.0	3.6	14.0	55.8	169.5	480.2
2000 [4]	7.6	*	*	*	*	0.3	1.1	3.3	14.3	53.1	181.9	481.1
1999 [4]	8.4	*	*	*	*	0.5	1.6	3.9	14.6	62.6	194.8	536.0
Male												
2001	7.8	*	*	*	*	*	0.7	3.2	16.1	75.5	224.0	676.1
2000 [4]	8.0	*	*	*	*	*	0.9	3.4	18.2	72.4	250.3	671.1
1999 [4]	8.9	*	*	*	*	*	1.4	3.6	17.6	81.9	272.7	836.8
Female												
2001	7.5	*	*	*	*	*	1.2	4.1	12.1	40.3	132.7	384.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)												
Hispanic												
Female												
2000 [4]	7.2	*	*	*	*	*	1.3	3.3	10.8	38.0	136.0	387.8
1999 [4]	8.0	*	*	*	*	*	1.8	4.2	12.0	47.5	142.9	391.0
Non-Hispanic [5]												
Both sexes												
2001	48.4	1.1	0.3	0.3	0.5	0.8	2.3	8.9	46.3	174.8	388.6	649.4
2000 [4]	48.3	1.0	0.3	0.4	0.5	0.8	2.2	9.0	46.3	176.1	393.8	653.2
1999 [4]	49.4	1.0	0.4	0.4	0.6	0.9	2.0	8.9	49.8	183.5	405.1	648.0
Male												
2001	48.0	*	0.4	0.3	0.6	0.8	2.2	9.3	49.0	198.1	485.7	924.8
2000 [4]	48.7	*	0.4	0.4	0.6	0.9	2.1	9.4	49.7	201.9	497.6	978.0
1999 [4]	50.8	*	0.4	0.5	0.7	0.8	1.9	9.0	54.5	217.4	521.9	1,000.2
Female												
2001	48.7	*	*	0.2	0.4	0.8	2.4	8.5	43.8	155.4	324.9	534.8
2000 [4]	48.0	*	*	0.3	0.4	0.7	2.3	8.7	43.2	154.8	326.2	521.9
1999 [4]	48.0	*	*	0.2	0.5	0.9	2.1	8.7	45.4	155.7	329.3	507.0
Non-Hispanic white												
Both sexes												
2001	55.9	*	*	0.2	0.3	0.5	1.9	8.7	48.6	185.9	408.6	676.9
2000 [4]	55.7	*	*	0.2	0.4	0.6	1.8	8.6	48.2	186.5	413.2	680.8
1999 [4]	56.6	*	0.2	0.2	0.4	0.6	1.6	8.5	51.8	193.6	422.4	675.6
Male												
2001	54.6	*	*	0.2	0.3	0.5	1.9	9.1	50.5	206.1	500.9	951.5

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)												
Non-Hispanic white												
Male												
2000 [4]	55.1	*	*	0.2	0.4	0.5	1.7	8.9	50.8	208.8	513.6	1,008.6
1999 [4]	57.3	*	*	0.3	0.5	0.5	1.5	8.5	55.6	224.9	534.9	1,033.5
Female												
2001	57.2	*	*	*	0.3	0.6	2.0	8.3	46.8	168.8	347.3	562.7
2000 [4]	56.2	*	*	0.2	0.3	0.6	1.9	8.3	45.8	167.6	347.2	548.7
1999 [4]	55.9	*	*	*	0.4	0.6	1.8	8.5	48.3	167.4	348.8	532.8
Non-Hispanic black												
Both sexes												
2001	21.1	*	0.9	0.8	1.3	2.5	4.9	12.7	41.3	120.4	236.3	379.3
2000 [4]	21.5	*	0.9	1.1	1.4	2.2	4.9	14.5	44.7	123.2	236.5	376.2
1999 [4]	22.6	*	*	1.0	1.4	2.7	4.7	14.1	46.7	129.9	265.3	377.0
Male												
2001	24.5	*	*	1.0	1.7	2.8	4.7	13.5	50.4	170.1	385.6	682.6
2000 [4]	25.1	*	*	1.3	1.8	2.8	4.9	15.2	55.3	178.9	372.4	696.3
1999 [4]	27.0	*	*	1.4	1.9	2.7	4.7	15.6	60.4	186.1	438.4	707.2
Female												
2001	18.0	*	*	0.7	1.0	2.2	5.0	11.9	34.0	85.3	152.5	266.9
2000 [4]	18.2	*	*	0.8	1.0	1.6	4.9	13.8	36.1	84.1	159.9	257.3
1999 [4]	18.6	*	*	0.6	0.9	2.7	4.7	12.8	35.8	90.3	168.4	253.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bronchitis, chronic and unspecified (J40-J42)												
All origins [3]												
Both sexes												
2001	0.3	0.5	*	*	*	*	0.0	0.1	0.2	0.7	2.1	9.1
2000 [4]	0.4	*	*	*	*	*	0.0	0.1	0.3	0.9	2.7	11.8
1999 [4]	0.4	0.5	*	*	*	*	*	0.1	0.4	0.9	2.8	10.7
Male												
2001	0.3	*	*	*	*	*	*	0.1	0.3	0.9	2.3	9.2
2000 [4]	0.4	*	*	*	*	*	*	*	0.3	1.0	3.6	12.9
1999 [4]	0.4	*	*	*	*	*	*	*	0.4	1.1	3.3	11.9
Female												
2001	0.4	*	*	*	*	*	*	0.1	0.2	0.6	1.9	9.1
2000 [4]	0.5	*	*	*	*	*	*	0.1	0.3	0.8	2.1	11.4
1999 [4]	0.5	*	*	*	*	*	*	*	0.4	0.8	2.5	10.3
Hispanic												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bronchitis, chronic and unspecified (J40-J42)												
Hispanic												
Female												
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.4	*	*	*	*	*	*	0.1	0.2	0.8	2.1	9.1
2000 [4]	0.5	*	*	*	*	*	*	0.1	0.3	0.9	2.7	11.8
1999 [4]	0.5	*	*	*	*	*	*	0.1	0.4	1.0	2.8	10.7
Male												
2001	0.3	*	*	*	*	*	*	0.1	0.3	1.0	2.3	9.3
2000 [4]	0.4	*	*	*	*	*	*	*	0.3	0.9	3.6	12.8
1999 [4]	0.4	*	*	*	*	*	*	*	0.5	1.1	3.3	11.9
Female												
2001	0.4	*	*	*	*	*	*	*	0.2	0.6	2.0	9.1
2000 [4]	0.5	*	*	*	*	*	*	0.1	0.3	0.8	2.1	11.4
1999 [4]	0.5	*	*	*	*	*	*	*	0.4	0.8	2.4	10.2
Non-Hispanic white												
Both sexes												
2001	0.4	*	*	*	*	*	*	0.1	0.3	0.8	2.2	9.7
2000 [4]	0.5	*	*	*	*	*	*	0.1	0.3	0.9	2.8	12.6
1999 [4]	0.5	*	*	*	*	*	*	0.1	0.5	1.0	2.8	11.2
Male												
2001	0.3	*	*	*	*	*	*	*	0.3	1.0	2.3	9.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bronchitis, chronic and unspecified (J40-J42)												
Non-Hispanic white												
Male												
2000 [4]	0.4	*	*	*	*	*	*	*	0.3	0.9	3.8	13.8
1999 [4]	0.4	*	*	*	*	*	*	*	0.5	1.2	3.2	12.4
Female												
2001	0.5	*	*	*	*	*	*	*	0.2	0.6	2.1	9.7
2000 [4]	0.6	*	*	*	*	*	*	*	0.3	0.8	2.1	12.2
1999 [4]	0.6	*	*	*	*	*	*	*	0.4	0.8	2.5	10.7
Non-Hispanic black												
Both sexes												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	2.2	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	2.9	*
Male												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Emphysema (J43)												
All origins [3]												
Both sexes												
2001	5.7	*	*	*	*	*	0.2	1.1	7.2	26.2	50.4	62.1
2000 [4]	5.9	*	*	*	*	*	0.2	1.3	7.4	27.8	52.7	63.3
1999 [4]	6.4	*	*	*	*	*	0.2	1.3	8.5	30.7	56.9	62.1
Male												
2001	6.0	*	*	*	*	*	0.2	1.4	7.9	30.8	65.3	101.2
2000 [4]	6.3	*	*	*	*	*	0.2	1.5	8.1	33.0	69.1	104.2
1999 [4]	6.9	*	*	*	*	*	0.2	1.5	9.5	37.8	76.4	105.4
Female												
2001	5.4	*	*	*	*	*	0.1	0.9	6.6	22.5	40.6	45.8
2000 [4]	5.6	*	*	*	*	*	0.2	1.0	6.8	23.6	41.9	46.7
1999 [4]	5.8	*	*	*	*	*	0.2	1.1	7.5	24.9	44.2	44.7
Hispanic												
Both sexes												
2001	0.8	*	*	*	*	*	*	*	1.4	6.6	20.4	39.4
2000 [4]	0.8	*	*	*	*	*	*	*	1.5	6.0	22.3	31.2
1999 [4]	0.9	*	*	*	*	*	*	*	1.3	7.1	23.5	44.2
Male												
2001	0.9	*	*	*	*	*	*	*	*	9.2	34.0	60.9
2000 [4]	0.9	*	*	*	*	*	*	*	*	9.1	32.5	58.4
1999 [4]	1.0	*	*	*	*	*	*	*	*	11.2	36.0	77.6
Female												
2001	0.6	*	*	*	*	*	*	*	*	4.6	11.2	28.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Emphysema (J43)												
Hispanic												
Female												
2000 [4]	0.6	*	*	*	*	*	*	*	*	3.6	15.5	*
1999 [4]	0.7	*	*	*	*	*	*	*	*	4.0	15.3	28.1
Non-Hispanic [5]												
Both sexes												
2001	6.4	*	*	*	*	*	0.2	1.2	7.6	27.5	51.7	62.8
2000 [4]	6.7	*	*	*	*	*	0.2	1.3	7.9	29.1	53.8	64.3
1999 [4]	7.1	*	*	*	*	*	0.2	1.3	9.0	32.0	58.1	62.5
Male												
2001	6.8	*	*	*	*	*	0.3	1.5	8.4	32.1	66.6	102.7
2000 [4]	7.1	*	*	*	*	*	0.2	1.6	8.5	34.3	70.6	105.9
1999 [4]	7.8	*	*	*	*	*	0.2	1.5	10.0	39.2	77.8	106.1
Female												
2001	6.1	*	*	*	*	*	0.1	1.0	7.0	23.6	41.9	46.3
2000 [4]	6.3	*	*	*	*	*	0.2	1.1	7.3	24.8	42.9	47.5
1999 [4]	6.5	*	*	*	*	*	0.2	1.1	8.1	26.1	45.3	45.0
Non-Hispanic white												
Both sexes												
2001	7.5	*	*	*	*	*	0.2	1.3	8.2	29.6	54.9	65.9
2000 [4]	7.8	*	*	*	*	*	0.2	1.3	8.4	31.2	57.0	66.9
1999 [4]	8.3	*	*	*	*	*	0.2	1.3	9.7	34.5	61.2	65.2
Male												
2001	7.8	*	*	*	*	*	0.3	1.5	8.7	33.8	69.6	106.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Emphysema (J43)												
Non-Hispanic white												
Male												
2000 [4]	8.1	*	*	*	*	*	0.2	1.4	8.9	35.8	73.9	107.3
1999 [4]	8.9	*	*	*	*	*	0.2	1.4	10.4	41.1	80.6	110.0
Female												
2001	7.2	*	*	*	*	*	0.1	1.0	7.7	26.1	45.2	48.9
2000 [4]	7.4	*	*	*	*	*	0.2	1.1	8.0	27.3	45.8	50.6
1999 [4]	7.7	*	*	*	*	*	0.2	1.2	9.0	29.0	48.5	47.3
Non-Hispanic black												
Both sexes												
2001	2.2	*	*	*	*	*	*	1.4	5.6	15.6	23.6	35.9
2000 [4]	2.5	*	*	*	*	*	*	2.0	6.1	17.1	27.0	37.5
1999 [4]	2.7	*	*	*	*	*	0.4	1.9	6.3	17.3	31.8	37.8
Male												
2001	2.8	*	*	*	*	*	*	1.6	7.9	23.0	40.2	69.3
2000 [4]	3.3	*	*	*	*	*	*	3.0	8.2	27.4	41.3	89.2
1999 [4]	3.6	*	*	*	*	*	*	3.0	9.1	29.3	55.7	73.7
Female												
2001	1.7	*	*	*	*	*	*	1.1	3.7	10.4	14.3	23.5
2000 [4]	1.8	*	*	*	*	*	*	1.3	4.5	9.8	18.9	18.3
1999 [4]	1.8	*	*	*	*	*	*	1.0	4.0	8.9	18.4	24.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Asthma (J45-J46)												
All origins [3]												
Both sexes												
2001	1.5	*	0.2	0.2	0.4	0.6	1.0	1.5	2.2	3.5	6.4	16.1
2000 [4]	1.6	*	0.2	0.3	0.4	0.6	1.0	1.6	2.2	4.0	6.9	16.7
1999 [4]	1.7	*	0.2	0.3	0.5	0.7	1.0	1.7	2.5	4.2	7.6	17.3
Male												
2001	1.1	*	0.3	0.3	0.4	0.6	0.9	1.1	1.5	2.7	4.5	12.1
2000 [4]	1.2	*	0.3	0.4	0.5	0.7	0.9	1.3	1.7	3.0	5.0	12.9
1999 [4]	1.2	*	*	0.4	0.5	0.6	0.8	1.1	1.8	3.2	5.7	12.7
Female												
2001	1.9	*	*	0.2	0.3	0.6	1.1	1.9	2.8	4.1	7.6	17.8
2000 [4]	2.0	*	*	0.3	0.3	0.5	1.2	2.0	2.7	4.8	8.1	18.2
1999 [4]	2.1	*	*	0.2	0.4	0.7	1.2	2.2	3.0	5.0	8.8	19.2
Hispanic												
Both sexes												
2001	0.7	*	*	*	*	*	0.6	1.4	2.4	3.8	7.9	13.8
2000 [4]	0.8	*	*	*	*	*	0.6	1.3	2.8	5.6	7.9	16.6
1999 [4]	0.9	*	*	*	*	0.3	0.9	1.6	2.4	5.8	10.4	17.5
Male												
2001	0.5	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.7	*	*	*	*	*	*	1.3	*	5.3	*	*
1999 [4]	0.7	*	*	*	*	*	0.9	1.4	*	*	*	*
Female												
2001	1.0	*	*	*	*	*	*	2.0	2.9	3.9	9.4	18.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Asthma (J45-J46)												
Hispanic												
Female												
2000 [4]	1.0	*	*	*	*	*	*	1.3	3.2	5.8	9.2	*
1999 [4]	1.2	*	*	*	*	*	1.0	1.9	3.1	7.1	13.2	*
Non-Hispanic [5]												
Both sexes												
2001	1.6	*	0.2	0.3	0.4	0.7	1.1	1.5	2.2	3.4	6.3	16.1
2000 [4]	1.7	*	0.2	0.3	0.5	0.7	1.1	1.6	2.2	3.9	6.8	16.6
1999 [4]	1.8	*	0.2	0.3	0.5	0.7	1.0	1.7	2.5	4.1	7.4	17.3
Male												
2001	1.1	*	*	0.3	0.4	0.6	0.9	1.2	1.5	2.6	4.4	12.2
2000 [4]	1.2	*	*	0.4	0.6	0.8	0.9	1.3	1.6	2.9	4.9	12.7
1999 [4]	1.3	*	*	0.4	0.6	0.7	0.8	1.1	1.8	3.1	5.6	12.6
Female												
2001	2.1	*	*	0.2	0.4	0.7	1.2	1.9	2.8	4.1	7.5	17.7
2000 [4]	2.1	*	*	0.3	0.4	0.6	1.2	2.0	2.6	4.7	8.0	18.2
1999 [4]	2.2	*	*	0.2	0.4	0.7	1.2	2.2	3.0	4.9	8.6	19.1
Non-Hispanic white												
Both sexes												
2001	1.4	*	*	0.1	0.2	0.4	0.7	1.1	1.5	2.9	5.8	15.7
2000 [4]	1.4	*	*	0.2	0.3	0.4	0.7	1.1	1.5	3.2	6.4	15.8
1999 [4]	1.5	*	*	0.2	0.4	0.4	0.7	1.2	1.9	3.4	7.1	16.5
Male												
2001	0.9	*	*	0.2	0.2	0.3	0.6	0.8	1.0	2.1	3.9	11.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Asthma (J45-J46)												
Non-Hispanic white												
Male												
2000 [4]	1.0	*	*	0.2	0.3	0.5	0.6	0.9	1.0	2.3	4.6	11.1
1999 [4]	1.0	*	*	0.2	0.4	0.4	0.5	0.7	1.4	2.4	5.2	10.6
Female												
2001	1.9	*	*	*	0.2	0.4	0.8	1.4	2.1	3.6	7.1	17.5
2000 [4]	1.9	*	*	0.2	0.3	0.4	0.9	1.3	2.0	4.0	7.6	17.7
1999 [4]	2.1	*	*	*	0.3	0.5	0.9	1.6	2.3	4.2	8.3	18.8
Non-Hispanic black												
Both sexes												
2001	3.1	*	*	0.8	1.2	2.2	3.3	4.7	6.9	7.7	10.7	16.5
2000 [4]	3.3	*	*	1.1	1.3	2.0	3.2	5.5	6.9	9.4	10.4	22.3
1999 [4]	3.3	*	*	1.0	1.3	2.4	3.0	5.4	7.0	9.8	10.6	20.7
Male												
2001	2.7	*	*	0.9	1.4	2.5	3.0	4.1	4.7	6.8	8.7	*
2000 [4]	2.8	*	*	1.3	1.6	2.5	3.0	3.7	6.1	7.9	6.5	24.7
1999 [4]	2.9	*	*	1.4	1.7	2.7	2.8	4.0	5.5	9.2	8.4	*
Female												
2001	3.4	*	*	0.7	0.9	2.0	3.6	5.1	8.6	8.4	11.8	17.1
2000 [4]	3.7	*	*	0.8	1.0	1.5	3.4	7.0	7.6	10.4	12.7	21.4
1999 [4]	3.6	*	*	0.6	0.9	2.2	3.2	6.5	8.1	10.2	11.7	21.3

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)												
All origins [3]												
Both sexes												
2001	35.7	*	*	*	0.1	0.1	0.9	5.7	34.5	137.5	321.0	557.4
2000 [4]	35.4	*	*	*	*	0.1	0.8	5.6	34.3	136.7	323.8	556.7
1999 [4]	36.0	*	*	*	*	0.1	0.7	5.5	36.2	141.4	330.6	555.8
Male												
2001	35.3	*	*	*	*	0.1	0.9	6.2	37.2	157.0	403.0	794.5
2000 [4]	35.6	*	*	*	*	*	0.8	6.1	37.7	158.3	410.9	838.0
1999 [4]	37.1	*	*	*	*	*	0.8	5.9	40.5	168.6	427.9	866.7
Female												
2001	36.0	*	*	*	*	0.1	1.0	5.2	32.0	121.3	267.1	458.2
2000 [4]	35.2	*	*	*	*	*	0.8	5.2	31.2	119.0	267.1	442.2
1999 [4]	35.0	*	*	*	*	0.2	0.7	5.0	32.2	119.1	267.4	430.5
Hispanic												
Both sexes												
2001	6.0	*	*	*	*	*	*	1.9	10.0	45.2	139.6	418.9
2000 [4]	5.9	*	*	*	*	*	0.4	1.7	9.9	40.7	149.1	422.7
1999 [4]	6.5	*	*	*	*	*	0.4	1.9	10.8	49.3	158.1	462.3
Male												
2001	6.3	*	*	*	*	*	*	2.1	12.2	62.5	181.5	605.7
2000 [4]	6.2	*	*	*	*	*	*	1.6	13.6	56.6	209.0	584.5
1999 [4]	7.0	*	*	*	*	*	*	1.6	13.9	65.9	228.9	735.4
Female												
2001	5.8	*	*	*	*	*	*	1.8	8.1	31.5	111.2	327.7

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)												
Hispanic												
Female												
2000 [4]	5.6	*	*	*	*	*	*	1.7	6.6	28.2	109.0	343.3
1999 [4]	5.9	*	*	*	*	*	*	2.1	8.1	36.3	111.0	330.7
Non-Hispanic [5]												
Both sexes												
2001	40.0	*	*	*	0.1	0.1	1.0	6.0	36.2	143.2	328.6	561.3
2000 [4]	39.5	*	*	*	*	0.1	0.9	6.0	36.0	142.3	330.5	560.5
1999 [4]	40.0	*	*	*	*	0.1	0.8	5.8	37.9	146.4	336.8	557.6
Male												
2001	39.8	*	*	*	*	0.1	1.0	6.6	38.9	162.6	412.4	800.6
2000 [4]	40.0	*	*	*	*	*	0.9	6.5	39.3	163.8	418.4	846.6
1999 [4]	41.4	*	*	*	*	*	0.8	6.3	42.2	173.9	435.1	869.6
Female												
2001	40.1	*	*	*	*	*	1.0	5.5	33.8	127.1	273.6	461.7
2000 [4]	39.1	*	*	*	*	*	0.9	5.5	32.9	124.5	273.2	444.8
1999 [4]	38.7	*	*	*	*	0.2	0.7	5.2	33.9	123.9	272.9	432.7
Non-Hispanic white												
Both sexes												
2001	46.6	*	*	*	*	0.1	1.0	6.2	38.6	152.6	345.7	585.7
2000 [4]	46.0	*	*	*	*	*	0.8	6.1	37.9	151.2	347.1	585.5
1999 [4]	46.3	*	*	*	*	0.1	0.7	5.9	39.9	154.8	351.4	582.8
Male												
2001	45.6	*	*	*	*	*	1.0	6.7	40.5	169.3	425.2	823.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)												
Non-Hispanic white												
Male												
2000 [4]	45.7	*	*	*	*	*	0.9	6.4	40.5	169.8	431.4	876.5
1999 [4]	47.1	*	*	*	*	*	0.8	6.2	43.4	180.3	445.9	900.5
Female												
2001	47.6	*	*	*	*	*	1.0	5.7	36.8	138.5	292.9	486.6
2000 [4]	46.3	*	*	*	*	*	0.8	5.8	35.5	135.5	291.6	468.2
1999 [4]	45.5	*	*	*	*	*	0.7	5.5	36.6	133.4	289.5	456.1
Non-Hispanic black												
Both sexes												
2001	15.6	*	*	*	*	*	1.3	6.5	28.7	96.1	200.8	323.4
2000 [4]	15.4	*	*	*	*	*	1.3	6.8	31.4	95.8	196.8	312.6
1999 [4]	16.4	*	*	*	*	*	1.2	6.6	33.2	101.9	220.1	314.0
Male												
2001	18.8	*	*	*	*	*	1.4	7.5	37.6	139.1	334.9	592.5
2000 [4]	18.7	*	*	*	*	*	1.3	8.5	40.7	142.7	322.4	578.9
1999 [4]	20.3	*	*	*	*	*	1.5	8.3	45.4	146.5	369.3	610.9
Female												
2001	12.7	*	*	*	*	*	1.3	5.7	21.5	65.8	125.5	223.7
2000 [4]	12.4	*	*	*	*	*	1.3	5.3	24.0	62.8	126.1	213.7
1999 [4]	12.9	*	*	*	*	*	1.0	5.0	23.3	70.4	136.6	203.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumoconioses and chemical effects (J60-J66,J68)												
All origins [3]												
Both sexes												
2001	0.4	*	*	*	*	*	*	*	0.3	1.5	4.1	6.2
2000 [4]	0.4	*	*	*	*	*	*	*	0.2	1.5	3.9	7.5
1999 [4]	0.4	*	*	*	*	*	*	*	0.2	1.5	4.1	6.1
Male												
2001	0.8	*	*	*	*	*	*	*	0.5	3.1	9.9	20.2
2000 [4]	0.8	*	*	*	*	*	*	*	0.5	3.2	9.6	24.9
1999 [4]	0.8	*	*	*	*	*	*	*	0.5	3.2	10.1	20.4
Female												
2001	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumoconioses and chemical effects (J60-J66,J68)												
Hispanic												
Female												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.4	*	*	*	*	*	*	*	0.3	1.5	4.2	6.2
2000 [4]	0.5	*	*	*	*	*	*	*	0.2	1.6	4.0	7.6
1999 [4]	0.4	*	*	*	*	*	*	*	0.3	1.5	4.3	6.2
Male												
2001	0.9	*	*	*	*	*	*	*	0.5	3.3	10.2	20.5
2000 [4]	0.9	*	*	*	*	*	*	*	0.5	3.3	9.7	25.7
1999 [4]	0.9	*	*	*	*	*	*	*	0.5	3.4	10.5	21.1
Female												
2001	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2001	0.5	*	*	*	*	*	*	*	0.3	1.6	4.5	6.6
2000 [4]	0.5	*	*	*	*	*	*	*	0.3	1.7	4.1	8.1
1999 [4]	0.5	*	*	*	*	*	*	*	0.2	1.6	4.6	6.4
Male												
2001	1.0	*	*	*	*	*	*	*	0.5	3.5	10.8	21.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumoconioses and chemical effects (J60-J66,J68)												
Non-Hispanic white												
Male												
2000 [4]	1.1	*	*	*	*	*	*	*	0.5	3.6	10.1	27.2
1999 [4]	1.0	*	*	*	*	*	*	*	0.5	3.6	11.1	21.8
Female												
2001	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2001	0.2	*	*	*	*	*	*	*	*	*	2.2	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	2.6	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	0.3	*	*	*	*	*	*	*	*	*	6.0	*
2000 [4]	0.3	*	*	*	*	*	*	*	*	*	6.5	*
1999 [4]	0.4	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)												
All origins [3]												
Both sexes												
2001	6.1	*	*	*	0.1	0.2	0.4	1.0	2.6	10.0	45.8	189.4
2000 [4]	5.9	*	*	*	0.1	0.2	0.4	1.0	2.5	10.3	44.5	187.6
1999 [4]	5.5	*	*	*	0.1	0.2	0.4	0.8	2.5	9.5	41.1	175.6
Male												
2001	6.1	*	*	*	0.1	0.2	0.5	1.2	3.2	13.3	63.1	270.3
2000 [4]	6.1	*	*	*	0.1	0.2	0.5	1.2	3.0	14.4	62.3	276.0
1999 [4]	5.7	*	*	*	0.1	0.2	0.4	0.9	3.2	13.1	59.7	258.4
Female												
2001	6.0	*	*	*	*	0.2	0.3	0.7	2.1	7.2	34.5	155.6
2000 [4]	5.8	*	*	*	*	0.1	0.3	0.7	1.9	7.0	33.0	151.6
1999 [4]	5.3	*	*	*	*	0.2	0.3	0.6	1.8	6.6	29.1	142.2
Hispanic												
Both sexes												
2001	1.5	*	*	*	*	*	*	*	1.8	6.1	27.4	161.3
2000 [4]	1.4	*	*	*	*	*	*	0.6	1.9	5.9	27.9	137.4
1999 [4]	1.2	*	*	*	*	*	*	*	1.6	5.5	24.6	130.5
Male												
2001	1.5	*	*	*	*	*	*	*	2.5	8.4	36.2	220.9
2000 [4]	1.4	*	*	*	*	*	*	*	2.5	7.4	42.8	169.3
1999 [4]	1.1	*	*	*	*	*	*	*	*	6.8	33.9	159.6
Female												
2001	1.5	*	*	*	*	*	*	*	*	4.3	21.5	132.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)												
Hispanic												
Female												
2000 [4]	1.4	*	*	*	*	*	*	*	*	4.8	17.8	121.7
1999 [4]	1.3	*	*	*	*	*	*	*	*	4.5	18.4	116.5
Non-Hispanic [5]												
Both sexes												
2001	6.7	*	*	*	0.1	0.2	0.4	1.0	2.7	10.2	46.5	190.0
2000 [4]	6.5	*	*	*	0.1	0.2	0.4	1.0	2.5	10.6	45.1	189.0
1999 [4]	6.0	*	*	*	0.1	0.2	0.4	0.8	2.5	9.7	41.7	176.8
Male												
2001	6.8	*	*	*	0.1	0.2	0.6	1.3	3.2	13.6	64.2	271.6
2000 [4]	6.8	*	*	*	0.1	0.2	0.5	1.2	3.1	14.8	62.8	279.7
1999 [4]	6.3	*	*	*	0.1	0.3	0.5	1.0	3.3	13.3	60.6	261.6
Female												
2001	6.6	*	*	*	*	0.2	0.3	0.8	2.2	7.4	34.9	156.0
2000 [4]	6.3	*	*	*	*	0.1	0.3	0.8	2.0	7.1	33.6	152.3
1999 [4]	5.8	*	*	*	*	0.2	0.3	0.7	1.8	6.8	29.4	142.9
Non-Hispanic white												
Both sexes												
2001	7.5	*	*	*	0.1	0.2	0.4	0.9	2.6	9.9	46.6	191.2
2000 [4]	7.3	*	*	*	0.1	0.2	0.4	0.9	2.3	10.0	45.0	192.5
1999 [4]	6.7	*	*	*	0.1	0.2	0.4	0.7	2.4	9.2	41.2	179.3
Male												
2001	7.6	*	*	*	*	0.3	0.6	1.2	3.1	13.2	64.4	276.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)												
Non-Hispanic white												
Male												
2000 [4]	7.5	*	*	*	*	0.2	0.5	1.1	2.9	13.7	62.7	288.1
1999 [4]	6.9	*	*	*	*	0.2	0.4	0.9	3.0	12.4	59.6	267.4
Female												
2001	7.4	*	*	*	*	0.2	0.3	0.7	2.1	7.0	34.7	155.5
2000 [4]	7.1	*	*	*	*	*	0.3	0.7	1.8	7.0	33.4	153.9
1999 [4]	6.5	*	*	*	*	0.2	0.3	0.6	1.7	6.5	29.3	144.2
Non-Hispanic black												
Both sexes												
2001	4.4	*	*	*	*	*	0.6	1.6	4.1	15.4	53.6	187.3
2000 [4]	4.3	*	*	*	*	*	0.7	1.9	4.3	17.9	52.0	160.6
1999 [4]	4.3	*	*	*	*	*	0.7	1.5	4.5	16.4	52.6	168.5
Male												
2001	4.3	*	*	*	*	*	0.8	1.9	5.5	20.8	75.6	243.7
2000 [4]	4.5	*	*	*	*	*	0.8	2.7	5.2	28.4	75.0	210.2
1999 [4]	4.7	*	*	*	*	*	0.9	1.8	6.6	25.3	81.6	229.4
Female												
2001	4.6	*	*	*	*	*	*	1.4	2.9	11.6	41.3	166.4
2000 [4]	4.1	*	*	*	*	*	*	1.2	3.6	10.4	39.1	142.2
1999 [4]	4.0	*	*	*	*	*	*	1.2	2.9	10.2	36.4	145.7

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of respiratory system (J00-J06,J30-J39, J67,J70-J98)												
All origins [3]												
Both sexes												
2001	9.2	7.3	0.6	0.2	0.4	0.5	1.3	3.5	9.6	30.1	73.2	142.7
2000 [4]	9.1	7.2	0.6	0.2	0.3	0.6	1.4	3.5	9.9	29.6	72.0	141.7
1999 [4]	9.0	6.6	0.6	0.2	0.3	0.6	1.3	3.3	9.5	29.4	70.7	147.6
Male												
2001	8.9	9.0	0.6	0.2	0.4	0.5	1.4	3.8	10.9	36.5	89.3	169.1
2000 [4]	9.0	7.9	0.6	0.2	0.4	0.6	1.5	3.8	11.4	35.7	90.5	179.7
1999 [4]	8.9	7.5	0.7	0.2	0.3	0.6	1.4	3.5	10.6	36.2	89.9	182.3
Female												
2001	9.4	5.5	0.6	0.2	0.3	0.6	1.2	3.1	8.4	24.8	62.6	131.6
2000 [4]	9.2	6.4	0.6	0.2	0.3	0.6	1.2	3.2	8.6	24.7	59.9	126.2
1999 [4]	9.1	5.8	0.5	0.2	0.2	0.7	1.1	3.1	8.5	23.7	58.3	133.6
Hispanic												
Both sexes												
2001	3.4	5.0	*	*	*	*	0.7	3.1	8.9	25.3	65.5	134.4
2000 [4]	3.5	5.6	*	*	*	0.5	1.1	2.6	9.8	25.4	64.2	135.4
1999 [4]	3.4	5.0	*	*	*	0.4	0.9	3.2	9.6	26.0	64.0	134.0
Male												
2001	3.4	5.0	*	*	*	*	0.9	3.4	10.6	30.2	78.3	167.6
2000 [4]	3.3	5.8	*	*	*	*	1.3	3.0	11.9	29.3	81.1	133.0
1999 [4]	3.3	5.2	*	*	*	*	1.1	3.1	11.6	30.5	82.4	161.8
Female												
2001	3.5	5.0	*	*	*	*	*	2.8	7.5	21.4	56.8	118.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of respiratory system (J00-J06,J30-J39, J67,J70-J98)												
Hispanic												
Female												
2000 [4]	3.6	5.3	*	*	*	*	0.9	2.2	7.8	22.2	52.8	136.5
1999 [4]	3.5	*	*	*	*	*	*	3.3	7.8	22.4	51.7	120.6
Non-Hispanic [5]												
Both sexes												
2001	10.0	7.8	0.6	0.2	0.4	0.6	1.3	3.5	9.7	30.3	73.3	142.7
2000 [4]	9.8	7.5	0.6	0.2	0.4	0.7	1.4	3.6	9.9	29.8	72.1	141.6
1999 [4]	9.7	6.9	0.6	0.2	0.3	0.7	1.3	3.3	9.4	29.5	70.8	147.7
Male												
2001	9.8	10.0	0.6	0.2	0.4	0.6	1.4	3.9	10.9	36.7	89.5	169.1
2000 [4]	9.8	8.4	0.6	0.2	0.4	0.6	1.5	3.9	11.3	36.0	90.7	181.3
1999 [4]	9.6	7.9	0.8	0.2	0.4	0.7	1.4	3.6	10.5	36.4	89.9	182.7
Female												
2001	10.2	5.5	0.6	0.2	0.3	0.6	1.3	3.1	8.5	25.0	62.7	131.8
2000 [4]	9.9	6.6	0.5	0.2	0.3	0.7	1.3	3.3	8.6	24.7	60.1	125.5
1999 [4]	9.8	5.9	0.5	0.2	0.2	0.7	1.2	3.1	8.5	23.8	58.4	133.7
Non-Hispanic white												
Both sexes												
2001	11.0	4.8	0.5	0.2	0.3	0.5	1.1	3.0	9.3	30.9	75.6	145.9
2000 [4]	10.8	5.4	0.5	0.2	0.3	0.6	1.2	3.2	9.6	30.2	73.8	145.7
1999 [4]	10.6	4.9	0.6	0.2	0.3	0.6	1.1	3.0	9.1	29.7	72.3	150.7
Male												
2001	10.8	6.2	0.4	0.2	0.4	0.4	1.3	3.4	10.5	37.7	92.4	173.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of respiratory system (J00-J06,J30-J39, J67,J70-J98)												
Non-Hispanic white												
Male												
2000 [4]	10.7	6.6	0.5	0.2	0.3	0.6	1.3	3.5	11.0	36.4	92.6	185.9
1999 [4]	10.5	5.6	0.6	0.2	0.4	0.7	1.2	3.2	10.0	37.0	92.0	186.1
Female												
2001	11.3	3.4	0.5	0.2	0.3	0.6	0.9	2.6	8.2	25.1	64.4	134.5
2000 [4]	11.0	4.1	0.5	*	0.3	0.6	1.1	3.0	8.4	25.0	61.5	129.5
1999 [4]	10.7	4.0	0.5	0.2	0.2	0.5	0.9	2.7	8.2	23.6	59.4	136.6
Non-Hispanic black												
Both sexes												
2001	6.9	19.0	1.1	*	0.7	1.3	2.9	7.5	13.9	29.6	56.9	110.8
2000 [4]	6.7	17.7	0.8	*	0.7	1.1	2.7	6.9	13.7	30.6	58.6	99.8
1999 [4]	6.7	15.8	1.1	*	0.4	1.4	2.9	6.6	13.6	31.3	58.7	112.8
Male												
2001	6.6	24.3	*	*	*	1.4	2.6	7.8	15.7	32.9	69.0	131.7
2000 [4]	6.8	17.1	*	*	0.8	1.2	3.2	8.0	15.6	36.2	74.0	135.0
1999 [4]	6.6	18.0	*	*	*	1.1	2.9	7.3	15.9	35.6	73.5	146.2
Female												
2001	7.2	13.4	*	*	*	1.2	3.2	7.2	12.4	27.2	50.0	103.1
2000 [4]	6.6	18.3	*	*	*	1.0	2.3	6.0	12.1	26.6	49.8	86.8
1999 [4]	6.9	13.6	*	*	*	1.7	2.9	5.9	11.9	28.2	50.4	100.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Peptic ulcer (K25-K28)												
All origins [3]												
Both sexes												
2001	1.6	*	*	*	*	0.1	0.3	0.7	1.7	3.9	11.4	33.4
2000 [4]	1.6	*	*	*	*	*	0.3	0.8	1.8	4.1	11.4	35.6
1999 [4]	1.6	*	*	*	*	0.1	0.3	0.7	1.8	4.5	12.3	34.6
Male												
2001	1.5	*	*	*	*	0.1	0.4	1.0	2.2	4.8	13.0	34.9
2000 [4]	1.6	*	*	*	*	*	0.4	1.2	2.4	5.3	13.7	40.7
1999 [4]	1.6	*	*	*	*	*	0.3	0.9	2.3	6.1	14.8	37.5
Female												
2001	1.7	*	*	*	*	*	0.1	0.5	1.1	3.2	10.4	32.7
2000 [4]	1.6	*	*	*	*	*	0.1	0.5	1.3	3.0	10.0	33.5
1999 [4]	1.7	*	*	*	*	*	0.2	0.5	1.3	3.3	10.6	33.5
Hispanic												
Both sexes												
2001	0.4	*	*	*	*	*	*	*	1.2	2.9	9.7	23.8
2000 [4]	0.5	*	*	*	*	*	*	*	1.6	3.3	9.9	30.5
1999 [4]	0.5	*	*	*	*	*	*	*	1.5	3.6	9.2	26.7
Male												
2001	0.5	*	*	*	*	*	*	*	*	*	16.5	*
2000 [4]	0.5	*	*	*	*	*	*	*	2.9	4.4	*	*
1999 [4]	0.6	*	*	*	*	*	*	*	*	4.8	14.1	*
Female												
2001	0.4	*	*	*	*	*	*	*	*	*	*	25.1

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Peptic ulcer (K25-K28)												
Hispanic												
Female												
2000 [4]	0.5	*	*	*	*	*	*	*	*	*	10.9	29.7
1999 [4]	0.4	*	*	*	*	*	*	*	*	*	*	22.9
Non-Hispanic [5]												
Both sexes												
2001	1.7	*	*	*	*	0.1	0.3	0.7	1.7	4.0	11.5	33.6
2000 [4]	1.8	*	*	*	*	*	0.3	0.8	1.8	4.1	11.5	35.7
1999 [4]	1.8	*	*	*	*	*	0.3	0.7	1.8	4.6	12.4	34.8
Male												
2001	1.6	*	*	*	*	0.1	0.4	1.0	2.2	4.8	12.8	35.5
2000 [4]	1.8	*	*	*	*	*	0.4	1.2	2.4	5.4	13.9	41.0
1999 [4]	1.7	*	*	*	*	*	0.4	1.0	2.3	6.1	14.8	37.7
Female												
2001	1.8	*	*	*	*	*	0.2	0.5	1.2	3.3	10.6	32.8
2000 [4]	1.8	*	*	*	*	*	0.1	0.5	1.3	3.1	9.9	33.5
1999 [4]	1.9	*	*	*	*	*	0.2	0.5	1.3	3.3	10.8	33.7
Non-Hispanic white												
Both sexes												
2001	1.9	*	*	*	*	*	0.2	0.7	1.5	3.9	11.6	34.7
2000 [4]	1.9	*	*	*	*	*	0.2	0.7	1.8	4.0	11.6	36.3
1999 [4]	2.0	*	*	*	*	*	0.2	0.6	1.7	4.5	12.4	35.6
Male												
2001	1.7	*	*	*	*	*	0.3	0.8	2.0	4.5	12.7	36.3

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Peptic ulcer (K25-K28)												
Non-Hispanic white												
Male												
2000 [4]	1.9	*	*	*	*	*	0.3	1.1	2.3	5.2	13.7	41.0
1999 [4]	1.8	*	*	*	*	*	0.3	0.8	2.1	5.9	14.9	38.6
Female												
2001	2.1	*	*	*	*	*	0.1	0.5	1.1	3.4	10.9	34.0
2000 [4]	2.0	*	*	*	*	*	0.1	0.4	1.3	3.1	10.2	34.4
1999 [4]	2.1	*	*	*	*	*	0.2	0.4	1.3	3.2	10.8	34.4
Non-Hispanic black												
Both sexes												
2001	1.1	*	*	*	*	*	0.6	1.4	2.8	4.5	9.0	20.9
2000 [4]	1.2	*	*	*	*	*	0.5	1.7	2.6	4.9	10.1	26.7
1999 [4]	1.3	*	*	*	*	*	0.8	1.5	2.8	5.7	11.0	25.5
Male												
2001	1.4	*	*	*	*	*	1.0	2.3	4.3	7.0	11.7	24.3
2000 [4]	1.4	*	*	*	*	*	0.8	2.4	3.8	7.2	13.9	31.7
1999 [4]	1.5	*	*	*	*	*	1.2	2.4	4.5	8.3	12.2	28.5
Female												
2001	0.9	*	*	*	*	*	*	*	1.5	2.7	7.4	19.7
2000 [4]	1.0	*	*	*	*	*	*	1.1	1.7	3.3	8.0	24.9
1999 [4]	1.1	*	*	*	*	*	*	*	*	3.9	10.3	24.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of appendix (K35-K38)												
All origins [3]												
Both sexes												
2001	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.7	2.1
2000 [4]	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.0	1.6
1999 [4]	0.1	*	*	0.1	*	*	0.1	0.1	0.2	0.4	0.8	1.5
Male												
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.7	1.0	2.8
2000 [4]	0.2	*	*	*	*	*	*	0.2	0.4	0.5	1.2	2.0
1999 [4]	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.0	2.9
Female												
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.5	1.8
2000 [4]	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.5
1999 [4]	0.1	*	*	*	*	*	*	*	*	0.3	0.7	0.9
Hispanic												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of appendix (K35-K38)												
Hispanic												
Female												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.7	2.2
2000 [4]	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.0	1.7
1999 [4]	0.1	*	*	*	*	*	0.0	0.1	0.2	0.4	0.8	1.5
Male												
2001	0.2	*	*	*	*	*	*	0.1	0.3	0.7	1.0	3.0
2000 [4]	0.2	*	*	*	*	*	*	0.2	0.4	0.5	1.2	2.0
1999 [4]	0.2	*	*	*	*	*	*	0.2	0.3	0.4	1.0	3.0
Female												
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.5	1.8
2000 [4]	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.5
1999 [4]	0.1	*	*	*	*	*	*	*	*	0.3	0.7	0.9
Non-Hispanic white												
Both sexes												
2001	0.2	*	*	*	*	*	*	0.1	0.1	0.5	0.7	2.1
2000 [4]	0.2	*	*	*	*	*	*	0.1	0.2	0.3	0.9	1.8
1999 [4]	0.1	*	*	*	*	*	*	0.1	0.1	0.4	0.8	1.6
Male												
2001	0.2	*	*	*	*	*	*	*	0.2	0.7	1.0	2.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of appendix (K35-K38)												
Non-Hispanic white												
Male												
2000 [4]	0.2	*	*	*	*	*	*	*	0.3	0.4	1.2	2.1
1999 [4]	0.2	*	*	*	*	*	*	0.2	0.2	0.4	0.9	3.1
Female												
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.4	2.0
2000 [4]	0.2	*	*	*	*	*	*	*	*	0.3	0.7	1.6
1999 [4]	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.0
Non-Hispanic black												
Both sexes												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hernia (K40-K46)												
All origins [3]												
Both sexes												
2001	0.5	1.1	*	*	*	*	0.1	0.2	0.5	1.2	3.5	12.6
2000 [4]	0.5	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.8	13.1
1999 [4]	0.5	1.2	*	*	*	*	0.1	0.2	0.4	1.2	3.7	12.7
Male												
2001	0.4	1.4	*	*	*	*	*	0.2	0.5	1.1	3.5	12.0
2000 [4]	0.4	1.4	*	*	*	*	*	0.2	0.4	1.1	4.0	12.6
1999 [4]	0.4	1.3	*	*	*	*	0.1	0.2	0.5	1.1	4.0	13.2
Female												
2001	0.6	*	*	*	*	*	*	0.2	0.5	1.2	3.6	12.8
2000 [4]	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.6	13.3
1999 [4]	0.6	*	*	*	*	*	*	0.1	0.4	1.2	3.5	12.6
Hispanic												
Both sexes												
2001	0.2	*	*	*	*	*	*	*	*	*	4.3	12.5
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	4.5	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.3	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hernia (K40-K46)												
Hispanic												
Female												
2000 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.6	1.0	*	*	*	*	0.1	0.2	0.5	1.2	3.5	12.5
2000 [4]	0.6	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.7	13.1
1999 [4]	0.6	1.3	*	*	*	*	0.1	0.1	0.5	1.1	3.7	12.7
Male												
2001	0.5	1.2	*	*	*	*	*	0.2	0.5	1.2	3.5	12.1
2000 [4]	0.5	1.4	*	*	*	*	*	0.2	0.4	1.1	3.9	12.8
1999 [4]	0.5	1.5	*	*	*	*	0.1	0.2	0.5	1.1	4.0	13.3
Female												
2001	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.5	12.7
2000 [4]	0.7	*	*	*	*	*	*	0.2	0.5	1.1	3.6	13.2
1999 [4]	0.7	*	*	*	*	*	*	0.1	0.4	1.2	3.5	12.4
Non-Hispanic white												
Both sexes												
2001	0.6	0.9	*	*	*	*	*	0.2	0.5	1.2	3.5	12.8
2000 [4]	0.6	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.7	13.3
1999 [4]	0.6	1.4	*	*	*	*	0.1	0.1	0.5	1.1	3.7	13.0
Male												
2001	0.5	*	*	*	*	*	*	0.2	0.5	1.2	3.5	11.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hernia (K40-K46)												
Non-Hispanic white												
Male												
2000 [4]	0.5	*	*	*	*	*	*	0.2	0.4	1.1	4.0	13.0
1999 [4]	0.5	*	*	*	*	*	*	0.2	0.6	1.1	4.1	13.0
Female												
2001	0.8	*	*	*	*	*	*	0.2	0.5	1.1	3.5	13.1
2000 [4]	0.8	*	*	*	*	*	*	0.2	0.5	1.1	3.6	13.4
1999 [4]	0.7	*	*	*	*	*	*	*	0.4	1.1	3.4	12.9
Non-Hispanic black												
Both sexes												
2001	0.4	*	*	*	*	*	*	*	0.9	1.5	3.6	11.2
2000 [4]	0.4	*	*	*	*	*	*	*	*	1.3	4.0	12.7
1999 [4]	0.4	*	*	*	*	*	*	*	*	1.5	4.2	10.0
Male												
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.4	*	*	*	*	*	*	*	*	*	*	9.0
2000 [4]	0.5	*	*	*	*	*	*	*	*	*	4.3	12.6
1999 [4]	0.5	*	*	*	*	*	*	*	*	*	4.2	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic liver disease and cirrhosis (K70,K73-K74)												
All origins [3]												
Both sexes												
2001	9.5	*	*	*	0.1	1.0	7.4	18.5	22.7	30.0	30.2	22.2
2000 [4]	9.4	*	*	*	0.1	1.0	7.5	17.7	23.8	29.8	31.0	23.1
1999 [4]	9.4	*	*	*	0.1	1.0	7.3	17.4	23.7	30.6	31.9	23.2
Male												
2001	12.4	*	*	*	*	1.3	9.8	27.5	33.1	40.2	36.8	32.0
2000 [4]	12.5	*	*	*	*	1.4	10.3	26.9	34.4	39.9	38.4	34.3
1999 [4]	12.5	*	*	*	*	1.3	10.3	26.9	34.6	41.6	39.6	33.1
Female												
2001	6.7	*	*	*	*	0.7	5.0	9.8	13.1	21.5	25.9	18.1
2000 [4]	6.5	*	*	*	*	0.7	4.7	8.8	14.0	21.5	26.1	18.5
1999 [4]	6.4	*	*	*	*	0.8	4.4	8.3	13.7	21.6	26.9	19.3
Hispanic												
Both sexes												
2001	8.9	*	*	*	*	1.2	9.7	28.0	39.0	54.4	60.4	36.9
2000 [4]	9.0	*	*	*	*	1.4	9.8	27.2	41.4	58.3	63.8	48.4
1999 [4]	8.8	*	*	*	*	1.3	10.5	26.6	39.0	56.3	59.4	51.2
Male												
2001	12.7	*	*	*	*	1.8	15.0	46.4	63.9	77.1	76.0	41.9
2000 [4]	12.7	*	*	*	*	2.3	15.0	45.6	66.1	83.4	80.1	60.5
1999 [4]	12.3	*	*	*	*	1.8	16.5	44.3	62.6	79.1	70.9	66.9
Female												
2001	5.0	*	*	*	*	*	4.0	9.9	17.0	36.5	49.9	34.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic liver disease and cirrhosis (K70,K73-K74)												
Hispanic												
Female												
2000 [4]	5.1	*	*	*	*	*	4.1	9.3	19.5	38.7	52.8	42.5
1999 [4]	5.0	*	*	*	*	*	4.1	9.3	18.1	38.4	51.7	43.7
Non-Hispanic [5]												
Both sexes												
2001	9.5	*	*	*	0.1	0.9	7.1	17.5	21.3	28.2	28.7	21.6
2000 [4]	9.4	*	*	*	0.1	1.0	7.1	16.7	22.3	27.9	29.5	22.1
1999 [4]	9.5	*	*	*	*	1.0	6.9	16.5	22.5	29.0	30.7	22.1
Male												
2001	12.3	*	*	*	*	1.2	9.1	25.6	30.6	37.6	34.9	31.4
2000 [4]	12.4	*	*	*	*	1.2	9.6	25.0	31.8	37.1	36.5	33.0
1999 [4]	12.5	*	*	*	*	1.2	9.5	25.2	32.4	39.3	38.2	31.6
Female												
2001	6.9	*	*	*	*	0.7	5.1	9.8	12.7	20.4	24.7	17.5
2000 [4]	6.7	*	*	*	*	0.7	4.7	8.7	13.6	20.3	24.9	17.7
1999 [4]	6.6	*	*	*	*	0.8	4.4	8.1	13.3	20.6	25.8	18.3
Non-Hispanic white												
Both sexes												
2001	10.1	*	*	*	*	0.9	7.0	17.0	21.0	28.9	29.9	22.5
2000 [4]	10.0	*	*	*	*	0.8	7.0	16.2	22.1	28.4	30.6	22.8
1999 [4]	9.9	*	*	*	*	0.9	6.6	15.8	21.8	29.5	31.7	22.6
Male												
2001	13.0	*	*	*	*	1.1	8.9	24.8	29.7	38.1	36.3	32.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic liver disease and cirrhosis (K70,K73-K74)												
Non-Hispanic white												
Male												
2000 [4]	13.1	*	*	*	*	1.1	9.4	24.4	31.5	37.7	37.8	34.1
1999 [4]	13.1	*	*	*	*	1.1	9.1	24.2	31.5	39.7	39.2	32.7
Female												
2001	7.4	*	*	*	*	0.7	5.1	9.3	12.8	21.1	25.7	18.2
2000 [4]	7.0	*	*	*	*	0.6	4.5	8.0	13.2	20.6	25.9	18.2
1999 [4]	6.9	*	*	*	*	0.7	4.1	7.5	12.7	21.0	26.7	18.6
Non-Hispanic black												
Both sexes												
2001	7.7	*	*	*	*	0.9	7.5	22.9	26.3	25.3	18.2	10.6
2000 [4]	7.7	*	*	*	*	1.1	8.0	22.6	26.1	26.5	18.1	14.0
1999 [4]	8.0	*	*	*	*	1.0	8.2	23.0	29.7	26.9	21.7	16.8
Male												
2001	10.7	*	*	*	*	1.0	10.4	35.1	42.0	39.7	22.5	*
2000 [4]	10.3	*	*	*	*	1.1	10.7	33.4	38.5	38.2	25.0	*
1999 [4]	11.0	*	*	*	*	1.1	11.3	35.8	43.8	41.0	30.0	*
Female												
2001	4.9	*	*	*	*	0.7	5.0	12.5	13.7	15.1	15.8	9.8
2000 [4]	5.4	*	*	*	*	1.1	5.7	13.4	16.2	18.3	14.2	11.8
1999 [4]	5.4	*	*	*	*	1.0	5.5	12.1	18.5	16.9	17.0	15.5

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcoholic liver disease (K70)												
All origins [3]												
Both sexes												
2001	4.3	*	*	*	*	0.7	4.8	10.7	11.5	10.1	5.7	2.4
2000 [4]	4.3	*	*	*	*	0.7	4.9	10.3	12.2	10.3	6.1	2.2
1999 [4]	4.3	*	*	*	*	0.6	4.8	10.4	12.0	10.5	6.5	2.8
Male												
2001	6.4	*	*	*	*	0.9	6.5	16.1	18.4	16.6	10.1	5.5
2000 [4]	6.5	*	*	*	*	1.0	6.8	16.0	19.0	17.3	11.5	5.3
1999 [4]	6.5	*	*	*	*	0.8	6.8	16.3	19.2	18.1	11.2	6.1
Female												
2001	2.3	*	*	*	*	0.4	3.1	5.5	5.1	4.7	2.8	1.0
2000 [4]	2.2	*	*	*	*	0.4	3.1	4.8	6.0	4.5	2.5	1.0
1999 [4]	2.1	*	*	*	*	0.5	2.9	4.7	5.4	4.3	3.4	1.5
Hispanic												
Both sexes												
2001	4.7	*	*	*	*	0.9	6.3	17.1	21.4	21.1	17.0	*
2000 [4]	4.8	*	*	*	*	1.0	6.9	16.8	23.4	22.9	15.4	*
1999 [4]	4.6	*	*	*	*	0.8	6.9	16.8	21.7	20.6	14.4	*
Male												
2001	7.7	*	*	*	*	1.5	10.0	29.3	38.9	41.9	30.4	*
2000 [4]	7.9	*	*	*	*	1.8	10.6	29.3	42.8	46.6	33.9	*
1999 [4]	7.4	*	*	*	*	1.2	10.8	29.1	40.0	39.5	27.1	*
Female												
2001	1.5	*	*	*	*	*	2.2	5.1	5.9	4.7	7.9	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcoholic liver disease (K70)												
Hispanic												
Female												
2000 [4]	1.4	*	*	*	*	*	2.9	4.5	6.2	4.3	*	*
1999 [4]	1.5	*	*	*	*	*	2.8	4.9	5.6	5.7	*	*
Non-Hispanic [5]												
Both sexes												
2001	4.2	*	*	*	*	0.6	4.6	10.1	10.7	9.3	5.2	2.2
2000 [4]	4.2	*	*	*	*	0.6	4.7	9.7	11.3	9.5	5.7	2.2
1999 [4]	4.2	*	*	*	*	0.6	4.6	9.8	11.3	9.9	6.1	2.5
Male												
2001	6.2	*	*	*	*	0.8	5.9	14.8	16.7	14.9	9.2	5.1
2000 [4]	6.2	*	*	*	*	0.8	6.3	14.7	17.1	15.4	10.5	5.0
1999 [4]	6.4	*	*	*	*	0.7	6.2	15.1	17.7	16.7	10.5	5.5
Female												
2001	2.4	*	*	*	*	0.5	3.2	5.5	5.0	4.7	2.5	1.0
2000 [4]	2.3	*	*	*	*	0.4	3.1	4.8	5.9	4.5	2.5	1.0
1999 [4]	2.2	*	*	*	*	0.5	2.9	4.7	5.3	4.2	3.3	1.4
Non-Hispanic white												
Both sexes												
2001	4.4	*	*	*	*	0.6	4.5	9.9	10.5	9.3	5.3	2.1
2000 [4]	4.3	*	*	*	*	0.6	4.5	9.4	11.1	9.5	5.6	2.2
1999 [4]	4.3	*	*	*	*	0.6	4.4	9.5	10.8	9.7	6.0	2.4
Male												
2001	6.4	*	*	*	*	0.8	5.8	14.7	16.4	14.6	9.3	4.8

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcoholic liver disease (K70)												
Non-Hispanic white												
Male												
2000 [4]	6.5	*	*	*	*	0.8	6.1	14.5	17.0	15.4	10.6	5.1
1999 [4]	6.5	*	*	*	*	0.7	6.0	14.7	17.1	16.3	10.3	5.1
Female												
2001	2.4	*	*	*	*	0.4	3.2	5.3	5.0	4.8	2.5	1.0
2000 [4]	2.2	*	*	*	*	0.4	3.0	4.4	5.5	4.6	2.4	1.0
1999 [4]	2.2	*	*	*	*	0.5	2.8	4.3	4.9	4.2	3.2	1.3
Non-Hispanic black												
Both sexes												
2001	3.7	*	*	*	*	0.5	4.4	11.7	13.1	10.8	4.6	*
2000 [4]	3.9	*	*	*	*	0.7	5.1	12.3	13.5	10.4	5.6	*
1999 [4]	4.2	*	*	*	*	0.5	5.1	12.9	15.4	11.9	7.2	*
Male												
2001	5.5	*	*	*	*	*	6.1	17.4	21.7	20.0	7.8	*
2000 [4]	5.5	*	*	*	*	*	7.1	17.9	19.6	18.4	10.2	*
1999 [4]	6.1	*	*	*	*	*	6.9	19.7	24.6	21.9	12.8	*
Female												
2001	2.2	*	*	*	*	*	2.8	6.8	6.2	4.3	*	*
2000 [4]	2.5	*	*	*	*	*	3.4	7.5	8.6	4.8	*	*
1999 [4]	2.4	*	*	*	*	*	3.5	7.0	8.1	4.8	4.0	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)												
All origins [3]												
Both sexes												
2001	5.2	*	*	*	*	0.3	2.6	7.8	11.2	19.9	24.5	19.9
2000 [4]	5.1	*	*	*	0.1	0.3	2.5	7.3	11.6	19.5	24.9	20.9
1999 [4]	5.1	*	*	*	*	0.4	2.5	7.0	11.7	20.1	25.5	20.4
Male												
2001	6.1	*	*	*	*	0.4	3.4	11.4	14.7	23.6	26.6	26.5
2000 [4]	6.0	*	*	*	*	0.4	3.5	10.9	15.4	22.7	26.9	29.0
1999 [4]	6.0	*	*	*	*	0.5	3.5	10.6	15.3	23.6	28.4	27.0
Female												
2001	4.4	*	*	*	*	0.2	1.9	4.3	8.0	16.8	23.1	17.1
2000 [4]	4.3	*	*	*	*	0.3	1.6	4.0	8.0	16.9	23.6	17.5
1999 [4]	4.3	*	*	*	*	0.3	1.4	3.6	8.3	17.3	23.5	17.7
Hispanic												
Both sexes												
2001	4.3	*	*	*	*	*	3.5	10.9	17.6	33.3	43.5	30.6
2000 [4]	4.3	*	*	*	*	0.4	2.9	10.5	18.0	35.4	48.4	44.5
1999 [4]	4.2	*	*	*	*	0.5	3.6	9.7	17.2	35.7	45.0	40.0
Male												
2001	5.0	*	*	*	*	*	5.1	17.1	25.0	35.1	45.6	*
2000 [4]	4.8	*	*	*	*	0.6	4.4	16.2	23.3	36.7	46.2	50.4
1999 [4]	4.9	*	*	*	*	0.6	5.7	15.1	22.6	39.5	43.8	45.3
Female												
2001	3.4	*	*	*	*	*	1.8	4.7	11.0	31.8	42.0	33.5

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)												
Hispanic												
Female												
2000 [4]	3.7	*	*	*	*	*	1.3	4.8	13.3	34.3	49.9	41.5
1999 [4]	3.5	*	*	*	*	*	1.3	4.5	12.4	32.7	45.8	37.4
Non-Hispanic [5]												
Both sexes												
2001	5.3	*	*	*	*	0.3	2.5	7.5	10.7	18.9	23.6	19.4
2000 [4]	5.2	*	*	*	0.1	0.3	2.5	7.0	11.0	18.4	23.8	19.9
1999 [4]	5.2	*	*	*	*	0.3	2.3	6.7	11.2	19.2	24.6	19.6
Male												
2001	6.2	*	*	*	*	0.4	3.1	10.8	13.8	22.8	25.7	26.4
2000 [4]	6.1	*	*	*	*	0.3	3.3	10.3	14.7	21.7	26.0	28.0
1999 [4]	6.1	*	*	*	*	0.4	3.3	10.1	14.7	22.6	27.7	26.2
Female												
2001	4.5	*	*	*	*	0.2	1.9	4.3	7.7	15.7	22.2	16.5
2000 [4]	4.4	*	*	*	*	0.3	1.6	3.9	7.6	15.8	22.4	16.7
1999 [4]	4.4	*	*	*	*	0.3	1.4	3.5	8.0	16.3	22.6	17.0
Non-Hispanic white												
Both sexes												
2001	5.8	*	*	*	*	0.3	2.5	7.1	10.5	19.6	24.7	20.4
2000 [4]	5.7	*	*	*	*	0.3	2.4	6.7	11.0	18.9	25.0	20.6
1999 [4]	5.6	*	*	*	*	0.3	2.3	6.4	11.0	19.8	25.6	20.2
Male												
2001	6.6	*	*	*	*	0.4	3.0	10.2	13.3	23.5	26.9	28.1

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)												
Non-Hispanic white												
Male												
2000 [4]	6.6	*	*	*	*	0.3	3.4	9.9	14.6	22.3	27.2	29.0
1999 [4]	6.6	*	*	*	*	0.4	3.2	9.5	14.4	23.3	28.9	27.5
Female												
2001	5.0	*	*	*	*	0.3	1.9	4.1	7.9	16.3	23.2	17.2
2000 [4]	4.8	*	*	*	*	0.3	1.5	3.6	7.6	16.0	23.5	17.2
1999 [4]	4.8	*	*	*	*	0.2	1.4	3.2	7.7	16.8	23.5	17.3
Non-Hispanic black												
Both sexes												
2001	3.9	*	*	*	*	*	3.2	11.2	13.2	14.5	13.6	7.8
2000 [4]	3.8	*	*	*	*	0.4	2.9	10.3	12.7	16.1	12.6	12.4
1999 [4]	3.9	*	*	*	*	0.5	3.1	10.1	14.3	15.0	14.5	12.3
Male												
2001	5.3	*	*	*	*	*	4.3	17.6	20.3	19.7	14.7	*
2000 [4]	4.8	*	*	*	*	*	3.6	15.5	18.9	19.8	14.8	*
1999 [4]	4.9	*	*	*	*	*	4.5	16.0	19.3	19.1	17.2	*
Female												
2001	2.7	*	*	*	*	*	2.1	5.7	7.5	10.8	13.0	*
2000 [4]	2.9	*	*	*	*	*	2.3	5.9	7.7	13.5	11.3	11.3
1999 [4]	2.9	*	*	*	*	*	1.9	5.0	10.4	12.1	13.0	12.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cholelithiasis and other disorders of gallbladder (K80-K82)												
All origins [3]												
Both sexes												
2001	1.0	*	*	*	*	*	0.1	0.3	1.0	2.9	7.5	23.7
2000 [4]	1.0	*	*	*	*	*	0.1	0.3	0.8	2.4	7.5	24.8
1999 [4]	1.0	*	*	*	*	*	0.1	0.3	0.9	2.6	7.8	23.9
Male												
2001	1.0	*	*	*	*	*	0.1	0.4	1.2	3.2	9.3	26.9
2000 [4]	0.9	*	*	*	*	*	0.1	0.3	0.9	2.9	9.4	27.6
1999 [4]	0.9	*	*	*	*	*	0.1	0.3	1.1	3.0	9.3	27.7
Female												
2001	1.1	*	*	*	*	*	*	0.3	0.8	2.6	6.3	22.4
2000 [4]	1.1	*	*	*	*	*	0.1	0.3	0.6	2.1	6.2	23.7
1999 [4]	1.1	*	*	*	*	*	0.1	0.4	0.7	2.2	6.8	22.4
Hispanic												
Both sexes												
2001	0.5	*	*	*	*	*	*	*	1.2	4.5	8.5	31.3
2000 [4]	0.5	*	*	*	*	*	*	*	*	3.8	10.1	29.2
1999 [4]	0.5	*	*	*	*	*	*	*	1.2	3.3	11.9	32.3
Male												
2001	0.5	*	*	*	*	*	*	*	*	5.0	11.2	*
2000 [4]	0.5	*	*	*	*	*	*	*	*	5.3	15.7	*
1999 [4]	0.5	*	*	*	*	*	*	*	*	*	15.1	*
Female												
2001	0.6	*	*	*	*	*	*	*	*	4.1	6.6	30.7

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Hispanic												
Female												
2000 [4]	0.5	*	*	*	*	*	*	*	*	*	*	31.7
1999 [4]	0.6	*	*	*	*	*	*	*	*	*	9.7	32.2
Non-Hispanic [5]												
Both sexes												
2001	1.1	*	*	*	*	*	0.1	0.3	0.9	2.8	7.5	23.4
2000 [4]	1.1	*	*	*	*	*	0.1	0.3	0.8	2.3	7.4	24.6
1999 [4]	1.1	*	*	*	*	*	0.1	0.3	0.8	2.5	7.6	23.6
Male												
2001	1.0	*	*	*	*	*	0.1	0.4	1.1	3.1	9.2	26.6
2000 [4]	1.0	*	*	*	*	*	0.1	0.3	0.9	2.7	9.2	27.7
1999 [4]	1.0	*	*	*	*	*	0.1	0.3	1.0	3.0	9.0	27.5
Female												
2001	1.2	*	*	*	*	*	*	0.3	0.8	2.5	6.3	22.1
2000 [4]	1.2	*	*	*	*	*	0.1	0.3	0.6	2.0	6.2	23.4
1999 [4]	1.2	*	*	*	*	*	0.1	0.3	0.7	2.1	6.6	22.0
Non-Hispanic white												
Both sexes												
2001	1.2	*	*	*	*	*	0.1	0.3	0.8	2.6	7.3	23.6
2000 [4]	1.2	*	*	*	*	*	0.1	0.2	0.7	2.2	7.3	25.3
1999 [4]	1.2	*	*	*	*	*	0.1	0.3	0.7	2.4	7.6	24.0
Male												
2001	1.1	*	*	*	*	*	*	0.4	1.0	2.9	9.2	26.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Non-Hispanic white												
Male												
2000 [4]	1.0	*	*	*	*	*	*	0.2	0.8	2.6	9.3	28.9
1999 [4]	1.1	*	*	*	*	*	*	0.3	0.9	2.9	9.3	27.8
Female												
2001	1.3	*	*	*	*	*	*	0.2	0.7	2.2	6.1	22.4
2000 [4]	1.3	*	*	*	*	*	*	0.2	0.6	1.8	6.0	23.9
1999 [4]	1.3	*	*	*	*	*	*	0.3	0.6	1.9	6.5	22.5
Non-Hispanic black												
Both sexes												
2001	0.9	*	*	*	*	*	*	0.7	1.7	4.8	8.3	20.0
2000 [4]	0.8	*	*	*	*	*	*	0.7	1.2	3.8	8.3	18.1
1999 [4]	0.7	*	*	*	*	*	0.4	0.5	1.5	3.7	7.3	15.5
Male												
2001	0.8	*	*	*	*	*	*	*	2.2	5.0	6.3	27.7
2000 [4]	0.6	*	*	*	*	*	*	*	*	3.4	8.0	*
1999 [4]	0.6	*	*	*	*	*	*	*	*	3.8	7.2	*
Female												
2001	1.0	*	*	*	*	*	*	*	*	4.6	9.4	17.1
2000 [4]	1.0	*	*	*	*	*	*	*	*	4.0	8.5	19.6
1999 [4]	0.8	*	*	*	*	*	*	*	*	3.6	7.4	15.5

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)												
All origins [3]												
Both sexes												
2001	13.9	3.3	*	*	0.2	0.6	1.7	4.6	13.0	40.2	104.2	287.7
2000 [4]	13.2	4.3	*	0.1	0.2	0.6	1.6	4.4	12.8	38.0	100.8	277.8
1999 [4]	12.7	4.4	*	0.1	0.2	0.6	1.6	4.0	12.0	37.1	97.6	268.9
Male												
2001	13.5	3.7	*	*	0.2	0.6	1.8	5.5	14.1	46.1	130.0	393.7
2000 [4]	12.9	5.3	*	*	0.2	0.6	1.9	5.3	14.2	42.9	127.3	380.7
1999 [4]	12.4	4.6	*	*	0.2	0.6	1.8	4.7	13.4	42.3	125.3	371.1
Female												
2001	14.2	2.9	*	*	0.2	0.6	1.6	3.9	11.9	35.2	87.1	243.3
2000 [4]	13.6	3.2	*	*	0.2	0.6	1.4	3.5	11.5	34.0	83.5	235.9
1999 [4]	13.0	4.1	*	*	0.2	0.6	1.3	3.4	10.8	33.0	79.6	227.7
Hispanic												
Both sexes												
2001	5.0	*	*	*	*	0.4	1.4	3.9	12.8	41.4	94.5	222.0
2000 [4]	4.7	3.2	*	*	*	0.4	1.0	3.5	13.6	37.5	84.7	229.6
1999 [4]	4.7	2.6	*	*	*	0.3	1.3	3.6	14.0	35.6	88.7	232.2
Male												
2001	4.7	*	*	*	*	*	1.6	3.8	15.2	43.3	112.2	281.9
2000 [4]	4.4	*	*	*	*	*	1.4	4.4	14.3	40.9	101.3	292.2
1999 [4]	4.7	*	*	*	*	*	1.2	4.4	17.8	39.1	110.5	336.4
Female												
2001	5.4	*	*	*	*	*	1.1	4.0	10.7	39.8	82.5	192.7

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)												
Hispanic												
Female												
2000 [4]	4.9	*	*	*	*	*	*	2.5	12.9	34.8	73.6	198.8
1999 [4]	4.7	*	*	*	*	*	1.3	2.9	10.7	32.9	74.2	182.0
Non-Hispanic [5]												
Both sexes												
2001	15.1	3.5	*	*	0.2	0.6	1.7	4.7	12.9	40.0	104.3	289.6
2000 [4]	14.4	4.4	*	0.1	0.2	0.7	1.7	4.4	12.6	37.9	101.2	279.0
1999 [4]	13.8	4.8	*	0.1	0.2	0.7	1.6	4.1	11.8	37.1	97.7	269.2
Male												
2001	14.8	3.7	*	*	0.2	0.6	1.9	5.6	14.0	46.1	130.6	397.6
2000 [4]	14.1	5.4	*	*	0.2	0.7	2.0	5.4	14.1	42.9	128.2	384.0
1999 [4]	13.5	4.8	*	*	0.2	0.7	1.9	4.7	13.0	42.3	125.7	370.9
Female												
2001	15.4	3.3	*	*	0.2	0.6	1.6	3.8	12.0	34.8	87.1	244.6
2000 [4]	14.7	3.4	*	*	0.2	0.6	1.5	3.6	11.3	33.8	83.6	236.6
1999 [4]	14.1	4.8	*	*	0.2	0.7	1.3	3.5	10.7	32.9	79.5	228.4
Non-Hispanic white												
Both sexes												
2001	14.9	2.4	*	*	0.2	0.4	1.1	3.1	9.3	33.5	96.6	279.7
2000 [4]	14.2	2.7	*	*	0.2	0.4	1.1	2.9	9.3	31.4	94.1	269.4
1999 [4]	13.4	3.5	*	*	0.1	0.4	1.0	2.6	8.7	31.1	89.9	255.3
Male												
2001	14.8	2.5	*	*	0.2	0.4	1.3	3.7	10.5	39.9	123.7	392.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)												
Non-Hispanic white												
Male												
2000 [4]	14.1	3.5	*	*	0.2	0.5	1.3	3.5	10.6	37.3	121.7	378.9
1999 [4]	13.3	3.9	*	*	*	0.5	1.2	2.9	9.9	36.6	117.3	359.6
Female												
2001	15.0	2.3	*	*	0.2	0.4	0.9	2.4	8.3	28.0	78.6	232.7
2000 [4]	14.2	1.8	*	*	0.2	0.4	1.0	2.3	8.0	26.5	76.0	225.2
1999 [4]	13.5	3.1	*	*	*	0.4	0.8	2.2	7.5	26.4	71.9	213.8
Non-Hispanic black												
Both sexes												
2001	20.2	8.5	*	*	0.5	1.8	5.5	16.2	42.5	104.2	204.0	432.9
2000 [4]	19.5	12.0	*	*	0.4	1.9	5.5	16.2	41.2	101.2	197.3	410.2
1999 [4]	19.2	10.2	*	*	0.5	2.0	5.6	15.6	37.8	96.6	196.4	436.8
Male												
2001	18.6	8.9	*	*	*	2.2	5.3	19.9	46.3	116.0	235.3	517.4
2000 [4]	18.3	13.6	*	*	*	2.0	6.7	20.8	45.6	105.7	234.2	496.7
1999 [4]	18.1	8.7	*	*	*	1.9	6.7	19.4	41.2	103.9	251.1	532.4
Female												
2001	21.7	8.1	*	*	*	1.4	5.6	13.0	39.4	95.8	186.5	401.7
2000 [4]	20.7	10.2	*	*	*	1.8	4.4	12.2	37.6	98.1	176.6	378.1
1999 [4]	20.2	11.8	*	*	*	2.0	4.7	12.3	35.1	91.5	165.8	401.1

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)												
All origins [3]												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.9
2000 [4]	0.1	*	*	*	*	*	*	*	*	0.2	0.4	1.2
1999 [4]	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.8
Male												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.5	1.8
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	0.7	*
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	0.7
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.3	0.9
1999 [4]	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.7
Hispanic												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)												
Hispanic												
Female												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.8
2000 [4]	0.1	*	*	*	*	*	*	*	*	0.2	0.4	1.2
1999 [4]	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.8
Male												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.5	1.8
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	0.7	*
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	0.7
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.4	0.9
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	0.4	0.7
Non-Hispanic white												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	0.8
2000 [4]	0.1	*	*	*	*	*	*	*	*	0.1	0.4	1.2
1999 [4]	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.8
Male												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)												
Non-Hispanic white												
Male												
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.5	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	0.6	*
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.4	1.0
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	0.4	*
Non-Hispanic black												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic glomerulonephritis, nephritis and nephropathy, not specified as acute or chronic, and renal sclerosis, unspecified (N02-N03,N05-N07,N26)												
All origins [3]												
Both sexes												
2001	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.2	2.8
2000 [4]	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.4	2.8
1999 [4]	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	1.6	3.2
Male												
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.8	1.7	3.7
2000 [4]	0.2	*	*	*	*	*	0.1	0.1	0.3	0.7	1.7	4.5
1999 [4]	0.2	*	*	*	*	*	*	0.1	0.3	0.5	2.2	5.0
Female												
2001	0.2	*	*	*	*	*	*	*	*	0.5	0.8	2.4
2000 [4]	0.2	*	*	*	*	*	*	*	0.2	0.5	1.2	2.1
1999 [4]	0.2	*	*	*	*	*	*	*	0.2	0.6	1.3	2.4
Hispanic												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic glomerulonephritis, nephritis and nephropathy, not specified as acute or chronic, and renal sclerosis, unspecified (N02-N03,N05-N07,N26)												
Hispanic												
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.2	2.8
2000 [4]	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.4	2.8
1999 [4]	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.7	3.2
Male												
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.8	1.7	3.7
2000 [4]	0.2	*	*	*	*	*	0.1	0.1	0.3	0.7	1.7	4.2
1999 [4]	0.2	*	*	*	*	*	*	0.1	0.3	0.5	2.3	5.1
Female												
2001	0.2	*	*	*	*	*	*	*	*	0.5	0.8	2.4
2000 [4]	0.2	*	*	*	*	*	*	*	0.2	0.5	1.2	2.2
1999 [4]	0.2	*	*	*	*	*	*	*	0.2	0.5	1.2	2.4
Non-Hispanic white												
Both sexes												
2001	0.2	*	*	*	*	*	*	*	0.1	0.6	1.1	2.8
2000 [4]	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.3	2.4
1999 [4]	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.6	2.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic glomerulonephritis, nephritis and nephropathy, not specified as acute or chronic, and renal sclerosis, unspecified (N02-N03,N05-N07,N26)												
Non-Hispanic white												
Male												
2001	0.2	*	*	*	*	*	*	*	*	0.8	1.7	4.0
2000 [4]	0.2	*	*	*	*	*	*	*	0.2	0.6	1.7	3.7
1999 [4]	0.2	*	*	*	*	*	*	*	*	0.5	2.2	4.6
Female												
2001	0.2	*	*	*	*	*	*	*	*	0.4	0.8	2.4
2000 [4]	0.2	*	*	*	*	*	*	*	*	0.5	1.1	1.9
1999 [4]	0.2	*	*	*	*	*	*	*	*	0.5	1.2	2.2
Non-Hispanic black												
Both sexes												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Renal failure (N17-N19)												
All origins [3]												
Both sexes												
2001	13.6	3.1	*	*	0.2	0.5	1.6	4.5	12.7	39.4	102.6	284.0
2000 [4]	13.0	4.0	*	*	0.2	0.6	1.6	4.2	12.5	37.2	98.9	273.7
1999 [4]	12.4	4.3	*	0.1	0.2	0.6	1.5	3.9	11.7	36.4	95.3	264.8
Male												
2001	13.2	3.5	*	*	0.2	0.5	1.8	5.3	13.8	45.2	128.0	388.7
2000 [4]	12.6	5.1	*	*	0.2	0.6	1.8	5.1	13.8	42.1	125.1	374.2
1999 [4]	12.1	4.5	*	*	0.1	0.5	1.7	4.5	13.1	41.5	122.3	364.6
Female												
2001	14.0	2.7	*	*	0.2	0.5	1.5	3.7	11.8	34.5	85.9	240.1
2000 [4]	13.3	2.8	*	*	0.2	0.5	1.3	3.4	11.2	33.3	81.8	232.8
1999 [4]	12.7	4.1	*	*	0.2	0.6	1.3	3.3	10.4	32.2	77.8	224.5
Hispanic												
Both sexes												
2001	4.9	*	*	*	*	0.3	1.2	3.7	12.5	40.5	93.2	217.6
2000 [4]	4.5	2.9	*	*	*	0.4	1.0	3.3	13.3	36.9	83.0	225.6
1999 [4]	4.5	*	*	*	*	*	1.2	3.5	13.6	34.4	86.9	228.0
Male												
2001	4.6	*	*	*	*	*	1.4	3.7	14.6	42.5	110.4	278.1
2000 [4]	4.3	*	*	*	*	*	1.3	4.2	13.8	40.5	99.3	280.1
1999 [4]	4.5	*	*	*	*	*	1.1	4.3	17.2	38.2	109.0	330.0
Female												
2001	5.2	*	*	*	*	*	1.0	3.7	10.5	38.9	81.6	188.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Renal failure (N17-N19)												
Hispanic												
Female												
2000 [4]	4.8	*	*	*	*	*	*	2.4	12.8	34.0	72.0	198.8
1999 [4]	4.5	*	*	*	*	*	1.3	2.7	10.4	31.3	72.2	178.8
Non-Hispanic [5]												
Both sexes												
2001	14.9	3.3	*	*	0.2	0.6	1.7	4.6	12.7	39.2	102.7	285.9
2000 [4]	14.1	4.2	*	*	0.2	0.6	1.6	4.3	12.3	37.1	99.3	274.9
1999 [4]	13.5	4.7	*	*	0.2	0.6	1.5	3.9	11.5	36.4	95.4	265.0
Male												
2001	14.6	3.5	*	*	0.2	0.6	1.8	5.4	13.7	45.3	128.5	392.7
2000 [4]	13.8	5.2	*	*	0.2	0.6	1.9	5.1	13.7	42.0	125.9	377.7
1999 [4]	13.2	4.6	*	*	0.1	0.6	1.8	4.5	12.7	41.5	122.6	364.4
Female												
2001	15.2	3.1	*	*	0.2	0.5	1.6	3.7	11.8	34.1	85.8	241.5
2000 [4]	14.4	3.1	*	*	0.2	0.5	1.4	3.5	11.0	33.2	82.0	233.4
1999 [4]	13.8	4.8	*	*	0.2	0.6	1.3	3.3	10.4	32.1	77.8	225.2
Non-Hispanic white												
Both sexes												
2001	14.6	2.2	*	*	0.2	0.4	1.1	3.0	9.2	32.7	95.1	276.1
2000 [4]	13.9	2.5	*	*	0.2	0.4	1.1	2.8	9.0	30.7	92.3	265.7
1999 [4]	13.1	3.5	*	*	0.1	0.4	0.9	2.4	8.4	30.4	87.8	251.6
Male												
2001	14.5	2.3	*	*	*	0.3	1.3	3.6	10.3	39.1	121.7	387.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Renal failure (N17-N19)												
Non-Hispanic white												
Male												
2000 [4]	13.7	3.3	*	*	0.2	0.4	1.2	3.3	10.3	36.5	119.4	373.4
1999 [4]	13.0	3.8	*	*	*	0.4	1.1	2.8	9.6	35.9	114.4	353.8
Female												
2001	14.7	2.1	*	*	0.2	0.4	0.9	2.4	8.1	27.3	77.5	229.7
2000 [4]	14.0	*	*	*	*	0.3	0.9	2.3	7.7	25.8	74.4	222.2
1999 [4]	13.2	3.1	*	*	*	0.4	0.8	2.1	7.3	25.8	70.3	210.8
Non-Hispanic black												
Both sexes												
2001	20.0	8.5	*	*	0.4	1.7	5.3	15.8	42.1	102.9	201.9	428.9
2000 [4]	19.2	11.4	*	*	0.4	1.8	5.3	15.6	40.7	99.9	194.6	404.2
1999 [4]	18.8	10.2	*	*	0.5	1.9	5.4	15.2	37.0	95.3	193.0	428.7
Male												
2001	18.4	8.9	*	*	*	2.1	5.1	19.3	45.7	114.3	233.5	514.0
2000 [4]	17.9	13.6	*	*	*	1.9	6.5	19.9	45.0	104.2	231.7	487.3
1999 [4]	17.7	8.7	*	*	*	1.8	6.4	19.0	40.5	102.5	246.7	519.4
Female												
2001	21.4	8.1	*	*	*	1.4	5.5	12.8	39.2	94.9	184.1	397.4
2000 [4]	20.3	9.2	*	*	*	1.7	4.1	11.8	37.3	96.9	173.7	373.3
1999 [4]	19.8	11.8	*	*	*	2.0	4.5	12.0	34.3	90.2	163.0	394.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of kidney (N25,N27)												
All origins [3]												
Both sexes												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of kidney (N25,N27)												
Hispanic												
Female												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of kidney (N25,N27)												
Non-Hispanic white												
Male												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Infections of kidney (N10-N12,N13.6,N15.1)												
All origins [3]												
Both sexes												
2001	0.3	*	*	*	*	*	0.1	0.1	0.2	0.5	2.0	6.0
2000 [4]	0.3	*	*	*	*	*	0.0	0.1	0.2	0.6	2.0	7.5
1999 [4]	0.3	*	*	*	*	*	0.1	0.2	0.2	0.7	2.0	7.5
Male												
2001	0.2	*	*	*	*	*	*	*	*	0.5	1.5	4.3
2000 [4]	0.2	*	*	*	*	*	*	*	*	0.3	1.5	5.6
1999 [4]	0.2	*	*	*	*	*	*	*	0.2	0.5	1.7	6.8
Female												
2001	0.4	*	*	*	*	*	0.1	0.2	0.3	0.5	2.3	6.7
2000 [4]	0.4	*	*	*	*	*	*	0.1	0.3	0.8	2.3	8.3
1999 [4]	0.4	*	*	*	*	*	0.1	0.2	0.3	0.9	2.1	7.8
Hispanic												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Infections of kidney (N10-N12,N13.6,N15.1)												
Hispanic												
Female												
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.3	*	*	*	*	*	0.1	0.1	0.2	0.5	2.0	6.1
2000 [4]	0.3	*	*	*	*	*	*	0.1	0.2	0.6	2.0	7.6
1999 [4]	0.3	*	*	*	*	*	0.0	0.1	0.2	0.7	1.9	7.5
Male												
2001	0.2	*	*	*	*	*	*	*	*	0.5	1.6	4.3
2000 [4]	0.2	*	*	*	*	*	*	*	*	0.3	1.5	5.8
1999 [4]	0.2	*	*	*	*	*	*	*	0.2	0.5	1.6	6.9
Female												
2001	0.4	*	*	*	*	*	0.1	0.2	0.3	0.4	2.3	6.9
2000 [4]	0.5	*	*	*	*	*	*	0.1	0.3	0.8	2.3	8.3
1999 [4]	0.4	*	*	*	*	*	*	0.2	0.2	0.9	2.1	7.7
Non-Hispanic white												
Both sexes												
2001	0.3	*	*	*	*	*	0.1	0.1	0.2	0.4	2.0	6.2
2000 [4]	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.9	7.6
1999 [4]	0.3	*	*	*	*	*	*	0.1	0.2	0.7	1.9	7.7
Male												
2001	0.2	*	*	*	*	*	*	*	*	0.4	1.5	4.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Infections of kidney (N10-N12,N13.6,N15.1)												
Non-Hispanic white												
Male												
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	1.5	6.0
1999 [4]	0.2	*	*	*	*	*	*	*	*	0.4	1.7	7.3
Female												
2001	0.4	*	*	*	*	*	*	0.2	0.3	0.5	2.3	7.1
2000 [4]	0.5	*	*	*	*	*	*	*	0.3	0.7	2.1	8.2
1999 [4]	0.5	*	*	*	*	*	*	0.2	0.2	0.9	2.1	7.9
Non-Hispanic black												
Both sexes												
2001	0.2	*	*	*	*	*	*	*	*	*	2.4	*
2000 [4]	0.3	*	*	*	*	*	*	*	*	*	3.1	7.9
1999 [4]	0.2	*	*	*	*	*	*	*	*	1.2	*	*
Male												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.4	*	*	*	*	*	*	*	*	*	3.6	9.2
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hyperplasia of prostate (N40)												
All origins [3]												
Both sexes												
2001	0.2	*	*	*	*	*	*	*	*	0.2	1.1	5.5
2000 [4]	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.2
1999 [4]	0.2	*	*	*	*	*	*	*	*	0.3	1.3	5.0
Male												
2001	0.3	*	*	*	*	*	*	*	*	0.5	2.8	18.6
2000 [4]	0.3	*	*	*	*	*	*	*	*	0.5	3.3	18.1
1999 [4]	0.3	*	*	*	*	*	*	*	*	0.6	3.3	17.5
Female												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Hispanic												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hyperplasia of prostate (N40)												
Hispanic												
Female												
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic [5]												
Both sexes												
2001	0.2	*	*	*	*	*	*	*	*	0.2	1.1	5.7
2000 [4]	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.2
1999 [4]	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.1
Male												
2001	0.4	*	*	*	*	*	*	*	*	0.5	2.8	19.3
2000 [4]	0.3	*	*	*	*	*	*	*	*	0.4	3.3	18.0
1999 [4]	0.3	*	*	*	*	*	*	*	*	0.5	3.4	18.0
Female												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic white												
Both sexes												
2001	0.2	*	*	*	*	*	*	*	*	0.2	1.1	5.9
2000 [4]	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.2
1999 [4]	0.2	*	*	*	*	*	*	*	*	0.2	1.4	5.2
Male												
2001	0.4	*	*	*	*	*	*	*	*	0.5	2.8	20.1

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hyperplasia of prostate (N40)												
Non-Hispanic white												
Male												
2000 [4]	0.4	*	*	*	*	*	*	*	*	0.4	3.4	18.0
1999 [4]	0.4	*	*	*	*	*	*	*	*	0.5	3.4	18.4
Female												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic black												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Inflammatory diseases of female pelvic organs (N70-N76)												
All origins [3]												
Both sexes												
2001	0.0	*	*	*	*	*	*	*	*	*	0.2	0.9
2000 [4]	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.6
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	0.3	*
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2000 [4]	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.9
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	0.4	*
Hispanic												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Inflammatory diseases of female pelvic organs (N70-N76)												
Hispanic												
Female												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.0	*	*	*	*	*	*	*	*	*	0.3	0.9
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	0.2	0.6
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	0.3	*
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.4	0.9
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	0.4	*
Non-Hispanic white												
Both sexes												
2001	0.0	*	*	*	*	*	*	*	*	*	0.3	0.9
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	0.2	0.6
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	0.2	*
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Inflammatory diseases of female pelvic organs (N70-N76)												
Non-Hispanic white												
Male												
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	0.4	0.9
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	0.4	*
Non-Hispanic black												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy, childbirth and the puerperium (O00-O99)												
All origins [3]												
Both sexes												
2001	0.1	. . .	. . .	*	0.3	0.5	0.2	*	*	*	*	*
2000 [4]	0.1	. . .	. . .	*	0.2	0.5	0.3	*	*	*	*	*
1999 [4]	0.1	. . .	. . .	*	0.3	0.5	0.3	*	*	*	*	*
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	0.3	. . .	. . .	*	0.6	1.0	0.5	*	*	*	*	*
2000 [4]	0.3	. . .	. . .	*	0.5	0.9	0.5	*	*	*	*	*
1999 [4]	0.3	. . .	. . .	*	0.5	0.9	0.5	*	*	*	*	*
Hispanic												
Both sexes												
2001	0.2	. . .	. . .	*	0.4	0.6	*	*	*	*	*	*
2000 [4]	0.2	. . .	. . .	*	0.4	0.6	0.4	*	*	*	*	*
1999 [4]	0.2	. . .	. . .	*	*	0.4	0.5	*	*	*	*	*
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	0.5	. . .	. . .	*	0.8	1.2	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy, childbirth and the puerperium (O00-O99)												
Hispanic												
Female												
2000 [4]	0.5	. . .	. . .	*	0.8	1.2	0.9	*	*	*	*	*
1999 [4]	0.4	. . .	. . .	*	*	0.9	1.0	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.1	. . .	. . .	*	0.3	0.5	0.2	*	*	*	*	*
2000 [4]	0.1	. . .	. . .	*	0.2	0.4	0.2	*	*	*	*	*
1999 [4]	0.1	. . .	. . .	*	0.2	0.5	0.2	*	*	*	*	*
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	0.3	. . .	. . .	*	0.5	0.9	0.4	*	*	*	*	*
2000 [4]	0.3	. . .	. . .	*	0.4	0.9	0.5	*	*	*	*	*
1999 [4]	0.3	. . .	. . .	*	0.5	0.9	0.5	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2001	0.1	. . .	. . .	*	0.2	0.3	0.1	*	*	*	*	*
2000 [4]	0.1	. . .	. . .	*	0.1	0.3	0.2	*	*	*	*	*
1999 [4]	0.1	. . .	. . .	*	0.1	0.3	0.1	*	*	*	*	*
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy, childbirth and the puerperium (O00-O99)												
Non-Hispanic white												
Male												
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	0.2	. . .	. . .	*	0.3	0.6	0.2	*	*	*	*	*
2000 [4]	0.2	. . .	. . .	*	0.3	0.6	0.3	*	*	*	*	*
1999 [4]	0.2	. . .	. . .	*	0.2	0.6	0.3	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2001	0.4	. . .	. . .	*	0.8	1.3	0.7	*	*	*	*	*
2000 [4]	0.4	. . .	. . .	*	0.6	1.3	0.7	*	*	*	*	*
1999 [4]	0.5	. . .	. . .	*	0.8	1.3	0.7	*	*	*	*	*
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	0.8	. . .	. . .	*	1.6	2.4	1.4	*	*	*	*	*
2000 [4]	0.8	. . .	. . .	*	1.2	2.4	1.3	*	*	*	*	*
1999 [4]	0.9	. . .	. . .	*	1.6	2.5	1.4	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy with abortive outcome (O00-O07)												
All origins [3]												
Both sexes												
2001	0.0	. . .	. . .	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	. . .	. . .	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	. . .	. . .	*	*	*	*	*	*	*	*	*
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	0.0	. . .	. . .	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	. . .	. . .	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	. . .	. . .	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2001	*	. . .	. . .	*	*	*	*	*	*	*	*	*
2000 [4]	*	. . .	. . .	*	*	*	*	*	*	*	*	*
1999 [4]	*	. . .	. . .	*	*	*	*	*	*	*	*	*
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	*	. . .	. . .	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy with abortive outcome (O00-O07)												
Hispanic												
Female												
2000 [4]	*	. . .	. . .	*	*	*	*	*	*	*	*	*
1999 [4]	*	. . .	. . .	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.0	. . .	. . .	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	. . .	. . .	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	. . .	. . .	*	*	*	*	*	*	*	*	*
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	0.0	. . .	. . .	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	. . .	. . .	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	. . .	. . .	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2001	*	. . .	. . .	*	*	*	*	*	*	*	*	*
2000 [4]	*	. . .	. . .	*	*	*	*	*	*	*	*	*
1999 [4]	*	. . .	. . .	*	*	*	*	*	*	*	*	*
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy with abortive outcome (O00-O07)												
Non-Hispanic white												
Male												
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	*	. . .	. . .	*	*	*	*	*	*	*	*	*
2000 [4]	*	. . .	. . .	*	*	*	*	*	*	*	*	*
1999 [4]	*	. . .	. . .	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2001	*	. . .	. . .	*	*	*	*	*	*	*	*	*
2000 [4]	*	. . .	. . .	*	*	*	*	*	*	*	*	*
1999 [4]	*	. . .	. . .	*	*	*	*	*	*	*	*	*
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	*	. . .	. . .	*	*	*	*	*	*	*	*	*
2000 [4]	*	. . .	. . .	*	*	*	*	*	*	*	*	*
1999 [4]	*	. . .	. . .	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other complications of pregnancy, childbirth and the puerperium (O10-O99)												
All origins [3]												
Both sexes												
2001	0.1	. . .	. . .	*	0.2	0.4	0.2	*	*	*	*	*
2000 [4]	0.1	. . .	. . .	*	0.2	0.4	0.2	*	*	*	*	*
1999 [4]	0.1	. . .	. . .	*	0.2	0.4	0.3	*	*	*	*	*
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	0.3	. . .	. . .	*	0.5	0.9	0.4	*	*	*	*	*
2000 [4]	0.3	. . .	. . .	*	0.5	0.8	0.5	*	*	*	*	*
1999 [4]	0.3	. . .	. . .	*	0.5	0.8	0.5	*	*	*	*	*
Hispanic												
Both sexes												
2001	0.2	. . .	. . .	*	0.4	0.6	*	*	*	*	*	*
2000 [4]	0.2	. . .	. . .	*	0.3	0.5	0.4	*	*	*	*	*
1999 [4]	0.2	. . .	. . .	*	*	0.4	0.5	*	*	*	*	*
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	0.5	. . .	. . .	*	0.8	1.2	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other complications of pregnancy, childbirth and the puerperium (O10-O99)												
Hispanic												
Female												
2000 [4]	0.5	. . .	. . .	*	0.7	1.1	0.9	*	*	*	*	*
1999 [4]	0.4	. . .	. . .	*	*	0.9	1.0	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.1	. . .	. . .	*	0.2	0.4	0.2	*	*	*	*	*
2000 [4]	0.1	. . .	. . .	*	0.2	0.4	0.2	*	*	*	*	*
1999 [4]	0.1	. . .	. . .	*	0.2	0.4	0.2	*	*	*	*	*
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	0.2	. . .	. . .	*	0.4	0.8	0.4	*	*	*	*	*
2000 [4]	0.2	. . .	. . .	*	0.4	0.8	0.4	*	*	*	*	*
1999 [4]	0.2	. . .	. . .	*	0.4	0.8	0.4	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2001	0.1	. . .	. . .	*	0.1	0.3	0.1	*	*	*	*	*
2000 [4]	0.1	. . .	. . .	*	0.1	0.3	0.1	*	*	*	*	*
1999 [4]	0.1	. . .	. . .	*	0.1	0.3	0.1	*	*	*	*	*
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other complications of pregnancy, childbirth and the puerperium (O10-O99)												
Non-Hispanic white												
Male												
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	0.1	. . .	. . .	*	0.3	0.6	0.2	*	*	*	*	*
2000 [4]	0.1	. . .	. . .	*	0.3	0.5	0.3	*	*	*	*	*
1999 [4]	0.1	. . .	. . .	*	0.2	0.5	0.3	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2001	0.4	. . .	. . .	*	0.7	1.2	0.6	*	*	*	*	*
2000 [4]	0.3	. . .	. . .	*	0.5	1.1	0.6	*	*	*	*	*
1999 [4]	0.4	. . .	. . .	*	0.8	1.2	0.7	*	*	*	*	*
Male												
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999 [4]	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female												
2001	0.7	. . .	. . .	*	1.3	2.2	1.2	*	*	*	*	*
2000 [4]	0.7	. . .	. . .	*	1.1	2.1	1.1	*	*	*	*	*
1999 [4]	0.8	. . .	. . .	*	1.5	2.3	1.3	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain conditions originating in the perinatal period (P00-P96)												
All origins [3]												
Both sexes												
2001	4.9	340.5	0.5	0.1	*	*	*	*	*	*	*	*
2000 [4]	5.0	365.7	0.5	0.1	*	*	*	*	*	*	*	*
1999 [4]	5.1	371.0	0.6	0.1	*	*	*	*	*	*	*	*
Male												
2001	5.6	376.6	0.6	0.1	*	*	*	*	*	*	*	*
2000 [4]	5.8	403.9	0.7	0.1	*	*	*	*	*	*	*	*
1999 [4]	5.9	408.6	0.7	0.1	*	*	*	*	*	*	*	*
Female												
2001	4.2	302.6	0.4	*	*	*	*	*	*	*	*	*
2000 [4]	4.3	325.6	0.4	*	*	*	*	*	*	*	*	*
1999 [4]	4.4	331.7	0.5	0.1	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2001	6.0	269.6	*	*	*	*	*	*	*	*	*	*
2000 [4]	6.1	274.9	*	*	*	*	*	*	*	*	*	*
1999 [4]	6.3	281.4	*	*	*	*	*	*	*	*	*	*
Male												
2001	6.7	302.0	*	*	*	*	*	*	*	*	*	*
2000 [4]	6.6	299.0	*	*	*	*	*	*	*	*	*	*
1999 [4]	6.8	305.4	*	*	*	*	*	*	*	*	*	*
Female												
2001	5.3	235.9	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain conditions originating in the perinatal period (P00-P96)												
Hispanic												
Female												
2000 [4]	5.5	249.7	*	*	*	*	*	*	*	*	*	*
1999 [4]	5.8	256.4	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	4.6	353.5	0.5	0.1	*	*	*	*	*	*	*	*
2000 [4]	4.7	380.5	0.5	0.1	*	*	*	*	*	*	*	*
1999 [4]	4.8	383.8	0.6	0.1	*	*	*	*	*	*	*	*
Male												
2001	5.4	390.1	0.6	0.1	*	*	*	*	*	*	*	*
2000 [4]	5.5	421.7	0.7	*	*	*	*	*	*	*	*	*
1999 [4]	5.6	423.2	0.8	0.1	*	*	*	*	*	*	*	*
Female												
2001	3.9	315.1	0.3	*	*	*	*	*	*	*	*	*
2000 [4]	4.0	337.2	0.4	*	*	*	*	*	*	*	*	*
1999 [4]	4.1	342.4	0.4	0.1	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2001	3.2	260.3	0.4	*	*	*	*	*	*	*	*	*
2000 [4]	3.2	275.5	0.4	0.1	*	*	*	*	*	*	*	*
1999 [4]	3.2	275.1	0.4	0.1	*	*	*	*	*	*	*	*
Male												
2001	3.7	286.6	0.5	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain conditions originating in the perinatal period (P00-P96)												
Non-Hispanic white												
Male												
2000 [4]	3.7	305.5	0.5	*	*	*	*	*	*	*	*	*
1999 [4]	3.8	304.4	0.5	*	*	*	*	*	*	*	*	*
Female												
2001	2.7	232.8	*	*	*	*	*	*	*	*	*	*
2000 [4]	2.7	243.9	*	*	*	*	*	*	*	*	*	*
1999 [4]	2.7	244.2	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2001	13.2	765.0	0.9	*	*	*	*	*	*	*	*	*
2000 [4]	13.9	838.3	1.1	*	*	*	*	*	*	*	*	*
1999 [4]	14.4	872.7	1.2	*	*	*	*	*	*	*	*	*
Male												
2001	15.8	846.6	*	*	*	*	*	*	*	*	*	*
2000 [4]	16.6	932.9	*	*	*	*	*	*	*	*	*	*
1999 [4]	17.1	968.0	*	*	*	*	*	*	*	*	*	*
Female												
2001	10.9	679.1	*	*	*	*	*	*	*	*	*	*
2000 [4]	11.5	740.7	*	*	*	*	*	*	*	*	*	*
1999 [4]	12.0	774.4	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
All origins [3]												
Both sexes												
2001	3.7	136.7	3.6	0.9	1.3	1.2	1.3	1.6	2.3	2.4	3.7	7.2
2000 [4]	3.8	150.9	3.2	1.0	1.1	1.2	1.3	1.6	2.3	2.5	3.9	7.5
1999 [4]	3.7	144.2	3.6	1.0	1.1	1.2	1.2	1.8	2.3	2.5	4.0	7.8
Male												
2001	3.9	143.1	3.6	1.0	1.5	1.2	1.4	1.7	2.3	2.7	4.1	7.7
2000 [4]	4.0	155.4	3.3	1.0	1.3	1.4	1.4	1.9	2.3	2.7	4.0	10.6
1999 [4]	3.9	147.9	4.0	1.0	1.3	1.2	1.3	1.9	2.5	2.5	4.2	8.5
Female												
2001	3.4	129.9	3.7	0.8	1.0	1.1	1.2	1.6	2.3	2.2	3.5	7.0
2000 [4]	3.5	146.2	3.1	1.0	0.9	1.0	1.1	1.4	2.4	2.4	3.9	6.3
1999 [4]	3.5	140.3	3.2	1.1	0.9	1.1	1.1	1.6	2.2	2.6	3.9	7.6
Hispanic												
Both sexes												
2001	4.4	146.2	3.4	0.8	1.1	0.7	0.8	1.1	1.3	1.9	4.3	*
2000 [4]	4.6	153.3	3.8	0.8	0.8	0.7	0.8	1.1	1.8	2.0	4.7	*
1999 [4]	4.4	141.8	3.5	0.9	0.9	0.9	0.7	1.4	1.6	2.9	*	*
Male												
2001	4.7	153.4	3.5	1.0	1.1	0.7	1.0	1.5	*	*	*	*
2000 [4]	4.7	153.3	4.0	0.8	1.1	0.7	1.0	1.4	*	*	*	*
1999 [4]	4.3	138.1	4.3	0.9	0.9	0.7	*	1.6	*	*	*	*
Female												
2001	4.1	138.8	3.3	0.6	1.1	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
Hispanic												
Female												
2000 [4]	4.4	153.3	3.5	0.7	*	0.7	*	*	*	*	*	*
1999 [4]	4.5	145.8	2.7	0.9	0.9	1.0	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	3.5	132.8	3.7	0.9	1.3	1.3	1.4	1.7	2.4	2.5	3.7	7.2
2000 [4]	3.6	148.8	3.1	1.0	1.2	1.3	1.3	1.7	2.4	2.6	3.9	7.5
1999 [4]	3.6	143.1	3.6	1.1	1.2	1.2	1.3	1.8	2.4	2.5	4.0	7.9
Male												
2001	3.7	139.4	3.6	1.0	1.6	1.4	1.5	1.7	2.3	2.7	4.0	7.6
2000 [4]	3.9	154.4	3.1	1.0	1.4	1.5	1.5	1.9	2.2	2.7	3.9	10.4
1999 [4]	3.8	148.4	3.9	1.1	1.4	1.3	1.4	1.9	2.6	2.4	4.2	8.5
Female												
2001	3.3	125.8	3.7	0.9	1.0	1.2	1.3	1.7	2.4	2.2	3.5	7.0
2000 [4]	3.4	142.8	3.0	1.0	1.0	1.1	1.2	1.4	2.5	2.4	3.9	6.4
1999 [4]	3.4	137.4	3.3	1.1	0.9	1.1	1.2	1.6	2.2	2.6	3.9	7.6
Non-Hispanic white												
Both sexes												
2001	3.4	127.4	3.1	0.9	1.2	1.3	1.4	1.7	2.5	2.5	3.8	7.4
2000 [4]	3.4	141.2	2.6	0.9	1.1	1.3	1.3	1.7	2.4	2.6	4.0	7.8
1999 [4]	3.4	136.8	3.0	0.9	1.1	1.3	1.3	1.8	2.5	2.5	4.0	8.0
Male												
2001	3.6	135.5	3.1	1.0	1.4	1.5	1.5	1.7	2.4	2.9	4.1	7.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
Non-Hispanic white												
Male												
2000 [4]	3.6	147.9	2.6	0.9	1.3	1.6	1.4	1.9	2.3	2.6	4.1	10.8
1999 [4]	3.6	144.4	3.2	0.9	1.4	1.3	1.4	2.0	2.7	2.4	4.1	8.6
Female												
2001	3.2	118.8	3.2	0.7	1.0	1.2	1.3	1.7	2.6	2.3	3.6	7.4
2000 [4]	3.2	134.2	2.6	1.0	1.0	1.1	1.2	1.6	2.5	2.6	4.0	6.6
1999 [4]	3.3	128.9	2.9	1.0	0.8	1.2	1.2	1.7	2.3	2.6	4.0	7.8
Non-Hispanic black												
Both sexes												
2001	4.7	163.5	5.4	1.1	2.0	1.3	1.6	1.8	1.9	2.1	3.0	*
2000 [4]	5.0	184.1	5.2	1.4	1.6	1.4	1.6	1.6	2.4	2.7	2.8	*
1999 [4]	4.8	174.8	5.8	1.6	1.5	1.1	1.4	1.9	2.0	2.9	4.3	7.4
Male												
2001	5.1	167.5	5.5	1.0	2.5	1.2	1.8	1.7	2.1	*	*	*
2000 [4]	5.5	186.0	5.6	1.5	2.1	1.7	1.9	2.2	*	3.9	*	*
1999 [4]	5.3	179.5	6.6	1.6	1.8	1.3	1.6	2.3	2.1	*	*	*
Female												
2001	4.4	159.2	5.4	1.3	1.4	1.4	1.5	1.8	1.8	2.4	*	*
2000 [4]	4.5	182.2	4.8	1.2	1.2	1.1	1.4	1.2	3.1	*	*	*
1999 [4]	4.4	170.0	5.0	1.6	1.1	1.0	1.2	1.5	1.9	3.2	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
All origins [3]												
Both sexes												
2001	11.4	86.5	1.8	0.4	2.4	3.9	6.9	8.9	9.1	15.1	39.8	206.4
2000 [4]	11.3	95.6	1.6	0.4	2.2	3.9	6.6	8.3	9.3	16.5	40.7	208.1
1999 [4]	9.6	93.2	1.5	0.3	1.6	2.7	4.6	5.9	7.8	14.6	36.1	187.7
Male												
2001	11.3	98.9	2.1	0.5	3.2	5.2	9.0	11.9	12.0	18.0	43.8	175.0
2000 [4]	11.5	110.6	1.7	0.5	2.9	5.3	8.9	11.5	12.3	20.6	46.0	182.7
1999 [4]	9.4	106.6	1.7	0.4	2.1	3.6	6.0	8.1	10.4	18.2	40.1	166.3
Female												
2001	11.4	73.6	1.5	0.4	1.5	2.5	4.9	6.1	6.3	12.7	37.2	219.5
2000 [4]	11.2	79.9	1.5	0.4	1.5	2.5	4.4	5.3	6.5	13.2	37.2	218.5
1999 [4]	9.7	79.2	1.3	0.3	1.0	1.9	3.2	3.8	5.5	11.7	33.5	196.4
Hispanic												
Both sexes												
2001	5.9	60.7	1.6	0.5	2.8	4.2	6.0	8.3	6.9	9.9	19.3	108.2
2000 [4]	5.9	65.6	1.7	0.5	2.1	4.0	6.7	8.0	6.7	9.9	20.5	102.2
1999 [4]	3.8	54.9	1.0	*	1.3	2.2	3.0	3.1	3.8	7.0	19.4	87.0
Male												
2001	7.4	67.2	1.8	*	3.9	6.0	8.5	12.6	10.6	14.2	25.0	87.6
2000 [4]	7.5	78.6	2.1	*	2.9	5.8	9.9	12.4	9.0	14.3	25.6	106.8
1999 [4]	4.4	61.3	*	*	1.9	2.9	4.3	4.7	5.5	8.3	24.0	77.6
Female												
2001	4.3	53.9	*	0.6	1.5	2.1	3.4	4.1	3.6	6.5	15.4	118.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
Hispanic												
Female												
2000 [4]	4.1	52.1	1.4	*	1.3	2.0	3.2	3.7	4.6	6.5	17.2	99.9
1999 [4]	3.1	48.2	*	*	*	1.3	1.6	1.7	2.4	6.0	16.3	91.5
Non-Hispanic [5]												
Both sexes												
2001	12.1	92.6	1.8	0.4	2.3	3.7	7.0	8.9	9.2	15.4	40.7	209.6
2000 [4]	12.0	102.4	1.6	0.4	2.2	3.8	6.5	8.2	9.3	16.8	41.4	211.6
1999 [4]	10.3	101.9	1.6	0.4	1.6	2.8	4.7	6.0	8.0	15.0	36.7	190.9
Male												
2001	11.8	106.2	2.1	0.5	3.0	4.9	9.0	11.7	12.0	18.2	44.6	178.2
2000 [4]	11.9	118.0	1.7	0.5	2.9	5.1	8.6	11.2	12.3	20.7	46.6	185.4
1999 [4]	10.0	116.7	1.9	0.4	2.2	3.7	6.1	8.2	10.6	18.7	40.6	169.5
Female												
2001	12.3	78.3	1.5	0.4	1.5	2.6	5.1	6.2	6.6	13.1	38.0	222.7
2000 [4]	12.1	86.1	1.5	0.4	1.5	2.6	4.5	5.4	6.6	13.6	38.0	222.3
1999 [4]	10.5	86.3	1.4	0.3	1.0	2.0	3.3	3.9	5.7	11.9	34.2	199.5
Non-Hispanic white												
Both sexes												
2001	12.4	73.1	1.5	0.3	2.2	3.6	6.9	8.5	8.6	14.7	40.4	213.3
2000 [4]	12.2	79.7	1.4	0.3	2.1	3.6	6.3	7.6	8.6	15.9	40.9	215.8
1999 [4]	10.2	79.3	1.4	0.3	1.4	2.5	4.1	5.2	7.3	14.3	35.7	193.8
Male												
2001	11.8	85.2	1.7	0.4	2.9	4.9	8.8	11.1	10.9	17.3	44.7	180.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
Non-Hispanic white												
Male												
2000 [4]	11.8	91.8	1.7	0.4	2.8	4.8	8.5	10.3	11.1	19.3	46.0	189.6
1999 [4]	9.6	92.3	1.7	0.3	1.9	3.3	5.5	6.9	9.5	17.7	39.2	171.1
Female												
2001	13.1	60.4	1.3	0.3	1.5	2.4	5.0	5.9	6.4	12.5	37.6	227.0
2000 [4]	12.6	66.9	1.2	0.3	1.3	2.3	4.1	4.9	6.2	13.0	37.5	226.4
1999 [4]	10.9	65.5	1.0	0.3	0.9	1.8	2.8	3.5	5.2	11.4	33.4	202.8
Non-Hispanic black												
Both sexes												
2001	12.9	176.9	3.0	0.6	2.7	5.2	9.3	14.0	15.1	24.6	50.3	194.2
2000 [4]	13.9	205.6	2.3	0.7	3.2	5.9	9.3	14.8	17.3	29.2	56.1	189.2
1999 [4]	13.0	202.0	2.9	0.5	2.7	5.0	9.1	13.2	16.0	24.6	54.8	182.0
Male												
2001	14.2	195.9	3.7	0.8	3.8	6.1	11.4	18.2	23.8	30.7	52.2	186.0
2000 [4]	15.7	236.8	2.2	0.9	3.8	7.8	10.9	20.6	25.1	40.0	66.0	163.2
1999 [4]	14.9	228.0	2.9	0.6	3.6	6.9	11.6	19.4	23.0	32.8	66.9	180.7
Female												
2001	11.7	156.8	2.2	*	1.7	4.3	7.4	10.3	8.1	20.3	49.2	197.2
2000 [4]	12.4	173.4	2.4	*	2.6	4.3	7.8	9.9	11.0	21.6	50.5	198.9
1999 [4]	11.2	175.3	2.8	*	1.9	3.2	7.0	7.9	10.3	18.8	48.0	182.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25, G31-H93,K00-K22,K29-K31,K50-K66,K71-K72, K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0, N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)												
All origins [3]												
Both sexes												
2001	65.5	28.0	4.2	2.1	4.5	7.8	18.0	36.2	63.4	140.9	395.3	1,476.7
2000 [4]	63.4	29.0	4.0	2.1	4.5	7.8	18.0	35.4	62.4	136.0	388.5	1,445.7
1999 [4]	61.5	30.2	4.2	2.3	4.6	7.6	17.7	33.9	60.5	136.6	373.0	1,410.7
Male												
2001	53.8	31.4	4.6	2.2	5.2	8.6	21.3	43.7	71.2	153.4	397.4	1,271.6
2000 [4]	52.3	32.3	4.3	2.3	5.3	9.0	21.4	42.8	70.7	148.4	391.5	1,266.2
1999 [4]	51.0	35.0	4.5	2.3	5.3	8.8	21.6	40.5	69.0	148.7	384.2	1,225.4
Female												
2001	76.7	24.5	3.8	2.0	3.8	6.9	14.6	29.1	56.2	130.5	394.0	1,562.6
2000 [4]	74.0	25.6	3.6	2.0	3.7	6.5	14.6	28.2	54.8	125.8	386.5	1,518.8
1999 [4]	71.6	25.3	3.9	2.2	3.8	6.4	13.9	27.4	52.7	126.7	365.7	1,485.3
Hispanic												
Both sexes												
2001	21.2	22.9	3.7	1.7	3.7	5.2	12.3	26.3	49.7	107.2	277.5	991.0
2000 [4]	20.7	25.4	3.7	1.7	3.3	5.9	13.8	27.5	47.5	103.4	284.0	913.0
1999 [4]	21.5	26.5	3.7	1.6	3.5	6.5	14.0	28.2	51.4	109.3	263.5	1,039.7
Male												
2001	20.2	25.4	4.3	1.8	4.1	6.0	15.9	34.7	59.1	117.2	273.7	939.0
2000 [4]	20.2	25.3	3.6	1.8	3.9	7.0	18.2	35.6	57.7	119.9	273.9	870.7
1999 [4]	21.0	32.1	3.4	1.6	3.8	8.1	18.8	37.0	62.6	123.9	276.3	923.1

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25, G31-H93,K00-K22,K29-K31,K50-K66,K71-K72, K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0, N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)												
Hispanic												
Female												
2001	22.3	20.2	3.1	1.7	3.2	4.2	8.4	18.0	41.4	99.4	280.2	1,016.5
2000 [4]	21.4	25.5	3.8	1.5	2.5	4.5	9.0	19.6	38.5	90.4	290.9	933.8
1999 [4]	22.0	20.6	4.0	1.7	3.1	4.6	8.8	19.7	41.6	97.9	255.0	1,096.0
Non-Hispanic [5]												
Both sexes												
2001	71.8	29.0	4.3	2.2	4.6	8.3	18.6	37.0	64.2	142.7	399.8	1,491.3
2000 [4]	69.3	29.6	4.0	2.2	4.7	8.1	18.4	35.8	63.3	137.5	391.9	1,462.0
1999 [4]	66.8	30.8	4.3	2.4	4.8	7.8	18.1	34.1	60.9	137.8	376.6	1,420.1
Male												
2001	58.8	32.7	4.6	2.2	5.4	9.2	21.9	44.2	71.7	155.2	402.2	1,282.3
2000 [4]	56.9	33.6	4.4	2.3	5.5	9.4	21.7	43.0	71.3	149.4	395.5	1,279.8
1999 [4]	55.2	35.1	4.7	2.4	5.6	9.0	21.7	40.5	69.1	149.6	387.8	1,234.0
Female												
2001	84.2	25.2	4.0	2.1	3.9	7.4	15.4	30.0	57.3	132.2	398.3	1,578.3
2000 [4]	81.0	25.4	3.5	2.1	3.9	6.8	15.2	28.9	55.9	127.6	389.5	1,535.7
1999 [4]	77.9	26.2	3.9	2.3	4.0	6.7	14.4	28.0	53.3	128.2	369.3	1,494.7
Non-Hispanic white												
Both sexes												
2001	77.3	23.2	3.9	2.1	4.3	7.2	16.4	33.2	60.1	139.7	399.5	1,512.9
2000 [4]	74.3	24.0	3.4	2.1	4.5	7.0	16.2	32.0	59.0	133.9	390.1	1,481.3

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25, G31-H93,K00-K22,K29-K31,K50-K66,K71-K72, K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0, N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)												
Non-Hispanic white												
Both sexes												
1999 [4]	71.5	24.2	4.0	2.2	4.5	7.0	15.5	30.1	56.9	133.5	375.1	1,443.8
Male												
2001	61.9	25.3	4.4	2.2	5.3	8.2	19.8	40.0	66.9	150.8	399.4	1,293.1
2000 [4]	59.6	26.7	3.8	2.3	5.4	8.5	19.4	39.2	66.0	144.1	390.9	1,288.3
1999 [4]	57.6	27.4	4.5	2.2	5.4	8.4	19.4	36.2	64.5	144.2	382.8	1,239.8
Female												
2001	92.1	21.0	3.4	2.1	3.3	6.2	13.1	26.4	53.7	130.2	399.5	1,604.4
2000 [4]	88.3	21.2	2.9	1.8	3.5	5.5	12.9	25.0	52.5	125.2	389.6	1,559.1
1999 [4]	84.9	20.9	3.5	2.2	3.5	5.6	11.7	24.1	49.8	124.5	370.2	1,525.2
Non-Hispanic black												
Both sexes												
2001	61.1	53.8	5.6	2.5	6.7	14.7	34.8	71.7	109.1	198.5	476.3	1,431.5
2000 [4]	60.1	51.3	6.1	2.9	6.6	15.6	35.1	70.6	109.6	195.0	472.0	1,403.3
1999 [4]	58.0	58.4	5.5	3.1	6.9	13.5	36.2	69.3	104.9	202.1	444.8	1,295.3
Male												
2001	56.4	61.3	5.2	2.5	6.8	15.6	39.3	85.6	128.3	232.9	530.2	1,329.4
2000 [4]	56.1	61.4	6.4	2.6	7.1	16.3	39.6	82.4	133.0	231.8	527.9	1,340.9
1999 [4]	54.4	66.5	5.6	3.6	7.3	13.9	40.3	81.9	122.9	228.2	509.7	1,303.8
Female												
2001	65.2	46.0	6.0	2.6	6.6	13.9	30.8	59.7	93.6	174.3	446.1	1,469.3

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25, G31-H93,K00-K22,K29-K31,K50-K66,K71-K72, K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0, N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)												
Non-Hispanic black												
Female												
2000 [4]	63.7	40.9	5.8	3.2	6.2	15.0	31.0	60.5	90.8	169.1	440.6	1,426.5
1999 [4]	61.3	50.0	5.4	2.6	6.5	13.1	32.7	58.5	90.5	183.7	408.5	1,292.1
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
All origins [3]												
Both sexes												
2001	35.7	24.2	11.2	6.9	36.1	29.9	35.4	34.1	30.3	42.8	100.9	276.4
2000 [4]	34.8	23.1	11.9	7.3	36.0	29.5	34.1	32.6	30.9	41.9	95.1	273.5
1999 [4]	35.1	22.3	12.4	7.6	35.3	29.6	33.8	31.8	30.6	44.6	100.5	282.4
Male												
2001	47.2	26.2	13.3	8.6	52.9	45.4	51.8	50.6	43.0	57.3	133.4	358.5
2000 [4]	46.2	25.7	14.0	8.8	52.1	45.0	51.0	49.0	44.1	55.9	124.3	359.7
1999 [4]	46.4	25.4	14.4	9.4	50.7	45.0	50.2	47.9	44.0	60.2	132.4	375.4
Female												
2001	24.5	22.1	8.9	5.2	18.4	14.0	19.2	18.0	18.5	30.8	79.6	242.0
2000 [4]	23.8	20.5	9.7	5.6	19.1	13.7	17.5	16.8	18.8	30.3	76.1	238.4
1999 [4]	24.1	19.0	10.3	5.6	19.2	14.0	17.6	16.3	18.3	31.8	79.7	244.8
Hispanic												
Both sexes												
2001	25.8	18.8	10.3	5.4	32.6	28.2	32.3	31.7	30.5	40.0	73.8	166.3
2000 [4]	25.0	15.3	10.4	5.6	31.8	27.3	31.5	30.4	28.8	38.8	72.1	184.5

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
Hispanic												
Both sexes												
1999 [4]	25.5	17.7	11.9	5.6	30.8	28.4	32.6	31.2	28.5	42.5	69.6	182.4
Male												
2001	37.6	18.0	12.7	6.2	49.2	43.8	50.1	48.8	45.0	54.9	105.1	198.1
2000 [4]	36.9	17.0	11.6	6.1	48.2	42.9	49.6	48.4	43.2	57.2	93.9	251.9
1999 [4]	37.7	20.7	13.5	6.8	46.2	44.1	51.3	50.2	43.5	63.3	94.9	258.8
Female												
2001	13.2	19.7	7.8	4.5	13.3	10.1	13.0	14.9	17.7	28.2	52.6	150.8
2000 [4]	12.4	13.5	9.2	5.2	12.5	9.2	12.2	12.8	16.0	24.4	57.4	151.3
1999 [4]	12.6	14.6	10.3	4.2	12.8	10.1	12.6	12.8	15.2	26.2	52.7	145.6
Non-Hispanic [5]												
Both sexes												
2001	36.9	25.4	11.3	7.2	36.6	30.1	35.6	34.0	30.0	42.8	101.9	279.8
2000 [4]	36.0	25.0	12.2	7.5	36.7	29.8	34.2	32.5	30.9	41.9	95.8	276.1
1999 [4]	36.2	23.4	12.4	7.9	36.0	29.6	33.7	31.6	30.5	44.5	101.3	285.0
Male												
2001	48.5	28.0	13.4	9.0	53.5	45.6	51.7	50.4	42.5	57.1	134.3	364.5
2000 [4]	47.3	27.7	14.5	9.4	52.7	45.2	50.8	48.6	43.8	55.5	125.2	363.1
1999 [4]	47.4	26.6	14.5	9.9	51.4	44.8	49.7	47.3	43.5	59.7	133.2	378.7
Female												
2001	26.0	22.6	9.2	5.3	19.3	14.7	19.9	18.3	18.5	30.8	80.6	244.6
2000 [4]	25.2	22.2	9.7	5.6	20.3	14.5	18.1	17.0	18.9	30.6	76.7	240.9
1999 [4]	25.5	20.0	10.2	5.9	20.2	14.6	18.0	16.5	18.5	32.0	80.6	247.5

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
Non-Hispanic white												
Both sexes												
2001	38.5	20.9	10.9	6.6	39.0	31.1	35.9	32.8	29.0	42.4	103.7	289.3
2000 [4]	37.3	20.1	11.1	7.2	39.2	30.0	34.2	31.0	30.0	41.0	97.2	285.6
1999 [4]	37.2	19.0	11.1	7.4	38.1	29.9	33.5	30.0	28.9	43.3	101.7	293.3
Male												
2001	49.7	22.9	13.2	8.1	56.6	46.7	51.8	48.0	40.3	56.1	135.2	374.7
2000 [4]	48.1	22.9	13.4	8.9	55.9	45.2	50.4	45.8	41.6	53.6	125.2	373.8
1999 [4]	47.8	22.0	12.9	9.1	53.6	44.9	49.0	44.2	40.7	57.3	132.7	387.1
Female												
2001	27.8	18.7	8.5	5.0	20.7	15.2	19.9	17.9	18.3	30.7	82.8	253.8
2000 [4]	26.9	17.1	8.7	5.3	21.9	14.6	18.0	16.3	19.1	30.4	78.8	250.0
1999 [4]	27.0	15.9	9.3	5.6	22.0	14.7	17.9	15.9	17.7	31.5	81.5	255.9
Non-Hispanic black												
Both sexes												
2001	34.4	44.5	14.2	9.7	29.6	31.1	40.9	48.5	42.4	51.3	89.6	183.9
2000 [4]	34.4	46.5	16.8	9.6	29.6	33.5	40.6	47.8	41.3	53.0	85.3	183.2
1999 [4]	36.1	44.1	18.2	10.9	30.5	33.2	40.7	48.3	45.1	58.6	103.8	208.9
Male												
2001	49.5	50.4	15.8	13.1	44.5	48.8	61.3	78.6	68.2	74.9	138.3	256.4
2000 [4]	50.3	49.5	19.4	12.3	44.2	53.5	63.3	76.7	69.4	80.4	133.3	277.1
1999 [4]	52.3	48.5	21.8	14.3	46.4	53.0	63.0	79.4	71.8	89.6	151.0	318.5
Female												
2001	20.6	38.3	12.4	6.2	14.8	15.1	22.7	22.7	21.6	34.6	62.2	157.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
Non-Hispanic black												
Female												
2000 [4]	20.1	43.3	14.2	6.7	15.1	15.5	20.5	23.0	18.7	33.7	58.4	148.3
1999 [4]	21.4	39.6	14.5	7.4	14.8	15.4	21.1	21.8	23.7	36.8	77.4	167.9
Transport accidents (V01-V99,Y85)												
All origins [3]												
Both sexes												
2001	16.6	3.7	4.4	4.4	27.9	18.8	17.4	16.0	15.3	17.9	27.9	28.9
2000 [4]	16.6	4.6	4.5	4.7	28.1	18.6	17.0	15.8	16.0	18.0	27.0	31.5
1999 [4]	16.6	5.0	4.6	4.8	27.6	18.5	16.8	15.4	16.0	19.4	28.4	32.0
Male												
2001	23.5	3.3	5.0	5.3	39.6	28.3	25.1	23.3	21.6	24.3	39.8	52.7
2000 [4]	23.3	4.8	5.0	5.4	39.3	27.7	24.8	23.0	22.5	24.3	38.1	59.8
1999 [4]	23.1	5.2	5.1	5.8	38.1	27.2	24.0	22.4	22.5	26.1	40.7	63.5
Female												
2001	10.0	4.2	3.8	3.5	15.6	9.2	9.8	8.9	9.4	12.7	20.1	18.9
2000 [4]	10.2	4.3	4.0	3.9	16.4	9.3	9.3	8.9	10.0	12.8	19.8	19.9
1999 [4]	10.4	4.8	4.2	3.9	16.6	9.6	9.6	8.7	10.1	14.0	20.4	19.3
Hispanic												
Both sexes												
2001	15.6	5.3	5.3	3.8	26.0	18.8	17.3	16.4	17.9	20.2	30.1	23.8
2000 [4]	14.8	5.4	5.2	4.1	24.6	17.3	16.2	16.2	15.7	20.4	23.9	25.2
1999 [4]	14.4	5.3	5.5	3.8	23.1	17.8	15.4	15.0	15.5	22.0	25.0	25.3

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Transport accidents (V01-V99,Y85)												
Hispanic												
Male												
2001	22.8	*	6.0	4.3	38.4	28.8	25.9	23.2	24.6	28.2	51.0	43.8
2000 [4]	21.6	6.1	5.6	4.5	36.1	26.4	24.1	24.1	23.5	29.5	36.4	44.3
1999 [4]	20.8	5.7	5.8	4.5	33.3	26.7	23.2	22.6	21.0	31.6	40.1	*
Female												
2001	8.0	6.7	4.6	3.2	11.6	7.1	8.0	9.7	12.0	13.9	16.0	*
2000 [4]	7.6	*	4.8	3.7	11.0	6.8	7.8	8.5	8.7	13.3	15.5	*
1999 [4]	7.6	*	5.2	3.1	11.0	7.6	7.0	7.7	10.6	14.5	14.9	*
Non-Hispanic [5]												
Both sexes												
2001	16.7	3.4	4.2	4.6	28.2	18.8	17.3	15.8	15.0	17.7	27.7	29.0
2000 [4]	16.8	4.3	4.3	4.8	28.8	18.7	17.0	15.7	15.9	17.8	27.1	31.7
1999 [4]	16.9	4.9	4.4	5.0	28.4	18.5	16.9	15.4	16.0	19.2	28.3	32.2
Male												
2001	23.5	3.2	4.7	5.5	39.7	28.1	24.8	23.2	21.3	24.0	39.2	52.9
2000 [4]	23.4	4.5	4.8	5.6	39.8	27.8	24.7	22.8	22.2	23.9	38.0	60.4
1999 [4]	23.3	5.1	4.9	6.0	39.0	27.2	24.0	22.2	22.4	25.6	40.4	64.3
Female												
2001	10.2	3.6	3.6	3.6	16.3	9.5	10.0	8.8	9.1	12.5	20.2	19.1
2000 [4]	10.5	4.2	3.8	3.9	17.4	9.7	9.4	8.9	10.1	12.8	20.0	20.1
1999 [4]	10.7	4.7	3.9	4.0	17.5	10.0	9.9	8.8	10.0	13.9	20.5	19.3

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Transport accidents (V01-V99,Y85)												
Non-Hispanic white												
Both sexes												
2001	17.0	3.1	3.8	4.4	29.9	19.0	17.4	15.6	14.7	17.6	28.3	29.9
2000 [4]	17.1	3.5	3.8	4.8	30.9	18.5	17.0	15.4	15.8	17.6	27.6	32.4
1999 [4]	17.1	4.3	3.7	5.0	30.3	18.6	16.7	15.0	15.6	19.1	28.6	32.5
Male												
2001	23.8	2.9	4.4	5.2	41.9	28.2	24.6	22.7	20.8	23.7	39.6	54.3
2000 [4]	23.7	3.8	4.3	5.7	42.5	27.1	24.5	22.2	21.6	23.2	38.1	62.3
1999 [4]	23.5	4.9	4.0	5.9	41.2	26.9	23.6	21.6	21.7	25.2	40.3	64.6
Female												
2001	10.6	3.3	3.2	3.5	17.4	9.7	10.2	8.6	9.1	12.5	20.8	19.8
2000 [4]	10.9	3.2	3.4	4.0	18.9	9.7	9.5	8.6	10.4	12.8	20.7	20.4
1999 [4]	11.0	3.6	3.4	4.0	19.1	10.2	9.9	8.6	9.9	13.9	20.9	19.7
Non-Hispanic black												
Both sexes												
2001	16.3	*	5.4	5.5	22.8	20.5	19.2	19.1	18.4	18.8	21.4	17.8
2000 [4]	16.4	6.9	5.9	4.8	22.4	22.2	19.1	19.7	17.5	19.9	19.9	21.9
1999 [4]	16.7	7.7	7.1	5.6	22.7	20.6	19.5	19.2	18.5	19.5	26.0	27.8
Male												
2001	24.5	*	5.8	7.1	33.4	32.6	29.7	30.7	30.2	27.4	36.9	35.8
2000 [4]	24.8	7.2	6.4	5.8	32.5	35.6	30.5	30.1	29.6	32.6	36.4	45.8
1999 [4]	24.9	*	8.0	7.0	32.9	33.0	29.8	30.7	29.3	29.5	43.5	68.9
Female												
2001	8.9	*	5.0	3.8	12.3	9.6	9.9	9.2	9.0	12.8	12.6	11.1

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Transport accidents (V01-V99,Y85)												
Non-Hispanic black												
Female												
2000 [4]	8.8	*	5.4	3.7	12.5	10.1	8.9	10.8	7.8	11.0	10.6	13.1
1999 [4]	9.4	9.3	6.2	4.1	12.6	9.4	10.4	9.5	9.8	12.5	16.3	12.4
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1, V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)												
All origins [3]												
Both sexes												
2001	15.4	3.6	4.1	4.1	26.8	17.5	15.7	14.2	13.7	16.3	26.3	27.8
2000 [4]	15.4	4.4	4.2	4.3	26.9	17.3	15.3	14.2	14.4	16.5	25.7	30.4
1999 [4]	15.2	4.8	4.2	4.3	26.2	16.9	14.9	13.6	14.2	17.8	26.5	30.4
Male												
2001	21.5	3.2	4.5	4.9	38.0	26.2	22.4	20.5	19.0	21.5	37.3	51.1
2000 [4]	21.3	4.6	4.7	5.0	37.4	25.5	22.0	20.4	19.8	21.7	35.6	57.5
1999 [4]	20.9	5.0	4.7	5.1	36.0	24.7	21.1	19.4	19.3	23.3	38.0	59.8
Female												
2001	9.5	4.0	3.6	3.3	15.1	8.7	9.1	8.1	8.8	12.0	19.2	18.0
2000 [4]	9.7	4.2	3.8	3.6	15.9	8.8	8.8	8.2	9.5	12.3	19.2	19.3
1999 [4]	9.7	4.7	3.8	3.6	15.9	8.9	8.8	8.0	9.5	13.3	19.1	18.5
Hispanic												
Both sexes												
2001	14.3	4.9	4.7	3.5	25.1	17.4	15.3	13.8	15.1	17.7	26.3	20.0
2000 [4]	13.9	5.2	4.9	3.8	23.7	16.4	14.7	15.1	14.4	18.4	22.3	22.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1, V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)												
Hispanic												
Both sexes												
1999 [4]	13.3	5.3	5.2	3.6	21.9	16.4	13.9	13.7	14.5	19.7	23.5	23.2
Male												
2001	21.0	*	5.3	4.0	36.9	26.8	23.1	20.2	21.2	24.6	44.3	*
2000 [4]	20.1	6.1	5.2	4.1	34.7	24.9	21.6	22.0	21.2	26.0	33.4	*
1999 [4]	19.1	5.7	5.6	4.1	31.6	24.4	20.8	20.1	19.0	27.2	37.0	*
Female												
2001	7.2	6.2	4.1	3.0	11.2	6.5	6.9	7.6	9.7	12.3	14.2	*
2000 [4]	7.2	*	4.6	3.6	10.6	6.5	7.3	8.3	8.4	12.4	14.9	*
1999 [4]	7.2	*	4.7	3.0	10.6	7.1	6.5	7.4	10.6	13.8	14.6	*
Non-Hispanic [5]												
Both sexes												
2001	15.5	3.2	3.9	4.2	27.1	17.5	15.7	14.2	13.5	16.2	26.3	28.0
2000 [4]	15.6	4.2	4.1	4.4	27.5	17.4	15.4	14.1	14.4	16.3	25.8	30.6
1999 [4]	15.4	4.7	4.0	4.5	26.9	16.9	15.0	13.5	14.0	17.6	26.5	30.5
Male												
2001	21.5	3.1	4.3	5.1	38.1	25.9	22.2	20.4	18.8	21.3	36.9	51.6
2000 [4]	21.4	4.2	4.5	5.1	37.9	25.5	22.0	20.2	19.6	21.3	35.7	58.3
1999 [4]	21.0	4.8	4.4	5.2	36.7	24.6	21.1	19.2	19.1	22.9	37.7	60.7
Female												
2001	9.8	3.4	3.5	3.3	15.8	9.1	9.4	8.2	8.7	12.0	19.3	18.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1, V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)												
Non-Hispanic [5]												
Female												
2000 [4]	10.0	4.2	3.6	3.6	16.9	9.2	8.9	8.2	9.6	12.2	19.3	19.4
1999 [4]	10.0	4.7	3.6	3.7	16.8	9.2	9.1	8.1	9.3	13.2	19.2	18.5
Non-Hispanic white												
Both sexes												
2001	15.8	3.0	3.5	4.1	28.7	17.6	15.7	13.9	13.3	16.1	26.9	29.0
2000 [4]	15.8	3.3	3.7	4.4	29.5	17.0	15.3	13.7	14.2	16.1	26.2	31.3
1999 [4]	15.6	4.1	3.4	4.4	28.7	16.8	14.8	13.1	13.7	17.5	26.8	30.9
Male												
2001	21.7	2.9	3.9	4.8	40.1	25.9	21.9	19.9	18.2	21.0	37.3	53.1
2000 [4]	21.5	3.4	4.1	5.1	40.3	24.7	21.6	19.6	18.9	20.7	35.6	60.0
1999 [4]	21.1	4.7	3.6	5.1	38.8	24.2	20.6	18.5	18.5	22.6	37.6	61.2
Female												
2001	10.1	3.1	3.1	3.3	17.0	9.3	9.6	8.1	8.7	11.9	20.0	19.0
2000 [4]	10.3	3.2	3.2	3.6	18.4	9.3	9.0	7.9	9.8	12.3	20.0	19.8
1999 [4]	10.3	3.5	3.1	3.7	18.3	9.4	9.0	7.9	9.2	13.2	19.7	18.8
Non-Hispanic black												
Both sexes												
2001	15.2	*	5.3	5.0	22.2	19.6	17.6	17.2	16.6	17.4	19.9	16.9
2000 [4]	15.5	6.9	5.6	4.5	21.6	21.1	17.6	18.3	16.3	18.4	18.8	20.7
1999 [4]	15.5	7.6	6.5	5.1	21.7	19.2	17.9	17.4	16.7	18.1	24.0	25.5

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1, V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)												
Non-Hispanic black												
Male												
2001	22.7	*	5.6	6.5	32.4	31.0	26.9	27.5	26.8	24.2	34.8	35.8
2000 [4]	23.1	7.2	6.0	5.4	31.1	33.6	27.9	27.6	27.2	29.8	34.6	44.6
1999 [4]	22.8	*	7.0	6.2	31.5	30.7	26.9	27.5	25.8	26.8	41.3	61.8
Female												
2001	8.5	*	4.9	3.4	12.1	9.2	9.4	8.4	8.4	12.6	11.5	9.8
2000 [4]	8.5	*	5.2	3.6	12.3	9.8	8.5	10.2	7.5	10.4	9.9	11.8
1999 [4]	8.9	9.3	5.9	4.0	12.1	8.9	9.9	8.8	9.3	12.0	14.4	12.0
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11, V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9, V88.9,V89.1,V89.3,V89.9)												
All origins [3]												
Both sexes												
2001	0.5	*	0.2	0.2	0.5	0.4	0.6	0.6	0.5	0.6	0.8	0.7
2000 [4]	0.5	*	0.2	0.2	0.6	0.5	0.7	0.6	0.5	0.5	0.7	0.8
1999 [4]	0.8	*	0.3	0.3	0.9	0.8	0.9	0.8	0.8	0.8	1.3	1.3
Male												
2001	0.8	*	0.3	0.2	0.8	0.8	1.0	1.0	0.9	1.0	1.1	*
2000 [4]	0.8	*	*	0.3	1.0	0.8	1.1	0.9	0.8	0.9	1.2	*
1999 [4]	1.1	*	0.3	0.4	1.4	1.2	1.4	1.3	1.2	1.2	1.8	2.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11, V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9, V88.9,V89.1,V89.3,V89.9)												
All origins [3]												
Female												
2001	0.2	*	*	0.2	0.2	0.1	0.2	0.3	0.2	0.3	0.6	*
2000 [4]	0.3	*	*	0.2	0.3	0.2	0.3	0.4	0.2	0.2	0.5	*
1999 [4]	0.4	*	0.3	0.2	0.4	0.4	0.4	0.4	0.4	0.5	1.0	*
Hispanic												
Both sexes												
2001	0.5	*	*	*	0.6	0.6	0.8	0.9	*	*	*	*
2000 [4]	0.6	*	*	*	0.5	0.6	0.9	0.8	*	*	*	*
1999 [4]	0.8	*	*	*	0.9	1.0	1.2	1.0	*	2.0	*	*
Male												
2001	0.9	*	*	*	1.0	0.9	1.4	1.3	*	*	*	*
2000 [4]	0.9	*	*	*	0.9	0.9	1.4	1.4	*	*	*	*
1999 [4]	1.3	*	*	*	1.3	1.6	1.8	1.8	*	*	*	*
Female												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.5	*	*	0.2	0.5	0.4	0.6	0.6	0.5	0.7	0.7	0.6
2000 [4]	0.5	*	*	0.2	0.6	0.5	0.7	0.6	0.5	0.5	0.7	0.8

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11, V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9, V88.9,V89.1,V89.3,V89.9)												
Non-Hispanic [5]												
Both sexes												
1999 [4]	0.7	*	0.3	0.3	0.9	0.7	0.9	0.8	0.8	0.7	1.3	1.3
Male												
2001	0.7	*	*	0.3	0.7	0.7	1.0	0.9	0.8	1.0	1.0	*
2000 [4]	0.8	*	*	0.3	1.0	0.8	1.1	0.8	0.8	0.8	1.1	*
1999 [4]	1.1	*	0.4	0.4	1.4	1.1	1.4	1.2	1.2	1.0	1.8	3.0
Female												
2001	0.2	*	*	0.2	0.2	0.1	0.2	0.3	0.2	0.3	0.5	*
2000 [4]	0.3	*	*	0.2	0.3	0.2	0.3	0.4	0.2	0.2	0.4	*
1999 [4]	0.4	*	*	0.2	0.5	0.4	0.4	0.4	0.4	0.5	1.0	*
Non-Hispanic white												
Both sexes												
2001	0.5	*	*	0.2	0.5	0.4	0.5	0.6	0.4	0.6	0.7	0.5
2000 [4]	0.5	*	*	0.3	0.7	0.5	0.6	0.6	0.4	0.5	0.7	0.8
1999 [4]	0.8	*	0.3	0.3	1.0	0.7	0.9	0.8	0.8	0.7	1.2	1.3
Male												
2001	0.7	*	*	0.2	0.8	0.7	0.8	0.9	0.8	0.9	0.9	*
2000 [4]	0.8	*	*	0.3	1.0	0.8	1.0	0.8	0.7	0.7	1.2	*
1999 [4]	1.1	*	*	0.4	1.4	1.1	1.3	1.2	1.2	1.0	1.7	2.8
Female												
2001	0.2	*	*	*	0.2	*	0.2	0.2	*	0.4	0.5	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11, V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9, V88.9,V89.1,V89.3,V89.9)												
Non-Hispanic white												
Female												
2000 [4]	0.3	*	*	0.2	0.3	0.2	0.2	0.4	*	*	0.4	*
1999 [4]	0.4	*	*	0.2	0.5	0.4	0.4	0.4	0.4	0.4	0.9	*
Non-Hispanic black												
Both sexes												
2001	0.6	*	*	0.4	0.4	0.5	1.1	0.9	0.9	*	*	*
2000 [4]	0.6	*	*	*	0.5	0.6	1.0	0.8	*	*	*	*
1999 [4]	0.7	*	*	0.3	0.6	0.8	0.9	1.0	*	*	*	*
Male												
2001	0.9	*	*	*	*	*	1.8	1.4	*	*	*	*
2000 [4]	0.9	*	*	*	0.9	1.0	1.6	1.4	*	*	*	*
1999 [4]	1.2	*	*	*	1.0	1.4	1.6	1.8	*	*	*	*
Female												
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)												
All origins [3]												
Both sexes												
2001	0.7	*	0.2	0.1	0.5	0.9	1.0	1.1	1.0	1.0	0.8	0.5
2000 [4]	0.7	*	*	0.1	0.6	0.8	0.9	1.0	1.1	1.0	0.6	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)												
All origins [3]												
Both sexes												
1999 [4]	0.7	*	*	0.2	0.5	0.8	0.9	1.0	1.1	0.9	0.5	*
Male												
2001	1.2	*	*	0.2	0.9	1.4	1.6	1.9	1.8	1.7	1.4	*
2000 [4]	1.2	*	*	0.2	0.9	1.3	1.6	1.7	1.8	1.7	1.3	*
1999 [4]	1.1	*	*	0.3	0.8	1.4	1.4	1.7	2.0	1.6	0.9	*
Female												
2001	0.3	*	*	0.1	0.2	0.3	0.4	0.4	0.3	0.3	0.4	*
2000 [4]	0.2	*	*	*	0.2	0.2	0.2	0.3	0.3	0.3	*	*
1999 [4]	0.3	*	*	0.1	0.2	0.3	0.4	0.3	0.3	0.2	0.3	*
Hispanic												
Both sexes												
2001	0.8	*	*	*	0.4	0.8	1.2	1.7	2.1	2.0	*	*
2000 [4]	0.3	*	*	*	0.4	0.4	0.6	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	0.5	*	*	*	*	*	*
Male												
2001	0.9	*	*	*	*	1.1	1.4	1.7	*	*	*	*
2000 [4]	0.5	*	*	*	*	0.6	1.1	*	*	*	*	*
1999 [4]	0.4	*	*	*	*	0.7	*	*	*	*	*	*
Female												
2001	0.7	*	*	*	*	*	1.0	1.8	2.0	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)												
Non-Hispanic [5]												
Both sexes												
2001	0.7	*	*	0.1	0.6	0.9	1.0	1.1	1.0	0.9	0.7	*
2000 [4]	0.7	*	*	0.1	0.6	0.8	1.0	1.0	1.1	1.0	0.6	*
1999 [4]	0.7	*	*	0.2	0.6	0.9	1.0	1.1	1.1	0.9	0.6	*
Male												
2001	1.2	*	*	0.2	0.9	1.4	1.6	1.9	1.8	1.7	1.3	*
2000 [4]	1.2	*	*	0.2	1.0	1.5	1.7	1.8	1.9	1.7	1.3	*
1999 [4]	1.2	*	*	0.3	0.8	1.5	1.6	1.8	2.1	1.7	1.0	*
Female												
2001	0.2	*	*	*	0.2	0.3	0.3	0.3	0.2	0.2	0.4	*
2000 [4]	0.2	*	*	*	0.2	0.2	0.3	0.3	0.4	0.3	*	*
1999 [4]	0.3	*	*	0.1	0.3	0.3	0.4	0.4	0.3	0.3	0.3	*
Non-Hispanic white												
Both sexes												
2001	0.8	*	*	0.1	0.7	1.0	1.1	1.1	1.0	0.9	0.7	*
2000 [4]	0.8	*	*	0.2	0.7	1.0	1.1	1.1	1.2	1.0	0.7	*
1999 [4]	0.8	*	*	0.2	0.6	1.0	1.1	1.1	1.1	0.9	0.5	*
Male												
2001	1.3	*	*	0.2	1.1	1.6	1.8	1.9	1.8	1.7	1.4	*
2000 [4]	1.4	*	*	0.2	1.1	1.7	1.8	1.9	2.0	1.8	1.3	*
1999 [4]	1.3	*	*	0.3	1.0	1.6	1.7	1.9	2.0	1.7	1.0	*
Female												
2001	0.3	*	*	*	0.3	0.3	0.4	0.3	0.2	0.2	0.3	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)												
Non-Hispanic white												
Female												
2000 [4]	0.3	*	*	*	0.3	0.3	0.3	0.3	0.4	0.3	*	*
1999 [4]	0.3	*	*	*	0.3	0.4	0.5	0.4	0.3	0.3	*	*
Non-Hispanic black												
Both sexes												
2001	0.5	*	*	*	*	0.5	0.5	1.0	0.9	*	*	*
2000 [4]	0.4	*	*	*	*	0.5	0.5	0.6	*	*	*	*
1999 [4]	0.5	*	*	*	0.4	0.5	0.7	0.8	1.1	*	*	*
Male												
2001	0.8	*	*	*	*	0.9	1.0	1.8	2.1	*	*	*
2000 [4]	0.7	*	*	*	*	0.9	1.0	1.1	*	*	*	*
1999 [4]	0.9	*	*	*	*	0.9	1.2	1.4	2.3	*	*	*
Female												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
Nontransport accidents (W00-X59,Y86)												
All origins [3]												
Both sexes												
2001	19.0	20.5	6.8	2.5	8.2	11.1	18.0	18.1	15.0	24.9	73.0	247.5
2000 [4]	18.2	18.6	7.4	2.6	7.9	11.0	17.2	16.8	14.9	23.8	68.1	242.0
1999 [4]	18.4	17.3	7.7	2.7	7.7	11.1	17.0	16.4	14.6	25.1	72.1	250.3

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)												
All origins [3]												
Male												
2001	23.8	22.9	8.3	3.3	13.3	17.1	26.8	27.3	21.4	33.0	93.6	305.8
2000 [4]	22.9	20.9	9.0	3.4	12.8	17.3	26.2	26.0	21.6	31.5	86.2	299.8
1999 [4]	23.3	20.2	9.3	3.7	12.5	17.7	26.2	25.6	21.5	34.1	91.7	311.9
Female												
2001	14.5	17.9	5.1	1.6	2.8	4.9	9.4	9.2	9.1	18.1	59.5	223.1
2000 [4]	13.6	16.2	5.7	1.7	2.7	4.5	8.2	7.9	8.8	17.5	56.3	218.4
1999 [4]	13.7	14.2	6.1	1.7	2.6	4.4	8.0	7.6	8.2	17.8	59.4	225.5
Hispanic												
Both sexes												
2001	10.1	13.6	5.0	1.6	6.6	9.4	15.0	15.3	12.6	19.8	43.6	142.6
2000 [4]	10.2	9.9	5.2	1.5	7.3	10.0	15.3	14.2	13.1	18.4	48.2	159.2
1999 [4]	11.1	12.4	6.4	1.7	7.7	10.5	17.2	16.2	13.0	20.5	44.6	157.2
Male												
2001	14.8	14.2	6.7	1.9	10.8	14.9	24.3	25.6	20.4	26.8	54.1	154.3
2000 [4]	15.3	10.9	5.9	1.6	12.1	16.5	25.6	24.3	19.7	27.6	57.5	207.6
1999 [4]	16.9	15.0	7.7	2.3	12.9	17.4	28.2	27.6	22.5	31.6	54.7	217.8
Female												
2001	5.2	13.0	3.3	1.3	1.7	2.9	5.0	5.2	5.7	14.3	36.6	136.8
2000 [4]	4.9	8.8	4.4	1.5	1.6	2.4	4.4	4.3	7.3	11.1	41.9	135.5
1999 [4]	5.0	9.8	5.2	1.1	1.7	2.6	5.5	5.1	4.6	11.7	37.8	127.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)												
Non-Hispanic [5]												
Both sexes												
2001	20.2	22.0	7.2	2.6	8.5	11.3	18.3	18.2	15.0	25.1	74.2	250.8
2000 [4]	19.2	20.7	7.9	2.8	8.0	11.1	17.3	16.8	14.9	24.1	68.7	244.4
1999 [4]	19.3	18.5	8.0	2.9	7.6	11.1	16.8	16.2	14.5	25.3	72.9	252.8
Male												
2001	25.0	24.9	8.7	3.5	13.7	17.5	26.9	27.2	21.2	33.2	95.1	311.6
2000 [4]	23.9	23.2	9.7	3.8	12.9	17.4	26.1	25.8	21.5	31.6	87.2	302.7
1999 [4]	24.0	21.5	9.6	3.9	12.4	17.6	25.7	25.0	21.1	34.1	92.8	314.3
Female												
2001	15.7	19.1	5.6	1.7	3.0	5.2	9.9	9.5	9.4	18.3	60.4	225.5
2000 [4]	14.7	18.0	6.0	1.7	2.9	4.8	8.6	8.1	8.8	17.8	56.7	220.8
1999 [4]	14.8	15.2	6.3	1.9	2.7	4.6	8.2	7.7	8.4	18.1	60.1	228.2
Non-Hispanic white												
Both sexes												
2001	21.5	17.8	7.1	2.2	9.1	12.1	18.5	17.2	14.2	24.7	75.4	259.4
2000 [4]	20.1	16.6	7.3	2.3	8.3	11.5	17.2	15.6	14.1	23.4	69.6	253.2
1999 [4]	20.0	14.7	7.4	2.4	7.8	11.3	16.7	14.9	13.3	24.2	73.2	260.8
Male												
2001	25.9	20.0	8.8	2.9	14.7	18.6	27.2	25.3	19.6	32.5	95.6	320.4
2000 [4]	24.5	19.1	9.1	3.2	13.5	18.1	26.0	23.6	19.9	30.4	87.2	311.5
1999 [4]	24.3	17.1	8.9	3.2	12.5	17.9	25.4	22.7	19.1	32.0	92.4	322.6
Female												
2001	17.2	15.5	5.3	1.5	3.3	5.5	9.8	9.3	9.2	18.1	62.0	234.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)												
Non-Hispanic white												
Female												
2000 [4]	16.0	14.0	5.4	1.3	3.0	4.9	8.4	7.7	8.7	17.6	58.1	229.7
1999 [4]	16.0	12.3	5.9	1.5	2.9	4.6	8.0	7.3	7.9	17.6	60.6	236.2
Non-Hispanic black												
Both sexes												
2001	18.1	41.6	8.7	4.3	6.8	10.5	21.7	29.4	23.9	32.5	68.2	166.1
2000 [4]	18.0	39.5	10.9	4.8	7.1	11.3	21.5	28.1	23.8	33.1	65.4	161.2
1999 [4]	19.4	36.4	11.1	5.3	7.8	12.7	21.2	29.1	26.6	39.1	77.8	181.1
Male												
2001	25.0	48.2	10.0	6.0	11.0	16.2	31.6	47.9	38.0	47.5	101.4	220.6
2000 [4]	25.5	42.3	12.9	6.5	11.7	17.9	32.7	46.6	39.8	47.8	96.9	231.3
1999 [4]	27.5	42.3	13.7	7.3	13.5	20.0	33.3	48.7	42.5	60.1	107.6	249.6
Female												
2001	11.8	34.6	7.4	2.4	2.5	5.4	12.9	13.5	12.6	21.9	49.5	145.9
2000 [4]	11.3	36.6	8.8	3.0	2.6	5.4	11.6	12.2	10.9	22.8	47.8	135.2
1999 [4]	12.0	30.3	8.3	3.3	2.1	6.1	10.7	12.3	13.8	24.3	61.1	155.5
Falls (W00-W19)												
All origins [3]												
Both sexes												
2001	5.3	0.6	0.2	0.2	0.6	0.9	1.4	2.6	4.0	10.0	35.3	121.5
2000 [4]	4.7	*	0.2	0.1	0.6	0.8	1.3	2.3	3.9	9.0	31.1	112.6
1999 [4]	4.7	*	0.4	0.1	0.6	0.9	1.4	2.3	3.7	9.0	31.5	110.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Falls (W00-W19)												
All origins [3]												
Male												
2001	5.8	*	0.3	0.2	1.1	1.5	2.4	4.3	5.9	13.1	45.7	155.7
2000 [4]	5.2	*	0.3	0.1	1.0	1.4	2.2	3.7	5.8	11.9	39.9	145.4
1999 [4]	5.2	*	0.5	0.2	1.1	1.5	2.4	3.7	5.7	12.4	40.5	142.5
Female												
2001	4.8	*	*	0.1	0.1	0.2	0.5	1.0	2.2	7.4	28.5	107.2
2000 [4]	4.3	*	*	*	0.2	0.2	0.5	0.9	2.1	6.6	25.3	99.2
1999 [4]	4.3	*	*	*	0.2	0.2	0.5	0.9	1.9	6.3	25.7	97.2
Hispanic												
Both sexes												
2001	2.0	*	*	*	0.7	1.1	1.8	2.6	4.2	9.1	23.6	70.0
2000 [4]	2.0	*	*	*	0.7	1.1	1.3	2.5	4.3	8.0	26.5	81.0
1999 [4]	1.9	*	*	*	0.7	1.1	1.4	2.2	3.8	8.1	24.2	86.3
Male												
2001	2.8	*	*	*	1.2	1.9	3.3	4.5	7.1	11.4	27.7	72.4
2000 [4]	2.7	*	*	*	1.2	2.0	2.3	4.4	6.6	12.7	30.5	104.8
1999 [4]	2.7	*	*	*	1.3	1.9	2.4	3.8	7.2	11.4	31.3	120.8
Female												
2001	1.3	*	*	*	*	*	*	*	*	7.2	20.9	68.9
2000 [4]	1.2	*	*	*	*	*	*	*	2.3	4.3	23.8	69.2
1999 [4]	1.1	*	*	*	*	*	*	*	*	5.5	19.4	69.7

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Falls (W00-W19)												
Non-Hispanic [5]												
Both sexes												
2001	5.7	*	0.2	0.1	0.6	0.8	1.4	2.6	3.9	10.0	35.8	123.1
2000 [4]	5.1	*	0.2	0.1	0.6	0.7	1.3	2.3	3.9	9.1	31.2	113.5
1999 [4]	5.1	*	0.4	0.1	0.6	0.8	1.4	2.2	3.7	9.0	31.7	110.7
Male												
2001	6.2	*	0.3	0.2	1.1	1.4	2.3	4.2	5.8	13.1	46.4	159.0
2000 [4]	5.5	*	*	0.2	1.0	1.2	2.2	3.6	5.7	11.9	40.2	146.5
1999 [4]	5.5	*	0.5	0.2	1.0	1.3	2.3	3.6	5.5	12.4	40.6	142.9
Female												
2001	5.3	*	*	*	0.1	0.3	0.5	1.0	2.2	7.4	28.8	108.1
2000 [4]	4.7	*	*	*	0.2	0.2	0.5	1.0	2.1	6.8	25.3	100.1
1999 [4]	4.6	*	*	*	0.2	0.3	0.5	0.9	2.0	6.3	25.9	97.8
Non-Hispanic white												
Both sexes												
2001	6.6	*	0.2	0.2	0.8	0.9	1.4	2.6	3.9	10.3	37.6	129.9
2000 [4]	5.8	*	*	0.1	0.7	0.7	1.4	2.3	3.8	9.2	32.7	120.2
1999 [4]	5.7	*	0.4	0.2	0.7	0.8	1.5	2.1	3.5	9.1	33.1	116.6
Male												
2001	7.0	*	*	0.2	1.3	1.5	2.3	4.1	5.7	13.5	48.2	166.0
2000 [4]	6.2	*	*	0.2	1.1	1.3	2.2	3.6	5.6	12.0	41.7	154.1
1999 [4]	6.1	*	0.5	0.2	1.1	1.4	2.3	3.4	5.1	12.2	42.0	149.7
Female												
2001	6.2	*	*	*	0.2	0.3	0.5	1.1	2.2	7.7	30.6	114.8

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Falls (W00-W19)												
Non-Hispanic white												
Female												
2000 [4]	5.5	*	*	*	0.2	0.2	0.5	1.0	2.1	6.9	26.8	106.5
1999 [4]	5.4	*	*	*	0.2	0.3	0.6	0.8	1.9	6.4	27.2	103.4
Non-Hispanic black												
Both sexes												
2001	2.3	*	*	*	*	0.5	1.6	3.0	4.4	8.8	18.7	44.0
2000 [4]	2.1	*	*	*	*	0.4	1.3	2.7	4.8	8.3	15.2	39.1
1999 [4]	2.5	*	*	*	0.4	0.8	1.3	3.5	5.5	9.3	17.2	45.3
Male												
2001	3.2	*	*	*	*	*	2.7	5.4	6.9	12.6	29.1	68.1
2000 [4]	2.8	*	*	*	*	*	2.1	4.6	8.2	11.4	21.6	61.1
1999 [4]	3.4	*	*	*	*	1.4	2.5	5.9	8.7	15.0	24.1	58.2
Female												
2001	1.6	*	*	*	*	*	*	*	2.3	6.1	12.8	35.1
2000 [4]	1.5	*	*	*	*	*	*	1.0	2.0	6.1	11.6	31.0
1999 [4]	1.7	*	*	*	*	*	*	1.4	2.9	5.3	13.3	40.4
Accidental discharge of firearms (W32-W34)												
All origins [3]												
Both sexes												
2001	0.3	*	*	0.1	0.5	0.3	0.3	0.3	0.3	0.2	0.3	*
2000 [4]	0.3	*	*	0.2	0.5	0.3	0.3	0.2	0.2	0.2	0.2	*
1999 [4]	0.3	*	*	0.2	0.6	0.4	0.3	0.3	0.2	0.2	0.2	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental discharge of firearms (W32-W34)												
All origins [3]												
Male												
2001	0.5	*	*	0.2	0.9	0.5	0.5	0.5	0.5	0.3	0.7	*
2000 [4]	0.5	*	*	0.3	0.9	0.5	0.6	0.4	0.4	0.4	0.5	*
1999 [4]	0.5	*	*	0.3	1.2	0.6	0.5	0.4	0.4	0.4	0.4	*
Female												
2001	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	0.1	0.1	*	*	*	*	*
1999 [4]	0.1	*	*	*	0.1	0.1	0.1	*	*	*	*	*
Hispanic												
Both sexes												
2001	0.2	*	*	*	0.5	0.3	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	0.4	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	0.5	*	*	*	*	*	*	*
Male												
2001	0.4	*	*	*	0.9	*	*	*	*	*	*	*
2000 [4]	0.3	*	*	*	0.7	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	0.9	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental discharge of firearms (W32-W34)												
Non-Hispanic [5]												
Both sexes												
2001	0.3	*	*	0.2	0.5	0.3	0.3	0.3	0.3	0.2	0.3	*
2000 [4]	0.3	*	*	0.2	0.5	0.4	0.4	0.2	0.2	0.2	0.2	*
1999 [4]	0.3	*	*	0.2	0.7	0.4	0.3	0.3	0.2	0.2	0.2	*
Male												
2001	0.5	*	*	0.3	0.9	0.5	0.6	0.5	0.5	0.3	0.7	*
2000 [4]	0.5	*	*	0.3	1.0	0.6	0.6	0.4	0.4	0.4	0.5	*
1999 [4]	0.5	*	*	0.3	1.2	0.6	0.5	0.5	0.4	0.4	0.4	*
Female												
2001	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	0.1	0.1	*	*	*	*	*
1999 [4]	0.1	*	*	*	0.1	0.1	0.1	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2001	0.3	*	*	0.2	0.5	0.3	0.4	0.3	0.3	0.2	0.3	*
2000 [4]	0.3	*	*	0.2	0.5	0.3	0.4	0.2	0.2	0.2	0.2	*
1999 [4]	0.3	*	*	0.2	0.6	0.4	0.3	0.3	0.2	0.2	0.2	*
Male												
2001	0.5	*	*	0.3	0.8	0.6	0.6	0.5	0.5	0.3	0.7	*
2000 [4]	0.5	*	*	0.3	0.8	0.5	0.6	0.4	0.4	0.4	0.5	*
1999 [4]	0.5	*	*	0.3	1.0	0.6	0.5	0.5	0.4	0.4	*	*
Female												
2001	0.1	*	*	*	*	*	0.1	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental discharge of firearms (W32-W34)												
Non-Hispanic white												
Female												
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	0.1	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2001	0.3	*	*	*	0.8	*	*	*	*	*	*	*
2000 [4]	0.4	*	*	*	0.9	0.6	*	*	*	*	*	*
1999 [4]	0.4	*	*	*	1.2	0.4	*	*	*	*	*	*
Male												
2001	0.6	*	*	*	1.6	*	*	*	*	*	*	*
2000 [4]	0.7	*	*	*	1.6	1.2	*	*	*	*	*	*
1999 [4]	0.8	*	*	*	2.2	0.9	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Accidental drowning and submersion (W65-W74)												
All origins [3]												
Both sexes												
2001	1.2	1.7	3.0	0.8	1.5	0.9	1.0	0.9	0.8	0.9	1.2	1.7
2000 [4]	1.2	2.0	3.2	0.9	1.6	1.1	1.1	0.9	0.9	1.0	1.3	1.5
1999 [4]	1.3	1.8	3.2	0.9	1.7	1.1	1.1	1.0	0.9	1.0	1.3	1.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental drowning and submersion (W65-W74)												
All origins [3]												
Male												
2001	1.8	1.5	3.7	1.1	2.7	1.6	1.7	1.6	1.3	1.5	2.0	3.5
2000 [4]	2.0	2.2	4.0	1.3	2.9	1.8	1.8	1.6	1.5	1.6	1.8	2.8
1999 [4]	2.0	2.0	4.0	1.3	3.0	2.0	1.8	1.7	1.5	1.6	2.3	2.3
Female												
2001	0.5	1.9	2.2	0.5	0.3	0.2	0.4	0.3	0.4	0.3	0.7	1.0
2000 [4]	0.5	1.7	2.4	0.5	0.3	0.3	0.3	0.3	0.3	0.5	0.9	1.0
1999 [4]	0.5	1.6	2.4	0.5	0.3	0.2	0.3	0.3	0.4	0.6	0.6	1.4
Hispanic												
Both sexes												
2001	1.1	*	2.6	0.6	1.5	1.1	1.0	0.7	*	*	*	*
2000 [4]	1.2	*	2.5	0.6	2.2	1.0	0.8	0.6	*	*	*	*
1999 [4]	1.4	*	2.7	0.7	2.3	1.3	0.9	0.8	*	*	*	*
Male												
2001	1.8	*	3.5	0.8	2.7	1.8	1.7	1.2	*	*	*	*
2000 [4]	1.9	*	2.9	0.6	3.8	1.7	1.4	*	*	*	*	*
1999 [4]	2.3	*	2.9	1.1	4.1	2.2	1.5	1.6	*	*	*	*
Female												
2001	0.4	*	1.7	*	*	*	*	*	*	*	*	*
2000 [4]	0.5	*	2.2	0.6	*	*	*	*	*	*	*	*
1999 [4]	0.5	*	2.5	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental drowning and submersion (W65-W74)												
Non-Hispanic [5]												
Both sexes												
2001	1.1	1.6	3.1	0.9	1.5	0.9	1.0	0.9	0.8	0.9	1.3	1.7
2000 [4]	1.2	2.1	3.4	1.0	1.5	1.1	1.1	1.0	0.9	1.0	1.3	1.5
1999 [4]	1.2	1.7	3.3	0.9	1.5	1.1	1.1	1.0	0.9	1.0	1.3	1.7
Male												
2001	1.8	1.3	3.8	1.2	2.6	1.6	1.7	1.6	1.3	1.5	2.1	3.4
2000 [4]	2.0	2.4	4.3	1.4	2.7	1.8	1.9	1.6	1.6	1.6	1.8	2.8
1999 [4]	2.0	1.7	4.2	1.4	2.7	1.9	1.8	1.7	1.5	1.6	2.3	2.4
Female												
2001	0.5	1.9	2.4	0.5	0.3	0.2	0.4	0.3	0.4	0.4	0.7	1.0
2000 [4]	0.5	1.9	2.4	0.5	0.3	0.3	0.3	0.3	0.3	0.5	0.9	1.0
1999 [4]	0.5	1.7	2.4	0.5	0.3	0.3	0.3	0.3	0.4	0.6	0.6	1.4
Non-Hispanic white												
Both sexes												
2001	1.1	1.5	3.5	0.5	1.3	0.8	1.0	0.9	0.8	0.9	1.2	1.7
2000 [4]	1.1	1.7	3.7	0.7	1.3	1.0	1.1	0.9	0.9	1.0	1.3	1.4
1999 [4]	1.1	1.3	3.6	0.5	1.3	1.0	1.1	1.0	0.9	1.0	1.2	1.6
Male												
2001	1.6	*	4.3	0.7	2.2	1.4	1.5	1.5	1.2	1.5	2.1	3.5
2000 [4]	1.8	1.9	4.7	1.0	2.2	1.7	1.8	1.5	1.6	1.5	1.8	2.7
1999 [4]	1.8	*	4.5	0.7	2.2	1.6	1.8	1.7	1.5	1.4	2.2	2.4
Female												
2001	0.5	1.7	2.6	0.4	0.3	0.2	0.4	0.3	0.4	0.3	0.7	1.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental drowning and submersion (W65-W74)												
Non-Hispanic white												
Female												
2000 [4]	0.5	*	2.6	0.4	0.3	0.3	0.4	0.3	0.3	0.5	0.9	0.9
1999 [4]	0.5	*	2.6	0.3	0.3	0.3	0.4	0.4	0.4	0.6	0.6	1.3
Non-Hispanic black												
Both sexes												
2001	1.6	*	2.0	2.1	2.3	1.2	1.3	1.2	1.3	*	*	*
2000 [4]	1.7	4.3	2.3	2.1	2.3	1.1	1.4	1.3	*	*	*	*
1999 [4]	1.8	3.7	2.5	2.5	2.5	1.6	1.1	0.9	1.2	1.3	*	*
Male												
2001	2.8	*	2.4	3.1	4.4	2.5	2.2	2.1	2.4	*	*	*
2000 [4]	2.8	*	2.8	3.1	4.4	2.1	2.6	2.4	*	*	*	*
1999 [4]	3.2	*	3.4	3.8	4.6	3.2	2.1	1.9	2.1	*	*	*
Female												
2001	0.5	*	*	1.0	*	*	*	*	*	*	*	*
2000 [4]	0.6	*	1.8	1.0	*	*	*	*	*	*	*	*
1999 [4]	0.6	*	*	1.1	*	*	*	*	*	*	*	*
Accidental exposure to smoke, fire and flames (X00-X09)												
All origins [3]												
Both sexes												
2001	1.2	1.2	1.5	0.6	0.5	0.6	1.0	1.1	1.5	2.2	3.4	5.7
2000 [4]	1.2	1.0	1.9	0.6	0.5	0.6	0.9	1.2	1.5	2.2	3.8	6.2
1999 [4]	1.2	1.1	2.0	0.6	0.5	0.7	0.9	1.0	1.3	2.3	4.0	5.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)												
All origins [3]												
Male												
2001	1.4	1.5	1.8	0.7	0.7	0.8	1.4	1.5	1.9	2.8	4.1	8.3
2000 [4]	1.5	1.0	2.2	0.7	0.6	0.8	1.2	1.6	2.1	2.8	5.0	9.4
1999 [4]	1.5	1.1	2.2	0.7	0.7	1.0	1.3	1.4	1.7	2.9	5.1	9.3
Female												
2001	0.9	1.0	1.1	0.5	0.4	0.4	0.6	0.7	1.1	1.8	2.9	4.6
2000 [4]	0.9	*	1.5	0.6	0.4	0.4	0.6	0.7	1.0	1.7	3.0	4.9
1999 [4]	0.9	1.1	1.7	0.6	0.4	0.4	0.5	0.6	1.0	1.7	3.2	4.5
Hispanic												
Both sexes												
2001	0.5	*	*	0.3	*	0.3	0.5	*	*	*	*	*
2000 [4]	0.5	*	0.7	*	0.3	*	0.5	*	*	*	*	*
1999 [4]	0.6	*	1.4	0.4	0.4	*	0.4	*	*	*	*	*
Male												
2001	0.6	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.6	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.7	*	1.6	*	*	*	*	*	*	*	*	*
Female												
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.4	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.4	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)												
Non-Hispanic [5]												
Both sexes												
2001	1.3	1.3	1.7	0.7	0.6	0.7	1.0	1.1	1.5	2.3	3.4	5.7
2000 [4]	1.3	1.0	2.2	0.7	0.5	0.7	0.9	1.2	1.5	2.2	3.8	6.3
1999 [4]	1.3	1.2	2.1	0.7	0.5	0.8	1.0	1.0	1.4	2.3	4.0	6.0
Male												
2001	1.6	1.5	2.0	0.8	0.8	0.9	1.5	1.6	2.0	2.8	4.2	8.3
2000 [4]	1.6	*	2.6	0.8	0.7	0.9	1.3	1.7	2.1	2.8	5.0	9.3
1999 [4]	1.6	*	2.3	0.8	0.7	1.1	1.4	1.4	1.7	3.0	5.1	9.2
Female												
2001	1.0	*	1.3	0.5	0.4	0.5	0.6	0.7	1.1	1.9	2.9	4.6
2000 [4]	1.0	*	1.7	0.7	0.4	0.5	0.6	0.7	1.0	1.7	3.1	5.0
1999 [4]	1.0	*	1.8	0.6	0.4	0.5	0.6	0.6	1.1	1.8	3.3	4.7
Non-Hispanic white												
Both sexes												
2001	1.1	1.1	1.3	0.5	0.5	0.6	0.9	1.0	1.3	2.0	2.9	4.7
2000 [4]	1.1	*	1.5	0.5	0.5	0.6	0.8	1.1	1.3	1.8	3.1	5.1
1999 [4]	1.1	0.9	1.5	0.5	0.5	0.8	0.9	0.9	1.2	1.9	3.2	4.7
Male												
2001	1.4	*	1.7	0.6	0.7	0.8	1.3	1.3	1.7	2.3	3.5	6.9
2000 [4]	1.4	*	1.8	0.6	0.6	0.8	1.1	1.6	1.7	2.3	4.1	8.1
1999 [4]	1.4	*	1.7	0.6	0.7	1.1	1.3	1.2	1.5	2.3	4.2	7.8
Female												
2001	0.9	*	0.9	0.4	0.3	0.4	0.6	0.6	1.0	1.7	2.5	3.8

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)												
Non-Hispanic white												
Female												
2000 [4]	0.9	*	1.1	0.5	0.3	0.4	0.5	0.7	0.8	1.5	2.4	3.9
1999 [4]	0.8	*	1.3	0.5	0.3	0.4	0.5	0.6	0.8	1.5	2.6	3.5
Non-Hispanic black												
Both sexes												
2001	2.3	*	3.4	1.3	0.7	1.2	1.8	2.4	3.6	6.1	9.9	17.8
2000 [4]	2.6	*	5.1	1.6	0.7	1.2	1.7	2.4	4.2	6.1	13.0	20.7
1999 [4]	2.6	*	4.7	1.6	0.7	1.0	1.8	2.2	3.4	7.0	13.5	22.0
Male												
2001	3.0	*	3.7	1.6	0.8	1.4	2.8	3.9	5.1	9.0	13.2	26.6
2000 [4]	3.2	*	5.8	1.7	0.9	1.6	2.4	3.5	6.3	9.0	17.9	28.2
1999 [4]	3.1	*	5.3	1.8	0.9	1.2	2.7	3.3	4.3	10.4	18.4	29.7
Female												
2001	1.8	*	3.1	1.0	*	1.1	1.0	1.1	2.4	4.0	8.1	14.5
2000 [4]	2.0	*	4.3	1.4	*	0.9	1.2	1.4	2.5	4.0	10.2	17.9
1999 [4]	2.1	*	4.1	1.4	*	0.8	1.1	1.3	2.8	4.7	10.7	19.1
Accidental poisoning and exposure to noxious substances (X40-X49)												
All origins [3]												
Both sexes												
2001	4.9	*	0.2	0.1	3.4	6.3	11.2	9.1	3.2	1.6	2.2	3.5
2000 [4]	4.5	*	0.2	0.1	3.0	6.0	10.3	8.1	2.8	1.5	2.0	4.3
1999 [4]	4.4	*	0.2	0.1	2.5	5.9	10.1	7.8	2.8	1.6	2.2	3.4

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)												
All origins [3]												
Male												
2001	7.1	*	0.3	0.1	5.3	9.3	15.7	13.0	4.1	2.0	2.7	3.8
2000 [4]	6.6	*	0.3	0.1	4.6	9.0	15.1	12.0	3.4	1.8	2.1	6.2
1999 [4]	6.5	*	*	0.1	3.8	9.0	14.9	11.7	3.9	2.1	2.4	4.4
Female												
2001	2.9	*	*	0.1	1.4	3.3	6.7	5.2	2.3	1.2	1.9	3.3
2000 [4]	2.5	*	*	*	1.3	2.8	5.6	4.4	2.3	1.3	1.9	3.6
1999 [4]	2.3	*	*	*	1.1	2.6	5.3	4.0	1.8	1.3	2.0	3.0
Hispanic												
Both sexes												
2001	3.6	*	*	*	2.2	4.9	8.7	7.7	3.5	*	*	*
2000 [4]	3.7	*	*	*	2.0	5.3	9.6	8.0	3.1	*	*	*
1999 [4]	4.4	*	*	*	2.3	6.1	11.9	9.8	3.5	*	*	*
Male												
2001	5.6	*	*	*	3.1	7.4	13.6	12.9	5.8	*	*	*
2000 [4]	6.0	*	*	*	3.2	8.4	15.6	13.6	4.4	*	*	*
1999 [4]	7.3	*	*	*	3.6	9.8	19.4	16.4	6.6	*	*	*
Female												
2001	1.5	*	*	*	1.1	1.9	3.4	2.6	*	*	*	*
2000 [4]	1.3	*	*	*	*	1.6	3.1	2.5	*	*	*	*
1999 [4]	1.4	*	*	*	0.8	1.7	3.9	3.3	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)												
Non-Hispanic [5]												
Both sexes												
2001	5.1	*	0.2	0.1	3.6	6.6	11.4	9.1	3.1	1.6	2.3	3.6
2000 [4]	4.6	*	0.2	0.1	3.1	6.0	10.3	8.0	2.8	1.5	2.0	4.4
1999 [4]	4.3	*	0.2	0.1	2.5	5.8	9.7	7.5	2.7	1.6	2.2	3.5
Male												
2001	7.2	*	*	0.1	5.7	9.6	15.9	12.9	3.9	2.0	2.8	3.9
2000 [4]	6.6	*	*	0.1	4.9	9.1	14.8	11.7	3.3	1.7	2.1	6.2
1999 [4]	6.3	*	*	0.1	3.8	8.8	14.2	11.1	3.6	2.0	2.5	4.4
Female												
2001	3.1	*	*	*	1.5	3.5	7.1	5.4	2.3	1.2	1.9	3.4
2000 [4]	2.7	*	*	*	1.4	3.0	5.9	4.4	2.3	1.3	2.0	3.6
1999 [4]	2.4	*	*	*	1.2	2.8	5.4	4.0	1.8	1.3	2.0	3.1
Non-Hispanic white												
Both sexes												
2001	5.2	*	*	0.1	4.3	7.3	11.7	8.3	2.7	1.5	2.1	3.4
2000 [4]	4.6	*	*	0.1	3.7	6.5	10.4	7.2	2.5	1.4	2.1	4.1
1999 [4]	4.2	*	*	0.1	2.8	6.0	9.6	6.7	2.4	1.5	2.3	3.3
Male												
2001	7.3	*	*	*	6.7	10.7	16.3	11.5	3.1	1.9	2.5	3.8
2000 [4]	6.6	*	*	*	5.8	9.8	14.9	10.3	2.7	1.5	2.2	5.8
1999 [4]	6.1	*	*	*	4.3	9.2	14.0	9.6	2.9	1.8	2.5	4.6
Female												
2001	3.1	*	*	*	1.7	3.8	7.1	5.2	2.4	1.2	1.9	3.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)												
Non-Hispanic white												
Female												
2000 [4]	2.7	*	*	*	1.5	3.2	5.8	4.2	2.3	1.4	2.0	3.4
1999 [4]	2.4	*	*	*	1.3	2.8	5.1	3.8	1.9	1.2	2.1	2.8
Non-Hispanic black												
Both sexes												
2001	6.0	*	*	*	1.6	5.3	13.4	17.0	6.4	2.4	3.7	6.6
2000 [4]	5.7	*	*	*	1.4	5.7	13.2	15.7	5.5	2.5	*	8.3
1999 [4]	5.8	*	*	*	1.7	6.3	13.2	15.6	6.0	3.0	*	*
Male												
2001	8.6	*	*	*	2.3	7.6	18.2	27.0	11.0	3.7	6.0	*
2000 [4]	8.4	*	*	*	1.8	8.5	19.0	25.3	9.5	4.3	*	*
1999 [4]	8.8	*	*	*	2.8	9.3	19.5	25.9	11.0	4.4	*	*
Female												
2001	3.6	*	*	*	0.9	3.2	9.2	8.4	2.6	*	*	*
2000 [4]	3.2	*	*	*	1.0	3.1	8.0	7.5	2.4	*	*	*
1999 [4]	3.1	*	*	*	*	3.5	7.7	6.7	1.9	2.1	*	*
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)												
All origins [3]												
Both sexes												
2001	6.2	16.6	1.8	0.6	1.6	2.0	3.1	4.2	5.3	10.0	30.5	115.0
2000 [4]	6.2	15.0	1.7	0.6	1.7	2.2	3.2	4.0	5.6	9.9	29.8	117.1
1999 [4]	6.6	13.8	1.9	0.8	1.7	2.2	3.2	4.1	5.6	11.0	32.9	128.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)												
All origins [3]												
Male												
2001	7.2	18.9	2.1	0.9	2.6	3.4	5.0	6.6	7.7	13.2	38.4	134.0
2000 [4]	7.2	16.8	2.1	0.9	2.8	3.8	5.4	6.6	8.3	13.0	36.9	135.2
1999 [4]	7.6	16.2	2.3	1.1	2.9	3.7	5.3	6.6	8.3	14.8	40.9	152.7
Female												
2001	5.3	14.2	1.5	0.4	0.6	0.6	1.1	1.9	3.1	7.3	25.4	107.0
2000 [4]	5.2	13.2	1.4	0.4	0.5	0.6	1.1	1.5	3.0	7.4	25.1	109.8
1999 [4]	5.6	11.3	1.5	0.5	0.5	0.7	1.2	1.7	3.1	8.0	27.7	119.3
Hispanic												
Both sexes												
2001	2.7	9.5	1.4	0.3	1.5	1.8	2.7	3.6	3.8	7.7	16.0	63.8
2000 [4]	2.7	7.5	1.4	0.4	1.6	2.2	2.9	2.7	4.0	6.2	16.6	70.3
1999 [4]	2.6	8.6	1.9	0.5	1.6	1.6	2.4	2.8	4.2	8.7	15.0	65.9
Male												
2001	3.7	9.6	1.8	*	2.6	3.0	4.6	6.1	5.6	10.8	20.6	68.6
2000 [4]	3.8	7.9	1.5	*	2.9	3.7	5.2	4.5	6.6	8.7	19.7	88.7
1999 [4]	3.6	9.0	2.5	*	2.6	2.7	4.0	4.8	6.2	13.2	15.6	84.1
Female												
2001	1.6	9.5	*	*	*	*	*	1.2	2.1	5.4	13.0	61.4
2000 [4]	1.4	7.2	*	*	*	*	*	*	*	4.3	14.5	61.3
1999 [4]	1.6	8.1	*	*	*	*	*	*	2.4	5.2	14.6	57.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)												
Non-Hispanic [5]												
Both sexes												
2001	6.7	18.3	1.9	0.7	1.6	2.0	3.1	4.2	5.4	10.1	31.1	116.6
2000 [4]	6.7	16.9	1.8	0.7	1.7	2.2	3.2	4.1	5.7	10.1	30.2	118.5
1999 [4]	7.1	15.0	1.9	0.8	1.7	2.3	3.3	4.2	5.6	11.1	33.6	130.7
Male												
2001	7.7	21.1	2.1	1.0	2.6	3.5	5.1	6.6	7.8	13.3	39.0	136.5
2000 [4]	7.7	19.0	2.2	1.0	2.7	3.8	5.3	6.7	8.4	13.2	37.5	136.9
1999 [4]	8.1	17.9	2.3	1.1	2.9	3.9	5.4	6.6	8.4	14.8	41.8	154.8
Female												
2001	5.8	15.4	1.6	0.4	0.6	0.6	1.2	2.0	3.2	7.4	25.9	108.4
2000 [4]	5.7	14.7	1.4	0.4	0.5	0.7	1.1	1.6	3.1	7.5	25.4	111.1
1999 [4]	6.2	12.0	1.5	0.5	0.5	0.7	1.2	1.8	3.1	8.1	28.2	121.1
Non-Hispanic white												
Both sexes												
2001	7.2	14.5	1.7	0.7	1.8	2.1	3.1	4.1	5.2	9.8	31.2	119.5
2000 [4]	7.1	13.7	1.7	0.7	1.7	2.3	3.2	3.9	5.4	9.7	30.3	122.0
1999 [4]	7.6	12.1	1.7	0.9	1.9	2.4	3.4	3.9	5.2	10.6	33.1	134.3
Male												
2001	8.1	16.8	2.0	1.0	2.9	3.6	5.1	6.3	7.4	13.0	38.6	139.7
2000 [4]	8.0	16.0	2.1	1.1	2.9	4.0	5.3	6.3	7.9	12.6	36.8	139.7
1999 [4]	8.4	14.2	2.0	1.2	3.1	4.0	5.5	6.2	7.7	13.9	41.0	157.3

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)												
Non-Hispanic white												
Female												
2001	6.4	12.1	1.5	0.4	0.7	0.6	1.1	1.9	3.1	7.1	26.3	111.2
2000 [4]	6.3	11.2	1.3	0.4	0.5	0.7	1.1	1.6	3.1	7.3	26.0	114.9
1999 [4]	6.8	9.8	1.5	0.5	0.6	0.7	1.3	1.7	2.8	7.9	28.0	125.2
Non-Hispanic black												
Both sexes												
2001	5.5	35.5	2.6	0.6	1.1	2.0	3.3	5.7	8.0	14.2	34.2	96.5
2000 [4]	5.6	31.7	2.3	0.7	1.6	2.3	3.6	5.8	8.4	15.1	34.1	90.9
1999 [4]	6.3	29.5	2.6	0.7	1.3	2.5	3.5	6.7	10.2	18.2	43.1	106.7
Male												
2001	7.0	41.8	2.9	0.9	1.6	3.3	5.2	9.2	12.2	19.9	49.5	115.5
2000 [4]	7.5	32.8	2.7	1.0	2.6	3.8	6.1	10.3	14.2	20.8	53.7	124.5
1999 [4]	8.3	35.0	3.5	1.0	2.4	4.1	5.9	11.2	16.0	27.4	57.8	155.7
Female												
2001	4.2	28.9	2.3	*	*	0.7	1.6	2.7	4.7	10.3	25.6	89.4
2000 [4]	3.9	30.7	2.0	*	*	0.9	1.3	1.9	3.7	11.1	23.1	78.5
1999 [4]	4.5	23.9	*	*	*	1.1	1.4	2.8	5.5	11.8	34.8	88.4
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
All origins [3]												
Both sexes												
2001 [6]	10.8	. . .	. . .	0.7	9.9	12.8	14.7	15.2	13.1	13.3	17.4	17.5
2000 [4]	10.4	. . .	. . .	0.7	10.2	12.0	14.5	14.4	12.1	12.5	17.6	19.6
1999 [4]	10.5	. . .	. . .	0.6	10.1	12.7	14.3	13.9	12.2	13.4	18.1	19.3

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
All origins [3]												
Male												
2001 [6]	17.6	. . .	. . .	1.0	16.6	21.0	23.1	23.4	21.1	24.6	37.8	51.1
2000 [4]	17.1	. . .	. . .	1.2	17.1	19.6	22.8	22.4	19.4	22.7	38.6	57.5
1999 [4]	17.1	. . .	. . .	0.9	16.8	20.7	22.4	21.5	19.8	24.7	38.8	57.1
Female												
2001 [6]	4.1	. . .	. . .	0.3	2.9	4.4	6.4	7.2	5.7	3.9	4.0	3.4
2000 [4]	4.0	. . .	. . .	0.3	3.0	4.3	6.4	6.7	5.4	4.0	4.0	4.2
1999 [4]	4.0	. . .	. . .	0.3	3.0	4.6	6.4	6.6	5.2	4.1	4.7	4.1
Hispanic												
Both sexes												
2001 [6]	5.0	. . .	. . .	0.5	6.2	7.1	7.3	7.0	6.2	7.6	8.8	13.8
2000 [4]	5.1	. . .	. . .	0.4	6.8	6.5	7.5	7.8	6.0	7.8	11.7	*
1999 [4]	5.0	. . .	. . .	0.5	6.2	6.7	7.5	7.1	6.8	7.0	13.1	15.4
Male												
2001 [6]	8.3	. . .	. . .	0.7	9.5	11.3	12.4	11.4	11.5	15.4	20.1	41.9
2000 [4]	8.4	. . .	. . .	0.6	10.9	10.6	12.1	12.5	11.1	15.6	26.1	*
1999 [4]	8.2	. . .	. . .	0.6	9.9	10.7	12.1	11.6	12.2	13.8	28.2	*
Female												
2001 [6]	1.5	. . .	. . .	*	2.3	2.2	1.8	2.7	*	*	*	*
2000 [4]	1.5	. . .	. . .	*	2.0	1.7	2.6	3.2	*	*	*	*
1999 [4]	1.6	. . .	. . .	*	1.8	2.0	2.6	2.7	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001 [6]	11.6	. . .	. . .	0.7	10.7	13.9	15.7	15.9	13.6	13.6	17.8	17.5
2000 [4]	11.1	. . .	. . .	0.8	10.8	13.0	15.4	15.0	12.5	12.7	17.9	19.9
1999 [4]	11.2	. . .	. . .	0.6	10.8	13.8	15.1	14.4	12.5	13.7	18.3	19.3

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Non-Hispanic [5]												
Male												
2001 [6]	19.0	. . .	. . .	1.1	18.1	23.1	24.6	24.5	21.7	25.0	38.5	51.3
2000 [4]	18.3	. . .	. . .	1.3	18.3	21.4	24.1	23.2	19.9	23.1	39.1	58.4
1999 [4]	18.4	. . .	. . .	1.0	18.2	22.7	23.6	22.2	20.2	25.2	39.1	57.6
Female												
2001 [6]	4.5	. . .	. . .	0.3	3.0	4.9	7.0	7.6	6.1	4.1	4.2	3.5
2000 [4]	4.3	. . .	. . .	0.3	3.1	4.8	6.8	7.0	5.7	4.1	4.1	4.3
1999 [4]	4.3	. . .	. . .	0.2	3.2	5.0	6.8	6.9	5.4	4.3	4.8	4.0
Non-Hispanic white												
Both sexes												
2001 [6]	13.0	. . .	. . .	0.7	11.6	15.4	17.7	17.9	15.0	14.7	19.0	18.7
2000 [4]	12.5	. . .	. . .	0.9	11.6	14.2	17.2	16.7	13.9	13.8	19.0	21.3
1999 [4]	12.4	. . .	. . .	0.7	11.5	15.0	16.9	16.0	13.7	14.8	19.5	20.8
Male												
2001 [6]	21.4	. . .	. . .	1.1	19.6	25.2	27.4	27.4	23.6	26.9	41.1	55.2
2000 [4]	20.4	. . .	. . .	1.4	19.5	23.1	26.6	25.5	21.8	24.7	41.7	62.7
1999 [4]	20.3	. . .	. . .	1.1	19.2	24.3	26.0	24.4	21.9	27.0	41.6	62.2
Female												
2001 [6]	5.0	. . .	. . .	0.3	3.3	5.5	8.1	8.6	6.8	4.4	4.4	3.5
2000 [4]	4.9	. . .	. . .	0.4	3.3	5.2	7.8	7.9	6.4	4.6	4.2	4.6
1999 [4]	4.9	. . .	. . .	0.3	3.4	5.5	7.8	7.8	6.0	4.6	5.0	4.2
Non-Hispanic black												
Both sexes												
2001 [6]	5.3	. . .	. . .	0.6	7.3	9.2	7.5	6.2	5.1	5.5	5.6	*
2000 [4]	5.5	. . .	. . .	0.6	8.4	9.0	7.7	6.5	4.2	5.3	5.9	*
1999 [4]	5.5	. . .	. . .	0.5	8.3	9.5	7.8	5.7	5.0	5.8	5.8	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Non-Hispanic black												
Male												
2001 [6]	9.3	. . .	. . .	0.9	13.4	16.6	12.5	9.8	9.3	10.8	13.2	*
2000 [4]	9.6	. . .	. . .	1.0	14.6	16.1	13.3	11.1	7.5	11.4	12.3	*
1999 [4]	9.9	. . .	. . .	0.7	14.8	17.5	13.7	10.1	9.2	11.7	14.1	*
Female												
2001 [6]	1.7	. . .	. . .	*	1.3	2.4	3.0	3.1	1.8	*	*	*
2000 [4]	1.7	. . .	. . .	*	2.3	2.6	2.7	2.5	1.6	*	*	*
1999 [4]	1.6	. . .	. . .	*	2.0	2.4	2.6	2.0	1.6	*	*	*
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
All origins [3]												
Both sexes												
2001	5.9	. . .	. . .	0.2	5.3	6.5	6.7	7.7	8.2	9.6	13.1	12.2
2000 [4]	5.9	. . .	. . .	0.3	5.8	6.1	7.1	7.6	7.6	9.2	13.0	13.2
1999 [4]	5.9	. . .	. . .	0.3	6.0	6.5	6.9	7.3	7.8	9.7	13.1	12.6
Male												
2001	10.6	. . .	. . .	0.3	9.4	11.2	11.4	13.3	14.6	19.2	30.7	38.9
2000 [4]	10.5	. . .	. . .	0.4	10.2	10.6	11.8	13.0	13.5	18.4	30.8	42.9
1999 [4]	10.6	. . .	. . .	0.4	10.5	11.2	11.5	12.5	14.0	19.5	31.1	41.6
Female												
2001	1.5	. . .	. . .	0.1	1.0	1.6	2.1	2.4	2.4	1.7	1.5	1.0
2000 [4]	1.5	. . .	. . .	0.1	1.2	1.7	2.4	2.3	2.1	1.7	1.4	1.1
1999 [4]	1.5	. . .	. . .	0.1	1.2	1.8	2.3	2.3	2.1	1.7	1.4	0.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
Hispanic												
Both sexes												
2001	2.2	. . .	. . .	*	2.9	3.3	2.5	3.0	3.2	4.0	4.7	*
2000 [4]	2.3	. . .	. . .	*	3.4	2.8	2.8	4.0	2.7	4.6	5.3	*
1999 [4]	2.3	. . .	. . .	*	3.5	2.9	3.0	3.2	2.9	3.6	8.1	*
Male												
2001	3.8	. . .	. . .	*	4.8	5.5	4.5	5.3	6.6	8.8	11.6	*
2000 [4]	4.1	. . .	. . .	*	5.8	4.7	4.7	6.8	5.7	9.7	12.8	*
1999 [4]	4.0	. . .	. . .	*	5.8	4.8	5.0	5.5	5.9	7.7	19.8	*
Female												
2001	0.4	. . .	. . .	*	0.7	0.7	*	*	*	*	*	*
2000 [4]	0.4	. . .	. . .	*	*	*	*	*	*	*	*	*
1999 [4]	0.5	. . .	. . .	*	0.8	*	0.9	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	6.5	. . .	. . .	0.2	5.8	7.1	7.3	8.1	8.6	9.9	13.5	12.4
2000 [4]	6.4	. . .	. . .	0.3	6.2	6.8	7.6	7.9	7.9	9.5	13.4	13.4
1999 [4]	6.4	. . .	. . .	0.3	6.5	7.2	7.3	7.7	8.2	10.1	13.3	12.7
Male												
2001	11.6	. . .	. . .	0.4	10.4	12.5	12.3	14.0	15.1	19.8	31.5	39.7
2000 [4]	11.4	. . .	. . .	0.5	11.1	11.7	12.7	13.5	14.0	18.9	31.6	43.8
1999 [4]	11.5	. . .	. . .	0.4	11.5	12.5	12.3	13.1	14.5	20.1	31.5	42.2
Female												
2001	1.6	. . .	. . .	0.1	1.1	1.8	2.4	2.5	2.5	1.8	1.6	1.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
Non-Hispanic [5]												
Female												
2000 [4]	1.6	. . .	. . .	0.1	1.3	1.9	2.6	2.4	2.3	1.8	1.5	1.1
1999 [4]	1.6	. . .	. . .	*	1.3	1.9	2.4	2.4	2.3	1.8	1.5	0.9
Non-Hispanic white												
Both sexes												
2001	7.4	. . .	. . .	0.3	6.2	8.0	8.3	9.3	9.6	10.9	14.5	13.4
2000 [4]	7.3	. . .	. . .	0.4	6.7	7.4	8.6	8.9	8.7	10.4	14.5	14.5
1999 [4]	7.3	. . .	. . .	0.3	6.8	7.9	8.3	8.6	9.0	11.0	14.3	13.7
Male												
2001	13.2	. . .	. . .	0.4	11.0	13.8	13.9	15.9	16.7	21.6	33.8	43.0
2000 [4]	12.9	. . .	. . .	0.6	11.8	12.7	14.3	15.1	15.3	20.4	34.2	47.6
1999 [4]	12.9	. . .	. . .	0.5	12.0	13.4	13.7	14.6	15.9	21.8	33.8	45.7
Female												
2001	1.9	. . .	. . .	*	1.2	2.1	2.7	2.9	2.9	1.9	1.7	1.1
2000 [4]	1.9	. . .	. . .	*	1.4	2.1	3.0	2.8	2.6	2.0	1.5	1.2
1999 [4]	1.9	. . .	. . .	*	1.4	2.3	2.9	2.7	2.6	2.0	1.6	1.0
Non-Hispanic black												
Both sexes												
2001	3.0	. . .	. . .	*	5.1	5.0	3.5	2.8	2.7	3.8	5.1	*
2000 [4]	3.1	. . .	. . .	*	5.4	5.2	3.5	3.2	3.0	3.8	3.6	*
1999 [4]	3.2	. . .	. . .	*	5.8	5.1	3.6	3.0	3.5	4.0	4.8	*
Male												
2001	5.6	. . .	. . .	*	9.6	9.5	6.4	4.9	5.5	7.9	12.0	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
Non-Hispanic black												
Male												
2000 [4]	5.7	. . .	. . .	*	9.9	9.6	6.4	5.6	5.7	9.0	8.3	*
1999 [4]	6.1	. . .	. . .	*	10.6	10.0	6.9	5.6	7.1	8.9	12.2	*
Female												
2001	0.6	. . .	. . .	*	*	0.9	1.0	1.0	*	*	*	*
2000 [4]	0.7	. . .	. . .	*	1.0	1.1	0.8	1.1	*	*	*	*
1999 [4]	0.6	. . .	. . .	*	1.0	0.8	*	*	*	*	*	*
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84, Y87.0)												
All origins [3]												
Both sexes												
2001 [6]	4.8	. . .	. . .	0.5	4.6	6.3	8.0	7.4	4.9	3.7	4.3	5.3
2000 [4]	4.5	. . .	. . .	0.5	4.4	5.9	7.5	6.8	4.6	3.2	4.6	6.5
1999 [4]	4.5	. . .	. . .	0.3	4.1	6.2	7.5	6.6	4.4	3.7	5.0	6.7
Male												
2001 [6]	7.1	. . .	. . .	0.7	7.2	9.8	11.7	10.2	6.5	5.4	7.1	12.2
2000 [4]	6.6	. . .	. . .	0.7	6.9	9.0	10.9	9.4	6.0	4.3	7.8	14.6
1999 [4]	6.6	. . .	. . .	0.5	6.3	9.5	10.9	8.9	5.8	5.3	7.7	15.5
Female												
2001 [6]	2.6	. . .	. . .	0.2	1.9	2.8	4.3	4.8	3.4	2.3	2.5	2.4
2000 [4]	2.5	. . .	. . .	0.2	1.8	2.7	4.0	4.4	3.3	2.3	2.6	3.2
1999 [4]	2.5	. . .	. . .	0.1	1.8	2.8	4.1	4.3	3.0	2.4	3.3	3.1
Hispanic												
Both sexes												
2001 [6]	2.8	. . .	. . .	0.5	3.3	3.9	4.8	4.0	3.0	3.6	4.1	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84, Y87.0)												
Hispanic												
Both sexes												
2000 [4]	2.8	. . .	. . .	0.3	3.4	3.7	4.7	3.8	3.2	3.3	6.3	*
1999 [4]	2.7	. . .	. . .	*	2.7	3.8	4.5	3.8	3.9	3.5	5.0	*
Male												
2001 [6]	4.4	. . .	. . .	0.6	4.7	5.8	7.9	6.1	4.9	6.6	*	*
2000 [4]	4.3	. . .	. . .	*	5.1	5.9	7.3	5.7	5.4	5.9	13.3	*
1999 [4]	4.2	. . .	. . .	*	4.2	5.9	7.1	6.0	6.4	6.2	*	*
Female												
2001 [6]	1.2	. . .	. . .	*	1.6	1.6	1.4	2.0	*	*	*	*
2000 [4]	1.1	. . .	. . .	*	1.4	1.1	1.9	2.0	*	*	*	*
1999 [4]	1.1	. . .	. . .	*	1.0	1.4	1.7	1.7	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001 [6]	5.1	. . .	. . .	0.5	4.8	6.8	8.4	7.7	5.0	3.7	4.3	5.1
2000 [4]	4.8	. . .	. . .	0.5	4.6	6.2	7.8	7.1	4.6	3.2	4.5	6.5
1999 [4]	4.7	. . .	. . .	0.4	4.4	6.6	7.8	6.8	4.4	3.7	5.0	6.6
Male												
2001 [6]	7.5	. . .	. . .	0.7	7.7	10.6	12.2	10.5	6.6	5.3	7.0	11.6
2000 [4]	6.9	. . .	. . .	0.8	7.2	9.6	11.4	9.7	6.0	4.2	7.5	14.6
1999 [4]	6.9	. . .	. . .	0.6	6.7	10.2	11.3	9.1	5.7	5.1	7.6	15.4
Female												
2001 [6]	2.9	. . .	. . .	0.2	1.9	3.1	4.7	5.1	3.5	2.3	2.6	2.5
2000 [4]	2.7	. . .	. . .	0.2	1.9	2.9	4.3	4.6	3.4	2.4	2.6	3.2
1999 [4]	2.7	. . .	. . .	0.1	1.9	3.0	4.4	4.5	3.1	2.5	3.3	3.1

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84, Y87.0)												
Non-Hispanic white												
Both sexes												
2001 [6]	5.6	. . .	. . .	0.4	5.4	7.5	9.4	8.6	5.3	3.8	4.5	5.3
2000 [4]	5.2	. . .	. . .	0.5	4.9	6.8	8.6	7.8	5.1	3.3	4.5	6.8
1999 [4]	5.2	. . .	. . .	0.4	4.6	7.1	8.6	7.4	4.6	3.8	5.1	7.0
Male												
2001 [6]	8.2	. . .	. . .	0.7	8.6	11.5	13.5	11.5	6.9	5.3	7.3	12.3
2000 [4]	7.5	. . .	. . .	0.8	7.7	10.3	12.4	10.4	6.5	4.3	7.5	15.2
1999 [4]	7.4	. . .	. . .	0.6	7.2	10.8	12.3	9.8	6.0	5.3	7.8	16.5
Female												
2001 [6]	3.2	. . .	. . .	0.2	2.0	3.4	5.3	5.7	3.9	2.4	2.7	2.4
2000 [4]	3.0	. . .	. . .	0.2	1.9	3.1	4.8	5.2	3.8	2.5	2.6	3.4
1999 [4]	3.0	. . .	. . .	*	2.0	3.3	4.9	5.1	3.4	2.6	3.4	3.2
Non-Hispanic black												
Both sexes												
2001 [6]	2.3	. . .	. . .	0.4	2.3	4.2	3.9	3.4	2.4	1.7	*	*
2000 [4]	2.4	. . .	. . .	0.5	3.0	3.9	4.2	3.3	1.3	1.4	2.3	*
1999 [4]	2.3	. . .	. . .	0.4	2.5	4.4	4.2	2.7	1.5	1.8	*	*
Male												
2001 [6]	3.7	. . .	. . .	0.7	3.8	7.1	6.1	4.9	3.7	3.0	*	*
2000 [4]	3.9	. . .	. . .	0.9	4.7	6.5	6.8	5.4	*	*	*	*
1999 [4]	3.8	. . .	. . .	*	4.1	7.5	6.8	4.5	2.1	*	*	*
Female												
2001 [6]	1.1	. . .	. . .	*	0.7	1.5	2.0	2.1	*	*	*	*
2000 [4]	1.1	. . .	. . .	*	1.3	1.5	1.9	1.5	*	*	*	*
1999 [4]	1.0	. . .	. . .	*	0.9	1.6	2.0	1.3	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
All origins [3]												
Both sexes												
2001 [6]	7.1	8.2	2.7	0.8	13.3	13.1	9.5	6.3	4.0	2.9	2.5	2.4
2000 [4]	6.0	9.2	2.3	0.9	12.6	10.4	7.1	4.7	3.0	2.4	2.4	2.4
1999 [4]	6.1	8.7	2.5	1.1	12.9	10.5	7.1	4.6	3.0	2.6	2.5	2.4
Male												
2001 [6]	11.1	9.5	3.0	0.9	22.2	20.8	13.9	9.5	5.9	4.0	3.1	3.2
2000 [4]	9.3	10.4	2.5	1.1	20.9	16.7	10.3	6.9	4.6	3.4	3.2	3.3
1999 [4]	9.3	9.6	2.5	1.2	21.1	16.5	10.3	6.8	4.7	3.7	3.4	3.4
Female												
2001 [6]	3.3	6.9	2.4	0.7	3.9	5.3	5.1	3.2	2.2	2.0	2.1	2.0
2000 [4]	2.8	7.9	2.1	0.7	3.9	4.1	4.0	2.5	1.6	1.6	2.0	2.0
1999 [4]	2.9	7.8	2.4	0.9	4.3	4.4	4.0	2.4	1.5	1.7	1.9	2.0
Hispanic												
Both sexes												
2001 [6]	9.0	6.7	2.3	0.8	18.3	14.0	10.4	7.3	4.4	4.0	*	*
2000 [4]	8.3	7.0	1.9	1.1	17.1	12.4	8.7	6.7	4.5	3.3	*	*
1999 [4]	8.4	7.0	1.8	1.4	17.7	12.7	8.8	6.6	4.2	3.2	*	*
Male												
2001 [6]	14.5	7.9	2.5	1.1	30.5	21.6	15.9	11.3	6.8	6.6	*	*
2000 [4]	13.4	6.6	2.0	1.6	28.5	19.9	13.5	10.2	7.1	4.6	*	*
1999 [4]	13.8	6.7	1.7	1.8	29.1	20.6	13.9	10.3	7.3	5.7	*	*
Female												
2001 [6]	3.2	5.5	2.1	*	4.0	5.1	4.4	3.4	2.4	*	*	*
2000 [4]	2.8	7.4	1.9	0.6	3.7	3.7	3.6	3.3	2.2	*	*	*
1999 [4]	2.8	7.3	1.9	0.9	4.4	3.5	3.3	3.1	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
Non-Hispanic [5]												
Both sexes												
2001 [6]	6.7	8.4	2.8	0.8	12.1	12.8	9.2	6.1	3.9	2.8	2.5	2.4
2000 [4]	5.6	9.6	2.4	0.9	11.6	10.0	6.9	4.4	2.9	2.3	2.4	2.3
1999 [4]	5.7	8.9	2.6	1.0	11.9	10.0	6.8	4.3	2.9	2.5	2.5	2.4
Male												
2001 [6]	10.4	9.8	3.2	0.9	20.1	20.3	13.4	9.2	5.8	3.8	3.1	3.1
2000 [4]	8.6	11.2	2.6	1.0	19.1	15.9	9.8	6.6	4.4	3.2	3.1	3.1
1999 [4]	8.6	10.1	2.6	1.1	19.4	15.6	9.7	6.4	4.5	3.5	3.3	3.4
Female												
2001 [6]	3.2	6.9	2.4	0.7	3.8	5.3	5.1	3.1	2.2	2.0	2.1	2.1
2000 [4]	2.7	7.9	2.2	0.7	3.8	4.1	4.0	2.3	1.5	1.6	1.9	2.0
1999 [4]	2.9	7.6	2.5	0.9	4.2	4.6	4.0	2.3	1.4	1.7	2.0	2.0
Non-Hispanic white												
Both sexes												
2001 [6]	4.0	5.8	1.7	0.6	4.2	6.5	6.6	4.6	3.3	2.3	2.1	2.2
2000 [4]	2.8	6.2	1.6	0.6	3.5	4.0	4.0	2.9	2.2	1.8	2.0	2.1
1999 [4]	2.9	6.3	1.7	0.7	3.9	4.3	4.1	2.9	2.2	2.1	2.0	2.1
Male												
2001 [6]	5.6	6.8	2.0	0.6	5.7	9.3	9.5	6.7	4.7	2.9	2.6	2.8
2000 [4]	3.6	8.3	1.8	0.7	4.7	5.2	5.2	3.9	3.1	2.2	2.4	2.6
1999 [4]	3.9	8.0	1.7	0.7	5.1	5.7	5.4	4.2	3.2	2.7	2.6	2.6
Female												
2001 [6]	2.4	4.7	1.5	0.7	2.7	3.7	3.7	2.6	1.9	1.8	1.8	2.0
2000 [4]	1.9	4.1	1.3	0.6	2.3	2.7	2.7	1.9	1.3	1.4	1.7	1.9
1999 [4]	2.0	4.4	1.6	0.7	2.6	2.9	2.8	1.7	1.3	1.6	1.6	1.9

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
Non-Hispanic black												
Both sexes												
2001 [6]	22.7	19.5	7.1	1.4	48.7	45.3	25.5	16.2	8.9	7.2	6.8	*
2000 [4]	22.1	24.1	6.1	1.9	49.3	42.2	24.5	15.0	8.4	7.6	7.1	*
1999 [4]	21.7	20.0	6.4	2.3	49.5	40.6	23.7	14.1	8.4	6.4	7.5	*
Male												
2001 [6]	39.3	21.7	8.4	2.0	88.3	81.7	39.5	27.2	15.0	12.6	10.2	*
2000 [4]	38.3	24.9	6.1	2.1	88.1	76.5	39.4	26.3	15.0	13.9	12.0	*
1999 [4]	37.2	20.1	6.2	2.7	88.1	71.5	37.9	23.7	15.4	11.9	12.2	*
Female												
2001 [6]	7.6	17.1	5.7	0.8	9.2	12.5	13.2	6.9	4.0	3.3	4.9	*
2000 [4]	7.4	23.3	6.0	1.6	11.0	11.4	11.3	5.3	3.2	3.2	4.3	*
1999 [4]	7.7	20.0	6.7	1.8	11.4	12.8	11.1	5.9	2.7	2.5	4.9	*
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
All origins [3]												
Both sexes												
2001	4.0	*	0.4	0.4	10.5	8.4	4.4	2.4	1.4	1.0	0.8	*
2000 [4]	3.8	*	0.2	0.5	10.1	7.7	4.2	2.5	1.5	1.0	0.7	0.7
1999 [4]	3.9	*	0.3	0.5	10.4	7.6	4.1	2.4	1.6	1.0	0.8	0.6
Male												
2001	6.8	*	0.3	0.6	18.5	14.3	6.8	3.8	2.2	1.5	1.1	*
2000 [4]	6.5	*	*	0.7	17.7	13.2	6.6	3.9	2.3	1.3	0.8	*
1999 [4]	6.5	*	0.3	0.7	18.0	12.9	6.3	3.8	2.6	1.5	1.0	*
Female												
2001	1.3	*	0.4	0.3	2.1	2.3	2.0	1.0	0.8	0.7	0.6	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
All origins [3]												
Female												
2000 [4]	1.3	*	*	0.2	2.2	2.1	1.8	1.2	0.8	0.8	0.6	*
1999 [4]	1.3	*	0.3	0.3	2.5	2.2	2.0	1.2	0.6	0.7	0.6	*
Hispanic												
Both sexes												
2001	5.7	*	*	0.4	14.6	9.5	5.6	3.0	1.1	*	*	*
2000 [4]	5.5	*	*	0.6	13.7	8.9	4.9	4.0	2.0	*	*	*
1999 [4]	5.7	*	*	0.8	14.4	8.7	5.2	3.8	2.1	*	*	*
Male												
2001	9.9	*	*	0.6	24.9	16.1	9.1	4.8	*	*	*	*
2000 [4]	9.6	*	*	1.0	23.5	15.0	7.8	6.4	3.2	*	*	*
1999 [4]	9.8	*	*	1.2	24.2	14.7	8.6	5.7	3.5	*	*	*
Female												
2001	1.3	*	*	*	2.5	1.8	1.9	1.2	*	*	*	*
2000 [4]	1.3	*	*	*	2.2	1.9	1.7	1.7	*	*	*	*
1999 [4]	1.4	*	*	*	2.9	1.7	1.7	1.9	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	3.7	*	0.3	0.4	9.6	8.0	4.2	2.3	1.5	1.0	0.8	*
2000 [4]	3.6	*	0.2	0.4	9.3	7.4	4.1	2.4	1.5	1.0	0.7	0.7
1999 [4]	3.6	*	0.4	0.5	9.6	7.3	4.0	2.3	1.5	1.1	0.8	0.5
Male												
2001	6.3	*	0.3	0.6	17.0	13.8	6.5	3.7	2.2	1.4	1.1	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
Non-Hispanic [5]												
Male												
2000 [4]	6.0	*	*	0.6	16.3	12.7	6.4	3.7	2.2	1.3	0.8	*
1999 [4]	6.0	*	0.4	0.7	16.7	12.4	6.0	3.6	2.5	1.5	0.9	*
Female												
2001	1.2	*	0.3	0.3	2.0	2.3	2.0	1.0	0.8	0.7	0.6	*
2000 [4]	1.2	*	*	0.2	2.2	2.2	1.9	1.1	0.8	0.7	0.6	*
1999 [4]	1.3	*	0.4	0.3	2.4	2.3	2.0	1.1	0.6	0.7	0.6	*
Non-Hispanic white												
Both sexes												
2001	1.6	*	0.2	0.4	2.4	2.7	2.4	1.6	1.2	0.8	0.7	*
2000 [4]	1.4	*	*	0.3	2.2	2.4	2.3	1.6	1.2	0.8	0.6	0.6
1999 [4]	1.5	*	0.2	0.3	2.4	2.6	2.3	1.5	1.1	0.9	0.6	0.6
Male												
2001	2.2	*	*	0.4	3.6	3.9	3.4	2.3	1.6	1.0	0.9	*
2000 [4]	2.0	*	*	0.4	3.2	3.4	3.2	2.2	1.6	0.9	0.6	*
1999 [4]	2.2	*	*	0.4	3.5	3.9	3.2	2.2	1.7	1.2	0.6	*
Female												
2001	0.9	*	*	0.3	1.2	1.5	1.4	0.8	0.8	0.7	0.6	*
2000 [4]	0.9	*	*	0.2	1.2	1.4	1.4	0.9	0.8	0.7	0.6	*
1999 [4]	0.9	*	*	0.2	1.2	1.4	1.5	0.8	0.6	0.7	0.6	*
Non-Hispanic black												
Both sexes												
2001	16.3	*	*	0.9	43.0	36.4	15.5	7.7	3.7	2.6	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
Non-Hispanic black												
Both sexes												
2000 [4]	16.0	*	*	1.0	42.6	34.5	15.4	8.5	3.6	2.8	*	*
1999 [4]	15.9	*	*	1.3	43.6	32.4	14.4	8.1	4.5	2.4	2.4	*
Male												
2001	30.6	*	*	1.3	80.6	69.4	26.7	14.0	7.1	5.5	*	*
2000 [4]	30.1	*	*	1.6	79.1	65.7	27.4	15.3	6.5	5.5	*	*
1999 [4]	29.3	*	*	1.8	80.0	61.1	24.6	14.1	9.1	4.7	*	*
Female												
2001	3.2	*	*	*	5.6	6.5	5.5	2.2	*	*	*	*
2000 [4]	3.3	*	*	*	6.7	6.4	4.7	2.6	*	*	*	*
1999 [4]	3.7	*	*	0.7	7.8	6.6	5.5	3.0	*	*	*	*
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)												
All origins [3]												
Both sexes												
2001 [6]	3.1	8.0	2.3	0.4	2.7	4.8	5.1	3.9	2.6	1.9	1.7	2.0
2000 [4]	2.1	8.9	2.1	0.4	2.5	2.7	2.9	2.1	1.5	1.4	1.8	1.7
1999 [4]	2.2	8.5	2.1	0.5	2.5	2.9	3.0	2.1	1.5	1.5	1.7	1.8
Male												
2001 [6]	4.3	9.2	2.7	0.3	3.6	6.5	7.1	5.7	3.8	2.5	2.0	2.7
2000 [4]	2.8	10.0	2.3	0.4	3.3	3.5	3.7	3.0	2.3	2.1	2.4	2.5
1999 [4]	2.8	9.4	2.1	0.5	3.1	3.6	4.0	3.1	2.2	2.2	2.4	2.6
Female												
2001 [6]	2.0	6.7	2.0	0.4	1.8	3.1	3.1	2.1	1.5	1.3	1.5	1.7

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)												
All origins [3]												
Female												
2000 [4]	1.5	7.7	2.0	0.5	1.7	1.9	2.1	1.3	0.8	0.9	1.4	1.4
1999 [4]	1.6	7.6	2.1	0.6	1.8	2.2	2.0	1.2	0.8	1.0	1.3	1.5
Hispanic												
Both sexes												
2001 [6]	3.3	6.2	1.9	0.4	3.7	4.5	4.8	4.3	3.4	2.7	*	*
2000 [4]	2.7	6.9	1.8	0.5	3.4	3.5	3.9	2.7	2.5	2.0	*	*
1999 [4]	2.7	6.7	1.6	0.6	3.3	4.0	3.5	2.9	2.1	2.3	*	*
Male												
2001 [6]	4.5	7.2	2.2	*	5.6	5.5	6.9	6.5	5.1	4.6	*	*
2000 [4]	3.8	6.6	1.8	0.6	5.0	5.0	5.7	3.8	3.9	*	*	*
1999 [4]	4.0	6.2	1.5	0.6	4.9	5.9	5.3	4.5	3.8	4.4	*	*
Female												
2001 [6]	1.9	5.2	1.6	*	1.5	3.2	2.6	2.2	*	*	*	*
2000 [4]	1.6	7.2	1.7	*	1.5	1.9	1.9	1.6	*	*	*	*
1999 [4]	1.4	7.3	1.8	*	1.5	1.8	1.6	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001 [6]	3.1	8.2	2.4	0.4	2.5	4.7	5.0	3.8	2.5	1.8	1.7	2.1
2000 [4]	2.0	9.2	2.2	0.4	2.3	2.6	2.8	2.0	1.4	1.4	1.7	1.6
1999 [4]	2.1	8.7	2.2	0.5	2.3	2.7	2.9	2.0	1.4	1.5	1.7	1.8
Male												
2001 [6]	4.2	9.5	2.8	0.3	3.2	6.5	7.0	5.5	3.6	2.4	2.0	2.7
2000 [4]	2.6	10.6	2.4	0.4	2.9	3.2	3.4	2.9	2.2	2.0	2.3	2.4
1999 [4]	2.6	10.0	2.3	0.4	2.8	3.2	3.8	2.9	2.0	2.0	2.4	2.6

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)												
Non-Hispanic [5]												
Female												
2001 [6]	2.0	6.8	2.0	0.4	1.9	2.9	3.1	2.1	1.4	1.3	1.5	1.8
2000 [4]	1.5	7.8	2.0	0.5	1.7	1.9	2.1	1.2	0.7	0.9	1.3	1.3
1999 [4]	1.6	7.3	2.2	0.6	1.8	2.3	2.0	1.2	0.8	1.0	1.3	1.5
Non-Hispanic white												
Both sexes												
2001 [6]	2.4	5.6	1.5	0.3	1.7	3.8	4.2	3.1	2.1	1.5	1.4	1.9
2000 [4]	1.3	6.0	1.4	0.4	1.3	1.6	1.7	1.4	1.0	1.0	1.4	1.5
1999 [4]	1.4	6.2	1.4	0.4	1.5	1.7	1.8	1.4	1.1	1.2	1.4	1.5
Male												
2001 [6]	3.4	6.7	1.8	0.2	2.0	5.4	6.1	4.5	3.1	1.9	1.7	2.3
2000 [4]	1.6	8.1	1.7	0.3	1.5	1.8	2.0	1.8	1.5	1.3	1.8	1.9
1999 [4]	1.7	7.8	1.5	0.3	1.6	1.8	2.3	2.0	1.5	1.5	2.0	*
Female												
2001 [6]	1.5	4.5	1.2	0.4	1.4	2.2	2.3	1.7	1.2	1.1	1.2	1.7
2000 [4]	1.0	3.9	1.2	0.4	1.2	1.3	1.4	1.0	0.5	0.7	1.1	1.3
1999 [4]	1.1	4.4	1.4	0.4	1.4	1.6	1.3	0.9	0.7	0.9	1.0	1.4
Non-Hispanic black												
Both sexes												
2001 [6]	6.4	19.1	6.3	0.6	5.7	9.0	10.1	8.6	5.1	4.5	4.9	*
2000 [4]	6.1	22.9	5.7	0.8	6.7	7.8	9.1	6.5	4.9	4.8	5.2	*
1999 [4]	5.8	19.7	5.7	1.0	5.9	8.2	9.2	6.0	3.9	4.0	5.2	*
Male												
2001 [6]	8.7	21.4	7.7	0.7	7.7	12.3	12.8	13.2	7.9	7.1	*	*
2000 [4]	8.2	22.5	5.7	*	9.1	10.8	12.0	11.0	8.5	8.4	8.3	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)												
Non-Hispanic black												
Male												
1999 [4]	7.9	19.7	5.6	0.9	8.1	10.5	13.3	9.6	6.3	7.2	7.2	*
Female												
2001 [6]	4.4	16.8	5.0	*	3.6	5.9	7.7	4.6	2.9	2.7	4.4	*
2000 [4]	4.1	23.3	5.8	1.1	4.3	5.0	6.5	2.7	2.0	2.3	3.5	*
1999 [4]	4.0	19.6	5.7	1.1	3.6	6.1	5.6	2.9	1.9	*	4.0	*
Legal intervention (Y35,Y89.0)												
All origins [3]												
Both sexes												
2001	0.1	*	*	*	0.2	0.3	0.3	0.1	0.1	*	*	*
2000 [4]	0.1	*	*	*	0.2	0.3	0.2	0.1	*	*	*	*
1999 [4]	0.1	*	*	*	0.2	0.3	0.3	0.1	0.1	*	*	*
Male												
2001	0.3	*	*	*	0.4	0.5	0.5	0.3	0.2	*	*	*
2000 [4]	0.3	*	*	*	0.4	0.5	0.4	0.3	*	*	*	*
1999 [4]	0.3	*	*	*	0.3	0.7	0.5	0.3	0.2	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2001	0.2	*	*	*	0.3	0.3	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Legal intervention (Y35,Y89.0)												
Hispanic												
Both sexes												
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	*	0.4	*	*	*	*	*	*
Male												
2001	0.3	*	*	*	0.6	0.6	*	*	*	*	*	*
2000 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	0.7	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.1	*	*	*	0.2	0.3	0.3	0.1	0.1	*	*	*
2000 [4]	0.1	*	*	*	0.2	0.3	0.2	0.1	*	*	*	*
1999 [4]	0.1	*	*	*	0.2	0.3	0.3	0.1	*	*	*	*
Male												
2001	0.3	*	*	*	0.4	0.5	0.5	0.3	0.2	*	*	*
2000 [4]	0.2	*	*	*	0.3	0.6	0.4	0.3	*	*	*	*
1999 [4]	0.3	*	*	*	0.3	0.7	0.5	0.3	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Legal intervention (Y35,Y89.0)												
Non-Hispanic [5]												
Female												
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2001	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*
2000 [4]	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*
1999 [4]	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*
Male												
2001	0.2	*	*	*	0.3	0.4	0.4	0.2	*	*	*	*
2000 [4]	0.2	*	*	*	0.2	0.4	0.3	0.3	*	*	*	*
1999 [4]	0.2	*	*	*	0.2	0.5	0.4	0.2	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2001	0.2	*	*	*	0.4	0.4	0.4	*	*	*	*	*
2000 [4]	0.3	*	*	*	0.5	0.6	0.6	*	*	*	*	*
1999 [4]	0.4	*	*	*	0.6	0.8	0.7	*	*	*	*	*
Male												
2001	0.5	*	*	*	0.8	0.8	0.8	*	*	*	*	*
2000 [4]	0.6	*	*	*	1.0	1.3	1.3	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Legal intervention (Y35,Y89.0)												
Non-Hispanic black												
Male												
1999 [4]	0.8	*	*	*	1.1	1.7	1.4	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
All origins [3]												
Both sexes												
2001	1.5	2.1	0.3	0.1	1.0	1.8	3.1	2.5	0.9	0.5	0.9	1.0
2000 [4]	1.4	1.7	0.3	0.1	0.9	1.7	2.8	2.2	0.9	0.6	0.6	1.3
1999 [4]	1.4	1.4	0.3	0.1	1.1	1.8	2.9	2.2	0.9	0.6	0.8	1.4
Male												
2001	2.0	2.4	*	0.2	1.5	2.6	4.2	3.4	1.2	0.8	1.1	1.5
2000 [4]	1.9	1.4	0.3	0.2	1.4	2.4	3.9	3.1	1.3	0.8	1.0	1.7
1999 [4]	2.0	1.7	0.3	0.2	1.7	2.6	4.0	3.0	1.3	0.9	1.1	2.1
Female												
2001	0.9	1.7	0.3	*	0.5	1.1	2.1	1.6	0.6	0.3	0.7	0.8
2000 [4]	0.8	2.0	*	*	0.4	1.0	1.8	1.4	0.6	0.3	0.4	1.2
1999 [4]	0.9	1.2	0.3	*	0.5	1.1	1.8	1.4	0.6	0.4	0.6	1.1
Hispanic												
Both sexes												
2001	0.8	*	*	*	0.5	0.9	1.7	1.6	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
Hispanic												
Both sexes												
2000 [4]	0.6	*	*	*	0.5	0.9	1.2	0.8	*	*	*	*
1999 [4]	0.7	*	*	*	0.8	1.1	1.2	0.9	*	*	*	*
Male												
2001	1.1	*	*	*	0.7	1.4	2.6	2.1	*	*	*	*
2000 [4]	0.9	*	*	*	0.8	1.4	1.8	1.3	*	*	*	*
1999 [4]	1.1	*	*	*	1.3	1.6	1.9	1.6	*	*	*	*
Female												
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	1.6	2.4	0.3	0.1	1.1	2.0	3.3	2.6	0.9	0.5	0.9	0.9
2000 [4]	1.5	1.9	0.3	0.2	1.0	1.8	3.0	2.4	0.9	0.6	0.6	1.4
1999 [4]	1.5	1.6	0.3	0.1	1.2	2.0	3.1	2.3	0.9	0.6	0.8	1.4
Male												
2001	2.1	2.7	*	0.2	1.7	2.8	4.4	3.5	1.2	0.7	1.2	*
2000 [4]	2.0	1.6	0.4	0.3	1.5	2.6	4.2	3.2	1.3	0.8	1.0	1.7
1999 [4]	2.1	1.9	*	0.2	1.8	2.7	4.2	3.1	1.3	0.9	1.1	2.1
Female												
2001	1.0	2.0	0.4	*	0.5	1.2	2.2	1.7	0.6	0.4	0.7	0.8
2000 [4]	0.9	2.2	*	*	0.5	1.1	1.9	1.5	0.6	0.3	0.4	1.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
Non-Hispanic [5]												
Female												
1999 [4]	0.9	*	*	*	0.5	1.2	2.0	1.5	0.6	0.4	0.6	1.1
Non-Hispanic white												
Both sexes												
2001	1.6	1.7	0.3	0.1	1.3	2.2	3.3	2.5	0.9	0.5	0.9	1.0
2000 [4]	1.4	1.5	0.2	0.1	1.1	1.9	3.0	2.3	0.8	0.5	0.7	1.3
1999 [4]	1.4	1.3	*	0.1	1.2	2.0	3.0	2.1	0.9	0.6	0.8	1.4
Male												
2001	2.1	2.0	*	0.2	1.8	3.0	4.3	3.3	1.1	0.7	1.2	*
2000 [4]	2.0	*	*	0.2	1.6	2.6	4.1	3.0	1.1	0.7	1.0	*
1999 [4]	2.0	*	*	0.2	1.8	2.7	4.0	2.6	1.1	0.8	1.1	2.1
Female												
2001	1.1	*	*	*	0.6	1.4	2.4	1.8	0.7	0.4	0.6	0.8
2000 [4]	0.9	*	*	*	0.6	1.2	1.9	1.5	0.6	0.3	0.4	1.2
1999 [4]	1.0	*	*	*	0.5	1.3	1.9	1.5	0.7	0.4	0.6	1.1
Non-Hispanic black												
Both sexes												
2001	1.8	5.9	*	*	0.7	1.7	4.0	3.8	1.2	*	*	*
2000 [4]	1.9	*	*	0.3	0.9	2.0	4.1	3.7	1.8	*	*	*
1999 [4]	2.0	*	*	*	1.3	2.3	4.3	4.2	1.2	*	*	*
Male												
2001	2.8	6.4	*	*	1.3	2.7	6.3	6.3	2.4	*	*	*
2000 [4]	2.9	*	*	*	1.6	3.2	6.1	6.0	3.0	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
Non-Hispanic black												
Male												
1999 [4]	3.1	*	*	*	2.0	3.7	6.6	7.3	2.6	*	*	*
Female												
2001	0.9	*	*	*	*	0.9	1.9	1.5	*	*	*	*
2000 [4]	1.0	*	*	*	*	0.9	2.4	1.8	*	*	*	*
1999 [4]	0.9	*	*	*	*	1.0	2.4	1.6	*	*	*	*
Discharge of firearms, undetermined intent (Y22-Y24)												
All origins [3]												
Both sexes												
2001	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2000 [4]	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
1999 [4]	0.1	*	*	*	0.3	0.1	0.1	0.1	*	*	*	*
Male												
2001	0.1	*	*	*	0.3	0.2	0.1	0.1	*	*	*	*
2000 [4]	0.1	*	*	*	0.3	0.2	0.1	*	*	*	*	*
1999 [4]	0.2	*	*	*	0.6	0.2	0.2	0.1	*	*	*	*
Female												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Discharge of firearms, undetermined intent (Y22-Y24)												
Hispanic												
Both sexes												
2000 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.1	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2000 [4]	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
1999 [4]	0.1	*	*	*	0.3	0.1	0.1	0.1	*	*	*	*
Male												
2001	0.1	*	*	*	0.3	0.2	0.2	*	*	*	*	*
2000 [4]	0.1	*	*	*	0.3	0.2	0.1	*	*	*	*	*
1999 [4]	0.2	*	*	*	0.6	0.2	0.2	0.1	*	*	*	*
Female												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Discharge of firearms, undetermined intent (Y22-Y24)												
Non-Hispanic [5]												
Female												
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2001	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2000 [4]	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
1999 [4]	0.1	*	*	*	0.3	0.1	0.1	0.1	*	*	*	*
Male												
2001	0.1	*	*	*	0.3	0.2	0.2	*	*	*	*	*
2000 [4]	0.1	*	*	*	0.2	0.2	0.1	*	*	*	*	*
1999 [4]	0.2	*	*	*	0.5	0.2	0.2	*	*	*	*	*
Female												
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.2	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.2	*	*	*	0.6	*	*	*	*	*	*	*
Male												
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Discharge of firearms, undetermined intent (Y22-Y24)												
Non-Hispanic black												
Male												
1999 [4]	0.3	*	*	*	1.1	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2, Y89.9)												
All origins [3]												
Both sexes												
2001	1.4	2.1	0.3	0.1	0.9	1.7	3.0	2.5	0.8	0.5	0.8	1.0
2000 [4]	1.3	1.7	0.3	0.1	0.8	1.6	2.8	2.2	0.9	0.5	0.6	1.3
1999 [4]	1.3	1.4	0.3	0.1	0.8	1.7	2.7	2.1	0.9	0.5	0.7	1.3
Male												
2001	1.9	2.4	*	0.2	1.3	2.4	4.0	3.3	1.1	0.7	1.0	*
2000 [4]	1.8	1.4	0.3	0.2	1.1	2.2	3.8	3.0	1.2	0.7	0.9	*
1999 [4]	1.8	1.7	*	0.1	1.2	2.4	3.8	2.8	1.2	0.7	0.9	1.8
Female												
2001	0.9	1.7	0.3	*	0.5	1.1	2.0	1.6	0.6	0.3	0.6	0.8
2000 [4]	0.8	2.0	*	*	0.4	0.9	1.7	1.4	0.6	0.3	0.4	1.2
1999 [4]	0.8	1.2	0.3	*	0.4	1.1	1.7	1.4	0.6	0.4	0.5	1.1

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2, Y89.9)												
Hispanic												
Both sexes												
2001	0.7	*	*	*	0.4	0.9	1.6	1.5	*	*	*	*
2000 [4]	0.6	*	*	*	0.3	0.9	1.2	0.8	*	*	*	*
1999 [4]	0.6	*	*	*	0.5	1.0	1.2	0.9	*	*	*	*
Male												
2001	1.1	*	*	*	*	1.3	2.5	2.0	*	*	*	*
2000 [4]	0.8	*	*	*	*	1.3	1.7	1.3	*	*	*	*
1999 [4]	1.0	*	*	*	0.9	1.5	1.8	1.5	*	*	*	*
Female												
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	1.5	2.4	0.3	0.1	1.0	1.9	3.2	2.5	0.8	0.5	0.8	0.9
2000 [4]	1.4	1.9	0.3	0.1	0.9	1.7	2.9	2.3	0.9	0.5	0.6	1.3
1999 [4]	1.4	1.6	0.3	0.1	0.8	1.8	2.9	2.2	0.9	0.5	0.7	1.3
Male												
2001	2.0	2.7	*	0.2	1.4	2.6	4.2	3.4	1.1	0.6	1.0	*
2000 [4]	1.9	1.6	0.4	0.2	1.2	2.4	4.1	3.1	1.2	0.7	0.9	*
1999 [4]	1.9	1.9	*	0.1	1.2	2.5	4.0	2.9	1.2	0.7	0.9	1.7

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2, Y89.9)												
Non-Hispanic [5]												
Female												
2001	1.0	2.0	0.4	*	0.5	1.2	2.2	1.6	0.6	0.3	0.7	0.8
2000 [4]	0.9	2.2	*	*	0.4	1.0	1.8	1.5	0.6	0.3	0.4	1.2
1999 [4]	0.9	*	*	*	0.4	1.2	1.9	1.4	0.6	0.4	0.5	1.1
Non-Hispanic white												
Both sexes												
2001	1.5	1.7	0.3	0.1	1.1	2.0	3.2	2.5	0.8	0.4	0.8	0.9
2000 [4]	1.4	1.5	0.2	0.1	0.9	1.8	2.9	2.2	0.8	0.5	0.6	1.2
1999 [4]	1.3	1.3	*	0.1	0.9	1.9	2.8	2.0	0.9	0.5	0.7	1.3
Male												
2001	2.0	2.0	*	0.2	1.6	2.7	4.1	3.2	1.0	0.6	1.0	*
2000 [4]	1.8	*	*	0.2	1.4	2.4	4.0	2.9	1.0	0.6	0.9	*
1999 [4]	1.8	*	*	0.2	1.4	2.5	3.7	2.5	1.0	0.7	0.9	*
Female												
2001	1.1	*	*	*	0.6	1.3	2.3	1.7	0.7	0.3	0.6	0.8
2000 [4]	0.9	*	*	*	0.5	1.1	1.9	1.5	0.6	0.3	0.4	1.2
1999 [4]	0.9	*	*	*	0.4	1.3	1.9	1.5	0.7	0.4	0.6	1.1
Non-Hispanic black												
Both sexes												
2001	1.7	5.9	*	*	0.5	1.7	3.9	3.7	1.1	*	*	*
2000 [4]	1.7	*	*	*	0.6	1.7	4.0	3.7	1.6	*	*	*
1999 [4]	1.8	*	*	*	0.6	2.0	4.2	4.1	1.2	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2, Y89.9)												
Non-Hispanic black												
Male												
2001	2.6	6.4	*	*	0.8	2.5	6.3	6.2	2.1	*	*	*
2000 [4]	2.6	*	*	*	1.0	2.7	5.9	5.9	2.7	*	*	*
1999 [4]	2.8	*	*	*	0.9	3.2	6.4	7.1	2.5	*	*	*
Female												
2001	0.9	*	*	*	*	0.9	1.9	1.5	*	*	*	*
2000 [4]	0.9	*	*	*	*	0.8	2.3	1.8	*	*	*	*
1999 [4]	0.9	*	*	*	*	0.9	2.3	1.6	*	*	*	*
Operations of war and their sequelae (Y36,Y89.1)												
All origins [3]												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Operations of war and their sequelae (Y36,Y89.1)												
Hispanic												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Operations of war and their sequelae (Y36,Y89.1)												
Non-Hispanic [5]												
Female												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.0	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black												
Both sexes												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	*	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Operations of war and their sequelae (Y36,Y89.1)												
Non-Hispanic black												
Male												
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	*	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	*	*	*	*	*	*	*	*	*	*	*	*
Complications of medical and surgical care (Y40-Y84,Y88)												
All origins [3]												
Both sexes												
2001	1.1	0.6	0.2	*	0.1	0.2	0.4	0.7	1.5	3.1	6.9	12.8
2000 [4]	1.1	*	0.1	0.1	0.1	0.2	0.3	0.8	1.5	3.5	6.7	13.6
1999 [4]	1.0	1.4	0.2	*	0.1	0.2	0.3	0.6	1.3	3.2	6.6	12.7
Male												
2001	1.0	*	*	*	*	0.2	0.4	0.7	1.6	3.2	7.7	14.5
2000 [4]	1.0	*	*	*	*	0.2	0.3	0.8	1.6	4.0	8.2	16.0
1999 [4]	1.0	1.6	*	*	0.1	0.2	0.3	0.5	1.4	3.4	7.8	16.5
Female												
2001	1.1	*	*	*	0.1	0.2	0.4	0.7	1.4	3.0	6.3	12.1
2000 [4]	1.1	*	*	*	*	0.2	0.3	0.8	1.5	3.1	5.7	12.7
1999 [4]	1.1	1.2	*	*	*	0.2	0.3	0.6	1.2	3.0	5.9	11.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)												
Hispanic												
Both sexes												
2001	0.4	*	*	*	*	*	*	0.6	*	2.2	5.2	*
2000 [4]	0.4	*	*	*	*	*	*	*	*	3.3	5.1	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
Male												
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.4	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
Female												
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000 [4]	0.4	*	*	*	*	*	*	*	*	*	*	*
1999 [4]	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic [5]												
Both sexes												
2001	1.2	0.7	*	*	0.1	0.2	0.4	0.7	1.6	3.1	6.9	12.9
2000 [4]	1.2	*	*	0.1	0.1	0.2	0.3	0.8	1.6	3.5	6.7	13.7
1999 [4]	1.1	1.6	0.2	*	0.1	0.2	0.4	0.6	1.3	3.3	6.8	12.8
Male												
2001	1.1	*	*	*	*	0.2	0.4	0.8	1.7	3.3	7.7	14.5
2000 [4]	1.1	*	*	*	*	0.3	0.4	0.8	1.6	4.0	8.2	16.1
1999 [4]	1.0	1.8	*	*	*	0.2	0.4	0.6	1.5	3.5	7.9	16.8
Female												
2001	1.2	*	*	*	*	0.2	0.4	0.7	1.5	3.0	6.4	12.2

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)												
Non-Hispanic [5]												
Female												
2000 [4]	1.2	*	*	*	*	0.2	0.3	0.8	1.6	3.1	5.7	12.7
1999 [4]	1.2	1.4	*	*	*	0.2	0.4	0.6	1.2	3.1	6.1	11.2
Non-Hispanic white												
Both sexes												
2001	1.2	*	*	*	0.1	0.2	0.3	0.6	1.4	2.8	6.8	12.2
2000 [4]	1.2	*	*	*	*	0.2	0.3	0.8	1.4	3.3	6.5	13.2
1999 [4]	1.1	1.3	*	*	0.1	0.2	0.3	0.5	1.2	3.1	6.5	12.1
Male												
2001	1.0	*	*	*	*	0.2	0.3	0.6	1.4	2.9	7.8	13.5
2000 [4]	1.2	*	*	*	*	0.2	0.4	0.8	1.4	3.7	8.2	15.9
1999 [4]	1.1	*	*	*	*	0.2	0.3	0.5	1.3	3.3	7.7	15.8
Female												
2001	1.3	*	*	*	*	0.2	0.3	0.6	1.4	2.7	6.2	11.6
2000 [4]	1.2	*	*	*	*	0.2	0.3	0.7	1.4	3.0	5.4	12.0
1999 [4]	1.2	*	*	*	*	*	0.3	0.6	1.1	3.0	5.7	10.7
Non-Hispanic black												
Both sexes												
2001	1.4	*	*	*	*	0.4	0.9	1.6	3.4	6.4	9.3	23.7
2000 [4]	1.4	*	*	*	*	0.4	0.6	1.4	3.6	6.2	10.6	21.3
1999 [4]	1.3	*	*	*	*	*	0.6	1.3	2.6	5.2	10.6	20.4
Male												
2001	1.5	*	*	*	*	*	1.0	1.9	4.3	7.9	9.0	30.0

See footnotes at end of table.

Work table Orig290R. Deaths Rates for 113 Selected Causes, by 10-year Age Groups, Hispanic Origin, Race for Non-Hispanic Population, and Sex: United States, 1999-2001 - Con.

[Rates per 100,000 in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001"]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages [1]	Under 1 year [2]	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)												
Non-Hispanic black												
Male												
2000 [4]	1.2	*	*	*	*	*	*	1.3	3.2	7.8	10.2	*
1999 [4]	1.2	*	*	*	*	*	*	1.4	2.9	6.0	11.9	23.8
Female												
2001	1.4	*	*	*	*	*	0.7	1.4	2.7	5.3	9.4	21.4
2000 [4]	1.6	*	*	*	*	*	*	1.5	3.9	5.2	10.8	22.2
1999 [4]	1.4	*	*	*	*	*	*	1.2	2.4	4.7	9.8	19.1

* Figure does not meet standards of reliability or precision.
0.0 Quantity more than zero but less than 0.05.
... Category not applicable.
[1] Figures for age not stated included in "All ages" but not distributed among age groups.
[2] Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).
[3] All origins include origin not stated; specified origins exclude origin not stated.
[4] Rates are revised and may differ from rates previously published; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."
[5] Includes races other than white and black.
[6] Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

Source: CDC/NCHS, National Vital Statistics System, Mortality.