

Number of deaths and age-adjusted death rates with standard errors for poisonings involving opioid analgesics, by state: United States, 2006

	<i># of deaths</i>	<i>Age-adjusted rate</i>	<i>SE</i>
1. West Virginia	275	15.6	0.9
2. Utah	305	12.5	0.7
3. New Mexico	222	11.6	0.8
4. Oklahoma	372	10.8	0.6
5. Nevada	265	10.5	0.6
6. Vermont	56	9.1	1.2
7. Washington	550	8.2	0.4
8. Kentucky	310	7.4	0.4
9. Oregon	280	7.3	0.4
10. North Carolina	640	7.2	0.3
11. Tennessee	438	7.1	0.3
12. Rhode Island	75	6.9	0.8
13. Maine	86	6.7	0.7
14. Florida	1,107	6.3	0.2
15. Missouri	348	6.1	0.3
16. New Hampshire	79	5.9	0.7
17. Massachusetts	355	5.4	0.3
18. Wisconsin	301	5.4	0.3
19. Maryland	307	5.3	0.3
20. Arizona	300	5	0.3
21. Arkansas	132	4.9	0.4
22. Montana	44	4.8	0.7
23. South Carolina	207	4.8	0.3
24. Michigan	471	4.6	0.2
25. Illinois	562	4.4	0.2
26. Ohio	503	4.4	0.2
27. Idaho	61	4.3	0.6
28. Kansas	114	4.2	0.4
29. Colorado	206	4.1	0.3
30. Louisiana	167	4	0.3
31. Virginia	309	4	0.2
32. Texas	830	3.6	0.1
33. Delaware	29	3.5	0.7
34. Alaska	24	3.4	0.7
35. New York	667	3.4	0.1
36. Hawaii	43	3.3	0.5
37. Georgia	308	3.2	0.2
38. South Dakota	23	3.2	0.7
39. California	1,151	3.1	0.1
40. Connecticut	103	2.9	0.3
41. Iowa	83	2.9	0.3
42. Alabama	118	2.6	0.2
43. Minnesota	135	2.6	0.2
44. New Jersey	221	2.5	0.2
45. Pennsylvania	303	2.5	0.1
46. Indiana	138	2.2	0.2
47. Mississippi	61	2.2	0.3
48. Nebraska	31	1.8	0.3
49. North Dakota	4	*	*
50. Wyoming	12	*	*
District of Columbia	24	4.2	0.9
United States	13,755	4.6	0