



## Gonorrhea - Reported rates for 15-19 year old females by race and ethnicity: United States, 1981-1997

Rate (per 100,000 population)




# Gonorrhea - Reported rates for 15-19 year old males by race and ethnicity: United States, 1981-1997

Rate (per 100,000 population)




# Gonorrhea - Age- and gender-specific rates: United States, 1997




# Gonorrhea - Age-specific rates among women 10-44 years of age: United States 1981-1997

Rate (per 100,000 population)




## Gonorrhea - Age-specific rates among men 10-44 years of age: United States, 1981-1997

Rate (per 100,000 population)



## Gonorrhea - Rates in selected U.S. cities of >200,000 population, 1981-1997 and the Healthy People year 2000 objective


Rate (per 100,000 population)



Note: Atlanta, GA did not report gonorrhea statistics in 1994.

## Gonorrhea - Reported rates: United States, 1970-1997 and the Healthy People year 2000 objective

Rate (per 100,000 population)



Note: Georgia did not report gonorrhea statistics in 1994.


## Gonorrhea - Rates by gender: United States, 1981-1997 and the Healthy People year 2000 objective



Note: Georgia did not report gonorrhea statistics in 1994.

## Gonorrhea - Rates by region: United States, 1981-1997 and the Healthy People year 2000 objective


Rate (per 100,000 population)



Note: Georgia did not report gonorrhea statistics in 1994.


## Gonorrhea - Rates by race and ethnicity: United States, 1981-1997 and the Healthy People year 2000 objective

Rate (per 100,000 population)



Note: "Other" includes Asian/Pacific Islander and American Indian/Alaska Native populations.  
Georgia did not report gonorrhea statistics in 1994.

## Gonococcal Isolate Surveillance Project (GISP) - Percent of *Neisseria gonorrhoeae* isolates obtained from men who have sex with men for STD clinics in nine cities, 1988-1997



Note: These nine clinics reported 89.5% (394/440) of GISP gonorrhea cases in men who have sex with men. Orange County first participated in GISP in 1991 and Chicago first participated in 1996.