

Resistance to Isoniazid with or without Rifampin Resistance in Reported TB Cases with No Previous TB by Origin: United States, 1993-2003

Year	Resistance to Isoniazid ¹						Resistance to Isoniazid and Rifampin ¹					
	Total Cases ²		U.S.-born		Foreign-born ³		Total Cases ²		U.S.-born		Foreign-born ³	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
1993	1,400	8.4	804	6.8	580	12.4	410	2.5	302	2.6	105	2.3
1994	1,353	8.3	708	6.4	631	12.1	352	2.2	238	2.2	109	2.1
1995	1,173	7.3	554	5.4	618	11.0	253	1.6	168	1.6	85	1.5
1996	1,136	7.4	494	5.2	639	11.3	206	1.3	104	1.1	101	1.8
1997	1,079	7.5	436	5.0	639	11.2	155	1.1	76	0.9	79	1.4
1998	1,011	7.5	366	4.7	643	11.3	130	1.0	54	0.7	75	1.3
1999	899	7.1	283	4.0	614	11.0	127	1.0	39	0.6	88	1.6
2000	894	7.6	269	4.4	622	11.1	121	1.0	38	0.6	83	1.5
2001	797	7.0	241	4.4	555	9.5	113	1.0	34	0.6	79	1.4
2002	801	7.6	200	4.1	598	10.7	124	1.2	35	0.7	89	1.6
2003	784	7.9	204	4.6	577	10.6	90	0.9	25	0.6	65	1.2

¹Isolates may be resistant to other drugs.

²Includes persons of unknown country of birth.

³Includes persons born outside the United States, American Samoa, the Federated States of Micronesia, Guam, the Republic of the Marshall Islands, Midway Island, the Commonwealth of the Northern Mariana Islands, Puerto Rico, the Republic of Palau, the U.S. Virgin Islands, and U.S. minor and outlying Pacific islands.

Note: Data for all years updated through March 30, 2004.

More than 85% of all cases in each group had drug susceptibility test results reported for an initial isolate.