

**Resistance to Isoniazid with or without Rifampin Resistance in Reported TB Cases with Previous TB by Origin: United States, 1993-2003**

Year	Resistance to Isoniazid <sup>1</sup>						Resistance to Isoniazid and Rifampin <sup>1</sup>					
	Total Cases <sup>2</sup>		U.S.-born		Foreign-born <sup>3</sup>		Total Cases <sup>2</sup>		U.S.-born		Foreign-born <sup>3</sup>	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
1993	164	16.6	85	12.7	76	25.0	75	7.7	30	4.5	45	15.0
1994	177	17.1	81	11.7	95	28.1	75	7.3	35	5.1	39	11.6
1995	168	17.6	77	13.0	91	25.1	70	7.3	28	4.7	42	11.6
1996	142	16.5	67	12.1	74	24.4	43	5.0	20	3.6	22	7.3
1997	109	14.7	35	7.7	74	25.9	44	5.9	12	2.6	32	11.2
1998	98	13.1	38	7.8	60	22.8	23	3.1	6	1.2	17	6.5
1999	82	12.3	25	6.5	55	19.4	28	4.2	6	1.6	22	7.8
2000	82	13.0	22	6.1	60	22.2	24	3.8	2	0.6	22	8.2
2001	82	13.2	27	8.4	55	18.5	29	4.7	7	2.2	22	7.4
2002	77	14.1	23	7.9	54	21.3	24	4.4	3	1.0	21	8.3
2003	63	12.6	17	7.1	46	17.6	18	3.6	3	1.3	15	5.8

<sup>1</sup>Isolates may be resistant to other drugs.

<sup>2</sup>Includes persons of unknown country of birth.

<sup>3</sup>Includes persons born outside the United States, American Samoa, the Federated States of Micronesia, Guam, the Republic of the Marshall Islands, Midway Island, the Commonwealth of the Northern Mariana Islands, Puerto Rico, the Republic of Palau, the U.S. Virgin Islands, and U.S. minor and outlying Pacific islands.

**Note:** Data for all years updated through March 30, 2004.

More than 85% of all cases in each group had drug susceptibility test results reported for an initial isolate.