

**Table 1: Requirements for routine vaccination of immigrants examined overseas who are not fully vaccinated or lack documentation.**

Vaccine	Age						
	Birth–1 month	2–11 months	12 months–6 years	7–10 years	11–17 years	18–64 years	≥65 years
DTP/DTaP/DT	NO	YES		NO			
Td/Tdap	NO			YES, ≥7 years old (for Td); 10–64 years old (for Tdap)			
Polio (IPV/OPV)	NO	YES				NO	
MMR	NO		YES, if born in 1957 or later			NO	
Rotavirus	NO	YES 2–6 months old	NO				
Hib	NO	YES 2–59 months old		NO			
Hepatitis A	NO		YES 12–23 months old	NO			
Hepatitis B	YES, through 18 years old					NO	
Meningococcal (MCV/MPSV)	NO			Yes 11–18 years old		NO	
Human papillomavirus	NO			YES, for females 11–26 years old		NO	
Varicella	NO		YES				
Zoster	NO					YES, ≥60 years old	
Pneumococcal	NO	YES, 2–59 months old (for PCV)		NO			YES (for PPV)
Influenza	NO		YES, 6 months–18 years old (annually each flu season)			NO	YES, ≥50 years old (annually each flu season)

DTP=diphtheria and tetanus toxoids and pertussis vaccine; DTaP=diphtheria and tetanus toxoids and acellular pertussis vaccine; DT=pediatric formulation diphtheria and tetanus toxoids; Td=adult formulation tetanus and diphtheria toxoids; Tdap=adolescent and adult formulation tetanus and diphtheria toxoids and acellular pertussis vaccine (Boostrix for persons 10–18 years old; Adacel for persons 11–64 years old); IPV=inactivated poliovirus vaccine (killed); OPV=oral poliovirus vaccine (live); MMR=combined measles, mumps, rubella vaccine; Hib=*Haemophilus influenzae* type b conjugate vaccine MCV=meningococcal conjugate vaccine; MPSV=meningococcal polysaccharide vaccine; PCV=pneumococcal conjugate vaccine; PPV=pneumococcal polysaccharide vaccine.