

Table 6D
2010 Pediatric Nutrition Surveillance (1)
National

Comparison of Growth and Anemia Indicators by Contributor
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Contributor	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Alabama	39,278	12.0 (48)	4.6 (4)	150,224	6.1 (36)	6.4 (49)	12.0 (16)	65,914	15.2 (13)	14.1 (24)	103,190	17.7 (47)
Arizona	40,534	7.9 (12)	6.3 (22)	188,497	5.9 (29)	2.7 (7)	14.1 (39)	87,897	15.7 (20)	14.2 (28)	126,611	13.7 (33)
Arkansas	18,343	10.3 (44)	5.3 (7)	75,557	7.3 (49)	4.4 (31)	11.7 (11)	32,615	16.2 (27)	14.1 (24)	50,193	16.9 (45)
AZ I/Tribal Council	2,600	7.5 (8)	8.8 (44)	12,361	5.2 (17)	1.5 (2)	22.5 (54)	6,187	20.5 (51)	24.5 (54)	7,489	12.9 (26)
California	183,516	6.1 (3)	7.4 (38)	856,841	5.1 (16)	5.8 (45)	15.1 (47)	274,730	16.0 (25)	17.2 (47)	421,890	13.5 (32)
Cheyenne River-SD	152	5.9 (2)	10.5 (49)	735	1.1 (2)	1.8 (3)	20.8 (52)	382	21.5 (52)	19.6 (51)	519	11.8 (22)
Colorado	27,885	9.7 (34)	4.3 (2)	119,309	7.7 (51)	3.5 (21)	8.4 (2)	52,292	14.1 (5)	9.1 (2)	71,978	12.4 (25)
Connecticut	15,205	8.7 (19)	6.8 (31)	60,897	3.4 (5)	7.6 (54)	12.2 (19)	28,401	14.9 (9)	15.8 (44)	42,733	8.3 (8)
D.C.	5,014	11.2 (47)	5.3 (7)	16,863	7.6 (50)	4.7 (34)	12.1 (18)	6,954	14.9 (9)	13.7 (20)	10,845	26.7 (55)
Florida	127,001	9.8 (36)	5.8 (12)	539,151	4.3 (7)	5.0 (39)	12.9 (26)	242,399	14.8 (8)	13.4 (16)	368,681	15.8 (42)
Georgia	82,253	10.5 (45)	5.2 (6)	324,332	6.2 (37)	3.2 (15)	13.4 (33)	136,379	15.8 (22)	13.5 (17)	215,798	15.3 (38)

- (1) Reporting period is January 1 through December 31.
 - (2) Analyses based on one record per child.
 - (3) Excludes records with unknown data or errors.
 - (4) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.
 - (5) Infants born during the reporting period included in the analysis.
 - (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
 - (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
 - (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6D
2010 Pediatric Nutrition Surveillance (1)
National

Comparison of Growth and Anemia Indicators by Contributor
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Contributor	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Hawaii	8,908	9.1 (30)	6.2 (20)	38,503	6.3 (39)	5.4 (41)	9.6 (3)	17,827	12.5 (3)	9.1 (2)	19,139	10.9 (18)
Idaho	10,982	7.4 (7)	6.7 (30)	47,740	6.5 (42)	4.3 (28)	9.7 (4)	22,973	15.3 (16)	11.4 (5)	31,335	10.5 (15)
Illinois	83,599	9.0 (27)	6.8 (31)	296,662	6.7 (44)	5.7 (44)	12.5 (22)	135,408	15.3 (16)	14.6 (35)	173,953	11.1 (19)
Indiana	43,579	8.9 (26)	6.3 (22)	176,939	5.9 (29)	3.4 (18)	11.9 (15)	78,634	16.6 (36)	14.2 (28)	124,826	14.4 (36)
Iowa	17,441	8.8 (22)	7.6 (40)	77,729	5.5 (21)	3.0 (12)	13.5 (35)	35,783	17.2 (45)	14.7 (36)	53,886	7.5 (5)
Kansas	18,864	8.5 (14)	6.5 (27)	80,951	6.4 (41)	3.2 (15)	11.8 (13)	37,838	15.7 (20)	13.0 (14)	54,397	10.2 (13)
Kentucky	51,846	9.9 (39)	6.3 (22)	194,714	5.9 (29)	7.3 (52)	12.6 (23)	75,189	16.0 (25)	15.6 (42)	99,280	12.2 (23)
Louisiana	30,719	12.1 (49)	3.3 (1)	117,792	8.1 (53)	4.6 (32)	12.9 (26)	53,969	15.5 (18)	12.5 (11)	69,886	19.0 (48)
Maine	0	* *	* *	9,306	3.4 (5)	2.8 (8)	14.3 (41)	9,306	17.1 (43)	14.3 (31)	15,990	13.1 (28)
Maryland	35,324	9.9 (39)	5.9 (13)	139,804	5.6 (23)	3.2 (15)	13.4 (33)	63,951	16.5 (34)	15.7 (43)	95,654	20.0 (50)
Massachusetts	26,431	8.7 (19)	7.4 (38)	121,763	4.8 (10)	5.5 (42)	14.4 (43)	60,433	16.3 (30)	16.1 (45)	65,534	10.7 (17)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6D
2010 Pediatric Nutrition Surveillance (1)
National

Comparison of Growth and Anemia Indicators by Contributor
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Contributor	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Michigan	44,244	9.0 (27)	7.0 (33)	231,205	5.2 (17)	3.7 (25)	12.4 (20)	106,019	16.3 (30)	13.3 (15)	137,121	15.6 (41)
Minnesota	28,188	7.5 (8)	9.2 (46)	136,436	4.9 (11)	2.6 (4)	12.0 (16)	68,594	16.4 (33)	12.7 (12)	91,874	10.5 (15)
Mississippi	0	* *	* *	109,325	8.5 (54)	6.5 (50)	13.3 (32)	52,112	14.9 (9)	13.7 (20)	84,042	16.0 (43)
Missouri	40,289	9.5 (33)	6.1 (17)	163,309	6.2 (37)	4.6 (32)	11.8 (13)	67,547	16.2 (27)	13.6 (18)	97,640	16.5 (44)
Montana	4,653	9.0 (27)	7.1 (36)	18,318	6.8 (46)	3.9 (26)	11.0 (6)	8,958	15.9 (24)	12.2 (8)	10,993	9.9 (11)
Navajo Nation	2,654	7.2 (5)	6.4 (25)	11,437	5.0 (13)	1.3 (1)	17.2 (49)	4,687	20.2 (50)	18.7 (50)	6,478	5.2 (1)
Nebraska	11,518	8.2 (13)	7.0 (33)	49,201	5.9 (29)	3.9 (26)	12.8 (24)	22,194	16.7 (38)	13.8 (23)	30,945	13.2 (29)
Nevada	17,056	8.7 (19)	5.9 (13)	73,674	6.8 (46)	3.5 (21)	14.1 (39)	32,080	14.6 (7)	13.6 (18)	48,557	9.7 (10)
New Hampshire	0	* *	* *	18,403	5.8 (27)	2.8 (8)	13.1 (29)	8,621	17.1 (43)	14.2 (28)	10,515	13.7 (33)
New Jersey	42,335	8.6 (15)	6.1 (17)	176,464	5.6 (23)	4.7 (34)	15.7 (48)	78,181	17.0 (41)	17.3 (48)	118,144	19.9 (49)
New Mexico	15,242	9.4 (32)	4.4 (3)	59,579	6.8 (46)	6.0 (46)	10.6 (5)	31,043	13.7 (4)	11.7 (6)	41,953	11.5 (21)

- (1) Reporting period is January 1 through December 31.
- (2) Analyses based on one record per child.
- (3) Excludes records with unknown data or errors.
- (4) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.
- (5) Infants born during the reporting period included in the analysis.
- (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
- (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
- (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6D
2010 Pediatric Nutrition Surveillance (1)
National

Comparison of Growth and Anemia Indicators by Contributor
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Contributor	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)		≥ 2 Yrs Over-weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
New York	125,391	8.8 (22)	6.6 (29)	491,642	4.3 (7)	4.8 (36)	13.1 (29)	224,243	17.0 (41)	14.5 (33)	367,485	12.3 (24)
North Carolina	54,076	9.8 (36)	6.5 (27)	208,406	5.6 (23)	7.5 (53)	14.6 (46)	105,392	16.2 (27)	15.5 (39)	164,551	13.8 (35)
North Dakota	3,551	9.1 (30)	8.0 (41)	15,367	4.7 (9)	2.6 (4)	13.5 (35)	6,836	16.8 (39)	14.1 (24)	11,347	9.6 (9)
Ohio	68,637	10.2 (42)	5.9 (13)	296,250	6.5 (42)	3.6 (24)	12.4 (20)	128,754	15.8 (22)	12.4 (10)	212,805	13.0 (27)
Oregon	21,182	7.1 (4)	9.1 (45)	110,212	5.0 (13)	2.8 (8)	13.1 (29)	54,150	17.7 (47)	15.1 (37)	64,989	13.3 (31)
Pennsylvania	59,354	10.2 (42)	6.2 (20)	248,178	6.3 (39)	4.3 (28)	11.7 (11)	117,337	14.9 (9)	12.0 (7)	166,493	20.7 (53)
Puerto Rico	0	* *	* *	172,098	10.5 (55)	6.0 (46)	18.3 (50)	89,438	17.5 (46)	18.3 (49)	121,421	5.6 (2)
Rhode Island	5,711	8.8 (22)	7.2 (37)	25,794	5.6 (23)	3.5 (21)	14.3 (41)	12,922	16.3 (30)	15.5 (39)	17,483	15.3 (38)
Rosebud Sioux-SD	216	5.6 (1)	6.0 (16)	1,104	0.8 (1)	3.4 (18)	20.5 (51)	598	23.9 (53)	21.4 (52)	772	26.6 (54)
South Carolina	32,157	9.9 (39)	8.0 (41)	120,688	6.0 (34)	5.6 (43)	11.6 (10)	54,506	15.2 (13)	12.8 (13)	61,673	17.0 (46)
South Dakota	5,137	7.7 (11)	8.0 (41)	21,834	5.9 (29)	3.1 (13)	14.0 (38)	10,106	19.3 (49)	16.1 (45)	14,833	8.1 (7)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6D
2010 Pediatric Nutrition Surveillance (1)
National

Comparison of Growth and Anemia Indicators by Contributor
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Contributor	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)		≥ 2 Yrs Over-weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Standing Rock-ND	89	* *	* *	712	1.3 (3)	4.8 (36)	22.2 (53)	436	28.2 (55)	23.6 (53)	544	20.6 (51)
Tennessee	37,307	10.6 (46)	5.3 (7)	152,989	6.0 (34)	6.1 (48)	12.8 (24)	71,349	15.2 (13)	14.5 (33)	108,485	7.6 (6)
Texas	219,949	8.8 (22)	5.5 (10)	976,663	6.7 (44)	2.8 (8)	14.4 (43)	458,461	16.8 (39)	15.3 (38)	743,808	20.6 (51)
Three Aff. Tribes-ND	73	* *	* *	288	1.4 (4)	6.9 (51)	28.1 (55)	123	24.4 (54)	38.2 (55)	220	15.0 (37)
Utah	17,304	8.6 (15)	5.6 (11)	74,895	7.8 (52)	4.3 (28)	7.8 (1)	34,217	12.0 (2)	8.7 (1)	54,762	11.1 (19)
Vermont	2,306	7.2 (5)	9.9 (48)	13,037	4.9 (11)	3.4 (18)	11.1 (7)	6,921	15.6 (19)	12.2 (8)	10,360	6.7 (4)
Virgin Islands	1,169	8.6 (15)	4.6 (4)	5,308	5.3 (19)	7.7 (55)	11.2 (8)	2,587	11.8 (1)	11.2 (4)	2,934	13.2 (29)
Virginia	34,666	9.7 (34)	6.4 (25)	152,334	5.8 (27)	5.0 (39)	13.9 (37)	67,133	16.5 (34)	15.5 (39)	103,865	15.3 (38)
Washington	44,913	7.5 (8)	9.2 (46)	224,097	5.0 (13)	2.6 (4)	14.5 (45)	105,886	18.4 (48)	14.4 (32)	123,410	10.2 (13)
West Virginia	12,613	9.8 (36)	6.1 (17)	54,166	5.4 (20)	4.8 (36)	11.3 (9)	23,205	14.4 (6)	13.7 (20)	40,097	6.0 (3)
Wisconsin	30,092	8.6 (15)	7.0 (33)	130,168	5.5 (21)	3.1 (13)	13.0 (28)	59,975	16.6 (36)	14.1 (24)	96,621	10.0 (12)

- (1) Reporting period is January 1 through December 31.
 - (2) Analyses based on one record per child.
 - (3) Excludes records with unknown data or errors.
 - (4) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.
 - (5) Infants born during the reporting period included in the analysis.
 - (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
 - (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
 - (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6D
2010 Pediatric Nutrition Surveillance (1)
National
Comparison of Growth and Anemia Indicators by Contributor
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Contributor	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% >95th (Rank)	Number
Nation	1,851,546	9.0	6.3	8,190,251	5.9	4.5	13.4	3,640,072	16.1	14.4	5,391,027	14.6

- (1) Reporting period is January 1 through December 31.
 - (2) Analyses based on one record per child.
 - (3) Excludes records with unknown data or errors.
 - (4) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.
 - (5) Infants born during the reporting period included in the analysis.
 - (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
 - (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
 - (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

