

Table 6D
2008 Pediatric Nutrition Surveillance (1)
National

Comparison of Growth and Anemia Indicators by Contributor
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Contributor (3,4)													
Contributor	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)		≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Alabama	36,968	12.2 (50)	4.3 (3)	140,873	6.3 (31)	7.2 (50)	11.9 (8)	56,813	15.1 (8)	13.8 (17)		92,890	18.9 (46)
Arizona	44,377	8.0 (11)	6.6 (24)	181,654	6.7 (37)	2.7 (10)	15.8 (43)	75,338	16.0 (17)	14.6 (27)		118,807	14.5 (34)
Arkansas	25,714	9.7 (33)	5.8 (11)	88,127	7.8 (50)	4.7 (33)	11.5 (7)	38,591	15.9 (16)	13.9 (20)		55,337	17.6 (43)
AZ I/Tribal Council	2,711	7.5 (10)	8.3 (40)	11,521	5.5 (18)	1.5 (3)	23.6 (52)	5,823	21.1 (49)	23.5 (52)		6,980	13.4 (28)
California	233,891	6.7 (2)	7.6 (34)	989,591	5.1 (14)	5.6 (40)	15.5 (41)	301,643	16.0 (17)	17.3 (45)		466,528	14.3 (31)
Cheyenne River-SD	160	9.4 (29)	9.4 (44)	800	1.3 (3)	1.3 (2)	21.5 (49)	423	21.7 (51)	18.4 (48)		531	13.0 (26)
Colorado	28,282	9.8 (35)	4.2 (2)	106,342	7.9 (51)	3.5 (21)	8.2 (1)	43,476	14.4 (6)	9.4 (2)		58,958	11.9 (19)
Connecticut	16,024	9.5 (30)	7.0 (30)	59,088	3.6 (5)	7.6 (51)	12.3 (13)	25,623	15.7 (14)	15.5 (34)		40,618	9.1 (10)
D.C.	3,045	11.0 (48)	5.0 (5)	16,484	7.7 (49)	3.9 (27)	12.0 (10)	6,195	16.1 (21)	13.3 (9)		10,615	30.3 (52)
Florida	133,354	9.7 (33)	5.8 (11)	507,850	4.6 (7)	4.8 (34)	13.5 (27)	209,671	15.4 (11)	14.1 (24)		329,165	17.3 (41)
Georgia	91,632	10.3 (42)	5.2 (7)	326,413	6.9 (43)	3.1 (17)	14.2 (32)	124,533	16.6 (25)	14.8 (32)		209,382	14.3 (31)

- (1) Reporting period is January 1 through December 31.
- (2) Analyses based on one record per child.
- (3) Excludes records with unknown data or errors.
- (4) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.
- (5) Infants born during the reporting period included in the analysis.
- (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
- (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
- (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6D
2008 Pediatric Nutrition Surveillance (1)
National

Comparison of Growth and Anemia Indicators by Contributor
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Contributor (3,4)												
Contributor	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Hawaii	9,263	9.0 (24)	6.6 (24)	36,444	6.4 (33)	5.4 (38)	10.8 (3)	16,106	12.1 (2)	9.3 (1)	17,980	13.0 (26)
Idaho	11,643	7.4 (8)	7.0 (30)	45,513	6.8 (41)	4.1 (29)	10.1 (2)	20,081	16.0 (17)	12.3 (6)	31,097	12.6 (24)
Illinois	59,819	8.8 (21)	6.9 (29)	152,392	7.6 (47)	6.2 (45)	11.9 (8)	63,414	15.3 (10)	14.7 (30)	67,391	12.0 (20)
Indiana	45,066	9.6 (32)	6.2 (18)	163,731	6.5 (34)	3.3 (18)	12.0 (10)	66,499	16.6 (25)	14.5 (26)	111,604	15.1 (35)
Iowa	18,189	8.5 (18)	7.9 (38)	76,257	5.6 (20)	2.8 (13)	14.5 (35)	33,548	17.5 (38)	15.1 (33)	52,672	8.1 (5)
Kansas	19,418	8.1 (12)	6.4 (22)	76,976	6.7 (37)	2.8 (13)	12.4 (16)	34,352	16.7 (29)	13.3 (9)	51,194	11.4 (15)
Kentucky	42,701	9.8 (35)	6.1 (17)	164,667	6.0 (24)	6.6 (48)	13.1 (23)	62,832	16.6 (25)	15.7 (36)	99,273	12.3 (22)
Maryland	35,679	10.3 (42)	6.0 (15)	131,211	6.0 (24)	3.3 (18)	14.0 (30)	54,866	16.8 (31)	15.7 (36)	87,135	21.9 (50)
Massachusetts	31,082	8.4 (16)	7.7 (37)	127,966	4.6 (7)	5.4 (38)	14.7 (37)	59,297	16.8 (31)	16.7 (43)	69,872	11.7 (18)
Michigan	57,235	9.5 (30)	7.2 (32)	240,409	6.9 (43)	3.8 (26)	12.9 (22)	103,523	16.6 (25)	13.9 (20)	143,054	15.7 (38)
Minnesota	33,584	7.4 (8)	8.8 (42)	142,629	4.9 (10)	2.6 (7)	12.8 (21)	65,607	17.0 (34)	13.4 (13)	96,428	11.2 (14)

- (1) Reporting period is January 1 through December 31.
- (2) Analyses based on one record per child.
- (3) Excludes records with unknown data or errors.
- (4) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.
- (5) Infants born during the reporting period included in the analysis.
- (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
- (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
- (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6D
2008 Pediatric Nutrition Surveillance (1)
National

Comparison of Growth and Anemia Indicators by Contributor
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Contributor (3,4)												
Contributor	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Mississippi	0	* *	* *	108,893	10.2 (52)	6.5 (47)	14.5 (35)	44,807	15.5 (13)	14.6 (27)	78,003	16.5 (39)
Missouri	43,541	9.2 (28)	6.3 (20)	147,488	6.7 (37)	4.5 (31)	12.3 (13)	60,908	16.8 (31)	13.9 (20)	96,831	16.8 (40)
Montana	5,505	8.2 (13)	7.4 (33)	22,487	5.3 (17)	2.6 (7)	12.7 (19)	10,428	16.0 (17)	12.4 (7)	16,305	7.6 (4)
Navajo Nation	2,994	7.0 (4)	6.7 (26)	13,830	5.0 (11)	1.1 (1)	15.5 (41)	6,824	20.6 (48)	16.9 (44)	10,773	8.7 (8)
Nebraska	12,246	8.5 (18)	6.7 (26)	48,403	6.2 (28)	3.7 (24)	13.4 (26)	20,658	17.0 (34)	13.9 (20)	29,322	14.3 (31)
Nevada	17,010	8.2 (13)	5.9 (14)	61,796	6.7 (37)	3.5 (21)	14.3 (33)	23,348	13.8 (3)	12.9 (8)	37,035	10.1 (11)
New Hampshire	4,457	7.1 (5)	9.6 (47)	18,123	6.2 (28)	2.7 (10)	14.4 (34)	8,082	17.8 (42)	15.5 (34)	10,532	12.6 (24)
New Jersey	44,281	9.0 (24)	6.2 (18)	166,774	6.0 (24)	3.9 (27)	16.4 (45)	68,163	17.5 (38)	17.9 (46)	112,586	18.8 (45)
New Mexico	6,619	9.8 (35)	4.7 (4)	49,421	7.5 (46)	5.9 (43)	11.2 (4)	22,295	13.9 (4)	12.0 (4)	24,780	15.5 (37)
New York	130,568	8.9 (22)	6.7 (26)	481,925	4.2 (6)	4.4 (30)	13.8 (28)	209,713	17.4 (37)	14.6 (27)	355,319	12.2 (21)
North Carolina	61,064	9.8 (35)	6.5 (23)	202,638	5.6 (20)	7.7 (52)	15.1 (40)	96,381	16.3 (22)	15.7 (36)	154,344	12.5 (23)

- (1) Reporting period is January 1 through December 31.
- (2) Analyses based on one record per child.
- (3) Excludes records with unknown data or errors.
- (4) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.
- (5) Infants born during the reporting period included in the analysis.
- (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
- (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
- (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6D
2008 Pediatric Nutrition Surveillance (1)
National

Comparison of Growth and Anemia Indicators by Contributor
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Contributor (3,4)												
Contributor	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
North Dakota	3,607	9.0 (24)	9.2 (43)	15,187	4.7 (9)	2.5 (6)	13.8 (28)	6,551	17.5 (38)	13.8 (17)	11,132	8.7 (8)
Ohio Child Hlth	0	* *	* *	20,150	7.6 (47)	4.9 (35)	14.1 (31)	7,660	17.2 (36)	15.9 (39)	0	* *
Ohio-WIC	72,168	10.5 (45)	5.7 (10)	297,403	6.5 (34)	3.7 (24)	12.4 (16)	125,011	15.8 (15)	12.2 (5)	211,108	14.1 (30)
Oregon	22,459	6.7 (2)	9.5 (45)	105,686	5.0 (11)	2.7 (10)	13.3 (24)	49,193	17.6 (41)	14.7 (30)	59,280	13.4 (28)
Pennsylvania	67,539	10.1 (40)	6.3 (20)	256,027	6.3 (31)	5.1 (36)	11.4 (6)	111,767	14.3 (5)	11.5 (3)	174,229	20.9 (49)
Puerto Rico	37,208	11.4 (49)	2.7 (1)	191,404	10.3 (53)	6.6 (48)	18.4 (48)	99,828	19.8 (46)	17.9 (46)	143,159	5.6 (1)
Rhode Island	5,215	9.1 (27)	7.6 (34)	23,640	5.9 (23)	3.0 (15)	16.3 (44)	11,466	16.7 (29)	16.2 (40)	11,587	17.5 (42)
Rosebud Sioux-SD	277	6.1 (1)	9.7 (48)	1,298	1.2 (2)	4.6 (32)	17.9 (47)	651	23.0 (52)	19.2 (50)	970	27.5 (51)
South Carolina	0	* *	* *	83,487	7.4 (45)	6.4 (46)	11.3 (5)	28,209	15.2 (9)	13.3 (9)	41,319	18.3 (44)
South Dakota	5,160	7.3 (7)	8.0 (39)	21,386	6.1 (27)	3.0 (15)	14.7 (37)	9,125	19.7 (45)	16.2 (40)	15,318	8.4 (6)
Standing Rock-ND	167	8.4 (16)	12.6 (50)	983	1.7 (4)	5.8 (42)	23.4 (50)	541	24.2 (53)	25.0 (53)	766	18.9 (46)

- (1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6D
2008 Pediatric Nutrition Surveillance (1)
National

Comparison of Growth and Anemia Indicators by Contributor
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Contributor (3,4)												
Contributor	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Tennessee	22,047	10.1 (40)	5.0 (5)	104,228	6.2 (28)	5.9 (43)	12.7 (19)	49,016	15.4 (11)	13.8 (17)	63,859	8.5 (7)
Texas	233,919	8.9 (22)	5.6 (9)	320,907	6.8 (41)	2.3 (5)	16.6 (46)	164,435	17.9 (43)	16.2 (40)	690,010	20.1 (48)
Three Aff. Tribes-ND	115	10.4 (44)	8.7 (41)	386	1.0 (1)	5.7 (41)	24.1 (53)	163	21.5 (50)	19.6 (51)	293	11.6 (17)
Vermont	2,447	8.3 (15)	11.0 (49)	13,525	5.0 (11)	3.3 (18)	12.5 (18)	7,009	16.3 (22)	13.3 (9)	10,655	5.7 (2)
Virgin Islands	1,272	10.6 (46)	5.3 (8)	5,118	5.8 (22)	8.2 (53)	12.3 (13)	2,339	11.8 (1)	13.6 (15)	2,905	11.5 (16)
Virginia	35,908	10.0 (39)	6.0 (15)	146,916	6.5 (34)	1.9 (4)	23.5 (51)	59,627	20.1 (47)	19.0 (49)	96,629	15.3 (36)
Washington	46,619	7.2 (6)	9.5 (45)	209,245	5.1 (14)	2.6 (7)	14.9 (39)	92,980	18.6 (44)	14.4 (25)	108,318	10.7 (12)
West Virginia	13,431	10.7 (47)	5.8 (11)	54,539	5.2 (16)	5.2 (37)	12.0 (10)	22,689	14.8 (7)	13.5 (14)	40,026	5.7 (2)
Wisconsin	31,940	8.5 (18)	7.6 (34)	126,769	5.5 (18)	3.5 (21)	13.3 (24)	55,875	16.3 (22)	13.6 (15)	91,619	11.0 (13)
Nation	1,909,620	9.0	6.4	7,107,000	6.0	4.5	14.1	2,943,996	16.5	14.8	4,916,523	14.9

- (1) Reporting period is January 1 through December 31.
- (2) Analyses based on one record per child.
- (3) Excludes records with unknown data or errors.
- (4) Rank compares this contributor's rate to other contributors. Rank 1 = best rate.
- (5) Infants born during the reporting period included in the analysis.
- (6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
- (7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.
- (8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
- * Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.