

# Tripler Army Medical Center, Hawaii – 2003 (N=24)

Figure A. Age of GISP participants, in years, 2003




Figure B. Race/ethnicity of GISP participants, 2003





Figure C. Percentage of GISP participants identifying as men who have sex with men, 1988–2003

Data not collected

Figure D. Reason for visit among GISP participants, 2003



# Tripler Army Medical Center, Hawaii – 2003 (N=24)

Figure E. Previous episode of gonorrhea among GISP participants, 1991 – 2003

Data not available

Figure F. Drugs used to treat gonorrhea among GISP participants, 2003




Figure G. Drugs used to treat *Chlamydia trachomatis* infection among GISP participants, 2003





Figure H. Resistance to penicillin and tetracycline among GISP isolates, 2003



# Tripler Army Medical Center, Hawaii – 2003 (N=24)


Figure I. Decreased susceptibility to ceftriaxone among GISP isolates, 1988–2003

No isolates with decreased susceptibility to ceftriaxone have been identified at this clinic.

Figure J. Decreased susceptibility to cefixime among GISP isolates, 1992–2003

No isolates with decreased susceptibility to cefixime have been identified at this clinic.

Figure K. Intermediate resistance and resistance to ciprofloxacin among GISP isolates, 1990–2003



Note: Susceptibility to ciprofloxacin first measured in 1990.

Figure L. Decreased susceptibility to azithromycin among GISP isolates, 1992–2003

No isolates with decreased susceptibility to azithromycin have been identified at this clinic.