

Index

Index

A

Age group, *18, 23, 27-30, 38, 76*

Appendices

A, *119*

B, *120*

Alcohol use, *58*

B

B (appendix), *120*

C

Case verification criterion, *21*

Cities and metropolitan statistical areas, *71-80*

Areas with >500,000 population, *72-80*

Extrapulmonary, *74*

Pulmonary, *74*

Selected cities, *71*

Completion of therapy (COT), *23, 65, 67, 68*

Correctional facilities, *53*

Country of origin, *32*

D

Deaths, *15*

Death rates, *15*

Directly observed therapy (DOT), *23, 64*

District of Columbia, *50*

Drug susceptibility results, *60*

Drug use

Injecting drug use, *56*

Noninjecting drug use, *57*

E

Executive commentary, *3*

Extrapulmonary disease, *47, 48, 74*

F

Foreign-born persons

Adult, *20*

Cities and metropolitan statistical areas, *80*

Country of origin, *44*

States, *42, 44*

Top 30 countries of origin of birth, *19, 20*

Years in the U.S. before TB diagnosis, *20*

Morbidity, *46*

H

Health care provider (type), *63*

HIV

By age group, *23, 61*

Coinfection, *23*

Status, *61*

Test results, *23*

Hispanic, *16, 27-30, 40, 67, 78*

Cases by race, sex, age group, *27*

Foreign-born, *30*

U.S.-born, *29*

Case rates by race, sex, and age group, *28*

Reporting areas, *40*

Homeless status, *54*

I

INH resistance, *22, 60*

Initial drug regimen, *23, 59*

L

Long-term care facilities, *55*

M

Morbidity

Cities and metropolitan statistical areas, *71-80*

Foreign-born persons, *46*

Reporting areas, *53-68*

States, *37-50*

Trend, *15-24*

Multidrug resistance, *22, 60*

N

Non-Hispanic, *16, 27-30, 40, 67, 78*

Cases by race, sex, age group, *27*

Foreign-born, *30*

U.S.-born, *29*

Case rates by race, sex, and age group, *28*

Reporting areas, *40*

O

Occupation, *62*

Origin of birth, *18-20, 22*

Index

- P**
Persons with no previous history of TB, 22
Persons with previous history of TB, 22
Preface, *ix*
Previous statistical reports in this series, *xi*
Pulmonary disease, 21, 47, 74
- R**
Race, 17, 27-30
Reason TB therapy stopped, 24, 66
Reporting areas (morbidity), 38, 40, 47, 48, 53-68
 Completion of therapy (COT), 65, 67, 68
 Directly observed therapy (DOT), 64
 Drug susceptibility results, 60
 Excess alcohol use, 58
 HIV status, 61
 Homeless status, 54
 Initial drug regimen, 59
 Injecting drug use, 56
 Multidrug resistance, 60
 Noninjecting drug use, 57
 Occupation, 62
 Reason therapy stopped, 66
 Residence in correctional facilities, 53
 Residence in long-term care facilities, 55
 Resistance to INH, 60
 Type of health care provider, 63
Resistance to INH, 22, 60
- S**
Sex, 27-30
Site of disease, 21, 48
Sputum culture, 21
Sputum smear, 21
State TB statistics on the internet, *xii*
States (morbidity), 37-50
 Age group, 38
 Cases and case rates, 37
 Reporting areas, 38
Surveillance slide set, 85
 Slides, 85-110
 Text, *III*
- T**
Technical notes, 9
Trend (morbidity), 15-24
 Age group, 18, 23
 Case verification criterion, 21
 Completion of therapy, 23
 Death and death rates, 15
 Directly observed therapy, 23
 Foreign-born persons, 19-20
 HIV coinfection, 23
 HIV test results, 23
 Hispanic/Non-Hispanic, 16
 Multidrug resistance, 22
 No previous history of TB, 22
 Origin of birth, 18, 22
 Previous history of TB, 22
 Race, 17
 Reason TB therapy stopped, 24
 Resistance to INH, 22
 Site of disease, 21
 Sputum culture/sputum smear, 21
Type of health care provider, 63
- U**
U.S.-born
 Cities and metropolitan statistical areas, 80
 Hispanic/Non-Hispanic, 29
 States, 42