


Number of D.C. youth projected to die prematurely from their smoking: 4,927

Adult Tobacco Use in the District of Columbia

Current Cigarette Smoking Among Adults Aged 18 and Older, 1999



Youth Tobacco Use

	GRADES 6-8		GRADES 9-12	
	Current Cigarette Smoking	Current Any Tobacco Use	Current Cigarette Smoking	Current Any Tobacco Use
National*	9.2%	12.8%	28.5%	34.8%
DC†	Data are not available		19.9%	23.3%
Boys†			21.0%	25.9%
Girls†			19.0%	21.2%

Current Cigarette Smoking = smoked cigarettes on ≥1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥1 of the 30 days preceding the survey.

*Source: National Youth Tobacco Survey, 1999.

†Source: District of Columbia Youth Risk Behavior Survey, 1999.


Health Impact and Costs

AVERAGE ANNUAL DEATHS RELATED TO SMOKING, 1990-1994		AVERAGE ANNUAL YEARS OF POTENTIAL LIFE LOST,* 1990-1994	MEDICAL COSTS RELATED TO SMOKING, 1993	
Overall	929	15,184 years or an average of 16.3 years for each death due to smoking. *Calculated to life expectancy	Ambulatory	\$41,620,000
Men	591		Hospital	\$85,330,000
Women	339		Nursing Home†	\$35,160,000
Death Rate	327/100,000		Drug	\$4,250,000
Rank (No. 1 is lowest death rate)	14		Other	\$12,120,000
			Total	\$178,490,000

Smoking-attributable Medicaid expenditures, District of Columbia, fiscal year 1993: \$35,830,000

†Preliminary estimates

Lung Cancer Death Rate*



*1997 deaths per 100,000 population, adjusted to the 1970 total U.S. population.

Scheduled 2001 settlement payment to the District of Columbia: \$42,034,805.86

Tobacco Control Funding, 2001

At 13% of the CDC Best Practices recommended lower estimate funding level, the District of Columbia ranks 37th (No. 1 is the highest) for tobacco control funding. D.C. ranks 14th (No. 1 is the lowest death rate) for average annual deaths related to smoking.



<i>FUNDING SOURCE</i>	<i>FY01 AMOUNT</i>	<i>FUNDING CYCLE</i>
State Appropriation—Settlement (Tobacco Only)	\$0	
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$0	
Federal—CDC Office on Smoking and Health	\$448,157	6/00–5/01
Federal—SAMHSA	n/a	
Non-Government Source—American Legacy Foundation	\$504,962	10/00–9/01
Non-Government Source—RWJF/AMA	n/a	
Subtotal: Federal/National Sources	\$953,119	
Total Per Capita Funding	\$953,119 \$1.67	

Funding as a Percentage of CDC Best Practices Recommendations

	<i>RECOMMENDED FOR TOTAL PROGRAM ANNUAL COST</i>	<i>RECOMMENDED PER CAPITA FUNDING LEVEL</i>	<i>PERCENT OF CDC BEST PRACTICES RECOMMENDATIONS</i>
Lower Estimate	\$7,479,000	\$14.14	13%
Upper Estimate	\$14,571,000	\$27.55	7%

Excise Tax

Cigarette tax per pack	65¢
Rank = 13 (No. 1 is highest tax)	
Number of packages of cigarettes sold and taxed, per capita, 1999	50.8
Rank = 4 (No. 1 is lowest cigarette sales)	
Federal and state taxes as a percentage of retail price	27.5%
Annual gross tax revenue from cigarettes	\$17,420,000
Smokeless tobacco taxNone