

Dose and Route of Administration for Selected Vaccines

Vaccines	Dose	Route
Diphtheria, tetanus, pertussis (DTaP, DT, Td, Tdap)	0.5 mL	Intramuscular (IM)
Diphtheria, tetanus, acellular pertussis, inactivated polio, hepatitis B vaccine (DTaP-IPV-HepB)	0.5 mL	IM
Diphtheria, tetanus, acellular pertussis, <i>haemophilus influenzae</i> type b vaccine (DTaP-Hib)	0.5 mL	IM
<i>Haemophilus influenzae</i> type b (Hib)	0.5 mL	IM
<i>Haemophilus influenzae</i> type b, hepatitis B (Hib-HepB)	0.5 mL	IM
Hepatitis A (HepA)	<u><18 yrs:</u> 0.5 mL <u>≥19 yrs:</u> 1.0 mL	IM
HepB	<u><19 yrs:</u> 0.5 mL* <u>≥20 yrs:</u> 1.0 mL	IM
HepA-HepB	<u>≥18 yrs:</u> 1.0 mL	IM
Influenza, live attenuated	0.5 mL	Intranasal spray
Influenza, trivalent inactivated	6–25 mos: 0.25 mL <u>≥3 yrs:</u> 0.5 mL	IM
Measles, mumps, rubella	0.5 mL	Subcutaneous (SC)
Measles, mumps, rubella, varicella	0.5 mL	SC
Meningococcal conjugate	0.5 mL	IM
Meningococcal polysaccharide	0.5 mL	SC
Pneumococcal conjugate	0.5 mL	IM
Pneumococcal polysaccharide	0.5 mL	IM or SC
Human papillomavirus	0.5 mL	IM
Polio, inactivated	0.5 mL	IM or SC
Rotavirus	2.0 mL	Oral
Varicella	0.5 mL	SC
Zoster	0.65 mL	SC

*Persons aged 11-15 years can be administered Recombivax HB® (Merck) 1.0 mL (adult formulation) on a 2-dose schedule.
Adapted from: Immunization Action Coalition (<http://www.immunize.org>).