ADVISORY COMMITTEE ON IMMUNIZATION PRACTICES VACCINES FOR CHILDREN PROGRAM VACCINES TO PREVENT POLIOMYELITIS

The purpose of this resolution is to update the Recommended Vaccination Schedule and Intervals section to reflect the currently available inactivated polio vaccine (IPV) and IPV- containing vaccines that can be used to prevent polio.

VFC resolution 06/09-1 is repealed and replaced by the following:

Eligible groups

All infants and children at least six weeks of age.

Recommended Schedule and Intervals for Polio-containing Vaccines

The table below indicates single antigen and combination vaccines licensed and currently available for routine use to prevent polio.

Vaccine	Brand(s)	Schedule	Comments
IPV	Ipol	2,4,6-18 mos; 4-6 yrs	Approved for use in infants and
			adults
DTaP-HepB-IPV	Pediarix	2,4,6 mos	Approved for first 3 doses of
			IPV through 6 yrs
DTaP-IPV/Hib	Pentacel	2,4,6 and 15-18 mos	Approved for 4 doses of IPV
			through 4 yrs
DTaP-IPV-Hib-HepB	Vaxelis	2,4,6 mos	Approved for first 3 doses of
			IPV through 4 yrs
DTaP-IPV	Kinrix,	4-6 yrs	Approved for booster dose at
	Quadracel		age 4-6 yrs

Additional details can be found at:

https://www.cdc.gov/mmwr/preview/mmwrhtml/mm5830a3.htm?s_cid=mm5830a3_e*

Recommended dosage

Refer to product package inserts.

Contraindications and Precautions

Contraindications and Precautions can be found in the package inserts available at: https://www.fda.gov/vaccines-blood-biologics/vaccines/vaccines-licensed-use-united-states

[If an ACIP recommendation or notice regarding polio vaccination is published within 6 months following this resolution, the relevant language above (except in the eligible groups sections) will be replaced with the language in the recommendation and incorporated by reference to the publication URL.]

Adopted and Effective: June 26, 2019

This document can be found on the CDC website at:

 $\underline{https://www.cdc.gov/vaccines/programs/vfc/providers/resolutions.html}$