

Year	DTP3+	DTP4+	Polio3+	MMR*	Hib3+	Var	PCV3+	HepB3+	Combined 4-3-1	Combined 4-3-1-3
1990†										
1991	68.8		53.2	82.0						
1992	83.0	59.0	72.4	82.5	28.2			8.0	68.7	55.3
1993	88.2	72.1	78.9	84.1	55.0			16.3	67.1	
1994	93.0	77.7	83.0	89.0	86.0			37.0	75.0	
1995	94.7	78.5	87.9	87.6	91.7			68.0	76.2	74.2
1996	95.0	81.1	91.1	90.7	91.7	16.0		81.8	78.4	76.5
1997	95.5	81.5	90.8	90.5	92.7	25.9		83.7	77.9	76.2
1998	95.6	83.9	90.8	92.0	93.4	43.2		87.0	80.6	79.2
1999	95.9	83.3	89.6	91.5	93.5	57.5		88.1	79.9	78.4
2000	94.1	81.7	89.5	90.5	93.4	67.8		90.3	77.6	76.2
2001	94.3	82.1	89.4	91.4	93.0	76.3		88.9	78.6	77.2
2002	94.9	81.6	90.2	91.6	93.1	80.6	40.8	89.9	78.5	77.5
2003	96.0	84.8	91.6	93.0	93.9	84.8	68.1	92.4	82.2	81.3
2004	95.9	85.5	91.6	93.0	93.5	87.5	73.2	92.4	83.5	82.5
2005	96.1	85.7	91.7	91.5	93.9	87.9	82.8	92.9	83.1	82.4
2006	95.8	85.2	92.9	92.4	93.4	89.3	87.0	93.4	83.2	82.3
2007	95.5	84.5	92.6	92.3	92.6	90.0	90.0	92.7	82.8	81.1

*Previously reported as measles-containing vaccine (MCV).

†No national coverage data were collected from 1986 through 1990.

Var: varicella vaccine

Combined 4-3-1: Four or more doses of DTP/DTaP/DT, three or more doses of poliovirus vaccine, and one or more doses of any measles-containing vaccine.

Combined 4-3-1-3: Four or more doses of DTP/DTaP/DT, three or more doses of poliovirus vaccine, one or more doses of any measles-containing vaccine, and three or more doses of *Haemophilus influenzae* type b vaccine.

Data prior to 1993 were collected by the National Health Interview Survey and represent 2-year-old children. Data from 1993 are from the National Immunization Survey and represent 19-35 month-old children. Different methods were used for the two surveys.

Most recent publication: CDC. National, State, and Local Area Vaccination Coverage Among Children Aged 19-35 Months – United States, 2007. *MMWR* 2008;57(35);961-66.