

9. Grad YH, Kirkcaldy RD, Trees D, Dordel J, Harris SR, Goldstein E, et al. Genomic epidemiology of *Neisseria gonorrhoeae* with reduced susceptibility to cefixime in the USA: a retrospective observational study. *Lancet Infect Dis*. 2014; 14:220–6. [http://dx.doi.org/10.1016/S1473-3099\(13\)70693-5](http://dx.doi.org/10.1016/S1473-3099(13)70693-5)
10. Martin IM, Ison CA, Aanensen DM, Fenton KA, Spratt BG. Rapid sequence-based identification of gonococcal transmission clusters in a large metropolitan area. *J Infect Dis*. 2004;189:1497–505. <http://dx.doi.org/10.1086/383047>
11. Demczuk W, Sidhu S, Unemo M, Whiley DM, Allen VG, Dillon JR, et al. *Neisseria gonorrhoeae* sequence typing for antimicrobial resistance, a novel antimicrobial resistance multilocus typing scheme for tracking global dissemination of *N. gonorrhoeae* strains. *J Clin Microbiol*. 2017;55:1454–68. <http://dx.doi.org/10.1128/JCM.00100-17>

Address for correspondence: Yue-Ping Yin, National Center for STD Control, Chinese CDC, Institute of Dermatology, Chinese Academy of Medical Sciences and Peking Union Medical College, 12 Jiangwangmiao St, Nanjing 210042, China; email: yinyp@ncstdlc.org

***Chlamydia trachomatis* in Cervical Lymph Node of Man with Lymphogranuloma Venereum, Croatia, 2014¹**

Branimir Gjurašin, Snježana Židovec Lepej, Michelle J. Cole, Rachel Pitt, Josip Begovac

Author affiliations: University Hospital for Infectious Diseases Dr. Fran Mihaljević, Zagreb, Croatia (B. Gjurašin, S.Ž. Lepej, J. Begovac); Public Health England, London, UK (M.J. Cole, R. Pitt); University of Zagreb School of Medicine, Zagreb (J. Begovac)

DOI: <https://doi.org/10.3201/eid2404.171872>

We report an HIV-infected person who was treated for lymphogranuloma venereum cervical lymphadenopathy and proctitis in Croatia in 2014. Infection with a variant L2b genotype of *Chlamydia trachomatis* was detected in a cervical lymph node aspirate. A prolonged course of doxycycline was required to cure the infection.

¹Results from this study were presented as a poster at the IDWEEK 2017 Conference, October 4–8, 2017, San Diego, CA, USA. Abstracts of the IDWEEK 2017 Conference have been published in a supplement issue of *Open Forum Infectious Diseases* (<https://idsa.confex.com/idsa/2017/webprogram/POSTER.html>).

Lymphogranuloma venereum (LGV) is a sexually transmitted infection caused by serovars L1, L2, and L3 of the bacterium *Chlamydia trachomatis*. The infection typically causes genital ulcers, proctitis, or femoral/inguinal lymphadenopathy with or without constitutional symptoms. In the past decade, outbreaks of LGV have been reported in North America, Australia, and Europe, mainly as proctitis among HIV-infected men who have sex with men (MSM) (1). We report a patient with pharyngitis, proctitis, and cervical lymphadenitis in whom LGV-specific DNA was detected by real-time reverse transcription PCR (RT-PCR) in a cervical lymph node fine-needle aspirate.

The patient was a 48-year-old, HIV-positive man in Croatia who came to an outpatient HIV clinic in August 2014 with perianal pain for 10 days and bloody rectal discharge with normal stool consistency. He also reported a painful, enlarged cervical lymph node but did not have a sore throat. On the first day of the illness, he had fever, which subsided the next day. He reported having unprotected receptive anal and oral sex with other men while visiting Berlin, Germany, 2 weeks earlier. Clinical examination demonstrated exudate on the right tonsil, a painful and enlarged right cervical lymph node (5 × 2 cm) (online Technical Appendix Figure, <https://wwwnc.cdc.gov/EID/article/24/4/17-1872-Techapp1.pdf>), perianal pain on palpation, and a purulent rectal discharge.

The patient was given a diagnosis of HIV infection in 2002 and had been receiving antiretroviral therapy since July 2002. Plasma viremia had been undetectable since October 2002, and his CD4+ T-cell count before this illness was 2,082 cells/mm³. His clinical history included treatment for neurosyphilis, epilepsy, and diarrhea caused by *Microsporidiae* spp., *Blastocystis hominis*, and *Entamoeba histolytica*.

During examination at the HIV clinic, specimens were obtained from the pharynx, rectum, and urine for culture and a nucleic acid amplification test (NAAT). During fine-needle aspiration of a cervical lymph node, ≈1 mL of pus was removed and analyzed. The lymph node aspirate and a rectal swab specimen were positive for *C. trachomatis* DNA by the *C. trachomatis/Neisseria gonorrhoeae* RT-PCR (Abbott Laboratories, Abbott Park, IL, USA).

Cytologic examination of the fine-needle aspirate of the affected lymph node predominantly showed elements of granulomatous inflammation. An indirect immunofluorescence assay serum test result for *C. trachomatis* antibodies was positive (IgG titer >1:512, IgA titer 1:256). Test results for *N. gonorrhoeae* were negative (culture of the rectal swab and NAAT for urine and rectum). Results of a throat culture for *Streptococcus pyogenes* and routine lymph node aspirate culture for bacteria were also negative. Serum serologic test results were negative for acute infection with *Treponema pallidum*, *Bartonella* spp., and *Toxoplasma gondii*.

Table. Characteristics of 8 patients with lymphogranuloma venereum and cervical lymphadenopathy*

Reference	Patient age, y/sex	Clinical presentation	Method of laboratory confirmation	Therapy/duration
Andrada et al. (7)	30/M	Mouth ulcer, weight loss, cervical lymphadenopathy	Serologic analysis	Tetracycline/5 wk
Thorsteinsson et al. (3)	31/M	Fever, supraclavicular, axillar, retroperitoneal, and cervical lymphadenopathy	Serologic analysis	Tetracycline/4 wk
Watson et al. (6)	19/F	Sore throat, tonsillitis, arthritis, erythema nodosum, cervical lymphadenopathy	Serologic analysis	Phenoxymethylpenicillin, indomethacin, erythromycin†
Albay and Mathisen (5)	18/F	Fever, cervical lymphadenopathy	Serologic analysis	Ampicillin/sulbactam, doxycycline†
Tchernev et al. (8)	36/M	Facial skin lesions, cervical and axillary lymphadenopathy	NAAT: <i>Chlamydia trachomatis</i> DNA in skin lesions and serologic analysis	Surgical excision of cervical lymph nodes; pentamidine and doxycycline/3 wk
Dosekun et al. (4)	32/M	Sore throat, cervical lymphadenopathy, odynophagia, mouth ulcer, proctitis, cervical lymphadenopathy	NAAT: LGV-specific DNA in pharyngeal swab specimen	Amoxicillin/1 wk, doxycycline/1 wk
Dosekun et al. (4)	27/M	Sore throat, cervical lymphadenopathy, odynophagia, mouth ulcer, proctitis, cervical lymphadenopathy	NAAT: LGV-specific DNA in pharyngeal and rectal swab specimens	Azithromycin/1 g, doxycycline/2 wk
This study	48/M	Fever, cervical lymphadenopathy, proctitis	NAAT: LGV-specific DNA in cervical lymph node sample obtained by fine-needle aspirate; serologic analysis	Ceftriaxone/5 d, doxycycline/6 wk

*LGV, lymphogranuloma venereum; NAAT, nucleic acid amplification test.

†Duration of therapy not reported.

DNA from the lymph node specimen was frozen and sent to Public Health England (London, UK) in February 2017. LGV-specific DNA was detected by using an in-house TaqMan RT-PCR. The sequence of the outer membrane protein gene from lymph node punctate was identical to that of the *C. trachomatis* L2 reference strain L2/434/Bu.

At the initial visit, the patient was treated with intravenous ceftriaxone (2 g) and oral doxycycline (2 × 100 mg). After NAATs showed *C. trachomatis* infection, only doxycycline therapy was continued. Symptoms of proctitis subsided in 2 days. However, because cervical lymphadenitis persisted, we treated the patient with a prolonged course (6 weeks) of doxycycline. Eventually, the patient showed a full recovery.

Our report indicates that LGV might be present in MSM in Croatia. The first NAAT-confirmed case of LGV in southeastern Europe was reported in Slovenia and described an HIV-negative MSM who was ill in 2015 (2). LGV is probably underdiagnosed in southeastern Europe because of lack of diagnostics and awareness of the infection.

There have been only a few case reports of LGV with associated cervical lymphadenopathy (3–8) (Table). Some cases had generalized lymphadenopathy (axillar, supraclavicular, and retroperitoneal) with constitutional symptoms (3); pharyngitis/odynophagia/proctitis/tongue soreness (4,7); constitutional symptoms (5,7); tonsillitis (6); or skin lesions (8). Case reports have also been described of LGV with supraclavicular and mediastinal lymphadenopathy

without cervical involvement (9). In all of these cases, infection with LGV caused by *C. trachomatis* was established by serologic testing or an NAAT for a pharyngeal specimen. It is essential to maintain a high level of clinical suspicion for LGV in MSM even if noninguinal/femoral lymph nodes are affected.

The recommended treatment for LGV is doxycycline for 21 days. However, several clinical observations have suggested that a 21-day course of doxycycline therapy is insufficient for treating inguinal bubonic LGV (2,10). Recommendations have been given to carefully follow up with patients and continue doxycycline treatment until symptoms resolve (10). We followed these recommendations for our patient who had bubonic cervical lymph node LGV.

Acknowledgment

We thank the patient for providing permission to publish the case.

This study was partially supported by the Croatian Science Foundation (project no. IP-2014-09-4461) and the European Centre for Disease Prevention and Control (service contract no. ECD.6300).

Dr. Gjurašin is a fourth-year resident in infectious diseases at the University Hospital for Infectious Diseases Dr. Fran Mihaljević, Zagreb, Croatia. His primary research interests are infectious diseases of the central nervous system, sexually transmitted infections, zoonoses, and implementation of antimicrobial drug stewardship in Croatia.

References

1. de Vrieze NH, de Vries HJ. Lymphogranuloma venereum among men who have sex with men. An epidemiological and clinical review. *Expert Rev Anti Infect Ther*. 2014;12:697–704. <http://dx.doi.org/10.1586/14787210.2014.901169>
2. Maticič M, Klavs I, Videčnik Zorman J, Vidmar Vovko D, Kogoj R, Keše D. Confirmed inguinal lymphogranuloma venereum genovar L2c in a man who had sex with men, Slovenia, 2015. *Euro Surveill*. 2016;21:2–5. <http://dx.doi.org/10.2807/1560-7917.ES.2016.21.5.30129>
3. Thorsteinsson SB, Musher DM, Min KW, Gyorkey F. Lymphogranuloma venereum. A cause of cervical lymphadenopathy. *JAMA*. 1976;235:1882. <http://dx.doi.org/10.1001/jama.1976.03260430052029>
4. Dosekun O, Edmonds S, Stockwell S, French P, White JA. Lymphogranuloma venereum detected from the pharynx in four London men who have sex with men. *Int J STD AIDS*. 2013;24:495–6. <http://dx.doi.org/10.1177/0956462412472830>
5. Albay DT, Mathisen GE. Head and neck manifestations of lymphogranuloma venereum. *Ear Nose Throat J*. 2008;87:478–80.
6. Watson DJ, Parker AJ, Macleod TI. Lymphogranuloma venereum of the tonsil. *J Laryngol Otol*. 1990;104:331–2. <http://dx.doi.org/10.1017/S0022215100112629>
7. Andrada MT, Dhar JK, Wilde H. Oral lymphogranuloma venereum and cervical lymphadenopathy. Case report. *Mil Med*. 1974;139:99–101.
8. Tehernev G, Salaro C, Costa MC, Patterson JW, Nenoff P. Lymphogranuloma venereum: “a clinical and histopathological chameleon?” *An Bras Dermatol*. 2010;85:525–30. <http://dx.doi.org/10.1590/S0365-05962010000400015>
9. Bernstein DI, Hubbard T, Wenman WM, Johnson BL Jr, Holmes KK, Liebhaver H, et al. Mediastinal and supraclavicular lymphadenitis and pneumonitis due to *Chlamydia trachomatis* serovars L1 and L2. *N Engl J Med*. 1984;311:1543–6. <http://dx.doi.org/10.1056/NEJM198412133112405>
10. Oud EV, de Vrieze NH, de Meij A, de Vries HJ. Pitfalls in the diagnosis and management of inguinal lymphogranuloma venereum: important lessons from a case series. *Sex Transm Infect*. 2014;90:279–82. <http://dx.doi.org/10.1136/sextrans-2013-051427>

Address for correspondence: Branimir Gjurašin, University Hospital for Infectious Diseases Dr. Fran Mihaljević, Mirogojska Cesta 8, 10000 Zagreb, Croatia; email: brangjur@gmail.com

Letters

Letters commenting on recent articles as well as letters reporting cases, outbreaks, or original research are welcome. Letters commenting on articles should contain no more than 300 words and 5 references; they are more likely to be published if submitted within 4 weeks of the original article’s publication. Letters reporting cases, outbreaks, or original research should contain no more than 800 words and 10 references. They may have 1 Figure or Table and should not be divided into sections. All letters should contain material not previously published and include a word count.

Zika Virus MB16-23 in Mosquitoes, Miami-Dade County, Florida, USA, 2016

**John-Paul Mutebi, Holly R. Hughes,
Kristen L. Burkhalter, Linda Kothera,
Chalmers Vasquez, Joan L. Kenney**

Author affiliations: Centers for Disease Control and Prevention, Fort Collins, Colorado, USA (J.-P. Mutebi, H.R. Hughes, K.L. Burkhalter, L. Kothera, J.L. Kenney); Miami-Dade County Mosquito Control, Miami, Florida, USA (C. Vasquez)

DOI: <https://doi.org/10.3201/eid2404.171919>

We isolated a strain of Zika virus, MB16-23, from *Aedes aegypti* mosquitoes collected in Miami Beach, Florida, USA, on September 2, 2016. Phylogenetic analysis suggests that MB16-23 most likely originated from the Caribbean region.

In 2016, outbreaks of locally transmitted Zika virus occurred in Miami (Wynwood neighborhood) and Miami Beach, in Miami-Dade County, Florida, USA (1). During these outbreaks, a Centers for Disease Control and Prevention (CDC) emergency response team was deployed to assist Miami-Dade County disease surveillance and control efforts. CDC entomologists within the CDC emergency response team worked with Miami-Dade County Mosquito Control and sampled mosquito populations using BG-Sentinel type-2 traps (Biogents AG, Regensburg, Germany) to determine basic entomological parameters. Routinely, mosquitoes were collected, identified to species on the basis of the morphologic characteristics described by Darsie and Ward (2), and shipped inactivated and preserved in RNAlater (Ambion Inc., Austin, TX, USA) to the Bronson Animal Disease Diagnostic Laboratory (Kissimmee, FL, USA) for Zika virus testing.

In addition to the routine outbreak protocol, 2 BG-Sentinel type-2 traps were placed at a construction site near the intersection of James Avenue and Lincoln Road (25°47'25.68"N, 80°07'50.24"W) in Miami Beach on September 1, 2016. This site was selected because it was adjacent to a site where Zika cases had been detected. On September 2, 2016, the mosquitoes captured were frozen and shipped on dry ice to the CDC laboratory in Fort Collins, Colorado, USA. In the laboratory, the mosquitoes were identified to species on chill tables; female *Aedes aegypti* mosquitoes were separated into pools of 50 mosquitoes or less. A total of 293 female *Ae. aegypti* mosquitoes were collected (146.5/trap/day), grouped into 7 pools, and processed for presence of arboviral agents by cytopathic effect (CPE) assay.

We triturated pools of mosquitoes in 500 µL of Dulbecco’s modified Eagle medium complete with penicillin

Chlamydia trachomatis in Cervical Lymph Node of Man with Lymphogranuloma Venereum, Croatia 2014

Technical Appendix.



Technical Appendix Figure. Right cervical lymphadenopathy in 48-year-old man with lymphogranuloma venereum, Croatia 2014.