

***Toxoplasma gondii* Infections and Associated Factors in Female Children and Adolescents, Germany**

Laura Giese,¹ Frank Seeber,¹ Anton Aebischer, Ronny Kuhnert, Martin Schlaud, Klaus Stark, Hendrik Wilking

In a representative sample of female children and adolescents in Germany, *Toxoplasma gondii* seroprevalence was 6.3% (95% CI 4.7%–8.0%). With each year of life, the chance of being seropositive increased by 1.2, indicating a strong force of infection. Social status and municipality size were found to be associated with seropositivity.

Toxoplasmosis, caused by the protozoan parasite *Toxoplasma gondii*, is the most common parasitic foodborne disease in Germany. The seroprevalence in adults is exceptionally high (50%) compared with other countries (1). After infection via either undercooked meat from infected animals (pathway 1) or uptake of infectious oocysts shed by infected cats (pathway 2), *T. gondii* persists lifelong in infected persons, posing a risk for reactivation of latent infections (2). The proportion of pathways 1 and 2 leading to infection is largely unknown. We previously argued that in Germany, eating habits like consumption of raw pork products are likely responsible for the high seroprevalence (1). Infections in immune-competent persons remain largely asymptomatic or cause only mild, influenza-like symptoms. However, severe disease manifestations can occur, including ocular toxoplasmosis with sequelae, severe and often fatal consequences in immunocompromised persons, and congenital toxoplasmosis (2).

Worldwide, the World Health Organization (WHO) considered the burden of disease from toxoplasmosis to be high. Nevertheless, routine disease and pathogen surveillance is inadequate, so the incidence of human infection and parasite occurrence in animals and food is underestimated (3). In Germany, *T. gondii* screening during pregnancy to detect and treat primary infection, which could prevent parasite transmission to the unborn, is not covered by the

statutory health insurance. Ongoing discussions to reevaluate this policy demand data specifically assessing risk estimates for women of reproductive age.

In adults, age, dietary habits, and region of residence are associated with seroprevalence (1). Comparable analyses for children and adolescents are missing but are needed to estimate the public health problem and to suggest countermeasures. To provide such baseline data, this study aimed to estimate seroprevalence and to determine associated factors of *T. gondii* infections in female children and adolescents in Germany. The Hannover Medical School ethics committee approved the study (4).

The Study

The second wave of the nationwide German Health Interview and Examination Survey for Children and Adolescents (KiGGS Wave 2) was conducted as a 2-stage sampling survey during 2014–2017 in 167 representative sample points in Germany (4). Serum specimens of participants 3–17 years of age were tested for the presence of *T. gondii* IgG using a highly sensitive and specific commercial enzyme linked fluorescence assay (ELFA), as described previously (1,5).

We determined overall and stratified seroprevalence. We used data from standardized interviews to assess factors associated with seropositivity. On the basis of our previous serosurvey results in adults (1), we tested potential associations between seropositivity and vegetarianism, residence (eastern or western Germany), municipality size, and socioeconomic status. We identified minimal adjustment sets for multivariable logistic regression models by using directed acyclic graphs (Appendix, <https://wwwnc.cdc.gov/EID/article/30/5/23-1045-App1.pdf>). We determined adjusted odds ratios (aOR) for each exposure variable with 95% CIs. We used sampling weights

Author affiliation: Robert Koch Institute, Berlin, Germany

DOI: <http://doi.org/10.3201/eid3005.231045>

¹These authors contributed equally to this article.

Table. Stratified weighted seroprevalence of *Toxoplasma gondii* and results of weighted logistic regression analyses of factors potentially associated with seropositivity in female children and adolescents, Germany, 2014–2017*

Characteristics	No. positive/no. total†	Prevalence (95% CI)	Multivariable analysis aOR (95% CI)
Age group, y			
3–6	11/272	3.5 (1.3–5.8)	Referent
7–10	11/339	3.0 (0.8–5.1)	0.8 (0.3–2.3)
11–13	23/363	7.0 (3.5–10.4)	2.0 (0.9–4.9)
14–17	49/481	11.3 (6.7–15.8)	3.5 (1.6–7.4)
Socioeconomic status‡			
Low	20/211	10.8 (4.7–16.9)	2.7 (1.3–5.9)
Middle	51/868	4.7 (3.1–6.3)	Referent
High	19/325	6.1 (3.1–9.2)	1.5 (0.7–3.0)
Municipality size§			
<5,000	26/308	9.1 (4.4–13.7)	2.6 (1.1–5.7)
5,000 to <20,000	24/411	4.0 (1.9–6.1)	Referent
20,000 to <100,000	23/401	6.4 (2.8–9.9)	1.9 (0.8–4.5)
≥100,000	21/333	6.7 (3.9–9.5)	2.2 (1.1–4.4)
Residence¶			
East	57/954	6.3 (4.3–8.3)	Referent
West	37/499	6.5 (4.0–8.9)	1.1 (0.6–1.8)
Vegetarian			
Yes	7/129	6.4 (4.6–8.2)	1.3 (0.5–3.3)
No	86/1,273	7.4 (1.3–13.5)	Referent
Total	94/1,453	6.3 (4.7–8.0)	

*We tested 5 hypothesized associations toward seropositivity. Minimum adjustment sets were identified based on directed acyclic graphs (Appendix, <https://wwwnc.cdc.gov/EID/article/30/5/23-1045-App1.pdf>). aOR, adjusted odds ratio.

†Unweighted.

‡Based on the multidimensional socioeconomic index according to Lampert et al. (4).

§By number of inhabitants.

¶Western states: Baden-Württemberg, Bavaria, Bremen, Hamburg, Hesse, Lower Saxony, North Rhine-Westfalia, Rhineland-Palatinate, Saarland, Schleswig-Holstein; Eastern states: Berlin, Brandenburg, Mecklenburg-West Pomerania, Saxony, Saxony-Anhalt, Thuringia.

for all statistical analyses accounting for the study design. In addition, we calculated survey weights based on age, sex, residence, nationality, and education to correct for deviations from national population statistics.

We included 1,453 girls and adolescents (mean age 10.3 years) in the analyses. Of those, 1,359 tested

negative and 94 tested positive for *T. gondii*-specific IgG. We estimated overall weighted seroprevalence at 6.3% (95% CI 4.7%–8.0%) (Table).

With each year of life, the chance of being seropositive increased significantly, by 1.2 (95% CI 1.1–1.3). When we combined the data from the girls with those of female adults, seroprevalence steadily increased




Figure 1. Weighted seroprevalence of *Toxoplasma gondii* infections in female children and adolescents by age, Germany, 2014–2017 (red). For comparison, results of Wilking et al. (3), a previous study among adults, were added to the graph (black)




Figure 2. Weighted seroprevalence of *Toxoplasma gondii* infections in female children and adolescents by federal state, Germany, 2014–2017.

with age (Figure 1). This increase is a result of cumulative seropositivity because *T. gondii* infection is persistent and shows little seroreversion.

Girls living in families with low socioeconomic status (SES) showed the highest prevalence (10.8%, 95% CI 4.7%–16.9%). Their chance of being seropositive is 2.7 (95% CI 1.3–5.9) times higher than that of girls with middle SES. Low social status is often found to be associated with various disease risks, which may be a result of lower health literacy and reduced options to avoid health-related risks (4).

The seroprevalence among girls living in rural areas (<5,000 inhabitants) was significantly higher than seroprevalence for girls living in small towns (5,000–<20,000 inhabitants) (aOR 2.6, 95% CI 1.1–5.7). Similarly, girls living in urban areas (>100,000 inhabitants) were also 2.2 (95% CI 1.1–4.4) times more likely to be seropositive than those living in small towns. Greater exposure to natural habitats, including sand and soil contaminated due to free-roaming cats (6), as well as cats and children using the same limited spaces (e.g., sandboxes) (7), might explain the respective higher

risks of infection. We did not observe any regional distribution patterns (e.g., differences between eastern and western Germany) (Figure 2).

Overall, 9.3% of the participants reported being vegetarian; of those, 6.4% (95% CI 4.6%–8.2%) were seropositive (aOR 1.3, 95% CI 0.5–3.3). This result is not significantly different from results for nonvegetarians and is consistent with other studies (8–10). Diet appeared to have less effect on the risk for infection in children and adolescents than in adults. This finding may indicate that the relevance of the 2 transmission pathways differs significantly between age groups, and more environmentally associated infections occur in children and adolescents. Alternatively, risk factors associated with transmission pathway 1 may have shifted over time, for example, from improvements in the production and preparation of meat. However, more detailed information on the type (raw or undercooked) and quantity of meat consumed would have been beneficial but were not available in our dataset.

Conclusions

Overall, 6 of every 100 girls in Germany become infected with *T. gondii* during the first 18 years of life, corresponding to ≈340,504 total infections in this population group. Internationally comparable studies are limited. Seroprevalence estimates vary worldwide, from <10% to >60% for girls and 10%–80% for adults (11,12). Toxoplasmosis causes a higher infection pressure for girls and young women in Germany than in countries with similar socioeconomic conditions (11,13). In the United States, for example, the seroprevalence is significantly lower in age groups 6–11 years (0.9%, 95% CI 0.5%–1.5%) and 12–19 years (3.1%, 95% CI 2.0%–4.6%) (11).

Independent risk factors identified in our study were age, low social status, and growing up in rural or urban areas. Those factors have also been associated with seropositivity in children and adolescents in other countries (12,14). Further risk factors include contact with cats (12,14,15), contact with soil or sand (8,9,15), and consumption of unwashed vegetables (8). Growing up on a farm or keeping farm animals was also associated with increased seroconversions (8,10). Regular handwashing showed protective effects (9,15).

Our data may be helpful as an empirical basis for prevention guidelines. Implementing screening of pregnant women is one possibility and should be reevaluated using current data. Furthermore, our results indicate that meat consumption does not appear to be a driving force in children and adolescents, which calls for different prevention strategies in this

population than in adults (Appendix). However, our serosurvey is cross-sectional and represents the cumulated lifetime risk for infection. Therefore, misclassification of exposures (e.g., participants reported vegetarianism but consumed raw meat earlier in life) and unmeasured confounding is likely. Thus, our data are not appropriate for establishing causal relationships. Future studies should use longitudinal data containing detailed information on exposures and time of infection to disentangle different transmission pathways. The ultimate goal is efficient primary prevention of *T. gondii* infections; the goal requires integrating the fields of veterinary, human, and environmental medicine in a One Health approach.

Acknowledgments

We thank Daniela Heckmann, Petra Gosten-Heinrich, Elisabeth Kamal, Sandra Klein, Gudrun Kliem, Ilka McCormick-Smith, and Elke Radam for expert technical help.

This work was supported by the Robert Koch Institute, Berlin, Germany.

About the Author

Ms. Giese is an epidemiologist at the Robert Koch Institute with strong interest in gastrointestinal infections, zoonoses and tropical infections.

References


1. Wilking H, Thamm M, Stark K, Aebischer T, Seeber F. Prevalence, incidence estimations and risk factors of *Toxoplasma gondii* infection in Germany: a representative, cross-sectional, serological study. *Sci Rep*. 2016;6:22551. <https://doi.org/10.1038/srep22551>
2. Pleyer U, Gross U, Schlüter D, Wilking H, Seeber F. Toxoplasmosis in Germany. *Dtsch Arztebl Int*. 2019; 116:435–44.
3. EFSA Panel on Biological Hazards (BIOHAZ); Koutsoumanis K, Allende A, Alvarez-Ordóñez A, Bolton D, Bover-Cid S, Chemaly M, et al. Public health risks associated with food-borne parasites. *EFSA J*. 2018;16:5495.
4. Lampert T, Hoebel J, Kuntz B, Müters S, Kroll LE. Socioeconomic status and subjective social status measurement in KiGGS Wave 2. *J Health Monit*. 2018;3:108–25.
5. Murat JB, Dard C, Fricker Hidalgo H, Dardé ML, Brenier-Pinchart MP, Pelloux H. Comparison of the Vidas system and two recent fully automated assays for diagnosis and follow-up of toxoplasmosis in pregnant women and newborns. *Clin Vaccine Immunol*. 2013;20:1203–12. <https://doi.org/10.1128/CVI.00089-13>
6. Candela MG, Fanelli A, Carvalho J, Serrano E, Domenech G, Alonso F, et al. Urban landscape and infection risk in free-roaming cats. *Zoonoses Public Health*. 2022;69:295–311. <https://doi.org/10.1111/zph.12919>
7. Pacheco-Ortega GA, Chan-Pérez JI, Ortega-Pacheco A, Guzmán-Marín E, Edwards M, Brown MA, et al. Screening of zoonotic parasites in playground sandboxes of public parks

- from subtropical Mexico. *J Parasitol Res.* 2019;2019:7409076. <https://doi.org/10.1155/2019/7409076>
8. Hofhuis A, van Pelt W, van Duynhoven YTHP, Nijhuis CDM, Mollema L, van der Klis FRM, et al. Decreased prevalence and age-specific risk factors for *Toxoplasma gondii* IgG antibodies in the Netherlands between 1995/1996 and 2006/2007. *Epidemiol Infect.* 2011;139:530–8. <https://doi.org/10.1017/S0950268810001044>
 9. Sharif M, Daryani A, Barzegar G, Nasrolahei M. A seroepidemiological survey for toxoplasmosis among schoolchildren of Sari, Northern Iran. *Trop Biomed.* 2010;27:220–5.
 10. Nash JQ, Chissel S, Jones J, Warburton F, Verlander NQ. Risk factors for toxoplasmosis in pregnant women in Kent, United Kingdom. *Epidemiol Infect.* 2005;133:475–83. <https://doi.org/10.1017/S0950268804003620>
 11. Jones JL, Kruszon-Moran D, Elder S, Rivera HN, Press C, Montoya JG, et al. *Toxoplasma gondii* infection in the United States, 2011–2014. *Am J Trop Med Hyg.* 2018;98:551–7. <https://doi.org/10.4269/ajtmh.17-0677>
 12. Fan CK, Lee LW, Liao CW, Huang YC, Lee YL, Chang YT, et al. *Toxoplasma gondii* infection: relationship between seroprevalence and risk factors among primary schoolchildren in the capital areas of Democratic Republic of São Tomé and Príncipe, West Africa. *Parasit Vectors.* 2012;5:141. <https://doi.org/10.1186/1756-3305-5-141>
 13. Liassides M, Christodoulou V, Moschandreas J, Karagiannis C, Mitis G, Koliou M, et al. Toxoplasmosis in female high school students, pregnant women, and ruminants in Cyprus. *Trans R Soc Trop Med Hyg.* 2016;110:359–66. <https://doi.org/10.1093/trstmh/trw038>
 14. Cabral Monica T, Evers F, de Souza Lima Nino B, Pinto-Ferreira F, Breganó JW, Ragassi Urbano M, et al. Socioeconomic factors associated with infection by *Toxoplasma gondii* and *Toxocara canis* in children. *Transbound Emerg Dis.* 2022;69:1589–95. <https://doi.org/10.1111/tbed.14129>
 15. Wang S, Yao Z, Li H, Li P, Wang D, Zhang H, et al. Seroprevalence and risk factors of *Toxoplasma gondii* infection in primary school children in Henan province, central China. *Parasite.* 2020;27:23. <https://doi.org/10.1051/parasite/2020018>
- Address for correspondence: Laura Giese, Robert Koch Institute, Department of Infectious Disease Epidemiology, Seestraße 10, 13353 Berlin, Germany; email: giesel@rki.de

October 2023

Congenital Syphilis

- Serotype Distribution and Disease Severity in Adults Hospitalized with *Streptococcus pneumoniae* Infection, Bristol and Bath, UK, 2006–2022
- Spike in Congenital Syphilis, Mississippi, USA, 2016–2022
- Carbapenem-Resistant *Klebsiella pneumoniae* in Large Public Acute-Care Healthcare System, New York, New York, USA, 2016–2022
- Posttransfusion Sepsis Attributable to Bacterial Contamination in Platelet Collection Set Manufacturing Facility, United States
- Effects of COVID-19 on Maternal and Neonatal Outcomes and Access to Antenatal and Postnatal Care, Malawi
- Emergence of SARS-CoV-2 Delta Variant and Effect of Nonpharmaceutical Interventions, British Columbia, Canada
- Community Outbreak of *Pseudomonas aeruginosa* Infections Associated with Contaminated Piercing Aftercare Solution, Australia, 2021
- Stability of Monkeypox Virus in Body Fluids and Wastewater



- Cycle Threshold Values as Indication of Increasing SARS-CoV-2 New Variants, England, 2020–2022
- Comprehensive Case–Control Study of Protective and Risk Factors for Buruli Ulcer, Southeastern Australia
- Human-to-Human Transmission of Andes Virus Modeled in Syrian Hamsters

- *Candida auris* Clinical Isolates Associated with Outbreak in Neonatal Unit of Tertiary Academic Hospital, South Africa
- Ancestral Origin and Dissemination Dynamics of Reemerging Toxigenic *Vibrio cholerae*, Haiti
- *Treponema pallidum* Detection at Asymptomatic Oral, Anal, and Vaginal Sites in Adults Reporting Sexual Contact with Persons with Syphilis
- Managing Risk for Congenital Syphilis, Perth, Western Australia, Australia
- Human Tularemia Epididymo-Orchitis Caused by *Francisella tularensis* Subspecies *holartica*, Austria
- *Listeria monocytogenes* Transmission from Donated Blood to Platelet Transfusion Recipient, Italy
- Imported Toxigenic *Corynebacterium Diphtheriae* in Refugees with Polymicrobial Skin Infections, Germany, 2022
- Expansion of Invasive Group A *Streptococcus* M1UK Lineage in Active Bacterial Core Surveillance, United States, 2019–2021

**EMERGING
INFECTIOUS DISEASES**

To revisit the October 2023 issue, go to:
<https://wwwnc.cdc.gov/eid/articles/issue/29/10/table-of-contents>

Toxoplasma gondii Infections and Associated Factors in Female Children and Adolescents, Germany

Appendix

Additional Discussion: Practical recommendations for the prevention of *Toxoplasma gondii*

To effectively prevent toxoplasmosis, the various transmission routes and risk factors must be considered in a One Health approach combining the fields of human, animal and environmental health. There are currently three main opportunities for prevention:

Opportunity 1: Regarding congenital toxoplasmosis, regular testing during pregnancy is the only way to timely detect and treat infections in the unborn child. The implementation of mandatory screening programs during pregnancy should therefore be systematically evaluated. This includes the specificity and sensitivity of diagnostic tests and the consequences of a positive test result, including treatment options and effects (1).

Opportunity 2: Population groups that may be at a higher risk of primary infection and disease (e.g., pregnant women, people with low social status or people living in rural/urban areas) should be particularly addressed in health education programs. Although not apparent from our results, cat owners and raw meat consumers should also be targeted (2). The overall aim should be to increase the level of knowledge about toxoplasmosis, its transmission and prevention. Also, the use of new media should be considered to specifically address children and adolescents.


Practical recommendations should include how to reduce the risk of infection from the environment, for example by i. following hygiene measures when cleaning cat litter or when being in contact with sand or soil (e.g., handwashing or wearing gloves), ii. feeding cats with commercial food, iii. keeping cats away from public playgrounds or sand boxes. They should also include how to prevent foodborne infections, for example by i. avoiding the consumption of raw or undercooked meat or by ii. washing fruits and vegetables before consumption (3).

Opportunity 3: Measures to reduce the spread of *T. gondii* in livestock and the environment, for example through rodent control, monitoring of livestock and freshly produced meat or parasite inactivation, are discussed by experts of biologic hazards (4). These have to be evaluated and implemented in interdisciplinary cooperation between infectious disease experts and experts in environmental and veterinary health.


Additional References

1. Ahmed M, Sood A, Gupta J. Toxoplasmosis in pregnancy. Eur J Obstet Gynecol Reprod Biol. 2020;255:44–50. [PubMed https://doi.org/10.1016/j.ejogrb.2020.10.003](https://doi.org/10.1016/j.ejogrb.2020.10.003)
2. Wilking H, Thamm M, Stark K, Aebischer T, Seeber F. Prevalence, incidence estimations and risk factors of *Toxoplasma gondii* infection in Germany: a representative, cross-sectional, serological study. Sci Rep. 2016;6:22551. [PubMed https://doi.org/10.1038/srep22551](https://doi.org/10.1038/srep22551)
3. Centers for Disease Control and Prevention. Parasites – Toxoplasmosis (Toxoplasma infection). 2018 Sep 27 [cited 2024 Feb 14]. <https://www.cdc.gov/parasites/toxoplasmosis/>
4. EFSA Panel on Biological Hazards (BIOHAZ). Koutsoumanis K, Allende A, Alvarez-Ordóñez A, Bolton D, Bover-Cid S, Chemaly M, et al. Public health risks associated with food-borne parasites. EFSA J. 2018;16:5495.


Directed Acyclic Graphs




Appendix Figure 1. Directed Acyclic Graph (DAG) used to select the minimum adjustment set in the analysis of the hypothesized association between age and seropositivity in female children and adolescents in Germany. The graph shows correlations between variables included in our dataset based on current evidence. Green paths represent causal paths, red paths are biasing paths for which we have adjusted in our multivariable model. The DAG was created using Dagitty.net.




Appendix Figure 2. Directed Acyclic Graph (DAG) used to select the minimum adjustment set in the analysis of the hypothesized association between socioeconomic status (SES) and seropositivity in female children and adolescents in Germany. The graph shows correlations between variables included in our dataset based on current evidence. Green paths represent causal paths, red paths are biasing paths for which we have adjusted in our multivariable model. The DAG was created using Dagitty.net.



Appendix Figure 3. Directed Acyclic Graph (DAG) used to select the minimum adjustment set in the analysis of the hypothesized association between municipality size and seropositivity in female children and adolescents in Germany. The graph shows correlations between variables included in our dataset based on current evidence. Green paths represent causal paths, red paths are biasing paths for which we have adjusted in our multivariable model. The DAG was created using Dagitty.net.



Appendix Figure 4. Directed Acyclic Graph (DAG) used to select the minimum adjustment set in the analysis of the hypothesized association between East-West-Germany (East-West-GER) and seropositivity in female children and adolescents in Germany. The graph shows correlations between variables included in our dataset based on current evidence. Green paths represent causal paths, red paths are biasing paths for which we have adjusted in our multivariable model. The DAG was created using Dagitty.net.



Appendix Figure 5. Directed Acyclic Graph (DAG) used to select the minimum adjustment set in the analysis of the hypothesized association between Vegetarianism and seropositivity in female children and adolescents in Germany. The graph shows correlations between variables included in our dataset based on current evidence. Green paths represent causal paths, red paths are biasing paths for which we have adjusted in our multivariable model. The DAG was created using Dagitty.net.