



# National Health Interview Survey

## Current Asthma Among Adults Aged 18–64, by State

Percentages (and adjusted standard errors) of adults aged 18-64 years with current asthma, by state: United States, 2014-2016

States	Percent (adjusted standard error)
All states and the District of Columbia	8.0 (0.14)
Alabama	7.1 (1.27)
Alaska	8.3 (2.02)
Arizona	11.0 (1.53)
Arkansas	5.9 (1.19)
California	6.9 (0.38)
Colorado	8.7 (1.15)
Connecticut	9.8 (1.45)
Delaware	9.0 (1.48)
District of Columbia	8.0 (1.58)
Florida	6.1 (0.49)
Georgia	6.8 (0.64)
Hawaii	4.9 (0.91)
Idaho	7.7 (1.54)
Illinois	7.4 (0.69)
Indiana	8.5 (1.22)
Iowa	8.8 (1.65)
Kansas	9.6 (1.43)
Kentucky	10.8 (1.33)
Louisiana	6.2 (1.20)
Maine	10.8 (1.89)
Maryland	10.3 (1.72)
Massachusetts	9.8 (1.30)
Michigan	10.2 (0.89)
Minnesota	7.2 (0.90)
Mississippi	7.0 (1.11)
Missouri	8.1 (1.45)
Montana	7.6 (1.70)
Nebraska	7.5 (0.70)
Nevada	9.0 (1.36)
New Hampshire	12.7 (1.20)
New Jersey	7.4 (1.25)
New Mexico	10.4 (1.57)
New York	8.2 (0.64)
North Carolina	6.2 (0.74)
North Dakota	5.7 (0.95)
Ohio	9.6 (0.79)
Oklahoma	8.1 (1.27)
Oregon	9.5 (1.78)

See footnotes at the end of the table.

Percentages (and adjusted standard errors) of adults aged 18-64 with current asthma: United States, 2014-2016—Continued

States	Percent (adjusted standard error)
Pennsylvania	8.4 (0.69)
Rhode Island	9.3 (1.62)
South Carolina	6.2 (1.09)
South Dakota	7.3 (1.95)
Tennessee	7.7 (1.02)
Texas	7.4 (0.61)
Utah	9.1 (1.13)
Vermont	12.3 (4.50)
Virginia	9.0 (1.29)
Washington	7.8 (0.98)
West Virginia	9.8 (1.57)
Wisconsin	8.6 (1.18)
Wyoming	8.7 (0.43)

NOTES: Current asthma is based on a positive response to the survey questions "Have you ever been told by a doctor or other health professional that you had asthma?" and "Do you still have asthma?" Estimates are based on household interviews of a sample of the civilian, noninstitutionalized U.S. adult population and are shown for sample adults aged 18-64 years. All estimates and their variances were calculated using SUDAAN software to account for the complex sample design of NHIS. The standard errors were adjusted as described in Cohen RA, Makuc DM. State, regional, and national estimates of health insurance coverage for people under 65 years of age: National Health Interview Survey, 2004-2006. National Center for Health Statistics. November 2008. Available from: <https://www.cdc.gov/nchs/data/nhsr/nhsr001.pdf>.

SOURCE: NCHS, National Health Interview Survey, 2014-2016, Sample Adult component.

Suggested citation: Villarroel, MA, and Blackwell, DL. Current Asthma Among Adults Aged 18-64, by State: National Health Interview Survey, United States, 2014-2016. National Center for Health Statistics. May 2018. Available from: [https://www.cdc.gov/nchs/data/health\\_policy/asthma\\_tables\\_SEs.pdf](https://www.cdc.gov/nchs/data/health_policy/asthma_tables_SEs.pdf).