

Appendix Table I. Number and percent of persons uninsured, with public or private coverage, based on the 1990 and 2000 census population estimates, by age group: United States, 2000-2002

Health insurance coverage and age group	2000		2001		2002	
	1990-based ¹	2000-based ²	1990-based ¹	2000-based ²	1990-based ¹	2000-based ²
Number of uninsured in millions						
All ages	40.4	41.3	39.1	40.2	40.4	41.5
Under 65 years	39.9	40.8	38.7	39.8	40.0	41.1
18-64 years	31.2	32.0	30.9	31.9	32.4	33.5
Under 18 years	8.8	8.9	7.8	7.9	7.6	7.6
Percent (95% confidence interval)						
Uninsured						
All ages	14.7 (14.3-15.1)	14.9 (14.5-15.3)	14.1 (13.7-14.6)	14.3 (13.8-14.8)	14.5 (14.1-14.9)	14.7 (14.3-15.1)
Under 65 years						
Crude	16.6 (16.1-17.1)	16.8 (16.3-17.2)	15.9 (15.4-16.4)	16.2 (15.7-16.7)	16.3 (15.8-16.7)	16.5 (16.0-16.9)
Age-adjusted	16.6 (16.1-17.1)	16.8 (16.3-17.3)	16.0 (15.5-16.5)	16.2 (15.7-16.7)	16.4 (15.9-16.9)	16.6 (16.1-17.1)
18-64 years	18.4 (17.9-18.9)	18.7 (18.1-19.2)	18.1 (17.5-18.6)	18.3 (17.8-18.8)	18.8 (18.3-19.3)	19.1 (18.6-19.6)
Under 18 years	12.2 (11.6-12.8)	12.3 (11.7-12.9)	10.8 (10.1-11.5)	11.0 (10.3-11.7)	10.4 (9.8-11.0)	10.5 (9.9-11.1)
Public						
Under 65 years	12.9 (12.4-13.4)	12.9 (12.4-13.4)	13.6 (13.1-14.1)	13.6 (13.1-14.1)	15.2 (14.6-15.8)	15.2 (14.6-15.8)
18-64 years	9.0 (8.7-9.4)	9.1 (8.7-9.4)	9.4 (9.0-9.8)	9.4 (9.0-9.8)	10.2 (9.8-10.7)	10.3 (9.9-10.7)
Under 18 years	21.8 (20.8-22.8)	22.0 (21.0-23.0)	23.4 (22.4-24.3)	23.6 (22.6-24.5)	26.8 (25.7-27.9)	27.1 (26.0-28.2)
Private						
Under 65 years	72.0 (71.4-72.7)	71.8 (71.1-72.5)	71.9 (71.2-72.6)	71.6 (70.9-72.3)	70.1 (69.3-70.9)	69.8 (69.0-70.6)
18-64 years	74.0 (73.4-74.6)	73.8 (73.2-74.4)	73.9 (73.3-74.6)	73.7 (73.1-74.4)	72.6 (71.9-73.2)	72.3 (71.6-72.9)
Under 18 years	67.5 (66.4-68.5)	67.1 (66.1-68.2)	67.1 (66.0-68.2)	66.7 (66.4-68.6)	64.3 (63.1-65.5)	63.9 (62.7-65.1)

¹The estimates are calculated based on data with sample weights derived from the 1990 census.

²The estimates are calculated based on data with sample weights derived from the 2000 census.

Appendix Table II. Estimates of selected measures based on the 1990 and 2000 census population estimates: United States, 2000-2002

Indicator	2000		2001		2002	
	1990 based ¹	2000 based ²	1990 based ¹	2000 based ²	1990 based ¹	2000 based ²
Percent (95% confidence interval)						
With a usual place to go for medical care	87.1 (86.7-87.5)	87.0 (86.5-87.4)	88.2 (87.7-88.6)	88.0 (87.6-88.5)	87.8 (87.3-88.2)	87.6 (87.1-88.0)
Failed to obtain needed care due to cost	4.5 (4.3-4.7)	4.5 (4.3-4.7)	4.7 (4.5-4.9)	4.7 (4.5-4.9)	4.7 (4.5-4.9)	4.7 (4.5-4.9)
Had influenza vaccination in the past year	64.3 (62.9-65.7)	64.4 (63.0-65.9)	63.0 (61.6-64.4)	63.1 (61.7-64.5)	65.6 (64.1-67.0)	65.7 (64.3-67.2)
Had pneumococcal vaccination in the past year	52.9 (51.3-54.5)	53.1 (51.5-54.7)	53.8 (52.2-55.4)	54.0 (52.4-55.6)	55.7 (54.2-57.2)	56.0 (54.5-57.5)
With obesity	21.8 (21.2-22.4)	21.8 (21.2-22.4)	23.0 (22.4-23.5)	23.0 (22.4-23.6)	23.9 (23.3-24.6)	23.9 (23.3-24.6)
Engaged in regular leisure-time physical activity	32.0 (31.3-32.7)	31.9 (31.2-32.6)	32.1 (31.3-32.8)	32.1 (31.3-32.8)	32.0 (31.2-32.8)	31.9 (31.1-32.7)
Current smoking	23.3 (22.7-23.9)	23.2 (22.5-23.8)	22.8 (22.2-23.4)	22.7 (22.1-23.3)	22.5 (21.9-23.1)	22.4 (21.7-23.0)
Had 5 or more drinks in one day at least once in the past year	19.3 (18.7-20.0)	19.2 (18.6-19.9)	20.1 (19.5-20.7)	20.0 (19.4-20.6)	19.9 (19.3-20.6)	19.9 (19.2-20.5)
Ever had HIV test	32.2 (31.5-32.9)	32.1 (31.4-32.8)	33.3 (32.6-34.0)	33.2 (32.5-34.0)	34.5 (33.8-35.3)	34.5 (33.8-35.2)
With excellent or very good health status	68.4 (67.9-69.0)	68.3 (67.7-68.8)	68.6 (68.1-69.2)	68.4 (67.9-69.0)	67.6 (67.0-68.2)	67.4 (66.8-68.0)
With need for personal care	6.3 (5.9-6.8)	6.4 (5.9-6.9)	6.5 (6.0-7.0)	6.5 (6.0-7.0)	6.2 (5.6-6.7)	6.2 (5.7-6.7)
Serious psychological distress	2.7 (2.5-2.9)	2.7 (2.5-2.9)	3.2 (2.9-3.4)	3.2 (2.9-3.4)	3.0 (2.8-3.3)	3.0 (2.8-3.2)
Diagnosed Diabetes	5.9 (5.6-6.2)	5.9 (5.6-6.2)	6.4 (6.1-6.7)	6.4 (6.1-6.7)	6.5 (6.2-6.8)	6.5 (6.2-6.8)
Asthma episodes in the past year	40.1 (37.9-42.4)	40.0 (37.7-42.2)	43.4 (41.0-45.8)	43.2 (40.8-45.5)	42.8 (40.4-45.2)	42.6 (40.2-44.9)

¹The estimates are calculated based on data with sample weights derived from the 1990 census.

²The estimates are calculated based on data with sample weights derived from the 2000 census.

Appendix Table III. The NHIS Questions used to define selected health measures

Lack of health insurance and type of coverage¹

- FHI.010 {Are you/Is anyone} covered by any kind of health insurance or some other kind of health care plan?
- FHI.070 What kind of health insurance or health care coverage {do/does} {you/subject name} have? INCLUDE those that pay for only one type of service (nursing home care, accidents, or dental care), exclude private plans that only provide extra cash while hospitalized.

Usual place to go for medical care

- AAU.020 Is there a place that you USUALLY go to when you are sick or need advice about your health?
- AAU.030 What kind of place is it - a clinic, doctor's office, emergency room, or some other place?

Obtaining needed medical care

- AAU.111 DURING THE PAST 12 MONTHS, was there any time when {you/someone in the family} needed medical care, but did not get it because {you/the family} couldn't afford it?

Influenza vaccination

- AAU.310 DURING THE PAST 12 MONTHS, have you had a flu shot? A flu shot is usually given in the fall and protects against influenza for the flu season.

Pneumococcal vaccination

- AAU.320 Have you EVER had a pneumonia shot? This shot is usually given only once or twice in a person's lifetime and is different from the flu shot. It is also called the pneumococcal vaccine.

Obesity

- AHB.190 How tall are you without shoes?
- AHB.200 How much do you weigh without shoes?

Leisure-time physical activity

- AHB.110 How often do you do LIGHT OR MODERATE activities for AT LEAST 10 MINUTES that cause ONLY LIGHT sweating or a SLIGHT to MODERATE increase in breathing or heart rate?
- AHB.120 About how long do you do these light or moderate activities each time?
- AHB.090 How often do you do VIGOROUS activities for AT LEAST 10 MINUTES that cause HEAVY sweating or LARGE increases in breathing or heart rate?

AHB.100 About how long do you do these vigorous activities each time?

Current smoking

AHB.010 Have you smoked at least 100 cigarettes in your ENTIRE LIFE?

AHB.030 Do you NOW smoke cigarettes every day, some days or not at all?

Alcohol consumption

AHB.150 In your ENTIRE LIFE, have you had at least 12 drinks of any type of alcoholic beverage?

AHB.180 In the PAST YEAR, on how many DAYS did you have 5 or more drinks of any alcoholic beverage?

HIV testing

ADS.040 Except for tests you may have had as part of blood donations, have you ever been tested for HIV?

General health status

FHS.500 Would you say {subject name's} health in general is excellent, very good, good, fair, or poor?

Personal care needs

FHS.070 Because of a physical, mental, or emotional problem, {do/does} {you/anyone in the family} need the help of other persons with PERSONAL CARE NEEDS, such as eating, bathing, dressing, or getting around inside this home?

Serious psychological distress

ACN.471 During the PAST 30 DAYS, how often did you feel... ..So sad that nothing could cheer you up? ...Nervous? ...Restless or fidgety? ...Hopeless? ...That everything was an effort? ...Worthless?

Diabetes

ACN.160 Other than during pregnancy, have you EVER been told by a doctor or health professional that you have diabetes or sugar diabetes?

Asthma

ACN.090 DURING THE PAST 12 MONTHS, have you had an episode of asthma or an asthma attack?

¹Information from follow-up questions such as plan name(s) was used to reassign insurance status and type of coverage to avoid misclassification.