


**Figure 10.1. Percentage of adults aged 18 years and over who had ever been tested for human immunodeficiency virus (HIV): United States, 1997-June 2009**




NOTES: Individuals who received human immunodeficiency virus (HIV) testing solely as a result of blood donation were considered as not having been tested for HIV. The analyses excluded those with unknown HIV test status (about 5% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000-2002 were recalculated using weights derived from the 2000 census. See "About This Early Release" for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997-June 2009, Sample Adult Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For the period January through June 2009, the percentage of U.S. adults who had ever been tested for HIV was 39.8% (95% confidence interval = 38.15%-41.46%), which was higher than, but not significantly different from, the 2008 estimate of 38.7%.
- The percentage of adults who had ever had an HIV test increased from 30.2% in 1998 to 39.8% for the period January through June 2009.

**Figure 10.2. Percentage of adults aged 18 years and over who had ever been tested for human immunodeficiency virus (HIV), by age group and sex: United States, January-June 2009**




NOTES: Individuals who received human immunodeficiency virus (HIV) testing solely as a result of blood donation were considered as not having been tested for HIV. The analyses excluded 341 adults (3.3%) with unknown HIV test status.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January-June 2009, Sample Adult Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For both sexes combined, the percentage of persons who ever had an HIV test was highest among adults aged 25-34 years (59.2%) and 35-44 years (55.1%) and lowest among adults aged 65 and over (12.9%).
- For age groups 18-24 years, 25-34 years and 35-44 years, women were more likely than men to have ever had an HIV test.

**Figure 10.3. Age-sex-adjusted percentage of adults aged 18 years and over who had ever been tested for human immunodeficiency virus (HIV), by race/ethnicity: United States, January-June 2009**



NOTES: Individuals who received human immunodeficiency virus (HIV) testing solely as a result of blood donation were considered as not having been tested for HIV. The analyses excluded 341 adults (3.3%) with unknown HIV test status. Estimates are age-adjusted using the projected 2000 U.S. population as the standard population and using five age groups: 18-24 years, 25-34 years, 35-44 years, 45-64 years, and 65 years and over.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January-June 2009, Sample Adult Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- The age-sex-adjusted percentages of persons who ever had an HIV test were 43.9% for Hispanic persons, 37.0% for non-Hispanic white persons, and 58.3% for non-Hispanic black persons.
- Of the three race/ethnicity groups, non-Hispanic black persons were the most likely to have ever had an HIV test.

## Data tables for Figures 10.1-10.3:

### Data table for Figure 10.1. Percentage of adults aged 18 years and over who had ever been tested for human immunodeficiency virus (HIV): United States, 1997-June 2009

Year	Crude <sup>1</sup> percent (95% confidence interval)	Age-adjusted <sup>2</sup> percent (95% confidence interval)
1997	31.8 (31.1-32.5)	31.0 (30.4-31.6)
1998	30.2 (29.5-30.9)	29.6 (28.9-30.2)
1999	32.3 (31.6-33.0)	31.8 (31.1-32.5)
2000	32.1 (31.4-32.8)	31.8 (31.2-32.5)
2001	33.2 (32.5-34.0)	33.1 (32.4-33.7)
2002	34.5 (33.8-35.2)	34.5 (33.8-35.1)
2003	35.4 (34.7-36.1)	35.3 (34.7-36.0)
2004	34.6 (33.9-35.3)	34.8 (34.1-35.5)
2005	35.3 (34.58-36.01)	35.4 (34.73-36.10)
2006	35.8 (34.94-36.70)	36.1 (35.25-36.92)
2007	36.6 (35.59-37.62)	37.0 (35.28-38.79)
2008	38.7 (37.73-39.59)	39.2 (38.40-40.09)
Jan.-June 2009	39.8 (38.15-41.46)	40.3 (38.75-41.84)

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates for this *Healthy People 2010* Leading Health Indicator are age adjusted using the projected 2000 U.S. population as the standard population and using five age groups: 18-24 years, 25-34 years, 35-44 years, 45-64 years, and 65 years and over.

NOTES: Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000-2002 were recalculated using weights derived from the 2000 census. See "About This Early Release" for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997-June 2009, Sample Adult Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

**Data table for Figure 10.2. Percentage of adults aged 18 years and over who had ever been tested for human immunodeficiency virus (HIV), by age group and sex: United States, January-June 2009**

Age and sex	Percent	95% confidence interval
18-24 years, total	34.0	30.16-37.91
18-24 years, male	24.2	19.67-28.70
18-24 years, female	43.9	37.24-50.66
25-34 years, total	59.2	56.05-62.30
25-34 years, male	47.9	43.58-52.22
25-34 years, female	70.4	66.63-74.24
35-44 years, total	55.1	51.78-58.35
35-44 years, male	47.4	43.15-51.61
35-44 years, female	62.6	58.47-66.69
45-64 years, total	36.2	34.02-38.46
45-64 years, male	37.6	34.49-40.77
45-64 years, female	35.0	32.16-37.75
65 years and over, total	12.9	10.41-15.43
65 years and over, male	14.7	11.33-18.15
65 years and over, female	11.5	8.84-14.21
18-64 years, total	44.9	43.03-46.84
18-64 years, male	39.9	37.51-42.33
18-64 years, female	49.8	47.46-52.07
18 years and over (crude <sup>1</sup> ), total	39.8	38.15-41.46
18 years and over (crude <sup>1</sup> ), male	36.3	34.16-38.40
18 years and over (crude <sup>1</sup> ), female	43.1	41.09-45.06
18 years and over (age-adjusted <sup>2</sup> ), total	40.3	38.75-41.84
18 years and over (age-adjusted <sup>2</sup> ), male	36.0	34.04-37.99
18 years and over (age-adjusted <sup>2</sup> ), female	44.7	42.84-46.47

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age adjusted using the projected 2000 U.S. population as the standard population and using five age groups: 18-24 years, 25-34 years, 35-44 years, 45-64 years, and 65 years and over.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January-June 2009, Sample Adult Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.



**Data table for Figure 10.3. Age-sex-adjusted percentage of adults aged 18 years and over who had ever been tested for human immunodeficiency virus (HIV), by race/ethnicity: United States, January-June 2009**

<b>Race/ethnicity</b>	<b>Percent<sup>1</sup></b>	<b>95% confidence interval</b>
Hispanic or Latino	43.9	40.67-47.11
Not Hispanic or Latino, single race, white	37.0	35.17-38.76
Not Hispanic or Latino, single race, black	58.3	55.05-61.46

<sup>1</sup>Estimates are age-sex adjusted using the projected 2000 U.S. population as the standard population and using five age groups: 18-24 years, 25-34 years, 35-44 years, 45-64 years, and 65 years and over.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January-June 2009, Sample Adult Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.