


Figure 7.1. Percentage of adults aged 18 years and over who engaged in regular leisure-time physical activity: United States, 1997–June 2010




NOTES: This measure reflects the definition used for the physical activity Leading Health Indicator in *Healthy People 2010* (3). Regular leisure-time physical activity is defined as engaging in light-moderate leisure-time physical activity for greater than or equal to 30 minutes at a frequency greater than or equal to five times per week or engaging in vigorous leisure-time physical activity for greater than or equal to 20 minutes at a frequency greater than or equal to three times per week. In Early Releases before September 2005 (based on the 2004 National Health Interview Survey (NHIS)), regular physical activity was calculated slightly differently than for *Healthy People 2010*. The earlier Early Release estimates excluded from the analysis persons with unknown duration of light-moderate or vigorous leisure-time physical activity who were known to have not met the frequency recommendations for light-moderate or vigorous leisure-time physical activity (i.e., partial unknowns). With the current release, persons who were known to have not met the frequency recommendations are classified as "not regular," regardless of duration. All estimates have been rerun using the revised denominator. The impact of the change on the estimates was minimal (typically 0.1 percentage points or less). The analyses excluded persons with unknown physical activity participation (about 3% of respondents each year). Beginning with the 2003 data, NHIS transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. See "About This Early Release" for more details

- For January–June 2010, 34.4% (95% confidence interval = 33.26%–35.51%) of U.S. adults aged 18 years and over engaged in regular leisure-time physical activity, which was higher than, but not significantly different from, the January–June 2009 estimate of 33.6%.
- The annual percentages of adults aged 18 years and over who engaged in regular leisuretime physical activity ranged from 29.8% in 1998 to 34.7% in 2009.



Figure 7.2. Percentage of adults aged 18 years and over who engaged in regular leisure-time physical activity, by age group and sex: United States, January–June 2010




NOTES: This measure reflects the definition used for the physical activity Leading Health Indicator in *Healthy People 2010* (3). Regular leisure-time physical activity is defined as engaging in light-moderate leisure-time physical activity for greater than or equal to 30 minutes at a frequency greater than or equal to five times per week or engaging in vigorous leisure-time physical activity for greater than or equal to 20 minutes at a frequency greater than or equal to three times per week. In Early Releases before September 2005 (based on the 2004 National Health Interview Survey), regular physical activity was calculated slightly differently than for *Healthy People 2010*. The earlier Early Release estimates excluded from the analysis persons with unknown duration of light-moderate or vigorous leisure-time physical activity who were known to have not met the frequency recommendations for light-moderate or vigorous leisure-time physical activity (i.e., partial unknowns). With the current release, persons who were known to have not met the frequency recommendations are classified as "not regular," regardless of duration. The analyses excluded 261 persons (1.8%) with unknown physical activity participation.

- For both sexes combined, adults aged 75 years and over were less likely than adults in younger age groups to engage in regular leisure-time physical activity.
- For adults aged 18–24 years, 25–64 years, and 75 years and over, women were less likely than men to engage in regular leisure-time physical activity.



Figure 7.3. Age-sex-adjusted percentage of adults aged 18 years and over who engaged in regular leisure-time physical activity, by race/ethnicity: United States, January-June 2010



NOTES: This measure reflects the definition used for the physical activity Leading Health Indicator in *Healthy People 2010* (3). Regular leisure-time physical activity is defined as engaging in light-moderate leisure-time physical activity for greater than or equal to 30 minutes at a frequency greater than or equal to five times per week or engaging in vigorous leisure-time physical activity for greater than or equal to 20 minutes at a frequency greater than or equal to three times per week. In Early Releases before September 2005 (based on the 2004 National Health Interview Survey), regular physical activity was calculated slightly differently than for *Healthy People 2010*. The earlier Early Release estimates excluded from the analysis persons with unknown duration of light-moderate or vigorous leisure-time physical activity who were known to have not met the frequency recommendations for light-moderate or vigorous leisure-time physical activity (i.e., partial unknowns). With the current release, persons who were known to have not met the frequency recommendations are classified as "not regular," regardless of duration. The analyses excluded 261 persons (1.8%) with unknown physical activity participation. Estimates are age-sex-adjusted using the projected 2000 U.S. population as the standard population and using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

- The age-sex-adjusted percentage of adults who engaged in regular leisure-time physical activity was 27.1% for Hispanic adults, 37.3% for non-Hispanic white adults, and 28.4% for non-Hispanic black adults.
- Non-Hispanic white adults were more likely to engage in regular leisure-time physical activity than Hispanic adults and non-Hispanic black adults.



Data tables for Figures 7.1–7.3:

Data table for Figure 7.1. Percentage of adults aged 18 years and over who engaged in regular leisure-time physical activity: United States, 1997–June 2010

| Year | Crude ¹ percent (95% confidence interval) | Age-adjusted ² percent (95% confidence interval) |
|-------------------|---|---|
| 1997 yearly | 32.0 (31.3-32.6) | 31.8 (31.1-32.4) |
| 1997 January-June | 31.3 (30.1-32.6) | 31.1 (29.9-32.5) |
| 1998 yearly | 29.8 (29.1-30.4) | 29.6 (28.9-30.3) |
| 1998 January-June | 28.8 (27.4-30.1) | 28.5 (27.2-29.9) |
| 1999 yearly | 30.2 (29.5-31.0) | 30.1 (29.4-30.8) |
| 1999 January-June | 27.6 (26.2-29.1) | 27.4 (26.0-28.9) |
| 2000 yearly | 31.7 (31.0-32.4) | 31.7 (31.0-32.4) |
| 2000 January–June | 31.3 (29.9-32.7) | 31.2 (29.8-32.6) |
| 2001 yearly | 31.9 (31.2-32.7) | 31.8 (31.1-32.6) |
| 2001 January-June | 30.4 (29.1-31.8) | 30.3 (29.0-31.7) |
| 2002 yearly | 31.7 (31.0-32.5) | 31.7 (30.9-32.5) |
| 2002 January-June | 31.3 (29.9-32.8) | 31.3 (29.9-32.7) |
| 2003 yearly | 32.8 (32.1-33.6) | 32.8 (32.0-33.6) |
| 2003 January-June | 31.2 (29.7-32.6) | 31.1 (29.7-32.5) |
| 2004 yearly | 30.2 (29.4-31.0) | 30.1 (29.3-30.9) |
| 2004 January-June | 29.6 (28.2-31.0) | 29.5 (28.1-30.9) |
| 2005 yearly | 30.1 (29.38-30.92) | 30.1 (29.37-30.90) |
| 2005 January–June | 30.8 (29.20-32.39) | 30.7 (29.20-32.34) |
| 2006 yearly | 30.9 (29.91-31.80) | 30.9 (29.96-31.85) |
| 2006 January-June | 30.0 (28.57-31.46) | 30.0 (28.58-31.48) |
| 2007 yearly | 30.8 (29.73-31.82) | 30.8 (29.74-31.82) |
| 2007 January-June | 30.8 (29.13-32.50) | 30.7 (29.02-32.40) |
| 2008 yearly | 31.9 (30.92-32.97) | 32.0 (30.95-33.01) |
| 2008 January-June | 29.5 (27.96-31.07) | 29.7 (28.15-31.16) |
| 2009 yearly | 34.7 (33.73-35.58) | 34.8 (33.89-35.76) |
| 2009 January-June | 33.6 (31.52-35.76) | 33.8 (31.59-35.92) |
| 2010 January-June | 34.4 (33.26-35.51) | 34.5 (33.36-35.65) |

¹Crude estimates are presented in the figure.

NOTES: Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. See "About This Early Release" for more details.

²Estimates for this *Healthy People 2010* Leading Health Indicator are age-adjusted using the projected 2000 U.S. population as the standard population and using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.



Data table for Figure 7.2. Percentage of adults aged 18 years and over who engaged in regular leisure-time physical activity, by age group and sex: United States, January–June 2010

| Age and sex | Percent | 95% confidence interval |
|--|---------|-------------------------|
| 18-24 years, total | 42.2 | 38.65-45.77 |
| 18-24 years, male | 51.4 | 46.63-56.18 |
| 18-24 years, female | 32.9 | 28.83-36.88 |
| 25-64 years, total | 35.9 | 34.60-37.15 |
| 25-64 years, male | 37.5 | 35.72-39.29 |
| 25-64 years, female | 34.3 | 32.68-35.92 |
| 65-74 years, total | 26.3 | 23.62-29.03 |
| 65-74 years, male | 30.8 | 26.82-34.81 |
| 65-74 years, female | 22.5 | 18.94-25.97 |
| 75 years and over, total | 17.1 | 14.72-19.41 |
| 75 years and over, male | 23.7 | 19.32-28.04 |
| 75 years and over, female | 12.7 | 10.36-15.14 |
| 18 years and over (crude ¹), total | 34.4 | 33.26-35.51 |
| 18 years and over (crude ¹), male | 37.9 | 36.37-39.48 |
| 18 years and over (crude ¹), female | 31.1 | 29.76-32.39 |
| 18 years and over (age-adjusted ²), total | 34.5 | 33.36-35.65 |
| 18 years and over (age-adjusted ²), male | 37.7 | 36.16-39.26 |
| 18 years and over (age-adjusted ²), female | 31.5 | 30.19-32.89 |

¹Crude estimates are presented in the figure.

²Estimates for this *Healthy People 2010* Leading Health Indicator are age-adjusted using the projected 2000 U.S. population as the standard population and using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.



Data table for Figure 7.3. Age-sex-adjusted percentage of adults aged 18 years and over who engaged in regular leisure-time physical activity, by race/ethnicity: United States, January-June 2010

| Race/ethnicity | Age-sex-adjusted ¹ percent (95% confidence interval) | Age-adjusted ² percent (95% confidence interval) |
|--|---|--|
| Hispanic or Latino | 27.1 (25.09-29.06) | 27.2 (25.13-29.19) |
| Not Hispanic or Latino, single race, white | 37.3 (35.84-38.76) | 37.3 (35.89-38.81) |
| Not Hispanic or Latino, single race, black | 28.4 (25.99-30.71) | 28.1 (25.68-30.46) |

¹Age-sex-adjusted estimates are presented in the figure. Estimates are age-sex-adjusted using the projected 2000 U.S. population as the standard population and using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

²Estimates for this *Healthy People 2010* Leading Health Indicator are age-adjusted using the projected 2000 U.S. population as the standard population and using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over