


Figure 4.1. Percentage of adults aged 18 years and over who had received an influenza vaccination during the past 12 months, by age group and quarter: United States, 1997–September 2010



NOTES: NHIS questions related to influenza vaccination have changed since 1997, as follows. Starting in 1997, respondents were asked if they had received a flu shot during the past 12 months. Beginning in 2003, respondents were also asked if they had received a flu vaccine sprayed in their nose during the past 12 months. In August 2010, NHIS influenza vaccination questions were modified to reflect that for the first time, the widely available influenza vaccine included protection against both seasonal and H1N1 types of influenza. NHIS Early Release influenza vaccination estimates have changed since 1997, as follows. Starting in 1997, Early Release influenza vaccination estimates covered receipt of an influenza shot only. Starting in 2005, Early Release influenza vaccination estimates covered seasonal influenza shot and/or seasonal intranasal influenza vaccination. When interpreting influenza vaccination estimates, readers should take into account changes made to the influenza vaccination questions noted above. An error in calculating influenza vaccination rates occurred for the first quarter of 2005 to the first quarter of 2007. The effect of this error on estimates was small. Compared with the original estimates, corrected estimates are slightly higher, usually by no more than 0.3 percentage point. The error has been corrected for all estimates in this Early Release, and the correction of estimates had no perceptible impact on the graphs. Responses to the previously mentioned influenza vaccination questions cannot be used to determine when during the preceding 12 months the subject received the influenza vaccination. In addition, estimates are subject to recall error, which will vary depending on when the question is asked because the receipt of an influenza vaccination is seasonal. The recommendations of the Advisory Committee on Immunization Practices regarding who should receive an influenza vaccination have changed over the years and changes in coverage estimates may reflect changes in recommendations (11–13). Influenza vaccination shortages have occurred during several influenza seasons (12–14). The analyses excluded those with unknown influenza vaccination status (about 3% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. See "About This Early Release" for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997–September 2010, Sample Adult Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.



- In the third quarter of 2010, the percentage of adults who had received an influenza vaccination during the past 12 months was 63.1% for persons aged 65 years and over, 37.9% for persons aged 50–64, and 26.7% for persons aged 18–49.
- For the age group 18–49 years, the third-quarter estimate in 2010 was higher than the third-quarter estimate in 2009. For the age group 50–64, the third-quarter estimate from 2010 was not significantly different from the third-quarter estimate from 2009. For the age group 65 and over, the third-quarter estimate from 2010 was lower than, but not significantly different from the third-quarter estimate from 2009. For all three age groups, third-quarter estimates increased from 2005 to 2010. An influenza vaccination shortage occurred during the 2004–2005 influenza season (13). Delays in the availability of influenza shots also occurred in fall 2000 and, to a lesser extent, in fall 2001 (11,13).

Table 4.1a. Annual percentage of adults aged 50–64 years who had received an influenza vaccination during the past 12 months, by sex: United States, 1997–2009

Year	Percent (95% confidence interval): total	Percent (95% confidence interval): men	Percent (95% confidence interval): women
1997	31.9 (30.5-33.3)	28.0 (26.1-29.9)	35.5 (33.6-37.4)
1998	33.1 (31.7-34.5)	29.0 (27.0-31.0)	37.0 (35.1-38.9)
1999	34.1 (32.8-35.4)	30.5 (28.6-32.4)	37.4 (35.5-39.3)
2000	34.6 (33.1-36.1)	31.9 (29.9-33.9)	37.2 (35.2-39.1)
2001	32.2 (30.9-33.5)	30.3 (28.3-32.2)	34.0 (32.2-35.8)
2002	34.0 (32.7-35.3)	30.7 (28.8-32.5)	37.2 (35.4-38.9)
2003	36.8 (35.4-38.2)	34.5 (32.6-36.3)	38.9 (37.0-40.9)
2004	35.9 (34.6-37.3)	33.3 (31.3-35.3)	38.5 (36.7-40.3)
2005	23.0 (21.93-24.10)	19.7 (18.11-21.36)	26.1 (24.61-27.52)
2006	33.2 (31.59-34.82)	29.9 (27.58-32.18)	36.3 (34.23-38.36)
2007	36.2 (34.56-37.93)	33.0 (30.94-35.05)	39.3 (36.93-41.64)
2008	39.4 (37.79-41.10)	36.3 (34.04-38.56)	42.4 (40.18-44.68)
2009	40.7 (39.31-42.07)	38.3 (36.23-40.30)	43.0 (40.92-45.03)

NOTES: NHIS questions related to influenza vaccination have changed since 1997, as follows. Starting in 1997, respondents were asked if they had received a flu shot during the past 12 months. Beginning in 2003, respondents were also asked if they had received a flu vaccine sprayed in their nose during the past 12 months. NHIS Early Release influenza vaccination estimates have changed since 1997, as follows. Starting in 1997, Early Release influenza vaccination estimates covered receipt of an influenza shot only. Starting in 2005, Early Release influenza vaccination estimates covered seasonal influenza shot and/or seasonal intranasal influenza vaccination. When interpreting influenza vaccination estimates, readers should take into account changes made to the influenza vaccination questions noted above. An error in calculating influenza vaccination rates occurred for the first quarter of 2005 to the first quarter of 2007. The effect of this error on estimates was small. Compared with the original estimates, corrected estimates are slightly higher, usually by no more than 0.3 percentage point. The error has been corrected for all estimates in this Early Release, and the correction of estimates had no perceptible impact on the graphs. Responses to the previously mentioned influenza vaccination questions cannot be used to determine when during the preceding 12 months the subject received the influenza vaccination. In addition, estimates are subject to recall error, which will vary depending on when the question is asked because the receipt of an influenza vaccination is seasonal. The recommendations of the Advisory Committee on Immunization Practices regarding who should receive an influenza vaccination have changed over the years and changes in coverage estimates may reflect changes in recommendations (11–13). Influenza vaccination shortages have occurred during several influenza seasons (12–14). The analyses excluded those with unknown influenza vaccination status (about 3% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. See "About This Early Release" for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997–2009, Sample Adult Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

Table 4.1b. Annual percentage of adults aged 65 years and over who had received an influenza vaccination during the past 12 months, by sex: United States, 1997–2009

Year	Crude percent (95% confidence interval): total	Age-adjusted percent (95% confidence interval): total	Percent (95% confidence interval): men	Percent (95% confidence interval): women
1997	63.2(61.9-64.6)	63.1 (61.7-64.4)	64.8 (62.5-67.1)	62.1 (60.5-63.7)
1998	63.3(61.9-64.7)	63.3 (61.9-64.6)	63.7 (61.5-65.9)	63.0 (61.2-64.8)
1999	65.7(64.3-67.2)	65.1 (63.6-66.5)	67.2 (65.0-69.4)	64.6 (62.7-66.5)
2000	64.4 (63.0-65.9)	64.6 (63.2-66.0)	66.0 (63.8-68.3)	63.3 (61.6-65.0)
2001	63.1 (61.7-64.5)	63.2 (61.8-64.6)	64.8 (62.5-67.1)	61.8 (60.1-63.5)
2002	65.7 (64.3-67.2)	65.9 (64.5-67.3)	67.1 (64.7-69.5)	64.7 (62.8-66.6)
2003	65.5 (64.1-66.9)	65.6 (64.2-66.9)	66.0 (63.9-68.1)	65.1 (63.2-67.0)
2004	64.6 (63.2-66.1)	64.7 (63.2-66.1)	64.1 (61.9-66.3)	65.0 (63.3-66.7)
2005	59.7 (58.16-61.15)	59.7 (58.24-61.23)	58.9 (56.64-61.17)	60.2 (58.22-62.20)
2006	64.3 (62.39-66.19)	64.4 (62.51-66.32)	64.7 (62.04-67.43)	63.9 (61.65-66.24)
2007	66.7 (64.90-68.59)	66.8 (65.00-68.68)	66.7 (64.06-69.31)	66.8 (64.62-68.96)
2008	66.9 (65.08-68.80)	67.1 (65.31-68.89)	65.5 (62.74-68.33)	68.0 (65.94-70.07)
2009	66.7 (64.99-68.48)	67.0 (65.32-68.69)	67.3 (64.82-69.82)	66.3 (64.11-68.46)


NOTES: NHIS questions related to influenza vaccination have changed since 1997, as follows. Starting in 1997, respondents were asked if they had received a flu shot during the past 12 months. Beginning in 2003, respondents were also asked if they had received a flu vaccine sprayed in their nose during the past 12 months. NHIS Early Release influenza vaccination estimates have changed since 1997, as follows. Starting in 1997, Early Release influenza vaccination estimates covered receipt of an influenza shot only. Starting in 2005, Early Release influenza vaccination estimates covered seasonal influenza shot and/or seasonal intranasal influenza vaccination. When interpreting influenza vaccination estimates, readers should take into account changes made to the influenza vaccination questions noted above. An error in calculating influenza vaccination rates occurred for the first quarter of 2005 to the first quarter of 2007. The effect of this error on estimates was small. Compared with the original estimates, corrected estimates are slightly higher, usually by no more than 0.3 percentage point. The error has been corrected for all estimates in this Early Release, and the correction of estimates had no perceptible impact on the graphs. Responses to the previously mentioned influenza vaccination questions cannot be used to determine when during the preceding 12 months the subject received the influenza vaccination. In addition, estimates are subject to recall error, which will vary depending on when the question is asked because the receipt of an influenza vaccination is seasonal. The recommendations of the Advisory Committee on Immunization Practices regarding who should receive an influenza vaccination have changed over the years and changes in coverage estimates may reflect changes in recommendations (11–13). Influenza vaccination shortages have occurred during several influenza seasons (12–14). The analyses excluded those with unknown influenza vaccination status (about 3% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. See "About This Early Release" for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997–2009, Sample Adult Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.



- For adults aged 50–64 years, the annual percentage of persons who received an influenza vaccination during the past 12 months was 40.7% in 2009. This estimate was higher than, but not significantly different from, the estimate in 2008 (39.4%). This pattern was seen in both men and women. Following the influenza vaccination shortage during the 2004–2005 influenza season, estimates for this age group increased from 2005 to 2008, with the 2007 estimates being similar to the estimates in 2004 (13).
- For adults aged 65 years and over, the annual percentage of persons who received an influenza vaccination during the past 12 months was 66.7% in 2009. This estimate was not significantly different from the 2008 estimate (66.9%). The annual percentage of men who received an influenza vaccination during the past 12 months was 67.3% in 2009. This estimate was higher than, but not significantly different from, the 2008 estimate (65.5%). The annual percentage of women who received an influenza vaccination during the past 12 months was 66.3% in 2009. This estimate was lower than, but not significantly different from, the 2008 estimate (68.0%). Following the influenza vaccination shortage during the 2004–2005 influenza season, estimates for this age group increased from 2005 to 2008, with the 2006 estimates being similar to the estimates in 2004 (13).

Figure 4.2. Percentage of persons who had received an influenza vaccination during the past 12 months, by age group and sex: United States, January–September 2010




NOTES: Respondents were asked if they had received a flu vaccine sprayed in their nose during the past 12 months, in addition to a question regarding receipt of a flu shot during the past 12 months. These questions do not indicate whether the vaccination was a child's first or second dose. In August 2010, NHIS influenza vaccination questions were modified to reflect that for the first time, the widely available influenza vaccine included protection against both seasonal and H1N1 types of influenza. When interpreting influenza vaccination estimates, readers should take into account changes made to the influenza vaccination questions noted above. An error in calculating influenza vaccination rates occurred for the first quarter of 2005 to the first quarter of 2007. The effect of this error on estimates was small. Compared with the original estimates, corrected estimates are slightly higher, usually by no more than 0.3 percentage point. The error has been corrected for all estimates in this Early Release, and the correction of estimates had no perceptible impact on the graphs. Responses to the previously mentioned influenza vaccination questions cannot be used to determine when during the preceding 12 months the subject received the influenza vaccination. In addition, estimates are subject to recall error, which will vary depending on when the question is asked because the receipt of an influenza vaccination is seasonal. The recommendations of the Advisory Committee on Immunization Practices regarding who should receive an influenza vaccination have changed over the years and changes in coverage estimates may reflect changes in recommendations (11,15). The analyses excluded 1,597 persons (5.4%) with unknown influenza vaccination status.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–September 2010, combined Sample Adult and Sample Child Core components. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For both sexes combined, the percentage of persons who had an influenza vaccination during the past 12 months was highest among persons aged 65 years and over (63.7%), followed by persons aged 6 months–17 years (46.9%), 50–64 years (40.3%), and 18–49 years (24.8%).
- For adults aged 18–49 and 50–64 years, women were more likely than men to have received an influenza vaccination during the past 12 months.

Figure 4.3. Percentage of adults aged 65 years and over who had received an influenza vaccination during the past 12 months, by race/ethnicity: United States, January–September 2010



NOTES: Respondents were asked if they had received a flu vaccine sprayed in their nose during the past 12 months, in addition to a question regarding receipt of a flu shot during the past 12 months. In August 2010, NHIS influenza vaccination questions were modified to reflect that for the first time, the widely available influenza vaccine included protection against both seasonal and H1N1 types of influenza. When interpreting influenza vaccination estimates, readers should take into account changes made to the influenza vaccination questions noted above. An error in calculating influenza vaccination rates occurred for the first quarter of 2005 to the first quarter of 2007. The effect of this error on estimates was small. Compared with the original estimates, corrected estimates are slightly higher, usually by no more than 0.3 percentage point. The error has been corrected for all estimates in this Early Release, and the correction of estimates had no perceptible impact on the graphs. Responses to the previously mentioned influenza vaccination questions cannot be used to determine when during the preceding 12 months the subject received the influenza vaccination. In addition, estimates are subject to recall error, which will vary depending on when the question is asked because the receipt of an influenza vaccination is seasonal. The recommendations of the Advisory Committee on Immunization Practices regarding who should receive an influenza vaccination have changed over the years and changes in coverage estimates may reflect changes in recommendations (11). The analyses excluded 123 adults (3.0%) aged 65 years and over with unknown influenza vaccination status.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–September 2010, Sample Adult Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For adults aged 65 years and over, the percentage of persons receiving an influenza vaccination during the past 12 months was 53.7% for Hispanic persons, 65.8% for non-Hispanic white persons, and 53.1% for non-Hispanic black persons.
- Hispanic persons and non-Hispanic black persons were less likely than non-Hispanic white persons to have received an influenza vaccination during the past 12 months.

Data tables for Figures 4.1–4.3:

Data table for Figure 4.1. Percentage of adults aged 18 years and over who had received an influenza vaccination during the past 12 months, by age group and quarter: United States, 1997–September 2010

Year and quarter	Percent (95% confidence interval): 18-49 years	Percent (95% confidence interval): 50-64 years	Percent (95% confidence interval): 65 years and over
1997, quarter 1	13.6 (12.5-14.6)	30.5 (27.8-33.2)	65.0 (62.3-67.6)
1997, quarter 2	14.5 (13.4-15.5)	31.3 (28.7-34.0)	63.7 (61.1-66.2)
1997, quarter 3	13.6 (12.6-14.6)	32.0 (29.3-34.6)	63.1 (60.3-65.9)
1997, quarter 4	15.6 (14.5-16.7)	33.6 (31.1-36.2)	61.2 (58.7-63.8)
1998, quarter 1	16.1 (14.8-17.3)	34.2 (31.3-37.1)	66.3 (63.2-69.4)
1998, quarter 2	15.3 (14.1-16.5)	32.8 (30.1-35.5)	64.0 (61.3-66.8)
1998, quarter 3	14.5 (13.3-15.6)	32.0 (29.3-34.6)	61.3 (58.5-64.0)
1998, quarter 4	16.0 (14.8-17.2)	33.5 (30.8-36.1)	61.6 (58.7-64.5)
1999, quarter 1	16.3 (14.8-17.7)	34.2 (31.1-37.3)	67.0 (64.0-70.1)
1999, quarter 2	16.0 (14.7-17.3)	32.6 (29.8-35.4)	66.1 (63.4-68.8)
1999, quarter 3	15.8 (14.5-17.1)	32.8 (30.1-35.5)	64.1 (61.2-67.0)
1999, quarter 4	17.6 (16.2-18.9)	36.7 (34.2-39.2)	65.7 (62.7-68.6)
2000, quarter 1	18.6 (17.2-19.9)	36.6 (33.7-39.4)	68.2 (65.3-71.0)
2000, quarter 2	18.0 (16.7-19.4)	38.5 (35.7-41.4)	68.1 (65.6-70.7)
2000, quarter 3	18.2 (16.9-19.4)	36.6 (33.7-39.5)	67.1 (64.4-69.8)
2000, quarter 4	13.6 (12.4-14.8)	26.6 (24.2-29.0)	54.3 (51.6-57.1)
2001, quarter 1	14.7 (13.4-16.0)	32.3 (29.6-35.0)	63.3 (60.2-66.3)
2001, quarter 2	15.9 (14.7-17.1)	32.6 (30.1-35.1)	65.4 (62.8-68.0)
2001, quarter 3	14.9 (13.9-15.9)	33.3 (30.7-35.8)	64.0 (61.1-66.8)
2001, quarter 4	14.5 (13.6-15.9)	30.6 (28.0-33.1)	59.6 (56.7-62.4)
2002, quarter 1	16.4 (15.2-17.7)	36.3 (33.6-38.9)	66.6 (63.8-69.4)
2002, quarter 2	16.0 (14.8-17.2)	33.0 (30.5-35.5)	67.8 (65.3-70.3)
2002, quarter 3	16.2 (14.9-17.5)	33.1 (30.6-35.6)	63.1 (60.5-65.8)
2002, quarter 4	16.4 (15.1-17.8)	33.8 (31.0-36.6)	65.5 (62.4-68.6)
2003, quarter 1	17.1 (15.7-18.4)	36.8 (34.2-39.4)	67.8 (65.0-70.6)
2003, quarter 2	17.2 (15.8-18.6)	36.8 (33.9-39.7)	65.4 (62.6-68.3)
2003, quarter 3	16.4 (15.2-17.6)	37.4 (34.9-39.9)	65.4 (62.8-67.9)
2003, quarter 4	16.7 (15.2-18.1)	36.1 (33.3-39.0)	63.3 (60.1-66.5)
2004, quarter 1	19.3 (17.9-20.8)	40.6 (38.0-43.3)	70.3 (67.5-73.0)
2004, quarter 2	20.9 (19.1-22.6)	40.0 (37.1-43.0)	69.5 (66.7-72.3)
2004, quarter 3	19.4 (18.2-20.7)	41.0 (38.4-43.6)	66.4 (63.6-69.2)
2004, quarter 4	12.0 (10.9-13.1)	22.3 (20.2-24.5)	52.4 (49.5-55.4)

See notes at end of table.



Year and quarter	Percent (95% confidence interval): 18-49 years	Percent (95% confidence interval): 50-64 years	Percent (95% confidence interval): 65 years and over
2005, quarter 1	10.2 (9.03-11.41)	21.2 (19.05-23.42)	59.8 (56.66-62.90)
2005, quarter 2	10.3 (9.25-11.37)	21.1 (19.05-23.19)	62.0 (59.02-64.91)
2005, quarter 3	10.0 (9.03-11.03)	21.8 (19.64-24.01)	58.2 (55.42-60.97)
2005, quarter 4	12.4 (11.28-13.51)	27.8 (25.47-30.19)	58.7 (55.68-61.71)
2006, quarter 1	15.0 (13.69-16.36)	33.1 (29.95-36.20)	64.9 (61.65-68.15)
2006, quarter 2	16.2 (14.78-17.68)	31.8 (29.05-34.50)	66.6 (63.60-69.51)
2006, quarter 3	15.5 (13.56-17.47)	33.5 (29.45-37.51)	63.9 (58.90-68.84)
2006, quarter 4	15.4 (14.07-16.82)	34.5 (31.81-37.11)	61.9 (58.72-65.00)
2007, quarter 1	18.0 (16.36-19.63)	38.3 (35.28-41.39)	67.5 (64.47-70.62)
2007, quarter 2	16.7 (15.07-18.31)	36.2 (33.37-39.03)	68.8 (65.62-71.91)
2007, quarter 3	17.6 (15.47-19.74)	36.5 (32.10-40.95)	65.6 (60.90-70.39)
2007, quarter 4	18.7 (17.11-20.38)	34.0 (31.14-36.79)	65.1 (62.17-68.02)
2008, quarter 1	19.0 (17.41-20.56)	42.3 (39.12-45.57)	68.0 (64.78-71.21)
2008, quarter 2	19.5 (17.87-21.08)	39.8 (37.01-42.57)	67.4 (64.60-70.20)
2008, quarter 3	21.6 (19.95-23.26)	36.0 (33.10-38.86)	64.5 (61.09-68.00)
2008, quarter 4	19.4 (17.29-21.60)	39.7 (35.45-43.92)	67.8 (63.70-71.91)
2009, quarter 1	23.3 (20.81-25.78)	40.9 (37.05-44.67)	70.3 (66.08-74.59)
2009, quarter 2	22.5 (20.92-24.06)	42.5 (40.09-44.93)	64.8 (62.04-67.55)
2009, quarter 3	21.4 (19.81-22.98)	40.0 (37.13-42.82)	66.3 (63.03-69.47)
2009, quarter 4	23.9 (22.39-25.48)	40.6 (37.77-43.37)	65.8 (62.95-68.72)
2010, quarter 1	24.8 (22.03-27.58)	39.7 (35.70-43.66)	65.2 (60.50-69.85)
2010, quarter 2	23.3 (21.56-25.02)	42.7 (39.64-45.74)	64.6 (61.74-67.39)
2010, quarter 3	26.7 (24.81-28.64)	37.9 (35.02-40.69)	63.1 (59.87-66.39)

NOTES: Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. See "About This Early Release" for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997–September 2010, Sample Adult Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

Data table for Figure 4.2. Percentage of persons who had received an influenza vaccination during the past 12 months, by age group and sex: United States, January–September 2010

Age and sex	Percent	95% confidence interval
6 months-4 years, total	57.2	54.01-60.46
6 months-4 years, male	57.2	53.12-61.30
6 months-4 years, female	57.3	53.08-61.46
5-11 years, total	47.7	45.38-50.12
5-11 years, male	47.7	44.41-50.98
5-11 years, female	47.8	44.25-51.35
12-17 years, total	37.5	35.09-40.00
12-17 years, male	35.3	32.30-38.38
12-17 years, female	39.8	36.01-43.68
6 months-17 years, total	46.9	45.16-48.67
6 months-17 years, male	46.1	43.97-48.28
6 months-17 years, female	47.7	45.42-50.03
18-49 years, total	24.8	23.65-25.89
18-49 years, male	20.0	18.61-21.43
18-49 years, female	29.5	27.97-30.97
50-64 years, total	40.3	38.67-41.97
50-64 years, male	35.5	33.14-37.76
50-64 years, female	44.9	42.69-47.04
65 years and over, total	63.7	61.82-65.49
65 years and over, male	64.1	61.27-66.86
65 years and over, female	63.3	60.83-65.85
18 years and over (crude ¹), total	35.2	34.17-36.17
18 years and over (crude ¹), male	30.5	29.12-31.89
18 years and over (crude ¹), female	39.5	38.30-40.76
65 years and over (age-adjusted ²), total	63.9	62.11-65.72
65 years and over (age-adjusted ²), male	64.8	62.02-67.62
65 years and over (age-adjusted ²), female	63.4	60.89-65.85

¹Crude estimates are presented in the figure.

²Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted using the projected 2000 U.S. population as the standard population and using two age groups: 65–74 years and 75 years and over.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–September 2010, combined Sample Adult and Sample Child Core components. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

Data table for Figure 4.3. Percentage of adults aged 65 years and over who had received an influenza vaccination during the past 12 months, by race/ethnicity: United States, January–September 2010

Race/ethnicity	Crude¹ percent (95% confidence interval)	Age-adjusted² percent (95% confidence interval)
Hispanic or Latino	53.7 (47.39-59.97)	54.3 (47.91-60.76)
Not Hispanic or Latino, single race, white	65.8 (63.75-67.86)	66.0 (63.93-67.97)
Not Hispanic or Latino, single race, black	53.1 (48.45-57.78)	53.1 (48.31-57.85)

¹Crude estimates are presented in the figure.

²Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted using the projected 2000 U.S. population as the standard population and using two age groups: 65–74 years and 75 years and over.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–September 2010, Sample Adult Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.