


Figure 7.1. Percentage of adults aged 18 years and over who engaged in regular leisure-time physical activity: United States, 1997–September 2010




NOTES: This measure reflects the definition used for the physical activity Leading Health Indicator in *Healthy People 2010* (3). Regular leisure-time physical activity is defined as engaging in light-moderate leisure-time physical activity for greater than or equal to 30 minutes at a frequency greater than or equal to five times per week or engaging in vigorous leisure-time physical activity for greater than or equal to 20 minutes at a frequency greater than or equal to three times per week. In Early Releases before September 2005 (based on the 2004 National Health Interview Survey (NHIS)), regular physical activity was calculated slightly differently than for *Healthy People 2010*. The earlier Early Release estimates excluded from the analysis persons with unknown duration of light-moderate or vigorous leisure-time physical activity who were known to have not met the frequency recommendations for light-moderate or vigorous leisure-time physical activity (i.e., partial unknowns). With the current release, persons who were known to have not met the frequency recommendations are classified as "not regular," regardless of duration. All estimates have been rerun using the revised denominator. The impact of the change on the estimates was minimal (typically 0.1 percentage point or less). The analyses excluded persons with unknown physical activity participation (about 3% of respondents each year). Beginning with the 2003 data, NHIS transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. See "About This Early Release" for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997–September 2010, Sample Adult Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For January–September 2010, 34.8% (95% confidence interval = 33.79%–35.78%) of U.S. adults aged 18 years and over engaged in regular leisure-time physical activity, which was lower than, but not significantly different from, the January–September 2009 estimate of 35.4%.
- The annual percentages of adults aged 18 years and over who engaged in regular leisuretime physical activity ranged from 29.8% in 1998 to 34.7% in 2009.



Figure 7.2. Percentage of adults aged 18 years and over who engaged in regular leisure-time physical activity, by age group and sex: United States, January–September 2010




NOTES: This measure reflects the definition used for the physical activity Leading Health Indicator in *Healthy People 2010* (3). Regular leisure-time physical activity is defined as engaging in light-moderate leisure-time physical activity for greater than or equal to 30 minutes at a frequency greater than or equal to five times per week or engaging in vigorous leisure-time physical activity for greater than or equal to 20 minutes at a frequency greater than or equal to three times per week. In Early Releases before September 2005 (based on the 2004 National Health Interview Survey), regular physical activity was calculated slightly differently than for Healthy People 2010. The earlier Early Release estimates excluded from the analysis persons with unknown duration of light-moderate or vigorous leisure-time physical activity who were known to have not met the frequency recommendations for light-moderate or vigorous leisure-time physical activity (i.e., partial unknowns). With the current release, persons who were known to have not met the frequency recommendations are classified as "not regular," regardless of duration. The analyses excluded 361 persons (1.7%) with unknown physical activity participation.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–September 2010, Sample Adult Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For both sexes combined, adults aged 75 years and over were less likely than adults in younger age groups to engage in regular leisure-time physical activity.
- For adults aged 18–24 years, 25–64 years, 65–74 years, and 75 years and over, women were less likely than men to engage in regular leisure-time physical activity.



Figure 7.3. Age-sex-adjusted percentage of adults aged 18 years and over who engaged in regular leisure-time physical activity, by race/ethnicity: United States, January-September 2010



NOTES: This measure reflects the definition used for the physical activity Leading Health Indicator in *Healthy People 2010* (3). Regular leisure-time physical activity is defined as engaging in light-moderate leisure-time physical activity for greater than or equal to 30 minutes at a frequency greater than or equal to five times per week or engaging in vigorous leisure-time physical activity for greater than or equal to 20 minutes at a frequency greater than or equal to three times per week. In Early Releases before September 2005 (based on the 2004 National Health Interview Survey), regular physical activity was calculated slightly differently than for Healthy People 2010. The earlier Early Release estimates excluded from the analysis persons with unknown duration of light-moderate or vigorous leisure-time physical activity who were known to have not met the frequency recommendations for light-moderate or vigorous leisure-time physical activity (i.e., partial unknowns). With the current release, persons who were known to have not met the frequency recommendations are classified as "not regular," regardless of duration. The analyses excluded 361 persons (1.7%) with unknown physical activity participation. Estimates are age-sex-adjusted using the projected 2000 U.S. population as the standard population and using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–September 2010, Sample Adult Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- The age-sex-adjusted percentage of adults who engaged in regular leisure-time physical activity was 26.9% for Hispanic adults, 38.3% for non-Hispanic white adults, and 27.3% for non-Hispanic black adults.
- Non-Hispanic white adults were more likely to engage in regular leisure-time physical activity than Hispanic adults and non-Hispanic black adults.



Data tables for Figures 7.1–7.3:

Data table for Figure 7.1. Percentage of adults aged 18 years and over who engaged in regular leisure-time physical activity: United States, 1997–September 2010

Year	Crude ¹ percent (95% confidence interval)	Age-adjusted ² percent (95% confidence interval)
1997 yearly	32.0 (31.3–32.6)	31.8 (31.1–32.4)
1997 JanSept.	32.6 (31.7–33.5)	32.4 (31.4–33.3)
1998 yearly	29.8 (29.1–30.4)	29.6 (28.9–30.3)
1998 JanSept.	29.6 (28.6–30.5)	29.6 (28.7–30.6)
1999 yearly	30.2 (29.5–31.0)	30.1 (29.4–30.8)
1999 JanSept.	29.4 (28.4–30.5)	29.7 (28.6–30.7)
2000 yearly	31.7 (31.0-32.4)	31.7 (31.0-32.4)
2000 JanSept.	32.4 (31.7–33.2)	32.4 (31.6–33.2)
2001 yearly	31.9 (31.2-32.7)	31.8 (31.1-32.6)
2001 JanSept.	32.7 (31.9–33.6)	32.7 (31.8–33.5)
2002 yearly	31.7 (31.0-32.5)	31.7 (30.9-32.5)
2002 JanSept.	32.5 (31.6–33.4)	32.4 (31.6–33.3)
2003 yearly	32.8 (32.1–33.6)	32.8 (32.0–33.6)
2003 JanSept.	32.7 (31.6–33.8)	33.0 (31.9–34.1)
2004 yearly	30.2 (29.4–31.0)	30.1 (29.3–30.9)
2004 JanSept.	30.4 (29.3–31.5)	30.3 (29.2–31.4)
2005 yearly	30.1 (29.38-30.92)	30.1 (29.37-30.90)
2005 JanSept.	30.8 (29.79–31.84)	30.8 (29.78–31.81)
2006 yearly	30.9 (29.91-31.80)	30.9 (29.96-31.85)
2006 JanSept.	31.1 (29.91-32.24)	31.1 (29.93-32.27)
2007 yearly	30.8 (29.73-31.82)	30.8 (29.74-31.82)
2007 JanSept.	30.7 (29.47-32.02)	30.7 (29.40-31.93)
2008 yearly	31.9 (30.92-32.97)	32.0 (30.95-33.01)
2008 JanSept.	31.7 (30.69-32.74)	31.8 (30.74-32.78)
2009 yearly	34.7 (33.73-35.58)	34.8 (33.89-35.76)
2009 JanSept.	35.4 (34.26-36.62)	35.6 (34.43-36.84)
2010 JanSept.	34.8 (33.79-35.78)	35.0 (33.97-35.97)

¹Crude estimates are presented in the figure.

NOTES: Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. See "About This Early Release" for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997–September 2010, Sample Adult Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

²Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted using the projected 2000 U.S. population as the standard population and using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.



Data table for Figure 7.2. Percentage of adults aged 18 years and over who engaged in regular leisure-time physical activity, by age group and sex: United States, January—September 2010

Age and sex	Percent	95% confidence interval
18-24 years, total	42.4	39.27-45.46
18-24 years, male	50.9	46.88-54.94
18-24 years, female	33.7	30.13-37.26
25-64 years, total	36.3	35.15-37.40
25-64 years, male	37.9	36.38-39.44
25-64 years, female	34.7	33.37-36.02
65-74 years, total	26.8	24.52-29.07
65-74 years, male	31.8	28.11-35.40
65-74 years, female	22.5	19.70-25.38
75 years and over, total	17.8	15.61-19.99
75 years and over, male	24.3	20.33-28.35
75 years and over, female	13.4	11.35-15.53
18 years and over (crude ¹), total	34.8	33.79-35.78
18 years and over (crude ¹), male	38.3	36.95-39.60
18 years and over (crude ¹), female	31.5	30.41-32.67
18 years and over (age-adjusted ²), total	35.0	33.97-35.97
18 years and over (age-adjusted ²), male	38.2	36.90-39.51
18 years and over (age-adjusted ²), female	32.0	30.84-33.12

¹Crude estimates are presented in the figure.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January—September 2010, Sample Adult Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

²Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted using the projected 2000 U.S. population as the standard population and using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.



Data table for Figure 7.3. Age-sex-adjusted percentage of adults aged 18 years and over who engaged in regular leisure-time physical activity, by race/ethnicity: United States, January-September 2010

Race/ethnicity	Age-sex-adjusted ¹ percent (95% confidence interval)	Age-adjusted ² percent (95% confidence interval)
Hispanic or Latino	26.9 (25.21-28.66)	27.1 (25.33-28.86)
Not Hispanic or Latino, single race, white	38.3 (36.94-39.63)	38.3 (36.98-39.67)
Not Hispanic or Latino, single race, black	27.3 (25.59-29.04)	27.0 (25.26-28.72)

¹Age-sex-adjusted estimates are presented in the figure. Estimates are age-sex-adjusted using the projected 2000 U.S. population as the standard population and using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January—September 2010, Sample Adult Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

²Estimates for this Healthy People 2010 Leading Health Indicator are age-adjusted using the projected 2000 U.S. population as the standard population and using five age groups: 18–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.