


## Human immunodeficiency virus testing

**Figure 10.1. Percentage of adults aged 18 and over who had ever been tested for human immunodeficiency virus (HIV): United States, 1997–2013**



\*The AIDS Knowledge and Attitudes section of the National Health Interview Survey (NHIS) was dropped in 2011; only the HIV testing question was retained, and it was moved to the Adult Access to Health Care and Utilization section of the Sample Adult questionnaire and is not comparable to the years 2010 and earlier, or 2013 and later. This change is indicated by the dashed line in the figure. Differences observed in estimates based on the 2011–2012 NHIS and 2010 and earlier or 2013 and later NHIS may be partially or fully attributable to these changes in placement of the HIV testing question on the NHIS questionnaire.


§In 2013, the HIV testing question was moved from the Adult Access to Care and Utilization section of the Sample Adult questionnaire to the Adult Sexual Identity and Lifestyle section of the Sample Adult questionnaire and is not comparable to the years 2012 and earlier. This change is indicated by the dotted line in the figure. Differences observed in estimates based on the 2012 and earlier NHIS and the 2013 and later NHIS may be partially or fully attributable to these changes in placement of the HIV testing question on the NHIS questionnaire.

NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. • Individuals who received HIV testing solely as a result of blood donation were considered not to have been tested for HIV. • The analyses excluded those with unknown HIV test status (about 5% of respondents each year). • See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997–2013, Sample Adult Core component.

- For 2013, the percentage of adults who had ever been tested for HIV was 37.3% (95% confidence interval = 36.33%–38.20%).
- The percentage of adults who had ever had an HIV test increased from 31.8% in 1997 to 39.5% in 2010.

**Figure 10.2. Percentage of adults aged 18 and over who had ever been tested for human immunodeficiency virus (HIV), by age group and sex: United States, 2013**




NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. • Individuals who received HIV testing solely as a result of blood donation were considered not to have been tested for HIV. • The AIDS Knowledge and Attitudes section of the National Health Interview Survey (NHIS) was dropped in 2011; only the HIV testing question was retained, and it was moved to the Adult Access to Health Care and Utilization section of the Sample Adult questionnaire. In 2013, the HIV testing question was moved to the Adult Sexual Identity and Lifestyle section of the Sample Adult questionnaire and is not comparable to the years 2011–2012. Differences observed in estimates based on the 2010 and earlier NHIS and the 2011 and later NHIS may be partially or fully attributable to these changes in placement of the HIV testing question on the NHIS questionnaire. • The analyses excluded the 4.4% of adults with unknown HIV test status. • See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 2013, Sample Adult Core component.

- For both sexes combined, the percentage of persons who ever had an HIV test was highest among adults aged 25–34 (50.6%) and 35–44 (51.8%) and lowest among adults aged 65 and over (14.9%).
- For adults aged 18 and over and for age groups 18–24, 25–34, and 35–44, women were more likely than men to have ever had an HIV test. For adults aged 65 and over, women were less likely than men to have ever had an HIV test.

**Figure 10.3. Age-sex-adjusted percentage of adults aged 18 and over who had ever been tested for human immunodeficiency virus (HIV), by race/ethnicity: United States, 2013**



NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. • Individuals who received HIV testing solely as a result of blood donation were considered not to have been tested for HIV. • The AIDS Knowledge and Attitudes section of the National Health Interview Survey (NHIS) was dropped in 2011; only the HIV testing question was retained, and it was moved to the Adult Access to Health Care and Utilization section of the Sample Adult questionnaire. In 2013, the HIV testing question was moved to the Adult Sexual Identity and Lifestyle section of the Sample Adult questionnaire and is not comparable to the years 2011–2012. Differences observed in estimates based on the 2010 and earlier NHIS and the 2011 and later NHIS may be partially or fully attributable to these changes in placement of the HIV testing question on the NHIS questionnaire. • The analyses excluded the 4.4% of adults with unknown HIV test status. • Estimates are age adjusted using the projected 2000 U.S. population as the standard population and using five age groups: 18–24, 25–34, 35–44, 45–64, and 65 and over. • See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 2013, Sample Adult Core component.

- The age-sex-adjusted percentages of persons who ever had an HIV test were 39.9% for Hispanic persons, 35.2% for non-Hispanic white persons, and 57.0% for non-Hispanic black persons.
- Of the three race/ethnicity groups, non-Hispanic black persons were more likely to have ever had an HIV test than were Hispanic persons and non-Hispanic white persons.

## Data tables for Figures 10.1–10.3:

## Data table for Figure 10.1. Percentage of adults aged 18 and over who had ever been tested for human immunodeficiency virus (HIV): United States, 1997–2013

Year	Crude <sup>1</sup> percent (95% confidence interval)	Age-adjusted <sup>2</sup> percent (95% confidence interval)
1997	31.8 (31.1-32.5)	31.0 (30.4-31.6)
1998	30.2 (29.5-30.9)	29.6 (28.9-30.2)
1999	32.3 (31.6-33.0)	31.8 (31.1-32.5)
2000	32.1 (31.4-32.8)	31.8 (31.2-32.5)
2001	33.2 (32.5-34.0)	33.1 (32.4-33.7)
2002	34.5 (33.8-35.2)	34.5 (33.8-35.1)
2003	35.4 (34.7-36.1)	35.3 (34.7-36.0)
2004	34.6 (33.9-35.3)	34.8 (34.1-35.5)
2005	35.3 (34.58-36.01)	35.4 (34.73-36.10)
2006	35.8 (34.94-36.70)	36.1 (35.25-36.92)
2007	36.6 (35.59-37.62)	37.0 (35.28-38.79)
2008	38.7 (37.73-39.59)	39.2 (38.40-40.09)
2009	39.8 (38.70-40.84)	40.4 (39.36-41.40)
2010	39.5 (38.65-40.41)	40.3 (39.52-41.15)
2011 <sup>3</sup>	35.9 (35.14-36.72)	36.7 (35.97-37.51)
2012	34.8 (34.04-35.64)	35.8 (34.98-36.62)
2013 <sup>3</sup>	37.3 (36.33-38.20)	38.3 (37.42-39.25)

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age adjusted using the projected 2000 U.S. population as the standard population and using five age groups: 18–24, 25–34, 35–44, 45–64, and 65 and over.

<sup>3</sup>The AIDS Knowledge and Attitudes section of the National Health Interview Survey (NHIS) was dropped in 2011; only the HIV testing question was retained, and it was moved to the Adult Access to Health Care and Utilization section of the Sample Adult questionnaire. In 2013, the HIV testing question was moved to the Adult Sexual Identity and Lifestyle section of the Sample Adult questionnaire and is not comparable to the years 2011–2012. Differences observed in estimates based on the 2010 and earlier NHIS and the 2011 and later NHIS may be partially or fully attributable to these changes in placement of the HIV testing question on the NHIS questionnaire.

NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Beginning with the 2012 data, the National Health Interview Survey transitioned to weights derived from the 2010 census. For 2003–2011 data, weights were derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. For 1997–1999 data, weights were derived from the 1990 census. See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997–2013, Sample Adult Core component.

**Data table for Figure 10.2. Percentage of adults aged 18 and over who had ever been tested for human immunodeficiency virus (HIV), by age group and sex: United States, 2013**

Age (years) and sex	Percent	95% confidence interval
18–24, total	32.9	30.71-35.19
18–24, male	25.4	22.20-28.50
18–24, female	40.6	37.52-43.68
25–34, total	50.6	48.67-52.48
25–34, male	41.4	38.67-44.16
25–34, female	59.4	56.92-61.82
35–44, total	51.8	49.98-53.55
35–44, male	43.4	40.75-46.02
35–44, female	59.9	57.69-62.04
45–64, total	36.7	35.34-38.01
45–64, male	36.2	34.44-38.02
45–64, female	37.1	35.42-38.78
65 and over, total	14.9	13.72-16.08
65 and over, male	17.8	15.89-19.74
65 and over, female	12.6	11.19-14.04
18–64, total	42.2	41.13-43.23
18–64, male	37.1	35.68-38.46
18–64, female	47.1	45.85-48.35
18 and over (crude <sup>1</sup> ), total	37.3	36.33-38.20
18 and over (crude <sup>1</sup> ), male	33.9	32.68-35.11
18 and over (crude <sup>1</sup> ), female	40.4	39.28-41.51
18 and over (age-adjusted <sup>2</sup> ), total	38.3	37.42-39.25
18 and over (age-adjusted <sup>2</sup> ), male	34.2	33.01-35.40
18 and over (age-adjusted <sup>2</sup> ), female	42.4	41.36-43.52

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age adjusted using the projected 2000 U.S. population as the standard population and using five age groups: 18–24, 25–34, 35–44, 45–64, and 65 and over.

NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. The AIDS Knowledge and Attitudes section of the National Health Interview Survey (NHIS) was dropped in 2011; only the HIV testing question was retained, and it was moved to the Adult Access to Health Care and Utilization section of the Sample Adult questionnaire. In 2013, the HIV testing question was moved to the Adult Sexual Identity and Lifestyle section of the Sample Adult questionnaire and is not comparable to the years 2011–2012. Differences observed in estimates based on the 2010 and earlier NHIS and the 2011 and later NHIS may be partially or fully attributable to these changes in placement of the HIV testing question on the NHIS questionnaire.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 2013, Sample Adult Core component.

**Data table for Figure 10.3. Age-sex-adjusted percentage of adults aged 18 and over who had ever been tested for human immunodeficiency virus (HIV), by race/ethnicity: United States, 2013**

Race/ethnicity	Percent <sup>1</sup>	95% confidence interval
Hispanic or Latino	39.9	38.04-41.74
Not Hispanic or Latino, single race, white	35.2	34.13-36.20
Not Hispanic or Latino, single race, black	57.0	54.89-59.02

<sup>1</sup>Estimates are age-sex adjusted using the projected 2000 U.S. population as the standard population and using five age groups: 18–24, 25–34, 35–44, 45–64, and 65 and over.

NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. The AIDS Knowledge and Attitudes section of the National Health Interview Survey (NHIS) was dropped in 2011; only the HIV testing question was retained, and it was moved to the Adult Access to Health Care and Utilization section of the Sample Adult questionnaire. In 2013, the HIV testing question was moved to the Adult Sexual Identity and Lifestyle section of the Sample Adult questionnaire and is not comparable to the years 2011–2012. Differences observed in estimates based on the 2010 and earlier NHIS and the 2011 and later NHIS may be partially or fully attributable to these changes in placement of the HIV testing question on the NHIS questionnaire.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 2013, Sample Adult Core component.