


Receipt of influenza vaccination

Figure 4.1. Percentage of adults aged 18 and over who received an influenza vaccination during the past 12 months, by age group and quarter: United States, 1997–March 2014



NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. National Health Interview Survey (NHIS) questions related to influenza vaccination have changed since 1997. Starting in 1997, respondents were asked if they had received a flu shot during the past 12 months. Beginning in 2003, respondents were also asked if they had received a flu vaccine sprayed in their nose during the past 12 months. In August 2010, NHIS influenza vaccination questions were modified to reflect that, for the first time, the widely available influenza vaccine included protection against both seasonal and H1N1 types of influenza. NHIS Early Release influenza vaccination estimates have changed since 1997. Starting in 1997, Early Release influenza vaccination estimates covered receipt of an influenza shot only. Starting in 2005, Early Release influenza vaccination estimates covered seasonal influenza shot and/or seasonal intranasal influenza vaccination. Estimates based on data collected in quarters three and four of 2010 and one and two of 2011 could be affected, to an unknown extent, by reports of H1N1 immunization without seasonal flu immunization for the period when the two were administered separately (October 2009–May 2010). Prevalence of influenza vaccination during the past 12 months is different from season-specific coverage (see http://www.cdc.gov/mmwr/preview/mmwrhtml/ss6204a1.htm?s_cid=ss6204a1_w; estimates available from: <http://www.cdc.gov/flu/fluview>). The recommendations of the Advisory Committee on Immunization Practices regarding who should receive an influenza vaccination have changed over the years, and changes in coverage estimates may reflect changes in recommendations (4–6). An influenza vaccine shortage occurred during the 2004–2005 influenza season (4). Delays in the availability of influenza shots also occurred in fall 2000 and, to a lesser extent, in fall 2001 (4–7). The analyses excluded those with unknown influenza vaccination status (about 3% of respondents each year). See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997–March 2014, Sample Adult Core component.

- In the first quarter of 2014, the percentage of adults who had received an influenza vaccination during the past 12 months was 68.8% for adults aged 65 and over, 45.0% for those aged 50–64, and 31.2% for those aged 18–49.
- For the age groups 50–64 and 65 and over, first-quarter estimates in 2014 were lower than, but not significantly different from, first-quarter estimates in 2013. For the age group 18–49, the first-quarter estimate in 2014 was higher than, but not significantly different from, the first-quarter estimate in 2013.

Table 4.1a. Percentage of adults aged 50–64 who received an influenza vaccination during the past 12 months, by sex: United States, 1997–March 2014

Year	Total	Men	Women
	Percent (95% confidence interval)		
1997	31.9 (30.5-33.3)	28.0 (26.1-29.9)	35.5 (33.6-37.4)
1998	33.1 (31.7-34.5)	29.0 (27.0-31.0)	37.0 (35.1-38.9)
1999	34.1 (32.8-35.4)	30.5 (28.6-32.4)	37.4 (35.5-39.3)
2000	34.6 (33.1-36.1)	31.9 (29.9-33.9)	37.2 (35.2-39.1)
2001	32.2 (30.9-33.5)	30.3 (28.3-32.2)	34.0 (32.2-35.8)
2002	34.0 (32.7-35.3)	30.7 (28.8-32.5)	37.2 (35.4-38.9)
2003	36.8 (35.4-38.2)	34.5 (32.6-36.3)	38.9 (37.0-40.9)
2004	35.9 (34.6-37.3)	33.3 (31.3-35.3)	38.5 (36.7-40.3)
2005	23.0 (21.93-24.10)	19.7 (18.11-21.36)	26.1 (24.61-27.52)
2006	33.2 (31.59-34.82)	29.9 (27.58-32.18)	36.3 (34.23-38.36)
2007	36.2 (34.56-37.93)	33.0 (30.94-35.05)	39.3 (36.93-41.64)
2008	39.4 (37.79-41.10)	36.3 (34.04-38.56)	42.4 (40.18-44.68)
2009	40.7 (39.31-42.07)	38.3 (36.23-40.30)	43.0 (40.92-45.03)
2010	41.2 (39.73-42.66)	37.5 (35.41-39.56)	44.7 (42.78-46.56)
2011	42.7 (41.31-44.08)	39.0 (36.95-41.00)	46.2 (44.23-48.16)
2012	42.7 (41.44-43.99)	38.5 (36.87-40.11)	46.7 (44.94-48.50)
2013	46.5 (45.05-47.97)	43.1 (40.92-45.35)	49.7 (47.88-51.43)
Jan.–Mar. 2014	45.0 (42.03-47.95)	41.1 (36.49-45.71)	48.6 (44.84-52.41)

See notes at end of table.

Table 4.1b. Percentage of adults aged 65 and over who received an influenza vaccination during the past 12 months, by sex: United States, 1997–March 2014


Year	Total	Total	Men	Women
	Crude percent (95% confidence interval)	Age-adjusted percent (95% confidence interval)	Percent (95% confidence interval)	Percent (95% confidence interval)
1997	63.2 (61.9-64.6)	63.1 (61.7-64.4)	64.8 (62.5-67.1)	62.1 (60.5-63.7)
1998	63.3 (61.9-64.7)	63.3 (61.9-64.6)	63.7 (61.5-65.9)	63.0 (61.2-64.8)
1999	65.7 (64.3-67.2)	65.1 (63.6-66.5)	67.2 (65.0-69.4)	64.6 (62.7-66.5)
2000	64.4 (63.0-65.9)	64.6 (63.2-66.0)	66.0 (63.8-68.3)	63.3 (61.6-65.0)
2001	63.1 (61.7-64.5)	63.2 (61.8-64.6)	64.8 (62.5-67.1)	61.8 (60.1-63.5)
2002	65.7 (64.3-67.2)	65.9 (64.5-67.3)	67.1 (64.7-69.5)	64.7 (62.8-66.6)
2003	65.5 (64.1-66.9)	65.6 (64.2-66.9)	66.0 (63.9-68.1)	65.1 (63.2-67.0)
2004	64.6 (63.2-66.1)	64.7 (63.2-66.1)	64.1 (61.9-66.3)	65.0 (63.3-66.7)
2005	59.7 (58.16-61.15)	59.7 (58.24-61.23)	58.9 (56.64-61.17)	60.2 (58.22-62.20)
2006	64.3 (62.39-66.19)	64.4 (62.51-66.32)	64.7 (62.04-67.43)	63.9 (61.65-66.24)
2007	66.7 (64.90-68.59)	66.8 (65.00-68.68)	66.7 (64.06-69.31)	66.8 (64.62-68.96)
2008	66.9 (65.08-68.80)	67.1 (65.31-68.89)	65.5 (62.74-68.33)	68.0 (65.94-70.07)
2009	66.7 (64.99-68.48)	67.0 (65.32-68.69)	67.3 (64.82-69.82)	66.3 (64.11-68.46)
2010	63.6 (61.99-65.29)	63.9 (62.26-65.50)	63.1 (60.76-65.52)	64.0 (61.79-66.26)
2011	67.0 (65.45-68.50)	67.2 (65.74-68.75)	66.3 (64.12-68.53)	67.5 (65.68-69.30)
2012	66.5 (64.98-67.96)	66.9 (65.46-68.38)	65.2 (63.17-67.29)	67.4 (65.56-69.34)
2013	67.9 (66.38-69.48)	68.4 (66.89-69.92)	66.4 (63.89-68.87)	69.2 (67.42-70.88)
Jan.–Mar. 2014	68.8 (65.96-71.63)	69.3 (66.58-72.10)	69.7 (65.40-74.00)	68.1 (64.82-71.33)

NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. National Health Interview Survey (NHIS) questions related to influenza vaccination have changed since 1997. Starting in 1997, respondents were asked if they had received a flu shot during the past 12 months. Beginning in 2003, respondents were also asked if they had received a flu vaccine sprayed in their nose during the past 12 months. In August 2010, NHIS influenza vaccination questions were modified to reflect that, for the first time, the widely available influenza vaccine included protection against both seasonal and H1N1 types of influenza. NHIS Early Release influenza vaccination estimates have changed since 1997. Starting in 1997, Early Release influenza vaccination estimates covered receipt of an influenza shot only. Starting in 2005, Early Release influenza vaccination estimates covered seasonal influenza shot and/or seasonal intranasal influenza vaccination. Estimates based on data collected in quarters three and four of 2010 and one and two of 2011 could be affected, to an unknown extent, by reports of H1N1 immunization without seasonal flu immunization for the period when the two were administered separately (October 2009–May 2010). Prevalence of influenza vaccination during the past 12 months is different from season-specific coverage (see http://www.cdc.gov/mmwr/preview/mmwrhtml/ss6204a1.htm?s_cid=ss6204a1_w; estimates available from: <http://www.cdc.gov/flu/fluavaxview>). The recommendations of the Advisory Committee on Immunization Practices regarding who should receive an influenza vaccination have changed over the years and changes in coverage estimates may reflect changes in recommendations (4–6). An influenza vaccine shortage occurred during the 2004–2005 influenza season (4). Delays in the availability of influenza shots also occurred in fall 2000 and, to a lesser extent, in fall 2001 (4–7). The analyses excluded those with unknown influenza vaccination status (about 3% of respondents each year). See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997–March 2014, Sample Adult Core component.

- For January–March 2014, the percentage of adults who had received an influenza vaccination during the past 12 months was 45.0% for persons aged 50–64 (Table 4.1a). This estimate was lower than, but not significantly different from the 2013 estimate of 46.5%.
- For January–March 2014, the percentage of adults who had received an influenza vaccination during the past 12 months was 68.8% for persons aged 65 and over (Table 4.1b). This estimate was higher than, but not significantly different from, the 2013 estimate of 67.9%.
- For the age groups 50–64 and 65 and over, the percentage of adults who had received an influenza vaccination during the past 12 months increased from 1997 to January–March 2014.

Figure 4.2. Percentage of persons who received an influenza vaccination during the past 12 months, by age group and sex: United States, January–March 2014




NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Respondents were asked in separate questions if they had received a flu shot during the past 12 months or a flu vaccine sprayed in their nose during the past 12 months. These questions do not indicate whether the vaccination was a child’s first or second dose. In August 2010, National Health Interview Survey influenza vaccination questions were modified to reflect that, for the first time, the widely available influenza vaccine included protection against both seasonal and H1N1 types of influenza. When interpreting influenza vaccination estimates, readers should take into account changes made to the influenza vaccination questions noted above. Estimates based on data collected in quarters one and two of 2011 could be affected, to an unknown extent, by reports of H1N1 immunization without seasonal flu immunization for the period when the two were administered separately (October 2009–May 2010). Prevalence of influenza vaccination during the past 12 months is different from season-specific coverage (see http://www.cdc.gov/mmwr/preview/mmwrhtml/ss6204a1.htm?s_cid=ss6204a1_w; estimates available from: <http://www.cdc.gov/flu/fluview>). The recommendations of the Advisory Committee on Immunization Practices regarding who should receive an influenza vaccination have changed over the years, and changes in coverage estimates may reflect changes in recommendations (5,8). The analyses excluded the 2.0% of persons with unknown influenza vaccination status. See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–March 2014, combined Sample Adult and Sample Child Core components.

- For both sexes combined, the percentage of persons who had received an influenza vaccination during the past 12 months was highest among persons aged 65 and over (68.8%), followed by persons aged 6 months–17 years (49.7%), 50–64 years (45.0%), and 18–49 years (31.2%).
- For persons aged 6 months and over, for children aged 6 months–17 years, and for adults aged 18–49 and 50–64, females were more likely than males to have received an influenza vaccination during the past 12 months.

Figure 4.3. Percentage of persons aged 6 months and over who received an influenza vaccination during the past 12 months, by age group and race/ethnicity: United States, January–March 2014



NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Respondents were asked in separate questions if they had received a flu shot during the past 12 months or a flu vaccine sprayed in their nose during the past 12 months. In August 2010, National Health Interview Survey influenza vaccination questions were modified to reflect that, for the first time, the widely available influenza vaccine included protection against both seasonal and H1N1 types of influenza. When interpreting influenza vaccination estimates, readers should take into account changes made to the influenza vaccination questions noted above. Estimates based on data collected in quarters one and two of 2011 could be affected, to an unknown extent, by reports of H1N1 immunization without seasonal flu immunization for the period when the two were administered separately (October 2009–May 2010). Prevalence of influenza vaccination during the past 12 months is different from season-specific coverage (see http://www.cdc.gov/mmwr/preview/mmwrhtml/ss6204a1.htm?s_cid=ss6204a1_w; estimates available from: <http://www.cdc.gov/flu/fluview>). The recommendations of the Advisory Committee on Immunization Practices regarding who should receive an influenza vaccination have changed over the years, and changes in coverage estimates may reflect changes in recommendations (5). The analyses excluded the 2.0% of adults aged 65 and over with unknown influenza vaccination status. See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–March 2014, combined Sample Adult and Sample Child Core components.

- For children aged 6 months–17 years, the percentage who had received an influenza vaccination during the past 12 months was 51.2% for Hispanic children, 48.5% for non-Hispanic white children, and 51.6% for non-Hispanic black children.
- For adults aged 18 and over, the percentage who had received an influenza vaccination during the past 12 months was 31.4% for Hispanic adults, 45.0% for non-Hispanic white adults, and 35.5% for non-Hispanic black adults. Non-Hispanic white adults were more likely to have received an influenza vaccination than were Hispanic and non-Hispanic black adults.

Data tables for Figures 4.1–4.3:

Data table for Figure 4.1. Percentage of adults aged 18 and over who received an influenza vaccination during the past 12 months, by age group and quarter: United States, 1997–March 2014

Year and quarter	18–49 years	50–64 years	65 years and over
	Percent (95% confidence interval)		
1997, quarter 1	13.6 (12.5-14.6)	30.5 (27.8-33.2)	65.0 (62.3-67.6)
1997, quarter 2	14.5 (13.4-15.5)	31.3 (28.7-34.0)	63.7 (61.1-66.2)
1997, quarter 3	13.6 (12.6-14.6)	32.0 (29.3-34.6)	63.1 (60.3-65.9)
1997, quarter 4	15.6 (14.5-16.7)	33.6 (31.1-36.2)	61.2 (58.7-63.8)
1998, quarter 1	16.1 (14.8-17.3)	34.2 (31.3-37.1)	66.3 (63.2-69.4)
1998, quarter 2	15.3 (14.1-16.5)	32.8 (30.1-35.5)	64.0 (61.3-66.8)
1998, quarter 3	14.5 (13.3-15.6)	32.0 (29.3-34.6)	61.3 (58.5-64.0)
1998, quarter 4	16.0 (14.8-17.2)	33.5 (30.8-36.1)	61.6 (58.7-64.5)
1999, quarter 1	16.3 (14.8-17.7)	34.2 (31.1-37.3)	67.0 (64.0-70.1)
1999, quarter 2	16.0 (14.7-17.3)	32.6 (29.8-35.4)	66.1 (63.4-68.8)
1999, quarter 3	15.8 (14.5-17.1)	32.8 (30.1-35.5)	64.1 (61.2-67.0)
1999, quarter 4	17.6 (16.2-18.9)	36.7 (34.2-39.2)	65.7 (62.7-68.6)
2000, quarter 1	18.6 (17.2-19.9)	36.6 (33.7-39.4)	68.2 (65.3-71.0)
2000, quarter 2	18.0 (16.7-19.4)	38.5 (35.7-41.4)	68.1 (65.6-70.7)
2000, quarter 3	18.2 (16.9-19.4)	36.6 (33.7-39.5)	67.1 (64.4-69.8)
2000, quarter 4	13.6 (12.4-14.8)	26.6 (24.2-29.0)	54.3 (51.6-57.1)
2001, quarter 1	14.7 (13.4-16.0)	32.3 (29.6-35.0)	63.3 (60.2-66.3)
2001, quarter 2	15.9 (14.7-17.1)	32.6 (30.1-35.1)	65.4 (62.8-68.0)
2001, quarter 3	14.9 (13.9-15.9)	33.3 (30.7-35.8)	64.0 (61.1-66.8)
2001, quarter 4	14.5 (13.6-15.9)	30.6 (28.0-33.1)	59.6 (56.7-62.4)
2002, quarter 1	16.4 (15.2-17.7)	36.3 (33.6-38.9)	66.6 (63.8-69.4)
2002, quarter 2	16.0 (14.8-17.2)	33.0 (30.5-35.5)	67.8 (65.3-70.3)
2002, quarter 3	16.2 (14.9-17.5)	33.1 (30.6-35.6)	63.1 (60.5-65.8)
2002, quarter 4	16.4 (15.1-17.8)	33.8 (31.0-36.6)	65.5 (62.4-68.6)
2003, quarter 1	17.1 (15.7-18.4)	36.8 (34.2-39.4)	67.8 (65.0-70.6)
2003, quarter 2	17.2 (15.8-18.6)	36.8 (33.9-39.7)	65.4 (62.6-68.3)
2003, quarter 3	16.4 (15.2-17.6)	37.4 (34.9-39.9)	65.4 (62.8-67.9)
2003, quarter 4	16.7 (15.2-18.1)	36.1 (33.3-39.0)	63.3 (60.1-66.5)
2004, quarter 1	19.3 (17.9-20.8)	40.6 (38.0-43.3)	70.3 (67.5-73.0)
2004, quarter 2	20.9 (19.1-22.6)	40.0 (37.1-43.0)	69.5 (66.7-72.3)
2004, quarter 3	19.4 (18.2-20.7)	41.0 (38.4-43.6)	66.4 (63.6-69.2)
2004, quarter 4	12.0 (10.9-13.1)	22.3 (20.2-24.5)	52.4 (49.5-55.4)
2005, quarter 1	10.2 (9.03-11.41)	21.2 (19.05-23.42)	59.8 (56.66-62.90)
2005, quarter 2	10.3 (9.25-11.37)	21.1 (19.05-23.19)	62.0 (59.02-64.91)
2005, quarter 3	10.0 (9.03-11.03)	21.8 (19.64-24.01)	58.2 (55.42-60.97)
2005, quarter 4	12.4 (11.28-13.51)	27.8 (25.47-30.19)	58.7 (55.68-61.71)

See notes at end of table.

Data table for Figure 4.1. Percentage of adults aged 18 and over who received an influenza vaccination during the past 12 months, by age group and quarter: United States, 1997–March 2014 (Cont.)

Year and quarter	18–49 years	50–64 years	65 years and over
	Percent (95% confidence interval)		
2006, quarter 1	15.0 (13.69-16.36)	33.1 (29.95-36.20)	64.9 (61.65-68.15)
2006, quarter 2	16.2 (14.78-17.68)	31.8 (29.05-34.50)	66.6 (63.60-69.51)
2006, quarter 3	15.5 (13.56-17.47)	33.5 (29.45-37.51)	63.9 (58.90-68.84)
2006, quarter 4	15.4 (14.07-16.82)	34.5 (31.81-37.11)	61.9 (58.72-65.00)
2007, quarter 1	18.0 (16.36-19.63)	38.3 (35.28-41.39)	67.5 (64.47-70.62)
2007, quarter 2	16.7 (15.07-18.31)	36.2 (33.37-39.03)	68.8 (65.62-71.91)
2007, quarter 3	17.6 (15.47-19.74)	36.5 (32.10-40.95)	65.6 (60.90-70.39)
2007, quarter 4	18.7 (17.11-20.38)	34.0 (31.14-36.79)	65.1 (62.17-68.02)
2008, quarter 1	19.0 (17.41-20.56)	42.3 (39.12-45.57)	68.0 (64.78-71.21)
2008, quarter 2	19.5 (17.87-21.08)	39.8 (37.01-42.57)	67.4 (64.60-70.20)
2008, quarter 3	21.6 (19.95-23.26)	36.0 (33.10-38.86)	64.5 (61.09-68.00)
2008, quarter 4	19.4 (17.29-21.60)	39.7 (35.45-43.92)	67.8 (63.70-71.91)
2009, quarter 1	23.3 (20.81-25.78)	40.9 (37.05-44.67)	70.3 (66.08-74.59)
2009, quarter 2	22.5 (20.92-24.06)	42.5 (40.09-44.93)	64.8 (62.04-67.55)
2009, quarter 3	21.4 (19.81-22.98)	40.0 (37.13-42.82)	66.3 (63.03-69.48)
2009, quarter 4	24.5 (23.06-25.93)	39.5 (37.24-41.75)	65.5 (62.96-68.03)
2010, quarter 1	24.3 (22.45-26.17)	40.4 (37.81-43.05)	63.3 (60.51-66.04)
2010, quarter 2	23.3 (21.56-25.02)	42.7 (39.65-45.74)	64.6 (61.74-67.39)
2010, quarter 3	26.7 (24.81-28.64)	37.9 (35.02-40.69)	63.1 (59.87-66.39)
2010, quarter 4	25.1 (23.24-27.04)	43.8 (40.80-46.82)	63.6 (60.19-66.99)
2011, quarter 1	26.4 (24.86-27.93)	43.4 (40.61-46.17)	68.7 (66.01-71.37)
2011, quarter 2	26.9 (25.32-28.46)	41.7 (38.97-44.39)	69.4 (67.00-71.78)
2011, quarter 3	28.2 (26.47-29.87)	41.9 (39.23-44.57)	65.5 (62.59-68.50)
2011, quarter 4	27.5 (25.48-29.47)	43.8 (41.26-46.32)	64.3 (61.52-67.14)
2012, quarter 1	25.5 (23.78-27.25)	43.1 (40.28-46.00)	68.6 (65.62-71.63)
2012, quarter 2	27.9 (26.16-29.55)	42.5 (39.69-45.21)	69.4 (66.84-72.03)
2012, quarter 3	26.6 (24.68-28.54)	43.3 (40.95-45.73)	64.6 (61.86-67.32)
2012, quarter 4	25.2 (23.49-26.84)	41.9 (39.03-44.78)	63.4 (60.28-66.44)
2013, quarter 1	28.7 (26.90-30.51)	47.2 (44.43-49.88)	69.8 (66.84-72.70)
2013, quarter 2	31.5 (29.65-33.44)	48.1 (45.22-50.95)	69.1 (66.21-71.97)
2013, quarter 3	31.2 (29.45-32.96)	46.2 (43.58-48.90)	69.3 (66.30-72.20)
2013, quarter 4	27.0 (25.21-28.71)	44.6 (41.64-47.54)	63.7 (60.59-66.77)
2014, quarter 1	31.2 (29.29-33.06)	45.0 (42.03-47.95)	68.8 (65.96-71.63)

NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Beginning with the 2012 data, the National Health Interview Survey transitioned to weights derived from the 2010 census. For 2003–2011 data, weights were derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. For 1997–1999 data, weights were derived from the 1990 census. See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997–March 2014, Sample Adult Core component.

Data table for Figure 4.2. Percentage of persons who received an influenza vaccination during the past 12 months, by age group and sex: United States, January–March 2014

Age (years) and sex	Percent	95% confidence interval
6 months–4 years, total	55.8	51.22-60.30
6 months–4 years, male	51.7	45.24-58.10
6 months–4 years, female	60.2	53.99-66.43
5–11, total	50.2	46.22-54.19
5–11, male	48.6	43.48-53.81
5–11, female	51.8	46.34-57.30
12–17, total	44.9	41.10-48.68
12–17, male	42.5	37.28-47.72
12–17, female	47.5	41.63-53.29
6 months–17 years, total	49.7	47.35-52.09
6 months–17 years, male	47.3	43.87-50.64
6 months–17 years, female	52.3	49.13-55.56
18–49, total	31.2	29.29-33.06
18–49, male	26.8	24.10-29.59
18–49, female	35.4	32.77-37.93
50–64, total	45.0	42.03-47.95
50–64, male	41.1	36.49-45.71
50–64, female	48.6	44.84-52.41
65 and over, total	68.8	65.96-71.63
65 and over, male	69.7	65.40-74.00
65 and over, female	68.1	64.82-71.33
6 months and over (crude ¹), total	43.5	41.92-45.05
6 months and over (crude ¹), male	40.1	37.87-42.27
6 months and over (crude ¹), female	46.7	44.86-48.62
18 and over (crude ¹), total	41.7	39.88-43.45
18 and over (crude ¹), male	37.8	35.24-40.41
18 and over (crude ¹), female	45.2	43.20-47.22
65 and over (age-adjusted ²), total	69.3	66.58-72.10
65 and over (age-adjusted ²), male	71.0	66.73-75.17
65 and over (age-adjusted ²), female	68.3	65.06-71.50

¹Crude estimates are presented in the figure.

²Estimates are age adjusted using the projected 2000 U.S. population as the standard population and using two age groups: 65–74 and 75 and over.

NOTE: Data are based on household interviews of a sample of the civilian noninstitutionalized population.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–March 2014, combined Sample Adult and Sample Child Core components.

Data table for Figure 4.3. Percentage of persons aged 6 months and over who received an influenza vaccination during the past 12 months, by age group and race/ethnicity: United States, January–March 2014

Age and race/ethnicity	Percent ¹	95% confidence interval
6 months–17 years, Hispanic or Latino	51.2	46.24-56.19
6 months–17 years, not Hispanic or Latino, single race, white	48.5	45.12-51.86
6 months–17 years, not Hispanic or Latino, single race, black	51.6	45.03-58.25
18 years and over, Hispanic or Latino	31.4	28.41-34.46
18 years and over, not Hispanic or Latino, single race, white	45.0	42.84-47.20
18 years and over, not Hispanic or Latino, single race, black	35.5	32.05-39.01

NOTE: Data are based on household interviews of a sample of the civilian noninstitutionalized population.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–March 2014, combined Sample Adult and Sample Child Core components.