


## Failure to obtain needed medical care

**Figure 3.1. Percentage of persons of all ages who failed to obtain needed medical care due to cost at some time during the past 12 months: United States, 1997–June 2014**




NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. The analyses excluded persons with unknown responses to the question on failure to obtain needed medical care due to cost (about 0.2% of respondents each year). See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997–June 2014, Family Core component.

- For January–June 2014, 5.4% (95% confidence interval = 5.09%–5.77%) of the population failed to obtain needed medical care due to cost at some time during the past 12 months, which was lower than the 2013 estimate of 5.9%.
- The percentage of persons who failed to obtain needed medical care due to cost increased from 4.3% in 1999 to 6.9% in 2009 and 2010 then decreased to 5.4% in January–June 2014.

**Figure 3.2. Percentage of persons of all ages who failed to obtain needed medical care due to cost at some time during the past 12 months, by age group and sex: United States, January–June 2014**




NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. The analyses excluded the 0.1% persons with unknown responses to the question on failure to obtain needed medical care due to cost. See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–June 2014, Family Core component.

- For both sexes combined, the percentage of persons who failed to obtain needed medical care due to cost at some time during the past 12 months was 1.7% for those under age 18, 7.5% for those aged 18–64, and 2.4% for those aged 65 and over. Adults aged 18–64 were more likely than children under age 18 and adults aged 65 and over to have failed to obtain needed medical care due to cost.
- For persons of all ages combined, as well as for the age group 18–64, females were more likely than males to have failed to obtain needed medical care due to cost during the past 12 months.

**Figure 3.3. Age-sex-adjusted percentage of persons of all ages who failed to obtain needed medical care due to cost at some time during the past 12 months, by race/ethnicity: United States, January–June 2014**



NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. The analyses excluded the 0.1% of persons with unknown responses to the question on failure to obtain needed medical care due to cost. Estimates are age-sex adjusted using the projected 2000 U.S. population as the standard population and using three age groups: under 18, 18–64, and 65 and over. See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–June 2014, Family Core component.

- After adjustment for age and sex, 5.9% of Hispanic persons, 5.1% of non-Hispanic white persons, and 7.2% of non-Hispanic black persons failed to obtain needed medical care due to cost at some time during the year preceding the interview.
- Non-Hispanic black persons were more likely than Hispanic and non-Hispanic white persons to have failed to obtain needed medical care due to cost.

## Data tables for Figures 3.1–3.3:

### Data table for Figure 3.1. Percentage of persons of all ages who failed to obtain needed medical care due to cost at some time during the past 12 months: United States, 1997–June 2014

Year	Crude <sup>1</sup> percent (95% confidence interval)	Age-adjusted <sup>2</sup> percent (95% confidence interval)
1997	4.5 (4.3-4.7)	4.5 (4.4-4.7)
1998	4.2 (4.0-4.4)	4.2 (4.0-4.4)
1999	4.3 (4.1-4.5)	4.3 (4.1-4.5)
2000	4.5 (4.3-4.7)	4.5 (4.2-4.7)
2001	4.7 (4.5-4.9)	4.7 (4.5-4.9)
2002	4.7 (4.5-4.9)	4.7 (4.5-4.9)
2003	5.3 (5.1-5.5)	5.2 (5.0-5.5)
2004	5.5 (5.2-5.7)	5.4 (5.2-5.6)
2005	5.3 (5.05-5.45)	5.2 (4.99-5.40)
2006	5.8 (5.49-6.07)	5.7 (5.42-6.00)
2007	5.8 (5.54-6.11)	5.8 (5.47-6.03)
2008	6.5 (6.17-6.87)	6.4 (6.09-6.78)
2009	6.9 (6.62-7.21)	6.8 (6.54-7.12)
2010	6.9 (6.64-7.20)	6.8 (6.56-7.10)
2011	6.5 (6.26-6.77)	6.4 (6.18-6.68)
2012	6.2 (5.96-6.45)	6.1 (5.89-6.36)
2013	5.9 (5.66-6.16)	5.8 (5.60-6.10)
January–June 2014	5.4 (5.09-5.77)	5.4 (5.04-5.71)

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age adjusted using the projected 2000 U.S. population as the standard population and using three age groups: under 18, 18–64, and 65 and over.

NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Beginning with the 2012 data, the National Health Interview Survey transitioned to weights derived from the 2010 census. For 2003–2011 data, weights were derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. For 1997–1999 data, weights were derived from the 1990 census. See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997–June 2014, Family Core component.

**Data table for Figure 3.2. Percentage of persons of all ages who failed to obtain needed medical care due to cost at some time during the past 12 months, by age group and sex: United States, January–June 2014**

Age (years) and sex	Percent	95% confidence interval
Under 18, total	1.7	1.29-2.10
Under 18, male	1.7	1.20-2.24
Under 18, female	1.7	1.16-2.18
18–64, total	7.5	7.04-8.01
18–64, male	6.7	6.10-7.26
18–64, female	8.3	7.78-8.89
65 and over, total	2.4	1.99-2.87
65 and over, male	2.2	1.54-2.76
65 and over, female	2.7	2.06-3.25
All ages (crude <sup>1</sup> ), total	5.4	5.09-5.77
All ages (crude <sup>1</sup> ), male	4.9	4.47-5.28
All ages (crude <sup>1</sup> ), female	6.0	5.55-6.36
All ages (age-adjusted <sup>2</sup> ), total	5.4	5.04-5.71
All ages (age-adjusted <sup>2</sup> ), male	4.8	4.43-5.23
All ages (age-adjusted <sup>2</sup> ), female	5.9	5.51-6.29

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age adjusted using the projected 2000 U.S. population as the standard population and using three age groups: under 18, 18–64, and 65 and over.

NOTE: Data are based on household interviews of a sample of the civilian noninstitutionalized population.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–June 2014, Family Core component.

**Data table for Figure 3.3. Age-sex-adjusted percentage of persons of all ages who failed to obtain needed medical care due to cost at some time during the past 12 months, by race/ethnicity: United States, January–June 2014**

Race/ethnicity	Percent <sup>1</sup>	95% confidence interval
Hispanic or Latino	5.9	5.10-6.62
Not Hispanic or Latino, single race, white	5.1	4.67-5.45
Not Hispanic or Latino, single race, black	7.2	6.27-8.05

<sup>1</sup>Estimates are age-sex adjusted using the projected 2000 U.S. population as the standard population and using three age groups: under 18, 18–64, and 65 and over.

NOTE: Data are based on household interviews of a sample of the civilian noninstitutionalized population.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–June 2014, Family Core component.