


## Receipt of influenza vaccination

**Figure 4.1. Percentage of adults aged 18 and over who received an influenza vaccination during the past 12 months, by age group and quarter: United States, 1997–September 2014**



NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. National Health Interview Survey (NHIS) questions related to influenza vaccination have changed since 1997. Starting in 1997, respondents were asked if they had received a flu shot during the past 12 months. Beginning in 2003, respondents were also asked if they had received a flu vaccine sprayed in their nose during the past 12 months. In August 2010, questions were modified to reflect that, for the first time, the widely available influenza vaccine included protection against both seasonal and H1N1 types of influenza. NHIS Early Release influenza vaccination estimates have changed since 1997. Starting in 1997, Early Release influenza vaccination estimates covered receipt of an influenza shot only. Starting in 2005, Early Release influenza vaccination estimates covered seasonal influenza shot and/or seasonal intranasal influenza vaccination. Estimates based on data collected in quarters three and four of 2010 and one and two of 2011 could be affected, to an unknown extent, by reports of H1N1 immunization without seasonal flu immunization for the period when the two were administered separately (October 2009–May 2010). Prevalence of influenza vaccination during the past 12 months is different from season-specific coverage (see [http://www.cdc.gov/mmwr/preview/mmwrhtml/ss6204a1.htm?s\\_cid=ss6204a1\\_w](http://www.cdc.gov/mmwr/preview/mmwrhtml/ss6204a1.htm?s_cid=ss6204a1_w); estimates available from: <http://www.cdc.gov/flu/fluview>). Advisory Committee on Immunization Practices recommendations regarding who should receive an influenza vaccination have changed over the years, and changes in coverage estimates may reflect changes in recommendations (4–6). An influenza vaccine shortage occurred during the 2004–2005 influenza season (4). Delays in the availability of influenza shots also occurred in fall 2000 and, to a lesser extent, in fall 2001 (4–7). The analyses excluded those with unknown influenza vaccination status (about 3% of respondents each year). See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997–September 2014, Sample Adult Core component.

- In the third quarter of 2014, the percentages of adults who had received an influenza vaccination during the past 12 months were 69.9% for adults aged 65 and over, 45.1% for those aged 50–64, and 32.7% for those aged 18–49.
- For age groups 18–49, 50–64, and 65 and over, third-quarter estimates in 2014 were not significantly different from third-quarter estimates in 2013.

**Table 4.1a. Percentage of adults aged 50–64 who received an influenza vaccination during the past 12 months, by sex: United States, 1997–September 2014**

Year	Total	Men	Women
	Percent (95% confidence interval)		
1997	31.9 (30.5-33.3)	28.0 (26.1-29.9)	35.5 (33.6-37.4)
1998	33.1 (31.7-34.5)	29.0 (27.0-31.0)	37.0 (35.1-38.9)
1999	34.1 (32.8-35.4)	30.5 (28.6-32.4)	37.4 (35.5-39.3)
2000	34.6 (33.1-36.1)	31.9 (29.9-33.9)	37.2 (35.2-39.1)
2001	32.2 (30.9-33.5)	30.3 (28.3-32.2)	34.0 (32.2-35.8)
2002	34.0 (32.7-35.3)	30.7 (28.8-32.5)	37.2 (35.4-38.9)
2003	36.8 (35.4-38.2)	34.5 (32.6-36.3)	38.9 (37.0-40.9)
2004	35.9 (34.6-37.3)	33.3 (31.3-35.3)	38.5 (36.7-40.3)
2005	23.0 (21.93-24.10)	19.7 (18.11-21.36)	26.1 (24.61-27.52)
2006	33.2 (31.59-34.82)	29.9 (27.58-32.18)	36.3 (34.23-38.36)
2007	36.2 (34.56-37.93)	33.0 (30.94-35.05)	39.3 (36.93-41.64)
2008	39.4 (37.79-41.10)	36.3 (34.04-38.56)	42.4 (40.18-44.68)
2009	40.7 (39.31-42.07)	38.3 (36.23-40.30)	43.0 (40.92-45.03)
2010	41.2 (39.73-42.66)	37.5 (35.41-39.56)	44.7 (42.78-46.56)
2011	42.7 (41.31-44.08)	39.0 (36.95-41.00)	46.2 (44.23-48.16)
2012	42.7 (41.44-43.99)	38.5 (36.87-40.11)	46.7 (44.94-48.50)
2013	46.5 (45.05-47.97)	43.1 (40.92-45.35)	49.7 (47.88-51.43)
January–September 2014	46.1 (44.19-48.01)	41.7 (38.90-44.55)	50.2 (47.85-52.54)

See notes at end of table.

**Table 4.1b. Percentage of adults aged 65 and over who received an influenza vaccination during the past 12 months, by sex: United States, 1997–September 2014**


Year	Total	Total	Men	Women
	Crude percent (95% confidence interval)	Age-adjusted percent (95% confidence interval)	Percent (95% confidence interval)	Percent (95% confidence interval)
1997	63.2 (61.9-64.6)	63.1 (61.7-64.4)	64.8 (62.5-67.1)	62.1 (60.5-63.7)
1998	63.3 (61.9-64.7)	63.3 (61.9-64.6)	63.7 (61.5-65.9)	63.0 (61.2-64.8)
1999	65.7 (64.3-67.2)	65.1 (63.6-66.5)	67.2 (65.0-69.4)	64.6 (62.7-66.5)
2000	64.4 (63.0-65.9)	64.6 (63.2-66.0)	66.0 (63.8-68.3)	63.3 (61.6-65.0)
2001	63.1 (61.7-64.5)	63.2 (61.8-64.6)	64.8 (62.5-67.1)	61.8 (60.1-63.5)
2002	65.7 (64.3-67.2)	65.9 (64.5-67.3)	67.1 (64.7-69.5)	64.7 (62.8-66.6)
2003	65.5 (64.1-66.9)	65.6 (64.2-66.9)	66.0 (63.9-68.1)	65.1 (63.2-67.0)
2004	64.6 (63.2-66.1)	64.7 (63.2-66.1)	64.1 (61.9-66.3)	65.0 (63.3-66.7)
2005	59.7 (58.16-61.15)	59.7 (58.24-61.23)	58.9 (56.64-61.17)	60.2 (58.22-62.20)
2006	64.3 (62.39-66.19)	64.4 (62.51-66.32)	64.7 (62.04-67.43)	63.9 (61.65-66.24)
2007	66.7 (64.90-68.59)	66.8 (65.00-68.68)	66.7 (64.06-69.31)	66.8 (64.62-68.96)
2008	66.9 (65.08-68.80)	67.1 (65.31-68.89)	65.5 (62.74-68.33)	68.0 (65.94-70.07)
2009	66.7 (64.99-68.48)	67.0 (65.32-68.69)	67.3 (64.82-69.82)	66.3 (64.11-68.46)
2010	63.6 (61.99-65.29)	63.9 (62.26-65.50)	63.1 (60.76-65.52)	64.0 (61.79-66.26)
2011	67.0 (65.45-68.50)	67.2 (65.74-68.75)	66.3 (64.12-68.53)	67.5 (65.68-69.30)
2012	66.5 (64.98-67.96)	66.9 (65.46-68.38)	65.2 (63.17-67.29)	67.4 (65.56-69.34)
2013	67.9 (66.38-69.48)	68.4 (66.89-69.92)	66.4 (63.89-68.87)	69.2 (67.42-70.88)
January–September 2014	70.3 (68.72-71.89)	70.7 (69.17-72.31)	70.6 (67.96-73.14)	70.1 (68.26-71.96)

NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. National Health Interview Survey (NHIS) questions related to influenza vaccination have changed since 1997. Starting in 1997, respondents were asked if they had received a flu shot during the past 12 months. Beginning in 2003, respondents were also asked if they had received a flu vaccine sprayed in their nose during the past 12 months. In August 2010, questions were modified to reflect that, for the first time, the widely available influenza vaccine included protection against both seasonal and H1N1 types of influenza. NHIS Early Release influenza vaccination estimates have changed since 1997. Starting in 1997, Early Release influenza vaccination estimates covered receipt of an influenza shot only. Starting in 2005, Early Release influenza vaccination estimates covered seasonal influenza shot and/or seasonal intranasal influenza vaccination. Estimates based on data collected in quarters three and four of 2010 and one and two of 2011 could be affected, to an unknown extent, by reports of H1N1 immunization without seasonal flu immunization for the period when the two were administered separately (October 2009–May 2010). Prevalence of influenza vaccination during the past 12 months is different from season-specific coverage (see [http://www.cdc.gov/mmwr/preview/mmwrhtml/ss6204a1.htm?s\\_cid=ss6204a1\\_w](http://www.cdc.gov/mmwr/preview/mmwrhtml/ss6204a1.htm?s_cid=ss6204a1_w); estimates available from: <http://www.cdc.gov/flu/fluview>). Advisory Committee on Immunization Practices recommendations regarding who should receive an influenza vaccination have changed over the years and changes in coverage estimates may reflect changes in recommendations (4–6). An influenza vaccine shortage occurred during the 2004–2005 influenza season (4). Delays in the availability of influenza shots also occurred in fall 2000 and, to a lesser extent, in fall 2001 (4–7). The analyses excluded those with unknown influenza vaccination status (about 3% of respondents each year). See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997–September 2014, Sample Adult Core component.

- For January–September 2014, the percentage of adults who had received an influenza vaccination during the past 12 months was 46.1% for persons aged 50–64 (Table 4.1a). This estimate was not significantly different from the 2013 estimate of 46.5%.
- For January–September 2014, the percentage of adults who had received an influenza vaccination during the past 12 months was 70.3% for persons aged 65 and over (Table 4.1b). This estimate was higher than the 2013 estimate of 67.9%.
- For age group 50–64, the percentage of adults who had received an influenza vaccination during the past 12 months increased from 1997 to 2013. For age group 65 and over, the percentage of adults who had received an influenza vaccination during the past 12 months increased from 1997 to January–September 2014.

**Figure 4.2. Percentage of persons who received an influenza vaccination during the past 12 months, by age group and sex: United States, January–September 2014**




NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Respondents were asked in separate questions if they had received a flu shot during the past 12 months or a flu vaccine sprayed in their nose during the past 12 months. These questions do not indicate whether the vaccination was a child’s first or second dose. In August 2010, National Health Interview Survey influenza vaccination questions were modified to reflect that, for the first time, the widely available influenza vaccine included protection against both seasonal and H1N1 types of influenza. When interpreting influenza vaccination estimates, changes made to the influenza vaccination questions noted above should be taken into account. Estimates based on data collected in quarters one and two of 2011 could be affected, to an unknown extent, by reports of H1N1 immunization without seasonal flu immunization for the period when the two were administered separately (October 2009–May 2010). Prevalence of influenza vaccination during the past 12 months is different from season-specific coverage (see [http://www.cdc.gov/mmwr/preview/mmwrhtml/ss6204a1.htm?s\\_cid=ss6204a1\\_w](http://www.cdc.gov/mmwr/preview/mmwrhtml/ss6204a1.htm?s_cid=ss6204a1_w); estimates available from: <http://www.cdc.gov/flu/fluview>). Advisory Committee on Immunization Practices recommendations regarding who should receive an influenza vaccination have changed over the years, and changes in coverage estimates may reflect changes in recommendations (5,8). The analyses excluded the 2.1% of persons with unknown influenza vaccination status. See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–September 2014, combined Sample Adult and Sample Child Core components.

- For both sexes combined, the percentage of persons who had received an influenza vaccination during the past 12 months was highest among persons aged 65 and over (70.3%), followed by persons aged 6 months–17 years (51.2%), 50–64 years (46.1%), and 18–49 years (31.8%).
- For persons aged 6 months and over, and for adults aged 18–49 and 50–64, females were more likely than males to have received an influenza vaccination during the past 12 months.

**Figure 4.3. Percentage of persons aged 6 months and over who received an influenza vaccination during the past 12 months, by age group and race/ethnicity: United States, January–September 2014**



NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Respondents were asked in separate questions if they had received a flu shot during the past 12 months or a flu vaccine sprayed in their nose during the past 12 months. In August 2010, questions were modified to reflect that, for the first time, the widely available influenza vaccine included protection against both seasonal and H1N1 types of influenza. When interpreting influenza vaccination estimates, changes made to the influenza vaccination questions noted above should be taken into account. Estimates based on data collected in quarters one and two of 2011 could be affected, to an unknown extent, by reports of H1N1 immunization without seasonal flu immunization for the period when the two were administered separately (October 2009–May 2010). Prevalence of influenza vaccination during the past 12 months is different from season-specific coverage (see [http://www.cdc.gov/mmwr/preview/mmwrhtml/ss6204a1.htm?s\\_cid=ss6204a1\\_w](http://www.cdc.gov/mmwr/preview/mmwrhtml/ss6204a1.htm?s_cid=ss6204a1_w); estimates available from: <http://www.cdc.gov/flu/fluview>). Advisory Committee on Immunization Practices recommendations regarding who should receive an influenza vaccination have changed over the years, and changes in coverage estimates may reflect changes in recommendations (5). The analyses excluded the 2.1% of adults aged 65 and over with unknown influenza vaccination status. See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–September 2014, combined Sample Adult and Sample Child Core components.

- For children aged 6 months–17 years, the percentages who had received an influenza vaccination during the past 12 months were 53.2% for Hispanic children, 49.8% for non-Hispanic white children, and 49.6% for non-Hispanic black children. Hispanic children were more likely to have received an influenza vaccination than were non-Hispanic white children.
- For adults aged 18 and over, the percentages who had received an influenza vaccination during the past 12 months were 32.1% for Hispanic adults, 46.5% for non-Hispanic white adults, and 34.4% for non-Hispanic black adults. Non-Hispanic white adults were more likely to have received an influenza vaccination than were Hispanic and non-Hispanic black adults.

## Data tables for Figures 4.1–4.3:

**Data table for Figure 4.1. Percentage of adults aged 18 and over who received an influenza vaccination during the past 12 months, by age group and quarter: United States, 1997–September 2014**

Year and quarter	18–49 years	50–64 years	65 years and over
	Percent (95% confidence interval)		
1997, quarter 1	13.6 (12.5-14.6)	30.5 (27.8-33.2)	65.0 (62.3-67.6)
1997, quarter 2	14.5 (13.4-15.5)	31.3 (28.7-34.0)	63.7 (61.1-66.2)
1997, quarter 3	13.6 (12.6-14.6)	32.0 (29.3-34.6)	63.1 (60.3-65.9)
1997, quarter 4	15.6 (14.5-16.7)	33.6 (31.1-36.2)	61.2 (58.7-63.8)
1998, quarter 1	16.1 (14.8-17.3)	34.2 (31.3-37.1)	66.3 (63.2-69.4)
1998, quarter 2	15.3 (14.1-16.5)	32.8 (30.1-35.5)	64.0 (61.3-66.8)
1998, quarter 3	14.5 (13.3-15.6)	32.0 (29.3-34.6)	61.3 (58.5-64.0)
1998, quarter 4	16.0 (14.8-17.2)	33.5 (30.8-36.1)	61.6 (58.7-64.5)
1999, quarter 1	16.3 (14.8-17.7)	34.2 (31.1-37.3)	67.0 (64.0-70.1)
1999, quarter 2	16.0 (14.7-17.3)	32.6 (29.8-35.4)	66.1 (63.4-68.8)
1999, quarter 3	15.8 (14.5-17.1)	32.8 (30.1-35.5)	64.1 (61.2-67.0)
1999, quarter 4	17.6 (16.2-18.9)	36.7 (34.2-39.2)	65.7 (62.7-68.6)
2000, quarter 1	18.6 (17.2-19.9)	36.6 (33.7-39.4)	68.2 (65.3-71.0)
2000, quarter 2	18.0 (16.7-19.4)	38.5 (35.7-41.4)	68.1 (65.6-70.7)
2000, quarter 3	18.2 (16.9-19.4)	36.6 (33.7-39.5)	67.1 (64.4-69.8)
2000, quarter 4	13.6 (12.4-14.8)	26.6 (24.2-29.0)	54.3 (51.6-57.1)
2001, quarter 1	14.7 (13.4-16.0)	32.3 (29.6-35.0)	63.3 (60.2-66.3)
2001, quarter 2	15.9 (14.7-17.1)	32.6 (30.1-35.1)	65.4 (62.8-68.0)
2001, quarter 3	14.9 (13.9-15.9)	33.3 (30.7-35.8)	64.0 (61.1-66.8)
2001, quarter 4	14.5 (13.6-15.9)	30.6 (28.0-33.1)	59.6 (56.7-62.4)
2002, quarter 1	16.4 (15.2-17.7)	36.3 (33.6-38.9)	66.6 (63.8-69.4)
2002, quarter 2	16.0 (14.8-17.2)	33.0 (30.5-35.5)	67.8 (65.3-70.3)
2002, quarter 3	16.2 (14.9-17.5)	33.1 (30.6-35.6)	63.1 (60.5-65.8)
2002, quarter 4	16.4 (15.1-17.8)	33.8 (31.0-36.6)	65.5 (62.4-68.6)
2003, quarter 1	17.1 (15.7-18.4)	36.8 (34.2-39.4)	67.8 (65.0-70.6)
2003, quarter 2	17.2 (15.8-18.6)	36.8 (33.9-39.7)	65.4 (62.6-68.3)
2003, quarter 3	16.4 (15.2-17.6)	37.4 (34.9-39.9)	65.4 (62.8-67.9)
2003, quarter 4	16.7 (15.2-18.1)	36.1 (33.3-39.0)	63.3 (60.1-66.5)
2004, quarter 1	19.3 (17.9-20.8)	40.6 (38.0-43.3)	70.3 (67.5-73.0)
2004, quarter 2	20.9 (19.1-22.6)	40.0 (37.1-43.0)	69.5 (66.7-72.3)
2004, quarter 3	19.4 (18.2-20.7)	41.0 (38.4-43.6)	66.4 (63.6-69.2)
2004, quarter 4	12.0 (10.9-13.1)	22.3 (20.2-24.5)	52.4 (49.5-55.4)
2005, quarter 1	10.2 (9.03-11.41)	21.2 (19.05-23.42)	59.8 (56.66-62.90)
2005, quarter 2	10.3 (9.25-11.37)	21.1 (19.05-23.19)	62.0 (59.02-64.91)
2005, quarter 3	10.0 (9.03-11.03)	21.8 (19.64-24.01)	58.2 (55.42-60.97)
2005, quarter 4	12.4 (11.28-13.51)	27.8 (25.47-30.19)	58.7 (55.68-61.71)

See notes at end of table.

**Data table for Figure 4.1. Percentage of adults aged 18 and over who received an influenza vaccination during the past 12 months, by age group and quarter: United States, 1997–September 2014 (Cont.)**

Year and quarter	18–49 years	50–64 years	65 years and over
	Percent (95% confidence interval)		
2006, quarter 1	15.0 (13.69-16.36)	33.1 (29.95-36.20)	64.9 (61.65-68.15)
2006, quarter 2	16.2 (14.78-17.68)	31.8 (29.05-34.50)	66.6 (63.60-69.51)
2006, quarter 3	15.5 (13.56-17.47)	33.5 (29.45-37.51)	63.9 (58.90-68.84)
2006, quarter 4	15.4 (14.07-16.82)	34.5 (31.81-37.11)	61.9 (58.72-65.00)
2007, quarter 1	18.0 (16.36-19.63)	38.3 (35.28-41.39)	67.5 (64.47-70.62)
2007, quarter 2	16.7 (15.07-18.31)	36.2 (33.37-39.03)	68.8 (65.62-71.91)
2007, quarter 3	17.6 (15.47-19.74)	36.5 (32.10-40.95)	65.6 (60.90-70.39)
2007, quarter 4	18.7 (17.11-20.38)	34.0 (31.14-36.79)	65.1 (62.17-68.02)
2008, quarter 1	19.0 (17.41-20.56)	42.3 (39.12-45.57)	68.0 (64.78-71.21)
2008, quarter 2	19.5 (17.87-21.08)	39.8 (37.01-42.57)	67.4 (64.60-70.20)
2008, quarter 3	21.6 (19.95-23.26)	36.0 (33.10-38.86)	64.5 (61.09-68.00)
2008, quarter 4	19.4 (17.29-21.60)	39.7 (35.45-43.92)	67.8 (63.70-71.91)
2009, quarter 1	23.3 (20.81-25.78)	40.9 (37.05-44.67)	70.3 (66.08-74.59)
2009, quarter 2	22.5 (20.92-24.06)	42.5 (40.09-44.93)	64.8 (62.04-67.55)
2009, quarter 3	21.4 (19.81-22.98)	40.0 (37.13-42.82)	66.3 (63.03-69.48)
2009, quarter 4	24.5 (23.06-25.93)	39.5 (37.24-41.75)	65.5 (62.96-68.03)
2010, quarter 1	24.3 (22.45-26.17)	40.4 (37.81-43.05)	63.3 (60.51-66.04)
2010, quarter 2	23.3 (21.56-25.02)	42.7 (39.65-45.74)	64.6 (61.74-67.39)
2010, quarter 3	26.7 (24.81-28.64)	37.9 (35.02-40.69)	63.1 (59.87-66.39)
2010, quarter 4	25.1 (23.24-27.04)	43.8 (40.80-46.82)	63.6 (60.19-66.99)
2011, quarter 1	26.4 (24.86-27.93)	43.4 (40.61-46.17)	68.7 (66.01-71.37)
2011, quarter 2	26.9 (25.32-28.46)	41.7 (38.97-44.39)	69.4 (67.00-71.78)
2011, quarter 3	28.2 (26.47-29.87)	41.9 (39.23-44.57)	65.5 (62.59-68.50)
2011, quarter 4	27.5 (25.48-29.47)	43.8 (41.26-46.32)	64.3 (61.52-67.14)
2012, quarter 1	25.5 (23.78-27.25)	43.1 (40.28-46.00)	68.6 (65.62-71.63)
2012, quarter 2	27.9 (26.16-29.55)	42.5 (39.69-45.21)	69.4 (66.84-72.03)
2012, quarter 3	26.6 (24.68-28.54)	43.3 (40.95-45.73)	64.6 (61.86-67.32)
2012, quarter 4	25.2 (23.49-26.84)	41.9 (39.03-44.78)	63.4 (60.28-66.44)
2013, quarter 1	28.7 (26.90-30.51)	47.2 (44.43-49.88)	69.8 (66.84-72.70)
2013, quarter 2	31.5 (29.65-33.44)	48.1 (45.22-50.95)	69.1 (66.21-71.97)
2013, quarter 3	31.2 (29.45-32.96)	46.2 (43.58-48.90)	69.3 (66.30-72.20)
2013, quarter 4	27.0 (25.21-28.71)	44.6 (41.64-47.54)	63.7 (60.59-66.77)
2014, quarter 1	31.2 (29.29-33.06)	45.0 (42.03-47.95)	68.8 (65.96-71.63)
2014, quarter 2	31.6 (29.55-33.72)	48.2 (45.02-51.40)	72.2 (69.54-74.86)
2014, quarter 3	32.7 (30.64-34.67)	45.1 (42.11-48.11)	69.9 (67.35-72.48)

NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Beginning with 2012 data, the National Health Interview Survey transitioned to weights derived from the 2010 census. For 2003–2011 data, weights were derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. For 1997–1999 data, weights were derived from the 1990 census. See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997–September 2014, Sample Adult Core component.

**Data table for Figure 4.2. Percentage of persons who received an influenza vaccination during the past 12 months, by age group and sex: United States, January–September 2014**

Age (years) and sex	Percent	95% confidence interval
6 months–4 years, total	58.8	56.15-61.50
6 months–4 years, male	56.0	52.34-59.73
6 months–4 years, female	61.8	58.45-65.14
5–11, total	52.0	49.64-54.27
5–11, male	52.1	48.81-55.35
5–11, female	51.8	48.54-55.10
12–17, total	44.9	42.66-47.05
12–17, male	44.1	41.07-47.22
12–17, female	45.6	42.56-48.66
6 months–17 years, total	51.2	49.71-52.65
6 months–17 years, male	50.3	48.31-52.24
6 months–17 years, female	52.1	50.14-54.11
18–49, total	31.8	30.64-32.99
18–49, male	27.3	25.62-29.01
18–49, female	36.2	34.54-37.83
50–64, total	46.1	44.19-48.01
50–64, male	41.7	38.90-44.55
50–64, female	50.2	47.85-52.54
65 and over, total	70.3	68.72-71.89
65 and over, male	70.6	67.96-73.14
65 and over, female	70.1	68.26-71.96
6 months and over (crude <sup>1</sup> ), total	44.6	43.69-45.55
6 months and over (crude <sup>1</sup> ), male	41.3	39.89-42.70
6 months and over (crude <sup>1</sup> ), female	47.8	46.72-48.85
18 and over (crude <sup>1</sup> ), total	42.7	41.64-43.74
18 and over (crude <sup>1</sup> ), male	38.5	36.82-40.15
18 and over (crude <sup>1</sup> ), female	46.6	45.40-47.76
65 and over (age-adjusted <sup>2</sup> ), total	70.7	69.17-72.31
65 and over (age-adjusted <sup>2</sup> ), male	71.5	68.96-74.09
65 and over (age-adjusted <sup>2</sup> ), female	70.3	68.43-72.12

<sup>1</sup>Crude estimates are presented in the figure.

<sup>2</sup>Estimates are age-adjusted using the projected 2000 U.S. population as the standard population and using two age groups: 65–74 and 75 and over.

NOTE: Data are based on household interviews of a sample of the civilian noninstitutionalized population.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–September 2014, combined Sample Adult and Sample Child Core components.



**Data table for Figure 4.3. Percentage of persons aged 6 months and over who received an influenza vaccination during the past 12 months, by age group and race/ethnicity: United States, January–September 2014**

Age and race/ethnicity	Percent <sup>1</sup>	95% confidence interval
6 months–17 years, Hispanic or Latino	53.2	50.45-56.01
6 months–17 years, not Hispanic or Latino, single race, white	49.8	47.95-51.69
6 months–17 years, not Hispanic or Latino, single race, black	49.6	45.82-53.43
18 years and over, Hispanic or Latino	32.1	29.96-34.24
18 years and over, not Hispanic or Latino, single race, white	46.5	45.30-47.72
18 years and over, not Hispanic or Latino, single race, black	34.4	32.41-36.46

NOTE: Data are based on household interviews of a sample of the civilian noninstitutionalized population.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–September 2014, combined Sample Adult and Sample Child Core components.