


## Receipt of influenza vaccination

**Figure 4.1. Percentage of adults aged 18 and over who received an influenza vaccination during the past 12 months, by age group and quarter: United States, 1997– June 2017**



NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Respondents were asked if they received a flu vaccination during the past 12 months. From August 2010 through 2011, questions were modified to reflect that the seasonal influenza vaccine included protection for the 2009 pandemic H1N1 virus. Prevalence of influenza vaccination during the past 12 months is different from season-specific coverage (see <https://www.cdc.gov/flu/fluview>). Advisory Committee on Immunization Practices recommendations regarding who should receive an influenza vaccination have changed over the years, and changes in coverage estimates may reflect changes in recommendations (4–8). The analyses exclude the 2.0% of persons with unknown influenza vaccination status. See [Technical Notes](#) for more details.

DATA SOURCE: NCHS, National Health Interview Survey, 1997–June 2017, Sample Adult Core component.

- In the second quarter of 2017, the percentage of adults who received an influenza vaccination during the past 12 months by age group was 70.0% for adults aged 65 and over, 45.7% for those aged 50–64, and 33.6% for those aged 18–49.
- For the age group 18–49, the second-quarter estimate in 2017 (33.6%) was higher than, but not significantly different from, the second-quarter estimate in 2016 (32.7%).
- For the age group 50–64, the second-quarter estimate in 2017 (45.7%) was lower than, but not significantly different from, the second-quarter estimate in 2016 (46.4%).
- For adults 65 and over, the second-quarter estimate in 2017 (70.0%) was not significantly different from the second-quarter estimate in 2016 (69.8%).

**Table 4.1a. Percentage of adults aged 50–64 who received an influenza vaccination during the past 12 months, by sex: United States, 1997–June 2017**

Year	Total	Men	Women
	Percent (95% confidence interval)		
1997	31.9 (30.5–33.3)	28.0 (26.1–29.9)	35.5 (33.6–37.4)
1998	33.1 (31.7–34.5)	29.0 (27.0–31.0)	37.0 (35.1–38.9)
1999	34.1 (32.8–35.4)	30.5 (28.6–32.4)	37.4 (35.5–39.3)
2000	34.6 (33.1–36.1)	31.9 (29.9–33.9)	37.2 (35.2–39.1)
2001	32.2 (30.9–33.5)	30.3 (28.3–32.2)	34.0 (32.2–35.8)
2002	34.0 (32.7–35.3)	30.7 (28.8–32.5)	37.2 (35.4–38.9)
2003	36.8 (35.4–38.2)	34.5 (32.6–36.3)	38.9 (37.0–40.9)
2004	35.9 (34.6–37.3)	33.3 (31.3–35.3)	38.5 (36.7–40.3)
2005	23.0 (21.93–24.10)	19.7 (18.11–21.36)	26.1 (24.61–27.52)
2006	33.2 (31.59–34.82)	29.9 (27.58–32.18)	36.3 (34.23–38.36)
2007	36.2 (34.56–37.93)	33.0 (30.94–35.05)	39.3 (36.93–41.64)
2008	39.4 (37.79–41.10)	36.3 (34.04–38.56)	42.4 (40.18–44.68)
2009	40.7 (39.31–42.07)	38.3 (36.23–40.30)	43.0 (40.92–45.03)
2010	41.2 (39.73–42.66)	37.5 (35.41–39.56)	44.7 (42.78–46.56)
2011	42.7 (41.31–44.08)	39.0 (36.95–41.00)	46.2 (44.23–48.16)
2012	42.7 (41.44–43.99)	38.5 (36.87–40.11)	46.7 (44.94–48.50)
2013	46.5 (45.05–47.97)	43.1 (40.92–45.35)	49.7 (47.88–51.43)
2014	45.5 (43.91–47.03)	41.0 (38.65–43.38)	49.6 (47.70–51.59)
2015	48.1 (46.46–49.79)	45.3 (43.12–47.50)	50.8 (48.59–52.92)
2016	45.2 (43.88–46.47)	42.0 (40.16–43.77)	48.2 (46.30–50.09)
January–June 2017	46.8 (44.84–48.82)	43.6 (41.08–46.06)	49.8 (46.88–52.74)

See notes at end of table 4.1b.

**Table 4.1b. Percentage of adults aged 65 and over who received an influenza vaccination during the past 12 months, by sex: United States, 1997–June 2017**

Year	Total	Total	Men	Women
	Crude percent (95% confidence interval)	Age-adjusted percent (95% confidence interval) <sup>1</sup>	Percent (95% confidence interval)	Percent (95% confidence interval)
1997	63.2 (61.9–64.6)	63.1 (61.7–64.4)	64.8 (62.5–67.1)	62.1 (60.5–63.7)
1998	63.3 (61.9–64.7)	63.3 (61.9–64.6)	63.7 (61.5–65.9)	63.0 (61.2–64.8)
1999	65.7 (64.3–67.2)	65.1 (63.6–66.5)	67.2 (65.0–69.4)	64.6 (62.7–66.5)
2000	64.4 (63.0–65.9)	64.6 (63.2–66.0)	66.0 (63.8–68.3)	63.3 (61.6–65.0)
2001	63.1 (61.7–64.5)	63.2 (61.8–64.6)	64.8 (62.5–67.1)	61.8 (60.1–63.5)
2002	65.7 (64.3–67.2)	65.9 (64.5–67.3)	67.1 (64.7–69.5)	64.7 (62.8–66.6)
2003	65.5 (64.1–66.9)	65.6 (64.2–66.9)	66.0 (63.9–68.1)	65.1 (63.2–67.0)
2004	64.6 (63.2–66.1)	64.7 (63.2–66.1)	64.1 (61.9–66.3)	65.0 (63.3–66.7)
2005	59.7 (58.16–61.15)	59.7 (58.24–61.23)	58.9 (56.64–61.17)	60.2 (58.22–62.20)
2006	64.3 (62.39–66.19)	64.4 (62.51–66.32)	64.7 (62.04–67.43)	63.9 (61.65–66.24)
2007	66.7 (64.90–68.59)	66.8 (65.00–68.68)	66.7 (64.06–69.31)	66.8 (64.62–68.96)
2008	66.9 (65.08–68.80)	67.1 (65.31–68.89)	65.5 (62.74–68.33)	68.0 (65.94–70.07)
2009	66.7 (64.99–68.48)	67.0 (65.32–68.69)	67.3 (64.82–69.82)	66.3 (64.11–68.46)
2010	63.6 (61.99–65.29)	63.9 (62.26–65.50)	63.1 (60.76–65.52)	64.0 (61.79–66.26)
2011	67.0 (65.45–68.50)	67.2 (65.74–68.75)	66.3 (64.12–68.53)	67.5 (65.68–69.30)
2012	66.5 (64.98–67.96)	66.9 (65.46–68.38)	65.2 (63.17–67.29)	67.4 (65.56–69.34)
2013	67.9 (66.38–69.48)	68.4 (66.89–69.92)	66.4 (63.89–68.87)	69.2 (67.42–70.88)
2014	70.0 (68.62–71.38)	70.5 (69.09–71.86)	70.1 (67.94–72.31)	69.9 (68.23–71.59)
2015	69.1 (67.79–70.33)	69.4 (68.14–70.63)	70.4 (68.66–72.18)	68.0 (66.16–69.75)
2016	67.2 (65.77–68.65)	67.7 (66.33–69.17)	66.7 (64.72–68.75)	67.6 (65.97–69.21)
January–June 2017	70.2 (68.15–72.22)	71.1 (69.16–73.04)	71.9 (69.17–74.60)	68.8 (65.92–71.57)


<sup>1</sup>Estimates are age-adjusted using the projected 2000 U.S. population as the standard population and two age groups: 65–74 and 75 and over.

NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Respondents were asked if they received a flu vaccination during the past 12 months. From August 2010 through 2011, questions were modified to reflect that the seasonal influenza vaccine included protection for the 2009 pandemic H1N1 virus. Prevalence of influenza vaccination during the past 12 months is different from season-specific coverage (see <https://www.cdc.gov/flu/fluview>). Advisory Committee on Immunization Practices recommendations regarding who should receive an influenza vaccination have changed over the years, and changes in coverage estimates may reflect changes in recommendations (4–8). The analyses exclude the 2.0% of persons with unknown influenza vaccination status. See [Technical Notes](#) for more details.

DATA SOURCE: NCHS, National Health Interview Survey, 1997–June 2017, Sample Adult Core component.

- For January–June 2017, the percentage of adults aged 50–64 who received an influenza vaccination during the past 12 months was 46.8% (95% confidence interval = 44.84–48.82) (Table 4.1a). This estimate was higher than, but not significantly different from, the 2016 estimate of 45.2%. For this age group, the percentage of adults who received an influenza vaccination during the past 12 months generally increased from 1997 to January–June 2017.
- For January–June 2017, the percentage of adults aged 65 and over who received an influenza vaccination during the past 12 months was 70.2%, which was higher than, but not significantly different from, the 2016 estimate of 67.2% (Table 4.1b). For this age group, the percentage of adults who received an influenza vaccination during the past 12 months increased from 1997 to January–June 2017.

**Figure 4.2. Percentage of persons who received an influenza vaccination during the past 12 months, by age group and sex: United States, January–June 2017**




NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Respondents were asked if they received a flu vaccination during the past 12 months. From August 2010 through 2011, questions were modified to reflect that the seasonal influenza vaccine included protection for the 2009 pandemic H1N1 virus. Prevalence of influenza vaccination during the past 12 months is different from season-specific coverage (see <https://www.cdc.gov/flu/fluview>). Advisory Committee on Immunization Practices recommendations regarding who should receive an influenza vaccination have changed over the years, and changes in coverage estimates may reflect changes in recommendations (4–8). The analyses exclude the 2.0% of persons with unknown influenza vaccination status. See [Technical Notes](#) for more details.

DATA SOURCE: NCHS, National Health Interview Survey, January–June 2017, combined Sample Adult and Sample Child Core components.

- For both sexes combined, the percentage of persons who received an influenza vaccination during the past 12 months was highest among persons aged 65 and over (70.2%) and lowest among persons aged 18–49 (33.9%). This pattern held true for males and females.
- For persons aged 6 months and over and for adults aged 18–49 and 50–64, females were more likely than males to have received an influenza vaccination during the past 12 months.

**Figure 4.3. Percentage of persons aged 6 months and over who received an influenza vaccination during the past 12 months, by age group and race and ethnicity: United States, January–June 2017**



NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Respondents were asked if they received a flu vaccination during the past 12 months. From August 2010 through 2011, questions were modified to reflect that the seasonal influenza vaccine included protection for the 2009 pandemic H1N1 virus. Prevalence of influenza vaccination during the past 12 months is different from season-specific coverage (see <https://www.cdc.gov/flu/fluview>). Advisory Committee on Immunization Practices recommendations regarding who should receive an influenza vaccination have changed over the years, and changes in coverage estimates may reflect changes in recommendations (4–8). The analyses exclude the 2.0% of persons with unknown influenza vaccination status. See [Technical Notes](#) for more details.

DATA SOURCE: NCHS, National Health Interview Survey, January–June 2017, combined Sample Adult and Sample Child Core components.

- For children aged 6 months–17 years, the percentage by race and ethnicity who received an influenza vaccination during the past 12 months was 50.2% for Hispanic children, 45.5% for non-Hispanic white children, and 43.2% for non-Hispanic black children.
- No significant differences in the prevalence of receiving an influenza vaccination during the past 12 months by race and ethnicity groups were found for children aged 6 months–17 years.
- For adults aged 18 and over, the percentage who received an influenza vaccination during the past 12 months was 36.7% for Hispanic adults, 47.4% for non-Hispanic white adults, and 36.3% for non-Hispanic black adults.
- Non-Hispanic white adults were more likely to have received an influenza vaccination than non-Hispanic black adults and Hispanic adults.

Data tables for Figures 4.1–4.3:

Data table for Figure 4.1. Percentage of adults aged 18 and over who received an influenza vaccination during the past 12 months, by age group and quarter: United States, 1997–June 2017

Year and quarter	18–49	50–64	65 and over
	Percent (95% confidence interval)		
1997, quarter 1	13.6 (12.5–14.6)	30.5 (27.8–33.2)	65.0 (62.3–67.6)
1997, quarter 2	14.5 (13.4–15.5)	31.3 (28.7–34.0)	63.7 (61.1–66.2)
1997, quarter 3	13.6 (12.6–14.6)	32.0 (29.3–34.6)	63.1 (60.3–65.9)
1997, quarter 4	15.6 (14.5–16.7)	33.6 (31.1–36.2)	61.2 (58.7–63.8)
1998, quarter 1	16.1 (14.8–17.3)	34.2 (31.3–37.1)	66.3 (63.2–69.4)
1998, quarter 2	15.3 (14.1–16.5)	32.8 (30.1–35.5)	64.0 (61.3–66.8)
1998, quarter 3	14.5 (13.3–15.6)	32.0 (29.3–34.6)	61.3 (58.5–64.0)
1998, quarter 4	16.0 (14.8–17.2)	33.5 (30.8–36.1)	61.6 (58.7–64.5)
1999, quarter 1	16.3 (14.8–17.7)	34.2 (31.1–37.3)	67.0 (64.0–70.1)
1999, quarter 2	16.0 (14.7–17.3)	32.6 (29.8–35.4)	66.1 (63.4–68.8)
1999, quarter 3	15.8 (14.5–17.1)	32.8 (30.1–35.5)	64.1 (61.2–67.0)
1999, quarter 4	17.6 (16.2–18.9)	36.7 (34.2–39.2)	65.7 (62.7–68.6)
2000, quarter 1	18.6 (17.2–19.9)	36.6 (33.7–39.4)	68.2 (65.3–71.0)
2000, quarter 2	18.0 (16.7–19.4)	38.5 (35.7–41.4)	68.1 (65.6–70.7)
2000, quarter 3	18.2 (16.9–19.4)	36.6 (33.7–39.5)	67.1 (64.4–69.8)
2000, quarter 4	13.6 (12.4–14.8)	26.6 (24.2–29.0)	54.3 (51.6–57.1)
2001, quarter 1	14.7 (13.4–16.0)	32.3 (29.6–35.0)	63.3 (60.2–66.3)
2001, quarter 2	15.9 (14.7–17.1)	32.6 (30.1–35.1)	65.4 (62.8–68.0)
2001, quarter 3	14.9 (13.9–15.9)	33.3 (30.7–35.8)	64.0 (61.1–66.8)
2001, quarter 4	14.5 (13.6–15.9)	30.6 (28.0–33.1)	59.6 (56.7–62.4)
2002, quarter 1	16.4 (15.2–17.7)	36.3 (33.6–38.9)	66.6 (63.8–69.4)
2002, quarter 2	16.0 (14.8–17.2)	33.0 (30.5–35.5)	67.8 (65.3–70.3)
2002, quarter 3	16.2 (14.9–17.5)	33.1 (30.6–35.6)	63.1 (60.5–65.8)
2002, quarter 4	16.4 (15.1–17.8)	33.8 (31.0–36.6)	65.5 (62.4–68.6)
2003, quarter 1	17.1 (15.7–18.4)	36.8 (34.2–39.4)	67.8 (65.0–70.6)
2003, quarter 2	17.2 (15.8–18.6)	36.8 (33.9–39.7)	65.4 (62.6–68.3)
2003, quarter 3	16.4 (15.2–17.6)	37.4 (34.9–39.9)	65.4 (62.8–67.9)
2003, quarter 4	16.7 (15.2–18.1)	36.1 (33.3–39.0)	63.3 (60.1–66.5)
2004, quarter 1	19.3 (17.9–20.8)	40.6 (38.0–43.3)	70.3 (67.5–73.0)
2004, quarter 2	20.9 (19.1–22.6)	40.0 (37.1–43.0)	69.5 (66.7–72.3)
2004, quarter 3	19.4 (18.2–20.7)	41.0 (38.4–43.6)	66.4 (63.6–69.2)
2004, quarter 4	12.0 (10.9–13.1)	22.3 (20.2–24.5)	52.4 (49.5–55.4)
2005, quarter 1	10.2 (9.03–11.41)	21.2 (19.05–23.42)	59.8 (56.66–62.90)
2005, quarter 2	10.3 (9.25–11.37)	21.1 (19.05–23.19)	62.0 (59.02–64.91)
2005, quarter 3	10.0 (9.03–11.03)	21.8 (19.64–24.01)	58.2 (55.42–60.97)
2005, quarter 4	12.4 (11.28–13.51)	27.8 (25.47–30.19)	58.7 (55.68–61.71)

See notes at end of table.

**Data table for Figure 4.1. Percentage of adults aged 18 and over who received an influenza vaccination during the past 12 months, by age group and quarter: United States, 1997–June 2017 (Cont.)**

Year and quarter	18–49	50–64	65 and over
	Percent (95% confidence interval)		
2006, quarter 1	15.0 (13.69–16.36)	33.1 (29.95–36.20)	64.9 (61.65–68.15)
2006, quarter 2	16.2 (14.78–17.68)	31.8 (29.05–34.50)	66.6 (63.60–69.51)
2006, quarter 3	15.5 (13.56–17.47)	33.5 (29.45–37.51)	63.9 (58.90–68.84)
2006, quarter 4	15.4 (14.07–16.82)	34.5 (31.81–37.11)	61.9 (58.72–65.00)
2007, quarter 1	18.0 (16.36–19.63)	38.3 (35.28–41.39)	67.5 (64.47–70.62)
2007, quarter 2	16.7 (15.07–18.31)	36.2 (33.37–39.03)	68.8 (65.62–71.91)
2007, quarter 3	17.6 (15.47–19.74)	36.5 (32.10–40.95)	65.6 (60.90–70.39)
2007, quarter 4	18.7 (17.11–20.38)	34.0 (31.14–36.79)	65.1 (62.17–68.02)
2008, quarter 1	19.0 (17.41–20.56)	42.3 (39.12–45.57)	68.0 (64.78–71.21)
2008, quarter 2	19.5 (17.87–21.08)	39.8 (37.01–42.57)	67.4 (64.60–70.20)
2008, quarter 3	21.6 (19.95–23.26)	36.0 (33.10–38.86)	64.5 (61.09–68.00)
2008, quarter 4	19.4 (17.29–21.60)	39.7 (35.45–43.92)	67.8 (63.70–71.91)
2009, quarter 1	23.3 (20.81–25.78)	40.9 (37.05–44.67)	70.3 (66.08–74.59)
2009, quarter 2	22.5 (20.92–24.06)	42.5 (40.09–44.93)	64.8 (62.04–67.55)
2009, quarter 3	21.4 (19.81–22.98)	40.0 (37.13–42.82)	66.3 (63.03–69.48)
2009, quarter 4	24.5 (23.06–25.93)	39.5 (37.24–41.75)	65.5 (62.96–68.03)
2010, quarter 1	24.3 (22.45–26.17)	40.4 (37.81–43.05)	63.3 (60.51–66.04)
2010, quarter 2	23.3 (21.56–25.02)	42.7 (39.65–45.74)	64.6 (61.74–67.39)
2010, quarter 3	26.7 (24.81–28.64)	37.9 (35.02–40.69)	63.1 (59.87–66.39)
2010, quarter 4	25.1 (23.24–27.04)	43.8 (40.80–46.82)	63.6 (60.19–66.99)
2011, quarter 1	26.4 (24.86–27.93)	43.4 (40.61–46.17)	68.7 (66.01–71.37)
2011, quarter 2	26.9 (25.32–28.46)	41.7 (38.97–44.39)	69.4 (67.00–71.78)
2011, quarter 3	28.2 (26.47–29.87)	41.9 (39.23–44.57)	65.5 (62.59–68.50)
2011, quarter 4	27.5 (25.48–29.47)	43.8 (41.26–46.32)	64.3 (61.52–67.14)
2012, quarter 1	25.5 (23.78–27.25)	43.1 (40.28–46.00)	68.6 (65.62–71.63)
2012, quarter 2	27.9 (26.16–29.55)	42.5 (39.69–45.21)	69.4 (66.84–72.03)
2012, quarter 3	26.6 (24.68–28.54)	43.3 (40.95–45.73)	64.6 (61.86–67.32)
2012, quarter 4	25.2 (23.49–26.84)	41.9 (39.03–44.78)	63.4 (60.28–66.44)
2013, quarter 1	28.7 (26.90–30.51)	47.2 (44.43–49.88)	69.8 (66.84–72.70)
2013, quarter 2	31.5 (29.65–33.44)	48.1 (45.22–50.95)	69.1 (66.21–71.97)
2013, quarter 3	31.2 (29.45–32.96)	46.2 (43.58–48.90)	69.3 (66.30–72.20)
2013, quarter 4	27.0 (25.21–28.71)	44.6 (41.64–47.54)	63.7 (60.59–66.77)
2014, quarter 1	31.2 (29.29–33.06)	45.0 (42.03–47.95)	68.8 (65.96–71.63)
2014, quarter 2	31.6 (29.55–33.72)	48.2 (45.02–51.40)	72.2 (69.54–74.86)
2014, quarter 3	32.7 (30.64–34.67)	45.1 (42.11–48.11)	69.9 (67.35–72.48)
2014, quarter 4	29.3 (27.66–30.96)	43.6 (40.90–46.26)	69.1 (66.38–71.85)
2015, quarter 1	32.5 (30.48–34.43)	50.7 (47.82–53.52)	71.9 (69.48–74.34)
2015, quarter 2	33.3 (31.49–35.03)	47.6 (44.68–50.47)	71.4 (68.76–74.13)
2015, quarter 3	29.7 (27.83–31.62)	48.2 (45.07–51.38)	67.2 (64.35–69.96)
2015, quarter 4	31.2 (28.84–33.52)	46.0 (43.07–48.90)	65.8 (63.05–68.60)
2016, quarter 1	30.5 (28.28–32.69)	44.3 (41.52–47.17)	68.9 (66.52–71.24)

See notes at end of table.

**Data table for Figure 4.1. Percentage of adults aged 18 and over who received an influenza vaccination during the past 12 months, by age group and quarter: United States, 1997–June 2017 (Cont.)**

Year and quarter	18–49	50–64	65 and over
	Percent (95% confidence interval)		
2016, quarter 2	32.7 (31.07–34.43)	46.4 (43.37–49.52)	69.8 (67.11–72.54)
2016, quarter 3	33.3 (31.18–35.39)	47.7 (45.12–50.25)	68.0 (65.15–70.81)
2016, quarter 4	30.6 (28.46–32.73)	42.2 (39.40–44.92)	62.3 (59.49–65.05)
2017, quarter 1	34.1 (31.91–36.44)	47.9 (45.10–50.75)	70.3 (68.12–72.33)
2017, quarter 2	33.6 (31.12–36.19)	45.7 (42.73–48.63)	70.0 (66.72–73.09)

NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Beginning with 2012 data, the National Health Interview Survey transitioned to weights derived from the 2010 census. For 2003–2011 data, weights were derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. For 1997–1999 data, weights were derived from the 1990 census. See [Technical Notes](#) for more details.

DATA SOURCE: NCHS, National Health Interview Survey, 1997–June 2017, Sample Adult Core component.



**Data table for Figure 4.2. Percentage of persons who received an influenza vaccination during the past 12 months, by age group and sex: United States, January–June 2017**

Age (months and years) and sex	Percent	95% confidence interval
6 months–4 years, total	55.5	51.44–59.45
6 months–4 years, male	56.3	51.53–60.92
6 months–4 years, female	54.6	47.44–61.66
5–11, total	49.7	46.50–52.99
5–11, male	49.6	45.78–53.51
5–11, female	49.9	45.46–54.24
12–17, total	40.8	37.72–43.93
12–17, male	41.5	37.36–45.73
12–17, female	40.1	35.61–44.66
6 months–17 years, total	48.0	46.20–49.87
6 months–17 years, male	48.5	45.89–51.07
6 months–17 years, female	47.6	44.99–50.16
18–49, total	33.9	32.35–35.52
18–49, male	28.9	26.63–31.17
18–49, female	38.9	36.79–41.12
50–64, total	46.8	44.84–48.82
50–64, male	43.6	41.08–46.06
50–64, female	49.8	46.88–52.74
65 and over, total	70.2	68.15–72.22
65 and over, male	71.9	69.17–74.60
65 and over, female	68.8	65.92–71.57
6 months and over (crude <sup>1</sup> ), total	45.3	44.32–46.21
6 months and over (crude <sup>1</sup> ), male	42.4	41.06–43.78
6 months and over (crude <sup>1</sup> ), female	48.0	46.60–49.39
18 and over (crude <sup>1</sup> ), total	44.5	43.48–45.48
18 and over (crude <sup>1</sup> ), male	40.6	39.07–42.13
18 and over (crude <sup>1</sup> ), female	48.1	46.55–49.67
65 and over (age–adjusted <sup>2</sup> ), total	71.1	69.16–73.04
65 and over (age–adjusted <sup>2</sup> ), male	73.2	70.73–75.53
65 and over (age–adjusted <sup>2</sup> ), female	69.5	66.65–72.24

<sup>1</sup>Crude estimates are presented.

<sup>2</sup>Estimates are age–adjusted using the projected 2000 U.S. population as the standard population and two age groups: 65–74 and 75 and over.

NOTE: Data are based on household interviews of a sample of the civilian noninstitutionalized population.

DATA SOURCE: NCHS, National Health Interview Survey, January–June 2017, combined Sample Adult and Sample Child Core components.

**Data table for Figure 4.3. Percentage of persons aged 6 months and over who received an influenza vaccination during the past 12 months, by age group and race and ethnicity: United States, January–June 2017**

Age and race and ethnicity	Percent <sup>1</sup>	95% confidence interval
6 months–17 years, Hispanic or Latino	50.2	45.31–55.04
6 months–17 years, not Hispanic or Latino, single race, white	45.5	42.96–48.00
6 months–17 years, not Hispanic or Latino, single race, black	43.2	37.91–48.65
18 and over, Hispanic or Latino	36.7	34.16–39.31
18 and over, not Hispanic or Latino, single race, white	47.4	46.28–48.59
18 and over, not Hispanic or Latino, single race, black	36.3	33.49–39.22

<sup>1</sup>Crude estimates are presented.

NOTE: Data are based on household interviews of a sample of the civilian noninstitutionalized population.

DATA SOURCE: NCHS, National Health Interview Survey, January–June 2017, combined Sample Adult and Sample Child Core components.