


Alcohol consumption

Figure 9.1. Percentage of adults aged 18 and over who had at least 1 heavy drinking day in the past year, by sex: United States, 1997–September 2017



† For 1997–2013, the alcohol consumption estimates presented were for the percentage of adults aged 18 and over who had five or more drinks in 1 day at least once in the past year, regardless of sex. However, in the 2014 National Health Interview Survey (NHIS), the survey questions were changed; male and female respondents were asked about a different quantity of drinks consumed in a day in the past year. As a result, the estimates presented for 2014 and later (dashed line) were for men aged 18 and over who had five or more drinks in 1 day at least once in the past year and for women aged 18 and over who had four or more drinks in 1 day at least once in the past year. Differences observed in estimates for women based on the 2014 and later NHIS may be partially or fully attributable to these changes in the survey questions on alcohol consumption.


NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. The analyses exclude adults with unknown alcohol consumption (about 1% of respondents each year). See [Technical Notes](#) for more details.

SOURCE: NCHS, National Health Interview Survey, 1997–September 2017, Sample Adult Core component.

- For January–September 2017, the percentage of men who had at least 1 heavy drinking day in the past year was 30.6% (95% confidence interval = 29.48%–31.73%). This was lower than, but not significantly different from, the 2016 estimate of 31.6%.
- The percentage of women who had at least one heavy drinking day in the past year was 20.1% (95% confidence interval = 19.02%–21.26%). This was higher than, but not significantly different from, the 2016 estimate of 18.9%.
- The percentage of men who had at least 1 heavy drinking day in the past year decreased, from 31.6% in 1997 to 27.8% in 2006, and then increased to 32.4% in 2009. From 2009 through January–September 2017, the percentage of men who had at least 1 heavy drinking day in the past year remained stable.

- The percentage of women who had at least 1 heavy drinking day in the past year remained stable from 1997 to 2004, increased from 11.2% in 2004 to 14.5% in 2013, and then remained stable from 2014 through January–September 2017.

Figure 9.2. Percentage of adults aged 18 and over who had at least 1 heavy drinking day in the past year, by age group and sex: United States, January–September 2017




NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Heavy drinking days are defined as days in which men consumed five or more drinks and women consumed four or more drinks. The analyses exclude the 1.3% of adults with unknown alcohol consumption. See [Technical Notes](#) for more details.

SOURCE: NCHS, National Health Interview Survey, January–September 2017, Sample Adult Core component.

- For January–September 2017, the percentage of adults aged 18 and over who had at least 1 heavy drinking day in the past year was 25.2% (95% confidence interval = 24.36%–25.98%).
- For both sexes combined, the percentage of adults who had at least 1 heavy drinking day in the past year was highest among adults aged 18–24 (31.7%) and 25–44 (35.9%). The percentage decreased with increasing age after age 45. This pattern was observed for both men and women.
- For adults aged 18 and over and those aged 25–44, 45–64, and 65 and over, men were more likely than women to have had at least 1 heavy drinking day in the past year.

Figure 9.3. Age-sex-adjusted percentage of adults aged 18 and over who had at least 1 heavy drinking day in the past year, by race and ethnicity: United States, January–September 2017



NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Heavy drinking days are defined as days in which men consumed five or more drinks and women consumed four or more drinks. The analyses exclude the 1.3% of adults with unknown alcohol consumption. Estimates are age-sex-adjusted using the projected 2000 U.S. population as the standard population and four age groups: 18–24, 25–44, 45–64, and 65 and over. See [Technical Notes](#) for more details.

SOURCE: NCHS, National Health Interview Survey, January–September 2017, Sample Adult Core component.

- The age-sex-adjusted percentage of adults by race and ethnicity who had at least 1 heavy drinking day in the past year was 22.0% for Hispanic adults, 31.6% for non-Hispanic white adults, and 14.8% for non-Hispanic black adults.
- Non-Hispanic white adults were the most likely to have had at least 1 heavy drinking day in the past year compared with Hispanic and non-Hispanic black adults. Hispanic adults were more likely to have had at least 1 heavy drinking day in the past year compared with non-Hispanic black adults.

Data tables for Figures 9.1–9.3:

Data table for Figure 9.1. Percentage of adults aged 18 and over who had at least 1 heavy drinking day in the past year, by sex: United States, 1997–September 2017

Year and sex	Crude ¹ percent (95% confidence interval)	Age-adjusted ² percent (95% confidence interval)
1997, male	31.6 (30.6–32.5)	30.6 (29.7–31.5)
1997, female	12.1 (11.5–12.7)	12.2 (11.6–12.7)
1998, male	29.9 (28.9–30.9)	29.0 (28.1–29.9)
1998, female	11.4 (10.8–12.1)	11.5 (10.9–12.1)
1999, male	29.8 (28.7–30.9)	29.0 (28.0–30.0)
1999, female	11.7 (11.1–12.4)	11.8 (11.2–12.5)
2000, male	28.9 (27.9–29.9)	28.2 (27.2–29.2)
2000, female	10.6 (10.0–11.2)	10.8 (10.2–11.4)
2001, male	29.2 (28.3–30.2)	28.6 (27.7–29.4)
2001, female	11.6 (11.0–12.3)	11.9 (11.3–12.5)
2002, male	29.2 (28.2–30.3)	28.7 (27.7–29.7)
2002, female	11.5 (10.9–12.1)	11.8 (11.2–12.4)
2003, male	27.8 (26.7–28.8)	27.3 (26.3–28.3)
2003, female	11.3 (10.7–12.0)	11.6 (10.9–12.3)
2004, male	27.7 (26.6–28.8)	27.3 (26.2–28.4)
2004, female	11.2 (10.7–11.9)	11.6 (11.0–12.2)
2005, male	28.1 (27.14–29.15)	27.9 (26.91–28.81)
2005, female	11.6 (10.95–12.28)	12.0 (11.38–12.72)
2006, male	27.8 (26.54–29.01)	27.6 (26.38–28.72)
2006, female	12.1 (11.41–12.86)	12.6 (11.83–13.29)
2007, male	29.0 (27.58–30.44)	28.9 (27.56–30.27)
2007, female	12.4 (11.53–13.32)	13.0 (12.05–13.88)
2008, male	30.8 (29.38–32.24)	30.8 (29.42–32.19)
2008, female	13.8 (12.91–14.72)	14.5 (13.56–15.47)
2009, male	32.4 (31.05–33.80)	32.6 (31.20–33.91)
2009, female	13.8 (13.01–14.66)	14.5 (13.72–15.36)
2010, male	31.5 (30.36–32.66)	31.8 (30.64–32.89)
2010, female	14.6 (13.81–15.42)	15.4 (14.56–16.19)
2011, male	31.2 (30.08–32.28)	31.6 (30.54–32.67)
2011, female	13.6 (12.91–14.38)	14.4 (13.69–15.20)
2012, male	31.4 (30.28–32.52)	31.9 (30.80–33.06)
2012, female	14.2 (13.45–15.01)	15.2 (14.41–16.00)
2013, male	30.9 (29.82–32.01)	31.7 (30.60–32.71)
2013, female	14.5 (13.75–15.25)	15.6 (14.73–16.37)
2014, male ³	31.4 (30.25–32.63)	32.3 (31.13–33.53)
2014, female ³	18.9 (17.96–19.86)	20.2 (19.27–21.20)
2015, male ³	29.9 (28.68–31.18)	30.8 (29.61–32.06)
2015, female ³	17.4 (16.56–18.29)	18.6 (17.70–19.54)
2016, male ³	31.6 (30.46–32.79)	32.7 (31.52–33.84)
2016, female ³	18.9 (18.08–19.73)	20.3 (19.39–21.15)
January–September 2017, male ³	30.6 (29.48–31.73)	31.8 (30.63–32.92)
January–September 2017, female ³	20.1 (19.02–21.26)	21.6 (20.38–22.81)

¹Crude estimates are presented.

²Estimates are age-adjusted using the projected 2000 U.S. population as the standard population and four age groups: 18–24, 25–44, 45–64, and 65 and over.

³For 1997–2013, the alcohol consumption estimates presented were for the percentage of adults aged 18 and over who had five or more drinks in 1 day at least once in the past year, regardless of sex. However, in the 2014 National Health Interview Survey (NHIS), the survey questions were changed; male and female respondents were asked about a different quantity of drinks consumed in a day in the past year. As a result, the estimates presented for 2014 and later were for men aged 18 and over who had five or more drinks in 1 day at least once in the past year and for women aged 18 and over who had four or more drinks in 1 day at least once in the past year. Differences observed in estimates for women based on the 2014 and earlier NHIS may be partially or fully attributable to these changes in the survey questions on alcohol consumption.

NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Beginning with 2012 data, NHIS transitioned to weights derived from the 2010 census. For 2003–2011 data, weights were derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. For 1997–1999 data, weights were derived from the 1990 census. See [Technical Notes](#) for more details.

SOURCE: NCHS, National Health Interview Survey, 1997–September 2017, Sample Adult Core component.

Data table for Figure 9.2. Percentage of adults aged 18 and over who had at least 1 heavy drinking day in the past year, by age group and sex: United States, January–September 2017

Age (years) and sex	Percent	95% confidence interval
18–24 , total	31.7	28.76–34.80
18–24 , male	33.3	28.98–37.94
18–24 , female	30.1	26.57–33.77
25–44 , total	35.9	34.63–37.25
25–44 , male	43.5	41.65–45.37
25–44 , female	28.6	26.73–30.54
45–64 , total	22.2	21.01–23.45
45–64 , male	27.2	25.55–28.97
45–64 , female	17.5	15.86–19.28
65 and over, total	8.0	7.28–8.75
65 and over, male	10.8	9.47–12.31
65 and over, female	5.7	4.88–6.59
18 and over (crude ¹), total	25.2	24.36–25.98
18 and over (crude ¹), male	30.6	29.48–31.73
18 and over (crude ¹), female	20.1	19.02–21.26
18 and over (age-adjusted ²), total	26.5	25.64–27.42
18 and over (age-adjusted ²), male	31.8	30.63–32.92
18 and over (age-adjusted ²), female	21.6	20.38–22.81

¹Crude estimates are presented.

²Estimates are age-adjusted using the projected 2000 U.S. population as the standard population and four age groups: 18–24, 25–44, 45–64, and 65 and over.

NOTE: Data are based on household interviews of a sample of the civilian noninstitutionalized population.

SOURCE: NCHS, National Health Interview Survey, January–September 2017, Sample Adult Core component.

Data table for Figure 9.3. Age-sex-adjusted percentage of adults aged 18 and over who had at least 1 heavy drinking day in the past year, by race and ethnicity: United States, January–September 2017

Race and ethnicity	Percent ¹	95% confidence interval
Hispanic or Latino	22.0	19.60–24.55
Not Hispanic or Latino, single race, white	31.6	30.49–32.76
Not Hispanic or Latino, single race, black	14.8	13.10–16.56

¹Estimates are age-sex-adjusted using the projected 2000 U.S. population as the standard population and four age groups: 18–24, 25–44, 45–64, and 65 and over.

NOTE: Data are based on household interviews of a sample of the civilian noninstitutionalized population.

SOURCE: NCHS, National Health Interview Survey, January–September 2017, Sample Adult Core component.