

Deaths: Leading Causes for 2020

Supplemental Tables

Table I-1. Deaths, percentage of total deaths, and death rates for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020

Table I-2. Infant, neonatal, and postneonatal deaths, percentage of total deaths, and mortality rates for the 10 leading causes of infant death, by bridged race and Hispanic origin and sex: United States, 2020

Table I-3. Estimated population, by age group, bridged race and Hispanic origin, and sex: United States, July 1, 2020

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black non-Hispanic and White non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
All races and origins, both sexes, all ages ⁴				
...	All causes	3,383,729	100.0	1,027.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	696,962	20.6	211.5
2	Malignant neoplasms (C00-C97)	602,350	17.8	182.8
3	COVID-19 (†U07.1)	350,831	10.4	106.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	200,955	5.9	61.0
5	Cerebrovascular diseases (I60-I69)	160,264	4.7	48.6
6	Chronic lower respiratory diseases (J40-J47)	152,657	4.5	46.3
7	Alzheimer disease (G30)	134,242	4.0	40.7
8	Diabetes mellitus (E10-E149)	102,188	3.0	31.0
9	Influenza and pneumonia (J09-J18)	53,544	1.6	16.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52,547	1.6	15.9
...	All other causes (residual)	877,189	25.9	266.2
All races and origins, both sexes, 1-4 years				
...	All causes	3,529	100.0	22.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,153	32.7	7.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	382	10.8	2.5
3	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	311	8.8	2.0
4	Malignant neoplasms (C00-C97)	307	8.7	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	112	3.2	0.7
6	Influenza and pneumonia (J09-J18)	84	2.4	0.5
7	Cerebrovascular diseases (I60-I69)	55	1.6	0.4
8	Certain conditions originating in the perinatal period (P00-P96)	54	1.5	0.3
9	Septicemia (A40-A41)	43	1.2	0.3
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	1.0	0.2
...	All other causes (residual)	993	28.1	6.4
All races and origins, both sexes, 5-9 years				
...	All causes	2,210	100.0	10.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	685	31.0	3.4
2	Malignant neoplasms (C00-C97)	382	17.3	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	171	7.7	0.8
4	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	169	7.6	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	56	2.5	0.3
6	Influenza and pneumonia (J09-J18)	55	2.5	0.3
7	Chronic lower respiratory diseases (J40-J47)	54	2.4	0.3
8	Cerebrovascular diseases (I60-I69)	32	1.4	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	1.3	0.1
10	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	20	0.9	0.1
...	All other causes (residual)	558	25.2	2.8
All races and origins, both sexes, 10-14 years				
...	All causes	3,413	100.0	16.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	881	25.8	4.2
2	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	581	17.0	2.8
3	Malignant neoplasms (C00-C97)	410	12.0	2.0
4	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	285	8.4	1.4
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	150	4.4	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	111	3.3	0.5
7	Chronic lower respiratory diseases (J40-J47)	93	2.7	0.4
8	Influenza and pneumonia (J09-J18)	50	1.5	0.2
8	Diabetes mellitus (E10-E149)	50	1.5	0.2
10	Cerebrovascular diseases (I60-I69)	44	1.3	0.2
...	All other causes (residual)	758	22.2	3.7

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
All races and origins, both sexes, 15-19 years				
...	All causes	12,278	100.0	58.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,654	37.9	22.2
2	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	2,572	20.9	12.3
3	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	2,216	18.0	10.6
4	Malignant neoplasms (C00-C97)	549	4.5	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	294	2.4	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	184	1.5	0.9
7	COVID-19 (†U07.1)	118	1.0	0.6
8	Diabetes mellitus (E10-E149)	93	0.8	0.4
9	Chronic lower respiratory diseases (J40-J47)	90	0.7	0.4
10	Cerebrovascular diseases (I60-I69)	75	0.6	0.4
...	All other causes (residual)	1,433	11.7	6.8
All races and origins, both sexes, 20-24 years				
...	All causes	23,538	100.0	109.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,463	44.5	48.5
2	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	3,894	16.5	18.0
3	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	3,846	16.3	17.8
4	Malignant neoplasms (C00-C97)	757	3.2	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	576	2.4	2.7
6	COVID-19 (†U07.1)	383	1.6	1.8
7	Diabetes mellitus (E10-E149)	219	0.9	1.0
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	200	0.8	0.9
9	Pregnancy, childbirth and the puerperium (O00-O99)	155	0.7	0.7
10	Chronic lower respiratory diseases (J40-J47)	130	0.6	0.6
...	All other causes (residual)	2,915	12.4	13.5
All races and origins, both sexes, 25-34 years				
...	All causes	73,486	100.0	159.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31,315	42.6	68.0
2	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	8,454	11.5	18.4
3	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	7,125	9.7	15.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,984	5.4	8.6
5	Malignant neoplasms (C00-C97)	3,573	4.9	7.8
6	COVID-19 (†U07.1)	2,254	3.1	4.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,631	2.2	3.5
8	Diabetes mellitus (E10-E149)	1,168	1.6	2.5
9	Cerebrovascular diseases (I60-I69)	600	0.8	1.3
10	Pregnancy, childbirth and the puerperium (O00-O99)	594	0.8	1.3
...	All other causes (residual)	12,788	17.4	27.8
All races and origins, both sexes, 35-44 years				
...	All causes	104,490	100.0	248.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31,057	29.7	73.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,177	11.7	28.9
3	Malignant neoplasms (C00-C97)	10,730	10.3	25.5
4	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	7,314	7.0	17.4
5	COVID-19 (†U07.1)	6,079	5.8	14.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,938	4.7	11.7
7	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	4,482	4.3	10.6
8	Diabetes mellitus (E10-E149)	2,904	2.8	6.9
9	Cerebrovascular diseases (I60-I69)	2,008	1.9	4.8
10	Influenza and pneumonia (J09-J18)	1,148	1.1	2.7
...	All other causes (residual)	21,653	20.7	51.4

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
All races and origins, both sexes, 45-54 years				
...	All causes	191,142	100.0	473.5
1	Malignant neoplasms	34,589	18.1	85.7
2	Diseases of heart	34,169	17.9	84.6
3	Accidents (unintentional injuries)	27,819	14.6	68.9
4	COVID-19	16,964	8.9	42.0
5	Chronic liver disease and cirrhosis	9,503	5.0	23.5
6	Diabetes mellitus	7,546	3.9	18.7
7	Intentional self-harm (suicide)	7,249	3.8	18.0
8	Cerebrovascular diseases	5,686	3.0	14.1
9	Chronic lower respiratory diseases	3,538	1.9	8.8
10	Assault (homicide)	2,542	1.3	6.3
...	All other causes	41,537	21.7	102.9
All races and origins, both sexes, 55-64 years				
...	All causes	440,549	100.0	1,038.9
1	Malignant neoplasms	110,243	25.0	260.0
2	Diseases of heart	88,551	20.1	208.8
3	COVID-19	42,090	9.6	99.3
4	Accidents (unintentional injuries)	28,915	6.6	68.2
5	Chronic lower respiratory diseases	18,816	4.3	44.4
6	Diabetes mellitus	18,002	4.1	42.5
7	Chronic liver disease and cirrhosis	16,151	3.7	38.1
8	Cerebrovascular diseases	14,153	3.2	33.4
9	Intentional self-harm (suicide)	7,160	1.6	16.9
10	Influenza and pneumonia	6,295	1.4	14.8
...	All other causes	90,173	20.5	212.7
All races and origins, both sexes, 65 years and older				
...	All causes	2,509,396	100.0	4,508.5
1	Diseases of heart	556,665	22.2	1,000.1
2	Malignant neoplasms	440,753	17.6	791.9
3	COVID-19	282,836	11.3	508.2
4	Cerebrovascular diseases	137,392	5.5	246.8
5	Alzheimer disease	132,741	5.3	238.5
6	Chronic lower respiratory diseases	128,712	5.1	231.2
7	Diabetes mellitus	72,194	2.9	129.7
8	Accidents (unintentional injuries)	62,796	2.5	112.8
9	Nephritis, nephrotic syndrome and nephrosis	42,675	1.7	76.7
10	Influenza and pneumonia	42,511	1.7	76.4
...	All other causes	610,121	24.3	1,096.2
All races and origins, both sexes, 65-74 years				
...	All causes	674,507	100.0	2,072.3
1	Malignant neoplasms	175,464	26.0	539.1
2	Diseases of heart	134,827	20.0	414.2
3	COVID-19	76,277	11.3	234.3
4	Chronic lower respiratory diseases	38,559	5.7	118.5
5	Diabetes mellitus	27,213	4.0	83.6
6	Cerebrovascular diseases	26,363	3.9	81.0
7	Accidents (unintentional injuries)	18,609	2.8	57.2
8	Chronic liver disease and cirrhosis	11,895	1.8	36.5
9	Nephritis, nephrotic syndrome and nephrosis	11,461	1.7	35.2
10	Influenza and pneumonia	10,460	1.6	32.1
...	All other causes	143,379	21.3	440.5

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
All races and origins, both sexes, 75–84 years				
...	All causes	822,084	100.0	4,997.0
1	Diseases of heart (I00–I09,I11,I13,I20–I51)	167,346	20.4	1,017.2
2	Malignant neoplasms (C00–C97)	162,876	19.8	990.0
3	COVID-19 (†U07.1)	97,030	11.8	589.8
4	Chronic lower respiratory diseases (J40–J47)	49,363	6.0	300.1
5	Cerebrovascular diseases (I60–I69)	43,252	5.3	262.9
6	Alzheimer disease (G30)	37,728	4.6	229.3
7	Diabetes mellitus (E10–E149)	25,445	3.1	154.7
8	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	18,646	2.3	113.3
9	Parkinson disease (G20–G21)	17,323	2.1	105.3
10	Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)	14,696	1.8	89.3
...	All other causes (residual)	188,379	22.9	1,145.1
All races and origins, both sexes, 85 years and older				
...	All causes	1,012,805	100.0	15,210.9
1	Diseases of heart (I00–I09,I11,I13,I20–I51)	254,492	25.1	3,822.1
2	COVID-19 (†U07.1)	109,529	10.8	1,645.0
3	Malignant neoplasms (C00–C97)	102,413	10.1	1,538.1
4	Alzheimer disease (G30)	85,713	8.5	1,287.3
5	Cerebrovascular diseases (I60–I69)	67,777	6.7	1,017.9
6	Chronic lower respiratory diseases (J40–J47)	40,790	4.0	612.6
7	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	25,541	2.5	383.6
8	Diabetes mellitus (E10–E149)	19,536	1.9	293.4
9	Influenza and pneumonia (J09–J18)	18,219	1.8	273.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16,663	1.6	250.3
...	All other causes (residual)	272,132	26.9	4,087.0
All races and origins, male, all ages ⁴				
...	All causes	1,769,884	100.0	1,090.8
1	Diseases of heart (I00–I09,I11,I13,I20–I51)	382,776	21.6	235.9
2	Malignant neoplasms (C00–C97)	317,731	18.0	195.8
3	COVID-19 (†U07.1)	192,512	10.9	118.6
4	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	133,205	7.5	82.1
5	Chronic lower respiratory diseases (J40–J47)	72,942	4.1	45.0
6	Cerebrovascular diseases (I60–I69)	69,637	3.9	42.9
7	Diabetes mellitus (E10–E149)	57,532	3.3	35.5
8	Alzheimer disease (G30)	41,273	2.3	25.4
9	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	36,551	2.1	22.5
10	Chronic liver disease and cirrhosis (K70,K73–K74)	32,546	1.8	20.1
...	All other causes (residual)	433,179	24.5	267.0
All races and origins, male, 1–4 years				
...	All causes	2,026	100.0	25.5
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	733	36.2	9.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	197	9.7	2.5
3	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	181	8.9	2.3
4	Malignant neoplasms (C00–C97)	159	7.8	2.0
5	Diseases of heart (I00–I09,I11,I13,I20–I51)	58	2.9	0.7
6	Influenza and pneumonia (J09–J18)	51	2.5	0.6
7	Certain conditions originating in the perinatal period (P00–P96)	31	1.5	0.4
8	Cerebrovascular diseases (I60–I69)	30	1.5	0.4
9	Septicemia (A40–A41)	20	1.0	0.3
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00–D48)	15	0.7	*
...	All other causes (residual)	551	27.2	6.9

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
All races and origins, male, 5–9 years				
...	All causes	1,245	100.0	12.0
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	413	33.2	4.0
2	Malignant neoplasms (C00–C97)	203	16.3	2.0
3	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	88	7.1	0.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	88	7.1	0.9
5	Chronic lower respiratory diseases (J40–J47)	32	2.6	0.3
6	Diseases of heart (I00–I09,I11,I13,I20–I51)	31	2.5	0.3
7	Influenza and pneumonia (J09–J18)	28	2.2	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00–D48)	17	1.4	*
9	Cerebrovascular diseases (I60–I69)	14	1.1	*
10	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	13	1.0	*
...	All other causes (residual)	318	25.5	3.1
All races and origins, male, 10–14 years				
...	All causes	2,117	100.0	20.0
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	565	26.7	5.3
2	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	377	17.8	3.6
3	Malignant neoplasms (C00–C97)	219	10.3	2.1
4	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	210	9.9	2.0
5	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	81	3.8	0.8
6	Chronic lower respiratory diseases (J40–J47)	63	3.0	0.6
7	Diseases of heart (I00–I09,I11,I13,I20–I51)	57	2.7	0.5
8	Diabetes mellitus (E10–E149)	25	1.2	0.2
8	Influenza and pneumonia (J09–J18)	25	1.2	0.2
10	Cerebrovascular diseases (I60–I69)	23	1.1	0.2
...	All other causes (residual)	472	22.3	4.5
All races and origins, male, 15–19 years				
...	All causes	8,951	100.0	83.7
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	3,350	37.4	31.3
2	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	2,235	25.0	20.9
3	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	1,689	18.9	15.8
4	Malignant neoplasms (C00–C97)	329	3.7	3.1
5	Diseases of heart (I00–I09,I11,I13,I20–I51)	188	2.1	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	99	1.1	0.9
7	COVID-19 (†U07.1)	59	0.7	0.6
8	Diabetes mellitus (E10–E149)	55	0.6	0.5
9	Chronic lower respiratory diseases (J40–J47)	48	0.5	0.4
10	Legal intervention (Y35,Y89.0)	41	0.5	0.4
...	All other causes (residual)	858	9.6	8.0
All races and origins, male, 20–24 years				
...	All causes	17,533	100.0	158.9
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	7,770	44.3	70.4
2	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	3,359	19.2	30.4
3	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	3,170	18.1	28.7
4	Malignant neoplasms (C00–C97)	445	2.5	4.0
5	Diseases of heart (I00–I09,I11,I13,I20–I51)	360	2.1	3.3
6	COVID-19 (†U07.1)	250	1.4	2.3
7	Diabetes mellitus (E10–E149)	141	0.8	1.3
8	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	111	0.6	1.0
9	Chronic lower respiratory diseases (J40–J47)	73	0.4	0.7
10	Legal intervention (Y35,Y89.0)	68	0.4	0.6
...	All other causes (residual)	1,786	10.2	16.2

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
All races and origins, male, 25–34 years				
...	All causes	51,832	100.0	221.1
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	22,993	44.4	98.1
2	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	6,882	13.3	29.4
3	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	6,088	11.7	26.0
4	Diseases of heart (I00–I09,I11,I13,I20–I51)	2,661	5.1	11.4
5	Malignant neoplasms (C00–C97)	1,707	3.3	7.3
6	COVID-19 (†U07.1)	1,466	2.8	6.3
7	Chronic liver disease and cirrhosis (K70,K73–K74)	1,028	2.0	4.4
8	Diabetes mellitus (E10–E149)	718	1.4	3.1
9	Human immunodeficiency virus (HIV) disease (B20–B24)	367	0.7	1.6
10	Cerebrovascular diseases (I60–I69)	338	0.7	1.4
10	Influenza and pneumonia (J09–J18)	338	0.7	1.4
...	All other causes (residual)	7,246	14.0	30.9
All races and origins, male, 35–44 years				
...	All causes	68,494	100.0	325.5
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	22,499	32.8	106.9
2	Diseases of heart (I00–I09,I11,I13,I20–I51)	8,362	12.2	39.7
3	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	5,723	8.4	27.2
4	Malignant neoplasms (C00–C97)	4,707	6.9	22.4
5	COVID-19 (†U07.1)	4,068	5.9	19.3
6	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	3,658	5.3	17.4
7	Chronic liver disease and cirrhosis (K70,K73–K74)	3,086	4.5	14.7
8	Diabetes mellitus (E10–E149)	1,844	2.7	8.8
9	Cerebrovascular diseases (I60–I69)	1,204	1.8	5.7
10	Influenza and pneumonia (J09–J18)	631	0.9	3.0
...	All other causes (residual)	12,712	18.6	60.4
All races and origins, male, 45–54 years				
...	All causes	119,844	100.0	601.5
1	Diseases of heart (I00–I09,I11,I13,I20–I51)	24,066	20.1	120.8
2	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	19,846	16.6	99.6
3	Malignant neoplasms (C00–C97)	16,483	13.8	82.7
4	COVID-19 (†U07.1)	11,385	9.5	57.1
5	Chronic liver disease and cirrhosis (K70,K73–K74)	6,079	5.1	30.5
6	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	5,514	4.6	27.7
7	Diabetes mellitus (E10–E149)	4,858	4.1	24.4
8	Cerebrovascular diseases (I60–I69)	3,311	2.8	16.6
9	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	1,974	1.6	9.9
10	Chronic lower respiratory diseases (J40–J47)	1,667	1.4	8.4
...	All other causes (residual)	24,661	20.6	123.8
All races and origins, male, 55–64 years				
...	All causes	271,127	100.0	1,323.3
1	Diseases of heart (I00–I09,I11,I13,I20–I51)	61,700	22.8	301.1
2	Malignant neoplasms (C00–C97)	59,412	21.9	290.0
3	COVID-19 (†U07.1)	26,997	10.0	131.8
4	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	20,589	7.6	100.5
5	Diabetes mellitus (E10–E149)	11,330	4.2	55.3
6	Chronic liver disease and cirrhosis (K70,K73–K74)	10,755	4.0	52.5
7	Chronic lower respiratory diseases (J40–J47)	9,499	3.5	46.4
8	Cerebrovascular diseases (I60–I69)	8,368	3.1	40.8
9	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	5,539	2.0	27.0
10	Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)	3,655	1.3	17.8
...	All other causes (residual)	53,283	19.7	260.1

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
All races and origins, male, 65 years and older				
...	All causes	1,215,777	100.0	4,898.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	285,159	23.5	1,148.8
2	Malignant neoplasms (C00-C97)	234,044	19.3	942.9
3	COVID-19 (†U07.1)	148,222	12.2	597.2
4	Chronic lower respiratory diseases (J40-J47)	60,925	5.0	245.5
5	Cerebrovascular diseases (I60-I69)	56,204	4.6	226.4
6	Alzheimer disease (G30)	40,640	3.3	163.7
7	Diabetes mellitus (E10-E149)	38,556	3.2	155.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33,742	2.8	135.9
9	Parkinson disease (G20-G21)	23,837	2.0	96.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21,456	1.8	86.4
...	All other causes (residual)	272,992	22.5	1,099.8
All races and origins, male, 65-74 years				
...	All causes	391,999	100.0	2,581.7
1	Malignant neoplasms (C00-C97)	96,939	24.7	638.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	86,738	22.1	571.3
3	COVID-19 (†U07.1)	47,014	12.0	309.6
4	Chronic lower respiratory diseases (J40-J47)	20,247	5.2	133.3
5	Diabetes mellitus (E10-E149)	16,176	4.1	106.5
6	Cerebrovascular diseases (I60-I69)	14,124	3.6	93.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,543	3.2	82.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	7,536	1.9	49.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,230	1.6	41.0
10	Influenza and pneumonia (J09-J18)	5,916	1.5	39.0
...	All other causes (residual)	78,536	20.0	517.2
All races and origins, male, 75-84 years				
...	All causes	428,858	100.0	5,937.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	93,977	21.9	1,301.0
2	Malignant neoplasms (C00-C97)	87,107	20.3	1,205.9
3	COVID-19 (†U07.1)	54,719	12.8	757.5
4	Chronic lower respiratory diseases (J40-J47)	23,977	5.6	331.9
5	Cerebrovascular diseases (I60-I69)	20,064	4.7	277.8
6	Alzheimer disease (G30)	13,939	3.3	193.0
7	Diabetes mellitus (E10-E149)	13,788	3.2	190.9
8	Parkinson disease (G20-G21)	11,014	2.6	152.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,394	2.4	143.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,671	1.8	106.2
...	All other causes (residual)	92,208	21.5	1,276.5
All races and origins, male, 85 years and older				
...	All causes	394,920	100.0	16,354.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	104,444	26.4	4,325.4
2	Malignant neoplasms (C00-C97)	49,998	12.7	2,070.6
3	COVID-19 (†U07.1)	46,489	11.8	1,925.3
4	Alzheimer disease (G30)	23,009	5.8	952.9
5	Cerebrovascular diseases (I60-I69)	22,016	5.6	911.8
6	Chronic lower respiratory diseases (J40-J47)	16,701	4.2	691.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,805	2.7	447.5
8	Parkinson disease (G20-G21)	8,722	2.2	361.2
9	Diabetes mellitus (E10-E149)	8,592	2.2	355.8
10	Influenza and pneumonia (J09-J18)	8,134	2.1	336.9
...	All other causes (residual)	96,010	24.3	3,976.1

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
All races and origins, female, all ages ⁴				
...	All causes	1,613,845	100.0	965.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	314,186	19.5	187.9
2	Malignant neoplasms (C00-C97)	284,619	17.6	170.2
3	COVID-19 (†U07.1)	158,319	9.8	94.7
4	Alzheimer disease (G30)	92,969	5.8	55.6
5	Cerebrovascular diseases (I60-I69)	90,627	5.6	54.2
6	Chronic lower respiratory diseases (J40-J47)	79,715	4.9	47.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	67,750	4.2	40.5
8	Diabetes mellitus (E10-E149)	44,656	2.8	26.7
9	Influenza and pneumonia (J09-J18)	25,799	1.6	15.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25,254	1.6	15.1
...	All other causes (residual)	429,951	26.6	257.1
All races and origins, female, 1-4 years				
...	All causes	1,503	100.0	19.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	420	27.9	5.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	185	12.3	2.4
3	Malignant neoplasms (C00-C97)	148	9.8	1.9
4	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	130	8.6	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	3.6	0.7
6	Influenza and pneumonia (J09-J18)	33	2.2	0.4
7	Cerebrovascular diseases (I60-I69)	25	1.7	0.3
8	Certain conditions originating in the perinatal period (P00-P96)	23	1.5	0.3
8	Septicemia (A40-A41)	23	1.5	0.3
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.3	0.3
...	All other causes (residual)	442	29.4	5.8
All races and origins, female, 5-9 years				
...	All causes	965	100.0	9.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	272	28.2	2.7
2	Malignant neoplasms (C00-C97)	179	18.5	1.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	83	8.6	0.8
4	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	81	8.4	0.8
5	Influenza and pneumonia (J09-J18)	27	2.8	0.3
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	2.6	0.3
7	Chronic lower respiratory diseases (J40-J47)	22	2.3	0.2
8	Cerebrovascular diseases (I60-I69)	18	1.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.1	*
10	Septicemia (A40-A41)	9	0.9	*
...	All other causes (residual)	238	24.7	2.4
All races and origins, female, 10-14 years				
...	All causes	1,296	100.0	12.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	316	24.4	3.1
2	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	204	15.7	2.0
3	Malignant neoplasms (C00-C97)	191	14.7	1.9
4	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	75	5.8	0.7
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	69	5.3	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	4.2	0.5
7	Chronic lower respiratory diseases (J40-J47)	30	2.3	0.3
8	Influenza and pneumonia (J09-J18)	25	1.9	0.2
8	Diabetes mellitus (E10-E149)	25	1.9	0.2
10	Cerebrovascular diseases (I60-I69)	21	1.6	0.2
...	All other causes (residual)	286	22.1	2.8

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
All races and origins, female, 15-19 years				
...	All causes	3,327	100.0	32.4
1	Accidents (unintentional injuries)..... (V01-X59,Y85-Y86)	1,304	39.2	12.7
2	Intentional self-harm (suicide)..... (†U03,X60-X84,Y87.0)	527	15.8	5.1
3	Assault (homicide)..... (†U01-†U02.9,X85-Y09,Y87.1)	337	10.1	3.3
4	Malignant neoplasms..... (C00-C97)	220	6.6	2.1
5	Diseases of heart..... (I00-I09,I11,I13,I20-I51)	106	3.2	1.0
6	Congenital malformations, deformations and chromosomal abnormalities..... (Q00-Q99)	85	2.6	0.8
7	COVID-19..... (†U07.1)	59	1.8	0.6
8	Cerebrovascular diseases..... (I60-I69)	45	1.4	0.4
9	Chronic lower respiratory diseases..... (J40-J47)	42	1.3	0.4
10	Diabetes mellitus..... (E10-E149)	38	1.1	0.4
...	All other causes..... (residual)	564	17.0	5.5
All races and origins, female, 20-24 years				
...	All causes.....	6,005	100.0	56.9
1	Accidents (unintentional injuries)..... (V01-X59,Y85-Y86)	2,693	44.8	25.5
2	Intentional self-harm (suicide)..... (†U03,X60-X84,Y87.0)	676	11.3	6.4
3	Assault (homicide)..... (†U01-†U02.9,X85-Y09,Y87.1)	535	8.9	5.1
4	Malignant neoplasms..... (C00-C97)	312	5.2	3.0
5	Diseases of heart..... (I00-I09,I11,I13,I20-I51)	216	3.6	2.0
6	Pregnancy, childbirth and the puerperium..... (O00-O99)	155	2.6	1.5
7	COVID-19..... (†U07.1)	133	2.2	1.3
8	Congenital malformations, deformations and chromosomal abnormalities..... (Q00-Q99)	89	1.5	0.8
9	Diabetes mellitus..... (E10-E149)	78	1.3	0.7
10	Influenza and pneumonia..... (J09-J18)	58	1.0	0.5
...	All other causes..... (residual)	1,060	17.7	10.0
All races and origins, female, 25-34 years				
...	All causes.....	21,654	100.0	95.7
1	Accidents (unintentional injuries)..... (V01-X59,Y85-Y86)	8,322	38.4	36.8
2	Malignant neoplasms..... (C00-C97)	1,866	8.6	8.2
3	Intentional self-harm (suicide)..... (†U03,X60-X84,Y87.0)	1,572	7.3	6.9
4	Diseases of heart..... (I00-I09,I11,I13,I20-I51)	1,323	6.1	5.8
5	Assault (homicide)..... (†U01-†U02.9,X85-Y09,Y87.1)	1,037	4.8	4.6
6	COVID-19..... (†U07.1)	788	3.6	3.5
7	Chronic liver disease and cirrhosis..... (K70,K73-K74)	603	2.8	2.7
8	Pregnancy, childbirth and the puerperium..... (O00-O99)	594	2.7	2.6
9	Diabetes mellitus..... (E10-E149)	450	2.1	2.0
10	Cerebrovascular diseases..... (I60-I69)	262	1.2	1.2
...	All other causes..... (residual)	4,837	22.3	21.4
All races and origins, female, 35-44 years				
...	All causes.....	35,996	100.0	170.7
1	Accidents (unintentional injuries)..... (V01-X59,Y85-Y86)	8,558	23.8	40.6
2	Malignant neoplasms..... (C00-C97)	6,023	16.7	28.6
3	Diseases of heart..... (I00-I09,I11,I13,I20-I51)	3,815	10.6	18.1
4	COVID-19..... (†U07.1)	2,011	5.6	9.5
5	Chronic liver disease and cirrhosis..... (K70,K73-K74)	1,852	5.1	8.8
6	Intentional self-harm (suicide)..... (†U03,X60-X84,Y87.0)	1,591	4.4	7.5
7	Diabetes mellitus..... (E10-E149)	1,060	2.9	5.0
8	Assault (homicide)..... (†U01-†U02.9,X85-Y09,Y87.1)	824	2.3	3.9
9	Cerebrovascular diseases..... (I60-I69)	804	2.2	3.8
10	Influenza and pneumonia..... (J09-J18)	517	1.4	2.5
...	All other causes..... (residual)	8,941	24.8	42.4

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
All races and origins, female, 45-54 years				
...	All causes	71,298	100.0	348.8
1	Malignant neoplasms (C00-C97)	18,106	25.4	88.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,103	14.2	49.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,973	11.2	39.0
4	COVID-19 (†U07.1)	5,579	7.8	27.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,424	4.8	16.8
6	Diabetes mellitus (E10-E149)	2,688	3.8	13.1
7	Cerebrovascular diseases (I60-I69)	2,375	3.3	11.6
8	Chronic lower respiratory diseases (J40-J47)	1,871	2.6	9.2
9	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	1,735	2.4	8.5
10	Septicemia (A40-A41)	1,215	1.7	5.9
...	All other causes (residual)	16,229	22.8	79.4
All races and origins, female, 55-64 years				
...	All causes	169,422	100.0	773.1
1	Malignant neoplasms (C00-C97)	50,831	30.0	232.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,851	15.8	122.5
3	COVID-19 (†U07.1)	15,093	8.9	68.9
4	Chronic lower respiratory diseases (J40-J47)	9,317	5.5	42.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,326	4.9	38.0
6	Diabetes mellitus (E10-E149)	6,672	3.9	30.4
7	Cerebrovascular diseases (I60-I69)	5,785	3.4	26.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	5,396	3.2	24.6
9	Septicemia (A40-A41)	2,934	1.7	13.4
10	Influenza and pneumonia (J09-J18)	2,641	1.6	12.1
...	All other causes (residual)	35,576	21.0	162.3
All races and origins, female, 65 years and older				
...	All causes	1,293,619	100.0	4,194.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	271,506	21.0	880.4
2	Malignant neoplasms (C00-C97)	206,709	16.0	670.3
3	COVID-19 (†U07.1)	134,614	10.4	436.5
4	Alzheimer disease (G30)	92,101	7.1	298.7
5	Cerebrovascular diseases (I60-I69)	81,188	6.3	263.3
6	Chronic lower respiratory diseases (J40-J47)	67,787	5.2	219.8
7	Diabetes mellitus (E10-E149)	33,638	2.6	109.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29,054	2.2	94.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21,219	1.6	68.8
10	Influenza and pneumonia (J09-J18)	21,115	1.6	68.5
...	All other causes (residual)	334,688	25.9	1,085.3
All races and origins, female, 65-74 years				
...	All causes	282,508	100.0	1,626.8
1	Malignant neoplasms (C00-C97)	78,525	27.8	452.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,089	17.0	276.9
3	COVID-19 (†U07.1)	29,263	10.4	168.5
4	Chronic lower respiratory diseases (J40-J47)	18,312	6.5	105.4
5	Cerebrovascular diseases (I60-I69)	12,239	4.3	70.5
6	Diabetes mellitus (E10-E149)	11,037	3.9	63.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,066	2.1	34.9
8	Alzheimer disease (G30)	5,608	2.0	32.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,231	1.9	30.1
10	Septicemia (A40-A41)	4,693	1.7	27.0
...	All other causes (residual)	63,445	22.5	365.3

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
All races and origins, female, 75–84 years				
...	All causes	393,226	100.0	4,261.1
1	Malignant neoplasms (C00–C97)	75,769	19.3	821.1
2	Diseases of heart (I00–I09,I11,I13,I20–I51)	73,369	18.7	795.0
3	COVID-19 (†U07.1)	42,311	10.8	458.5
4	Chronic lower respiratory diseases (J40–J47)	25,386	6.5	275.1
5	Alzheimer disease (G30)	23,789	6.0	257.8
6	Cerebrovascular diseases (I60–I69)	23,188	5.9	251.3
7	Diabetes mellitus (E10–E149)	11,657	3.0	126.3
8	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	8,252	2.1	89.4
9	Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)	7,025	1.8	76.1
10	Influenza and pneumonia (J09–J18)	6,486	1.6	70.3
...	All other causes (residual)	95,994	24.4	1,040.2
All races and origins, female, 85 years and older				
...	All causes	617,885	100.0	14,560.0
1	Diseases of heart (I00–I09,I11,I13,I20–I51)	150,048	24.3	3,535.8
2	COVID-19 (†U07.1)	63,040	10.2	1,485.5
3	Alzheimer disease (G30)	62,704	10.1	1,477.6
4	Malignant neoplasms (C00–C97)	52,415	8.5	1,235.1
5	Cerebrovascular diseases (I60–I69)	45,761	7.4	1,078.3
6	Chronic lower respiratory diseases (J40–J47)	24,089	3.9	567.6
7	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	14,736	2.4	347.2
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,394	1.8	268.5
9	Diabetes mellitus (E10–E149)	10,944	1.8	257.9
10	Influenza and pneumonia (J09–J18)	10,085	1.6	237.6
...	All other causes (residual)	172,669	27.9	4,068.8
All races, Hispanic, both sexes, all ages ⁴				
...	All causes	305,708	100.0	498.6
1	COVID-19 (†U07.1)	65,237	21.3	106.4
2	Diseases of heart (I00–I09,I11,I13,I20–I51)	48,323	15.8	78.8
3	Malignant neoplasms (C00–C97)	43,942	14.4	71.7
4	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	23,152	7.6	37.8
5	Cerebrovascular diseases (I60–I69)	13,370	4.4	21.8
6	Diabetes mellitus (E10–E149)	12,832	4.2	20.9
7	Alzheimer disease (G30)	9,867	3.2	16.1
8	Chronic liver disease and cirrhosis (K70,K73–K74)	8,049	2.6	13.1
9	Chronic lower respiratory diseases (J40–J47)	5,949	1.9	9.7
10	Influenza and pneumonia (J09–J18)	4,875	1.6	8.0
...	All other causes (residual)	70,112	22.9	114.4
All races, Hispanic, both sexes, 1–4 years				
...	All causes	742	100.0	18.4
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	240	32.3	6.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	90	12.1	2.2
3	Malignant neoplasms (C00–C97)	78	10.5	1.9
4	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	54	7.3	1.3
5	Diseases of heart (I00–I09,I11,I13,I20–I51)	22	3.0	0.5
6	Influenza and pneumonia (J09–J18)	19	2.6	*
7	Certain conditions originating in the perinatal period (P00–P96)	11	1.5	*
7	Septicemia (A40–A41)	11	1.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00–D48)	8	1.1	*
10	Cerebrovascular diseases (I60–I69)	6	0.8	*
...	All other causes (residual)	203	27.4	5.0

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
All races, Hispanic, both sexes, 5-9 years				
...	All causes	468	100.0	9.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	144	30.8	2.8
2	Malignant neoplasms (C00-C97)	96	20.5	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	7.5	0.7
4	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	25	5.3	0.5
5	Cerebrovascular diseases (I60-I69)	12	2.6	*
6	Influenza and pneumonia (J09-J18)	10	2.1	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	2.1	*
8	COVID-19 (†U07.1)	8	1.7	*
9	Chronic lower respiratory diseases (J40-J47)	6	1.3	*
9	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	6	1.3	*
...	All other causes (residual)	116	24.8	2.2
All races, Hispanic, both sexes, 10-14 years				
...	All causes	772	100.0	14.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	215	27.8	4.0
2	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	120	15.5	2.2
3	Malignant neoplasms (C00-C97)	113	14.6	2.1
4	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	50	6.5	0.9
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	5.2	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	3.0	0.4
7	Cerebrovascular diseases (I60-I69)	12	1.6	*
7	Influenza and pneumonia (J09-J18)	12	1.6	*
9	COVID-19 (†U07.1)	10	1.3	*
10	Diabetes mellitus (E10-E149)	8	1.0	*
10	Chronic lower respiratory diseases (J40-J47)	8	1.0	*
...	All other causes (residual)	161	20.9	3.0
All races, Hispanic, both sexes, 15-19 years				
...	All causes	2,803	100.0	54.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,184	42.2	23.2
2	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	549	19.6	10.8
3	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	429	15.3	8.4
4	Malignant neoplasms (C00-C97)	127	4.5	2.5
5	COVID-19 (†U07.1)	50	1.8	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	1.7	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	1.3	0.7
8	Cerebrovascular diseases (I60-I69)	23	0.8	0.5
9	Diabetes mellitus (E10-E149)	19	0.7	*
10	Chronic lower respiratory diseases (J40-J47)	15	0.5	*
...	All other causes (residual)	324	11.6	6.3
All races, Hispanic, both sexes, 20-24 years				
...	All causes	4,906	100.0	100.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,348	47.9	47.8
2	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	678	13.8	13.8
3	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	660	13.5	13.4
4	Malignant neoplasms (C00-C97)	231	4.7	4.7
5	COVID-19 (†U07.1)	165	3.4	3.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	103	2.1	2.1
7	Diabetes mellitus (E10-E149)	37	0.8	0.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	0.8	0.8
7	Cerebrovascular diseases (I60-I69)	37	0.8	0.8
10	Pregnancy, childbirth and the puerperium (O00-O99)	27	0.6	0.6
...	All other causes (residual)	583	11.9	11.9

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
All races, Hispanic, both sexes, 25-34 years				
...	All causes	12,403	100.0	128.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,129	41.4	53.3
2	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	1,267	10.2	13.2
3	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	1,063	8.6	11.0
4	COVID-19 (†U07.1)	947	7.6	9.8
5	Malignant neoplasms (C00-C97)	704	5.7	7.3
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	527	4.2	5.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	352	2.8	3.7
8	Diabetes mellitus (E10-E149)	167	1.3	1.7
9	Pregnancy, childbirth and the puerperium (O00-O99)	120	1.0	1.2
10	Influenza and pneumonia (J09-J18)	114	0.9	1.2
...	All other causes (residual)	2,013	16.2	20.9
All races, Hispanic, both sexes, 35-44 years				
...	All causes	17,252	100.0	197.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,259	24.7	48.7
2	COVID-19 (†U07.1)	2,866	16.6	32.8
3	Malignant neoplasms (C00-C97)	1,873	10.9	21.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,479	8.6	16.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,057	6.1	12.1
6	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	810	4.7	9.3
7	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	781	4.5	8.9
8	Diabetes mellitus (E10-E149)	479	2.8	5.5
9	Cerebrovascular diseases (I60-I69)	379	2.2	4.3
10	Influenza and pneumonia (J09-J18)	197	1.1	2.3
...	All other causes (residual)	3,072	17.8	35.1
All races, Hispanic, both sexes, 45-54 years				
...	All causes	28,652	100.0	395.2
1	COVID-19 (†U07.1)	7,010	24.5	96.7
2	Malignant neoplasms (C00-C97)	4,476	15.6	61.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,535	12.3	48.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,318	11.6	45.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,676	5.8	23.1
6	Diabetes mellitus (E10-E149)	1,239	4.3	17.1
7	Cerebrovascular diseases (I60-I69)	950	3.3	13.1
8	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	583	2.0	8.0
9	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	408	1.4	5.6
10	Influenza and pneumonia (J09-J18)	379	1.3	5.2
...	All other causes (residual)	5,078	17.7	70.0
All races, Hispanic, both sexes, 55-64 years				
...	All causes	47,899	100.0	911.3
1	COVID-19 (†U07.1)	12,853	26.8	244.5
2	Malignant neoplasms (C00-C97)	9,036	18.9	171.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,156	14.9	136.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,717	5.7	51.7
5	Diabetes mellitus (E10-E149)	2,457	5.1	46.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,266	4.7	43.1
7	Cerebrovascular diseases (I60-I69)	1,537	3.2	29.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	776	1.6	14.8
9	Influenza and pneumonia (J09-J18)	721	1.5	13.7
10	Chronic lower respiratory diseases (J40-J47)	631	1.3	12.0
...	All other causes (residual)	7,749	16.2	147.4

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
All races, Hispanic, both sexes, 65 years and older				
...	All causes	185,562	100.0	3,801.1
1	COVID-19 (†U07.1)	41,309	22.3	846.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,368	19.1	724.5
3	Malignant neoplasms (C00-C97)	27,194	14.7	557.0
4	Cerebrovascular diseases (I60-I69)	10,283	5.5	210.6
5	Alzheimer disease (G30)	9,722	5.2	199.1
6	Diabetes mellitus (E10-E149)	8,424	4.5	172.6
7	Chronic lower respiratory diseases (J40-J47)	4,935	2.7	101.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,430	1.8	70.3
9	Influenza and pneumonia (J09-J18)	3,372	1.8	69.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,324	1.8	68.1
...	All other causes (residual)	38,201	20.6	782.5
All races, Hispanic, both sexes, 65-74 years				
...	All causes	58,239	100.0	1,955.3
1	COVID-19 (†U07.1)	16,281	28.0	546.6
2	Malignant neoplasms (C00-C97)	11,362	19.5	381.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,351	16.1	313.9
4	Diabetes mellitus (E10-E149)	3,266	5.6	109.6
5	Cerebrovascular diseases (I60-I69)	2,159	3.7	72.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,610	2.8	54.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,278	2.2	42.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,124	1.9	37.7
9	Chronic lower respiratory diseases (J40-J47)	1,119	1.9	37.6
10	Influenza and pneumonia (J09-J18)	971	1.7	32.6
...	All other causes (residual)	9,718	16.7	326.3
All races, Hispanic, both sexes, 75-84 years				
...	All causes	62,231	100.0	4,547.5
1	COVID-19 (†U07.1)	14,443	23.2	1,055.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,306	18.2	826.2
3	Malignant neoplasms (C00-C97)	9,941	16.0	726.4
4	Cerebrovascular diseases (I60-I69)	3,263	5.2	238.4
5	Diabetes mellitus (E10-E149)	3,030	4.9	221.4
6	Alzheimer disease (G30)	2,677	4.3	195.6
7	Chronic lower respiratory diseases (J40-J47)	1,744	2.8	127.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,155	1.9	84.4
9	Influenza and pneumonia (J09-J18)	1,118	1.8	81.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,035	1.7	75.6
...	All other causes (residual)	12,519	20.1	914.8
All races, Hispanic, both sexes, 85 years and older				
...	All causes	65,092	100.0	12,172.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,711	22.6	2,750.9
2	COVID-19 (†U07.1)	10,585	16.3	1,979.4
3	Alzheimer disease (G30)	6,361	9.8	1,189.5
4	Malignant neoplasms (C00-C97)	5,891	9.1	1,101.6
5	Cerebrovascular diseases (I60-I69)	4,861	7.5	909.0
6	Diabetes mellitus (E10-E149)	2,128	3.3	397.9
7	Chronic lower respiratory diseases (J40-J47)	2,072	3.2	387.5
8	Influenza and pneumonia (J09-J18)	1,283	2.0	239.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,268	1.9	237.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,117	1.7	208.9
...	All other causes (residual)	14,815	22.8	2,770.4

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
All races, Hispanic, male, all ages ⁴				
...	All causes	175,585	100.0	567.8
1	COVID-19 (†U07.1)	40,830	23.3	132.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,700	15.8	89.6
3	Malignant neoplasms (C00-C97)	22,700	12.9	73.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,606	10.0	56.9
5	Diabetes mellitus (E10-E149)	7,136	4.1	23.1
6	Cerebrovascular diseases (I60-I69)	6,374	3.6	20.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	5,526	3.1	17.9
8	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	3,701	2.1	12.0
9	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	3,277	1.9	10.6
10	Alzheimer disease (G30)	3,065	1.7	9.9
...	All other causes (residual)	37,670	21.5	121.8
All races, Hispanic, male, 1-4 years				
...	All causes	418	100.0	20.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	148	35.4	7.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	11.0	2.2
3	Malignant neoplasms (C00-C97)	38	9.1	1.9
4	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	35	8.4	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	2.9	*
5	Influenza and pneumonia (J09-J18)	12	2.9	*
7	Cerebrovascular diseases (I60-I69)	6	1.4	*
8	Certain conditions originating in the perinatal period (P00-P96)	5	1.2	*
8	Septicemia (A40-A41)	5	1.2	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
10	COVID-19 (†U07.1)	3	0.7	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
...	All other causes (residual)	102	24.4	5.0
All races, Hispanic, male, 5-9 years				
...	All causes	253	100.0	9.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73	28.9	2.8
2	Malignant neoplasms (C00-C97)	48	19.0	1.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	7.5	*
4	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	17	6.7	*
5	COVID-19 (†U07.1)	7	2.8	*
5	Influenza and pneumonia (J09-J18)	7	2.8	*
7	Cerebrovascular diseases (I60-I69)	6	2.4	*
8	Chronic lower respiratory diseases (J40-J47)	5	2.0	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	2.0	*
8	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	5	2.0	*
...	All other causes (residual)	61	24.1	2.3
All races, Hispanic, male, 10-14 years				
...	All causes	440	100.0	16.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	128	29.1	4.7
2	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	65	14.8	2.4
3	Malignant neoplasms (C00-C97)	58	13.2	2.1
4	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	28	6.4	1.0
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	4.5	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	2.3	*
7	Cerebrovascular diseases (I60-I69)	7	1.6	*
7	Chronic lower respiratory diseases (J40-J47)	7	1.6	*
9	Diabetes mellitus (E10-E149)	6	1.4	*
9	Influenza and pneumonia (J09-J18)	6	1.4	*
...	All other causes (residual)	105	23.9	3.8

See footnotes at end of table.

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
All races, Hispanic, male, 15-19 years				
...	All causes	2,076	100.0	79.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	885	42.6	34.0
2	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	472	22.7	18.1
3	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	313	15.1	12.0
4	Malignant neoplasms (C00-C97)	86	4.1	3.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	1.4	1.2
6	COVID-19 (†U07.1)	27	1.3	1.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	1.0	0.8
8	Diabetes mellitus (E10-E149)	9	0.4	*
8	Legal intervention (Y35,Y89.0)	9	0.4	*
10	Cerebrovascular diseases (I60-I69)	8	0.4	*
10	Chronic lower respiratory diseases (J40-J47)	8	0.4	*
...	All other causes (residual)	209	10.1	8.0
All races, Hispanic, male, 20-24 years				
...	All causes	3,837	100.0	152.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,915	49.9	76.2
2	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	587	15.3	23.4
3	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	551	14.4	21.9
4	Malignant neoplasms (C00-C97)	151	3.9	6.0
5	COVID-19 (†U07.1)	110	2.9	4.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	1.7	2.5
7	Diabetes mellitus (E10-E149)	27	0.7	1.1
8	Cerebrovascular diseases (I60-I69)	24	0.6	1.0
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	0.5	*
10	Legal intervention (Y35,Y89.0)	14	0.4	*
...	All other causes (residual)	375	9.8	14.9
All races, Hispanic, male, 25-34 years				
...	All causes	9,311	100.0	185.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,148	44.5	82.7
2	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	1,037	11.1	20.7
3	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	920	9.9	18.4
4	COVID-19 (†U07.1)	667	7.2	13.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	390	4.2	7.8
6	Malignant neoplasms (C00-C97)	371	4.0	7.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	250	2.7	5.0
8	Diabetes mellitus (E10-E149)	100	1.1	2.0
9	Influenza and pneumonia (J09-J18)	73	0.8	1.5
10	Legal intervention (Y35,Y89.0)	70	0.8	1.4
...	All other causes (residual)	1,285	13.8	25.6
All races, Hispanic, male, 35-44 years				
...	All causes	12,372	100.0	274.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,429	27.7	76.0
2	COVID-19 (†U07.1)	2,144	17.3	47.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,132	9.1	25.1
4	Malignant neoplasms (C00-C97)	815	6.6	18.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	794	6.4	17.6
6	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	660	5.3	14.6
7	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	658	5.3	14.6
8	Diabetes mellitus (E10-E149)	313	2.5	6.9
9	Cerebrovascular diseases (I60-I69)	265	2.1	5.9
10	Influenza and pneumonia (J09-J18)	125	1.0	2.8
...	All other causes (residual)	2,037	16.5	45.1

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
All races, Hispanic, male, 45–54 years				
...	All causes	19,654	100.0	538.1
1	COVID-19 (†U07.1)	5,188	26.4	142.0
2	Diseases of heart (I00–I09,I11,I13,I20–I51)	2,686	13.7	73.5
3	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	2,625	13.4	71.9
4	Malignant neoplasms (C00–C97)	2,043	10.4	55.9
5	Chronic liver disease and cirrhosis (K70,K73–K74)	1,285	6.5	35.2
6	Diabetes mellitus (E10–E149)	839	4.3	23.0
7	Cerebrovascular diseases (I60–I69)	623	3.2	17.1
8	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	478	2.4	13.1
9	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	327	1.7	9.0
10	Influenza and pneumonia (J09–J18)	252	1.3	6.9
...	All other causes (residual)	3,308	16.8	90.6
All races, Hispanic, male, 55–64 years				
...	All causes	31,415	100.0	1,220.6
1	COVID-19 (†U07.1)	8,902	28.3	345.9
2	Diseases of heart (I00–I09,I11,I13,I20–I51)	5,192	16.5	201.7
3	Malignant neoplasms (C00–C97)	4,804	15.3	186.7
4	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	2,135	6.8	83.0
5	Chronic liver disease and cirrhosis (K70,K73–K74)	1,684	5.4	65.4
6	Diabetes mellitus (E10–E149)	1,561	5.0	60.7
7	Cerebrovascular diseases (I60–I69)	962	3.1	37.4
8	Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)	491	1.6	19.1
9	Influenza and pneumonia (J09–J18)	460	1.5	17.9
10	Chronic lower respiratory diseases (J40–J47)	337	1.1	13.1
...	All other causes (residual)	4,887	15.6	189.9
All races, Hispanic, male, 65 years and older				
...	All causes	93,432	100.0	4,373.5
1	COVID-19 (†U07.1)	23,768	25.4	1,112.6
2	Diseases of heart (I00–I09,I11,I13,I20–I51)	18,153	19.4	849.7
3	Malignant neoplasms (C00–C97)	14,280	15.3	668.4
4	Cerebrovascular diseases (I60–I69)	4,392	4.7	205.6
5	Diabetes mellitus (E10–E149)	4,280	4.6	200.3
6	Alzheimer disease (G30)	3,002	3.2	140.5
7	Chronic lower respiratory diseases (J40–J47)	2,488	2.7	116.5
8	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	2,009	2.2	94.0
9	Influenza and pneumonia (J09–J18)	1,698	1.8	79.5
10	Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)	1,650	1.8	77.2
...	All other causes (residual)	17,712	19.0	829.1
All races, Hispanic, male, 65–74 years				
...	All causes	35,047	100.0	2,562.1
1	COVID-19 (†U07.1)	10,471	29.9	765.5
2	Malignant neoplasms (C00–C97)	6,272	17.9	458.5
3	Diseases of heart (I00–I09,I11,I13,I20–I51)	6,108	17.4	446.5
4	Diabetes mellitus (E10–E149)	1,909	5.4	139.6
5	Cerebrovascular diseases (I60–I69)	1,227	3.5	89.7
6	Chronic liver disease and cirrhosis (K70,K73–K74)	1,005	2.9	73.5
7	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	928	2.6	67.8
8	Chronic lower respiratory diseases (J40–J47)	637	1.8	46.6
9	Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)	608	1.7	44.4
10	Influenza and pneumonia (J09–J18)	549	1.6	40.1
...	All other causes (residual)	5,333	15.2	389.9

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
All races, Hispanic, male, 75–84 years				
...	All causes	32,212	100.0	5,615.5
1	COVID-19 (†U07.1)	8,265	25.7	1,440.8
2	Diseases of heart (I00–I09,I11,I13,I20–I51)	6,085	18.9	1,060.8
3	Malignant neoplasms (C00–C97)	5,193	16.1	905.3
4	Diabetes mellitus (E10–E149)	1,505	4.7	262.4
5	Cerebrovascular diseases (I60–I69)	1,500	4.7	261.5
6	Alzheimer disease (G30)	940	2.9	163.9
7	Chronic lower respiratory diseases (J40–J47)	925	2.9	161.3
8	Parkinson disease (G20–G21)	614	1.9	107.0
9	Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)	578	1.8	100.8
10	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	574	1.8	100.1
...	All other causes (residual)	6,033	18.7	1,051.7
All races, Hispanic, male, 85 years and older				
...	All causes	26,173	100.0	13,436.2
1	Diseases of heart (I00–I09,I11,I13,I20–I51)	5,960	22.8	3,059.6
2	COVID-19 (†U07.1)	5,032	19.2	2,583.2
3	Malignant neoplasms (C00–C97)	2,815	10.8	1,445.1
4	Alzheimer disease (G30)	1,771	6.8	909.2
5	Cerebrovascular diseases (I60–I69)	1,665	6.4	854.7
6	Chronic lower respiratory diseases (J40–J47)	926	3.5	475.4
7	Diabetes mellitus (E10–E149)	866	3.3	444.6
8	Influenza and pneumonia (J09–J18)	582	2.2	298.8
9	Parkinson disease (G20–G21)	537	2.1	275.7
10	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	507	1.9	260.3
...	All other causes (residual)	5,512	21.1	2,829.6
All races, Hispanic, female, all ages ⁴				
...	All causes	130,123	100.0	428.2
1	COVID-19 (†U07.1)	24,407	18.8	80.3
2	Malignant neoplasms (C00–C97)	21,242	16.3	69.9
3	Diseases of heart (I00–I09,I11,I13,I20–I51)	20,623	15.8	67.9
4	Cerebrovascular diseases (I60–I69)	6,996	5.4	23.0
5	Alzheimer disease (G30)	6,802	5.2	22.4
6	Diabetes mellitus (E10–E149)	5,696	4.4	18.7
7	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	5,546	4.3	18.2
8	Chronic lower respiratory diseases (J40–J47)	2,918	2.2	9.6
9	Chronic liver disease and cirrhosis (K70,K73–K74)	2,523	1.9	8.3
10	Influenza and pneumonia (J09–J18)	2,214	1.7	7.3
...	All other causes (residual)	31,156	23.9	102.5
All races, Hispanic, female, 1–4 years				
...	All causes	324	100.0	16.4
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	92	28.4	4.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	44	13.6	2.2
3	Malignant neoplasms (C00–C97)	40	12.3	2.0
4	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	19	5.9	*
5	Diseases of heart (I00–I09,I11,I13,I20–I51)	10	3.1	*
6	Influenza and pneumonia (J09–J18)	7	2.2	*
7	Certain conditions originating in the perinatal period (P00–P96)	6	1.9	*
7	Septicemia (A40–A41)	6	1.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00–D48)	5	1.5	*
10	Diseases of appendix (K35–K38)	2	0.6	*
10	COVID-19 (†U07.1)	2	0.6	*
10	Complications of medical and surgical care (Y40–Y84,Y88)	2	0.6	*
...	All other causes (residual)	89	27.5	4.5

See footnotes at end of table.

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
All races, Hispanic, female, 5-9 years				
...	All causes	215	100.0	8.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	71	33.0	2.8
2	Malignant neoplasms (C00-C97)	48	22.3	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	7.4	*
4	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	8	3.7	*
5	Cerebrovascular diseases (I60-I69)	6	2.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	2.8	*
7	Influenza and pneumonia (J09-J18)	3	1.4	*
8	Septicemia (A40-A41)	2	0.9	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.5	*
9	Anemias (D50-D64)	1	0.5	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	0.5	*
9	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	1	0.5	*
9	COVID-19 (†U07.1)	1	0.5	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
9	Diseases of appendix (K35-K38)	1	0.5	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.5	*
...	All other causes (residual)	47	21.9	1.9
All races, Hispanic, female, 10-14 years				
...	All causes	332	100.0	12.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	87	26.2	3.3
2	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	55	16.6	2.1
2	Malignant neoplasms (C00-C97)	55	16.6	2.1
4	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	22	6.6	0.8
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	6.0	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	3.9	*
7	Influenza and pneumonia (J09-J18)	6	1.8	*
8	Cerebrovascular diseases (I60-I69)	5	1.5	*
8	COVID-19 (†U07.1)	5	1.5	*
10	Certain conditions originating in the perinatal period (P00-P96)	3	0.9	*
...	All other causes (residual)	61	18.4	2.3
All races, Hispanic, female, 15-19 years				
...	All causes	727	100.0	29.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	299	41.1	11.9
2	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	116	16.0	4.6
3	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	77	10.6	3.1
4	Malignant neoplasms (C00-C97)	41	5.6	1.6
5	COVID-19 (†U07.1)	23	3.2	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	2.3	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	2.2	*
8	Cerebrovascular diseases (I60-I69)	15	2.1	*
9	Diabetes mellitus (E10-E149)	10	1.4	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	7	1.0	*
10	Chronic lower respiratory diseases (J40-J47)	7	1.0	*
...	All other causes (residual)	99	13.6	4.0

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
All races, Hispanic, female, 20–24 years				
...	All causes	1,069	100.0	44.6
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	433	40.5	18.1
2	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	109	10.2	4.6
3	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	91	8.5	3.8
4	Malignant neoplasms (C00–C97)	80	7.5	3.3
5	COVID-19 (†U07.1)	55	5.1	2.3
6	Diseases of heart (I00–I09,I11,I13,I20–I51)	39	3.6	1.6
7	Pregnancy, childbirth and the puerperium (O00–O99)	27	2.5	1.1
8	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	18	1.7	*
9	Cerebrovascular diseases (I60–I69)	13	1.2	*
10	Chronic lower respiratory diseases (J40–J47)	11	1.0	*
10	Influenza and pneumonia (J09–J18)	11	1.0	*
...	All other causes (residual)	182	17.0	7.6
All races, Hispanic, female, 25–34 years				
...	All causes	3,092	100.0	67.0
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	981	31.7	21.3
2	Malignant neoplasms (C00–C97)	333	10.8	7.2
3	COVID-19 (†U07.1)	280	9.1	6.1
4	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	230	7.4	5.0
5	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	143	4.6	3.1
6	Diseases of heart (I00–I09,I11,I13,I20–I51)	137	4.4	3.0
7	Pregnancy, childbirth and the puerperium (O00–O99)	120	3.9	2.6
8	Chronic liver disease and cirrhosis (K70,K73–K74)	102	3.3	2.2
9	Diabetes mellitus (E10–E149)	67	2.2	1.5
10	Cerebrovascular diseases (I60–I69)	41	1.3	0.9
10	Influenza and pneumonia (J09–J18)	41	1.3	0.9
...	All other causes (residual)	617	20.0	13.4
All races, Hispanic, female, 35–44 years				
...	All causes	4,880	100.0	115.4
1	Malignant neoplasms (C00–C97)	1,058	21.7	25.0
2	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	830	17.0	19.6
3	COVID-19 (†U07.1)	722	14.8	17.1
4	Diseases of heart (I00–I09,I11,I13,I20–I51)	347	7.1	8.2
5	Chronic liver disease and cirrhosis (K70,K73–K74)	263	5.4	6.2
6	Diabetes mellitus (E10–E149)	166	3.4	3.9
7	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	150	3.1	3.5
8	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	123	2.5	2.9
9	Cerebrovascular diseases (I60–I69)	114	2.3	2.7
10	Pregnancy, childbirth and the puerperium (O00–O99)	81	1.7	1.9
...	All other causes (residual)	1,026	21.0	24.3

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
All races, Hispanic, female, 45-54 years				
...	All causes	8,998	100.0	250.1
1	Malignant neoplasms (C00-C97)	2,433	27.0	67.6
2	COVID-19 (†U07.1)	1,822	20.2	50.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	849	9.4	23.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	693	7.7	19.3
5	Diabetes mellitus (E10-E149)	400	4.4	11.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	391	4.3	10.9
7	Cerebrovascular diseases (I60-I69)	327	3.6	9.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	135	1.5	3.8
9	Influenza and pneumonia (J09-J18)	127	1.4	3.5
10	Septicemia (A40-A41)	121	1.3	3.4
...	All other causes (residual)	1,700	18.9	47.3
All races, Hispanic, female, 55-64 years				
...	All causes	16,484	100.0	614.6
1	Malignant neoplasms (C00-C97)	4,232	25.7	157.8
2	COVID-19 (†U07.1)	3,951	24.0	147.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,964	11.9	73.2
4	Diabetes mellitus (E10-E149)	896	5.4	33.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	582	3.5	21.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	582	3.5	21.7
7	Cerebrovascular diseases (I60-I69)	575	3.5	21.4
8	Chronic lower respiratory diseases (J40-J47)	294	1.8	11.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	285	1.7	10.6
10	Influenza and pneumonia (J09-J18)	261	1.6	9.7
...	All other causes (residual)	2,862	17.4	106.7
All races, Hispanic, female, 65 years and older				
...	All causes	92,130	100.0	3,355.7
1	COVID-19 (†U07.1)	17,541	19.0	638.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,215	18.7	627.0
3	Malignant neoplasms (C00-C97)	12,914	14.0	470.4
4	Alzheimer disease (G30)	6,720	7.3	244.8
5	Cerebrovascular diseases (I60-I69)	5,891	6.4	214.6
6	Diabetes mellitus (E10-E149)	4,144	4.5	150.9
7	Chronic lower respiratory diseases (J40-J47)	2,447	2.7	89.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,674	1.8	61.0
8	Influenza and pneumonia (J09-J18)	1,674	1.8	61.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,600	1.7	58.3
...	All other causes (residual)	20,310	22.0	739.8
All races, Hispanic, female, 65-74 years				
...	All causes	23,192	100.0	1,439.9
1	COVID-19 (†U07.1)	5,810	25.1	360.7
2	Malignant neoplasms (C00-C97)	5,090	21.9	316.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,243	14.0	201.3
4	Diabetes mellitus (E10-E149)	1,357	5.9	84.3
5	Cerebrovascular diseases (I60-I69)	932	4.0	57.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	605	2.6	37.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	516	2.2	32.0
8	Chronic lower respiratory diseases (J40-J47)	482	2.1	29.9
9	Influenza and pneumonia (J09-J18)	422	1.8	26.2
10	Alzheimer disease (G30)	393	1.7	24.4
...	All other causes (residual)	4,342	18.7	269.6

See footnotes at end of table.

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
All races, Hispanic, female, 75–84 years				
...	All causes	30,019	100.0	3,776.7
1	COVID-19 (†U07.1)	6,178	20.6	777.3
2	Diseases of heart (I00–I09,I11,I13,I20–I51)	5,221	17.4	656.9
3	Malignant neoplasms (C00–C97)	4,748	15.8	597.4
4	Cerebrovascular diseases (I60–I69)	1,763	5.9	221.8
5	Alzheimer disease (G30)	1,737	5.8	218.5
6	Diabetes mellitus (E10–E149)	1,525	5.1	191.9
7	Chronic lower respiratory diseases (J40–J47)	819	2.7	103.0
8	Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)	577	1.9	72.6
9	Influenza and pneumonia (J09–J18)	551	1.8	69.3
10	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	461	1.5	58.0
...	All other causes (residual)	6,439	21.4	810.1
All races, Hispanic, female, 85 years and older				
...	All causes	38,919	100.0	11,447.7
1	Diseases of heart (I00–I09,I11,I13,I20–I51)	8,751	22.5	2,574.0
2	COVID-19 (†U07.1)	5,553	14.3	1,633.4
3	Alzheimer disease (G30)	4,590	11.8	1,350.1
4	Cerebrovascular diseases (I60–I69)	3,196	8.2	940.1
5	Malignant neoplasms (C00–C97)	3,076	7.9	904.8
6	Diabetes mellitus (E10–E149)	1,262	3.2	371.2
7	Chronic lower respiratory diseases (J40–J47)	1,146	2.9	337.1
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	827	2.1	243.3
9	Influenza and pneumonia (J09–J18)	701	1.8	206.2
10	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	610	1.6	179.4
...	All other causes (residual)	9,207	23.7	2,708.2
American Indian and Alaska Native, non-Hispanic, both sexes, all ages ⁴				
...	All causes	26,755	100.0	967.1
1	COVID-19 (†U07.1)	4,436	16.6	160.4
2	Diseases of heart (I00–I09,I11,I13,I20–I51)	3,868	14.5	139.8
3	Malignant neoplasms (C00–C97)	3,385	12.7	122.4
4	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	2,670	10.0	96.5
5	Chronic liver disease and cirrhosis (K70,K73–K74)	1,522	5.7	55.0
6	Diabetes mellitus (E10–E149)	1,356	5.1	49.0
7	Chronic lower respiratory diseases (J40–J47)	861	3.2	31.1
8	Cerebrovascular diseases (I60–I69)	848	3.2	30.7
9	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	663	2.5	24.0
10	Influenza and pneumonia (J09–J18)	451	1.7	16.3
...	All other causes (residual)	6,695	25.0	242.0
American Indian and Alaska Native, non-Hispanic, both sexes, 1–4 years				
...	All causes	70	100.0	48.4
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	25	35.7	17.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	11	15.7	*
3	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	8	11.4	*
4	Malignant neoplasms (C00–C97)	3	4.3	*
5	Cerebrovascular diseases (I60–I69)	2	2.9	*
5	Influenza and pneumonia (J09–J18)	2	2.9	*
5	Diseases of heart (I00–I09,I11,I13,I20–I51)	2	2.9	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00–D48)	1	1.4	*
8	COVID-19 (†U07.1)	1	1.4	*
...	All other causes (residual)	15	21.4	*

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
American Indian and Alaska Native, non-Hispanic, both sexes, 5-9 years				
...	All causes	41	100.0	20.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	36.6	*
2	Malignant neoplasms (C00-C97)	7	17.1	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	7.3	*
3	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	3	7.3	*
5	Cerebrovascular diseases (I60-I69)	2	4.9	*
6	Chronic lower respiratory diseases (J40-J47)	1	2.4	*
6	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	1	2.4	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.4	*
...	All other causes (residual)	8	19.5	*
American Indian and Alaska Native, non-Hispanic, both sexes, 10-14 years				
...	All causes	57	100.0	27.8
1	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	20	35.1	9.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	22.8	*
3	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	4	7.0	*
4	Malignant neoplasms (C00-C97)	3	5.3	*
4	Chronic lower respiratory diseases (J40-J47)	3	5.3	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	5.3	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	3.5	*
8	Anemias (D50-D64)	1	1.8	*
8	Influenza and pneumonia (J09-J18)	1	1.8	*
8	Pneumonitis due to solids and liquids (J69)	1	1.8	*
8	Cerebrovascular diseases (I60-I69)	1	1.8	*
...	All other causes (residual)	5	8.8	*
American Indian and Alaska Native, non-Hispanic, both sexes, 15-19 years				
...	All causes	215	100.0	105.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	78	36.3	38.4
2	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	70	32.6	34.4
3	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	25	11.6	12.3
4	Malignant neoplasms (C00-C97)	4	1.9	*
4	Chronic lower respiratory diseases (J40-J47)	4	1.9	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	1.9	*
7	COVID-19 (†U07.1)	3	1.4	*
8	Influenza and pneumonia (J09-J18)	2	0.9	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	2	0.9	*
8	Diabetes mellitus (E10-E149)	2	0.9	*
...	All other causes (residual)	21	9.8	10.3
American Indian and Alaska Native, non-Hispanic, both sexes, 20-24 years				
...	All causes	416	100.0	198.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	196	47.1	93.6
2	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	103	24.8	49.2
3	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	40	9.6	19.1
4	COVID-19 (†U07.1)	13	3.1	*
5	Malignant neoplasms (C00-C97)	8	1.9	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	1.2	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	5	1.2	*
8	Cerebrovascular diseases (I60-I69)	3	0.7	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.5	*
...	All other causes (residual)	38	9.1	18.2

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
American Indian and Alaska Native, non-Hispanic, both sexes, 25-34 years				
...	All causes	1,755	100.0	407.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	609	34.7	141.5
2	Chronic liver disease and cirrhosis (K70,K73-K74)	214	12.2	49.7
3	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	207	11.8	48.1
4	COVID-19 (†U07.1)	127	7.2	29.5
5	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	117	6.7	27.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	3.4	13.7
7	Malignant neoplasms (C00-C97)	28	1.6	6.5
8	Diabetes mellitus (E10-E149)	25	1.4	5.8
9	Septicemia (A40-A41)	14	0.8	*
10	Influenza and pneumonia (J09-J18)	13	0.7	*
...	All other causes (residual)	342	19.5	79.5
American Indian and Alaska Native, non-Hispanic, both sexes, 35-44 years				
...	All causes	2,214	100.0	645.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	549	24.8	159.9
2	Chronic liver disease and cirrhosis (K70,K73-K74)	338	15.3	98.5
3	COVID-19 (†U07.1)	221	10.0	64.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	179	8.1	52.1
5	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	112	5.1	32.6
6	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	99	4.5	28.8
7	Malignant neoplasms (C00-C97)	98	4.4	28.5
8	Diabetes mellitus (E10-E149)	78	3.5	22.7
9	Influenza and pneumonia (J09-J18)	26	1.2	7.6
10	Septicemia (A40-A41)	22	1.0	6.4
...	All other causes (residual)	492	22.2	143.3
American Indian and Alaska Native, non-Hispanic, both sexes, 45-54 years				
...	All causes	3,063	100.0	974.7
1	COVID-19 (†U07.1)	470	15.3	149.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	455	14.9	144.8
3	Chronic liver disease and cirrhosis (K70,K73-K74)	383	12.5	121.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	360	11.8	114.6
5	Malignant neoplasms (C00-C97)	272	8.9	86.6
6	Diabetes mellitus (E10-E149)	205	6.7	65.2
7	Cerebrovascular diseases (I60-I69)	78	2.5	24.8
8	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	72	2.4	22.9
9	Influenza and pneumonia (J09-J18)	52	1.7	16.5
10	Chronic lower respiratory diseases (J40-J47)	40	1.3	12.7
...	All other causes (residual)	676	22.1	215.1
American Indian and Alaska Native, non-Hispanic, both sexes, 55-64 years				
...	All causes	5,054	100.0	1,519.4
1	COVID-19 (†U07.1)	897	17.7	269.7
2	Malignant neoplasms (C00-C97)	822	16.3	247.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	771	15.3	231.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	360	7.1	108.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	359	7.1	107.9
6	Diabetes mellitus (E10-E149)	309	6.1	92.9
7	Chronic lower respiratory diseases (J40-J47)	188	3.7	56.5
8	Cerebrovascular diseases (I60-I69)	130	2.6	39.1
9	Influenza and pneumonia (J09-J18)	79	1.6	23.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	78	1.5	23.4
...	All other causes (residual)	1,061	21.0	319.0

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
American Indian and Alaska Native, non-Hispanic, both sexes, 65 years and older				
...	All causes	13,646	100.0	3,890.2
1	COVID-19 (†U07.1)	2,704	19.8	770.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,489	18.2	709.6
3	Malignant neoplasms (C00-C97)	2,138	15.7	609.5
4	Diabetes mellitus (E10-E149)	737	5.4	210.1
5	Chronic lower respiratory diseases (J40-J47)	610	4.5	173.9
6	Cerebrovascular diseases (I60-I69)	609	4.5	173.6
7	Alzheimer disease (G30)	429	3.1	122.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	341	2.5	97.2
9	Influenza and pneumonia (J09-J18)	271	2.0	77.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	243	1.8	69.3
...	All other causes (residual)	3,075	22.5	876.6
American Indian and Alaska Native, non-Hispanic, both sexes, 65-74 years				
...	All causes	5,551	100.0	2,488.9
1	COVID-19 (†U07.1)	1,207	21.7	541.2
2	Malignant neoplasms (C00-C97)	1,051	18.9	471.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	935	16.8	419.2
4	Diabetes mellitus (E10-E149)	359	6.5	161.0
5	Chronic lower respiratory diseases (J40-J47)	243	4.4	109.0
6	Cerebrovascular diseases (I60-I69)	178	3.2	79.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	173	3.1	77.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	162	2.9	72.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	93	1.7	41.7
10	Influenza and pneumonia (J09-J18)	87	1.6	39.0
...	All other causes (residual)	1,063	19.1	476.6
American Indian and Alaska Native, non-Hispanic, both sexes, 75-84 years				
...	All causes	4,842	100.0	5,075.0
1	COVID-19 (†U07.1)	953	19.7	998.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	914	18.9	958.0
3	Malignant neoplasms (C00-C97)	773	16.0	810.2
4	Chronic lower respiratory diseases (J40-J47)	263	5.4	275.7
5	Diabetes mellitus (E10-E149)	258	5.3	270.4
6	Cerebrovascular diseases (I60-I69)	233	4.8	244.2
7	Alzheimer disease (G30)	162	3.3	169.8
8	Influenza and pneumonia (J09-J18)	91	1.9	95.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	88	1.8	92.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	81	1.7	84.9
...	All other causes (residual)	1,026	21.2	1,075.4
American Indian and Alaska Native, non-Hispanic, both sexes, 85 years and older				
...	All causes	3,253	100.0	10,058.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	640	19.7	1,978.9
2	COVID-19 (†U07.1)	544	16.7	1,682.0
3	Malignant neoplasms (C00-C97)	314	9.7	970.9
4	Alzheimer disease (G30)	220	6.8	680.2
5	Cerebrovascular diseases (I60-I69)	198	6.1	612.2
6	Diabetes mellitus (E10-E149)	120	3.7	371.0
7	Chronic lower respiratory diseases (J40-J47)	104	3.2	321.6
8	Influenza and pneumonia (J09-J18)	93	2.9	287.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	2.5	247.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	69	2.1	213.3
...	All other causes (residual)	871	26.8	2,693.1

Table I–1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*]

Rank ¹	Cause of death (based on ICD–10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
American Indian and Alaska Native, non-Hispanic, male, all ages ⁴				
...	All causes	14,827	100.0	1,094.3
1	COVID-19 (†U07.1)	2,424	16.3	178.9
2	Diseases of heart (I00–I09,I11,I13,I20–I51)	2,334	15.7	172.3
3	Malignant neoplasms (C00–C97)	1,790	12.1	132.1
4	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	1,729	11.7	127.6
5	Chronic liver disease and cirrhosis (K70,K73–K74)	802	5.4	59.2
6	Diabetes mellitus (E10–E149)	759	5.1	56.0
7	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	512	3.5	37.8
8	Chronic lower respiratory diseases (J40–J47)	436	2.9	32.2
9	Cerebrovascular diseases (I60–I69)	376	2.5	27.7
10	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	300	2.0	22.1
...	All other causes (residual)	3,365	22.7	248.3
American Indian and Alaska Native, non-Hispanic, male, 1–4 years				
...	All causes	41	100.0	55.8
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	16	39.0	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	5	12.2	*
2	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	5	12.2	*
4	Diseases of heart (I00–I09,I11,I13,I20–I51)	1	2.4	*
4	COVID-19 (†U07.1)	1	2.4	*
4	Malignant neoplasms (C00–C97)	1	2.4	*
...	All other causes (residual)	12	29.3	*
American Indian and Alaska Native, non-Hispanic, male, 5–9 years				
...	All causes	19	100.0	*
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	8	42.1	*
2	Malignant neoplasms (C00–C97)	4	21.1	*
3	Cerebrovascular diseases (I60–I69)	1	5.3	*
3	Chronic lower respiratory diseases (J40–J47)	1	5.3	*
3	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	1	5.3	*
3	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	1	5.3	*
3	Diseases of heart (I00–I09,I11,I13,I20–I51)	1	5.3	*
...	All other causes (residual)	2	10.5	*
American Indian and Alaska Native, non-Hispanic, male, 10–14 years				
...	All causes	35	100.0	33.8
1	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	12	34.3	*
2	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	9	25.7	*
3	Chronic lower respiratory diseases (J40–J47)	3	8.6	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	2	5.7	*
5	Anemias (D50–D64)	1	2.9	*
5	Cerebrovascular diseases (I60–I69)	1	2.9	*
5	Influenza and pneumonia (J09–J18)	1	2.9	*
5	Pneumonitis due to solids and liquids (J69)	1	2.9	*
5	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	1	2.9	*
5	Malignant neoplasms (C00–C97)	1	2.9	*
...	All other causes (residual)	3	8.6	*

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
American Indian and Alaska Native, non-Hispanic, male, 15-19 years				
...	All causes	144	100.0	139.9
1	Intentional self-harm (suicide)..... (†U03,X60-X84,Y87.0)	51	35.4	49.5
2	Accidents (unintentional injuries)..... (V01-X59,Y85-Y86)	49	34.0	47.6
3	Assault (homicide)..... (†U01-†U02.9,X85-Y09,Y87.1)	21	14.6	20.4
4	Congenital malformations, deformations and chromosomal abnormalities..... (Q00-Q99)	3	2.1	*
4	Chronic lower respiratory diseases..... (J40-J47)	3	2.1	*
6	Diabetes mellitus..... (E10-E149)	2	1.4	*
6	Influenza and pneumonia..... (J09-J18)	2	1.4	*
6	Malignant neoplasms..... (C00-C97)	2	1.4	*
9	Viral hepatitis..... (B15-B19)	1	0.7	*
9	Legal intervention..... (Y35,Y89.0)	1	0.7	*
9	COVID-19..... (†U07.1)	1	0.7	*
...	All other causes..... (residual)	8	5.6	*
American Indian and Alaska Native, non-Hispanic, male, 20-24 years				
...	All causes	293	100.0	276.9
1	Accidents (unintentional injuries)..... (V01-X59,Y85-Y86)	130	44.4	122.9
2	Intentional self-harm (suicide)..... (†U03,X60-X84,Y87.0)	81	27.6	76.6
3	Assault (homicide)..... (†U01-†U02.9,X85-Y09,Y87.1)	31	10.6	29.3
4	COVID-19..... (†U07.1)	12	4.1	*
5	Malignant neoplasms..... (C00-C97)	5	1.7	*
6	Diseases of heart..... (I00-I09,I11,I13,I20-I51)	4	1.4	*
7	Cerebrovascular diseases..... (I60-I69)	2	0.7	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior..... (D00-D48)	1	0.3	*
8	Complications of medical and surgical care..... (Y40-Y84,Y88)	1	0.3	*
...	All other causes..... (residual)	26	8.9	24.6
American Indian and Alaska Native, non-Hispanic, male, 25-34 years				
...	All causes	1,111	100.0	510.4
1	Accidents (unintentional injuries)..... (V01-X59,Y85-Y86)	388	34.9	178.2
2	Intentional self-harm (suicide)..... (†U03,X60-X84,Y87.0)	157	14.1	72.1
3	Chronic liver disease and cirrhosis..... (K70,K73-K74)	122	11.0	56.0
4	Assault (homicide)..... (†U01-†U02.9,X85-Y09,Y87.1)	97	8.7	44.6
5	COVID-19..... (†U07.1)	72	6.5	33.1
6	Diseases of heart..... (I00-I09,I11,I13,I20-I51)	40	3.6	18.4
7	Diabetes mellitus..... (E10-E149)	15	1.4	*
8	Malignant neoplasms..... (C00-C97)	14	1.3	*
9	Legal intervention..... (Y35,Y89.0)	10	0.9	*
10	Septicemia..... (A40-A41)	7	0.6	*
...	All other causes..... (residual)	189	17.0	86.8
American Indian and Alaska Native, non-Hispanic, male, 35-44 years				
...	All causes	1,372	100.0	813.2
1	Accidents (unintentional injuries)..... (V01-X59,Y85-Y86)	367	26.7	217.5
2	Chronic liver disease and cirrhosis..... (K70,K73-K74)	180	13.1	106.7
3	COVID-19..... (†U07.1)	123	9.0	72.9
4	Diseases of heart..... (I00-I09,I11,I13,I20-I51)	120	8.7	71.1
5	Intentional self-harm (suicide)..... (†U03,X60-X84,Y87.0)	85	6.2	50.4
6	Assault (homicide)..... (†U01-†U02.9,X85-Y09,Y87.1)	80	5.8	47.4
7	Diabetes mellitus..... (E10-E149)	50	3.6	29.6
8	Malignant neoplasms..... (C00-C97)	41	3.0	24.3
9	Influenza and pneumonia..... (J09-J18)	21	1.5	12.4
10	Nephritis, nephrotic syndrome and nephrosis..... (N00-N07,N17-N19,N25-N27)	14	1.0	*
...	All other causes..... (residual)	291	21.2	172.5

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
American Indian and Alaska Native, non-Hispanic, male, 45–54 years				
...	All causes	1,854	100.0	1,218.8
1	COVID-19	304	16.4	199.8
2	Accidents (unintentional injuries)	296	16.0	194.6
3	Diseases of heart	248	13.4	163.0
4	Chronic liver disease and cirrhosis	198	10.7	130.2
5	Diabetes mellitus	131	7.1	86.1
6	Malignant neoplasms	122	6.6	80.2
7	Intentional self-harm (suicide)	59	3.2	38.8
8	Cerebrovascular diseases	39	2.1	25.6
9	Influenza and pneumonia	33	1.8	21.7
10	Assault (homicide)	27	1.5	17.7
...	All other causes	397	21.4	261.0
American Indian and Alaska Native, non-Hispanic, male, 55–64 years				
...	All causes	2,955	100.0	1,894.4
1	Diseases of heart	531	18.0	340.4
2	COVID-19	501	17.0	321.2
3	Malignant neoplasms	454	15.4	291.0
4	Accidents (unintentional injuries)	258	8.7	165.4
5	Diabetes mellitus	199	6.7	127.6
6	Chronic liver disease and cirrhosis	187	6.3	119.9
7	Chronic lower respiratory diseases	85	2.9	54.5
8	Cerebrovascular diseases	60	2.0	38.5
9	Influenza and pneumonia	39	1.3	25.0
9	Nephritis, nephrotic syndrome and nephrosis	39	1.3	25.0
...	All other causes	602	20.4	385.9
American Indian and Alaska Native, non-Hispanic, male, 65 years and older				
...	All causes	6,873	100.0	4,393.5
1	COVID-19	1,410	20.5	901.3
2	Diseases of heart	1,389	20.2	887.9
3	Malignant neoplasms	1,145	16.7	731.9
4	Diabetes mellitus	362	5.3	231.4
5	Chronic lower respiratory diseases	318	4.6	203.3
6	Cerebrovascular diseases	266	3.9	170.0
7	Accidents (unintentional injuries)	193	2.8	123.4
8	Influenza and pneumonia	125	1.8	79.9
9	Alzheimer disease	123	1.8	78.6
10	Chronic liver disease and cirrhosis	115	1.7	73.5
...	All other causes	1,427	20.8	912.2
American Indian and Alaska Native, non-Hispanic, male, 65–74 years				
...	All causes	3,144	100.0	3,069.9
1	COVID-19	668	21.2	652.3
2	Malignant neoplasms	605	19.2	590.7
3	Diseases of heart	602	19.1	587.8
4	Diabetes mellitus	193	6.1	188.5
5	Chronic lower respiratory diseases	128	4.1	125.0
6	Accidents (unintentional injuries)	112	3.6	109.4
7	Cerebrovascular diseases	89	2.8	86.9
8	Chronic liver disease and cirrhosis	83	2.6	81.0
9	Influenza and pneumonia	47	1.5	45.9
10	Nephritis, nephrotic syndrome and nephrosis	43	1.4	42.0
...	All other causes	574	18.3	560.5

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
American Indian and Alaska Native, non-Hispanic, male, 75–84 years				
...	All causes	2,439	100.0	5,817.0
1	COVID-19 (†U07.1)	511	21.0	1,218.7
2	Diseases of heart (I00–I09,I11,I13,I20–I51)	508	20.8	1,211.6
3	Malignant neoplasms (C00–C97)	395	16.2	942.1
4	Chronic lower respiratory diseases (J40–J47)	141	5.8	336.3
5	Diabetes mellitus (E10–E149)	120	4.9	286.2
6	Cerebrovascular diseases (I60–I69)	111	4.6	264.7
7	Alzheimer disease (G30)	48	2.0	114.5
8	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	47	1.9	112.1
9	Influenza and pneumonia (J09–J18)	44	1.8	104.9
10	Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)	37	1.5	88.2
...	All other causes (residual)	477	19.6	1,137.6
American Indian and Alaska Native, non-Hispanic, male, 85 years and older				
...	All causes	1,290	100.0	10,666.4
1	Diseases of heart (I00–I09,I11,I13,I20–I51)	279	21.6	2,306.9
2	COVID-19 (†U07.1)	231	17.9	1,910.0
3	Malignant neoplasms (C00–C97)	145	11.2	1,198.9
4	Cerebrovascular diseases (I60–I69)	66	5.1	545.7
5	Alzheimer disease (G30)	59	4.6	487.8
6	Diabetes mellitus (E10–E149)	49	3.8	405.2
6	Chronic lower respiratory diseases (J40–J47)	49	3.8	405.2
8	Influenza and pneumonia (J09–J18)	34	2.6	281.1
8	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	34	2.6	281.1
10	Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)	32	2.5	264.6
...	All other causes (residual)	312	24.2	2,579.8
American Indian and Alaska Native, non-Hispanic, female, all ages ⁴				
...	All causes	11,928	100.0	845.1
1	COVID-19 (†U07.1)	2,012	16.9	142.5
2	Malignant neoplasms (C00–C97)	1,595	13.4	113.0
3	Diseases of heart (I00–I09,I11,I13,I20–I51)	1,534	12.9	108.7
4	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	941	7.9	66.7
5	Chronic liver disease and cirrhosis (K70,K73–K74)	720	6.0	51.0
6	Diabetes mellitus (E10–E149)	597	5.0	42.3
7	Cerebrovascular diseases (I60–I69)	472	4.0	33.4
8	Chronic lower respiratory diseases (J40–J47)	425	3.6	30.1
9	Alzheimer disease (G30)	310	2.6	22.0
10	Influenza and pneumonia (J09–J18)	221	1.9	15.7
...	All other causes (residual)	3,101	26.0	219.7
American Indian and Alaska Native, non-Hispanic, female, 1–4 years				
...	All causes	29	100.0	40.7
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	9	31.0	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	6	20.7	*
3	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	3	10.3	*
4	Malignant neoplasms (C00–C97)	2	6.9	*
4	Cerebrovascular diseases (I60–I69)	2	6.9	*
4	Influenza and pneumonia (J09–J18)	2	6.9	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00–D48)	1	3.4	*
7	Diseases of heart (I00–I09,I11,I13,I20–I51)	1	3.4	*
...	All other causes (residual)	3	10.3	*

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
American Indian and Alaska Native, non-Hispanic, female, 5-9 years				
...	All causes	22	100.0	22.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7	31.8	*
2	Malignant neoplasms (C00-C97)	3	13.6	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	13.6	*
4	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	2	9.1	*
5	Cerebrovascular diseases (I60-I69)	1	4.5	*
...	All other causes (residual)	6	27.3	*
American Indian and Alaska Native, non-Hispanic, female, 10-14 years				
...	All causes	22	100.0	21.6
1	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	8	36.4	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4	18.2	*
3	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	3	13.6	*
4	Malignant neoplasms (C00-C97)	2	9.1	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	9.1	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	4.5	*
...	All other causes (residual)	2	9.1	*
American Indian and Alaska Native, non-Hispanic, female, 15-19 years				
...	All causes	71	100.0	70.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	40.8	28.9
2	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	19	26.8	*
3	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	4	5.6	*
4	Pregnancy, childbirth and the puerperium (O00-O99)	2	2.8	*
4	Malignant neoplasms (C00-C97)	2	2.8	*
4	COVID-19 (†U07.1)	2	2.8	*
7	Septicemia (A40-A41)	1	1.4	*
7	Cerebrovascular diseases (I60-I69)	1	1.4	*
7	Chronic lower respiratory diseases (J40-J47)	1	1.4	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	1.4	*
...	All other causes (residual)	9	12.7	*
American Indian and Alaska Native, non-Hispanic, female, 20-24 years				
...	All causes	123	100.0	118.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	66	53.7	63.8
2	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	22	17.9	21.3
3	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	9	7.3	*
4	Pregnancy, childbirth and the puerperium (O00-O99)	5	4.1	*
5	Malignant neoplasms (C00-C97)	3	2.4	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2	1.6	*
6	Complications of medical and surgical care (Y40-Y84,Y88)	2	1.6	*
8	Cerebrovascular diseases (I60-I69)	1	0.8	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	0.8	*
8	Influenza and pneumonia (J09-J18)	1	0.8	*
8	COVID-19 (†U07.1)	1	0.8	*
...	All other causes (residual)	10	8.1	*

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
American Indian and Alaska Native, non-Hispanic, female, 25-34 years				
...	All causes	644	100.0	302.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	221	34.3	103.9
2	Chronic liver disease and cirrhosis (K70,K73-K74)	92	14.3	43.3
3	COVID-19 (†U07.1)	55	8.5	25.9
4	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	50	7.8	23.5
5	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	20	3.1	9.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.0	*
7	Malignant neoplasms (C00-C97)	14	2.2	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	10	1.6	*
8	Diabetes mellitus (E10-E149)	10	1.6	*
10	Septicemia (A40-A41)	7	1.1	*
10	Influenza and pneumonia (J09-J18)	7	1.1	*
...	All other causes (residual)	139	21.6	65.4
American Indian and Alaska Native, non-Hispanic, female, 35-44 years				
...	All causes	842	100.0	482.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	182	21.6	104.3
2	Chronic liver disease and cirrhosis (K70,K73-K74)	158	18.8	90.5
3	COVID-19 (†U07.1)	98	11.6	56.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	7.0	33.8
5	Malignant neoplasms (C00-C97)	57	6.8	32.7
6	Diabetes mellitus (E10-E149)	28	3.3	16.0
7	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	27	3.2	15.5
8	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	19	2.3	*
9	Septicemia (A40-A41)	12	1.4	*
10	Cerebrovascular diseases (I60-I69)	9	1.1	*
...	All other causes (residual)	193	22.9	110.6
American Indian and Alaska Native, non-Hispanic, female, 45-54 years				
...	All causes	1,209	100.0	745.6
1	Chronic liver disease and cirrhosis (K70,K73-K74)	185	15.3	114.1
2	COVID-19 (†U07.1)	166	13.7	102.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	159	13.2	98.1
4	Malignant neoplasms (C00-C97)	150	12.4	92.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	112	9.3	69.1
6	Diabetes mellitus (E10-E149)	74	6.1	45.6
7	Cerebrovascular diseases (I60-I69)	39	3.2	24.1
8	Chronic lower respiratory diseases (J40-J47)	22	1.8	13.6
9	Influenza and pneumonia (J09-J18)	19	1.6	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	1.3	*
...	All other causes (residual)	267	22.1	164.7
American Indian and Alaska Native, non-Hispanic, female, 55-64 years				
...	All causes	2,099	100.0	1,188.3
1	COVID-19 (†U07.1)	396	18.9	224.2
2	Malignant neoplasms (C00-C97)	368	17.5	208.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	240	11.4	135.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	172	8.2	97.4
5	Diabetes mellitus (E10-E149)	110	5.2	62.3
6	Chronic lower respiratory diseases (J40-J47)	103	4.9	58.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	102	4.9	57.7
8	Cerebrovascular diseases (I60-I69)	70	3.3	39.6
9	Influenza and pneumonia (J09-J18)	40	1.9	22.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	1.9	22.1
...	All other causes (residual)	459	21.9	259.9

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
American Indian and Alaska Native, non-Hispanic, female, 65 years and older				
...	All causes	6,773	100.0	3,485.0
1	COVID-19 (†U07.1)	1,294	19.1	665.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,100	16.2	566.0
3	Malignant neoplasms (C00-C97)	993	14.7	510.9
4	Diabetes mellitus (E10-E149)	375	5.5	193.0
5	Cerebrovascular diseases (I60-I69)	343	5.1	176.5
6	Alzheimer disease (G30)	306	4.5	157.5
7	Chronic lower respiratory diseases (J40-J47)	292	4.3	150.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	148	2.2	76.2
9	Influenza and pneumonia (J09-J18)	146	2.2	75.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	131	1.9	67.4
...	All other causes (residual)	1,645	24.3	846.4
American Indian and Alaska Native, non-Hispanic, female, 65-74 years				
...	All causes	2,407	100.0	1,995.6
1	COVID-19 (†U07.1)	539	22.4	446.9
2	Malignant neoplasms (C00-C97)	446	18.5	369.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	333	13.8	276.1
4	Diabetes mellitus (E10-E149)	166	6.9	137.6
5	Chronic lower respiratory diseases (J40-J47)	115	4.8	95.3
6	Cerebrovascular diseases (I60-I69)	89	3.7	73.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	79	3.3	65.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	61	2.5	50.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	2.1	41.5
10	Septicemia (A40-A41)	40	1.7	33.2
10	Influenza and pneumonia (J09-J18)	40	1.7	33.2
...	All other causes (residual)	449	18.7	372.2
American Indian and Alaska Native, non-Hispanic, female, 75-84 years				
...	All causes	2,403	100.0	4,493.3
1	COVID-19 (†U07.1)	442	18.4	826.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	406	16.9	759.2
3	Malignant neoplasms (C00-C97)	378	15.7	706.8
4	Diabetes mellitus (E10-E149)	138	5.7	258.0
5	Cerebrovascular diseases (I60-I69)	122	5.1	228.1
5	Chronic lower respiratory diseases (J40-J47)	122	5.1	228.1
7	Alzheimer disease (G30)	114	4.7	213.2
8	Influenza and pneumonia (J09-J18)	47	2.0	87.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	1.8	82.3
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41	1.7	76.7
...	All other causes (residual)	549	22.8	1,026.6
American Indian and Alaska Native, non-Hispanic, female, 85 years and older				
...	All causes	1,963	100.0	9,694.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	361	18.4	1,782.9
2	COVID-19 (†U07.1)	313	15.9	1,545.8
3	Malignant neoplasms (C00-C97)	169	8.6	834.7
4	Alzheimer disease (G30)	161	8.2	795.1
5	Cerebrovascular diseases (I60-I69)	132	6.7	651.9
6	Diabetes mellitus (E10-E149)	71	3.6	350.7
7	Influenza and pneumonia (J09-J18)	59	3.0	291.4
8	Chronic lower respiratory diseases (J40-J47)	55	2.8	271.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	50	2.5	246.9
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	46	2.3	227.2
...	All other causes (residual)	546	27.8	2,696.6

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
Asian and Pacific Islander, non-Hispanic, both sexes, all ages ⁴				
...	All causes	99,574	100.0	467.8
1	Malignant neoplasms (C00–C97)	19,869	20.0	93.3
2	Diseases of heart (I00–I09,I11,I13,I20–I51)	18,992	19.1	89.2
3	COVID-19 (†U07.1)	13,597	13.7	63.9
4	Cerebrovascular diseases (I60–I69)	6,655	6.7	31.3
5	Diabetes mellitus (E10–E149)	4,123	4.1	19.4
6	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	3,927	3.9	18.4
7	Alzheimer disease (G30)	3,574	3.6	16.8
8	Chronic lower respiratory diseases (J40–J47)	2,176	2.2	10.2
9	Influenza and pneumonia (J09–J18)	2,169	2.2	10.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,841	1.8	8.6
...	All other causes (residual)	22,651	22.7	106.4
Asian and Pacific Islander, non-Hispanic, both sexes, 1–4 years				
...	All causes	132	100.0	12.8
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	30	22.7	2.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	16	12.1	*
3	Malignant neoplasms (C00–C97)	11	8.3	*
4	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	9	6.8	*
5	Diseases of heart (I00–I09,I11,I13,I20–I51)	4	3.0	*
6	Influenza and pneumonia (J09–J18)	3	2.3	*
7	Cerebrovascular diseases (I60–I69)	2	1.5	*
7	Acute bronchitis and bronchiolitis (J20–J21)	2	1.5	*
7	Chronic lower respiratory diseases (J40–J47)	2	1.5	*
7	Certain conditions originating in the perinatal period (P00–P96)	2	1.5	*
...	All other causes (residual)	51	38.6	4.9
Asian and Pacific Islander, non-Hispanic, both sexes, 5–9 years				
...	All causes	87	100.0	6.6
1	Malignant neoplasms (C00–C97)	18	20.7	*
2	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	11	12.6	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	10	11.5	*
4	Diseases of heart (I00–I09,I11,I13,I20–I51)	7	8.0	*
5	Influenza and pneumonia (J09–J18)	4	4.6	*
6	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	3	3.4	*
7	Diseases of appendix (K35–K38)	2	2.3	*
7	Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)	2	2.3	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00–D48)	2	2.3	*
10	Meningitis (G00,G03)	1	1.1	*
10	Cerebrovascular diseases (I60–I69)	1	1.1	*
10	Complications of medical and surgical care (Y40–Y84,Y88)	1	1.1	*
10	Septicemia (A40–A41)	1	1.1	*
10	Certain conditions originating in the perinatal period (P00–P96)	1	1.1	*
10	Chronic lower respiratory diseases (J40–J47)	1	1.1	*
...	All other causes (residual)	22	25.3	1.7

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
Asian and Pacific Islander, non-Hispanic, both sexes, 10-14 years				
...	All causes	121	100.0	9.8
1	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	21	17.4	1.7
2	Malignant neoplasms (C00-C97)	17	14.0	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	14.0	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	6.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	3.3	*
5	Chronic lower respiratory diseases (J40-J47)	4	3.3	*
5	Influenza and pneumonia (J09-J18)	4	3.3	*
5	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	4	3.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	2.5	*
10	Cerebrovascular diseases (I60-I69)	2	1.7	*
10	Septicemia (A40-A41)	2	1.7	*
10	COVID-19 (†U07.1)	2	1.7	*
...	All other causes (residual)	33	27.3	2.7
Asian and Pacific Islander, non-Hispanic, both sexes, 15-19 years				
...	All causes	344	100.0	27.1
1	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	117	34.0	9.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94	27.3	7.4
3	Malignant neoplasms (C00-C97)	33	9.6	2.6
4	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	17	4.9	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	2.0	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	2.0	*
7	Influenza and pneumonia (J09-J18)	5	1.5	*
8	Cerebrovascular diseases (I60-I69)	4	1.2	*
9	COVID-19 (†U07.1)	3	0.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.9	*
9	Diabetes mellitus (E10-E149)	3	0.9	*
...	All other causes (residual)	51	14.8	4.0
Asian and Pacific Islander, non-Hispanic, both sexes, 20-24 years				
...	All causes	660	100.0	47.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	231	35.0	16.6
2	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	172	26.1	12.4
3	Malignant neoplasms (C00-C97)	44	6.7	3.2
4	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	43	6.5	3.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	4.7	2.2
6	COVID-19 (†U07.1)	19	2.9	*
7	Septicemia (A40-A41)	11	1.7	*
8	Diabetes mellitus (E10-E149)	6	0.9	*
9	Cerebrovascular diseases (I60-I69)	5	0.8	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.8	*
...	All other causes (residual)	93	14.1	6.7

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
Asian and Pacific Islander, non-Hispanic, both sexes, 25-34 years				
...	All causes	1,858	100.0	52.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	525	28.3	14.9
2	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	325	17.5	9.2
3	Malignant neoplasms (C00-C97)	189	10.2	5.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	136	7.3	3.9
5	COVID-19 (†U07.1)	116	6.2	3.3
6	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	91	4.9	2.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	33	1.8	0.9
8	Cerebrovascular diseases (I60-I69)	29	1.6	0.8
9	Diabetes mellitus (E10-E149)	25	1.3	0.7
10	Pregnancy, childbirth and the puerperium (O00-O99)	22	1.2	0.6
...	All other causes (residual)	367	19.8	10.4
Asian and Pacific Islander, non-Hispanic, both sexes, 35-44 years				
...	All causes	3,049	100.0	91.2
1	Malignant neoplasms (C00-C97)	588	19.3	17.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	506	16.6	15.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	400	13.1	12.0
4	COVID-19 (†U07.1)	297	9.7	8.9
5	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	235	7.7	7.0
6	Cerebrovascular diseases (I60-I69)	138	4.5	4.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	122	4.0	3.6
8	Diabetes mellitus (E10-E149)	87	2.9	2.6
9	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	64	2.1	1.9
10	Influenza and pneumonia (J09-J18)	31	1.0	0.9
...	All other causes (residual)	581	19.1	17.4
Asian and Pacific Islander, non-Hispanic, both sexes, 45-54 years				
...	All causes	6,028	100.0	211.1
1	Malignant neoplasms (C00-C97)	1,670	27.7	58.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,032	17.1	36.1
3	COVID-19 (†U07.1)	748	12.4	26.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	462	7.7	16.2
5	Cerebrovascular diseases (I60-I69)	339	5.6	11.9
6	Diabetes mellitus (E10-E149)	267	4.4	9.4
7	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	232	3.8	8.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	161	2.7	5.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	71	1.2	2.5
10	Septicemia (A40-A41)	65	1.1	2.3
...	All other causes (residual)	981	16.3	34.4
Asian and Pacific Islander, non-Hispanic, both sexes, 55-64 years				
...	All causes	11,672	100.0	506.0
1	Malignant neoplasms (C00-C97)	3,715	31.8	161.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,008	17.2	87.1
3	COVID-19 (†U07.1)	1,811	15.5	78.5
4	Cerebrovascular diseases (I60-I69)	565	4.8	24.5
5	Diabetes mellitus (E10-E149)	528	4.5	22.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	418	3.6	18.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	202	1.7	8.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	187	1.6	8.1
9	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	179	1.5	7.8
10	Influenza and pneumonia (J09-J18)	174	1.5	7.5
...	All other causes (residual)	1,885	16.1	81.7

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
Asian and Pacific Islander, non-Hispanic, both sexes, 65 years and older				
...	All causes	74,967	100.0	2,705.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,359	20.5	554.2
2	Malignant neoplasms (C00-C97)	13,582	18.1	490.1
3	COVID-19 (†U07.1)	10,598	14.1	382.4
4	Cerebrovascular diseases (I60-I69)	5,567	7.4	200.9
5	Alzheimer disease (G30)	3,545	4.7	127.9
6	Diabetes mellitus (E10-E149)	3,207	4.3	115.7
7	Chronic lower respiratory diseases (J40-J47)	1,954	2.6	70.5
8	Influenza and pneumonia (J09-J18)	1,856	2.5	67.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,644	2.2	59.3
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,617	2.2	58.4
...	All other causes (residual)	16,038	21.4	578.8
Asian and Pacific Islander, non-Hispanic, both sexes, 65-74 years				
...	All causes	18,889	100.0	1,139.7
1	Malignant neoplasms (C00-C97)	5,317	28.1	320.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,366	17.8	203.1
3	COVID-19 (†U07.1)	3,221	17.1	194.4
4	Cerebrovascular diseases (I60-I69)	1,100	5.8	66.4
5	Diabetes mellitus (E10-E149)	986	5.2	59.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	428	2.3	25.8
7	Chronic lower respiratory diseases (J40-J47)	355	1.9	21.4
8	Influenza and pneumonia (J09-J18)	350	1.9	21.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	343	1.8	20.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	260	1.4	15.7
...	All other causes (residual)	3,163	16.7	190.9
Asian and Pacific Islander, non-Hispanic, both sexes, 75-84 years				
...	All causes	23,982	100.0	3,037.9
1	Malignant neoplasms (C00-C97)	4,979	20.8	630.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,611	19.2	584.1
3	COVID-19 (†U07.1)	3,483	14.5	441.2
4	Cerebrovascular diseases (I60-I69)	1,785	7.4	226.1
5	Diabetes mellitus (E10-E149)	1,150	4.8	145.7
6	Alzheimer disease (G30)	721	3.0	91.3
7	Chronic lower respiratory diseases (J40-J47)	659	2.7	83.5
8	Influenza and pneumonia (J09-J18)	573	2.4	72.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	536	2.2	67.9
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	524	2.2	66.4
...	All other causes (residual)	4,961	20.7	628.4
Asian and Pacific Islander, non-Hispanic, both sexes, 85 years and older				
...	All causes	32,096	100.0	9,893.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,382	23.0	2,275.5
2	COVID-19 (†U07.1)	3,894	12.1	1,200.3
3	Malignant neoplasms (C00-C97)	3,286	10.2	1,012.9
4	Cerebrovascular diseases (I60-I69)	2,682	8.4	826.7
5	Alzheimer disease (G30)	2,658	8.3	819.3
6	Diabetes mellitus (E10-E149)	1,071	3.3	330.1
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	952	3.0	293.4
8	Chronic lower respiratory diseases (J40-J47)	940	2.9	289.7
9	Influenza and pneumonia (J09-J18)	933	2.9	287.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	665	2.1	205.0
...	All other causes (residual)	7,633	23.8	2,352.8

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
Asian and Pacific Islander, non-Hispanic, male, all ages ⁴				
...	All causes	52,442	100.0	513.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,573	20.2	103.6
2	Malignant neoplasms (C00-C97)	9,958	19.0	97.6
3	COVID-19 (†U07.1)	8,060	15.4	79.0
4	Cerebrovascular diseases (I60-I69)	2,981	5.7	29.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,611	5.0	25.6
6	Diabetes mellitus (E10-E149)	2,257	4.3	22.1
7	Chronic lower respiratory diseases (J40-J47)	1,259	2.4	12.3
8	Influenza and pneumonia (J09-J18)	1,213	2.3	11.9
9	Alzheimer disease (G30)	1,079	2.1	10.6
10	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	1,063	2.0	10.4
...	All other causes (residual)	11,388	21.7	111.6
Asian and Pacific Islander, non-Hispanic, male, 1-4 years				
...	All causes	71	100.0	13.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	25.4	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	11.3	*
3	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	6	8.5	*
4	Malignant neoplasms (C00-C97)	4	5.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.8	*
6	Tuberculosis (A16-A19)	1	1.4	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.4	*
6	Septicemia (A40-A41)	1	1.4	*
6	Chronic lower respiratory diseases (J40-J47)	1	1.4	*
6	Anemias (D50-D64)	1	1.4	*
6	Certain conditions originating in the perinatal period (P00-P96)	1	1.4	*
...	All other causes (residual)	27	38.0	5.1
Asian and Pacific Islander, non-Hispanic, male, 5-9 years				
...	All causes	44	100.0	6.5
1	Malignant neoplasms (C00-C97)	11	25.0	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	11.4	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	9.1	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	6.8	*
4	Influenza and pneumonia (J09-J18)	3	6.8	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	4.5	*
7	Septicemia (A40-A41)	1	2.3	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.3	*
7	Meningitis (G00,G03)	1	2.3	*
7	Cerebrovascular diseases (I60-I69)	1	2.3	*
7	Certain conditions originating in the perinatal period (P00-P96)	1	2.3	*
7	Complications of medical and surgical care (Y40-Y84,Y88)	1	2.3	*
...	All other causes (residual)	10	22.7	*

Table I–1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*]

Rank ¹	Cause of death (based on ICD–10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
Asian and Pacific Islander, non-Hispanic, male, 10–14 years				
...	All causes	73	100.0	11.6
1	Intentional self-harm (suicide)..... (†U03,X60–X84,Y87.0)	14	19.2	*
2	Accidents (unintentional injuries)..... (V01–X59,Y85–Y86)	9	12.3	*
3	Malignant neoplasms..... (C00–C97)	5	6.8	*
3	Congenital malformations, deformations and chromosomal abnormalities..... (Q00–Q99)	5	6.8	*
5	Assault (homicide)..... (†U01–†U02.9,X85–Y09,Y87.1)	3	4.1	*
5	Influenza and pneumonia..... (J09–J18)	3	4.1	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior..... (D00–D48)	3	4.1	*
5	Diseases of heart..... (I00–I09,I11,I13,I20–I51)	3	4.1	*
5	Chronic lower respiratory diseases..... (J40–J47)	3	4.1	*
10	COVID-19..... (†U07.1)	2	2.7	*
...	All other causes..... (residual)	23	31.5	3.7
Asian and Pacific Islander, non-Hispanic, male, 15–19 years				
...	All causes.....	211	100.0	33.1
1	Intentional self-harm (suicide)..... (†U03,X60–X84,Y87.0)	79	37.4	12.4
2	Accidents (unintentional injuries)..... (V01–X59,Y85–Y86)	65	30.8	10.2
3	Malignant neoplasms..... (C00–C97)	14	6.6	*
4	Assault (homicide)..... (†U01–†U02.9,X85–Y09,Y87.1)	13	6.2	*
5	Diseases of heart..... (I00–I09,I11,I13,I20–I51)	4	1.9	*
5	Congenital malformations, deformations and chromosomal abnormalities..... (Q00–Q99)	4	1.9	*
7	Diabetes mellitus..... (E10–E149)	3	1.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior..... (D00–D48)	2	0.9	*
8	Influenza and pneumonia..... (J09–J18)	2	0.9	*
8	Cerebrovascular diseases..... (I60–I69)	2	0.9	*
8	Septicemia..... (A40–A41)	2	0.9	*
...	All other causes..... (residual)	21	10.0	3.3
Asian and Pacific Islander, non-Hispanic, male, 20–24 years				
...	All causes.....	469	100.0	66.9
1	Accidents (unintentional injuries)..... (V01–X59,Y85–Y86)	184	39.2	26.3
2	Intentional self-harm (suicide)..... (†U03,X60–X84,Y87.0)	116	24.7	16.6
3	Assault (homicide)..... (†U01–†U02.9,X85–Y09,Y87.1)	40	8.5	5.7
4	Malignant neoplasms..... (C00–C97)	27	5.8	3.9
5	Diseases of heart..... (I00–I09,I11,I13,I20–I51)	20	4.3	2.9
6	COVID-19..... (†U07.1)	15	3.2	*
7	Septicemia..... (A40–A41)	6	1.3	*
8	Diabetes mellitus..... (E10–E149)	4	0.9	*
9	Congenital malformations, deformations and chromosomal abnormalities..... (Q00–Q99)	2	0.4	*
9	Cerebrovascular diseases..... (I60–I69)	2	0.4	*
9	Influenza and pneumonia..... (J09–J18)	2	0.4	*
9	Legal intervention..... (Y35,Y89.0)	2	0.4	*
...	All other causes..... (residual)	49	10.4	7.0

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
Asian and Pacific Islander, non-Hispanic, male, 25-34 years				
...	All causes	1,297	100.0	74.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	412	31.8	23.7
2	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	253	19.5	14.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	108	8.3	6.2
4	Malignant neoplasms (C00-C97)	96	7.4	5.5
5	COVID-19 (†U07.1)	73	5.6	4.2
6	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	71	5.5	4.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	25	1.9	1.4
8	Cerebrovascular diseases (I60-I69)	17	1.3	*
9	Influenza and pneumonia (J09-J18)	15	1.2	*
10	Diabetes mellitus (E10-E149)	13	1.0	*
...	All other causes (residual)	214	16.5	12.3
Asian and Pacific Islander, non-Hispanic, male, 35-44 years				
...	All causes	2,015	100.0	127.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	396	19.7	25.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	314	15.6	19.9
3	Malignant neoplasms (C00-C97)	254	12.6	16.1
4	COVID-19 (†U07.1)	203	10.1	12.9
5	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	173	8.6	11.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	99	4.9	6.3
7	Cerebrovascular diseases (I60-I69)	89	4.4	5.6
8	Diabetes mellitus (E10-E149)	65	3.2	4.1
9	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	45	2.2	2.9
10	Influenza and pneumonia (J09-J18)	20	1.0	1.3
...	All other causes (residual)	357	17.7	22.6
Asian and Pacific Islander, non-Hispanic, male, 45-54 years				
...	All causes	3,758	100.0	281.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	814	21.7	61.0
2	Malignant neoplasms (C00-C97)	701	18.7	52.5
3	COVID-19 (†U07.1)	526	14.0	39.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	359	9.6	26.9
5	Cerebrovascular diseases (I60-I69)	194	5.2	14.5
6	Diabetes mellitus (E10-E149)	183	4.9	13.7
7	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	163	4.3	12.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	121	3.2	9.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	1.2	3.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	42	1.1	3.1
...	All other causes (residual)	609	16.2	45.6
Asian and Pacific Islander, non-Hispanic, male, 55-64 years				
...	All causes	7,323	100.0	692.8
1	Malignant neoplasms (C00-C97)	1,883	25.7	178.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,497	20.4	141.6
3	COVID-19 (†U07.1)	1,316	18.0	124.5
4	Diabetes mellitus (E10-E149)	354	4.8	33.5
5	Cerebrovascular diseases (I60-I69)	341	4.7	32.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	300	4.1	28.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	150	2.0	14.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	118	1.6	11.2
8	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	118	1.6	11.2
10	Influenza and pneumonia (J09-J18)	100	1.4	9.5
...	All other causes (residual)	1,146	15.6	108.4

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
Asian and Pacific Islander, non-Hispanic, male, 65 years and older				
...	All causes	36,810	100.0	3,070.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,804	21.2	651.1
2	Malignant neoplasms (C00-C97)	6,961	18.9	580.7
3	COVID-19 (†U07.1)	5,922	16.1	494.0
4	Cerebrovascular diseases (I60-I69)	2,333	6.3	194.6
5	Diabetes mellitus (E10-E149)	1,635	4.4	136.4
6	Chronic lower respiratory diseases (J40-J47)	1,135	3.1	94.7
7	Alzheimer disease (G30)	1,064	2.9	88.8
8	Influenza and pneumonia (J09-J18)	1,026	2.8	85.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	853	2.3	71.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	730	2.0	60.9
...	All other causes (residual)	7,347	20.0	612.9
Asian and Pacific Islander, non-Hispanic, male, 65-74 years				
...	All causes	11,042	100.0	1,512.2
1	Malignant neoplasms (C00-C97)	2,741	24.8	375.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,229	20.2	305.3
3	COVID-19 (†U07.1)	2,157	19.5	295.4
4	Diabetes mellitus (E10-E149)	593	5.4	81.2
5	Cerebrovascular diseases (I60-I69)	562	5.1	77.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	257	2.3	35.2
7	Chronic lower respiratory diseases (J40-J47)	244	2.2	33.4
8	Influenza and pneumonia (J09-J18)	228	2.1	31.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	189	1.7	25.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	142	1.3	19.4
...	All other causes (residual)	1,700	15.4	232.8
Asian and Pacific Islander, non-Hispanic, male, 75-84 years				
...	All causes	12,894	100.0	3,730.5
1	Malignant neoplasms (C00-C97)	2,653	20.6	767.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,533	19.6	732.8
3	COVID-19 (†U07.1)	2,080	16.1	601.8
4	Cerebrovascular diseases (I60-I69)	829	6.4	239.8
5	Diabetes mellitus (E10-E149)	639	5.0	184.9
6	Chronic lower respiratory diseases (J40-J47)	402	3.1	116.3
7	Influenza and pneumonia (J09-J18)	337	2.6	97.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	306	2.4	88.5
9	Alzheimer disease (G30)	296	2.3	85.6
10	Parkinson disease (G20-G21)	287	2.2	83.0
...	All other causes (residual)	2,532	19.6	732.6
Asian and Pacific Islander, non-Hispanic, male, 85 years and older				
...	All causes	12,874	100.0	10,482.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,042	23.6	2,476.9
2	COVID-19 (†U07.1)	1,685	13.1	1,372.0
3	Malignant neoplasms (C00-C97)	1,567	12.2	1,275.9
4	Cerebrovascular diseases (I60-I69)	942	7.3	767.0
5	Alzheimer disease (G30)	707	5.5	575.7
6	Chronic lower respiratory diseases (J40-J47)	489	3.8	398.2
7	Influenza and pneumonia (J09-J18)	461	3.6	375.4
8	Diabetes mellitus (E10-E149)	403	3.1	328.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	327	2.5	266.3
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	290	2.3	236.1
...	All other causes (residual)	2,961	23.0	2,410.9

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
Asian and Pacific Islander, non-Hispanic, female, all ages ⁴				
...	All causes	47,132	100.0	425.5
1	Malignant neoplasms (C00–C97)	9,911	21.0	89.5
2	Diseases of heart (I00–I09,I11,I13,I20–I51)	8,419	17.9	76.0
3	COVID-19 (†U07.1)	5,537	11.7	50.0
4	Cerebrovascular diseases (I60–I69)	3,674	7.8	33.2
5	Alzheimer disease (G30)	2,495	5.3	22.5
6	Diabetes mellitus (E10–E149)	1,866	4.0	16.8
7	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	1,316	2.8	11.9
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,018	2.2	9.2
9	Influenza and pneumonia (J09–J18)	956	2.0	8.6
10	Chronic lower respiratory diseases (J40–J47)	917	1.9	8.3
...	All other causes (residual)	11,023	23.4	99.5
Asian and Pacific Islander, non-Hispanic, female, 1–4 years				
...	All causes	61	100.0	12.2
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	12	19.7	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	8	13.1	*
3	Malignant neoplasms (C00–C97)	7	11.5	*
4	Influenza and pneumonia (J09–J18)	3	4.9	*
4	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	3	4.9	*
6	Diseases of heart (I00–I09,I11,I13,I20–I51)	2	3.3	*
6	Cerebrovascular diseases (I60–I69)	2	3.3	*
6	Acute bronchitis and bronchiolitis (J20–J21)	2	3.3	*
9	Chronic lower respiratory diseases (J40–J47)	1	1.6	*
9	Certain conditions originating in the perinatal period (P00–P96)	1	1.6	*
9	Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)	1	1.6	*
...	All other causes (residual)	19	31.1	*
Asian and Pacific Islander, non-Hispanic, female, 5–9 years				
...	All causes	43	100.0	6.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	7	16.3	*
1	Malignant neoplasms (C00–C97)	7	16.3	*
3	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	6	14.0	*
4	Diseases of heart (I00–I09,I11,I13,I20–I51)	3	7.0	*
4	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	3	7.0	*
6	Diseases of appendix (K35–K38)	2	4.7	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00–D48)	1	2.3	*
7	Influenza and pneumonia (J09–J18)	1	2.3	*
7	Chronic lower respiratory diseases (J40–J47)	1	2.3	*
...	All other causes (residual)	12	27.9	*
Asian and Pacific Islander, non-Hispanic, female, 10–14 years				
...	All causes	48	100.0	7.9
1	Malignant neoplasms (C00–C97)	12	25.0	*
2	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	8	16.7	*
3	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	7	14.6	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	3	6.3	*
5	Septicemia (A40–A41)	2	4.2	*
6	Diseases of heart (I00–I09,I11,I13,I20–I51)	1	2.1	*
6	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	1	2.1	*
6	Influenza and pneumonia (J09–J18)	1	2.1	*
6	Cerebrovascular diseases (I60–I69)	1	2.1	*
6	Chronic lower respiratory diseases (J40–J47)	1	2.1	*
...	All other causes (residual)	11	22.9	*

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
Asian and Pacific Islander, non-Hispanic, female, 15-19 years				
...	All causes	133	100.0	21.0
1	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	38	28.6	6.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	21.8	4.6
3	Malignant neoplasms (C00-C97)	19	14.3	*
4	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	4	3.0	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	2.3	*
5	Influenza and pneumonia (J09-J18)	3	2.3	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	2.3	*
5	COVID-19 (†U07.1)	3	2.3	*
9	Cerebrovascular diseases (I60-I69)	2	1.5	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.8	*
10	Tuberculosis (A16-A19)	1	0.8	*
10	Hernia (K40-K46)	1	0.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.8	*
10	Chronic lower respiratory diseases (J40-J47)	1	0.8	*
10	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.8	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.8	*
...	All other causes (residual)	22	16.5	3.5
Asian and Pacific Islander, non-Hispanic, female, 20-24 years				
...	All causes	191	100.0	27.8
1	Intentional self-harm (suicide) (†U03,X60-TX84,Y87.0)	56	29.3	8.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	24.6	6.8
3	Malignant neoplasms (C00-C97)	17	8.9	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	5.8	*
5	Septicemia (A40-A41)	5	2.6	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	4	2.1	*
6	COVID-19 (†U07.1)	4	2.1	*
8	Cerebrovascular diseases (I60-I69)	3	1.6	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.6	*
8	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	3	1.6	*
...	All other causes (residual)	38	19.9	5.5
Asian and Pacific Islander, non-Hispanic, female, 25-34 years				
...	All causes	561	100.0	31.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	113	20.1	6.3
2	Malignant neoplasms (C00-C97)	93	16.6	5.2
3	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	72	12.8	4.0
4	COVID-19 (†U07.1)	43	7.7	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	5.0	1.6
6	Pregnancy, childbirth and the puerperium (O00-O99)	22	3.9	1.2
7	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	20	3.6	1.1
8	Diabetes mellitus (E10-E149)	12	2.1	*
8	Cerebrovascular diseases (I60-I69)	12	2.1	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	8	1.4	*
...	All other causes (residual)	138	24.6	7.7

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
Asian and Pacific Islander, non-Hispanic, female, 35-44 years				
...	All causes	1,034	100.0	58.6
1	Malignant neoplasms..... (C00-C97)	334	32.3	18.9
2	Accidents (unintentional injuries)..... (V01-X59,Y85-Y86)	110	10.6	6.2
3	COVID-19..... (†U07.1)	94	9.1	5.3
4	Diseases of heart..... (I00-I09,I11,I13,I20-I51)	86	8.3	4.9
5	Intentional self-harm (suicide)..... (†U03,X60-X84,Y87.0)	62	6.0	3.5
6	Cerebrovascular diseases..... (I60-I69)	49	4.7	2.8
7	Pregnancy, childbirth and the puerperium..... (O00-O99)	30	2.9	1.7
8	Chronic liver disease and cirrhosis..... (K70,K73-K74)	23	2.2	1.3
9	Diabetes mellitus..... (E10-E149)	22	2.1	1.2
10	Assault (homicide)..... (†U01-†U02.9,X85-Y09,Y87.1)	19	1.8	*
...	All other causes..... (residual)	205	19.8	11.6
Asian and Pacific Islander, non-Hispanic, female, 45-54 years				
...	All causes.....	2,270	100.0	149.3
1	Malignant neoplasms..... (C00-C97)	969	42.7	63.7
2	COVID-19..... (†U07.1)	222	9.8	14.6
3	Diseases of heart..... (I00-I09,I11,I13,I20-I51)	218	9.6	14.3
4	Cerebrovascular diseases..... (I60-I69)	145	6.4	9.5
5	Accidents (unintentional injuries)..... (V01-X59,Y85-Y86)	103	4.5	6.8
6	Diabetes mellitus..... (E10-E149)	84	3.7	5.5
7	Intentional self-harm (suicide)..... (†U03,X60-X84,Y87.0)	69	3.0	4.5
8	Chronic liver disease and cirrhosis..... (K70,K73-K74)	40	1.8	2.6
9	Septicemia..... (A40-A41)	37	1.6	2.4
10	Chronic lower respiratory diseases..... (J40-J47)	27	1.2	1.8
...	All other causes..... (residual)	356	15.7	23.4
Asian and Pacific Islander, non-Hispanic, female, 55-64 years				
...	All causes.....	4,349	100.0	348.1
1	Malignant neoplasms..... (C00-C97)	1,832	42.1	146.6
2	Diseases of heart..... (I00-I09,I11,I13,I20-I51)	511	11.7	40.9
3	COVID-19..... (†U07.1)	495	11.4	39.6
4	Cerebrovascular diseases..... (I60-I69)	224	5.2	17.9
5	Diabetes mellitus..... (E10-E149)	174	4.0	13.9
6	Accidents (unintentional injuries)..... (V01-X59,Y85-Y86)	118	2.7	9.4
7	Influenza and pneumonia..... (J09-J18)	74	1.7	5.9
8	Nephritis, nephrotic syndrome and nephrosis..... (N00-N07,N17-N19,N25-N27)	69	1.6	5.5
9	Intentional self-harm (suicide)..... (†U03,X60-X84,Y87.0)	61	1.4	4.9
10	Chronic lower respiratory diseases..... (J40-J47)	53	1.2	4.2
...	All other causes..... (residual)	738	17.0	59.1
Asian and Pacific Islander, non-Hispanic, female, 65 years and older				
...	All causes.....	38,157	100.0	2,426.6
1	Diseases of heart..... (I00-I09,I11,I13,I20-I51)	7,555	19.8	480.5
2	Malignant neoplasms..... (C00-C97)	6,621	17.4	421.1
3	COVID-19..... (†U07.1)	4,676	12.3	297.4
4	Cerebrovascular diseases..... (I60-I69)	3,234	8.5	205.7
5	Alzheimer disease..... (G30)	2,481	6.5	157.8
6	Diabetes mellitus..... (E10-E149)	1,572	4.1	100.0
7	Essential hypertension and hypertensive renal disease..... (I10,I12,I15)	960	2.5	61.1
8	Influenza and pneumonia..... (J09-J18)	830	2.2	52.8
9	Chronic lower respiratory diseases..... (J40-J47)	819	2.1	52.1
10	Accidents (unintentional injuries)..... (V01-X59,Y85-Y86)	764	2.0	48.6
...	All other causes..... (residual)	8,645	22.7	549.8

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
Asian and Pacific Islander, non-Hispanic, female, 65–74 years				
...	All causes	7,847	100.0	846.4
1	Malignant neoplasms	2,576	32.8	277.9
2	Diseases of heart	1,137	14.5	122.6
3	COVID-19	1,064	13.6	114.8
4	Cerebrovascular diseases	538	6.9	58.0
5	Diabetes mellitus	393	5.0	42.4
6	Accidents (unintentional injuries)	171	2.2	18.4
7	Nephritis, nephrotic syndrome and nephrosis	154	2.0	16.6
8	Influenza and pneumonia	122	1.6	13.2
9	Essential hypertension and hypertensive renal disease	118	1.5	12.7
10	Chronic lower respiratory diseases	111	1.4	12.0
...	All other causes	1,463	18.6	157.8
Asian and Pacific Islander, non-Hispanic, female, 75–84 years				
...	All causes	11,088	100.0	2,498.5
1	Malignant neoplasms	2,326	21.0	524.1
2	Diseases of heart	2,078	18.7	468.2
3	COVID-19	1,403	12.7	316.1
4	Cerebrovascular diseases	956	8.6	215.4
5	Diabetes mellitus	511	4.6	115.1
6	Alzheimer disease	425	3.8	95.8
7	Chronic lower respiratory diseases	257	2.3	57.9
8	Nephritis, nephrotic syndrome and nephrosis	256	2.3	57.7
9	Influenza and pneumonia	236	2.1	53.2
10	Accidents (unintentional injuries)	218	2.0	49.1
...	All other causes	2,422	21.8	545.8
Asian and Pacific Islander, non-Hispanic, female, 85 years and older				
...	All causes	19,222	100.0	9,534.6
1	Diseases of heart	4,340	22.6	2,152.7
2	COVID-19	2,209	11.5	1,095.7
3	Alzheimer disease	1,951	10.1	967.7
4	Cerebrovascular diseases	1,740	9.1	863.1
5	Malignant neoplasms	1,719	8.9	852.7
6	Diabetes mellitus	668	3.5	331.3
7	Essential hypertension and hypertensive renal disease	625	3.3	310.0
8	Influenza and pneumonia	472	2.5	234.1
9	Chronic lower respiratory diseases	451	2.3	223.7
10	Accidents (unintentional injuries)	375	2.0	186.0
...	All other causes	4,672	24.3	2,317.4
Black, non-Hispanic, both sexes, all ages ⁴				
...	All causes	451,792	100.0	1,035.7
1	Diseases of heart	93,320	20.7	213.9
2	Malignant neoplasms	70,963	15.7	162.7
3	COVID-19	56,528	12.5	129.6
4	Accidents (unintentional injuries)	29,133	6.4	66.8
5	Cerebrovascular diseases	22,404	5.0	51.4
6	Diabetes mellitus	19,495	4.3	44.7
7	Assault (homicide)	13,594	3.0	31.2
8	Chronic lower respiratory diseases	12,361	2.7	28.3
9	Alzheimer disease	10,893	2.4	25.0
10	Nephritis, nephrotic syndrome and nephrosis	10,581	2.3	24.3
...	All other causes	112,520	24.9	258.0

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
Black, non-Hispanic, both sexes, 1-4 years				
...	All causes	966	100.0	39.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	273	28.3	11.3
2	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	147	15.2	6.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	95	9.8	3.9
4	Malignant neoplasms (C00-C97)	46	4.8	1.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	4.8	1.9
6	Certain conditions originating in the perinatal period (P00-P96)	22	2.3	0.9
7	Cerebrovascular diseases (I60-I69)	18	1.9	*
8	Influenza and pneumonia (J09-J18)	17	1.8	*
9	Septicemia (A40-A41)	13	1.3	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.9	*
...	All other causes (residual)	280	29.0	11.5
Black, non-Hispanic, both sexes, 5-9 years				
...	All causes	594	100.0	19.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	180	30.3	5.8
2	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	81	13.6	2.6
3	Malignant neoplasms (C00-C97)	56	9.4	1.8
4	Chronic lower respiratory diseases (J40-J47)	35	5.9	1.1
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	5.1	1.0
6	Influenza and pneumonia (J09-J18)	19	3.2	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	2.9	*
8	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	9	1.5	*
9	COVID-19 (†U07.1)	8	1.3	*
10	Cerebrovascular diseases (I60-I69)	7	1.2	*
...	All other causes (residual)	152	25.6	4.9
Black, non-Hispanic, both sexes, 10-14 years				
...	All causes	813	100.0	25.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	170	20.9	5.4
2	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	138	17.0	4.4
3	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	89	10.9	2.8
4	Malignant neoplasms (C00-C97)	72	8.9	2.3
5	Chronic lower respiratory diseases (J40-J47)	58	7.1	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	3.9	1.0
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	3.6	0.9
8	Diabetes mellitus (E10-E149)	19	2.3	*
9	Cerebrovascular diseases (I60-I69)	10	1.2	*
10	Influenza and pneumonia (J09-J18)	9	1.1	*
10	COVID-19 (†U07.1)	9	1.1	*
...	All other causes (residual)	178	21.9	5.6
Black, non-Hispanic, both sexes, 15-19 years				
...	All causes	3,389	100.0	109.1
1	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	1,619	47.8	52.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	797	23.5	25.7
3	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	267	7.9	8.6
4	Malignant neoplasms (C00-C97)	94	2.8	3.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	87	2.6	2.8
6	Chronic lower respiratory diseases (J40-J47)	41	1.2	1.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	1.1	1.2
8	COVID-19 (†U07.1)	30	0.9	1.0
9	Diabetes mellitus (E10-E149)	27	0.8	0.9
10	Legal intervention (Y35,Y89.0)	25	0.7	0.8
...	All other causes (residual)	366	10.8	11.8

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
Black, non-Hispanic, both sexes, 20–24 years				
...	All causes	6,461	100.0	198.0
1	Assault (homicide) (†U01-†U02.9,X85–Y09,Y87.1)	2,598	40.2	79.6
2	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	1,905	29.5	58.4
3	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	572	8.9	17.5
4	Diseases of heart (I00–I09,I11,I13,I20–I51)	162	2.5	5.0
5	Malignant neoplasms (C00–C97)	114	1.8	3.5
6	COVID-19 (†U07.1)	104	1.6	3.2
7	Diabetes mellitus (E10–E149)	75	1.2	2.3
8	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	56	0.9	1.7
9	Pregnancy, childbirth and the puerperium (O00–O99)	50	0.8	1.5
10	Chronic lower respiratory diseases (J40–J47)	45	0.7	1.4
...	All other causes (residual)	780	12.1	23.9
Black, non-Hispanic, both sexes, 25–34 years				
...	All causes	17,458	100.0	252.7
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	5,250	30.1	76.0
2	Assault (homicide) (†U01-†U02.9,X85–Y09,Y87.1)	4,536	26.0	65.7
3	Diseases of heart (I00–I09,I11,I13,I20–I51)	1,238	7.1	17.9
4	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	1,011	5.8	14.6
5	Malignant neoplasms (C00–C97)	650	3.7	9.4
6	COVID-19 (†U07.1)	583	3.3	8.4
7	Diabetes mellitus (E10–E149)	398	2.3	5.8
8	Human immunodeficiency virus (HIV) disease (B20–B24)	289	1.7	4.2
9	Pregnancy, childbirth and the puerperium (O00–O99)	192	1.1	2.8
10	Cerebrovascular diseases (I60–I69)	181	1.0	2.6
...	All other causes (residual)	3,130	17.9	45.3
Black, non-Hispanic, both sexes, 35–44 years				
...	All causes	22,216	100.0	399.1
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	4,874	21.9	87.6
2	Diseases of heart (I00–I09,I11,I13,I20–I51)	3,560	16.0	64.0
3	Assault (homicide) (†U01-†U02.9,X85–Y09,Y87.1)	2,292	10.3	41.2
4	Malignant neoplasms (C00–C97)	1,901	8.6	34.1
5	COVID-19 (†U07.1)	1,491	6.7	26.8
6	Diabetes mellitus (E10–E149)	903	4.1	16.2
7	Cerebrovascular diseases (I60–I69)	585	2.6	10.5
8	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	575	2.6	10.3
9	Human immunodeficiency virus (HIV) disease (B20–B24)	439	2.0	7.9
10	Chronic liver disease and cirrhosis (K70,K73–K74)	361	1.6	6.5
...	All other causes (residual)	5,235	23.6	94.0
Black, non-Hispanic, both sexes, 45–54 years				
...	All causes	38,609	100.0	749.0
1	Diseases of heart (I00–I09,I11,I13,I20–I51)	8,491	22.0	164.7
2	Malignant neoplasms (C00–C97)	5,519	14.3	107.1
3	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	4,835	12.5	93.8
4	COVID-19 (†U07.1)	4,034	10.4	78.3
5	Diabetes mellitus (E10–E149)	1,963	5.1	38.1
6	Cerebrovascular diseases (I60–I69)	1,491	3.9	28.9
7	Assault (homicide) (†U01-†U02.9,X85–Y09,Y87.1)	1,111	2.9	21.6
8	Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)	823	2.1	16.0
9	Chronic liver disease and cirrhosis (K70,K73–K74)	775	2.0	15.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	650	1.7	12.6
...	All other causes (residual)	8,917	23.1	173.0

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
Black, non-Hispanic, both sexes, 55–64 years				
...	All causes	82,593	100.0	1,627.5
1	Diseases of heart (I00–I09,I11,I13,I20–I51)	18,785	22.7	370.2
2	Malignant neoplasms (C00–C97)	16,982	20.6	334.6
3	COVID-19 (†U07.1)	9,899	12.0	195.1
4	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	5,480	6.6	108.0
5	Diabetes mellitus (E10–E149)	4,258	5.2	83.9
6	Cerebrovascular diseases (I60–I69)	3,530	4.3	69.6
7	Chronic lower respiratory diseases (J40–J47)	2,339	2.8	46.1
8	Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)	1,927	2.3	38.0
9	Chronic liver disease and cirrhosis (K70,K73–K74)	1,516	1.8	29.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,465	1.8	28.9
...	All other causes (residual)	16,412	19.9	323.4
Black, non-Hispanic, both sexes, 65 years and older				
...	All causes	272,667	100.0	5,177.2
1	Diseases of heart (I00–I09,I11,I13,I20–I51)	60,831	22.3	1,155.0
2	Malignant neoplasms (C00–C97)	45,519	16.7	864.3
3	COVID-19 (†U07.1)	40,356	14.8	766.2
4	Cerebrovascular diseases (I60–I69)	16,516	6.1	313.6
5	Diabetes mellitus (E10–E149)	11,848	4.3	225.0
6	Alzheimer disease (G30)	10,722	3.9	203.6
7	Chronic lower respiratory diseases (J40–J47)	8,854	3.2	168.1
8	Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)	7,343	2.7	139.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,606	2.1	106.4
10	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	4,930	1.8	93.6
...	All other causes (residual)	60,142	22.1	1,141.9
Black, non-Hispanic, both sexes, 65–74 years				
...	All causes	103,700	100.0	3,144.0
1	Diseases of heart (I00–I09,I11,I13,I20–I51)	22,953	22.1	695.9
2	Malignant neoplasms (C00–C97)	22,367	21.6	678.1
3	COVID-19 (†U07.1)	15,370	14.8	466.0
4	Cerebrovascular diseases (I60–I69)	5,203	5.0	157.7
5	Diabetes mellitus (E10–E149)	5,182	5.0	157.1
6	Chronic lower respiratory diseases (J40–J47)	3,581	3.5	108.6
7	Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)	2,794	2.7	84.7
8	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	2,726	2.6	82.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,952	1.9	59.2
10	Septicemia (A40–A41)	1,823	1.8	55.3
...	All other causes (residual)	19,749	19.0	598.7
Black, non-Hispanic, both sexes, 75–84 years				
...	All causes	89,555	100.0	6,238.4
1	Diseases of heart (I00–I09,I11,I13,I20–I51)	19,307	21.6	1,344.9
2	Malignant neoplasms (C00–C97)	15,240	17.0	1,061.6
3	COVID-19 (†U07.1)	14,244	15.9	992.2
4	Cerebrovascular diseases (I60–I69)	5,504	6.1	383.4
5	Diabetes mellitus (E10–E149)	4,092	4.6	285.1
6	Alzheimer disease (G30)	3,317	3.7	231.1
7	Chronic lower respiratory diseases (J40–J47)	3,177	3.5	221.3
8	Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)	2,599	2.9	181.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,758	2.0	122.5
10	Septicemia (A40–A41)	1,520	1.7	105.9
...	All other causes (residual)	18,797	21.0	1,309.4

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
Black, non-Hispanic, both sexes, 85 years and older				
...	All causes	79,412	100.0	14,905.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,571	23.4	3,485.8
2	COVID-19 (†U07.1)	10,742	13.5	2,016.3
3	Malignant neoplasms (C00-C97)	7,912	10.0	1,485.1
4	Alzheimer disease (G30)	6,350	8.0	1,191.9
5	Cerebrovascular diseases (I60-I69)	5,809	7.3	1,090.3
6	Diabetes mellitus (E10-E149)	2,574	3.2	483.1
7	Chronic lower respiratory diseases (J40-J47)	2,096	2.6	393.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,950	2.5	366.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,896	2.4	355.9
10	Influenza and pneumonia (J09-J18)	1,364	1.7	256.0
...	All other causes (residual)	20,148	25.4	3,781.8
Black, non-Hispanic, male, all ages ⁴				
...	All causes	239,016	100.0	1,143.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	50,199	21.0	240.1
2	Malignant neoplasms (C00-C97)	35,693	14.9	170.7
3	COVID-19 (†U07.1)	29,946	12.5	143.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,617	8.6	98.6
5	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	11,785	4.9	56.4
6	Cerebrovascular diseases (I60-I69)	10,276	4.3	49.2
7	Diabetes mellitus (E10-E149)	10,080	4.2	48.2
8	Chronic lower respiratory diseases (J40-J47)	6,203	2.6	29.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,176	2.2	24.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,895	1.6	18.6
...	All other causes (residual)	55,146	23.1	263.8
Black, non-Hispanic, male, 1-4 years				
...	All causes	553	100.0	45.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	178	32.2	14.5
2	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	80	14.5	6.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	9.4	4.2
4	Malignant neoplasms (C00-C97)	26	4.7	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	4.0	1.8
6	Certain conditions originating in the perinatal period (P00-P96)	17	3.1	*
7	Influenza and pneumonia (J09-J18)	11	2.0	*
8	Cerebrovascular diseases (I60-I69)	10	1.8	*
9	Chronic lower respiratory diseases (J40-J47)	5	0.9	*
10	Septicemia (A40-A41)	3	0.5	*
10	Meningitis (G00,G03)	3	0.5	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
10	Acute bronchitis and bronchiolitis (J20-J21)	3	0.5	*
...	All other causes (residual)	140	25.3	11.4

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
Black, non-Hispanic, male, 5–9 years				
...	All causes	343	100.0	21.8
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	115	33.5	7.3
2	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	36	10.5	2.3
3	Malignant neoplasms (C00–C97)	32	9.3	2.0
4	Chronic lower respiratory diseases (J40–J47)	21	6.1	1.3
5	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	16	4.7	*
6	Influenza and pneumonia (J09–J18)	10	2.9	*
6	Diseases of heart (I00–I09,I11,I13,I20–I51)	10	2.9	*
8	Certain conditions originating in the perinatal period (P00–P96)	6	1.7	*
9	Anemias (D50–D64)	4	1.2	*
9	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	4	1.2	*
...	All other causes (residual)	89	25.9	5.6
Black, non-Hispanic, male, 10–14 years				
...	All causes	535	100.0	33.3
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	122	22.8	7.6
2	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	109	20.4	6.8
3	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	59	11.0	3.7
4	Malignant neoplasms (C00–C97)	39	7.3	2.4
5	Chronic lower respiratory diseases (J40–J47)	37	6.9	2.3
6	Diseases of heart (I00–I09,I11,I13,I20–I51)	17	3.2	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	17	3.2	*
8	Diabetes mellitus (E10–E149)	8	1.5	*
9	Cerebrovascular diseases (I60–I69)	6	1.1	*
9	COVID-19 (†U07.1)	6	1.1	*
...	All other causes (residual)	115	21.5	7.2
Black, non-Hispanic, male, 15–19 years				
...	All causes	2,618	100.0	166.2
1	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	1,453	55.5	92.3
2	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	568	21.7	36.1
3	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	195	7.4	12.4
4	Diseases of heart (I00–I09,I11,I13,I20–I51)	61	2.3	3.9
5	Malignant neoplasms (C00–C97)	53	2.0	3.4
6	Legal intervention (Y35,Y89.0)	22	0.8	1.4
7	Chronic lower respiratory diseases (J40–J47)	21	0.8	1.3
8	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	20	0.8	1.3
9	COVID-19 (†U07.1)	18	0.7	*
10	Diabetes mellitus (E10–E149)	16	0.6	*
...	All other causes (residual)	191	7.3	12.1
Black, non-Hispanic, male, 20–24 years				
...	All causes	4,943	100.0	299.1
1	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	2,309	46.7	139.7
2	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	1,361	27.5	82.4
3	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	485	9.8	29.4
4	Diseases of heart (I00–I09,I11,I13,I20–I51)	96	1.9	5.8
5	Malignant neoplasms (C00–C97)	59	1.2	3.6
6	COVID-19 (†U07.1)	56	1.1	3.4
7	Diabetes mellitus (E10–E149)	41	0.8	2.5
8	Chronic lower respiratory diseases (J40–J47)	33	0.7	2.0
9	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	31	0.6	1.9
10	Human immunodeficiency virus (HIV) disease (B20–B24)	28	0.6	1.7
...	All other causes (residual)	444	9.0	26.9

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
Black, non-Hispanic, male, 25–34 years				
...	All causes	12,417	100.0	360.1
1	Assault (homicide)..... (†U01-†U02.9,X85–Y09,Y87.1)	4,052	32.6	117.5
2	Accidents (unintentional injuries)..... (V01–X59,Y85–Y86)	3,736	30.1	108.4
3	Intentional self-harm (suicide)..... (†U03,X60–X84,Y87.0)	837	6.7	24.3
4	Diseases of heart..... (I00–I09,I11,I13,I20–I51)	776	6.2	22.5
5	COVID-19..... (†U07.1)	349	2.8	10.1
6	Malignant neoplasms..... (C00–C97)	275	2.2	8.0
7	Diabetes mellitus..... (E10–E149)	224	1.8	6.5
8	Human immunodeficiency virus (HIV) disease..... (B20–B24)	216	1.7	6.3
9	Chronic lower respiratory diseases..... (J40–J47)	103	0.8	3.0
10	Cerebrovascular diseases..... (I60–I69)	99	0.8	2.9
...	All other causes..... (residual)	1,750	14.1	50.8
Black, non-Hispanic, male, 35–44 years				
...	All causes	14,008	100.0	528.7
1	Accidents (unintentional injuries)..... (V01–X59,Y85–Y86)	3,586	25.6	135.4
2	Diseases of heart..... (I00–I09,I11,I13,I20–I51)	2,271	16.2	85.7
3	Assault (homicide)..... (†U01-†U02.9,X85–Y09,Y87.1)	1,982	14.1	74.8
4	COVID-19..... (†U07.1)	864	6.2	32.6
5	Malignant neoplasms..... (C00–C97)	756	5.4	28.5
6	Diabetes mellitus..... (E10–E149)	567	4.0	21.4
7	Intentional self-harm (suicide)..... (†U03,X60–X84,Y87.0)	457	3.3	17.2
8	Cerebrovascular diseases..... (I60–I69)	313	2.2	11.8
9	Human immunodeficiency virus (HIV) disease..... (B20–B24)	267	1.9	10.1
10	Chronic liver disease and cirrhosis..... (K70,K73–K74)	188	1.3	7.1
...	All other causes..... (residual)	2,757	19.7	104.1
Black, non-Hispanic, male, 45–54 years				
...	All causes	23,225	100.0	964.2
1	Diseases of heart..... (I00–I09,I11,I13,I20–I51)	5,520	23.8	229.2
2	Accidents (unintentional injuries)..... (V01–X59,Y85–Y86)	3,546	15.3	147.2
3	COVID-19..... (†U07.1)	2,426	10.4	100.7
4	Malignant neoplasms..... (C00–C97)	2,425	10.4	100.7
5	Diabetes mellitus..... (E10–E149)	1,161	5.0	48.2
6	Assault (homicide)..... (†U01-†U02.9,X85–Y09,Y87.1)	936	4.0	38.9
7	Cerebrovascular diseases..... (I60–I69)	834	3.6	34.6
8	Nephritis, nephrotic syndrome and nephrosis..... (N00–N07,N17–N19,N25–N27)	495	2.1	20.6
9	Chronic liver disease and cirrhosis..... (K70,K73–K74)	439	1.9	18.2
10	Essential hypertension and hypertensive renal disease..... (I10,I12,I15)	414	1.8	17.2
...	All other causes..... (residual)	5,029	21.7	208.8
Black, non-Hispanic, male, 55–64 years				
...	All causes	48,903	100.0	2,104.5
1	Diseases of heart..... (I00–I09,I11,I13,I20–I51)	12,110	24.8	521.1
2	Malignant neoplasms..... (C00–C97)	8,751	17.9	376.6
3	COVID-19..... (†U07.1)	5,771	11.8	248.3
4	Accidents (unintentional injuries)..... (V01–X59,Y85–Y86)	4,064	8.3	174.9
5	Diabetes mellitus..... (E10–E149)	2,495	5.1	107.4
6	Cerebrovascular diseases..... (I60–I69)	2,070	4.2	89.1
7	Chronic lower respiratory diseases..... (J40–J47)	1,157	2.4	49.8
8	Nephritis, nephrotic syndrome and nephrosis..... (N00–N07,N17–N19,N25–N27)	1,074	2.2	46.2
9	Chronic liver disease and cirrhosis..... (K70,K73–K74)	956	2.0	41.1
10	Essential hypertension and hypertensive renal disease..... (I10,I12,I15)	884	1.8	38.0
...	All other causes..... (residual)	9,571	19.6	411.9

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
Black, non-Hispanic, male, 65 years and older				
...	All causes	128,151	100.0	6,002.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,281	22.8	1,371.6
2	Malignant neoplasms (C00-C97)	23,272	18.2	1,090.1
3	COVID-19 (†U07.1)	20,444	16.0	957.6
4	Cerebrovascular diseases (I60-I69)	6,910	5.4	323.7
5	Diabetes mellitus (E10-E149)	5,566	4.3	260.7
6	Chronic lower respiratory diseases (J40-J47)	4,430	3.5	207.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,348	2.6	156.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,094	2.4	144.9
9	Alzheimer disease (G30)	3,003	2.3	140.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,400	1.9	112.4
...	All other causes (residual)	26,403	20.6	1,236.8
Black, non-Hispanic, male, 65-74 years				
...	All causes	58,206	100.0	4,116.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,563	23.3	959.1
2	Malignant neoplasms (C00-C97)	11,971	20.6	846.6
3	COVID-19 (†U07.1)	8,876	15.2	627.7
4	Cerebrovascular diseases (I60-I69)	2,856	4.9	202.0
5	Diabetes mellitus (E10-E149)	2,765	4.8	195.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,985	3.4	140.4
7	Chronic lower respiratory diseases (J40-J47)	1,983	3.4	140.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,454	2.5	102.8
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,061	1.8	75.0
10	Influenza and pneumonia (J09-J18)	992	1.7	70.2
...	All other causes (residual)	10,700	18.4	756.7
Black, non-Hispanic, male, 75-84 years				
...	All causes	43,220	100.0	7,757.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,508	22.0	1,706.5
2	Malignant neoplasms (C00-C97)	7,781	18.0	1,396.6
3	COVID-19 (†U07.1)	7,443	17.2	1,335.9
4	Cerebrovascular diseases (I60-I69)	2,436	5.6	437.2
5	Diabetes mellitus (E10-E149)	1,900	4.4	341.0
6	Chronic lower respiratory diseases (J40-J47)	1,611	3.7	289.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,192	2.8	213.9
8	Alzheimer disease (G30)	1,121	2.6	201.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	792	1.8	142.2
10	Influenza and pneumonia (J09-J18)	766	1.8	137.5
...	All other causes (residual)	8,670	20.1	1,556.1
Black, non-Hispanic, male, 85 years and older				
...	All causes	26,725	100.0	16,335.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,210	23.2	3,795.9
2	COVID-19 (†U07.1)	4,125	15.4	2,521.4
3	Malignant neoplasms (C00-C97)	3,520	13.2	2,151.6
4	Cerebrovascular diseases (I60-I69)	1,618	6.1	989.0
5	Alzheimer disease (G30)	1,449	5.4	885.7
6	Diabetes mellitus (E10-E149)	901	3.4	550.7
7	Chronic lower respiratory diseases (J40-J47)	836	3.1	511.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	702	2.6	429.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	547	2.0	334.4
10	Influenza and pneumonia (J09-J18)	532	2.0	325.2
...	All other causes (residual)	6,285	23.5	3,841.8

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
Black, non-Hispanic, female, all ages ⁴				
...	All causes	212,776	100.0	936.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,121	20.3	189.8
2	Malignant neoplasms (C00-C97)	35,270	16.6	155.3
3	COVID-19 (†U07.1)	26,582	12.5	117.0
4	Cerebrovascular diseases (I60-I69)	12,128	5.7	53.4
5	Diabetes mellitus (E10-E149)	9,415	4.4	41.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,516	4.0	37.5
7	Alzheimer disease (G30)	7,807	3.7	34.4
8	Chronic lower respiratory diseases (J40-J47)	6,158	2.9	27.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,405	2.5	23.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,156	2.0	18.3
...	All other causes (residual)	54,218	25.5	238.7
Black, non-Hispanic, female, 1-4 years				
...	All causes	413	100.0	34.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	23.0	8.0
2	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	67	16.2	5.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	10.4	3.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	5.8	2.0
5	Malignant neoplasms (C00-C97)	20	4.8	1.7
6	Septicemia (A40-A41)	10	2.4	*
7	Cerebrovascular diseases (I60-I69)	8	1.9	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	7	1.7	*
9	Influenza and pneumonia (J09-J18)	6	1.5	*
10	Certain conditions originating in the perinatal period (P00-P96)	5	1.2	*
...	All other causes (residual)	128	31.0	10.7
Black, non-Hispanic, female, 5-9 years				
...	All causes	251	100.0	16.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	65	25.9	4.3
2	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	45	17.9	2.9
3	Malignant neoplasms (C00-C97)	24	9.6	1.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	5.6	*
4	Chronic lower respiratory diseases (J40-J47)	14	5.6	*
6	Influenza and pneumonia (J09-J18)	9	3.6	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	2.8	*
8	Cerebrovascular diseases (I60-I69)	5	2.0	*
8	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	5	2.0	*
8	COVID-19 (†U07.1)	5	2.0	*
...	All other causes (residual)	58	23.1	3.8
Black, non-Hispanic, female, 10-14 years				
...	All causes	278	100.0	17.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	48	17.3	3.1
2	Malignant neoplasms (C00-C97)	33	11.9	2.1
3	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	30	10.8	1.9
4	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	29	10.4	1.9
5	Chronic lower respiratory diseases (J40-J47)	21	7.6	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	5.4	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	4.3	*
8	Diabetes mellitus (E10-E149)	11	4.0	*
9	Influenza and pneumonia (J09-J18)	6	2.2	*
10	Cerebrovascular diseases (I60-I69)	4	1.4	*
...	All other causes (residual)	69	24.8	4.4

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
Black, non-Hispanic, female, 15-19 years				
...	All causes	771	100.0	50.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	229	29.7	14.9
2	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	166	21.5	10.8
3	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	72	9.3	4.7
4	Malignant neoplasms (C00-C97)	41	5.3	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	3.4	1.7
6	Chronic lower respiratory diseases (J40-J47)	20	2.6	1.3
7	Pregnancy, childbirth and the puerperium (O00-O99)	19	2.5	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	2.1	*
9	COVID-19 (†U07.1)	12	1.6	*
10	Diabetes mellitus (E10-E149)	11	1.4	*
...	All other causes (residual)	159	20.6	10.4
Black, non-Hispanic, female, 20-24 years				
...	All causes	1,518	100.0	94.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	544	35.8	33.8
2	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	289	19.0	17.9
3	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	87	5.7	5.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	66	4.3	4.1
5	Malignant neoplasms (C00-C97)	55	3.6	3.4
6	Pregnancy, childbirth and the puerperium (O00-O99)	50	3.3	3.1
7	COVID-19 (†U07.1)	48	3.2	3.0
8	Diabetes mellitus (E10-E149)	34	2.2	2.1
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	1.6	1.6
10	Influenza and pneumonia (J09-J18)	17	1.1	*
...	All other causes (residual)	303	20.0	18.8
Black, non-Hispanic, female, 25-34 years				
...	All causes	5,041	100.0	145.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,514	30.0	43.8
2	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	484	9.6	14.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	462	9.2	13.4
4	Malignant neoplasms (C00-C97)	375	7.4	10.8
5	COVID-19 (†U07.1)	234	4.6	6.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	192	3.8	5.5
7	Diabetes mellitus (E10-E149)	174	3.5	5.0
7	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	174	3.5	5.0
9	Cerebrovascular diseases (I60-I69)	82	1.6	2.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	73	1.4	2.1
...	All other causes (residual)	1,277	25.3	36.9
Black, non-Hispanic, female, 35-44 years				
...	All causes	8,208	100.0	281.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,289	15.7	44.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,288	15.7	44.1
3	Malignant neoplasms (C00-C97)	1,145	13.9	39.2
4	COVID-19 (†U07.1)	627	7.6	21.5
5	Diabetes mellitus (E10-E149)	336	4.1	11.5
6	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	310	3.8	10.6
7	Cerebrovascular diseases (I60-I69)	272	3.3	9.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	173	2.1	5.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	172	2.1	5.9
10	Pregnancy, childbirth and the puerperium (O00-O99)	160	1.9	5.5
...	All other causes (residual)	2,436	29.7	83.5

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
Black, non-Hispanic, female, 45–54 years				
...	All causes	15,384	100.0	560.3
1	Malignant neoplasms..... (C00–C97)	3,094	20.1	112.7
2	Diseases of heart..... (I00–I09,I11,I13,I20–I51)	2,971	19.3	108.2
3	COVID-19..... (†U07.1)	1,608	10.5	58.6
4	Accidents (unintentional injuries)..... (V01–X59,Y85–Y86)	1,289	8.4	46.9
5	Diabetes mellitus..... (E10–E149)	802	5.2	29.2
6	Cerebrovascular diseases..... (I60–I69)	657	4.3	23.9
7	Chronic liver disease and cirrhosis..... (K70,K73–K74)	336	2.2	12.2
8	Nephritis, nephrotic syndrome and nephrosis..... (N00–N07,N17–N19,N25–N27)	328	2.1	11.9
9	Septicemia..... (A40–A41)	301	2.0	11.0
10	Chronic lower respiratory diseases..... (J40–J47)	291	1.9	10.6
...	All other causes..... (residual)	3,707	24.1	135.0
Black, non-Hispanic, female, 55–64 years				
...	All causes.....	33,690	100.0	1,224.7
1	Malignant neoplasms..... (C00–C97)	8,231	24.4	299.2
2	Diseases of heart..... (I00–I09,I11,I13,I20–I51)	6,675	19.8	242.6
3	COVID-19..... (†U07.1)	4,128	12.3	150.1
4	Diabetes mellitus..... (E10–E149)	1,763	5.2	64.1
5	Cerebrovascular diseases..... (I60–I69)	1,460	4.3	53.1
6	Accidents (unintentional injuries)..... (V01–X59,Y85–Y86)	1,416	4.2	51.5
7	Chronic lower respiratory diseases..... (J40–J47)	1,182	3.5	43.0
8	Nephritis, nephrotic syndrome and nephrosis..... (N00–N07,N17–N19,N25–N27)	853	2.5	31.0
9	Septicemia..... (A40–A41)	662	2.0	24.1
10	Essential hypertension and hypertensive renal disease..... (I10,I12,I15)	581	1.7	21.1
...	All other causes..... (residual)	6,739	20.0	245.0
Black, non-Hispanic, female, 65 years and older				
...	All causes.....	144,516	100.0	4,614.4
1	Diseases of heart..... (I00–I09,I11,I13,I20–I51)	31,550	21.8	1,007.4
2	Malignant neoplasms..... (C00–C97)	22,247	15.4	710.3
3	COVID-19..... (†U07.1)	19,912	13.8	635.8
4	Cerebrovascular diseases..... (I60–I69)	9,606	6.6	306.7
5	Alzheimer disease..... (G30)	7,719	5.3	246.5
6	Diabetes mellitus..... (E10–E149)	6,282	4.3	200.6
7	Chronic lower respiratory diseases..... (J40–J47)	4,424	3.1	141.3
8	Nephritis, nephrotic syndrome and nephrosis..... (N00–N07,N17–N19,N25–N27)	3,995	2.8	127.6
9	Essential hypertension and hypertensive renal disease..... (I10,I12,I15)	3,206	2.2	102.4
10	Septicemia..... (A40–A41)	2,450	1.7	78.2
...	All other causes..... (residual)	33,125	22.9	1,057.7
Black, non-Hispanic, female, 65–74 years				
...	All causes.....	45,494	100.0	2,414.3
1	Malignant neoplasms..... (C00–C97)	10,396	22.9	551.7
2	Diseases of heart..... (I00–I09,I11,I13,I20–I51)	9,390	20.6	498.3
3	COVID-19..... (†U07.1)	6,494	14.3	344.6
4	Diabetes mellitus..... (E10–E149)	2,417	5.3	128.3
5	Cerebrovascular diseases..... (I60–I69)	2,347	5.2	124.6
6	Chronic lower respiratory diseases..... (J40–J47)	1,598	3.5	84.8
7	Nephritis, nephrotic syndrome and nephrosis..... (N00–N07,N17–N19,N25–N27)	1,340	2.9	71.1
8	Septicemia..... (A40–A41)	901	2.0	47.8
9	Essential hypertension and hypertensive renal disease..... (I10,I12,I15)	891	2.0	47.3
10	Influenza and pneumonia..... (J09–J18)	746	1.6	39.6
...	All other causes..... (residual)	8,974	19.7	476.2

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
Black, non-Hispanic, female, 75–84 years				
...	All causes	46,335	100.0	5,275.1
1	Diseases of heart (I00–I09,I11,I13,I20–I51)	9,799	21.1	1,115.6
2	Malignant neoplasms (C00–C97)	7,459	16.1	849.2
3	COVID-19 (†U07.1)	6,801	14.7	774.3
4	Cerebrovascular diseases (I60–I69)	3,068	6.6	349.3
5	Alzheimer disease (G30)	2,196	4.7	250.0
6	Diabetes mellitus (E10–E149)	2,192	4.7	249.6
7	Chronic lower respiratory diseases (J40–J47)	1,566	3.4	178.3
8	Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)	1,407	3.0	160.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	966	2.1	110.0
10	Septicemia (A40–A41)	835	1.8	95.1
...	All other causes (residual)	10,046	21.7	1,143.7
Black, non-Hispanic, female, 85 years and older				
...	All causes	52,687	100.0	14,271.7
1	Diseases of heart (I00–I09,I11,I13,I20–I51)	12,361	23.5	3,348.3
2	COVID-19 (†U07.1)	6,617	12.6	1,792.4
3	Alzheimer disease (G30)	4,901	9.3	1,327.6
4	Malignant neoplasms (C00–C97)	4,392	8.3	1,189.7
5	Cerebrovascular diseases (I60–I69)	4,191	8.0	1,135.2
6	Diabetes mellitus (E10–E149)	1,673	3.2	453.2
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,349	2.6	365.4
8	Chronic lower respiratory diseases (J40–J47)	1,260	2.4	341.3
9	Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)	1,248	2.4	338.1
10	Influenza and pneumonia (J09–J18)	832	1.6	225.4
...	All other causes (residual)	13,863	26.3	3,755.2
White, non-Hispanic, both sexes, all ages ⁴				
...	All causes	2,491,026	100.0	1,242.4
1	Diseases of heart (I00–I09,I11,I13,I20–I51)	530,355	21.3	264.5
2	Malignant neoplasms (C00–C97)	463,160	18.6	231.0
3	COVID-19 (†U07.1)	209,606	8.4	104.5
4	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	141,335	5.7	70.5
5	Chronic lower respiratory diseases (J40–J47)	130,947	5.3	65.3
6	Cerebrovascular diseases (I60–I69)	116,752	4.7	58.2
7	Alzheimer disease (G30)	109,316	4.4	54.5
8	Diabetes mellitus (E10–E149)	64,098	2.6	32.0
9	Influenza and pneumonia (J09–J18)	39,003	1.6	19.5
10	Chronic liver disease and cirrhosis (K70,K73–K74)	36,919	1.5	18.4
...	All other causes (residual)	649,535	26.1	324.0
White, non-Hispanic, both sexes, 1–4 years				
...	All causes	1,614	100.0	20.3
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	585	36.2	7.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	170	10.5	2.1
3	Malignant neoplasms (C00–C97)	169	10.5	2.1
4	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	92	5.7	1.2
5	Influenza and pneumonia (J09–J18)	43	2.7	0.5
6	Diseases of heart (I00–I09,I11,I13,I20–I51)	38	2.4	0.5
7	Cerebrovascular diseases (I60–I69)	27	1.7	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00–D48)	21	1.3	0.3
9	Certain conditions originating in the perinatal period (P00–P96)	18	1.1	*
10	Septicemia (A40–A41)	17	1.1	*
...	All other causes (residual)	434	26.9	5.5

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
White, non-Hispanic, both sexes, 5–9 years				
...	All causes	1,019	100.0	9.8
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	335	32.9	3.2
2	Malignant neoplasms (C00–C97)	205	20.1	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	93	9.1	0.9
4	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	57	5.6	0.5
5	Influenza and pneumonia (J09–J18)	22	2.2	0.2
6	Diseases of heart (I00–I09,I11,I13,I20–I51)	21	2.1	0.2
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00–D48)	17	1.7	*
8	Chronic lower respiratory diseases (J40–J47)	11	1.1	*
9	Cerebrovascular diseases (I60–I69)	10	1.0	*
10	Septicemia (A40–A41)	9	0.9	*
...	All other causes (residual)	239	23.5	2.3
White, non-Hispanic, both sexes, 10–14 years				
...	All causes	1,645	100.0	15.3
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	465	28.3	4.3
2	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	331	20.1	3.1
3	Malignant neoplasms (C00–C97)	205	12.5	1.9
4	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	88	5.3	0.8
5	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	67	4.1	0.6
6	Diseases of heart (I00–I09,I11,I13,I20–I51)	53	3.2	0.5
7	Influenza and pneumonia (J09–J18)	24	1.5	0.2
8	Diabetes mellitus (E10–E149)	23	1.4	0.2
9	Chronic lower respiratory diseases (J40–J47)	20	1.2	0.2
10	Cerebrovascular diseases (I60–I69)	18	1.1	*
...	All other causes (residual)	351	21.3	3.3
White, non-Hispanic, both sexes, 15–19 years				
...	All causes	5,505	100.0	48.8
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	2,496	45.3	22.1
2	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	1,330	24.2	11.8
3	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	357	6.5	3.2
4	Malignant neoplasms (C00–C97)	288	5.2	2.6
5	Diseases of heart (I00–I09,I11,I13,I20–I51)	151	2.7	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	101	1.8	0.9
7	Diabetes mellitus (E10–E149)	41	0.7	0.4
8	Influenza and pneumonia (J09–J18)	35	0.6	0.3
9	Cerebrovascular diseases (I60–I69)	32	0.6	0.3
10	COVID-19 (†U07.1)	31	0.6	0.3
...	All other causes (residual)	643	11.7	5.7
White, non-Hispanic, both sexes, 20–24 years				
...	All causes	11,060	100.0	93.5
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	5,766	52.1	48.8
2	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	2,333	21.1	19.7
3	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	532	4.8	4.5
4	Malignant neoplasms (C00–C97)	358	3.2	3.0
5	Diseases of heart (I00–I09,I11,I13,I20–I51)	274	2.5	2.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	102	0.9	0.9
7	Diabetes mellitus (E10–E149)	100	0.9	0.8
8	COVID-19 (†U07.1)	82	0.7	0.7
9	Pregnancy, childbirth and the puerperium (O00–O99)	69	0.6	0.6
10	Influenza and pneumonia (J09–J18)	65	0.6	0.5
...	All other causes (residual)	1,379	12.5	11.7

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
White, non-Hispanic, both sexes, 25–34 years				
...	All causes	39,871	100.0	155.8
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	19,749	49.5	77.2
2	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	5,629	14.1	22.0
3	Diseases of heart (I00–I09,I11,I13,I20–I51)	2,013	5.0	7.9
4	Malignant neoplasms (C00–C97)	1,997	5.0	7.8
5	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	1,309	3.3	5.1
6	Chronic liver disease and cirrhosis (K70,K73–K74)	933	2.3	3.6
7	Diabetes mellitus (E10–E149)	550	1.4	2.1
8	COVID-19 (†U07.1)	473	1.2	1.8
9	Influenza and pneumonia (J09–J18)	305	0.8	1.2
10	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	286	0.7	1.1
...	All other causes (residual)	6,627	16.6	25.9
White, non-Hispanic, both sexes, 35–44 years				
...	All causes	59,483	100.0	246.4
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	20,787	34.9	86.1
2	Diseases of heart (I00–I09,I11,I13,I20–I51)	6,533	11.0	27.1
3	Malignant neoplasms (C00–C97)	6,252	10.5	25.9
4	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	5,568	9.4	23.1
5	Chronic liver disease and cirrhosis (K70,K73–K74)	3,051	5.1	12.6
6	Diabetes mellitus (E10–E149)	1,345	2.3	5.6
7	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	1,240	2.1	5.1
8	COVID-19 (†U07.1)	1,183	2.0	4.9
9	Cerebrovascular diseases (I60–I69)	887	1.5	3.7
10	Influenza and pneumonia (J09–J18)	608	1.0	2.5
...	All other causes (residual)	12,029	20.2	49.8
White, non-Hispanic, both sexes, 45–54 years				
...	All causes	114,109	100.0	460.3
1	Malignant neoplasms (C00–C97)	22,581	19.8	91.1
2	Diseases of heart (I00–I09,I11,I13,I20–I51)	20,648	18.1	83.3
3	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	18,586	16.3	75.0
4	Chronic liver disease and cirrhosis (K70,K73–K74)	6,480	5.7	26.1
5	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	5,963	5.2	24.1
6	COVID-19 (†U07.1)	4,629	4.1	18.7
7	Diabetes mellitus (E10–E149)	3,850	3.4	15.5
8	Cerebrovascular diseases (I60–I69)	2,814	2.5	11.4
9	Chronic lower respiratory diseases (J40–J47)	2,671	2.3	10.8
10	Septicemia (A40–A41)	1,517	1.3	6.1
...	All other causes (residual)	24,370	21.4	98.3
White, non-Hispanic, both sexes, 55–64 years				
...	All causes	291,512	100.0	990.4
1	Malignant neoplasms (C00–C97)	79,451	27.3	269.9
2	Diseases of heart (I00–I09,I11,I13,I20–I51)	59,407	20.4	201.8
3	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	19,700	6.8	66.9
4	COVID-19 (†U07.1)	16,406	5.6	55.7
5	Chronic lower respiratory diseases (J40–J47)	15,462	5.3	52.5
6	Chronic liver disease and cirrhosis (K70,K73–K74)	11,764	4.0	40.0
7	Diabetes mellitus (E10–E149)	10,388	3.6	35.3
8	Cerebrovascular diseases (I60–I69)	8,350	2.9	28.4
9	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	6,254	2.1	21.2
10	Septicemia (A40–A41)	4,128	1.4	14.0
...	All other causes (residual)	60,202	20.7	204.5

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
White, non-Hispanic, both sexes, 65 years and older				
...	All causes	1,956,915	100.0	4,616.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	441,089	22.5	1,040.6
2	Malignant neoplasms (C00-C97)	351,626	18.0	829.5
3	COVID-19 (†U07.1)	186,770	9.5	440.6
4	Chronic lower respiratory diseases (J40-J47)	112,082	5.7	264.4
5	Alzheimer disease (G30)	108,169	5.5	255.2
6	Cerebrovascular diseases (I60-I69)	104,245	5.3	245.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	52,312	2.7	123.4
8	Diabetes mellitus (E10-E149)	47,796	2.4	112.8
9	Parkinson disease (G20-G21)	33,689	1.7	79.5
10	Influenza and pneumonia (J09-J18)	32,316	1.7	76.2
...	All other causes (residual)	486,821	24.9	1,148.5
White, non-Hispanic, both sexes, 65-74 years				
...	All causes	485,693	100.0	1,991.2
1	Malignant neoplasms (C00-C97)	135,011	27.8	553.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	97,551	20.1	399.9
3	COVID-19 (†U07.1)	39,779	8.2	163.1
4	Chronic lower respiratory diseases (J40-J47)	33,124	6.8	135.8
5	Cerebrovascular diseases (I60-I69)	17,655	3.6	72.4
6	Diabetes mellitus (E10-E149)	17,325	3.6	71.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,893	2.9	57.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	8,889	1.8	36.4
9	Alzheimer disease (G30)	7,332	1.5	30.1
10	Influenza and pneumonia (J09-J18)	7,267	1.5	29.8
...	All other causes (residual)	107,867	22.2	442.2
White, non-Hispanic, both sexes, 75-84 years				
...	All causes	639,678	100.0	5,012.1
1	Malignant neoplasms (C00-C97)	131,708	20.6	1,032.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	130,730	20.4	1,024.3
3	COVID-19 (†U07.1)	63,506	9.9	497.6
4	Chronic lower respiratory diseases (J40-J47)	43,431	6.8	340.3
5	Cerebrovascular diseases (I60-I69)	32,415	5.1	254.0
6	Alzheimer disease (G30)	30,800	4.8	241.3
7	Diabetes mellitus (E10-E149)	16,857	2.6	132.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,742	2.5	123.3
9	Parkinson disease (G20-G21)	14,947	2.3	117.1
10	Influenza and pneumonia (J09-J18)	10,524	1.6	82.5
...	All other causes (residual)	149,018	23.3	1,167.6
White, non-Hispanic, both sexes, 85 years and older				
...	All causes	831,544	100.0	15,887.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	212,808	25.6	4,065.8
2	Malignant neoplasms (C00-C97)	84,907	10.2	1,622.2
3	COVID-19 (†U07.1)	83,485	10.0	1,595.0
4	Alzheimer disease (G30)	70,037	8.4	1,338.1
5	Cerebrovascular diseases (I60-I69)	54,175	6.5	1,035.0
6	Chronic lower respiratory diseases (J40-J47)	35,527	4.3	678.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22,677	2.7	433.3
8	Influenza and pneumonia (J09-J18)	14,525	1.7	277.5
9	Diabetes mellitus (E10-E149)	13,614	1.6	260.1
10	Parkinson disease (G20-G21)	13,552	1.6	258.9
...	All other causes (residual)	226,237	27.2	4,322.3

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
White, non-Hispanic, male, all ages ⁴				
...	All causes	1,282,214	100.0	1,296.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	290,527	22.7	293.9
2	Malignant neoplasms (C00-C97)	246,965	19.3	249.8
3	COVID-19 (†U07.1)	110,300	8.6	111.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	90,081	7.0	91.1
5	Chronic lower respiratory diseases (J40-J47)	61,801	4.8	62.5
6	Cerebrovascular diseases (I60-I69)	49,490	3.9	50.1
7	Diabetes mellitus (E10-E149)	37,120	2.9	37.5
8	Alzheimer disease (G30)	33,846	2.6	34.2
9	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	28,433	2.2	28.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	23,028	1.8	23.3
...	All other causes (residual)	310,623	24.2	314.2
White, non-Hispanic, male, 1-4 years				
...	All causes	940	100.0	23.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	373	39.7	9.2
2	Malignant neoplasms (C00-C97)	90	9.6	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	86	9.1	2.1
4	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	55	5.9	1.4
5	Influenza and pneumonia (J09-J18)	28	3.0	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	2.2	0.5
7	Cerebrovascular diseases (I60-I69)	14	1.5	*
8	Septicemia (A40-A41)	10	1.1	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.1	*
10	Certain conditions originating in the perinatal period (P00-P96)	7	0.7	*
...	All other causes (residual)	246	26.2	6.1
White, non-Hispanic, male, 5-9 years				
...	All causes	585	100.0	10.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	212	36.2	4.0
2	Malignant neoplasms (C00-C97)	108	18.5	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	8.5	0.9
4	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	34	5.8	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	2.1	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.4	*
6	Influenza and pneumonia (J09-J18)	8	1.4	*
8	Chronic lower respiratory diseases (J40-J47)	5	0.9	*
9	Cerebrovascular diseases (I60-I69)	4	0.7	*
10	Septicemia (A40-A41)	3	0.5	*
10	Anemias (D50-D64)	3	0.5	*
10	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	3	0.5	*
10	Pneumonitis due to solids and liquids (J69)	3	0.5	*
...	All other causes (residual)	132	22.6	2.5

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
White, non-Hispanic, male, 10–14 years				
...	All causes	1,030	100.0	18.7
1	Accidents (unintentional injuries)..... (V01–X59,Y85–Y86)	296	28.7	5.4
2	Intentional self-harm (suicide)..... (†U03,X60–X84,Y87.0)	227	22.0	4.1
3	Malignant neoplasms..... (C00–C97)	116	11.3	2.1
4	Assault (homicide)..... (†U01–†U02.9,X85–Y09,Y87.1)	68	6.6	1.2
5	Congenital malformations, deformations and chromosomal abnormalities..... (Q00–Q99)	37	3.6	0.7
6	Diseases of heart..... (I00–I09,I11,I13,I20–I51)	27	2.6	0.5
7	Chronic lower respiratory diseases..... (J40–J47)	13	1.3	*
8	Influenza and pneumonia..... (J09–J18)	12	1.2	*
9	Diabetes mellitus..... (E10–E149)	11	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior..... (D00–D48)	8	0.8	*
...	All other causes..... (residual)	215	20.9	3.9
White, non-Hispanic, male, 15–19 years				
...	All causes.....	3,889	100.0	67.3
1	Accidents (unintentional injuries)..... (V01–X59,Y85–Y86)	1,781	45.8	30.8
2	Intentional self-harm (suicide)..... (†U03,X60–X84,Y87.0)	1,049	27.0	18.2
3	Assault (homicide)..... (†U01–†U02.9,X85–Y09,Y87.1)	271	7.0	4.7
4	Malignant neoplasms..... (C00–C97)	173	4.4	3.0
5	Diseases of heart..... (I00–I09,I11,I13,I20–I51)	92	2.4	1.6
6	Congenital malformations, deformations and chromosomal abnormalities..... (Q00–Q99)	52	1.3	0.9
7	Diabetes mellitus..... (E10–E149)	25	0.6	0.4
8	Influenza and pneumonia..... (J09–J18)	19	0.5	*
9	Chronic lower respiratory diseases..... (J40–J47)	15	0.4	*
10	COVID-19..... (†U07.1)	13	0.3	*
...	All other causes..... (residual)	399	10.3	6.9
White, non-Hispanic, male, 20–24 years				
...	All causes.....	7,963	100.0	131.4
1	Accidents (unintentional injuries)..... (V01–X59,Y85–Y86)	4,167	52.3	68.7
2	Intentional self-harm (suicide)..... (†U03,X60–X84,Y87.0)	1,932	24.3	31.9
3	Assault (homicide)..... (†U01–†U02.9,X85–Y09,Y87.1)	389	4.9	6.4
4	Malignant neoplasms..... (C00–C97)	202	2.5	3.3
5	Diseases of heart..... (I00–I09,I11,I13,I20–I51)	175	2.2	2.9
6	Diabetes mellitus..... (E10–E149)	68	0.9	1.1
7	Congenital malformations, deformations and chromosomal abnormalities..... (Q00–Q99)	59	0.7	1.0
8	COVID-19..... (†U07.1)	57	0.7	0.9
9	Influenza and pneumonia..... (J09–J18)	38	0.5	0.6
10	Chronic lower respiratory diseases..... (J40–J47)	29	0.4	0.5
...	All other causes..... (residual)	847	10.6	14.0
White, non-Hispanic, male, 25–34 years				
...	All causes.....	27,596	100.0	211.8
1	Accidents (unintentional injuries)..... (V01–X59,Y85–Y86)	14,271	51.7	109.5
2	Intentional self-harm (suicide)..... (†U03,X60–X84,Y87.0)	4,586	16.6	35.2
3	Diseases of heart..... (I00–I09,I11,I13,I20–I51)	1,340	4.9	10.3
4	Malignant neoplasms..... (C00–C97)	948	3.4	7.3
5	Assault (homicide)..... (†U01–†U02.9,X85–Y09,Y87.1)	939	3.4	7.2
6	Chronic liver disease and cirrhosis..... (K70,K73–K74)	577	2.1	4.4
7	Diabetes mellitus..... (E10–E149)	364	1.3	2.8
8	COVID-19..... (†U07.1)	301	1.1	2.3
9	Influenza and pneumonia..... (J09–J18)	178	0.6	1.4
10	Congenital malformations, deformations and chromosomal abnormalities..... (Q00–Q99)	169	0.6	1.3
...	All other causes..... (residual)	3,923	14.2	30.1

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
White, non-Hispanic, male, 35-44 years				
...	All causes	38,538	100.0	317.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,657	38.0	120.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,508	11.7	37.1
3	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	4,338	11.3	35.7
4	Malignant neoplasms (C00-C97)	2,832	7.3	23.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,818	4.7	15.0
6	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	888	2.3	7.3
7	Diabetes mellitus (E10-E149)	843	2.2	6.9
8	COVID-19 (†U07.1)	719	1.9	5.9
9	Cerebrovascular diseases (I60-I69)	527	1.4	4.3
10	Influenza and pneumonia (J09-J18)	316	0.8	2.6
...	All other causes (residual)	7,092	18.4	58.4
White, non-Hispanic, male, 45-54 years				
...	All causes	70,872	100.0	572.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,717	20.8	118.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,894	18.2	104.2
3	Malignant neoplasms (C00-C97)	11,155	15.7	90.1
4	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	4,501	6.4	36.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,018	5.7	32.5
6	COVID-19 (†U07.1)	2,890	4.1	23.3
7	Diabetes mellitus (E10-E149)	2,527	3.6	20.4
8	Cerebrovascular diseases (I60-I69)	1,609	2.3	13.0
9	Chronic lower respiratory diseases (J40-J47)	1,238	1.7	10.0
10	Influenza and pneumonia (J09-J18)	811	1.1	6.6
...	All other causes (residual)	14,512	20.5	117.3
White, non-Hispanic, male, 55-64 years				
...	All causes	179,187	100.0	1,246.2
1	Malignant neoplasms (C00-C97)	43,358	24.2	301.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	42,047	23.5	292.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,645	7.6	94.9
4	COVID-19 (†U07.1)	10,337	5.8	71.9
5	Chronic lower respiratory diseases (J40-J47)	7,805	4.4	54.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	7,747	4.3	53.9
7	Diabetes mellitus (E10-E149)	6,677	3.7	46.4
8	Cerebrovascular diseases (I60-I69)	4,909	2.7	34.1
9	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	4,827	2.7	33.6
10	Influenza and pneumonia (J09-J18)	2,300	1.3	16.0
...	All other causes (residual)	35,535	19.8	247.1
White, non-Hispanic, male, 65 years and older				
...	All causes	947,020	100.0	4,933.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	227,523	24.0	1,185.3
2	Malignant neoplasms (C00-C97)	187,974	19.8	979.3
3	COVID-19 (†U07.1)	95,967	10.1	500.0
4	Chronic lower respiratory diseases (J40-J47)	52,390	5.5	272.9
5	Cerebrovascular diseases (I60-I69)	42,208	4.5	219.9
6	Alzheimer disease (G30)	33,377	3.5	173.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27,472	2.9	143.1
8	Diabetes mellitus (E10-E149)	26,603	2.8	138.6
9	Parkinson disease (G20-G21)	20,593	2.2	107.3
10	Influenza and pneumonia (J09-J18)	16,188	1.7	84.3
...	All other causes (residual)	216,725	22.9	1,129.1

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
White, non-Hispanic, male, 65–74 years				
...	All causes	282,869	100.0	2,445.1
1	Malignant neoplasms (C00–C97)	75,135	26.6	649.5
2	Diseases of heart (I00–I09,I11,I13,I20–I51)	63,741	22.5	551.0
3	COVID-19 (†U07.1)	24,548	8.7	212.2
4	Chronic lower respiratory diseases (J40–J47)	17,169	6.1	148.4
5	Diabetes mellitus (E10–E149)	10,653	3.8	92.1
6	Cerebrovascular diseases (I60–I69)	9,341	3.3	80.7
7	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	9,170	3.2	79.3
8	Chronic liver disease and cirrhosis (K70,K73–K74)	5,646	2.0	48.8
9	Influenza and pneumonia (J09–J18)	4,066	1.4	35.1
10	Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)	3,924	1.4	33.9
...	All other causes (residual)	59,476	21.0	514.1
White, non-Hispanic, male, 75–84 years				
...	All causes	336,942	100.0	5,906.2
1	Diseases of heart (I00–I09,I11,I13,I20–I51)	75,019	22.3	1,315.0
2	Malignant neoplasms (C00–C97)	70,939	21.1	1,243.5
3	COVID-19 (†U07.1)	36,151	10.7	633.7
4	Chronic lower respiratory diseases (J40–J47)	20,840	6.2	365.3
5	Cerebrovascular diseases (I60–I69)	15,160	4.5	265.7
6	Alzheimer disease (G30)	11,505	3.4	201.7
7	Parkinson disease (G20–G21)	9,595	2.8	168.2
8	Diabetes mellitus (E10–E149)	9,587	2.8	168.0
9	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	8,762	2.6	153.6
10	Influenza and pneumonia (J09–J18)	5,612	1.7	98.4
...	All other causes (residual)	73,772	21.9	1,293.1
White, non-Hispanic, male, 85 years and older				
...	All causes	327,209	100.0	17,029.8
1	Diseases of heart (I00–I09,I11,I13,I20–I51)	88,763	27.1	4,619.7
2	Malignant neoplasms (C00–C97)	41,900	12.8	2,180.7
3	COVID-19 (†U07.1)	35,268	10.8	1,835.5
4	Alzheimer disease (G30)	18,990	5.8	988.3
5	Cerebrovascular diseases (I60–I69)	17,707	5.4	921.6
6	Chronic lower respiratory diseases (J40–J47)	14,381	4.4	748.5
7	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	9,540	2.9	496.5
8	Parkinson disease (G20–G21)	7,527	2.3	391.7
9	Influenza and pneumonia (J09–J18)	6,510	2.0	338.8
10	Diabetes mellitus (E10–E149)	6,363	1.9	331.2
...	All other causes (residual)	80,260	24.5	4,177.2
White, non-Hispanic, female, all ages ⁴				
...	All causes	1,208,812	100.0	1,189.4
1	Diseases of heart (I00–I09,I11,I13,I20–I51)	239,828	19.8	236.0
2	Malignant neoplasms (C00–C97)	216,195	17.9	212.7
3	COVID-19 (†U07.1)	99,306	8.2	97.7
4	Alzheimer disease (G30)	75,470	6.2	74.3
5	Chronic lower respiratory diseases (J40–J47)	69,146	5.7	68.0
6	Cerebrovascular diseases (I60–I69)	67,262	5.6	66.2
7	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	51,254	4.2	50.4
8	Diabetes mellitus (E10–E149)	26,978	2.2	26.5
9	Influenza and pneumonia (J09–J18)	19,071	1.6	18.8
10	Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)	16,574	1.4	16.3
...	All other causes (residual)	327,728	27.1	322.5

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
White, non-Hispanic, female, 1-4 years				
...	All causes	674	100.0	17.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	212	31.5	5.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	84	12.5	2.2
3	Malignant neoplasms (C00-C97)	79	11.7	2.0
4	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	37	5.5	1.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	2.5	*
6	Influenza and pneumonia (J09-J18)	15	2.2	*
7	Cerebrovascular diseases (I60-I69)	13	1.9	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.6	*
8	Certain conditions originating in the perinatal period (P00-P96)	11	1.6	*
10	Septicemia (A40-A41)	7	1.0	*
...	All other causes (residual)	188	27.9	4.9
White, non-Hispanic, female, 5-9 years				
...	All causes	434	100.0	8.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	123	28.3	2.4
2	Malignant neoplasms (C00-C97)	97	22.4	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	9.9	0.8
4	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	23	5.3	0.5
5	Influenza and pneumonia (J09-J18)	14	3.2	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	2.1	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	2.1	*
8	Cerebrovascular diseases (I60-I69)	6	1.4	*
8	Chronic lower respiratory diseases (J40-J47)	6	1.4	*
8	Septicemia (A40-A41)	6	1.4	*
...	All other causes (residual)	98	22.6	1.9
White, non-Hispanic, female, 10-14 years				
...	All causes	615	100.0	11.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	169	27.5	3.2
2	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	104	16.9	2.0
3	Malignant neoplasms (C00-C97)	89	14.5	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	4.9	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	4.2	0.5
6	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	20	3.3	0.4
7	Influenza and pneumonia (J09-J18)	12	2.0	*
7	Diabetes mellitus (E10-E149)	12	2.0	*
9	Cerebrovascular diseases (I60-I69)	11	1.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
10	Chronic lower respiratory diseases (J40-J47)	7	1.1	*
...	All other causes (residual)	128	20.8	2.4
White, non-Hispanic, female, 15-19 years				
...	All causes	1,616	100.0	29.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	715	44.2	13.0
2	Intentional self-harm (suicide) (†U03,X60-X84,Y87.0)	281	17.4	5.1
3	Malignant neoplasms (C00-C97)	115	7.1	2.1
4	Assault (homicide) (†U01-†U02.9,X85-Y09,Y87.1)	86	5.3	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	3.7	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	49	3.0	0.9
7	Cerebrovascular diseases (I60-I69)	20	1.2	0.4
8	COVID-19 (†U07.1)	18	1.1	*
9	Diabetes mellitus (E10-E149)	16	1.0	*
9	Influenza and pneumonia (J09-J18)	16	1.0	*
...	All other causes (residual)	241	14.9	4.4

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
White, non-Hispanic, female, 20–24 years				
...	All causes	3,097	100.0	53.7
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	1,599	51.6	27.7
2	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	401	12.9	7.0
3	Malignant neoplasms (C00–C97)	156	5.0	2.7
4	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	143	4.6	2.5
5	Diseases of heart (I00–I09,I11,I13,I20–I51)	99	3.2	1.7
6	Pregnancy, childbirth and the puerperium (O00–O99)	69	2.2	1.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	43	1.4	0.7
8	Diabetes mellitus (E10–E149)	32	1.0	0.6
8	Chronic lower respiratory diseases (J40–J47)	32	1.0	0.6
10	Influenza and pneumonia (J09–J18)	27	0.9	0.5
10	Cerebrovascular diseases (I60–I69)	27	0.9	0.5
...	All other causes (residual)	469	15.1	8.1
White, non-Hispanic, female, 25–34 years				
...	All causes	12,275	100.0	97.8
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	5,478	44.6	43.6
2	Malignant neoplasms (C00–C97)	1,049	8.5	8.4
3	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	1,043	8.5	8.3
4	Diseases of heart (I00–I09,I11,I13,I20–I51)	673	5.5	5.4
5	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	370	3.0	2.9
6	Chronic liver disease and cirrhosis (K70,K73–K74)	356	2.9	2.8
7	Pregnancy, childbirth and the puerperium (O00–O99)	249	2.0	2.0
8	Diabetes mellitus (E10–E149)	186	1.5	1.5
9	COVID-19 (†U07.1)	172	1.4	1.4
10	Influenza and pneumonia (J09–J18)	127	1.0	1.0
...	All other causes (residual)	2,572	21.0	20.5
White, non-Hispanic, female, 35–44 years				
...	All causes	20,945	100.0	174.5
1	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	6,130	29.3	51.1
2	Malignant neoplasms (C00–C97)	3,420	16.3	28.5
3	Diseases of heart (I00–I09,I11,I13,I20–I51)	2,025	9.7	16.9
4	Chronic liver disease and cirrhosis (K70,K73–K74)	1,233	5.9	10.3
5	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	1,230	5.9	10.2
6	Diabetes mellitus (E10–E149)	502	2.4	4.2
7	COVID-19 (†U07.1)	464	2.2	3.9
8	Cerebrovascular diseases (I60–I69)	360	1.7	3.0
9	Assault (homicide) (†U01–†U02.9,X85–Y09,Y87.1)	352	1.7	2.9
10	Influenza and pneumonia (J09–J18)	292	1.4	2.4
...	All other causes (residual)	4,937	23.6	41.1
White, non-Hispanic, female, 45–54 years				
...	All causes	43,237	100.0	348.3
1	Malignant neoplasms (C00–C97)	11,426	26.4	92.0
2	Diseases of heart (I00–I09,I11,I13,I20–I51)	5,931	13.7	47.8
3	Accidents (unintentional injuries) (V01–X59,Y85–Y86)	5,692	13.2	45.8
4	Chronic liver disease and cirrhosis (K70,K73–K74)	2,462	5.7	19.8
5	COVID-19 (†U07.1)	1,739	4.0	14.0
6	Intentional self-harm (suicide) (†U03,X60–X84,Y87.0)	1,462	3.4	11.8
7	Chronic lower respiratory diseases (J40–J47)	1,433	3.3	11.5
8	Diabetes mellitus (E10–E149)	1,323	3.1	10.7
9	Cerebrovascular diseases (I60–I69)	1,205	2.8	9.7
10	Septicemia (A40–A41)	739	1.7	6.0
...	All other causes (residual)	9,825	22.7	79.1

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
White, non-Hispanic, female, 55-64 years				
...	All causes	112,325	100.0	746.1
1	Malignant neoplasms (C00-C97)	36,093	32.1	239.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,360	15.5	115.3
3	Chronic lower respiratory diseases (J40-J47)	7,657	6.8	50.9
4	COVID-19 (†U07.1)	6,069	5.4	40.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,055	5.4	40.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,017	3.6	26.7
7	Diabetes mellitus (E10-E149)	3,711	3.3	24.6
8	Cerebrovascular diseases (I60-I69)	3,441	3.1	22.9
9	Septicemia (A40-A41)	1,945	1.7	12.9
10	Influenza and pneumonia (J09-J18)	1,758	1.6	11.7
...	All other causes (residual)	24,219	21.6	160.9
White, non-Hispanic, female, 65 years and older				
...	All causes	1,009,895	100.0	4,354.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	213,566	21.1	920.8
2	Malignant neoplasms (C00-C97)	163,652	16.2	705.6
3	COVID-19 (†U07.1)	90,803	9.0	391.5
4	Alzheimer disease (G30)	74,792	7.4	322.5
5	Cerebrovascular diseases (I60-I69)	62,037	6.1	267.5
6	Chronic lower respiratory diseases (J40-J47)	59,692	5.9	257.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24,840	2.5	107.1
8	Diabetes mellitus (E10-E149)	21,193	2.1	91.4
9	Influenza and pneumonia (J09-J18)	16,128	1.6	69.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,650	1.5	63.2
...	All other causes (residual)	268,542	26.6	1,157.8
White, non-Hispanic, female, 65-74 years				
...	All causes	202,824	100.0	1,581.7
1	Malignant neoplasms (C00-C97)	59,876	29.5	466.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,810	16.7	263.7
3	Chronic lower respiratory diseases (J40-J47)	15,955	7.9	124.4
4	COVID-19 (†U07.1)	15,231	7.5	118.8
5	Cerebrovascular diseases (I60-I69)	8,314	4.1	64.8
6	Diabetes mellitus (E10-E149)	6,672	3.3	52.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,723	2.3	36.8
8	Alzheimer disease (G30)	4,450	2.2	34.7
9	Septicemia (A40-A41)	3,327	1.6	25.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	3,243	1.6	25.3
...	All other causes (residual)	47,223	23.3	368.3
White, non-Hispanic, female, 75-84 years				
...	All causes	302,736	100.0	4,289.4
1	Malignant neoplasms (C00-C97)	60,769	20.1	861.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	55,711	18.4	789.4
3	COVID-19 (†U07.1)	27,355	9.0	387.6
4	Chronic lower respiratory diseases (J40-J47)	22,591	7.5	320.1
5	Alzheimer disease (G30)	19,295	6.4	273.4
6	Cerebrovascular diseases (I60-I69)	17,255	5.7	244.5
7	Diabetes mellitus (E10-E149)	7,270	2.4	103.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,980	2.3	98.9
9	Parkinson disease (G20-G21)	5,352	1.8	75.8
10	Influenza and pneumonia (J09-J18)	4,912	1.6	69.6
...	All other causes (residual)	75,246	24.9	1,066.1

Table I-1. Deaths, percentage of total deaths, and death rate for the 10 leading causes of death in selected age groups, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are per 100,000 population in specified group. Race and Hispanic-origin categories are consistent with 1997 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black, non-Hispanic and White, non-Hispanic should be interpreted with caution because of misreporting of Hispanic origin and race on the death certificate. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), race and Hispanic origin, sex, and age	Number ^{2,3}	Percent of total deaths	Death rate ^{2,3}
White, non-Hispanic, female, 85 years and older				
...	All causes	504,335	100.0	15,224.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	124,045	24.6	3,744.5
2	Alzheimer disease (G30)	51,047	10.1	1,540.9
3	COVID-19 (†U07.1)	48,217	9.6	1,455.5
4	Malignant neoplasms (C00-C97)	43,007	8.5	1,298.2
5	Cerebrovascular diseases (I60-I69)	36,468	7.2	1,100.8
6	Chronic lower respiratory diseases (J40-J47)	21,146	4.2	638.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,137	2.6	396.6
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,524	1.7	257.3
9	Influenza and pneumonia (J09-J18)	8,015	1.6	241.9
10	Diabetes mellitus (E10-E149)	7,251	1.4	218.9
...	All other causes (residual)	143,478	28.4	4,331.1

... Category not applicable.

* Estimate does not meet National Center for Health Statistics standards of reliability.

¹Based on number of deaths.

²Data for age not stated are included in "all ages" but are not distributed among age groups.

³Data for two or more races and origin not stated are included in "all races and origins" but are not distributed among race and Hispanic-origin groups. For the race and Hispanic-origin categories shown, only one race was reported on the death certificate.

⁴Includes deaths for infants younger than 1 year.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Table I-2. Infant, neonatal, and postneonatal deaths, percentage of total deaths, and mortality rate for the 10 leading causes of infant death, by bridged race and Hispanic origin and sex: United States, 2020

[Rates are infant (younger than 1 year), neonatal (younger than 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic-origin categories are consistent with 1977 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated separately by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black non-Hispanic and White non-Hispanic should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on birth and death certificates. Deaths are based on race and Hispanic origin of decedent; live births are based on race and Hispanic origin of mother. National linked files should be used to measure infant mortality for specific subgroups. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), age, race and Hispanic origin, and sex	Number ²	Percent of total deaths	Mortality rate ²
Infants, all races and origins, both sexes				
...	All causes	19,582	100.0	541.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4,043	20.6	111.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	3,141	16.0	86.9
3	Sudden infant death syndrome (R95)	1,389	7.1	38.4
4	Accidents (unintentional injuries). (V01-X59)	1,194	6.1	33.0
5	Newborn affected by maternal complications of pregnancy. (P01)	1,116	5.7	30.9
6	Newborn affected by complications of placenta, cord and membranes (P02)	700	3.6	19.4
7	Bacterial sepsis of newborn (P36)	542	2.8	15.0
8	Respiratory distress of newborn (P22)	388	2.0	10.7
9	Diseases of the circulatory system. (I00-I99)	386	2.0	10.7
10	Neonatal hemorrhage. (P50-P52,P54)	317	1.6	8.8
...	All other causes (residual)	6,366	32.5	176.2
Infants, all races and origins, male				
...	All causes	10,859	100.0	587.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,140	19.7	115.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,741	16.0	94.2
3	Sudden infant death syndrome (R95)	786	7.2	42.5
4	Accidents (unintentional injuries). (V01-X59)	684	6.3	37.0
5	Newborn affected by maternal complications of pregnancy. (P01)	629	5.8	34.0
6	Newborn affected by complications of placenta, cord and membranes (P02)	380	3.5	20.6
7	Bacterial sepsis of newborn (P36)	313	2.9	16.9
8	Respiratory distress of newborn (P22)	213	2.0	11.5
9	Diseases of the circulatory system. (I00-I99)	192	1.8	10.4
10	Neonatal hemorrhage. (P50-P52,P54)	188	1.7	10.2
...	All other causes (residual)	3,593	33.1	194.4
Infants, all races and origins, female				
...	All causes	8,723	100.0	494.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,903	21.8	107.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,400	16.0	79.3
3	Sudden infant death syndrome (R95)	603	6.9	34.2
4	Accidents (unintentional injuries). (V01-X59)	510	5.8	28.9
5	Newborn affected by maternal complications of pregnancy. (P01)	487	5.6	27.6
6	Newborn affected by complications of placenta, cord and membranes (P02)	320	3.7	18.1
7	Bacterial sepsis of newborn (P36)	229	2.6	13.0
8	Diseases of the circulatory system. (I00-I99)	194	2.2	11.0
9	Respiratory distress of newborn (P22)	175	2.0	9.9
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	134	1.5	7.6
...	All other causes (residual)	2,768	31.7	156.8
Infants, all races, Hispanic, both sexes				
...	All causes	4,238	100.0	489.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,040	24.5	120.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	669	15.8	77.2
3	Newborn affected by maternal complications of pregnancy. (P01)	288	6.8	33.2
4	Sudden infant death syndrome (R95)	239	5.6	27.6
5	Accidents (unintentional injuries). (V01-X59)	162	3.8	18.7
6	Newborn affected by complications of placenta, cord and membranes (P02)	140	3.3	16.2
7	Bacterial sepsis of newborn (P36)	113	2.7	13.0
8	Diseases of the circulatory system. (I00-I99)	82	1.9	9.5
9	Respiratory distress of newborn (P22)	76	1.8	8.8
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	74	1.7	8.5
...	All other causes (residual)	1,355	32.0	156.3

Table I–2. Infant, neonatal, and postneonatal deaths, percentage of total deaths, and mortality rate for the 10 leading causes of infant death, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are infant (younger than 1 year), neonatal (younger than 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic-origin categories are consistent with 1977 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated separately by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black non-Hispanic and White non-Hispanic should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on birth and death certificates. Deaths are based on race and Hispanic origin of decedent; live births are based on race and Hispanic origin of mother. National linked files should be used to measure infant mortality for specific subgroups. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*]

Rank ¹	Cause of death (based on ICD–10), age, race and Hispanic origin, and sex	Number ²	Percent of total deaths	Mortality rate ²
Infants, all races, Hispanic, male				
...	All causes	2,366	100.0	536.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	550	23.2	124.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	367	15.5	83.1
3	Newborn affected by maternal complications of pregnancy (P01)	162	6.8	36.7
4	Sudden infant death syndrome (R95)	147	6.2	33.3
5	Accidents (unintentional injuries) (V01–X59)	105	4.4	23.8
6	Newborn affected by complications of placenta, cord and membranes (P02)	72	3.0	16.3
7	Bacterial sepsis of newborn (P36)	63	2.7	14.3
8	Intrauterine hypoxia and birth asphyxia (P20–P21)	48	2.0	10.9
9	Diseases of the circulatory system (I00–I99)	45	1.9	10.2
9	Neonatal hemorrhage (P50–P52,P54)	45	1.9	10.2
...	All other causes (residual)	762	32.2	172.6
Infants, all races, Hispanic, female				
...	All causes	1,872	100.0	440.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	490	26.2	115.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	302	16.1	71.0
3	Newborn affected by maternal complications of pregnancy (P01)	126	6.7	29.6
4	Sudden infant death syndrome (R95)	92	4.9	21.6
5	Newborn affected by complications of placenta, cord and membranes (P02)	68	3.6	16.0
6	Accidents (unintentional injuries) (V01–X59)	57	3.0	13.4
7	Bacterial sepsis of newborn (P36)	50	2.7	11.8
8	Diseases of the circulatory system (I00–I99)	37	2.0	8.7
9	Respiratory distress of newborn (P22)	35	1.9	8.2
10	Atelectasis (P28.0–P28.1)	33	1.8	7.8
...	All other causes (residual)	582	31.1	136.8
Infants, American Indian and Alaska Native, non-Hispanic, both sexes				
...	All causes	224	100.0	719.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	51	22.8	163.9
2	Accidents (unintentional injuries) (V01–X59)	29	12.9	93.2
2	Sudden infant death syndrome (R95)	29	12.9	93.2
4	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	28	12.5	90.0
5	Newborn affected by maternal complications of pregnancy (P01)	8	3.6	*
5	Necrotizing enterocolitis of newborn (P77)	8	3.6	*
7	Newborn affected by complications of placenta, cord and membranes (P02)	5	2.2	*
8	Influenza and pneumonia (J09–J18)	4	1.8	*
9	Intrauterine hypoxia and birth asphyxia (P20–P21)	3	1.3	*
9	Assault (homicide) (†U01,X85–Y09)	3	1.3	*
...	All other causes (residual)	56	25.0	180.0
Infants, American Indian and Alaska Native, non-Hispanic, male				
...	All causes	130	100.0	813.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	29	22.3	181.5
2	Sudden infant death syndrome (R95)	21	16.2	131.4
3	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	17	13.1	*
4	Accidents (unintentional injuries) (V01–X59)	15	11.5	*
5	Newborn affected by maternal complications of pregnancy (P01)	5	3.8	*
5	Necrotizing enterocolitis of newborn (P77)	5	3.8	*
7	Influenza and pneumonia (J09–J18)	3	2.3	*
7	Newborn affected by complications of placenta, cord and membranes (P02)	3	2.3	*
9	Intrauterine hypoxia and birth asphyxia (P20–P21)	2	1.5	*
9	Assault (homicide) (†U01,X85–Y09)	2	1.5	*
...	All other causes (residual)	28	21.5	175.3

Table I-2. Infant, neonatal, and postneonatal deaths, percentage of total deaths, and mortality rate for the 10 leading causes of infant death, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are infant (younger than 1 year), neonatal (younger than 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic-origin categories are consistent with 1977 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated separately by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black non-Hispanic and White non-Hispanic should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on birth and death certificates. Deaths are based on race and Hispanic origin of decedent; live births are based on race and Hispanic origin of mother. National linked files should be used to measure infant mortality for specific subgroups. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), age, race and Hispanic origin, and sex	Number ²	Percent of total deaths	Mortality rate ²
Infants, American Indian and Alaska Native, non-Hispanic, female				
...	All causes	94	100.0	620.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	23.4	145.3
2	Accidents (unintentional injuries) (V01-X59)	14	14.9	*
3	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	11	11.7	*
4	Sudden infant death syndrome (R95)	8	8.5	*
5	Newborn affected by maternal complications of pregnancy (P01)	3	3.2	*
5	Necrotizing enterocolitis of newborn (P77)	3	3.2	*
7	Newborn affected by complications of placenta, cord and membranes (P02)	2	2.1	*
7	Septicemia (A40-A41)	2	2.1	*
9	Slow fetal growth and fetal malnutrition (P05)	1	1.1	*
9	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	1	1.1	*
9	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	1	1.1	*
9	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	1	1.1	*
9	Assault (homicide) (†U01,X85-Y09)	1	1.1	*
9	Newborn affected by maternal hypertensive disorders (P00.0)	1	1.1	*
9	Malignant neoplasms (C00-C97)	1	1.1	*
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	1	1.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.1	*
9	Influenza and pneumonia (J09-J18)	1	1.1	*
9	Chronic respiratory disease originating in the perinatal period (P27)	1	1.1	*
9	Syndrome of infant of a diabetic mother and neonatal diabetes mellitus (P70.0-P70.2)	1	1.1	*
9	Hydrops fetalis not due to hemolytic disease (P83.2)	1	1.1	*
9	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	1	1.1	*
...	All other causes (residual)	15	16.0	*
Infants, Asian and Other Pacific Islander, non-Hispanic, both sexes				
...	All causes	655	100.0	271.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	170	26.0	70.4
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	103	15.7	42.7
3	Newborn affected by maternal complications of pregnancy (P01)	44	6.7	18.2
4	Newborn affected by complications of placenta, cord and membranes (P02)	33	5.0	13.7
5	Sudden infant death syndrome (R95)	24	3.7	9.9
6	Bacterial sepsis of newborn (P36)	21	3.2	8.7
7	Accidents (unintentional injuries) (V01-X59)	16	2.4	*
8	Neonatal hemorrhage (P50-P52,P54)	13	2.0	*
8	Respiratory distress of newborn (P22)	13	2.0	*
10	Diarrhea and gastroenteritis of infectious origin (A09)	12	1.8	*
...	All other causes (residual)	206	31.5	85.3
Infants, Asian and Other Pacific Islander, non-Hispanic, male				
...	All causes	370	100.0	297.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	21.9	65.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	62	16.8	49.9
3	Newborn affected by maternal complications of pregnancy (P01)	33	8.9	26.5
4	Newborn affected by complications of placenta, cord and membranes (P02)	18	4.9	*
5	Bacterial sepsis of newborn (P36)	14	3.8	*
6	Accidents (unintentional injuries) (V01-X59)	10	2.7	*
6	Sudden infant death syndrome (R95)	10	2.7	*
8	Diarrhea and gastroenteritis of infectious origin (A09)	9	2.4	*
9	Respiratory distress of newborn (P22)	8	2.2	*
9	Neonatal hemorrhage (P50-P52,P54)	8	2.2	*
...	All other causes (residual)	117	31.6	94.1

Table 1–2. Infant, neonatal, and postneonatal deaths, percentage of total deaths, and mortality rate for the 10 leading causes of infant death, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are infant (younger than 1 year), neonatal (younger than 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic-origin categories are consistent with 1977 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated separately by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black non-Hispanic and White non-Hispanic should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on birth and death certificates. Deaths are based on race and Hispanic origin of decedent; live births are based on race and Hispanic origin of mother. National linked files should be used to measure infant mortality for specific subgroups. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*]

Rank ¹	Cause of death (based on ICD–10), age, race and Hispanic origin, and sex	Number ²	Percent of total deaths	Mortality rate ²
Infants, Asian and Other Pacific Islander, non-Hispanic, female				
...	All causes	285	100.0	243.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	89	31.2	76.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	41	14.4	35.0
3	Newborn affected by complications of placenta, cord and membranes (P02)	15	5.3	*
4	Sudden infant death syndrome (R95)	14	4.9	*
5	Newborn affected by maternal complications of pregnancy (P01)	11	3.9	*
6	Bacterial sepsis of newborn (P36)	7	2.5	*
7	Accidents (unintentional injuries) (V01–X59)	6	2.1	*
8	Neonatal hemorrhage (P50–P52, P54)	5	1.8	*
8	Intrauterine hypoxia and birth asphyxia (P20–P21)	5	1.8	*
8	Respiratory distress of newborn (P22)	5	1.8	*
...	All other causes (residual)	87	30.5	74.3
Infants, Black, non-Hispanic, both sexes				
...	All causes	6,014	100.0	1,076.0
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,190	19.8	212.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	765	12.7	136.9
3	Sudden infant death syndrome (R95)	544	9.0	97.3
4	Accidents (unintentional injuries) (V01–X59)	437	7.3	78.2
5	Newborn affected by maternal complications of pregnancy (P01)	357	5.9	63.9
6	Newborn affected by complications of placenta, cord and membranes (P02)	217	3.6	38.8
7	Bacterial sepsis of newborn (P36)	199	3.3	35.6
8	Respiratory distress of newborn (P22)	125	2.1	22.4
9	Diseases of the circulatory system (I00–I99)	109	1.8	19.5
10	Necrotizing enterocolitis of newborn (P77)	105	1.7	18.8
...	All other causes (residual)	1,966	32.7	351.7
Infants, Black, non-Hispanic, male				
...	All causes	3,314	100.0	1,166.9
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	656	19.8	231.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	396	11.9	139.4
3	Sudden infant death syndrome (R95)	285	8.6	100.4
4	Accidents (unintentional injuries) (V01–X59)	246	7.4	86.6
5	Newborn affected by maternal complications of pregnancy (P01)	202	6.1	71.1
6	Newborn affected by complications of placenta, cord and membranes (P02)	129	3.9	45.4
7	Bacterial sepsis of newborn (P36)	109	3.3	38.4
8	Assault (homicide) (†TU01, X85–Y09)	65	2.0	22.9
9	Respiratory distress of newborn (P22)	64	1.9	22.5
10	Necrotizing enterocolitis of newborn (P77)	59	1.8	20.8
...	All other causes (residual)	1,103	33.3	388.4

Table 1–2. Infant, neonatal, and postneonatal deaths, percentage of total deaths, and mortality rate for the 10 leading causes of infant death, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are infant (younger than 1 year), neonatal (younger than 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic-origin categories are consistent with 1977 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated separately by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black non-Hispanic and White non-Hispanic should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on birth and death certificates. Deaths are based on race and Hispanic origin of decedent; live births are based on race and Hispanic origin of mother. National linked files should be used to measure infant mortality for specific subgroups. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*]

Rank ¹	Cause of death (based on ICD–10), age, race and Hispanic origin, and sex	Number ²	Percent of total deaths	Mortality rate ²
Infants, Black, non-Hispanic, female				
...	All causes	2,700	100.0	982.0
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	534	19.8	194.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	369	13.7	134.2
3	Sudden infant death syndrome (R95)	259	9.6	94.2
4	Accidents (unintentional injuries) (V01–X59)	191	7.1	69.5
5	Newborn affected by maternal complications of pregnancy (P01)	155	5.7	56.4
6	Bacterial sepsis of newborn (P36)	90	3.3	32.7
7	Newborn affected by complications of placenta, cord and membranes (P02)	88	3.3	32.0
8	Respiratory distress of newborn (P22)	61	2.3	22.2
9	Diseases of the circulatory system (I00–I99)	58	2.1	21.1
10	Intrauterine hypoxia and birth asphyxia (P20–P21)	46	1.7	16.7
10	Necrotizing enterocolitis of newborn (P77)	46	1.7	16.7
...	All other causes (residual)	803	29.7	292.1
Infants, White, non-Hispanic, both sexes				
...	All causes	8,250	100.0	438.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	1,997	24.2	106.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,090	13.2	57.9
3	Accidents (unintentional injuries) (V01–X59)	547	6.6	29.1
3	Sudden infant death syndrome (R95)	547	6.6	29.1
5	Newborn affected by maternal complications of pregnancy (P01)	397	4.8	21.1
6	Newborn affected by complications of placenta, cord and membranes (P02)	294	3.6	15.6
7	Bacterial sepsis of newborn (P36)	206	2.5	10.9
8	Diseases of the circulatory system (I00–I99)	185	2.2	9.8
9	Respiratory distress of newborn (P22)	167	2.0	8.9
10	Neonatal hemorrhage (P50–P52,P54)	157	1.9	8.3
...	All other causes (residual)	2,663	32.3	141.5
Infants, White, non-Hispanic, male				
...	All causes	4,567	100.0	473.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	1,072	23.5	111.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	601	13.2	62.3
3	Sudden infant death syndrome (R95)	320	7.0	33.2
4	Accidents (unintentional injuries) (V01–X59)	307	6.7	31.8
5	Newborn affected by maternal complications of pregnancy (P01)	215	4.7	22.3
6	Newborn affected by complications of placenta, cord and membranes (P02)	155	3.4	16.1
7	Bacterial sepsis of newborn (P36)	125	2.7	13.0
8	Respiratory distress of newborn (P22)	95	2.1	9.8
9	Neonatal hemorrhage (P50–P52,P54)	93	2.0	9.6
10	Diseases of the circulatory system (I00–I99)	91	2.0	9.4
...	All other causes (residual)	1,493	32.7	154.7

Table 1–2. Infant, neonatal, and postneonatal deaths, percentage of total deaths, and mortality rate for the 10 leading causes of infant death, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are infant (younger than 1 year), neonatal (younger than 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic-origin categories are consistent with 1977 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated separately by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black non-Hispanic and White non-Hispanic should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on birth and death certificates. Deaths are based on race and Hispanic origin of decedent; live births are based on race and Hispanic origin of mother. National linked files should be used to measure infant mortality for specific subgroups. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*]

Rank ¹	Cause of death (based on ICD–10), age, race and Hispanic origin, and sex	Number ²	Percent of total deaths	Mortality rate ²
Infants, White, non-Hispanic, female				
...	All causes	3,683	100.0	401.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	925	25.1	100.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	489	13.3	53.3
3	Accidents (unintentional injuries) (V01–X59)	240	6.5	26.2
4	Sudden infant death syndrome (R95)	227	6.2	24.8
5	Newborn affected by maternal complications of pregnancy (P01)	182	4.9	19.9
6	Newborn affected by complications of placenta, cord and membranes (P02)	139	3.8	15.2
7	Diseases of the circulatory system (I00–I99)	94	2.6	10.3
8	Bacterial sepsis of newborn (P36)	81	2.2	8.8
9	Respiratory distress of newborn (P22)	72	2.0	7.9
10	Neonatal hemorrhage (P50–P52,P54)	64	1.7	7.0
...	All other causes (residual)	1,170	31.8	127.6
Neonates, all races and origins, both sexes				
...	All causes	12,856	100.0	355.8
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	3,069	23.9	84.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	2,867	22.3	79.3
3	Newborn affected by maternal complications of pregnancy (P01)	1,102	8.6	30.5
4	Newborn affected by complications of placenta, cord and membranes (P02)	687	5.3	19.0
5	Bacterial sepsis of newborn (P36)	521	4.1	14.4
6	Respiratory distress of newborn (P22)	384	3.0	10.6
7	Neonatal hemorrhage (P50–P52,P54)	310	2.4	8.6
8	Intrauterine hypoxia and birth asphyxia (P20–P21)	296	2.3	8.2
9	Necrotizing enterocolitis of newborn (P77)	263	2.0	7.3
10	Atelectasis (P28.0–P28.1)	231	1.8	6.4
...	All other causes (residual)	3,126	24.3	86.5
Neonates, all races and origins, male				
...	All causes	7,098	100.0	384.1
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,701	24.0	92.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	1,561	22.0	84.5
3	Newborn affected by maternal complications of pregnancy (P01)	621	8.7	33.6
4	Newborn affected by complications of placenta, cord and membranes (P02)	370	5.2	20.0
5	Bacterial sepsis of newborn (P36)	300	4.2	16.2
6	Respiratory distress of newborn (P22)	211	3.0	11.4
7	Neonatal hemorrhage (P50–P52,P54)	184	2.6	10.0
8	Intrauterine hypoxia and birth asphyxia (P20–P21)	167	2.4	9.0
9	Necrotizing enterocolitis of newborn (P77)	153	2.2	8.3
10	Atelectasis (P28.0–P28.1)	133	1.9	7.2
...	All other causes (residual)	1,697	23.9	91.8
Neonates, all races and origins, female				
...	All causes	5,758	100.0	326.1
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,368	23.8	77.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	1,306	22.7	74.0
3	Newborn affected by maternal complications of pregnancy (P01)	481	8.4	27.2
4	Newborn affected by complications of placenta, cord and membranes (P02)	317	5.5	18.0
5	Bacterial sepsis of newborn (P36)	221	3.8	12.5
6	Respiratory distress of newborn (P22)	173	3.0	9.8
7	Intrauterine hypoxia and birth asphyxia (P20–P21)	129	2.2	7.3
8	Neonatal hemorrhage (P50–P52,P54)	126	2.2	7.1
9	Necrotizing enterocolitis of newborn (P77)	110	1.9	6.2
10	Atelectasis (P28.0–P28.1)	98	1.7	5.6
...	All other causes (residual)	1,429	24.8	80.9

Table 1–2. Infant, neonatal, and postneonatal deaths, percentage of total deaths, and mortality rate for the 10 leading causes of infant death, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are infant (younger than 1 year), neonatal (younger than 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic-origin categories are consistent with 1977 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated separately by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black non-Hispanic and White non-Hispanic should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on birth and death certificates. Deaths are based on race and Hispanic origin of decedent; live births are based on race and Hispanic origin of mother. National linked files should be used to measure infant mortality for specific subgroups. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*]

Rank ¹	Cause of death (based on ICD–10), age, race and Hispanic origin, and sex	Number ²	Percent of total deaths	Mortality rate ²
Neonates, all races, Hispanic, both sexes				
...	All causes	2,930	100.0	338.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	750	25.6	86.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	654	22.3	75.5
3	Newborn affected by maternal complications of pregnancy (P01)	286	9.8	33.0
4	Newborn affected by complications of placenta, cord and membranes (P02)	138	4.7	15.9
5	Bacterial sepsis of newborn (P36)	103	3.5	11.9
6	Respiratory distress of newborn (P22)	75	2.6	8.7
7	Neonatal hemorrhage (P50–P52,P54)	72	2.5	8.3
8	Intrauterine hypoxia and birth asphyxia (P20–P21)	71	2.4	8.2
9	Atelectasis (P28.0–P28.1)	62	2.1	7.2
10	Necrotizing enterocolitis of newborn (P77)	52	1.8	6.0
...	All other causes (residual)	667	22.8	77.0
Neonates, all races, Hispanic, male				
...	All causes	1,634	100.0	370.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	406	24.8	92.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	361	22.1	81.8
3	Newborn affected by maternal complications of pregnancy (P01)	160	9.8	36.2
4	Newborn affected by complications of placenta, cord and membranes (P02)	71	4.3	16.1
5	Bacterial sepsis of newborn (P36)	55	3.4	12.5
6	Intrauterine hypoxia and birth asphyxia (P20–P21)	48	2.9	10.9
7	Neonatal hemorrhage (P50–P52,P54)	44	2.7	10.0
8	Respiratory distress of newborn (P22)	40	2.4	9.1
9	Atelectasis (P28.0–P28.1)	35	2.1	7.9
10	Necrotizing enterocolitis of newborn (P77)	32	2.0	7.2
...	All other causes (residual)	382	23.4	86.5
Neonates, all races, Hispanic, female				
...	All causes	1,296	100.0	304.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	344	26.5	80.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	293	22.6	68.9
3	Newborn affected by maternal complications of pregnancy (P01)	126	9.7	29.6
4	Newborn affected by complications of placenta, cord and membranes (P02)	67	5.2	15.8
5	Bacterial sepsis of newborn (P36)	48	3.7	11.3
6	Respiratory distress of newborn (P22)	35	2.7	8.2
7	Neonatal hemorrhage (P50–P52,P54)	28	2.2	6.6
8	Atelectasis (P28.0–P28.1)	27	2.1	6.3
9	Intrauterine hypoxia and birth asphyxia (P20–P21)	23	1.8	5.4
10	Necrotizing enterocolitis of newborn (P77)	20	1.5	4.7
...	All other causes (residual)	285	22.0	67.0

Table 1–2. Infant, neonatal, and postneonatal deaths, percentage of total deaths, and mortality rate for the 10 leading causes of infant death, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are infant (younger than 1 year), neonatal (younger than 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic-origin categories are consistent with 1977 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated separately by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black non-Hispanic and White non-Hispanic should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on birth and death certificates. Deaths are based on race and Hispanic origin of decedent; live births are based on race and Hispanic origin of mother. National linked files should be used to measure infant mortality for specific subgroups. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*]

Rank ¹	Cause of death (based on ICD–10), age, race and Hispanic origin, and sex	Number ²	Percent of total deaths	Mortality rate ²
Neonates, American Indian and Alaska Native, non-Hispanic, both sexes				
...	All causes	113	100.0	363.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	32	28.3	102.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	27	23.9	86.8
3	Newborn affected by maternal complications of pregnancy (P01)	8	7.1	*
4	Necrotizing enterocolitis of newborn (P77)	7	6.2	*
5	Newborn affected by complications of placenta, cord and membranes (P02)	5	4.4	*
6	Sudden infant death syndrome (R95)	4	3.5	*
7	Intrauterine hypoxia and birth asphyxia (P20–P21)	3	2.7	*
8	Newborn affected by maternal hypertensive disorders (P00.0)	2	1.8	*
9	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1–P00.9)	1	0.9	*
9	Pulmonary hemorrhage originating in the perinatal period (P26)	1	0.9	*
9	Congenital pneumonia (P23)	1	0.9	*
9	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50–D89)	1	0.9	*
9	Slow fetal growth and fetal malnutrition (P05)	1	0.9	*
9	Respiratory distress of newborn (P22)	1	0.9	*
9	Chronic respiratory disease originating in the perinatal period (P27)	1	0.9	*
9	Accidents (unintentional injuries) (V01–X59)	1	0.9	*
9	Hydrops fetalis not due to hemolytic disease (P83.2)	1	0.9	*
9	Syndrome of infant of a diabetic mother and neonatal diabetes mellitus (P70.0–P70.2)	1	0.9	*
...	All other causes (residual)	15	13.3	*
Neonates, American Indian and Alaska Native, non-Hispanic, male				
...	All causes	68	100.0	425.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	21	30.9	131.4
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	17	25.0	*
3	Newborn affected by maternal complications of pregnancy (P01)	5	7.4	*
4	Necrotizing enterocolitis of newborn (P77)	4	5.9	*
5	Newborn affected by complications of placenta, cord and membranes (P02)	3	4.4	*
5	Sudden infant death syndrome (R95)	3	4.4	*
7	Intrauterine hypoxia and birth asphyxia (P20–P21)	2	2.9	*
8	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50–D89)	1	1.5	*
8	Newborn affected by maternal hypertensive disorders (P00.0)	1	1.5	*
8	Respiratory distress of newborn (P22)	1	1.5	*
8	Congenital pneumonia (P23)	1	1.5	*
8	Pulmonary hemorrhage originating in the perinatal period (P26)	1	1.5	*
...	All other causes (residual)	8	11.8	*

Table I–2. Infant, neonatal, and postneonatal deaths, percentage of total deaths, and mortality rate for the 10 leading causes of infant death, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are infant (younger than 1 year), neonatal (younger than 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic-origin categories are consistent with 1977 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated separately by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black non-Hispanic and White non-Hispanic should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on birth and death certificates. Deaths are based on race and Hispanic origin of decedent; live births are based on race and Hispanic origin of mother. National linked files should be used to measure infant mortality for specific subgroups. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*]

Rank ¹	Cause of death (based on ICD–10), age, race and Hispanic origin, and sex	Number ²	Percent of total deaths	Mortality rate ²
Neonates, American Indian and Alaska Native, non-Hispanic, female				
...	All causes	45	100.0	297.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	11	24.4	*
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	10	22.2	*
3	Necrotizing enterocolitis of newborn (P77)	3	6.7	*
3	Newborn affected by maternal complications of pregnancy (P01)	3	6.7	*
5	Newborn affected by complications of placenta, cord and membranes (P02)	2	4.4	*
6	Slow fetal growth and fetal malnutrition (P05)	1	2.2	*
6	Newborn affected by maternal hypertensive disorders (P00.0)	1	2.2	*
6	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1–P00.9)	1	2.2	*
6	Intrauterine hypoxia and birth asphyxia (P20–P21)	1	2.2	*
6	Chronic respiratory disease originating in the perinatal period (P27)	1	2.2	*
6	Syndrome of infant of a diabetic mother and neonatal diabetes mellitus (P70.0–P70.2)	1	2.2	*
6	Hydrops fetalis not due to hemolytic disease (P83.2)	1	2.2	*
6	Sudden infant death syndrome (R95)	1	2.2	*
6	Accidents (unintentional injuries) (V01–X59)	1	2.2	*
...	All other causes (residual)	7	15.6	*
Neonates, Asian and Other Pacific Islander, non-Hispanic, both sexes				
...	All causes	464	100.0	192.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	116	25.0	48.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	101	21.8	41.8
3	Newborn affected by maternal complications of pregnancy (P01)	43	9.3	17.8
4	Newborn affected by complications of placenta, cord and membranes (P02)	33	7.1	13.7
5	Bacterial sepsis of newborn (P36)	20	4.3	8.3
6	Neonatal hemorrhage (P50–P52,P54)	13	2.8	*
7	Respiratory distress of newborn (P22)	12	2.6	*
8	Intrauterine hypoxia and birth asphyxia (P20–P21)	10	2.2	*
9	Hydrops fetalis not due to hemolytic disease (P83.2)	7	1.5	*
10	Necrotizing enterocolitis of newborn (P77)	6	1.3	*
...	All other causes (residual)	103	22.2	42.7
Neonates, Asian and Other Pacific Islander, non-Hispanic, male				
...	All causes	259	100.0	208.3
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	60	23.2	48.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	54	20.8	43.4
3	Newborn affected by maternal complications of pregnancy (P01)	32	12.4	25.7
4	Newborn affected by complications of placenta, cord and membranes (P02)	18	6.9	*
5	Bacterial sepsis of newborn (P36)	13	5.0	*
6	Respiratory distress of newborn (P22)	8	3.1	*
6	Neonatal hemorrhage (P50–P52,P54)	8	3.1	*
8	Intrauterine hypoxia and birth asphyxia (P20–P21)	5	1.9	*
9	Hydrops fetalis not due to hemolytic disease (P83.2)	4	1.5	*
9	Atelectasis (P28.0–P28.1)	4	1.5	*
...	All other causes (residual)	53	20.5	42.6

Table 1–2. Infant, neonatal, and postneonatal deaths, percentage of total deaths, and mortality rate for the 10 leading causes of infant death, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are infant (younger than 1 year), neonatal (younger than 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic-origin categories are consistent with 1977 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated separately by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black non-Hispanic and White non-Hispanic should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on birth and death certificates. Deaths are based on race and Hispanic origin of decedent; live births are based on race and Hispanic origin of mother. National linked files should be used to measure infant mortality for specific subgroups. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*]

Rank ¹	Cause of death (based on ICD–10), age, race and Hispanic origin, and sex	Number ²	Percent of total deaths	Mortality rate ²
Neonates, Asian and Other Pacific Islander, non-Hispanic, female				
...	All causes	205	100.0	175.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	62	30.2	52.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	41	20.0	35.0
3	Newborn affected by complications of placenta, cord and membranes (P02)	15	7.3	*
4	Newborn affected by maternal complications of pregnancy (P01)	11	5.4	*
5	Bacterial sepsis of newborn (P36)	7	3.4	*
6	Intrauterine hypoxia and birth asphyxia (P20–P21)	5	2.4	*
6	Neonatal hemorrhage (P50–P52,P54)	5	2.4	*
8	Respiratory distress of newborn (P22)	4	2.0	*
9	Necrotizing enterocolitis of newborn (P77)	3	1.5	*
9	Hydrops fetalis not due to hemolytic disease (P83.2)	3	1.5	*
...	All other causes (residual)	49	23.9	41.8
Neonates, Black, non-Hispanic, both sexes				
...	All causes	3,724	100.0	666.3
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,159	31.1	207.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	502	13.5	89.8
3	Newborn affected by maternal complications of pregnancy (P01)	355	9.5	63.5
4	Newborn affected by complications of placenta, cord and membranes (P02)	212	5.7	37.9
5	Bacterial sepsis of newborn (P36)	193	5.2	34.5
6	Respiratory distress of newborn (P22)	124	3.3	22.2
7	Intrauterine hypoxia and birth asphyxia (P20–P21)	95	2.6	17.0
8	Necrotizing enterocolitis of newborn (P77)	89	2.4	15.9
9	Neonatal hemorrhage (P50–P52,P54)	66	1.8	11.8
10	Atelectasis (P28.0–P28.1)	58	1.6	10.4
...	All other causes (residual)	871	23.4	155.8
Neonates, Black, non-Hispanic, male				
...	All causes	2,048	100.0	721.1
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	639	31.2	225.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	269	13.1	94.7
3	Newborn affected by maternal complications of pregnancy (P01)	202	9.9	71.1
4	Newborn affected by complications of placenta, cord and membranes (P02)	125	6.1	44.0
5	Bacterial sepsis of newborn (P36)	106	5.2	37.3
6	Respiratory distress of newborn (P22)	63	3.1	22.2
7	Necrotizing enterocolitis of newborn (P77)	52	2.5	18.3
8	Intrauterine hypoxia and birth asphyxia (P20–P21)	49	2.4	17.3
9	Neonatal hemorrhage (P50–P52,P54)	37	1.8	13.0
10	Atelectasis (P28.0–P28.1)	28	1.4	9.9
...	All other causes (residual)	478	23.3	168.3
Neonates, Black, non-Hispanic, female				
...	All causes	1,676	100.0	609.6
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	520	31.0	189.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	233	13.9	84.7
3	Newborn affected by maternal complications of pregnancy (P01)	153	9.1	55.6
4	Bacterial sepsis of newborn (P36)	87	5.2	31.6
4	Newborn affected by complications of placenta, cord and membranes (P02)	87	5.2	31.6
6	Respiratory distress of newborn (P22)	61	3.6	22.2
7	Intrauterine hypoxia and birth asphyxia (P20–P21)	46	2.7	16.7
8	Necrotizing enterocolitis of newborn (P77)	37	2.2	13.5
9	Atelectasis (P28.0–P28.1)	30	1.8	10.9
10	Neonatal hemorrhage (P50–P52,P54)	29	1.7	10.5
...	All other causes (residual)	393	23.4	142.9

Table 1–2. Infant, neonatal, and postneonatal deaths, percentage of total deaths, and mortality rate for the 10 leading causes of infant death, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are infant (younger than 1 year), neonatal (younger than 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic-origin categories are consistent with 1977 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated separately by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black non-Hispanic and White non-Hispanic should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on birth and death certificates. Deaths are based on race and Hispanic origin of decedent; live births are based on race and Hispanic origin of mother. National linked files should be used to measure infant mortality for specific subgroups. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*]

Rank ¹	Cause of death (based on ICD–10), age, race and Hispanic origin, and sex	Number ²	Percent of total deaths	Mortality rate ²
Neonates, White, non-Hispanic, both sexes				
...	All causes	5,456	100.0	290.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	1,454	26.6	77.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,068	19.6	56.8
3	Newborn affected by maternal complications of pregnancy (P01)	388	7.1	20.6
4	Newborn affected by complications of placenta, cord and membranes (P02)	288	5.3	15.3
5	Bacterial sepsis of newborn (P36)	202	3.7	10.7
6	Respiratory distress of newborn (P22)	166	3.0	8.8
7	Neonatal hemorrhage (P50–P52,P54)	154	2.8	8.2
8	Intrauterine hypoxia and birth asphyxia (P20–P21)	115	2.1	6.1
9	Necrotizing enterocolitis of newborn (P77)	107	2.0	5.7
10	Hydrops fetalis not due to hemolytic disease (P83.2)	103	1.9	5.5
...	All other causes (residual)	1,411	25.9	75.0
Neonates, White, non-Hispanic, male				
...	All causes	2,995	100.0	310.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	804	26.8	83.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	586	19.6	60.7
3	Newborn affected by maternal complications of pregnancy (P01)	210	7.0	21.8
4	Newborn affected by complications of placenta, cord and membranes (P02)	150	5.0	15.5
5	Bacterial sepsis of newborn (P36)	124	4.1	12.9
6	Respiratory distress of newborn (P22)	95	3.2	9.8
7	Neonatal hemorrhage (P50–P52,P54)	92	3.1	9.5
8	Atelectasis (P28.0–P28.1)	64	2.1	6.6
9	Intrauterine hypoxia and birth asphyxia (P20–P21)	63	2.1	6.5
10	Necrotizing enterocolitis of newborn (P77)	61	2.0	6.3
...	All other causes (residual)	746	24.9	77.3
Neonates, White, non-Hispanic, female				
...	All causes	2,461	100.0	268.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	650	26.4	70.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	482	19.6	52.6
3	Newborn affected by maternal complications of pregnancy (P01)	178	7.2	19.4
4	Newborn affected by complications of placenta, cord and membranes (P02)	138	5.6	15.1
5	Bacterial sepsis of newborn (P36)	78	3.2	8.5
6	Respiratory distress of newborn (P22)	71	2.9	7.7
7	Neonatal hemorrhage (P50–P52,P54)	62	2.5	6.8
8	Intrauterine hypoxia and birth asphyxia (P20–P21)	52	2.1	5.7
9	Hydrops fetalis not due to hemolytic disease (P83.2)	49	2.0	5.3
10	Necrotizing enterocolitis of newborn (P77)	46	1.9	5.0
...	All other causes (residual)	655	26.6	71.5
Postneonates, all races and origins, both sexes				
...	All causes	6,726	100.0	186.1
1	Sudden infant death syndrome (R95)	1,247	18.5	34.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	1,176	17.5	32.5
3	Accidents (unintentional injuries) (V01–X59)	1,077	16.0	29.8
4	Diseases of the circulatory system (I00–I99)	306	4.5	8.5
5	Assault (homicide) (†U01,X85–Y09)	230	3.4	6.4
6	Diarrhea and gastroenteritis of infectious origin (A09)	134	2.0	3.7
7	Influenza and pneumonia (J09–J18)	124	1.8	3.4
8	Chronic respiratory disease originating in the perinatal period (P27)	120	1.8	3.3
9	Septicemia (A40–A41)	114	1.7	3.2
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	72	1.1	2.0
...	All other causes (residual)	2,126	31.6	58.8

Table I-2. Infant, neonatal, and postneonatal deaths, percentage of total deaths, and mortality rate for the 10 leading causes of infant death, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are infant (younger than 1 year), neonatal (younger than 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic-origin categories are consistent with 1977 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated separately by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black non-Hispanic and White non-Hispanic should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on birth and death certificates. Deaths are based on race and Hispanic origin of decedent; live births are based on race and Hispanic origin of mother. National linked files should be used to measure infant mortality for specific subgroups. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*]

Rank ¹	Cause of death (based on ICD-10), age, race and Hispanic origin, and sex	Number ²	Percent of total deaths	Mortality rate ²
Postneonates, all races and origins, male				
...	All causes	3,761	100.0	203.5
1	Sudden infant death syndrome (R95)	710	18.9	38.4
2	Accidents (unintentional injuries) (V01-X59)	623	16.6	33.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	579	15.4	31.3
4	Diseases of the circulatory system (I00-I99)	158	4.2	8.5
5	Assault (homicide) (†U01,X85-Y09)	144	3.8	7.8
6	Diarrhea and gastroenteritis of infectious origin (A09)	84	2.2	4.5
7	Influenza and pneumonia (J09-J18)	81	2.2	4.4
8	Chronic respiratory disease originating in the perinatal period (P27)	73	1.9	4.0
9	Septicemia (A40-A41)	63	1.7	3.4
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	40	1.1	2.2
...	All other causes (residual)	1,206	32.1	65.3
Postneonates, all races and origins, female				
...	All causes	2,965	100.0	167.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	597	20.1	33.8
2	Sudden infant death syndrome (R95)	537	18.1	30.4
3	Accidents (unintentional injuries) (V01-X59)	454	15.3	25.7
4	Diseases of the circulatory system (I00-I99)	148	5.0	8.4
5	Assault (homicide) (†U01,X85-Y09)	86	2.9	4.9
6	Septicemia (A40-A41)	51	1.7	2.9
7	Diarrhea and gastroenteritis of infectious origin (A09)	50	1.7	2.8
8	Chronic respiratory disease originating in the perinatal period (P27)	47	1.6	2.7
9	Influenza and pneumonia (J09-J18)	43	1.5	2.4
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	32	1.1	1.8
...	All other causes (residual)	920	31.0	52.1
Postneonates, all races, Hispanic, both sexes				
...	All causes	1,308	100.0	150.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	290	22.2	33.5
2	Sudden infant death syndrome (R95)	214	16.4	24.7
3	Accidents (unintentional injuries) (V01-X59)	148	11.3	17.1
4	Diseases of the circulatory system (I00-I99)	67	5.1	7.7
5	Assault (homicide) (†U01,X85-Y09)	37	2.8	4.3
6	Septicemia (A40-A41)	34	2.6	3.9
7	Diarrhea and gastroenteritis of infectious origin (A09)	28	2.1	3.2
8	Chronic respiratory disease originating in the perinatal period (P27)	21	1.6	2.4
9	Influenza and pneumonia (J09-J18)	17	1.3	*
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	15	1.1	*
...	All other causes (residual)	437	33.4	50.4
Postneonates, all races, Hispanic, male				
...	All causes	732	100.0	165.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	144	19.7	32.6
2	Sudden infant death syndrome (R95)	130	17.8	29.5
3	Accidents (unintentional injuries) (V01-X59)	96	13.1	21.7
4	Diseases of the circulatory system (I00-I99)	36	4.9	8.2
5	Diarrhea and gastroenteritis of infectious origin (A09)	18	2.5	*
5	Assault (homicide) (†U01,X85-Y09)	18	2.5	*
7	Septicemia (A40-A41)	17	2.3	*
8	Chronic respiratory disease originating in the perinatal period (P27)	14	1.9	*
9	Influenza and pneumonia (J09-J18)	10	1.4	*
10	Bacterial sepsis of newborn (P36)	8	1.1	*
...	All other causes (residual)	241	32.9	54.6

Table I–2. Infant, neonatal, and postneonatal deaths, percentage of total deaths, and mortality rate for the 10 leading causes of infant death, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are infant (younger than 1 year), neonatal (younger than 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic-origin categories are consistent with 1977 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated separately by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black non-Hispanic and White non-Hispanic should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on birth and death certificates. Deaths are based on race and Hispanic origin of decedent; live births are based on race and Hispanic origin of mother. National linked files should be used to measure infant mortality for specific subgroups. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*]

Rank ¹	Cause of death (based on ICD–10), age, race and Hispanic origin, and sex	Number ²	Percent of total deaths	Mortality rate ²
Postneonates, all races, Hispanic, female				
...	All causes	576	100.0	135.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	146	25.3	34.3
2	Sudden infant death syndrome (R95)	84	14.6	19.8
3	Accidents (unintentional injuries) (V01–X59)	52	9.0	12.2
4	Diseases of the circulatory system (I00–I99)	31	5.4	7.3
5	Assault (homicide) (†U01.X85–Y09)	19	3.3	*
6	Septicemia (A40–A41)	17	3.0	*
7	Diarrhea and gastroenteritis of infectious origin (A09)	10	1.7	*
8	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	9	1.6	*
9	Malignant neoplasms (C00–C97)	7	1.2	*
9	Influenza and pneumonia (J09–J18)	7	1.2	*
9	Chronic respiratory disease originating in the perinatal period (P27)	7	1.2	*
...	All other causes (residual)	187	32.5	44.0
Postneonates, American Indian and Alaska Native, non-Hispanic, both sexes				
...	All causes	111	100.0	356.7
1	Accidents (unintentional injuries) (V01–X59)	28	25.2	90.0
2	Sudden infant death syndrome (R95)	25	22.5	80.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	19	17.1	*
4	Influenza and pneumonia (J09–J18)	4	3.6	*
5	Assault (homicide) (†U01.X85–Y09)	3	2.7	*
6	Septicemia (A40–A41)	2	1.8	*
6	Malignant neoplasms (C00–C97)	2	1.8	*
8	Diseases of the circulatory system (I00–I99)	1	0.9	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00–D48)	1	0.9	*
8	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50–D89)	1	0.9	*
8	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86–E87)	1	0.9	*
8	Bronchitis, chronic and unspecified (J40–J42)	1	0.9	*
8	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50–K55)	1	0.9	*
8	Hernia of abdominal cavity and intestinal obstruction without hernia (K40–K46,K56)	1	0.9	*
8	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1	0.9	*
8	Necrotizing enterocolitis of newborn (P77)	1	0.9	*
...	All other causes (residual)	19	17.1	*
Postneonates, American Indian and Alaska Native, non-Hispanic, male				
...	All causes	62	100.0	388.1
1	Sudden infant death syndrome (R95)	18	29.0	*
2	Accidents (unintentional injuries) (V01–X59)	15	24.2	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	8	12.9	*
4	Influenza and pneumonia (J09–J18)	3	4.8	*
5	Assault (homicide) (†U01.X85–Y09)	2	3.2	*
6	Diseases of the circulatory system (I00–I99)	1	1.6	*
6	Malignant neoplasms (C00–C97)	1	1.6	*
6	Bronchitis, chronic and unspecified (J40–J42)	1	1.6	*
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50–K55)	1	1.6	*
6	Necrotizing enterocolitis of newborn (P77)	1	1.6	*
...	All other causes (residual)	11	17.7	*

Table 1–2. Infant, neonatal, and postneonatal deaths, percentage of total deaths, and mortality rate for the 10 leading causes of infant death, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are infant (younger than 1 year), neonatal (younger than 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic-origin categories are consistent with 1977 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated separately by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black non-Hispanic and White non-Hispanic should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on birth and death certificates. Deaths are based on race and Hispanic origin of decedent; live births are based on race and Hispanic origin of mother. National linked files should be used to measure infant mortality for specific subgroups. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*]

Rank ¹	Cause of death (based on ICD–10), age, race and Hispanic origin, and sex	Number ²	Percent of total deaths	Mortality rate ²
Postneonates, American Indian and Alaska Native, non-Hispanic, female				
...	All causes	49	100.0	323.6
1	Accidents (unintentional injuries) (V01–X59)	13	26.5	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	11	22.4	*
3	Sudden infant death syndrome (R95)	7	14.3	*
4	Septicemia (A40–A41)	2	4.1	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00–D48)	1	2.0	*
5	Malignant neoplasms (C00–C97)	1	2.0	*
5	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50–D89)	1	2.0	*
5	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86–E87)	1	2.0	*
5	Influenza and pneumonia (J09–J18)	1	2.0	*
5	Hernia of abdominal cavity and intestinal obstruction without hernia (K40–K46,K56)	1	2.0	*
5	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1	2.0	*
5	Assault (homicide) (†U01,X85–Y09)	1	2.0	*
...	All other causes (residual)	8	16.3	*
Postneonates, Asian and Other Pacific Islander, non-Hispanic, both sexes				
...	All causes	191	100.0	79.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	54	28.3	22.4
2	Sudden infant death syndrome (R95)	21	11.0	8.7
3	Accidents (unintentional injuries) (V01–X59)	16	8.4	*
4	Diarrhea and gastroenteritis of infectious origin (A09)	12	6.3	*
5	Influenza and pneumonia (J09–J18)	6	3.1	*
6	Assault (homicide) (†U01,X85–Y09)	5	2.6	*
7	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50–D89)	4	2.1	*
7	Diseases of the circulatory system (I00–I99)	4	2.1	*
7	Septicemia (A40–A41)	4	2.1	*
10	Meningitis (G00,G03)	3	1.6	*
10	Necrotizing enterocolitis of newborn (P77)	3	1.6	*
...	All other causes (residual)	59	30.9	24.4
Postneonates, Asian and Other Pacific Islander, non-Hispanic, male				
...	All causes	111	100.0	89.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	27	24.3	21.7
2	Accidents (unintentional injuries) (V01–X59)	10	9.0	*
3	Diarrhea and gastroenteritis of infectious origin (A09)	9	8.1	*
3	Sudden infant death syndrome (R95)	9	8.1	*
5	Influenza and pneumonia (J09–J18)	4	3.6	*
5	Assault (homicide) (†U01,X85–Y09)	4	3.6	*
7	Septicemia (A40–A41)	3	2.7	*
7	Diseases of the circulatory system (I00–I99)	3	2.7	*
9	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50–D89)	2	1.8	*
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	2	1.8	*
9	Necrotizing enterocolitis of newborn (P77)	2	1.8	*
9	Malignant neoplasms (C00–C97)	2	1.8	*
...	All other causes (residual)	34	30.6	27.3

Table 1–2. Infant, neonatal, and postneonatal deaths, percentage of total deaths, and mortality rate for the 10 leading causes of infant death, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are infant (younger than 1 year), neonatal (younger than 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic-origin categories are consistent with 1977 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated separately by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black non-Hispanic and White non-Hispanic should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on birth and death certificates. Deaths are based on race and Hispanic origin of decedent; live births are based on race and Hispanic origin of mother. National linked files should be used to measure infant mortality for specific subgroups. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*]

Rank ¹	Cause of death (based on ICD–10), age, race and Hispanic origin, and sex	Number ²	Percent of total deaths	Mortality rate ²
Postneonates, Asian and Other Pacific Islander, non-Hispanic, female				
...	All causes	80	100.0	68.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	27	33.8	23.0
2	Sudden infant death syndrome (R95)	12	15.0	*
3	Accidents (unintentional injuries) (V01–X59)	6	7.5	*
4	Diarrhea and gastroenteritis of infectious origin (A09)	3	3.8	*
5	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50–D89)	2	2.5	*
5	Meningitis (G00,G03)	2	2.5	*
5	Influenza and pneumonia (J09–J18)	2	2.5	*
8	Septicemia (A40–A41)	1	1.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00–D48)	1	1.3	*
8	Nutritional deficiencies (E40–E64)	1	1.3	*
8	Diseases of the circulatory system (I00–I99)	1	1.3	*
8	Pneumonitis due to solids and liquids (J69)	1	1.3	*
8	Hernia of abdominal cavity and intestinal obstruction without hernia (K40–K46,K56)	1	1.3	*
8	Renal failure and other disorders of kidney (N17–N19,N25,N27)	1	1.3	*
8	Respiratory distress of newborn (P22)	1	1.3	*
8	Atelectasis (P28.0–P28.1)	1	1.3	*
8	Assault (homicide) (†U01,X85–Y09)	1	1.3	*
8	Necrotizing enterocolitis of newborn (P77)	1	1.3	*
8	Infantile cerebral palsy (G80)	1	1.3	*
...	All other causes (residual)	14	17.5	*
Postneonates, Black, non-Hispanic, both sexes				
...	All causes	2,290	100.0	409.7
1	Sudden infant death syndrome (R95)	493	21.5	88.2
2	Accidents (unintentional injuries) (V01–X59)	397	17.3	71.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	263	11.5	47.1
4	Diseases of the circulatory system (I00–I99)	92	4.0	16.5
4	Assault (homicide) (†U01,X85–Y09)	92	4.0	16.5
6	Chronic respiratory disease originating in the perinatal period (P27)	55	2.4	9.8
7	Diarrhea and gastroenteritis of infectious origin (A09)	54	2.4	9.7
8	Septicemia (A40–A41)	42	1.8	7.5
9	Influenza and pneumonia (J09–J18)	39	1.7	7.0
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	31	1.4	5.5
...	All other causes (residual)	732	32.0	131.0
Postneonates, Black, non-Hispanic, male				
...	All causes	1,266	100.0	445.8
1	Sudden infant death syndrome (R95)	260	20.5	91.5
2	Accidents (unintentional injuries) (V01–X59)	226	17.9	79.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	127	10.0	44.7
4	Assault (homicide) (†U01,X85–Y09)	63	5.0	22.2
5	Diseases of the circulatory system (I00–I99)	45	3.6	15.8
6	Chronic respiratory disease originating in the perinatal period (P27)	31	2.4	10.9
7	Influenza and pneumonia (J09–J18)	30	2.4	10.6
8	Diarrhea and gastroenteritis of infectious origin (A09)	29	2.3	10.2
9	Septicemia (A40–A41)	24	1.9	8.5
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	17	1.3	*
...	All other causes (residual)	414	32.7	145.8

Table 1–2. Infant, neonatal, and postneonatal deaths, percentage of total deaths, and mortality rate for the 10 leading causes of infant death, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are infant (younger than 1 year), neonatal (younger than 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic-origin categories are consistent with 1977 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated separately by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black non-Hispanic and White non-Hispanic should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on birth and death certificates. Deaths are based on race and Hispanic origin of decedent; live births are based on race and Hispanic origin of mother. National linked files should be used to measure infant mortality for specific subgroups. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*]

Rank ¹	Cause of death (based on ICD–10), age, race and Hispanic origin, and sex	Number ²	Percent of total deaths	Mortality rate ²
Postneonates, Black, non-Hispanic, female				
...	All causes	1,024	100.0	372.4
1	Sudden infant death syndrome (R95)	233	22.8	84.7
2	Accidents (unintentional injuries) (V01–X59)	171	16.7	62.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	136	13.3	49.5
4	Diseases of the circulatory system (I00–I99)	47	4.6	17.1
5	Assault (homicide) (†U01.X85–Y09)	29	2.8	10.5
6	Diarrhea and gastroenteritis of infectious origin (A09)	25	2.4	9.1
7	Chronic respiratory disease originating in the perinatal period (P27)	24	2.3	8.7
8	Septicemia (A40–A41)	18	1.8	*
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	14	1.4	*
10	Anoxic brain damage, not elsewhere classified (G93.1)	9	0.9	*
10	Influenza and pneumonia (J09–J18)	9	0.9	*
10	Necrotizing enterocolitis of newborn (P77)	9	0.9	*
...	All other causes (residual)	300	29.3	109.1
Postneonates, White, non-Hispanic, both sexes				
...	All causes	2,794	100.0	148.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	543	19.4	28.9
2	Sudden infant death syndrome (R95)	488	17.5	25.9
3	Accidents (unintentional injuries) (V01–X59)	485	17.4	25.8
4	Diseases of the circulatory system (I00–I99)	140	5.0	7.4
5	Assault (homicide) (†U01.X85–Y09)	93	3.3	4.9
6	Influenza and pneumonia (J09–J18)	57	2.0	3.0
7	Chronic respiratory disease originating in the perinatal period (P27)	44	1.6	2.3
8	Diarrhea and gastroenteritis of infectious origin (A09)	37	1.3	2.0
9	Septicemia (A40–A41)	32	1.1	1.7
10	Malignant neoplasms (C00–C97)	22	0.8	1.2
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	22	0.8	1.2
...	All other causes (residual)	831	29.7	44.2
Postneonates, White, non-Hispanic, male				
...	All causes	1,572	100.0	162.9
1	Sudden infant death syndrome (R95)	290	18.4	30.1
2	Accidents (unintentional injuries) (V01–X59)	275	17.5	28.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	268	17.0	27.8
4	Diseases of the circulatory system (I00–I99)	73	4.6	7.6
5	Assault (homicide) (†U01.X85–Y09)	57	3.6	5.9
6	Influenza and pneumonia (J09–J18)	33	2.1	3.4
7	Chronic respiratory disease originating in the perinatal period (P27)	28	1.8	2.9
8	Diarrhea and gastroenteritis of infectious origin (A09)	25	1.6	2.6
9	Septicemia (A40–A41)	19	1.2	*
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	15	1.0	*
...	All other causes (residual)	489	31.1	50.7

Table 1–2. Infant, neonatal, and postneonatal deaths, percentage of total deaths, and mortality rate for the 10 leading causes of infant death, by bridged race and Hispanic origin and sex: United States, 2020—Con.

[Rates are infant (younger than 1 year), neonatal (younger than 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic-origin categories are consistent with 1977 Office of Management and Budget standards. Race and Hispanic origin are reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic people are not tabulated separately by race; data for non-Hispanic people are tabulated by race. Data for racial and ethnic groups other than Black non-Hispanic and White non-Hispanic should be interpreted with caution because of inconsistencies between reporting Hispanic origin and race on birth and death certificates. Deaths are based on race and Hispanic origin of decedent; live births are based on race and Hispanic origin of mother. National linked files should be used to measure infant mortality for specific subgroups. A dagger (†) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*]

Rank ¹	Cause of death (based on ICD–10), age, race and Hispanic origin, and sex	Number ²	Percent of total deaths	Mortality rate ²
	Postneonates, White, non-Hispanic, female			
...	All causes	1,222	100.0	133.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	275	22.5	30.0
2	Accidents (unintentional injuries) (V01–X59)	210	17.2	22.9
3	Sudden infant death syndrome (R95)	198	16.2	21.6
4	Diseases of the circulatory system (I00–I99)	67	5.5	7.3
5	Assault (homicide) (†U01,X85–Y09)	36	2.9	3.9
6	Influenza and pneumonia (J09–J18)	24	2.0	2.6
7	Malignant neoplasms (C00–C97)	16	1.3	*
7	Chronic respiratory disease originating in the perinatal period (P27)	16	1.3	*
9	Septicemia (A40–A41)	13	1.1	*
10	Diarrhea and gastroenteritis of infectious origin (A09)	12	1.0	*
...	All other causes (residual)	355	29.1	38.7

... Category not applicable.

* Estimate does not meet National Center for Health Statistics standards of reliability.

¹Based on number of deaths.

²Data for two or more races and origin not stated are included in "all races and origins" but are not distributed among race and Hispanic-origin groups. For the race and Hispanic-origin categories shown, only one race was reported on the death certificate.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Table I-3. Estimated population, by age group, bridged race and Hispanic origin, and sex: United States, July 1, 2020

Age group (years)	Non-Hispanic														
	Hispanic			American Indian and Alaska Native			Asian and Pacific Islander			Black			White		
	Total	Male	Female	Total	Male	Female	Total	Male	Female	Total	Male	Female	Total	Male	Female
Total	61,312,879	30,922,535	30,390,344	2,766,392	1,354,955	1,411,437	21,284,836	10,207,105	11,077,731	43,620,633	20,904,808	22,715,825	200,499,383	98,866,799	101,632,584
Younger than 1	970,381	494,752	475,629	34,541	17,521	17,020	237,088	121,629	115,459	589,846	300,818	289,028	1,903,154	973,421	929,733
1-4	4,029,420	2,051,303	1,978,117	144,764	73,468	71,296	1,034,813	533,089	501,724	2,424,518	1,230,226	1,194,292	7,932,767	4,064,930	3,867,837
5-9	5,172,470	2,633,961	2,538,509	197,914	100,808	97,106	1,322,302	679,680	642,622	3,102,465	1,576,621	1,525,844	10,442,560	5,355,683	5,086,877
10-14	5,368,702	2,736,509	2,632,193	205,209	103,490	101,719	1,232,969	627,726	605,243	3,163,709	1,605,603	1,558,106	10,783,834	5,521,640	5,262,194
15-19	5,106,426	2,602,741	2,503,685	203,344	102,942	100,402	1,270,545	637,420	633,125	3,107,185	1,574,744	1,532,441	11,273,429	5,776,750	5,496,679
20-24	4,907,287	2,512,676	2,394,611	209,328	105,803	103,525	1,388,775	700,795	687,980	3,262,982	1,652,385	1,610,597	11,826,383	6,061,187	5,765,196
25-34	9,627,323	5,013,447	4,613,876	430,352	217,673	212,679	3,521,572	1,738,283	1,783,289	6,907,314	3,447,824	3,459,490	25,583,085	13,027,152	12,555,933
35-44	8,742,890	4,514,577	4,228,313	343,264	168,706	174,558	3,344,077	1,578,366	1,765,711	5,566,714	2,649,354	2,917,360	24,139,247	12,134,865	12,004,382
45-54	7,250,232	3,652,520	3,597,712	314,263	152,117	162,146	2,855,047	1,334,467	1,520,580	5,154,467	2,408,645	2,745,822	24,792,124	12,376,943	12,415,181
55-64	5,255,928	2,573,717	2,682,211	332,630	155,990	176,640	2,306,507	1,056,978	1,249,529	5,074,739	2,323,764	2,750,975	29,433,873	14,378,985	15,054,888
65 and older	4,881,820	2,136,332	2,745,488	350,783	156,437	194,346	2,771,141	1,198,672	1,572,469	5,266,694	2,134,824	3,131,870	42,388,927	19,195,243	23,193,684
65-74	2,978,586	1,367,913	1,610,673	223,032	102,414	120,618	1,657,299	730,216	927,083	3,298,393	1,414,074	1,884,319	24,392,088	11,568,923	12,823,165
75-84	1,368,466	573,624	794,842	95,409	41,929	53,480	789,424	345,641	443,783	1,435,533	557,153	878,380	12,762,715	5,704,928	7,057,787
85 and older	534,768	194,795	339,973	32,342	12,094	20,248	324,418	122,815	201,603	532,768	163,597	369,171	5,234,124	1,921,392	3,312,732

NOTES: Population estimates are based on the U.S. Census 2010. Race and Hispanic-origin categories are consistent with 1977 Office of Management and Budget standards.

SOURCE: U.S. Census Bureau.