

APPENDIX



Objective Status: Mental Health and Mental Disorders


Target met O Improving

○ Little/No change ● Getting worse ○ Baseline only ○ Developmental

- SA-1 Suicide rate
- SA-2 Suicide attempts, grades 9–12
- SA-3 Engaging in disordered eating to control weight, grades 9-12
- SA-4.1 Major depressive episodes, 12–17 years
- SA-4.2 Major depressive episodes, 18+ years
- SA-5 Primary care facilities that provide mental health treatment
- SA-6 Children with mental health problems receiving treatment, 4—17 years
- SA-7 Juvenile residential facilities that screen admissions for mental health problems
- SA-8 Persons with serious mental illness who are employed, 18+ years

- SA-9.1 Adults with serious mental illness who receive treatment, 18+ years
- SA-9.2 Adults with major depressive episodes who receive treatment, 18+years
- SA-10 Persons with co-occurring substance abuse and mental disorders who receive treatment, 18+ years
- SA-11.1 Primary care physician office visits by adults that include screening for depression, 19+ years
- SA-11.2 Primary care physician office visits by youth that include screening for depression, 12-18 years
- SA-12 Homeless adults with mental health problems receiving mental health services, 18+ years

Current HP2020 Objective Status: Mental Health and Mental Disorders



Objective Status: Substance Abuse

Target met OImproving OLittle/No change OGetting worse OBaseline only ODevelopmental OInformational

- SA–1 Rode with a driver who has been drinking: 9–12th graders
- SA-2.1 Alcohol, never used: At risk youth 12-17 years
- SA-2.2 Marijuana, never used: At risk youth 12-17 years
- SA-2.3 Alcohol, never used: 12th graders
- SA–2.4 Illicit drugs, never used: 12th graders
- SA-3.1 Disapproval of 1-2 drinks every day: 8th graders
- SA–3.2 Disapproval of 1–2 drinks every day:10th graders
- SA–3.3 Disapproval of 1–2 drinks every day:12th graders
- SA-3.4 Disapproval of trying marijuana: 8th graders
- SA-3.5 Disapproval of trying marijuana:10th graders
- SA-3.6 Disapproval of trying marijuana:12th graders
- SA-4.1 Risk perception, binge drinking-weekly:12-17 years
- SA-4.2 Risk perception, marijuana use-monthly:12-17 years
- SA-4.3 Risk perception, cocaine use-monthly:12-17 years
- SA–5 DWI and other specialty courts: US
- SA-6 Mandatory ignition interlock laws: States
- SA–7 Injection drug use treatment, admissions
- SA-8.1 Illicit drug use treatment: 12+ years
- SA-8.2 Alcohol/illicit drug use treatment: 12+ years
- SA-8.3 Alcohol treatment: 12+ years
- SA–9 Alcohol/illicit drug use referral for care/treatment: ED visits
- SA–10 Alcohol Screening and Brief Intervention: Primary care and trauma centers

- SA-11 Cirrhosis deaths
- SA–12 Drug–induced deaths
- SA-13.1 Illicit drug use-past month: 12-17 years
- SA-13.2 Marijuana use-past month: 12-17 years
- SA–13.3 Illicit drug use–past month: 18+ years
- SA-14.1 Binge drinking-past 2 weeks: 12th graders
- SA-14.2 Binge drinking-past 2 weeks: College students
- SA-14.3 Binge drinking-past month: 18+ years
- SA-14.4 Binge drinking-past month: 12-17 years
- SA-15 Excessive drinking-past month: 18+ years
- SA-16 Average annual alcohol consumption: 14+ years
- SA–17 Alcohol–related driving deaths
- SA–18.1 Steroid use: 8th graders
- SA–18.2 Steroid use: 10th graders
- SA–18.3 Steroid use: 12th graders
- SA–19.1 Prescription pain reliever abuse–past year: 12+ years
- SA–19.2 Prescription tranquilizer abuse–past year: 12+ years
- SA–19.3 Prescription stimulants abuse–past year: 12+ years
- SA-19.4 Prescription sedatives abuse-past year: 12+ years
- SA–19.5 Prescription psychotherapeutic drug abuse–past year: 12+ years
- SA-20 Alcohol-attributable deaths
- SA-21Iinhalant use-past year: 12-17 years

Current HP2020 Objective Status: Substance Abuse

 \bigcirc

