
Common Epidemiology of *Rickettsia felis* Infection and Malaria, Africa

Oleg Mediannikov,¹ Cristina Socolovschi,¹ Sophie Edouard, Florence Fenollar, Nadjat Mouffok, Hubert Bassene, Georges Diatta, Adama Tall, Hamidou Niangaly, Ogobara Doumbo, Jean Bernard Lekana-Douki, Abir Znazen, M'hammed Sarih, Pavel Ratmanov, Herve Richet, Mamadou O. Ndiath, Cheikh Sokhna, Philippe Parola, and Didier Raoult

This study aimed to compare the epidemiology of *Rickettsia felis* infection and malaria in France, North Africa, and sub-Saharan Africa and to identify a common vector. Blood specimens from 3,122 febrile patients and from 500 nonfebrile persons were analyzed for *R. felis* and *Plasmodium* spp. We observed a significant linear trend ($p < 0.0001$) of increasing risk for *R. felis* infection. The risks were lowest in France, Tunisia, and Algeria (1%), and highest in rural Senegal (15%). Co-infections with *R. felis* and *Plasmodium* spp. and occurrences of *R. felis* relapses or reinfections were identified. This study demonstrates a correlation between malaria and *R. felis* infection regarding geographic distribution, seasonality, asymptomatic infections, and a potential vector. *R. felis* infection should be suspected in these geographical areas where malaria is endemic. Doxycycline chemoprophylaxis against malaria in travelers to sub-Saharan Africa also protects against rickettsioses; thus, empirical treatment strategies for febrile illness for travelers and residents in sub-Saharan Africa may require reevaluation.

Author affiliations: Aix Marseille Université, Marseille, Faculté de Médecine, Marseille, France (O. Mediannikov, C. Socolovschi, S. Edouard, F. Fenollar, P. Ratmanov, H. Richet, P. Parola, D. Raoult); Institut de Recherche pour le Développement, Dakar, Senegal (O. Mediannikov, F. Fenollar, H. Bassene, G. Diatta, M.O. Ndiath, C. Sokhna, D. Raoult); Centre Hospitalo-Universitaire d'Oran, Oran, Algeria (N. Mouffok); Institut Pasteur de Dakar, Dakar (A. Tall); University of Sciences, Techniques and Technology, Bamako, Mali (H. Niangaly, O. Doumbo); Unité de Parasitologie Médicale Centre International de Recherche Médicale de Franceville, Franceville, Gabon (J.B. Lekana-Douki); Université des Sciences de la santé de Libreville, Libreville, Gabon (J.B. Lekana-Douki); Habib Bourguiba University Hospital, Sfax, Tunisia (A. Znazen); Institut Pasteur du Maroc, Casablanca, Morocco (M. Sarih); and Far Eastern State Medical University, Khabarovsk, Russia (P. Ratmanov)

Investigations examining the etiologic spectrum of fever of unknown origin in Africa rapidly progressed during 2008–2011 (1–3), providing increased knowledge about bacterial infections. Bacterial agents that have been most frequently identified in North and sub-Saharan Africa by culture are non-typhoidal *Salmonella*, *Streptococcus pneumoniae*, *Staphylococcus aureus*, *Escherichia coli*, and *Mycobacterium tuberculosis* (2). Several studies have assessed the effect of fastidious bacterial infections in systemic febrile illness, including *Rickettsia felis* (4–6), *Coxiella burnetii* (7), *Tropheryma whippelii* (3), and *Borrelia* spp. (1,8). Tourism, immigration, international business travel, international aid work, and the deployment of troops overseas were documented as contributors to a tremendous increase in international travel during 1996–2004 (9). International tourist arrivals reached 940 million worldwide during 2010, an increase of 6.6% over 2009, and the current total number of international migrants has increased to an estimated 214 million persons in 2012 (10). Consequently, physicians in the Western hemisphere increasingly encounter febrile patients returning from international travel who were exposed to tropical infections that the physicians are unfamiliar with (9,10). Among international travelers, malaria, dengue, and rickettsiosis are among the most identified etiologies of febrile illness, and exposure to mosquitoes is reported as the most common source of fever (11).

Rickettsia felis, an obligate intracellular Gram-negative bacterium belonging to the spotted fever group of *Rickettsia*, has been shown to be a common agent of bloodstream infections in among humans Senegal and Kenya, identified in 7% of the population evaluated (4–6). However, the epidemiology (including vectors and reservoirs)

and clinical picture of this emerging infection in the rest of Africa is largely unknown (12,13). During 2011, a possibly primary infection with *R. felis*, named “yaaf,” was hypothesized in the case of an 8-month-old girl in Senegal with polymorphous skin lesions (12).

The considerable frequency of *R. felis* infections observed in febrile patients in malaria-endemic regions and the many relapses previously reported (4,5) led us to investigate the possible correlation of *R. felis* and that of the parasite, *Plasmodium falciparum*, a known vector of malaria. The reservoirs for malaria and many rickettsial species are mammals, including humans; humans have long been known to be a reservoir for malaria, and were documented as the reservoir for *R. prowazekii*, the agent of epidemic typhus (14). Vectors for both organisms are arthropods: for rickettsial diseases vectors are typically ticks, lice or mites, and infected humans are susceptible to relapse (such as epidemic and scrub typhus) (14).

The vectors for malaria are mosquitoes of the genus *Anopheles* that breed in warm and humid areas (15). Malaria is particularly common among young patients, because progressive immunity develops following multiple infections as the child grows older. Great apes in Cameroon were recently identified as targets or possibly the origin of malaria (16). *R. felis* has recently been detected in *Anopheles gambiae* mosquitoes in molecular form S, in *Aedes albopictus* mosquitoes, and in gorilla fecal samples (17–19). These elements suggest comparable features within the epidemiologic cycles of malaria and *R. felis* infection. In addition, co-infections by *R. felis* and *P. falciparum* have been reported in Kenya (5). To prove the hypothesis of the similar epidemiology of malaria and *R. felis* infection, target populations, clinical phenomena (relapses and bacteremia in apparently asymptomatic patients), and geographic and seasonal distribution should be compared. The objective of this work is to clarify the epidemiology of *R. felis* infection and to compare it with malarial epidemiology.

Materials and Methods

Study Areas and Participants

Febrile Patients

During June 2010–March 2012, a cohort of 2,075 patients (67% <15 years of age; sex ratio, 1:1) from 14 health centers distributed throughout rural Senegal (Senegal study sites S₁–S₆) were enrolled in this study. The study sites spanned various ecosystems, from dry regions in the north (Dielmo, Senegal study region 1,–S₁, Ndiop-S₂, Keur Momar Sarr-S₃, and Niakhar-S₄) to humid regions in the south (Basse-Casamance-S₅ and Kedougou-S₆) that had a rainy season during June through October (online Technical Appendix Table,

wwwnc.cdc.gov/EID/article/19/11/13-0361-Techapp1.pdf; Table). In addition, patients from various medical facilities were included: 100 from rural Mali dispensaries: Diakabou-Mali study site M₁ and Kole-Mali study site M₂; 50 from Franceville, in urban Gabon (pediatric consultation); 183 from Sfax, Tunisia (infectious diseases and pediatric departments); 266 from Oran, Algeria (department of infectious diseases); 48 from the Kenitra region, rural Morocco (dispensaries); and 400 from Marseille, France (hospital emergency units) (Figure 1). Questionnaires and informed consent forms were completed upon enrollment in the study. For each febrile patient (axillary temperature >37.5°C), an interview was conducted, a blood sample (200 µl blood containing EDTA) was collected, and a medical examination was performed. The national ethics committees of Senegal, Gabon, and France approved this project (No. 0–00.87MSP/DS/CNERS and No. 001380MSP/DS/CNERS).

Control Group

Samples were obtained from 400 afebrile persons (62% >15 years of age) from S_{1–2} who participated in a longitudinal study of malaria (20) and 100 persons from France who were under the medical care of 1 of the authors (D.R.) for conditions other than malaria.

Arthropod Collection in Senegal

Arthropod specimens collected in Senegal consisted of 949 adult mosquitoes from 3 locations (Table 1, 154 mosquito larvae from Mariste, Dakar, 370 ticks from 2 locations, 160 adult bed bugs from 6 locations, and 384 midges from 2 locations). The *Anopheles arabiensis* mosquito larvae were collected from breeding sites in Mariste, Dakar. The pooled larvae were maintained under laboratory conditions until they grew to the adult stage. In sites S_{1–2}, 144 adult ticks (2 *Rhipicephalus* spp., 4 *Argas persicus*, and 138 *Ornithodoros sonrai*) from 55 burrows inside of 16 human dwellings were collected. A total of 226 *Ornithodoros capensis* ticks were manually collected from the nests of great cormorants (*Phalacrocorax carbo*) in Sarpan Island (îles de la Madeleine) near Dakar. Bed bugs were manually captured from the beds of ill persons. The collection of *Culicoides* spp. was performed in S_{1–2} by using overnight posed CDC light traps with 0.7-mm mesh size. The arthropods were identified at the species level by using morphological characteristics according to identification keys.

Molecular Analysis

DNA was extracted by using the 2-stage protocol for a QIAamp kit (QIAGEN, Hilden, Germany) for the S_{4–6} groups (3,4,7), and a Biorobot EZ1 Workstation (QIAGEN, Courtaboeuf, France) was used to extract DNA from




Figure 1. Prevalence of *Rickettsia felis* infection (A) and *Plasmodium* spp. infection (malaria) (B) in febrile patients in Gabon, Senegal, Mali, Algeria, Morocco, Tunisia, and France, June 2010–April 2012.

samples from S₁₋₂, Algeria, Tunisia, Morocco, and France. In Gabon, the DNA Blood Omega Bio-tek-E.Z.N.A method (Omega Bio-tek, Norcross, GA, USA) was used according to the manufacturer’s protocol. For all locations, DNA was eluted in 100 µL of elution buffer, and 5 µL was used per reaction.

Quantitative real-time reverse transcription PCR (qRT-PCR) was performed by using a 7900HT-thermocycler (Applied Biosystems) with the QuantiTect-Probe PCR Kit (QIAGEN, Courtabeuf, France). Only samples positive for the β-actin gene product were considered reliable (3); thus, 51 and 9 samples from Senegal and Algeria, respectively, were excluded. All samples were screened by using a *Rickettsia* genus-specific qRT-PCR targeting the *gltA* gene and an *R. felis*-specific qRT-PCR targeting the *bioB* gene (4). The positive samples were tested by a second *R. felis*-specific qRT-PCR targeting the *orfB* gene (18). A sample was considered positive when the qRT-PCRs were positive for the 2 different specific genes. Positive samples from arthropods were further tested for plasmid pRFδ (21) and by a newly designed *R. felis*-specific qRT-PCR targeting the *vapB1* gene with the primers VapB1.R (5'-AGGCGAAAGCTTTGACGTG-3') and VapB1.F (5'-TGTCTTTCATGAATTGATCAGCA-3') and the probe VapB1.P (6-FAM-5'-AAGGCTTGGTTTCTGCGGGC-3'TAMRA).

Blood smears stained with Giemsa were examined for the samples collected in Gabon. All other samples were tested by using a *Plasmodium*-genus specific qRT-PCR targeting the *Cox-1* gene found in all *Plasmodium* species; the

primers Psp_15.F (5'-AGGAACTCGACTGGCCTACA-3') and Psp_16.R (5'-CCAGCGACAGCGGTTATACT-3') and the (6FAM-5'-CGAACGCTTTTAACGCCTGACATGG-3'TAMRA) probe were used. The positive samples were subsequently tested by *Plasmodium*-genus specific qRT-PCR targeting 18S rRNA with the primers Plasm_18S_2_MBF (5'-AGGCAACAACAGGTCTGTGA-3') and Plasm_18S_2_MBR (5'-GCAATAATCTATCCCCATCACG-3') and the (6FAM-5'- GAACTAGGCTGCACGCGTGCTACA-TAMRA-3') probe.

Statistical Analysis

Statistical analyses were performed by using the Statcalc module of Epi Info 3.5.3 (Centers for Disease Control and Prevention, Atlanta, GA, USA) to calculate the χ² values for the incidence rate trends calculated for each country. PASW Statistics software 17.0 (IBM, SPSS Inc., Armonk, NY, USA) was used to perform Pearson correlation analyses. The relative risk (RR) and the 95% CI of the risk were calculated by using either the Mantel-Haenszel χ² test or Fisher’s exact test. The statistical significance of the χ² values was evaluated at α = 0.05. The attack rates of *R. felis* infection and malaria were calculated for each country, site, sex, and age range. In contrast, the incidence rates of *R. felis* infection and malaria for S₁₋₂ were calculated monthly and yearly from June 13, 2010 through October 13, 2011. The data from a study performed in 2009 (4) were combined with those of this study to determine the frequency of relapses or re-infections of *R. felis* infections in S₁₋₂.

Table 1. Detection of *Rickettsia felis* DNA in mosquitoes, Senegal, March 2010–September 2012

Geographic location in Senegal	Period of collection	Collection method	Mosquito species, morphological identification	DNA samples	<i>Rickettsia felis</i> detection (%)
Ferlo, 15°52'N, 15°15'W	Mar 2010	CDC type light trap collections*	<i>Aedes luteocephalus</i>	203	1 (<1)
			<i>Culex quinquefasciatus</i>	186	0/186
Dakar, 14°41'N, 17°26'W	Dec 2011	Immature stages-lab conditions	<i>Anopheles arabiensis</i>	154 †	2 (<1)‡
Diélmo, 13°43'N, 16°24'W	Jul 2012	Human landing catches	<i>An. arabiensis</i>	8	0
			<i>An. welcomei</i>	6	0
			<i>Mansonia uniformis</i>	6	0
			<i>C. quinquefasciatus</i>	4	0
	Sep 2012	Pyrethrum spray catches	<i>An. ziemanni</i>	7	1 (14)
			<i>An. pharoensis</i>	10	1 (10)
			<i>M. uniformis</i>	8	2 (25)
			<i>An. welcomei</i>	8	0
			<i>An. funestus</i>	7	2 (29)
Elinkine, 12°30'N, 16°39'W	Sep 2012	CDC-type light trap collections	<i>An. gambiae</i>	50	0
			<i>Culex</i> sp.	10	0
		Pyrethrum spray catches	<i>Culex</i> sp.	10	0
			Human landing catches	<i>An. gambiae</i>	290
		<i>An. squamosus</i>		27	0
		<i>An. ziemanni</i>		31	0
		<i>Culex</i> sp.		35	0
		<i>Aedes</i> sp.		23	0
		<i>Mansonia</i> sp.		20	0
		Total			

*Manufactured by John W. Hock Company, Gainesville, FL, USA.

†Including 20 male mosquitoes.

‡*R. felis* DNA was detected in 1 male mosquito.

Results

Rickettsia felis Detection

Senegal

The attack rate of *R. felis* infections in febrile patients was 15% (312/2,024); those infections occurred primarily during the rainy season rather than the dry season (207/1,105 vs. 105/916, respectively; $p < 0.0001$). The risk of developing *R. felis* infection was 1.6× higher during the rainy period (95% CI 1.3–2) than during the dry period. When calculated by site, substantial differences in the rates of *R. felis* infection were observed (Table 2). The highest attack rates were observed in S_{5-6} , reaching 40% (92/231) from August–October 2011. The lowest attack rate was observed in S_{1-2} (7%–8%) and was significantly lower than that observed at the 4 other sites S_{3-6} ($p \geq 0.001$) (Table 2).

Incidence rates were obtained from 2 health centers (Figure 2). In 2011, the incidence rate of *R. felis* in S_1 was 6.7 (4.8–9.0) per 100 person-years or 0.55 (0.39–0.76) per 100 person-months; the incidence rate in S_2 was 3.1 (1.8–4.9) per 100 person-years or 0.26 (0.15–0.41) per 100 person-months during the same period. In S_{1-2} , a significant difference was found between the incidence of *R. felis* for patients <15 years of age, which was 0.23 (0.16–0.31) per 100 person-months, and the incidence in patients >15 years of age, which was 0.10 (0.06–0.15) per 100 person-months

(relative risk [RR] 2.38, 95% CI 1.34–4.28, $p = 0.003$). When the incidence rates by age group were calculated according to sex, a significant difference was observed only in the male group, in which the incidence rate was significantly higher in the patients <15 years of age than in the patients >15 years of age (0.29 vs. 0.07 per 100 person-months, RR 5.97, 95% CI 2.28–17.15, $p = 0.001$).

Table 3 shows the age distribution of *R. felis* infection. The occurrence of *R. felis* infection was significantly lower in patients 1–3 years of age (10%) than in patients >4 years of age ($p = 0.03$ for patients 4–6 years of age (15%); $p = 0.003$ for patients 7–15 years of age (16%); $p = 0.004$ for patients 16 to 29 years of age (16%); $p = 0.002$ for those >30 years of age (17%). The sex ratio for *R. felis* was 145M/162F (1:1.1). No deaths associated with *R. felis* infection were registered.

Combining these data with our preliminary report of 8 infected patients during 2008–2009 in S_{1-2} (4), we identified 61 patients with *R. felis* infections among a total of 456 villagers tested in S_{1-2} . A second *R. felis* infection was diagnosed in 5 patients after 44 to 911 days, and 1 patient was positive for *R. felis* infection a second and third time at days 378 and 441, respectively. The 6 patients (4 male, 2 female) who had relapses or re-infections were from S_1 , and 5 were <6 years of age.

Other Countries

Samples from 3 patients (3%, 3/100) in rural Mali (M_1 , 1/50; M_2 , 2/50), 5 patients (10%, 5/50) in urban

Table 2. Attack rate of *Rickettsia felis* infection and malaria by country and geographic site, Africa, 2010–2012

Participant status, country, and study site (site abbreviation)	Collection period	No. samples*	No. samples positive/no. tested (%)		
			<i>R. felis</i>	<i>Plasmodium</i> spp.	<i>R. felis/ Plasmodium</i> spp. co-infection
Febrile patients					
Senegal	Jun 2010–Mar 2012	2,024	312/2,024 (15)	400/1,867† (21)	66/285 (23)*
Dielmo (S ₁)	Jun 2010–Feb 2012	540	39/540 (7)	118/509 (23)	8/36 (22)
Ndiop (S ₂)	Jun 2010–Feb 2012	246	20/246 (8)	33/237 (14)	3/18 (17)
Keur-Momar Sarr (S ₃)	Mar–Nov 2011	223	36/223 (16)	44/196 (22)	9/33 (27)
Niakhar (S ₄)	Oct 2010–Mar 2012	316	76/316 (24)	74/303 (24)	18/74 (24)
Basse-Casamance (S ₅)	Jan 2011–Mar 2012	411	84/411 (20)	37/350 (11)	7/69 (10)
Kedougou (S ₆)	2011	288	57/288 (20)	94/272 (34)	21/55 (38)
Gabon					
Franceville	2011	50	5/50 (10)	19/50 (38)‡	2/5 (40)**
Mali	2011	100	3/100 (3)	90% (90/100)	3/3 (100)
Diakambou (M1)	Oct	50	1/50 (2)	82% (41/50)	1/1 (100)
Kole (M2)	Nov	50	2/50 (4)	98% (49/50)	2/2 (100)
Algeria					
Oran	Jul–Sep 2012	257	2/257 (1)	1/257 (0.4%)	0/1
Morocco					
Casablanca	May–Jun 2006	48	1/48 (2)	0/38†	0
Tunisia					
Sfax	2012	183	0/183	0/183	0
France					
Marseille	2012	400	0/400	0/400	0
Afebrile persons					
Senegal (S ₁ –S ₂)	Dec 2011–Apr 2012	391	17/391 (4)	5/391 (1)	0/5
France					
Marseille	2011–2012	100	0/100	0/100	0

*Reliable samples.

†There were insufficient DNA samples for the analysis of *Plasmodium* spp., as decided a posteriori.

‡Positive by blood smear.

Gabon, 1 patient (2%, 1/48) in rural Morocco, and 2 patients (1%, 2/257) in Algeria were positive for *R. felis* (Table 2). Conversely, *R. felis* DNA was not detected among the samples from febrile persons in France and Tunisia.

When the *R. felis* infection rates of the different countries were compared, *R. felis* was detected more often in countries with high malaria rates compared with countries with low malaria rates (Senegal, Gabon, and Mali vs. Algeria, Tunisia, Morocco, and France; $p < 0.001$) (Figure 1). The trend analysis showed a significant linear trend of increasing risk for *R. felis* infection; a lower risk was shown in northern countries (France, Tunisia, Morocco, and Algeria) and a higher risk in southern countries (Mali, Gabon, and Senegal) ($p < 0.0001$). The probability of *R. felis* infection was 1.00 for Algeria (baseline), 2.8 for Morocco, 4 for Mali, 14.5 for Gabon, and 24 for Senegal.

Malaria

Senegal

The attack rate of *Plasmodium* spp. in febrile persons from Senegal was 21% (400/1868, 206 females); those infections occurred significantly more often during the rainy season compared with the dry season (256/1042 vs. 144/822, respectively; $p = 0.0002$). The risk for ma-

laria was 1.4× higher during the rainy period than during the dry period (95% CI 1.2–1.7, $p < 0.0001$). The highest rate was in southeastern S₆, whereas the lowest rate, 11% (37/350), was in southwestern S₅ (Table 2). During the same time period, the incidence rate of malaria was 17.6 per 100 person-years or 1.46 per 100 person-months for S₁ and 5.1 per 100 person-years or 0.42 per 100 person-months for S₂. The highest incidence of malaria was among patients <15 years of age in S₁₋₂ (0.55 (0.44–0.67) versus 0.22 (0.16–0.30) per 100 person-months, RR 2.51, 95% CI 1.73–3.65, $p < 0.0001$).

When the incidence rate by age group was calculated according to sex, the highest incidence was found among girls <15 years of age: 0.47 (0.37–0.65) versus 0.25 (0.16–0.37) per 100 person-months, RR 1.92, 95% CI 1.15–3.18, $p = 0.01$; and boys (0.62 (0.47–0.81) versus 0.18 (0.10–0.30) per 100 person-months, RR 3.38, 95% CI 1.90–5.99, $p < 0.0001$) groups. Table 3 shows the age distribution of malaria. Patients 7–15 years of age (33%) were infected with *Plasmodium* spp. significantly more often than those in other age groups (15% for patients 1–3 years of age, $p < 0.0001$ to 23% for patients 4–6 years of age, $p = 0.004$). Co-infection of *Plasmodium* spp. and *R. felis* was found in 66 case-patients (23%, 66/285), mostly in women (61%) and in children 7–15 years of age (43%).




Figure 2. Incidence of *Rickettsia felis* and *Plasmodium* spp. infection (malaria) in patients, by age, in Dielmo and Ndiop, Senegal.

Other Countries

Plasmodium DNA was detected in 90% of the blood samples collected in Mali; 3 patients with malaria from Mali were co-infected with *R. felis* (Table 2). In Gabon, samples from 38% (19/50) of the patients tested positive for malaria by using blood smears; 2 of those patients were co-infected with *R. felis*. We most likely misdiagnosed malaria among the patients in Gabon, as based on the lower sensitivity and high specificity of microscopy versus PCR as the standard (22). *Plasmodium* DNA was not detected in the samples from Tunisia, Morocco, or France. However, 1 *Plasmodium* spp.-positive sample was collected in Algeria from a 21-year-old woman who was hospitalized for high fever, chills, and sweats after having spent >2 months visiting her family in Niger without malaria chemoprophylaxis.

Correlation of *R. felis* with Malaria

Using the Pearson correlation test, we found a significant correlation between the number of patients infected with *R. felis* and those infected with *Plasmodium* spp. ($p < 0.002$): a higher number of *R. felis* infections correlated with a higher number of malaria cases. A significant correlation was also found for seasonality for infection by both pathogens: most cases occurred during the rainy period ($p < 0.0001$). In addition, children <3 years of age were infected with both organisms less often than persons >4 years of age, and the Pearson test showed a significant correlation between *R. felis* and malaria ($p = 0.001$) for this age group.

Control Group

R. felis DNA was detected in 4% of the afebrile persons (17/391) from Senegal, 12 of whom were children (<15 years of age); malaria was detected in 5 afebrile persons, 3 of whom were children. Both pathogens were detected significantly less often in afebrile patients than in febrile patients ($p < 0.001$). DNA from *R. felis* and *Plasmodium* were not detected among persons in the control group in France.

Arthropod Study

Samples from 9 mosquitoes ($\approx 1\%$, 9/1,103) and 1 bed bug ($\approx 1\%$, 1/160) tested positive in 2 *R. felis*-specific qRT-PCRs (Tables 1,4). The pRF δ plasmid was detected in 8 mosquito samples (21). In Dakar, 1% (2/154) of the *An. arabiensis* mosquitoes collected were positive for *R. felis*, including 1 male, suggesting transovarian transmission. One *Aedes luteocephalus* from Ferlo 0.5% (1/203) was positive for *R. felis*. In S_1 , 15% (6/40) mosquitoes collected in September 2012 were positive for *R. felis*, including 1 *An. ziemanni*, 1 *An. pharoensis*, 2 *Mansonia uniformis*, and 2 *An. funestus*. None of the 24 mosquito samples collected from this region in July tested positive. In addition, 1 *Cimex hemipterus* bed bug (3%) (1/39), collected from a household in S_1 in February 2012, tested positive. No *R. felis* DNA was detected in soft or hard ticks or in *Culicoides* species.

Discussion

This study shows that *Rickettsia felis* is an emerging pathogen commonly detected in sub-Saharan rural

Table 3. Age distribution of infections with *Rickettsia felis* and *Plasmodium* spp. among patients in Senegal, by age group, 2010–2012*

Species	No. positive samples/no. tested samples (%)						ND	Total
	<1 y	1–3 y	4–6 y	7–15 y	16–29 y	>30 y		
<i>Plasmodium</i> spp.	20/169 (12)	60/327 (16)	57/251 (23)	134/401 (33)	56/271 (21)	58/310 (19)	15/78 (19)	400/1,867 (21)
<i>R. felis</i>	27/184 (15)	40/422 (10)	40/270 (15)	69/425 (16)	49/298 (16)	57/336 (17)	30/89 (34)	312/2,024 (15)

*ND, no age data available.

Table 4. Detection of *Rickettsia* species in arthropods collected in Senegal, 2008–2012

Group	Species	No. samples tested	Type of rickettsia (% positive samples)	Reference
Fleas	<i>Ctenocephalides felis</i>	48	None	(13)
	<i>Echidnophaga gallinacea</i>	150	None	
	<i>Synosternus pallidus</i>	41	<i>Rickettsia</i> sp., group <i>R. felis</i> (93)	
Tsetse flies	<i>Glossina morsitans submorsitans</i>	78	<i>Rickettsia</i> sp., group <i>R. felis</i> (100)	(30)
Hard ticks	<i>Amblyomma variegatum</i>	492	<i>Rickettsia africae</i> (87)	(29)
	<i>Rhipicephalus decoloratus</i>	40	Rickettsiae spotted fever group	(27)
	<i>R. annulatus</i>	5	(0–51)	
	<i>Hyalomma marginatum rufipes</i>	173		
	<i>H. truncatum</i>	141		
	<i>R. evertsi evertsi</i>	2358		
	<i>R. guilhoni</i>	50		
	<i>Rhipicephalus</i> sp.	2	None	This study
Soft ticks	<i>Ornithodoros sonrai</i>	138	None	This study
	<i>O. capensis</i>	40	<i>Rickettsia</i> sp., group <i>R. felis</i> (20)	This study
	<i>Argas persicus</i>	4	None	This study
Midges	<i>Culicoides</i> spp.	384	None	This study
Bed bugs	<i>Cimex hemipterus</i>	160	1/160, (0.6)	This study

Africa. We are confident that our molecular results are reliable and that the negative results in samples from France illustrate a correlation between *R. felis* infection and malaria with regard to the geographic distribution and seasonality. A trend of higher risk for *R. felis* infection in southern countries than in northern countries was revealed; the highest risk for *R. felis* infection was in rural Senegal (24 times than in Algeria). In Senegal, DNA from *Plasmodium* spp. and *R. felis* were detected at high levels, mostly during the rainy season and among children <15 years of age (Figure 2), but no coincidental relationship was found. The incidence of co-infection of *R. felis* and malaria was lower in Senegal (23%) than in Kenya (79%) (5), but higher than the rate of simultaneous bacterial bloodstream infections and malaria parasitemia, which ranged from 6% in rural Mozambique (23) to 11% in Nairobi (24). Mixed infections for rickettsioses, including co-infections with malaria or with other bacteria (*Leptospira* spp., *Coxiella burnetii*, and *Burkholderia pseudomallei*) have been described (25).

R. felis was detected in afebrile persons, most of whom were children <15 years of age, confirming the previously reported results in Kenya (5). Although rickettsioses have not previously been reported in afebrile persons, low-grade *Plasmodium* parasitemia has been reported among persons without a fever (26). This result should be confirmed by culture, but *R. felis* has never been isolated, even from acutely ill patients. Nonetheless, the absence of positive tests in the control group located in France confirmed the specificity of our tests. The S₁₋₂ population was screened serologically for *R. felis*, and low titers were identified in 1 of 479 serum samples tested (27), which is substantially lower than the seroprevalence of other spotted fever group rickettsiae. The mechanism of absence of a serologic response and the occurrence of multiple re-infections or relapses of *R. felis* should be investigated further.

In this work, we demonstrated a greater frequency of *R. felis* during the rainy season among children in the subtropical zones, a period coinciding with circulation of *P. falciparum*. There are other seasonal diseases, including influenza, which are most common during the rainy season in subtropical Africa, particularly in Senegal (28). Influenza is a disease found throughout the year, with seasonal peaks, in Africa; none of the tested patients had influenza symptoms. Furthermore, leptospirosis, for which rickettsial disease could be mistaken, has not been documented in Senegal. Last, the most common seasonal disease in the most northern part of the intertropical area is malaria; a disease, however, which is common in all seasons in equatorial wetlands. These data, for which confirmation is needed, show a seasonal correlation between *R. felis* and malaria; the correlation is related to the presence and activity of *Anopheles* mosquitoes. Although the cat flea, *Ctenocephalides felis*, is currently the only known vector of *R. felis*, a variety of other arthropods have been suspected, including different flea species, ticks, mites, and lice (13). In Senegal, the source of *R. felis* is yet to be determined. We did not detect *R. felis* in fleas that were screened during 1 year in S₁ and S₂ (13). In other studies, *R. felis* was not detected in soft or hard ticks (27,29), tsetse flies (30), or midges. These findings support the hypothesis of the role of *Anopheles* in the transmission of *R. felis*; this hypothesis should be confirmed or refuted by future studies.

The clinical findings for *R. felis* infection are often unclear and are typically misdiagnosed as other febrile illnesses (12,31). Recently, the primary infection was described in a patient with polymorphous skin lesions, including papules, vesicles, erosions, and ulcers (12), similar to patients from Mexico (32). In the current study, a high incidence of *R. felis* infection was identified in children <15 years of age, as described (4). Fortunately, such patients improve rapidly with doxycycline treatment (12). For travelers to sub-Saharan

Africa, the medications recommended for the chemoprophylaxis of malaria include doxycycline, which has the added advantage of being effective against rickettsioses (33).

This study showed the wide distribution and high incidence of *R. felis* infection; therefore, rickettsiosis should be considered one of the major causes of febrile diseases in sub-Saharan Africa. The demonstrated geographic distribution, seasonality, target population, incidence of relapses or re-infections, and asymptomatic infections of *R. felis* infection are similar to malaria. Further studies are needed to investigate the hypotheses that humans, as for epidemic typhus, another vector-borne relapsing rickettsiosis, or apes could be reservoirs and mosquitoes could be a vector for *R. felis* infection.

Acknowledgments

We thank the villagers who participated in this study. We also thank Masse Sambou, Aliou Diallo, Khadim Leye, Babacar Ndao, Malick Diop, Arsène Mabika, Marielle Bedotto, Denis Pyak and Annick Bernard for technical support.

This study was funded by the Agence National de Recherche grant 2010 (MALEMAF), Fondation Méditerranée Infection, Fondation Mérieux, and a collaborative grant to Josselin Thuilliez, University of Paris, Paris, France.

Dr Mediannikov is an infectious disease specialist and research scientist working at the Unit of Research on Emergent Infectious and Tropical Diseases in Marseille, France and Dakar, Senegal. His main research interests include vector-borne diseases and medical entomology.

Dr Socolovschi is a physician of infectious diseases and tropical medicine at the Medical School of Marseille, France. Her research interests focus on vector-borne infectious tropical diseases and medical entomology.

References

- Parola P, Diatta G, Socolovschi C, Mediannikov O, Tall A, Bassene H, et al. Tick-borne relapsing fever borreliosis, rural Senegal. *Emerg Infect Dis.* 2011;17:883–5. <http://dx.doi.org/10.3201/eid1705.100573>
- Reddy EA, Shaw AV, Crump JA. Community-acquired bloodstream infections in Africa: a systematic review and meta-analysis. *Lancet Infect Dis.* 2010;10:417–32. [http://dx.doi.org/10.1016/S1473-3099\(10\)70072-4](http://dx.doi.org/10.1016/S1473-3099(10)70072-4)
- Fenollar F, Mediannikov O, Socolovschi C, Bassene H, Diatta G, Richet H, et al. *Tropheryma whipplei* bacteremia during fever in rural west Africa. *Clin Infect Dis.* 2010;51:515–21. <http://dx.doi.org/10.1086/655677>
- Socolovschi C, Mediannikov O, Sokhna C, Tall A, Diatta G, Bassene H, et al. *Rickettsia felis*-associated unruptive fever, Senegal. *Emerg Infect Dis.* 2010;16:1140–2. <http://dx.doi.org/10.3201/eid1607.100070>
- Maina AN, Knobel DL, Jiang J, Halliday J, Feikin DR, Cleaveland S, et al. *Rickettsia felis* infection in febrile patients, Western Kenya, 2007–2010. *Emerg Infect Dis.* 2012;18:328–31. <http://dx.doi.org/10.3201/eid1802.111372>
- Richards AL, Jiang J, Omulo S, Dare R, Abdurahman K, Ali A, et al. Human infection with *Rickettsia felis*, Kenya. *Emerg Infect Dis.* 2010;16:1081–6. <http://dx.doi.org/10.3201/eid1607.091885>
- Mediannikov O, Fenollar F, Socolovschi C, Diatta G, Sokhna C, Bassene H, et al. *Coxiella burnetii* in humans and ticks in rural Senegal. *PLoS Negl Trop Dis.* 2010;4:e654. <http://dx.doi.org/10.1371/journal.pntd.0000654>
- Reller ME, Clemens EG, Schachterle SE, Mtove GA, Sullivan DJ, Dumler JS. Multiplex 5' nuclease-quantitative PCR for diagnosis of relapsing fever in a large Tanzanian cohort. *J Clin Microbiol.* 2011;49:3245–9. <http://dx.doi.org/10.1128/JCM.00940-11>
- Freedman DO, Weld LH, Kozarsky PE, Fisk T, Robins R, von Sonnenburg F, et al. Spectrum of disease and relation to place of exposure among ill returned travelers. *N Engl J Med.* 2006;354:119–30. <http://dx.doi.org/10.1056/NEJMoa051331>
- Tourism Highlights United Nations World Tourism Organization [cited 2013 May 15]. http://mkt.unwto.org/sites/all/files/docpdf/unwtohighlights11enhr_3.pdf. 2011
- Wilson ME, Weld LH, Boggild A, Keystone JS, Kain KC, von Sonnenburg F, et al. Fever in returned travelers: results from the GeoSentinel Surveillance Network. *Clin Infect Dis.* 2007;44:1560–8. <http://dx.doi.org/10.1086/518173>
- Mediannikov O, Fenollar F, Bassene H, Tall A, Sokhna C, Trape JF, et al. Description of “yaaf”, the vesicular fever caused by acute *Rickettsia felis* infection in Senegal. *J Infect.* 2013;66:536–40. <http://dx.doi.org/10.1016/j.jinf.2012.10.005>
- Roucher C, Mediannikov O, Diatta G, Trape JF, Raoult D. A new *Rickettsia* species found in fleas collected from human dwellings and from domestic cats and dogs in Senegal. *Vector Borne Zoonotic Dis.* 2012;12:360–5. <http://dx.doi.org/10.1089/vbz.2011.0734>
- Raoult D, Roux V. Rickettsioses as paradigms of new or emerging infectious diseases. *Clin Microbiol Rev.* 1997;10:694–719.
- Verhulst NO, Smallegange RC, Takken W. Mosquitoes as potential bridge vectors of malaria parasites from non-human primates to humans. *Front Physiol.* 2012;3:197. <http://dx.doi.org/10.3389/fphys.2012.00197>
- Duval L, Fourment M, Nerrienet E, Rousset D, Sadeuh SA, Goodman SM, et al. African apes as reservoirs of *Plasmodium falciparum* and the origin and diversification of the *Laverania* subgenus. *Proc Natl Acad Sci U S A.* 2010;107:10561–6. <http://dx.doi.org/10.1073/pnas.1005435107>
- Socolovschi C, Pages F, Ndiath MO, Ratmanov P, Raoult D. *Rickettsia* species in African *Anopheles* mosquitoes. *PLoS ONE.* 2012;7:e48254. <http://dx.doi.org/10.1371/journal.pone.0048254>
- Socolovschi C, Pages F, Raoult D. *Rickettsia felis* in *Aedes albopictus* mosquitoes, Libreville, Gabon. *Emerg Infect Dis.* 2012;18:1687–9. <http://dx.doi.org/10.3201/eid1810.120178>
- Keita AK, Socolovschi C, Ahuka-Mundeye S, Ratmanov P, Butel C, Ayouba A, et al. Molecular evidence for the presence of *Rickettsia felis* in the feces of wild-living African apes. *PLoS ONE.* 2013;8:e54679. <http://dx.doi.org/10.1371/journal.pone.0054679>
- Trape JF, Rogier C, Konate L, Diagne N, Bouganali H, Canque B, et al. The Dielmo project: a longitudinal study of natural malaria infection and the mechanisms of protective immunity in a community living in a holoendemic area of Senegal. *Am J Trop Med Hyg.* 1994;51:123–37.
- Rolain JM, Bitam I, Buffet S, Marie JL, Bourry O, Portelli-Clerc C, et al. Presence or absence of plasmid in *Rickettsia felis* depending on the source of fleas. *Clin Microbiol Infect.* 2009;Suppl 2:296–7. <http://dx.doi.org/10.1111/j.1469-0691.2008.02245.x>
- Ndao M, Bandyayera E, Kokoskin E, Gyorkos TW, MacLean JD, Ward BJ. Comparison of blood smear, antigen detection, and nested-PCR methods for screening refugees from regions where malaria

- is endemic after a malaria outbreak in Quebec, Canada. *J Clin Microbiol.* 2004;42:2694–700. <http://dx.doi.org/10.1128/JCM.42.6.2694-2700.2004>
23. Sigaúque B, Roca A, Mandomando I, Morais L, Quinto L, Sacarlal J, et al. Community-acquired bacteremia among children admitted to a rural hospital in Mozambique. *Pediatr Infect Dis J.* 2009;28:108–13. <http://dx.doi.org/10.1097/INF.0b013e318187a87d>
 24. Okwara FN, Obimbo EM, Wafula EM, Murila FV. Bacteraemia, urinary tract infection and malaria in hospitalised febrile children in Nairobi: is there an association? *East Afr Med J.* 2004;81:47–51. <http://dx.doi.org/10.4314/eamj.v81i1.8795>
 25. Phommason K, Haris DH, Anantatat T, Castonguay-Vanier J, Keomany S, Souvannasing P, et al. Concurrent infection with murine typhus and scrub typhus in southern Laos—the mixed and the unmixed. *PLoS Negl Trop Dis.* 2013;7:e2163. <http://dx.doi.org/10.1371/journal.pntd.0002163>
 26. Roucher C, Rogier C, Dieye-Ba F, Sokhna C, Tall A, Trape JF. Changing malaria epidemiology and diagnostic criteria for *Plasmodium falciparum* clinical malaria. *PLoS ONE.* 2012;7:e46188. <http://dx.doi.org/10.1371/journal.pone.0046188>
 27. Mediannikov O, Diatta G, Fenollar F, Sokhna C, Trape JF, Raoult D. Tick-borne rickettsioses, neglected emerging diseases in rural Senegal. *PLoS Negl Trop Dis.* 2010;4. <http://dx.doi.org/10.1371/journal.pntd.0000821>
 28. Niang MN, Dosseh A, Ndiaye K, Sagna M, Gregory V, Goudiaby D, et al. Sentinel surveillance for influenza in Senegal, 1996–2009. *J Infect Dis.* 2012;206(Suppl 1):S129–35. <http://dx.doi.org/10.1093/infdis/jis576>
 29. Mediannikov O, Trape JF, Diatta G, Parola P, Fournier PE, Raoult D. *Rickettsia africae*, a neglected pathogen in West Africa. *Emerg Infect Dis.* 2010;16:571–3. <http://dx.doi.org/10.3201/eid1603.090346>
 30. Mediannikov O, Audoly G, Diatta G, Trape JF, Raoult D. New *Rickettsia* sp. in tsetse flies from Senegal. *Comp Immunol Microbiol Infect Dis.* 2012;35:145–50. <http://dx.doi.org/10.1016/j.cimid.2011.12.011>
 31. Abdad MY, Stenos J, Graves S. *Rickettsia felis*, an emerging flea-transmitted human pathogen. *Emerg Health Threats J.* 2011;4:7168 <http://dx.doi.org/10.3402/ehjt.v4i0.7168>
 32. Zavala-Velázquez JE, Sosa-Ruiz JA, Zavala-Castro J, Jimenez-Delgado B, Vado-Solis IE, Sanchez-Elias RA, et al. *Rickettsia felis*—the etiologic agent of three cases of rickettsiosis in Yucatan. *Lancet.* 2000;356:1079–80. [http://dx.doi.org/10.1016/S0140-6736\(00\)02735-5](http://dx.doi.org/10.1016/S0140-6736(00)02735-5)
 33. Twartz JC, Shirai A, Selvaraju G, Saunders JP, Huxsoll DL, Groves MG. Doxycycline prophylaxis for human scrub typhus. *J Infect Dis.* 1982;146:811–8. <http://dx.doi.org/10.1093/infdis/146.6.811>

Address for correspondence: Didier Raoult, Université Aix-Marseille, URMITE, UMR CNRS 7278, IRD 198, INSERM 1095, Faculté de Médecine, 27 Bd Jean Moulin, 13385 Marseille Cedex 5 France; email: didier.raoult@gmail.com

The image shows a screenshot of the CDC's Facebook page. At the top, there is a large banner for the 'Solve the Outbreak' iPad app, featuring a map and the text 'CDC is on Facebook. To connect with CDC, sign up for Facebook today.' Below the banner, the page header shows 'CDC' with 263,397 likes and 3,144 talking about this. The 'About' section states: 'Government Organization. CDC is dedicated to protecting health & promoting quality of life through prevention and control of disease, injury, and disability. For official CDC info go to www.cdc.gov.' There are also links to 'Photos', 'Likes' (263k), 'Vital Signs', and 'Welcome'. The main feed shows a post from CDC: '#Heatwave safety tip: Muscle cramping might be the first sign of heat-related illness, and may lead to heat exhaustion or stroke. Learn how to recognize heat exhaustion and heat stroke and know what to do: Extreme Heat and Your Warning Signs and Symptoms of Heat Illness'. Another post from Carol Ferguson asks 'Did you know your ads are being aired on the Rush Limbaugh...'. A post from Thomas Roles says 'Thanks to the Freedom of Information Act (FOIA), we know...'. At the bottom, there is a large text overlay: 'Find emerging infectious disease information on facebook' with the URL 'http://www.facebook.com'.

Common Epidemiology of *Rickettsia felis* Infection and Malaria, Africa

Technical Appendix

Technical Appendix Table. Rural health centers and laboratories that participated in study of epidemiology of *Rickettsia felis* Infection and malaria, featuring the demography, geography, and climate of the study regions*

Country and study site (site abbreviation)	Region/department/district	Population size	Climate,vegetation	Annual precipitation, mm	Village	Coordinates	Other activities
Senegal Dielmo-Ndiop (S ₁ -S ₂)	Toubacouta/Foundiougne/Fatick	120,554	Sudanian†, wooded savannah	939	Dielmo	13°43'N, 16°24'W	Point-of-care laboratory
					Ndiop	13°41'N, 16°23'W	None
Keur Momar Sarr (S ₃)	Keur Momar Sarr/Louga/Louga	70,743	Sahelian‡,steppe-type	400	Keur Momar Sarr	15°55'N, 15°58'W	DNA extraction
					Loboudou	15°57'N, 15°55'W	None
					Ganket Balla	15°58'N, 15°55'W	None
Niakhar (S ₄)	Niakhar/Fatick/Fatick	69,446	Sahelo-Sudanian§, wooded steppe	757	Toucar	14°32'N, 16°28'W	None
					Diohine	14°30'N, 16°30'W	None
					Ngayokheme	14°32'N, 16°26'W	None
					Niakhar	14°28'N, 16°23'W	DNA extraction
Basse Casamance (S ₅)	Loudia-Oulof/Oussouye/Ziguinchor	57,505	Sub-Guinean¶, primary and secondary gallery forests	1,432	Mlomp	12°33'N, 16°34'W	DNA extraction
					Kagnout	12°33'N, 16°37'W	None
					Elinkine	12°30'N, 16°39'W	None
Kedougou (S ₆)	Bandafassi/Kedougou/Kedougou	20,021	Sudano-Guinean#, woodland, wooded savannah	1,189	Bandafassi	12°32'N, 12°18'W	DNA extraction
					Ibel	12°30'N, 12°22'W	None
					Tiabeji	12°38'N, 12°25'W	None

Country and study site (site abbreviation)	Region/department/district	Population size	Climate,vegetation	Annual precipitation, mm	Village	Coordinates	Other activities
Mali Diankambou (M ₁)	Mopti	11,333	Sahelian, savannah	580	–	14°35'N; 3°05'W	DNA extraction
Kole–M ₂	Sikasso	3,508	Sudano-Guinean, savannahs, forest	700–1,500	–	10°5'N; 7°20'W	DNA extraction
Gabon Franceville (G)	Mpassa/Haut-Ogooué	56,000	Equatorial**, savannah and tropical forest	1,862	–	1°37'S; 13°34'E	DNA extraction
Algeria Oran	Oran Province	18,682,000	Semi-arid climate, halophytic vegetation	326	–	35°41'N 0°37'W	None
Morocco Kenitra	Gharb-Chrarda-Béni Hssen	1,859,540	Mild Mediterranean climate, woody flora	750	Sidi Mohamed Lahmar Sidi Taybi	34°15'N 6°35'W	None
Tunisia Sfax	Sfax Governorate	2,256,320	Mediterranean climate, wild grasses	465	–	36°48'N 10°11'E	None
France Marseille	Provence-Alpes-Côte d'Azur	1,582,000	Mediterranean climate, Mediterranean maquis shrubland††	515	–	43°17'N 5°22'E	DNA extraction

*–, no participants at village level; NA, not applicable.

†Sudanian, tropical semi-arid climate with a mean annual precipitation between around 600 and 800 mm.

‡Sahelian, tropical semi-arid climate with a dry season mainly during the coldest 6 months of the year and a mean annual precipitation ≈200–400 mm.

§Sahelo-Sudanian tropical semi-arid climate with a mean annual precipitation ≈400–600 mm.

¶Sub-Guinean, tropical monsoon maritime climate characterized by the changing of dry and wet seasons; mean annual precipitation rate ≈1200–1500 mm.

#Sudano-Guinean, transitory climate between Sudanian and sub-Guinean climates; annual precipitation ≈ 800–1200 mm.

**Equatorial, a climate with no dry season; mean precipitation values of all months are ≥60 mm.

††Maquis, a shrubland biome in the Mediterranean region, typically consisting of densely growing evergreen shrubs such as holm oak, Kermes Oak, tree heath, strawberry tree, sage, juniper, buckthorn, spurge olive. and myrtle.