

11. Estrada-Peña A, Martínez M, Muñoz MJ. A population model to describe the distribution and seasonal dynamics of the tick *Hyalomma marginatum* in the Mediterranean basin. *Transbound Emerg Dis*. 2011;58:213–23. <http://dx.doi.org/10.1111/j.1865-1682.2010.01198.x>
12. Rumer L, Graser E, Hillebrand T, Talaska T, Dautel H, Mediannikov O, et al. *Rickettsia aeschlimannii* in *Hyalomma marginatum* ticks, Germany. *Emerg Infect Dis*. 2011;17:325–6. <http://dx.doi.org/10.3201/eid1702.100308>
13. Jameson LJ, Morgan PJ, Medlock JM, Watola G, Vaux AG. Importation of *Hyalomma marginatum*, vector of Crimean-Congo haemorrhagic fever virus, into the United Kingdom by migratory birds. *Ticks Tick Borne Dis*. 2012;3:95–9. <http://dx.doi.org/10.1016/j.ttbdis.2011.12.002>
14. Schmaljohann H, Liechti F, Bruderer B. Songbird migration across the Sahara: the non-stop hypothesis rejected! *Proc Biol Sci*. 2007;274:735–9. <http://dx.doi.org/10.1098/rspb.2006.0011>
15. Gale P, Stephenson B, Brouwer A, Martínez M, de la Torre A, Bosch J, et al. Impact of climate change on risk of incursion of Crimean-Congo haemorrhagic fever virus in livestock in Europe through migratory birds. *J Appl Microbiol*. 2012;112:246–57. <http://dx.doi.org/10.1111/j.1365-2672.2011.05203.x>

Address for correspondence: José A. Oteo, Área de Enfermedades Infecciosas, Hospital San Pedro–CIBIR, C/Piqueras 98-7ª NE, 26006 Logroño (La Rioja), Spain; email: jaoteo@riojasalud.es

Have you renewed your
print subscription to

**EMERGING
INFECTIOUS DISEASES®**



Subscribe or renew now at
<http://wwwnc.cdc.gov/eid/subscribe.htm>
and have the print issue delivered

etymologia

Quarantine

[kwor'ən-tēn]

From the Italian quaranta (forty), “quarantine” refers to the practice established in European port cities during the Black Death requiring vessels to lie at anchor for 40 days before landing. Isolation (from the Latin insula or island), the practice of separating sick persons from those who are healthy to prevent spread of disease, goes back a long time. “As long as they have the disease they remain unclean. They must live alone; they must live outside the camp” (Leviticus 13:46).

Quarantine, on the other hand, is the practice of separating persons who appear to be healthy but may have been exposed to a disease. In 1377, the Great Council of Ragusa (modern-day Dubrovnik) established a 30-day separation period (trentino) for visitors from plague-endemic areas. In the following decades, the practice spread to other cities and was extended from 30 to 40 days (quarantino). The longer period may have been more effective at preventing disease or just a nod to the 40-day duration of Biblical events—the Great Flood or Jesus’ fast in the wilderness.

Sources

1. Dorland’s Illustrated Medical Dictionary. 32nd ed. Philadelphia: Elsevier Saunders; 2012.
2. Holy Bible, New International Version. Colorado Springs (CO): Biblica; 2011.
3. Sehdev PS. The origin of quarantine. *Clin Infect Dis*. 2002;35:1071–2. PubMed <http://dx.doi.org/10.1086/344062>

Address for correspondence: Ronnie Henry, Centers for Disease Control and Prevention, 1600 Clifton Rd NE, Mailstop E03, Atlanta, GA 30333, USA; email: boq3@cdc.gov

DOI: <http://dx.doi.org/eid1902.ET1902>

Medscape Sign up to receive email announcements when
CME™ a new article is available.

Get an online subscription at wwwnc.cdc.gov/eid/subscribe.htm