

The *bdr* Gene Families of the Lyme Disease and Relapsing Fever Spirochetes: Potential Influence on Biology, Pathogenesis, and Evolution

David M. Roberts,* Jason A Carlyon,† Michael Theisen,‡
and Richard T. Marconi*

*Medical College of Virginia at Virginia Commonwealth University, School of Medicine, Richmond, Virginia, USA; †Yale School of Medicine, Yale University, New Haven, Connecticut, USA; and ‡Statens Serum Institute, Copenhagen, Denmark

Species of the genus *Borrelia* cause human and animal infections, including Lyme disease, relapsing fever, and epizootic bovine abortion. The borrelial genome is unique among bacterial genomes in that it is composed of a linear chromosome and a series of linear and circular plasmids. The plasmids exhibit significant genetic redundancy and carry 175 paralogous gene families, most of unknown function. Homologous alleles on different plasmids could influence the organization and evolution of the *Borrelia* genome by serving as foci for interplasmid homologous recombination. The plasmid-carried *Borrelia direct repeat (bdr)* gene family encodes polymorphic, acidic proteins with putative phosphorylation sites and transmembrane domains. These proteins may play regulatory roles in *Borrelia*. We describe recent progress in the characterization of the *Borrelia bdr* genes and discuss the possible influence of this gene family on the biology, pathogenesis, and evolution of the *Borrelia* genome.

Species of the genus *Borrelia* cause human and animal infections (1). In North America, Lyme disease and endemic relapsing fever pose the greatest threat to human health and have received the most attention of the borrelial diseases. Approximately 14,000 cases of Lyme disease are reported in the United States each year; however, the actual number of cases may be 10-fold higher (2). Lyme disease was not recognized as a distinct clinical entity in North America until the 1970s (3). The causative agent, a previously uncharacterized spirochete transmitted through the bite of infected ticks of the *Ixodes ricinus* complex (*I. scapularis* in the Northeast and Midwest and *I. pacificus* on the West Coast) (4,5), was classified in the genus *Borrelia* and named *B. burgdorferi*. With the emergence of Lyme disease and the identification of its etiologic agent, *Borrelia* research focused on

the development of reliable Lyme disease diagnostic assays and vaccines, and the phenotypic and genotypic diversity of *Borrelia* was thoroughly analyzed. Through modern molecular taxonomic techniques, several newly described species of *Borrelia* have emerged as possible causative agents of Lyme disease or at least as agents genetically related to *B. burgdorferi* (6-15). The *B. burgdorferi* sensu lato complex is composed of the following species: *B. turdae*, *B. tanukii*, *B. bissettii*, *B. valaisiana*, *B. lusitaniae*, *B. bissettii*, *B. andersonii*, *B. japonica*, *B. garinii*, and *B. afzelii*. Of these, *B. burgdorferi*, *B. garinii*, and *B. afzelii* are the dominant species associated with infection in humans.

Relapsing fever has been studied not only for its impact on human health but also as a model system for antigenic variation. There are two general forms of relapsing fever, epidemic (louse borne—*Pediculus humanus*) and endemic (tick borne—*Ornithodoros* spp.) (1). Epidemic relapsing fever tends to be associated with poor living conditions and social disruption (famine and war)

Address for correspondence: Richard T. Marconi, Department of Microbiology and Immunology, Medical College of Virginia at VCU, Richmond, VA 23298-0678, USA; fax: 804-828-9946; e-mail: Rmarconi@hsc.vcu.edu.

Synopses

and is rare in the United States. Endemic relapsing fever is more prevalent, predominantly in the western regions. Three closely related *Borrelia* species, *B. hermsii*, *B. turicatae*, and *B. parkeri*, are associated with this disease. Hallmark features of relapsing fever include cyclic fever and spirochetemia. The molecular basis for these features can be attributed to the differential production of dominant variable surface antigens of the Vmp protein families (16). The 40 or so plasmid-carried *vmp* related genes in the *B. hermsii* genome are expressed only one at a time. A single expression locus exists, and genes not at this site lack a promoter element and are therefore not transcribed (17). The expressed Vmp becomes a primary target of a vigorous humoral immune response that kills most of the spirochetal population. However, at a frequency of approximately of 1×10^{-3} to 1×10^{-4} per generation, the identity of the expressed Vmp changes (18) through gene conversion (19). The net effect of this nonreciprocal event is to replace the gene located in the expression locus with one that was previously silent. Production of a new antigenically distinct Vmp allows evasion of the humoral immune response. This ongoing change in Vmp synthesis allows the relapsing fever spirochete population to reestablish itself in the host, thus leading to spirochetemia and the relapse of fever. Antigenic variation systems have also been identified in the Lyme disease spirochetes; however, they appear to exert a more subtle effect (20).

While clinical relapsing fever and Lyme disease differ from each other in many ways, their causative agents share many similarities at both the biologic and genetic levels. At the biologic level, they are host associated and undergo similar environmental transitions in the course of cycling between mammals and arthropods. In view of the distinctly different characteristics of these environments, the spirochetes must be able to adapt rapidly. Evidence suggests that the relapsing fever and Lyme disease spirochetes use related proteins to adapt to or carry out similar functions in changing environments. For example, homologs of the plasmid-carried *ospC* gene of the Lyme disease spirochetes are carried by several other *Borrelia* species, including the relapsing fever spirochetes (21). Both *ospC* and its relapsing fever spirochete homolog (*vmp33*) are selectively expressed during the early stages of infection, which suggests that they play a

common functional role (22,23). The *B. burgdorferi* Rep or Bdr protein family is also distributed genuswide. Members of this polymorphic protein family possess highly conserved putative functional motifs and structural properties, which suggests that they may also carry out an important genuswide role (24,25).

The *Borrelia* Genome

At the molecular level, a unique feature of *Borrelia* is the unusual organization and structure of their genome. Unlike most bacteria, which carry their genetic material in the form of a single, circular DNA molecule, *Borrelia* have a segmented genome (26-28). Most genetic elements carried by these bacteria are linear with covalently closed termini or telomeres (27). The telomeres are characterized by short hairpin loops of DNA (29). If heat denatured, these linear molecules relax to form a single-stranded circular molecule. If reannealed, they base-pair upon themselves to form a double-stranded linear molecule that by physical necessity possesses a short single-stranded hairpin loop at each telomere. Genetic elements of this structure are rare in bacteria and are reminiscent of certain viral genomes. In *B. burgdorferi* (isolate B31), the largest of the linear genomic elements is the 911-kb chromosome (30). The chromosome carries 853 putative ORFs, most of which are thought to encode housekeeping functions. The remaining 12 linear and 8 circular genetic elements are plasmids. The plasmids might best be thought of as mini-chromosomes, since as a group they are indispensable in situ and may carry genes encoding proteins involved in housekeeping functions (31). In addition, they may further deviate from the true definition of a plasmid in that their replication may not be independent and may instead be tightly coordinated with the replication of the chromosome (32,33).

Nearly 50% of the plasmid-carried ORFs lack homology with known sequences, which suggests that their encoded proteins may define the unique biologic and pathogenetic aspects of *Borrelia* (30). Several of the proteins derived from these plasmid-carried genes of unknown function are antigenic or selectively expressed during infection, which indicates that they function in the mammalian environment (20,34-37). A striking feature of the plasmid-carried ORFs is that they are organized into 175 paralogous gene families of two or more members (30). Hence, the

Synopses

DNA content of the plasmids is highly redundant. Since the maintenance of DNA is energetically expensive, it is likely that this redundant DNA is of biologic importance to *Borrelia*. The paralogous gene families of *Borrelia* have been the focus of intensive research as they are thought to play important roles in pathogenesis and to influence genome organization and evolution (20,30,35,38-40).

Identification of *Borrelia* Direct Repeat (*bdr*) Related Genes

The *bdr* gene family is a large, polymorphic, plasmid-carried, paralogous gene family of unknown function that was originally identified in *B. burgdorferi* (41,42). Members of this gene family have been characterized in several *Borrelia* species and isolates (Table 1) and have

Table 1. *Borrelia* species carrying *bdr*-related genes or expressing proteins immunoreactive with anti-Bdr antisera

Species	Associated disease	Arthropod vector	<i>Bdr</i> -related information
<i>B. burgdorferi</i>	Lyme disease, endemic worldwide	<i>I. scapularis</i> , <i>I. ricinus</i> , <i>I. pacificus</i>	All <i>bdr</i> -gene family members (18 total) have been identified in isolate B31G (30), <i>bdr</i> -alleles that are organized into 3 subfamilies (D,E,F) (25), the genes are carried on variably sized linear and circular plasmids (30) Single <i>bdr</i> gene has been sequenced (43); several <i>Bdr</i> -related proteins have been detected by immunoblot analysis (this report)
<i>B. afzelii</i>	Lyme disease, Eurasia	<i>I. ricinus</i> , <i>I. persulcatus</i>	Bdr proteins detected by immunoblot analyses only (this report)
<i>B. garinii</i>	Lyme disease, Eurasia	<i>I. ricinus</i> , <i>I. persulcatus</i>	Bdr proteins detected by immunoblot analyses only (this report)
<i>B. tanukii</i>	Not associated with human disease	<i>I. tanuki</i>	Bdr proteins detected by immunoblot analyses only (this report)
<i>B. turdae</i>	Not associated with human disease	<i>I. turdus</i>	Bdr proteins detected by immunoblot analyses only (this report)
<i>B. bissettii</i>	Not associated with human disease	<i>I. pacificus</i> , <i>I. scapularis</i> , <i>I. spinipalpis</i>	Bdr proteins detected by immunoblot analyses only (this report)
<i>B. andersonii</i>	Not associated with human disease	<i>I. dentatus</i>	<i>bdr</i> genes detected by hybridization and Bdr-related proteins by immunoblot analyses (this report, data not shown)
<i>B. valaisiana</i>	Not associated with human disease	<i>I. columnae</i> , <i>I. ricinus</i>	Bdr proteins detected by immunoblot analyses only (this report)
<i>B. japonica</i>	Not associated with human disease	<i>I. ovatus</i>	Bdr proteins detected by immunoblot analyses only (this report)
<i>B. hermsii</i>	Endemic relapsing fever, United States	<i>Onithodoros hermsii</i>	Numerous <i>bdr</i> genes have been described and are carried on both linear and circular plasmids (25,46); several Bdr proteins have been detected by immunoblot analyses (this report) (44)
<i>B. parkeri</i>	Endemic relapsing fever, United States	<i>Onithodoros parkeri</i>	Two <i>bdr</i> -related genes have been sequenced (25), and others have been detected by hybridization with genes residing on both linear and circular plasmids (46); several Bdr proteins detected by immunoblot analyses (this report)
<i>B. turicatae</i>	Endemic relapsing fever, United States	<i>O. turicata</i>	At least nine <i>bdr</i> -related genes have been described and are present on linear plasmids ranging from 25 to 220 kb in size (24,46); several Bdr proteins have been detected by immunoblotting (this report) (24,44)
<i>B. miyamotoi</i>	Relapsing fever?	<i>I. persulcatus</i>	Bdr proteins detected by immunoblotting only (this report)
<i>B. coriaceae</i>	Epizootic bovine abortion, United States	<i>O. coriaceus</i>	<i>bdr</i> -related genes and proteins detected by hybridization (46) or immunoblotting (this report)
<i>B. anserina</i>	Avian spirochetosis, United States	<i>Argas persicus</i>	<i>bdr</i> -related sequences have been detected by hybridization (46); Bdr-related proteins were not detected in <i>in vitro</i> cultivated bacteria (this report)

Synopsis

been assigned various gene names (25,41-44) (Table 2). We have adopted the *bdr* designation in the context of a nomenclature system (25), summarized below. Genes belonging to the *bdr* gene family were first identified through the analysis of repeated DNA sequences in *B. burgdorferi* sensu lato complex isolates (41,42). Seven nonidentical but closely related copies of a plasmid-carried repeated element were identified in *B. burgdorferi* 297 (42). Three additional copies of this repeated sequence were further

identified in *B. burgdorferi* 297 (45). These loci carry several ORFs that were designated as *rep+*, *rep-*, LPA, LPB (the LP genes have recently been redesignated as *mlp* for multicopy lipoprotein [45]), *rev*, and the *orfABCD* operon (note: ORFs A and B have been redesignated as *blyA* and *blyB*). Some of these genes, particularly *rep* and *mlp*, exhibit allelic variation and encode polymorphic proteins, the functions of which are under investigation. Focusing specifically on the *rep* or *bdr* genes, the *rep* designation was originally

Table 2. *Borrelia* Bdr homology groups and gene nomenclature

<i>Bdr</i> subfamily designation	Species/ revised gene designation	Accession or TIGR number	Previous gene names	Ref.
Subfamily A				
<i>B. turicatae</i> OZ-1	<i>bdrA</i> ₁	AF062395	<i>repA</i>	(46)
<i>B. turicatae</i> OZ-1	<i>bdrA</i> ₂ , <i>A</i> ₃ , <i>A</i> ₄	AF128445-AF128447	none	(25)
<i>B. hermsii</i> YOR-1	<i>bdrA</i> ₁ , <i>A</i> ₂ , <i>A</i> ₃	AF143473-AF143475	none	(25)
<i>B. hermsii</i> HS1	<i>bdrA</i> ₁ , <i>A</i> ₂	AF143457-AF143458	none	(25)
<i>B. hermsii</i> MAN	<i>bdrA</i> ₁ , <i>A</i> ₂	AF143465, AF143467	none	(25)
<i>B. parkeri</i>	<i>bdrA</i> ₁	AF143455	none	(25)
Subfamily B				
<i>B. turicatae</i> OZ-1	<i>bdrB</i> ₁ , <i>B</i> ₂ , <i>B</i> ₃ , <i>B</i> ₄ , <i>B</i> ₅	AF128448-AF128452	none	(24)
<i>B. hermsii</i> MAN	<i>bdrB</i> ₁ , <i>B</i> ₂ , <i>B</i> ₃	AF143463, AF143464, AF143466	none	(25)
Subfamily C				
<i>B. parkeri</i>	<i>bdrC</i> ₁	AF143455	none	(25)
<i>B. hermsii</i> MAN	<i>bdrC</i> ₁ , <i>C</i> ₂ , <i>C</i> ₃ , <i>C</i> ₄ , <i>C</i> ₅	AF143468-AF143472	none	(25)
<i>B. hermsii</i> HS1	<i>bdrC</i> ₁ , <i>C</i> ₂ , <i>C</i> ₃ , <i>C</i> ₄	AF143459-AF143462	none	(25)
<i>B. hermsii</i> YOR-1	<i>bdrC</i> ₁	AF143476	none	(25)
<i>B. parkeri</i>	<i>bdrC</i> ₂	AF143456	none	(25)
Subfamily D				
<i>B. burgdorferi</i> B31G	<i>bdrD</i> ₁ , <i>D</i> ₂ , <i>D</i> ₃	BBL35, BBM34, BBO34	<i>bdrO</i> , <i>bdrK</i> , <i>bdrM</i>	(30)
<i>B. burgdorferi</i> B31G	<i>bdrD</i> ₄ , <i>D</i> ₅ , <i>D</i> ₆	BBP34, BBQ42, BBS37	<i>bdrA</i> , <i>bdrV</i> , <i>bdrE</i>	(30)
<i>B. burgdorferi</i> B31	<i>bdrD</i> ₇	X87201	ORF-E (lp50 allele)	(41)
<i>B. burgdorferi</i> B31	<i>bdrD</i> ₈	X87127	ORF-E (cp30.5 allele)	(41)
<i>B. burgdorferi</i> B31	<i>bdrD</i> ₉	U42599	ORF-E (cp18 allele)	(41)
<i>B. burgdorferi</i> B31	<i>bdrD</i> ₁₀	BBN34	<i>bdrQ</i>	(30)
<i>B. burgdorferi</i> B31	<i>bdrD</i> ₁₁	BBR35	<i>bdrG</i>	(30)
Subfamily E				
<i>B. burgdorferi</i> B31G	<i>bdrE</i> ₁ , <i>E</i> ₂ , <i>E</i> ₃	BBL27, BBN27, BBO27	<i>bdrP</i> , none, <i>bdrN</i>	(30)
<i>B. burgdorferi</i> B31G	<i>bdrE</i> ₄ , <i>E</i> ₅ , <i>E</i> ₆	BBR27, BBS29, BBQ34	<i>bdrH</i> , <i>bdrF</i> , <i>bdrW</i>	(30)
<i>B. burgdorferi</i> 297	<i>bdrE</i> ₁ , <i>E</i> ₂	U45421, U45422	<i>rep+2.9-1</i> , <i>rep+2.9-2</i>	(42)
<i>B. burgdorferi</i> 297	<i>bdrE</i> ₃ , <i>E</i> ₄	U45423, U45424	<i>rep+2.9-3</i> , <i>rep+2.9-4</i>	(42)
<i>B. burgdorferi</i> 297	<i>bdrE</i> ₅	U45425	<i>rep+2.9-5</i>	(42)
<i>B. burgdorferi</i> 297	<i>bdrE</i> ₆	AF046998	<i>rep+2.9-8</i>	(45)
<i>B. burgdorferi</i> 297	<i>bdrE</i> ₇	AF046999	<i>rep+2.9-9</i>	(45)
Subfamily F				
<i>B. afzelii</i> DK1	<i>bdrF</i> ₁	Y08143	<i>p21</i>	(43)
<i>B. burgdorferi</i> B31G	<i>bdrF</i> ₁ , <i>F</i> ₂ , <i>F</i> ₃	BBF03, BBG33, BBH13	<i>bdrS</i> , <i>bdrT</i> , <i>bdrU</i>	(30)

Synopses

chosen to reflect a central repeat motif carrying domains in the deduced amino acid sequences. The + and - designations were assigned to indicate that the overlapping *rep+* and *rep-* genes are located on opposing DNA strands. Plasmid-carried repeated DNA sequences were also identified in *B. burgdorferi* B31 and found to carry either all or a subset of seven ORFs, designated A through G (41). Of relevance to this discussion are the ORF-E sequences that are *rep* or *bdr* homologs. A *bdr*-related gene was also identified in *B. afzelii* DK1 and designated as *p21* (43). *B. afzelii* causes Lyme disease in Europe and Asia. The *rep+*, ORF-E, and *p21* designations have recently been replaced with *bdr* gene designations (24,25,44).

To assess and compare the composition and complexity of the *bdr* gene family among species and isolates of the *B. burgdorferi* sensu lato complex, restriction fragment length polymorphism (RFLP) patterns were determined (Appendix). Genomic DNA digested with *Xba*I was Southern blotted and probed with an oligonucleotide targeting the *bdr* genes (Figure 1). A variable number of hybridizing bands of different size were detected. These analyses demonstrate that extensive *bdr* gene families are carried by *B. burgdorferi* sensu lato complex isolates and that the RFLP patterns vary at the inter- and intraspecies level. Hybridization analyses of other *Borrelia* species showed that they also carry *bdr*-related gene families (24,25,46). *bdr*-related genes have been detected by hybridization in *B. turicatae*, *B. hermsii*, *B. parkeri*, *B. coriaceae*, and *B. anserina* (25,46). Isolates of these species also exhibit substantial variation in their *bdr* RFLP patterns at the intraspecies level. Table 1 lists the *Borrelia* species that carry *bdr*-related genes and indicates the methods by which these genes or proteins were detected.

Sequences flanking some *bdr* alleles also appear to be distributed genus wide. Some *bdr* alleles of *B. turicatae*, *B. parkeri*, and *B. hermsii* are flanked by genes that are homologs of genes carried by the Lyme disease spirochetes (24,25). As a specific example, the *B. turicatae* *bdrA*₁ gene is flanked by ORFs that are homologs of the BBG34 and BBG30 genes of *B. burgdorferi* (24,25). In the Lyme disease spirochetes, BBG34 is part of a three-member paralogous gene family, while BBG30 is a single-copy gene (30). Located between BBG30 and BBG34 is BBG33, a member of the *bdr* gene family (recently redesignated as




Figure 1. Restriction fragment length polymorphism pattern analysis of the *rep* or *bdr* genes of the Lyme disease spirochetes. Total DNA, isolated from *Borrelia* cultures, was digested with *Xba*I, fractionated by electrophoresis, and transferred onto membranes for hybridization. Hybridization was performed by the *bdr*AB-R1 oligonucleotide (46). The species and isolates analyzed are indicated above each lane. MW markers in kb are indicated.

*bdrF*₂) (25). Although these divergent *Borrelia* species carry related genes, their organization differs (24), which indicates that rearrangement has taken place in the ancestral plasmid that carried these homologs. Figure 2 compares the

Synopses




Figure 2. General organization of two *bdr* loci in *Borrelia turicatae* and *B. burgdorferi*. The gene arrangement depicted for *B. turicatae* was determined through cloning and sequence analysis of a 2,217 base-pair *Xba*I restriction fragment. The arrangement for the *bdr*-carrying locus of *B. burgdorferi* was previously determined through the sequencing of the *B. burgdorferi* B31 genome (30). The arrows indicate the direction of transcription. Genes exhibiting homology are indicated by similar shading or hatch marks. Genes indicated by unfilled arrows are not homologous. The numbering is indicated for scale and is not indicative of the positioning of these genes on the plasmids that carry them.

organization of two *bdr* loci from *B. turicatae* and *B. burgdorferi*.

Evolutionary Analyses of *bdr*-Related Sequences: Revised Nomenclature for the Bdr-Related Proteins

To simplify the complicated nomenclature of *bdr*-related genes, a *bdr* nomenclature system has been developed that assigns gene names on the basis of phylogenetic relationships inferred from comparative analysis of genetically stable regions of the *bdr* genes (25). This system, which is applicable genuswide, allows for a ready assessment of relationships among *bdr* paralogs and orthologs. The rationale for this system stemmed from the results of a comprehensive evolutionary analysis of >50 *bdr*-related sequences from five *Borrelia* species that demonstrated that *bdr* sequences are organized into six distinct subfamilies, designated A through F (25). Subfamilies are not necessarily species specific; some contain *bdr* alleles from different *Borrelia* species (25). Since members of a given subfamily are closely related to one another with identity values for the N terminal domain being >95%, each member is assigned the same gene name

designation, and paralogs are distinguished by a numerical subscript. In *B. turicatae* OZ-1, two *bdr* subfamilies, *bdrA* and *bdrB*, contain at least four and five members, respectively (24). Members of the *bdrA* subfamily are designated *bdrA*₁, *bdrA*₂, *bdrA*₃, and *bdrA*₄, while members of the *bdrB* family are designated *bdrB*₁ through *bdrB*₅. This revised Bdr nomenclature scheme was modeled after that proposed for bacterial polysaccharide synthesis genes (47) and is in accordance with the nomenclature guidelines established by Demerec (48).

The subfamily affiliation of *bdr* genes can be readily determined through comparative sequence analyses of the amino acid segment preceding the polymorphic repeat motif region of these proteins (described in detail below) (25). Relationship assessments based on the genetically stable N terminal domain (vs. complete sequences) are preferable because the calculated evolutionary distances and clustering relationships are not artificially skewed by the variable number of repeat motifs present in the repeat motif domain. Since the genetically unstable repeat motif domain comprises as much as 50% of the total coding sequence in some alleles, it can

Synopses

have a substantial impact on inferred relationships. In addition, extensive sequence variation in the carboxyl termini of the Bdr proteins at the inter-species level makes it difficult to align this domain with confidence, which further influences the inferred relationships.

bdr evolutionary analyses show that *Borrelia* species carry members of at least two *bdr* subfamilies (25,44). In fact, *B. burgdorferi* carries three distinct subfamilies. Multiple Bdr subfamilies in diverse *Borrelia* species suggest that there has been selective pressure to maintain multiple *bdr* alleles and *bdr* genetic diversity. This genetic diversity may increase the functional diversity of the Bdr proteins.

Molecular Features and Physical Properties of the Bdr Proteins

While early analyses of *Borrelia bdr* genes demonstrated their multicopy nature (41,42,46), the full extent of the complexity of the *bdr* gene family in the Lyme disease spirochetes was not fully recognized until the *B. burgdorferi* genome sequence was determined (30). *B. burgdorferi* B31 was found to carry 17 distinct *bdr*-related genes (and one truncated variant) distributed among different linear and circular plasmids. *B. turicatae*, which carries at least nine different *bdr* alleles, carries these genes exclusively on linear plasmids (24,25,46). Other relapsing fever spirochete species (*B. parkeri* and *B. hermsii*) are similar to the Lyme disease bacteria in that they carry *bdr* genes on both linear and circular

plasmids (25). In the Lyme disease spirochetes each of the 32-kb circular plasmids, with the exception of plasmids M and P, carry two different *bdr* genes separated by seven or eight ORFs. Each of these circular plasmids carries one *bdrD* subfamily member and one *bdrE* subfamily member. The maintenance of genes belonging to different subfamilies on a single plasmid is consistent with the possibility that each carries out a different function. In contrast, in the Lyme disease spirochetes, the *bdrF* subfamily members are localized to linear plasmids with only a single *bdr* gene per plasmid. These observations suggest that there has been selective pressure to maintain the association of specific subfamilies with specific types of plasmids. Less is known about the *bdr*-carrying plasmids and the organization of the *bdr* genes and subfamilies in the relapsing fever borreliae. However, as in the Lyme disease spirochetes, in *B. turicatae* most *bdr*-carrying plasmids carry two *bdr* genes, one from subfamily *bdrA* and one from subfamily *bdrB* (24).

The sequence of more than 50 *bdr* alleles from five different *Borrelia* species has been determined (Table 2) (24,25,41-43,46). These extensive comparative sequence analyses led to the identification of conserved features that provide insight into the possible biologic roles of the Bdr proteins. For example, all *bdr* alleles carry centrally located repeat motif domains (Figure 3). Although conserved in sequence, these domains vary in length among alleles as a result of varying numbers of the repeat motif. The core tripeptide




Figure 3. Key features and putative functional domains of the Bdr proteins. The schematic depicts a prototype Bdr protein with the characteristics of each domain indicated. The abbreviation, ID%, is for percentage amino acid identity at either the inter- or intra-family level as indicated in the figure. Standard amino acid abbreviations are used in the figure to denote the conserved C-terminal lysine (K) or asparagine (N) residues, which are thought to be exposed in the periplasm and the cytoplasmically located core tripeptide of the repeat motif (lysine-isoleucine-aspartic acid; KID).

the repeat is the sequence KID. The repeat motifs encode consensus casein kinase 2 phosphorylation (CK2P) motifs of the sequence T/SKID/E (43). While it may appear somewhat paradoxical for bacteria to carry casein kinases, casein kinase is a descriptive term broadly applied to at least two classes of ubiquitous protein kinases for which the substrates may include various enzymes and noncatalytic proteins involved in important cellular regulatory functions (49). Most proteins phosphorylated by CK2-like kinases are highly acidic, as are the *Borrelia* Bdr proteins (isoelectric points between 5 and 6). The phosphorylation site in CK2P motifs is either the Ser or Thr residue of the motif. Although histidine kinases have been known to exist in some bacteria, it has been widely held that bacteria lack Ser - Thr kinases. However, Ser - Thr kinases have recently been identified in several bacterial species, including *Myxococcus*, *Anabeana*, *Fremyella*, *Yersinia*, and *Streptomyces* (50). Most importantly, analysis of the *B. burgdorferi* genome sequence identified a putative Ser - Thr kinase designated BB0648 (30,50). This ORF carries a domain that exhibits homology with the active site of Ser - Thr kinases. *B. burgdorferi* also carries a homolog of the PPM family of eucaryotic protein Ser - Thr phosphatases (30,50). The presence of these genes in *B. burgdorferi* suggests that the *Borrelia* possess the machinery necessary for Ser - Thr phosphorylation and dephosphorylation.

Another important conserved feature identified through sequence analyses is the hydrophobic carboxyl terminal domain of approximately 20 amino acids. Computer analyses conducted with the Tmpred program indicate that this domain has a high propensity to form a transmembrane helix (24,25). The Tmpred values for the 20 aa C-terminal domains are 2,000 to 2,600. A value of 500 or greater is considered significant (24,25). Comparison of the Bdr putative transmembrane domain sequences from the Lyme disease spirochetes with those from the relapsing fever spirochetes indicates that, while there is conservation in physical properties, there is essentially no conservation of primary sequence. However, sequence conservation does exist at the subfamily level (24,25). Since the Bdr proteins lack an obvious export signal, membrane association would most likely be with the spirochetal inner membrane, with the rest of the protein, which is hydrophilic, extending into the

cytoplasm. The terminal residue of the protein is in almost all cases a positively charged amino acid (lysine or asparagine). This residue could extend into the periplasm and serve to anchor the Bdr proteins to other cellular components, such as the peptidoglycan.

Immunologic Analyses of the Bdr Proteins

The presence of multiple *bdr* alleles and *bdr* subfamilies within isogenic populations has prompted speculation that there may be differential expression at either the subfamily or individual allele level, possibly in response to environmental stimuli (46). Limited studies of *bdr* expression and production, based on either mRNA detection or immunoblot analyses, have been performed. Porcella et al. (42) used Northern hybridization to determine if expression of *B. burgdorferi* *bdr*-related genes occurs during cultivation in the laboratory under standard culture conditions (33°C in BSK media). *Bdr* transcripts were not detected by this approach. Similarly, in an earlier analysis, we also conducted Northern hybridization experiments to assess *bdr* expression (46). We detected expression of *B. turicatae* OZ1 *bdrA* subfamily members in bacteria cultivated under standard laboratory growth conditions (46). However, when reverse transcriptase (RT)-PCR methods were applied, transcription of a single *bdrA* allele was detected (46). *B. turicatae* OZ-1 was later demonstrated to carry at least nine *bdr* alleles, four of which belong to the *bdrA* subfamily. Analysis of the sequence of these alleles showed that all four should have been readily amplified by the RT-PCR primer set because of the conservation of the primer binding sites (24). The lack of detection of transcript derived from these alleles suggested that only a subset of the *bdr* A subfamily alleles is expressed. This raised the possibility that other *bdr* alleles are either nonfunctional genes or their expression requires different environmental stimuli. The transcriptional expression of the *bdrB* subfamily has not been specifically assessed. Thorough transcriptional analyses using allele-specific probes and primers are an important step, since they allow specific assessment of the expression of individual *bdr* alleles under differing environmental conditions. In addition, analyses of the upstream DNA sequences of individual *bdr* alleles and their genomic location may elucidate the molecular basis for *bdr* transcriptional regulation.

Synopses

Immunologic analyses have provided a somewhat different overall picture regarding Bdr production. Immunologic analyses described in this report and elsewhere (44) demonstrate that several members of the *bdr* gene family are expressed during in vitro cultivation. We conducted a comprehensive analysis of the expression of Bdr proteins among *Borrelia* species. When antisera raised against recombinant *B. afzelii* BdrF₁ (24) were used in immunoblot analyses, several immunoreactive proteins were detected in cell lysates of all *Borrelia* species tested (Figure 4). The only exception was *B. anserina*, a causative agent of avian spirochetosis. Although *bdr*-related sequences have been detected in *B. anserina* by hybridization techniques (46), immunoreactive proteins were not detected in immunoblot analyses. Additional analyses are required to determine if this indicates absence of translational expression or the lack of epitope conservation in this species. In any event, the fact that immunoreactive bands were not detected in this species attests to the specificity of the anti-Bdr antisera. As a further demonstration of the

specificity of the antisera and to highlight the fact that the Bdr proteins are unique to *Borrelia*, a cell lysate of *Leptospira interrogans* was included in the immunoblot analyses. Immunoreactivity with proteins in the Bdr size range was not observed with the anti-Bdr antisera in this spirochete species. *Borrelia* species that expressed immunoreactive proteins included *B. garinii*, *B. burgdorferi*, *B. turdae*, *B. tanukii*, *B. japonica*, *B. valaisiana*, *B. afzelii*, *B. coriaceae*, *B. bissetii*, *B. miyamotoi*, *B. parkeri*, *B. hermsii*, and *B. turicatae* (Table 1). Particularly striking was the extensive variation in the number and molecular weight of the immunoreactive proteins expressed, with up to 12 distinct Bdr proteins detected. Variation in expression patterns was observed at both the inter- and intraspecies level. Analysis of three *B. burgdorferi* isolates (B31G, cN40, and CA12) demonstrated variability in both the size and number of expressed Bdr proteins. Isolate B31G has been demonstrated by genomic sequencing to carry 18 distinct *bdr* alleles. Immunoblot analyses show that not all alleles are expressed during in vitro cultivation; therefore, some alleles may be differentially regulated.




Figure 4. Immunoblot analyses demonstrating the variation in Bdr protein expression in *Borrelia* species and isolates. Bacteria were cultivated and prepared for analysis as described in the methods. Proteins were fractionated by SDS-PAGE, immunoblotted and screened with anti-BdrF_{1-B.afzelii_DK1} antisera. The species and isolates analyzed are indicated above each lane in panels A, B and C. The migration positions of the protein standards are indicated in each panel.

The broad immunoreactivity of the antisera with diverse *Borrelia* species indicates that some epitopes are conserved genuswide. In view of the sequence divergence in the N and C terminal domains of the Bdr proteins derived from different subfamilies, it is likely that the cross-reactive epitopes reside in the conserved repeat motif region. Consistent with this, computer analyses of the repeat domain of all determined Bdr protein sequences predict them to be alpha helical and to have a surface exposed on the protein and a positive Jameson-Wolf antigenic index (24,25,44). The conservation and synthesis of these polymorphic proteins in such a diverse group of *Borrelia* species suggest that they play an important role in *Borrelia* biology genuswide.

The Bdr Proteins and *Borrelia* Biology: An Overview

Bdr genes and extensive *bdr* gene families have now been identified and characterized in several diverse *Borrelia* species (24,25,42-44,46). Comparative sequence analyses, which have identified conserved putative functional domains, have provided the basis for the development of hypotheses regarding Bdr function and cellular location. The Bdr proteins, which lack known consensus export signals, are likely anchored to the cytoplasmic membrane through their conserved, hydrophobic, putative transmembrane spanning domain. The C-terminal positively charged amino acid may be exposed to the periplasm, where it may interact with other cellular components that may include the peptidoglycan. The repeat motif domain, which is predicted by computer analyses to be hydrophilic and surface exposed on the protein, likely extends into the cytoplasm. The conserved repeat motif domain that carries the putative Ser - Thr phosphorylation motifs may then be accessible for phosphorylation or to interact with other cytoplasmic proteins or DNA to form a membrane anchored complex. As with numerous other proteins, phosphorylation and dephosphorylation could play a regulatory role, perhaps in signaling or sensing.

Multiple polymorphic *bdr* alleles may increase the functional range and diversity of the Bdr proteins. Functional partitioning among Bdr proteins could offer a possible explanation of why *Borrelia* expend such biologic energy to maintain these genes in large gene families and express variants of these proteins. The homology among

bdr alleles may also allow or lead to the continual modification of these genes through homologous recombination. In fact, the variable nature of the repeat motif region, which is clearly not evolutionarily stable, has likely arisen from slipped-strand mispairing, recombination, or rearrangement. In view of the extensive genetic redundancy of the plasmid component of the *Borrelia* genome, recombination in and among related sequences on different plasmids could affect the organization and evolution of the genome and ultimately host-pathogen interaction. Inter- or intra-plasmid exchange of DNA sequences could provide a mechanistic basis for the extensive genetic variability that has been widely described for *Borrelia* plasmids (28,29,51-59). In spite of the apparent necessity for at least most of the plasmids for survival, as inferred from their ubiquitous distribution among *Borrelia* isolates, these bacteria are able to tolerate remarkable genomic variability. Diversity in the plasmids and the genes they carry may actually be exploited as a tool for phenotypic diversity and rapid environmental adaptation.

Acknowledgments

We thank Michael Norgard, Steve Porcella, Justin Radolf, and the Molecular Pathogenesis research group at Virginia Commonwealth University for helpful discussions.

This work was supported in part by a grant from the Jeffress Memorial Trust.

Mr. Roberts is a 3rd-year Ph.D. student, Department of Microbiology and Immunology, Medical College of Virginia at Virginia Commonwealth University. His scientific interests are centered on the study of the molecular mechanisms of pathogenesis and adaptive responses of arthropod-carried spirochetes, specifically on the role of plasmid-carried gene families in *Borrelia* biology and pathogenesis.

References

1. Barbour AG, Hayes SF. Biology of *Borrelia* species. *Microbiological Reviews* 1986;50:381-400.
2. Young JD. Underreporting of Lyme disease. *N Engl J Med* 1998;338:1629.
3. Steere AC, Malawista SE, Snyderman DR, Shope RE, Andiman WA, Ross MR, et al. Lyme arthritis: an epidemic of oligoarticular arthritis in children and adults in three Connecticut communities. *Arthritis Rheum* 1977;20:7-17.
4. Burgdorfer W, Barbour AG, Hayes SF, Benach JL, Grunwaldt E, Davis JP. Lyme disease—a tick-borne spirochetosis? *Science* 1982;216:1317-9.

5. Benach JL, Bosler EM, Hanrahan JP, Coleman JL, Bast TF, Habicht GS, et al. Spirochetes isolated from the blood of two patients with Lyme disease. *N Engl J Med* 1983;308:740-2.
6. Baranton G, Postic D, Saint Girons I, Boerlin P, Piffaretti J-C, Assous M, et al. Delineation of *Borrelia burgdorferi* sensu stricto, *Borrelia garinii* sp. nov., and group VS461 associated with Lyme borreliosis. *Int J Syst Bacteriol* 1992;42:378-83.
7. Marconi RT, Garon CF. Identification of a third genomic group of *Borrelia burgdorferi* through signature nucleotide analysis and 16S rRNA sequence determination. *Journal of General Microbiology* 1992;138:533-6.
8. Marconi RT, Garon CF. Phylogenetic analysis of the genus *Borrelia*: a comparison of North American and European isolates of *B. burgdorferi*. *J Bacteriol* 1992;174:241-4.
9. Marconi RT, Liveris D, Schwartz I. Identification of novel insertion elements, restriction fragment length polymorphism patterns, and discontinuous 23S rRNA in Lyme Disease spirochetes: phylogenetic analyses of rRNA genes and their intergenic spacers in *Borrelia japonica* sp. nov. and genomic group 21038 (*Borrelia andersonii* sp. nov.) isolates. *J Clin Microbiol* 1995;33:2427-34.
10. Fukunaga M, Hamase A, Okada K, Inoue H, Tsuruta Y, Miyamoto K, et al. Characterization of spirochetes isolated from ticks (*Ixodes tanukin*, *I. turdus* and *Ixodes columnae*) and comparison of the sequences with those of *Borrelia burgdorferi* sensu lato strains. *Appl Environ Microbiol* 1996;62:2338-44.
11. Fukunaga M, Hamase A, Okada K, Nakao M. *Borrelia tanuki* sp. nov. and *Borrelia turdae* sp. nov. found from Ixodid ticks in Japan: rapid species identification by 16S rRNA gene-targeted PCR analysis. *Microbiol Immunol* 1996;40:877-81.
12. Fukunaga M, Takahashi Y, Tsuruta Y, Matsushita O, Ralph D, McClelland M, et al. Genetic and phenotypic analysis of *Borrelia miyamotoi* sp. nov., isolates from the Ixodid tick *Ixodes persulcatus*, the vector for Lyme disease in Japan. *Int J Syst Bacteriol* 1995;45:804-10.
13. Wang G, van Dam AP, Le Flecha A, Postic D, Peter O, Baranton G, et al. Genetic and phenotypic analysis of *Borrelia valaisiana* sp. nov. (*Borrelia* genomic groups VS116 and M19). *Int J Syst Bacteriol* 1997;47:926-32.
14. Postic D, Belfazia J, Isogai E, Saint Girons I, Grimont PAD, Baranton G. A new genomic species in *Borrelia burgdorferi* sensu lato isolated from Japanese ticks. *Res Microbiol* 1993;144:467-73.
15. Postic D, Edlinger C, Richaud C, Grimont F, Dufresne Y, Perolat P, et al. Two genomic species in *Borrelia burgdorferi*. *Res Microbiol* 1990;141:465-75.
16. Barbour AG. Antigenic variation of a relapsing fever *Borrelia* species. *Annu Rev Microbiol* 1990;44:155-71.
17. Barbour AG, Byrman N, Carter CJ, Kitten T, Berstrom S. Variable antigen genes of the relapsing fever agent *Borrelia hermsii* are activated by promoter addition. *Mol Microbiol* 1991;5:489-93.
18. Stoenner HG, Dodd T, Larsen C. Antigenic variation of *B. hermsii*. *J Exp Med* 1982;156:1297-311.
19. Plasterk RHA, Simon MI, Barbour AG. Transposition of structural genes to an expression sequence on a linear plasmid causes antigenic variation in the bacterium *B. hermsii*. *Nature* 1985;318:257-63.
20. Zhang J-R, Hardham JM, Barbour AG, Norris AG. Antigenic variation in Lyme disease *Borreliae* by promiscuous recombination of vmp like sequence cassettes. *Cell* 1997;89:275-85.
21. Marconi RT, Samuels DS, Schwan TG, Garon CF. Identification of a protein in several *Borrelia* species which is related to OspC of the Lyme disease spirochetes. *J Clin Microbiol* 1993;31:2577-83.
22. Schwan TG, Piesman J, Golde WT, Dolan MC, Rosa PA. Induction of an outer surface protein on *Borrelia burgdorferi* during tick feeding. *Proc Natl Acad Sci U S A* 1995;92:2909-13.
23. Schwan TG, Hinnebusch BJ. Bloodstream- versus tick-associated variants of a relapsing fever bacterium. *Science* 1998;280:1938-40.
24. Carlyon JA, Roberts DM, Theisen M, Marconi RT. Molecular analyses of the *B. turicatae* bdr genes: a polymorphic, linear plasmid carried, paralogous gene family. In press 1999.
25. Carlyon JA, Roberts DM, Marconi RT. Evolutionary and molecular analyses of the *Borrelia rep* super gene family: delineation of six distinct sub-families and demonstration of the genus wide conservation of putative functional domains, structural properties and repeat motifs. *Microb Pathog*. In press 2000.
26. Ferdows MS, Barbour AG. Megabase-sized linear DNA in the bacterium *Borrelia burgdorferi*, the Lyme disease agent. *Proc Natl Acad Sci U S A* 1989;86:5969-73.
27. Barbour AG, Garon CF. Linear plasmids of the bacterium *Borrelia burgdorferi* have covalently closed ends. *Science* 1987;237:409-11.
28. Barbour AG. Plasmid analysis of *Borrelia burgdorferi*, the Lyme disease agent. *J Clin Microbiol* 1988;26:475-8.
29. Hinnebusch J, Barbour AG. Linear plasmids of *Borrelia burgdorferi* have a telomeric structure and sequence similar to those of a eukaryotic virus. *J Bacteriol* 1991;173:7233-9.
30. Fraser C, Casjens S, Huang WM, Sutton GG, Clayton R, Lathigra R, et al. Genomic sequence of a Lyme disease spirochaete, *Borrelia burgdorferi*. *Nature* 1997;390:580-6.
31. Barbour AG. Linear DNA of *Borrelia* species and antigenic variation. *Trends Microbiol* 1993;1.
32. Hinnebusch J, Tilly K. Linear plasmids and chromosomes in bacteria. *Mol Microbiol* 1993;10:917-22.
33. Hinnebusch J, Barbour AG. Linear- and circular-plasmid copy numbers in *Borrelia burgdorferi*. *J Bacteriol* 1992;174:5251-7.
34. Champion CI, Blanco DR, Skare JT, Haake DA, Giladi M, Foley D, et al. A 9.0 kilobase-pair circular plasmid of *Borrelia burgdorferi* encodes an exported protein: evidence for expression only during infection. *Infect Immun* 1994;62:2653-61.
35. Akins D, Porcella SF, Popova TG, Shevchenko D, Baker SI, Li M, et al. Evidence for *in vivo* but not *in vitro* expression of a *Borrelia burgdorferi* outer surface protein F (OspF) homologue. *Mol Microbiol* 1995;18:507-20.
36. Suk K, Das S, Sun W, Jwang B, Barthold SW, Flavell RA, et al. *Borrelia burgdorferi* genes selectively expressed in the infected host. *Proc Natl Acad Sci U S A* 1995;92:4269-73.

37. Skare JT, Foley DM, Hernandez SR, Moore DC, Blanco DR, Miller JN, et al. Cloning and molecular characterization of plasmid-encoded antigens of *Borrelia burgdorferi*. *Infect Immun* 1999;67:4407-17.
38. Marconi RT, Sung SY, Hughes CN, Carlyon JA. Molecular and evolutionary analyses of a variable series of genes in *Borrelia burgdorferi* that are related to *ospE* and *ospF*, constitute a gene family, and share a common upstream homology box. *J Bacteriol* 1996;178:5615-26.
39. Sung S-Y, LaVoie C, Carlyon JA, Marconi RT. Evolutionary instability of *ospE* related members of the UHB gene family in *Borrelia burgdorferi* sensu lato complex isolates. *Infect Immun* 1998;66:4656-68.
40. Zhang JR, Norris SJ. Genetic variation of the *Borrelia burgdorferi* gene *vlsE* involves cassette-specific, segmental gene conversion. *Infect Immun* 1998;66:3698-704.
41. Zuckert WR, Meyer J. Circular and linear plasmids of Lyme Disease spirochetes have extensive homology: characterization of a repeated DNA element. *J Bacteriol* 1996;178:2287-98.
42. Porcella SF, Popova TG, Akins DR, Li M, Radolf JR, Norgard MV. *Borrelia burgdorferi* supercoiled plasmids encode multicopy open reading frames and a lipoprotein gene family. *J Bacteriol* 1996;178:3293-307.
43. Theisen M. Molecular cloning and characterization of *nlpH*, encoding a novel surface exposed, polymorphic, plasmid-encoded 33kD lipoprotein of *Borrelia afzelii*. *J Bacteriol* 1996;178:6435-42.
44. Zuckert WR, Meyer J, Barbour AG. Comparative analysis and immunological characterization of the *Borrelia* Bdr protein family. *Infect Immun* 1999;67:3257-66.
45. Yang X, Popova TG, Hagman KE, Wikel SK, Schoeler GB, Caimano MJ, et al. Identification, characterization, and expression of three new members of the *Borrelia burgdorferi* Mlp (2.9) lipoprotein gene family. *Infect Immun* 1999;67:6008-18.
46. Carlyon JA, Marconi RT. Cloning and molecular characterization of a multi-copy, linear plasmid-carried, repeat motif-containing gene from *Borrelia turicatae*, a causative agent of relapsing fever. *J Bacteriol* 1998;180:4974-81.
47. Reeves PR, Hobbs M, Valvano MA, Kido N, Klena J, Maskell D, et al. Bacterial polysaccharide synthesis and gene nomenclature. *Trends Microbiol* 1996;4:498-503.
48. Demerec M. A proposal for a uniform nomenclature in bacterial genetics. *Genetics* 1966;54:61-76.
49. Pinna LA. Casein kinase 2: an "eminence grise" in cellular regulation. *Biochim Biophys Acta* 1990;1054:267-84.
50. Shi L, Potts M, Kennelly PJ. The serine, threonine and/or tyrosine protein kinases and protein phosphatases of prokaryotic organisms: a family portrait. *FEMS Microbiol Rev* 1998;22:229-53.
51. Xu Y, Kodner C, Coleman L, Johnson RC. Correlation of plasmids with infectivity of *Borrelia burgdorferi* sensu stricto type strain B31. *Infect Immun* 1996;64:3870-6.
52. Xu Y, Johnson RC. Analysis and comparison of plasmid profiles of *Borrelia burgdorferi* sensu lato strains. *J Clin Microbiol* 1995;33:2679-85.
53. Ståhlhammar-Carlemalm M, Jenny E, Gern L, Aeschlimann A, Meyer J. Plasmid analysis and restriction fragment length polymorphisms of chromosomal DNA allow a distinction between *Borrelia burgdorferi* strains. *Zentralbl Bakteriol* 1990;274:28-39.
54. Simpson WJ, Garon CF, Schwan TG. Analysis of supercoiled circular plasmids in infectious and non-infectious *Borrelia burgdorferi*. *Microb Pathog* 1990;8:109-18.
55. Schwan TG, Schrupf ME, Karstens RH, Clover JR, Wong J, Daugherty M, et al. Distribution and molecular analysis of Lyme disease spirochetes, *Borrelia burgdorferi*, isolated from ticks throughout California. *J Clin Microbiol* 1993;31:3096-108.
56. Marconi RT, Casjens S, Munderloh UG, Samuels DS. Analysis of linear plasmid dimers in *Borrelia burgdorferi* sensu lato isolates: implications concerning the potential mechanism of linear plasmid replication. *J Bacteriol* 1996;178:3357-61.
57. Marconi RT, Samuels DS, Landry RK, Garon CF. Analysis of the distribution and molecular heterogeneity of the *ospD* gene among the Lyme disease spirochetes: evidence for lateral gene exchange. *J Bacteriol* 1994;176:4572-82.
58. Ferdows MS, Serwer P, Griess GA, Norris SJ, Barbour AG. Conversion of a linear to a circular plasmid in the relapsing fever agent *Borrelia hermsii*. *J Bacteriol* 1996;178:793-800.
59. Carlyon JA, LaVoie C, Sung SY, Marconi RT. Analysis of the organization of multi-copy linear and circular plasmid carried open reading frames in *Borrelia burgdorferi* sensu lato isolates. *Infect Immun* 1998;66:1149-58.

Appendix

Bacterial Cultivation, DNA Isolation, and Southern Hybridization Analyses

Isolates belonging to the *Borrelia burgdorferi* sensu lato complex were cultivated in complete BSK-H media (Sigma) at 33°C. To cultivate the relapsing fever borreliae and other *Borrelia* species, the complete BSK-H media were supplemented with additional rabbit sera (Sigma) to a final concentration of 12% (vol/vol). Bacteria were harvested by centrifugation and washed with phosphate buffered saline (pH 7.0), and DNA was extracted (25). For Southern hybridization analyses, 5 µg of DNA from each isolate was digested under standard conditions with *Xba*I and fractionated by electrophoresis in 0.8% GTG agarose gels. (The DNA was transferred onto membranes for hybridization by vacuum blotting using the VacuGene system as described by the manufacturer (Pharmacia). All other Southern hybridization methods were as previously described (39).

Synopses

Immunoblot Analyses

Bacterial cultures were grown and harvested as described above. One OD₆₀₀ equivalent of cells was pelleted and resuspended in 100 μ l of standard SDS-sample buffer with reducing agents. The cell lysates (7 μ l) were fractionated by electrophoresis in 15% SDS-PAGE gels and electroblotted onto Immobilon P membranes (38). The immunoblots were blocked overnight in blocking buffer (1X PBS, 0.2% Tween, 0.002% NaCl, and 5% nonfat dry milk) and then incubated with a 1:1,000 antisera dilutions. ImmunoPure Goat anti-mouse IgG (H+L) peroxidase conjugate served as the secondary antibody. The secondary antibody was incubated with the blots for 1 hour at room temperature at a 1:40,000-

fold dilution and then the blots were washed three times with wash buffer. For chemiluminescent detection, the Supersignal West Pico Stable Peroxide solution and the Supersignal West Pico Luminol/Enhancer solution were used. Both reagents were from Pierce Chemical Company, Rockford, IL and were used as described by the manufacturer. The immunoblots were exposed to film for time frames of 5 to 30 seconds.

The *bdr* Gene Families of the Lyme Disease and Relapsing Fever Spirochetes: Potential Influence on Biology, Pathogenesis, and Evolution

David M. Roberts,* Jason A Carlyon,† Michael Theisen,‡ and Richard T. Marconi*

*Medical College of Virginia at Virginia Commonwealth University, School of Medicine, Richmond, Virginia, USA;
†Yale School of Medicine, Yale University, New Haven, Connecticut, USA; and ‡Statens Serum Institute, Copenhagen, Denmark

Appendix

Bacterial Cultivation, DNA Isolation, and Southern Hybridization Analyses

Isolates belonging to the *Borrelia burgdorferi* sensu lato complex were cultivated in complete BSK-H media (Sigma) at 33°C. To cultivate the relapsing fever borreliae and other *Borrelia* species, the complete BSK-H media were supplemented with additional rabbit sera (Sigma) to a final concentration of 12% (vol/vol). Bacteria were harvested by centrifugation and washed with phosphate buffered saline (pH 7.0), and DNA was extracted (25). For Southern hybridization analyses, 5µg of DNA from each isolate was digested under standard conditions with *Xba*I and fractionated by electrophoresis in 0.8% GTG agarose gels. The DNA was transferred onto membranes for hybridization by vacuum blotting using the VacuGene system as described by the manufacturer (Pharmacia). All other Southern hybridization methods were as previously described (39).

Immunoblot Analyses

Bacterial cultures were grown and harvested as described above. One OD600 equivalent of cells was pelleted and resuspended in 100µl of standard SDS-sample buffer with reducing agents. The cell lysates (7µl) were fractionated by electrophoresis in 15% SDS-PAGE gels and electroblotted onto Immobilon P membranes (38). The immunoblots were blocked overnight in

Publisher: CDC; Journal: Emerging Infectious Diseases

Article Type: Synopsis; Volume: 6; Issue: 2; Year: 2000; Article ID: 00-0203

DOI: 10.321/eid0602.000203; TOC Head: Synopsis

blocking buffer (1X PBS, 0.2% Tween, 0.002% NaCl, and 5% nonfat dry milk) and then incubated with a 1:1,000 antisera dilutions. ImmunoPure Goat anti-mouse IgG (H+L) peroxidase conjugate served as the secondary antibody. The secondary antibody was incubated with the blots for 1 hour at room temperature at a 1:40,000-fold dilution and then the blots were washed three times with wash buffer. For chemiluminescent detection, the Supersignal West Pico Stable Peroxide solution and the Supersignal West Pico Luminol/Enhancer solution were used. Both reagents were from Pierce and were used as described by the manufacturer. The immunoblots were exposed to film for time frames of 5 to 30 seconds.

Suggested citation: Roberts DM, Carlyon JA, Theisen M, Marconi RT. The *bdr* Gene Families of the Lyme Disease and Relapsing Fever Spirochetes: Potential Influence on Biology, Pathogenesis, and Evolution. *Emerg Infect Dis* [serial on the Internet]. 2000, Apr [date cited]. Available from <http://www.cdc.gov/ncidod/eid/vol6no2/roberts.htm>