

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
All causes									
2007.....	760.2	905.6	643.4	749.4	890.5	634.8	958.0	1,184.4	793.8
2006.....	776.5	924.8	657.8	764.4	908.2	648.2	982.0	1,215.6	813.0
2005.....	798.8	951.1	677.6	785.3	933.2	666.5	1,016.5	1,252.9	845.7
2004.....	800.8	955.7	679.2	786.3	936.9	666.9	1,027.3	1,269.4	855.3
2003.....	832.7	994.3	706.2	817.0	973.9	693.1	1,065.9	1,319.1	885.6
2002.....	845.3	1,013.7	715.2	829.0	992.9	701.3	1,083.3	1,341.4	901.8
2001.....	854.5	1,029.1	721.8	836.5	1,006.1	706.7	1,101.2	1,375.0	912.5
2000.....	869.0	1,053.8	731.4	849.8	1,029.4	715.3	1,121.4	1,403.5	927.6
1999.....	875.6	1,067.0	734.0	854.6	1,040.0	716.6	1,135.7	1,432.6	933.6
I. Certain infectious and parasitic diseases (A00-B99)									
2007.....	21.1	25.1	17.6	18.3	21.6	15.3	46.5	58.7	36.9
2006.....	21.6	25.8	17.8	18.5	22.2	15.3	48.1	60.6	38.2
2005.....	21.8	26.3	17.9	18.7	22.4	15.4	49.6	63.7	38.6
2004.....	21.7	26.3	17.6	18.4	22.2	15.0	50.5	65.0	39.1
2003.....	22.1	26.7	17.9	18.6	22.4	15.2	52.5	68.3	40.0
2002.....	22.2	27.2	17.7	18.5	22.5	14.9	54.2	71.0	41.0
2001.....	21.3	26.3	16.9	17.6	21.7	14.1	53.3	70.1	40.2
2000.....	21.3	26.3	16.8	17.4	21.4	13.9	55.0	72.9	41.0
1999.....	21.8	27.2	17.2	17.8	22.0	14.1	56.9	75.9	42.0
Intestinal infectious diseases (A00-A09)									
2007.....	2.1	2.1	2.2	2.2	2.2	2.3	1.6	1.6	1.6
2006.....	2.1	2.1	2.1	2.2	2.2	2.2	1.4	1.4	1.4
2005.....	1.9	1.9	1.8	2.0	2.0	1.9	1.2	1.3	1.2
2004.....	1.5	1.4	1.5	1.5	1.5	1.6	0.9	1.0	0.9
2003.....	1.1	1.0	1.1	1.1	1.0	1.1	0.8	0.7	0.8
2002.....	0.9	0.9	0.9	0.9	0.9	0.9	0.6	0.7	0.6
2001.....	0.5	0.5	0.6	0.6	0.6	0.6	0.4	0.5	0.4
2000.....	0.5	0.5	0.5	0.5	0.5	0.5	0.3	0.4	0.3
1999.....	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Amebiasis (A06)									
2007.....	*	*	*	*	*	*	*	*	*
2006.....	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Intestinal infections due to other specified organisms (A04,A07-A08)									
2007.....	2.1	2.0	2.1	2.2	2.1	2.2	1.5	1.6	1.5
2006.....	2.1	2.1	2.1	2.2	2.1	2.2	1.3	1.3	1.3
2005.....	1.8	1.8	1.8	1.9	1.9	1.9	1.2	1.2	1.2
2004.....	1.4	1.4	1.4	1.5	1.4	1.5	0.9	0.9	0.9
2003.....	1.0	1.0	1.0	1.0	1.0	1.1	0.7	0.7	0.7
2002.....	0.8	0.8	0.8	0.8	0.9	0.8	0.6	0.7	0.6
2001.....	0.5	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.4
2000.....	0.5	0.5	0.4	0.5	0.5	0.5	0.3	0.4	0.3
1999.....	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3
Diarrhea and gastroenteritis of infectious origin (A09)									
2007.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2006.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2005.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Tuberculosis (A16-A19)										
2007.....	0.2	0.2	0.1	0.1	0.2	0.1	0.4	0.6	0.2	
2006.....	0.2	0.3	0.1	0.1	0.2	0.1	0.5	0.9	0.3	
2005.....	0.2	0.3	0.1	0.1	0.2	0.1	0.5	0.8	0.3	
2004.....	0.2	0.3	0.1	0.2	0.2	0.1	0.7	1.0	0.4	
2003.....	0.2	0.3	0.1	0.2	0.2	0.1	0.6	1.0	0.4	
2002.....	0.3	0.4	0.2	0.2	0.3	0.1	0.7	1.1	0.5	
2001.....	0.3	0.4	0.2	0.2	0.2	0.1	0.8	1.2	0.5	
2000.....	0.3	0.4	0.2	0.2	0.2	0.1	0.9	1.5	0.6	
1999.....	0.3	0.5	0.2	0.2	0.3	0.1	0.9	1.5	0.6	
Respiratory tuberculosis (A16)										
2007.....	0.1	0.2	0.1	0.1	0.1	0.0	0.3	0.5	0.2	
2006.....	0.2	0.3	0.1	0.1	0.2	0.1	0.4	0.7	0.2	
2005.....	0.1	0.2	0.1	0.1	0.1	0.1	0.4	0.6	0.2	
2004.....	0.2	0.2	0.1	0.1	0.2	0.1	0.5	0.8	0.3	
2003.....	0.2	0.3	0.1	0.1	0.2	0.1	0.5	0.8	0.3	
2002.....	0.2	0.3	0.1	0.2	0.2	0.1	0.6	0.9	0.4	
2001.....	0.2	0.3	0.1	0.1	0.2	0.1	0.6	1.0	0.3	
2000.....	0.2	0.3	0.1	0.1	0.2	0.1	0.7	1.3	0.4	
1999.....	0.3	0.4	0.2	0.2	0.3	0.1	0.8	1.2	0.5	
Other tuberculosis (A17-A19)										
2007.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.2	*	
2006.....	0.1	0.1	0.0	0.0	0.0	0.0	0.1	0.2	*	
2005.....	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.2	*	
2004.....	0.1	0.1	0.0	0.0	0.0	0.0	0.2	0.2	*	
2003.....	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.1	*	
2002.....	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.2	*	
2001.....	0.1	0.1	0.0	0.0	0.0	0.0	0.2	0.2	0.1	
2000.....	0.1	0.1	0.0	0.0	0.0	0.0	0.2	0.3	0.2	
1999.....	0.1	0.1	0.0	0.0	0.1	0.0	0.2	0.3	*	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Scarlet fever and erysipelas (A38,A46)									
2007.....	*	*	*	*	*	*	*	*	*
2006.....	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Meningococcal infection (A39)									
2007.....	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
2006.....	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
2005.....	0.0	0.0	0.0	0.0	0.0	0.0	0.0	*	*
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2003.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2002.....	0.0	0.1	0.0	0.0	0.1	0.0	0.1	*	*
2001.....	0.1	0.1	0.1	0.1	0.0	0.1	0.1	*	*
2000.....	0.0	0.1	0.1	0.0	0.1	0.1	0.1	0.1	*
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
Septicemia (A40-A41)									
2007.....	11.0	12.1	10.2	10.0	11.1	9.3	21.7	24.6	19.9
2006.....	11.0	12.1	10.2	10.1	11.2	9.3	21.6	24.2	19.9
2005.....	11.2	12.3	10.4	10.2	11.2	9.4	22.6	25.8	20.5
2004.....	11.2	12.3	10.3	10.2	11.3	9.4	22.5	25.7	20.4
2003.....	11.6	12.7	10.9	10.5	11.5	9.9	23.8	27.3	21.6
2002.....	11.7	12.9	10.9	10.7	11.7	9.9	24.2	27.8	22.1
2001.....	11.4	12.7	10.5	10.3	11.5	9.5	23.6	27.7	21.2
2000.....	11.3	12.4	10.5	10.2	11.2	9.4	24.6	28.0	22.5
1999.....	11.3	12.5	10.4	10.0	11.1	9.3	24.6	28.9	21.9

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)										
2007.....	0.5	0.6	0.4	0.5	0.6	0.4	0.8	1.0	0.7	
2006.....	0.5	0.6	0.4	0.5	0.6	0.4	0.8	0.9	0.7	
2005.....	0.5	0.7	0.4	0.5	0.6	0.4	0.8	1.0	0.6	
2004.....	0.5	0.6	0.5	0.5	0.6	0.4	0.8	1.1	0.7	
2003.....	0.5	0.6	0.4	0.5	0.6	0.4	0.8	1.1	0.5	
2002.....	0.5	0.6	0.4	0.5	0.6	0.4	0.8	1.0	0.7	
2001.....	0.5	0.6	0.4	0.5	0.6	0.4	0.8	1.0	0.7	
2000.....	0.5	0.6	0.4	0.4	0.5	0.4	0.7	1.0	0.6	
1999.....	0.7	0.8	0.6	0.6	0.7	0.5	1.2	1.6	1.0	
Infections with a predominately sexual mode of transmission (A50-A64)										
2007.....	0.0	0.0	0.0	0.0	*	*	0.1	*	*	
2006.....	0.0	0.0	0.0	*	*	*	0.1	*	*	
2005.....	0.0	0.0	*	0.0	*	*	0.1	0.2	*	
2004.....	0.0	0.0	0.0	0.0	*	*	0.1	*	*	
2003.....	0.0	0.0	0.0	0.0	*	*	*	*	*	
2002.....	0.0	0.0	0.0	0.0	*	*	0.1	*	*	
2001.....	0.0	0.0	0.0	0.0	*	*	*	*	*	
2000.....	0.0	0.0	0.0	0.0	*	0.0	0.1	0.2	*	
1999.....	0.0	*	0.0	0.0	*	*	*	*	*	
Syphilis (A50-A53)										
2007.....	0.0	0.0	*	0.0	*	*	0.1	*	*	
2006.....	0.0	0.0	*	*	*	*	0.1	*	*	
2005.....	0.0	0.0	*	*	*	*	0.1	0.2	*	
2004.....	0.0	0.0	0.0	*	*	*	0.1	*	*	
2003.....	0.0	0.0	*	*	*	*	*	*	*	
2002.....	0.0	0.0	0.0	*	*	*	0.1	*	*	
2001.....	0.0	*	*	0.0	*	*	*	*	*	
2000.....	0.0	0.0	*	0.0	*	*	0.1	*	*	
1999.....	0.0	*	0.0	0.0	*	*	*	*	*	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Measles (B05)									
2007.....	*	*	*	*	*	*	*	*	*
2006.....	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Rubella (German measles) (B06)									
2007.....	*	*	*	*	*	*	*	*	*
2006.....	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Viral hepatitis (B15-B19)									
2007.....	2.3	3.1	1.5	2.2	3.0	1.4	2.9	4.2	1.9
2006.....	2.3	3.1	1.5	2.1	2.9	1.4	3.2	4.5	2.1
2005.....	1.8	2.4	1.2	1.7	2.3	1.1	2.6	3.4	1.9
2004.....	1.8	2.4	1.2	1.7	2.3	1.1	2.6	3.8	1.7
2003.....	1.8	2.5	1.2	1.7	2.4	1.1	2.7	3.8	1.8
2002.....	2.0	2.7	1.3	1.9	2.6	1.2	2.8	3.8	1.9
2001.....	2.0	2.6	1.3	1.8	2.5	1.2	2.7	3.4	2.1
2000.....	1.9	2.6	1.3	1.8	2.4	1.2	2.7	3.8	1.8
1999.....	1.8	2.4	1.2	1.6	2.2	1.1	2.6	3.4	1.9

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			Both sexes		White 2/		Black 2/	
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Human immunodeficiency virus (HIV) disease (B20-B24)									
2007.....	3.7	5.4	2.1	1.9	3.1	0.7	17.3	24.5	11.3
2006.....	4.0	5.9	2.2	2.1	3.4	0.7	18.6	26.3	12.2
2005.....	4.2	6.2	2.3	2.2	3.6	0.8	19.4	28.2	12.0
2004.....	4.5	6.6	2.4	2.3	3.8	0.9	20.4	29.2	13.0
2003.....	4.7	7.1	2.4	2.5	4.2	0.9	21.3	31.3	12.8
2002.....	4.9	7.4	2.5	2.6	4.3	0.9	22.5	33.3	13.4
2001.....	5.0	7.5	2.5	2.6	4.4	0.9	22.8	33.8	13.4
2000.....	5.2	7.9	2.5	2.8	4.6	1.0	23.3	35.1	13.2
1999.....	5.3	8.2	2.5	2.9	4.9	1.0	23.6	36.1	13.1
Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)									
2007.....	1.4	1.9	0.8	0.8	1.2	0.3	5.9	8.4	3.9
2006.....	1.5	2.2	0.9	0.8	1.3	0.3	6.7	9.4	4.5
2005.....	1.6	2.4	0.9	0.9	1.4	0.3	7.2	10.4	4.6
2004.....	1.8	2.5	1.0	0.9	1.5	0.4	7.8	10.9	5.1
2003.....	1.8	2.7	0.9	1.0	1.7	0.4	7.7	11.5	4.5
2002.....	1.9	2.9	1.0	1.1	1.7	0.4	8.6	12.8	5.1
2001.....	1.9	2.9	1.0	1.1	1.8	0.4	8.6	12.6	5.1
2000.....	2.0	3.1	1.0	1.1	1.8	0.4	9.0	13.4	5.2
1999.....	2.1	3.3	1.1	1.2	1.9	0.4	9.3	14.1	5.2
Human immunodeficiency virus (HIV) disease resulting in malignant neoplasms (B21)									
2007.....	0.3	0.4	0.1	0.2	0.3	0.0	1.1	1.8	0.5
2006.....	0.3	0.5	0.1	0.2	0.3	0.0	1.0	1.6	0.5
2005.....	0.3	0.5	0.1	0.2	0.4	0.1	1.2	1.9	0.6
2004.....	0.3	0.5	0.1	0.2	0.4	0.0	1.1	1.8	0.5
2003.....	0.3	0.5	0.1	0.2	0.4	0.0	1.1	1.6	0.6
2002.....	0.3	0.5	0.1	0.2	0.4	0.0	1.2	1.8	0.6
2001.....	0.3	0.6	0.1	0.2	0.4	0.1	1.2	2.0	0.6
2000.....	0.4	0.6	0.1	0.3	0.5	0.0	1.3	2.1	0.5
1999.....	0.4	0.7	0.1	0.3	0.5	0.1	1.3	2.2	0.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/		Black 2/			
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)									
2007.....	0.4	0.6	0.2	0.2	0.3	0.1	1.7	2.5	1.0
2006.....	0.4	0.6	0.2	0.2	0.4	0.1	1.8	2.7	1.1
2005.....	0.4	0.6	0.2	0.2	0.4	0.1	1.8	2.6	1.2
2004.....	0.4	0.7	0.2	0.3	0.4	0.1	1.8	2.7	1.1
2003.....	0.4	0.7	0.2	0.3	0.5	0.1	1.8	2.9	0.9
2002.....	0.5	0.7	0.2	0.3	0.5	0.1	1.8	2.8	1.0
2001.....	0.5	0.8	0.2	0.3	0.5	0.1	2.1	3.1	1.2
2000.....	0.6	0.9	0.2	0.3	0.6	0.1	2.3	3.6	1.2
1999.....	0.6	0.9	0.2	0.4	0.6	0.1	2.3	3.6	1.2
Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)									
2007.....	0.6	0.9	0.4	0.3	0.5	0.1	3.5	4.9	2.3
2006.....	0.7	1.0	0.4	0.3	0.5	0.1	3.8	5.3	2.5
2005.....	0.7	1.0	0.4	0.3	0.5	0.1	3.7	5.4	2.2
2004.....	0.7	1.1	0.4	0.3	0.5	0.1	3.8	5.6	2.4
2003.....	1.1	1.6	0.6	0.5	0.8	0.2	5.8	8.2	3.8
2002.....	1.2	1.7	0.7	0.5	0.8	0.2	6.2	9.1	3.8
2001.....	1.2	1.8	0.7	0.6	0.9	0.2	6.2	9.1	3.8
2000.....	1.2	1.8	0.7	0.6	0.9	0.2	6.0	8.9	3.7
1999.....	1.2	1.8	0.7	0.6	0.9	0.2	6.4	9.5	3.7
Unspecified human immunodeficiency virus (HIV) disease (B24)									
2007.....	1.0	1.5	0.6	0.5	0.8	0.2	5.1	6.9	3.6
2006.....	1.1	1.6	0.6	0.5	0.9	0.2	5.3	7.4	3.6
2005.....	1.2	1.7	0.6	0.6	1.0	0.2	5.5	7.9	3.4
2004.....	1.2	1.7	0.7	0.6	1.0	0.2	5.8	8.2	3.9
2003.....	1.0	1.5	0.6	0.5	0.8	0.2	4.9	7.1	3.0
2002.....	1.0	1.4	0.5	0.5	0.8	0.2	4.6	6.8	2.9
2001.....	1.0	1.5	0.5	0.5	0.8	0.1	4.7	7.0	2.8
2000.....	1.0	1.5	0.5	0.5	0.8	0.2	4.7	7.1	2.6
1999.....	1.0	1.5	0.5	0.5	0.9	0.2	4.4	6.6	2.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
All other and unspecified viral diseases (A81,A85.0-A85.1,A85.8,A86-A89,B01,B04,B07-B09,B25-B34)										
2007.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.4	0.3
2006.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
2005.....	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.3	0.3	0.3
2004.....	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.3	0.3	0.3
2003.....	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.3	0.3	0.3
2002.....	0.4	0.4	0.3	0.4	0.4	0.3	0.3	0.3	0.4	0.3
2001.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
2000.....	0.4	0.4	0.3	0.4	0.4	0.3	0.4	0.4	0.4	0.4
1999.....	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)										
2007.....	0.8	1.0	0.7	0.8	1.0	0.7	1.1	1.4	0.9	
2006.....	0.9	1.1	0.8	0.9	1.1	0.7	1.4	1.8	1.2	
2005.....	1.5	1.9	1.2	1.4	1.8	1.1	1.9	2.4	1.5	
2004.....	1.5	1.9	1.1	1.5	1.9	1.1	2.0	2.5	1.5	
2003.....	1.5	1.8	1.2	1.4	1.7	1.1	1.9	2.4	1.6	
2002.....	1.3	1.7	0.9	1.2	1.6	0.9	1.8	2.5	1.3	
2001.....	1.1	1.3	0.9	1.0	1.2	0.8	1.7	2.1	1.4	
2000.....	1.1	1.3	0.8	1.0	1.2	0.8	1.7	2.2	1.3	
1999.....	1.5	1.8	1.2	1.3	1.6	1.1	2.9	3.3	2.6	
Mycoses (B35-B49)										
2007.....	0.3	0.4	0.3	0.3	0.4	0.2	0.5	0.6	0.5	
2006.....	0.3	0.4	0.3	0.3	0.4	0.2	0.7	0.8	0.6	
2005.....	0.3	0.4	0.3	0.3	0.4	0.2	0.7	0.8	0.6	
2004.....	0.4	0.5	0.3	0.3	0.4	0.3	0.7	0.9	0.5	
2003.....	0.4	0.4	0.3	0.3	0.4	0.3	0.7	0.8	0.6	
2002.....	0.4	0.6	0.3	0.4	0.5	0.3	0.8	1.2	0.5	
2001.....	0.4	0.5	0.3	0.3	0.4	0.3	0.8	1.1	0.6	
2000.....	0.4	0.6	0.3	0.4	0.5	0.3	0.8	1.1	0.7	
1999.....	0.7	0.9	0.5	0.6	0.8	0.4	1.3	1.6	1.1	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Trypanosomiasis (B56-B57)									
2007.....	*	*	*	*	*	*	*	*	*
2006.....	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Pneumocystosis (B59)									
2007.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2006.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*
2005.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*
2004.....	0.1	0.1	0.0	0.0	0.1	0.0	0.1	0.1	*
2003.....	0.1	0.1	0.0	0.0	0.0	0.0	0.1	0.1	*
2002.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	0.1
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.1
2000.....	0.1	0.1	0.1	0.1	0.1	0.0	0.2	0.2	0.1
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2
Other and unspecified protozoal diseases (B58,B60-B64)									
2007.....	*	*	*	*	*	*	*	*	*
2006.....	*	*	*	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*
1999.....	0.0	0.0	*	*	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Sequelae of tuberculosis (B90)									
2007.....	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2006.....	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2005.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.1	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
Sequelae of poliomyelitis (B91)									
2007.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2006.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2005.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2004.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2003.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99)									
2007.....	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.6	0.3
2006.....	0.4	0.4	0.3	0.4	0.4	0.3	0.5	0.7	0.4
2005.....	0.9	1.2	0.7	0.9	1.2	0.7	1.1	1.4	0.9
2004.....	0.9	1.2	0.7	0.9	1.2	0.7	1.0	1.3	0.8
2003.....	0.9	1.1	0.7	0.9	1.1	0.6	1.0	1.4	0.8
2002.....	0.6	0.8	0.4	0.6	0.8	0.4	0.7	1.0	0.6
2001.....	0.4	0.5	0.4	0.4	0.5	0.3	0.6	0.6	0.5
2000.....	0.3	0.4	0.3	0.3	0.4	0.3	0.5	0.7	0.4
1999.....	0.5	0.6	0.5	0.4	0.5	0.4	1.2	1.4	1.1

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
II. Neoplasms (C00-D48)									
2007.....	182.9	223.1	155.0	182.1	220.9	155.0	219.6	287.5	178.3
2006.....	185.3	225.6	157.5	184.6	223.5	157.5	221.7	290.2	180.0
2005.....	188.3	230.6	159.4	187.2	228.0	159.1	226.9	298.3	183.4
2004.....	190.4	233.3	161.2	189.0	230.1	160.9	231.6	306.3	186.4
2003.....	194.7	239.1	164.8	193.2	235.9	164.2	237.6	313.9	191.4
2002.....	198.1	244.6	167.0	196.4	241.0	166.4	243.2	324.7	194.4
2001.....	200.9	249.8	168.8	198.8	245.4	168.1	247.8	336.5	195.5
2000.....	204.5	255.0	171.8	202.2	250.1	171.2	253.1	345.7	197.9
1999.....	205.6	258.0	171.7	202.9	252.7	170.5	257.3	351.4	201.3
Malignant neoplasms (C00-C97)									
2007.....	178.4	217.5	151.3	177.5	215.1	151.2	215.5	282.3	174.9
2006.....	180.7	220.1	153.6	179.9	217.9	153.6	217.4	284.9	176.1
2005.....	183.8	225.1	155.6	182.6	222.3	155.2	222.7	293.7	179.6
2004.....	185.8	227.7	157.4	184.4	224.4	157.0	227.2	301.2	182.5
2003.....	190.1	233.3	160.9	188.5	230.1	160.2	233.3	308.8	187.7
2002.....	193.5	238.9	163.1	191.7	235.2	162.4	238.8	319.6	190.3
2001.....	196.0	243.7	164.7	193.9	239.2	163.9	243.1	330.9	191.3
2000.....	199.6	248.9	167.6	197.2	243.9	166.9	248.5	340.3	193.8
1999.....	200.8	251.9	167.6	197.9	246.5	166.4	252.5	346.1	196.9
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)									
2007.....	2.5	3.9	1.4	2.5	3.7	1.4	3.3	5.7	1.6
2006.....	2.5	3.8	1.4	2.4	3.6	1.4	3.2	5.7	1.5
2005.....	2.5	3.9	1.4	2.4	3.6	1.4	3.5	6.5	1.4
2004.....	2.6	4.0	1.5	2.5	3.7	1.5	3.8	6.9	1.6
2003.....	2.6	4.0	1.5	2.5	3.8	1.4	3.8	6.9	1.6
2002.....	2.7	4.1	1.5	2.5	3.8	1.5	3.6	6.3	1.7
2001.....	2.7	4.1	1.6	2.6	3.9	1.5	3.9	6.7	1.8
2000.....	2.7	4.1	1.6	2.5	3.7	1.6	4.1	7.3	1.8
1999.....	2.7	4.1	1.6	2.5	3.8	1.5	4.3	7.5	2.1

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Of lip (C00)									
2007.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2006.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2005.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2002.....	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2001.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2000.....	0.0	0.0	*	0.0	0.0	*	*	*	*
1999.....	0.0	0.0	*	0.0	0.0	*	*	*	*
Of tongue (C01-C02)									
2007.....	0.6	0.9	0.4	0.6	0.9	0.4	0.7	1.1	0.4
2006.....	0.6	0.9	0.4	0.6	0.9	0.4	0.6	1.0	0.3
2005.....	0.6	0.9	0.4	0.6	0.9	0.4	0.7	1.4	0.2
2004.....	0.6	0.9	0.4	0.6	0.9	0.4	0.7	1.2	0.3
2003.....	0.6	0.9	0.4	0.6	0.9	0.4	0.7	1.3	0.4
2002.....	0.7	0.9	0.4	0.6	0.9	0.4	0.7	1.3	0.3
2001.....	0.6	0.9	0.4	0.6	0.9	0.4	0.7	1.2	0.3
2000.....	0.6	0.9	0.4	0.6	0.9	0.4	0.8	1.4	0.3
1999.....	0.6	0.9	0.4	0.6	0.9	0.4	0.8	1.4	0.4
Of pharynx (C10-C13,C14.0)									
2007.....	1.0	1.7	0.5	0.9	1.5	0.5	1.7	3.1	0.7
2006.....	1.0	1.7	0.5	0.9	1.5	0.5	1.7	3.3	0.6
2005.....	1.0	1.7	0.5	0.9	1.5	0.5	1.9	3.6	0.6
2004.....	1.1	1.8	0.5	1.0	1.5	0.5	2.1	3.9	0.9
2003.....	1.1	1.8	0.5	1.0	1.5	0.5	2.0	3.8	0.7
2002.....	1.1	1.7	0.6	1.0	1.5	0.5	1.8	3.3	0.8
2001.....	1.1	1.8	0.6	1.0	1.6	0.5	2.0	3.6	0.9
2000.....	1.1	1.8	0.6	1.0	1.5	0.5	2.2	4.1	0.9
1999.....	1.1	1.8	0.6	1.0	1.6	0.5	2.3	4.0	1.0

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Of other and unspecified sites within the lip, oral cavity and pharynx (C03-C09,C14.2-C14.8)										
2007.....	0.8	1.2	0.5	0.8	1.2	0.5	0.9	1.5	0.5	
2006.....	0.8	1.2	0.5	0.8	1.2	0.5	0.9	1.4	0.5	
2005.....	0.8	1.2	0.5	0.8	1.2	0.5	0.9	1.5	0.5	
2004.....	0.9	1.2	0.5	0.8	1.2	0.6	1.0	1.8	0.4	
2003.....	0.9	1.3	0.5	0.9	1.2	0.6	1.1	1.8	0.5	
2002.....	0.9	1.3	0.6	0.9	1.3	0.6	1.0	1.7	0.5	
2001.....	0.9	1.3	0.6	0.9	1.3	0.6	1.2	1.9	0.6	
2000.....	0.9	1.3	0.6	0.9	1.3	0.6	1.1	1.9	0.5	
1999.....	0.9	1.3	0.6	0.9	1.3	0.6	1.2	2.0	0.7	
Malignant neoplasms of digestive organs (C15-C26)										
2007.....	42.7	55.1	32.9	41.3	53.3	31.6	58.2	76.8	45.2	
2006.....	43.2	55.3	33.6	41.7	53.4	32.3	58.7	76.9	46.0	
2005.....	43.5	56.0	33.7	41.9	54.0	32.3	59.8	78.6	46.7	
2004.....	43.9	56.4	34.2	42.4	54.4	32.8	60.4	78.4	47.7	
2003.....	45.1	57.7	35.3	43.4	55.6	33.7	62.4	80.8	49.6	
2002.....	45.7	58.8	35.6	44.0	56.6	34.1	63.4	82.9	50.1	
2001.....	46.4	59.4	36.5	44.6	57.2	34.9	64.8	83.3	52.1	
2000.....	47.2	60.5	37.2	45.3	58.1	35.6	66.2	86.2	52.5	
1999.....	47.3	60.7	37.3	45.4	58.4	35.7	66.7	87.1	52.7	
Of esophagus (C15)										
2007.....	4.3	7.7	1.6	4.3	7.8	1.5	4.7	8.3	2.2	
2006.....	4.4	7.8	1.7	4.5	7.9	1.6	4.8	8.3	2.3	
2005.....	4.4	7.9	1.7	4.5	8.0	1.6	5.0	8.8	2.3	
2004.....	4.3	7.7	1.7	4.3	7.7	1.6	5.4	9.0	2.8	
2003.....	4.4	7.7	1.7	4.3	7.7	1.7	5.7	9.7	2.9	
2002.....	4.4	7.8	1.7	4.3	7.7	1.6	5.9	10.1	2.9	
2001.....	4.4	7.8	1.8	4.3	7.6	1.7	6.4	11.3	3.0	
2000.....	4.4	7.7	1.8	4.3	7.6	1.7	6.5	11.0	3.3	
1999.....	4.4	7.7	1.8	4.2	7.4	1.6	6.5	11.5	3.1	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Of stomach (C16)										
2007.....	3.6	5.0	2.6	3.1	4.4	2.2	6.9	10.4	4.6	
2006.....	3.7	5.0	2.7	3.2	4.4	2.3	6.8	9.8	4.9	
2005.....	3.8	5.2	2.7	3.3	4.5	2.4	7.2	10.6	4.9	
2004.....	4.0	5.5	2.8	3.5	4.9	2.5	7.5	11.1	5.2	
2003.....	4.1	5.7	3.0	3.6	5.0	2.6	7.8	11.3	5.4	
2002.....	4.2	5.9	3.0	3.7	5.2	2.5	8.4	11.9	6.1	
2001.....	4.4	6.1	3.1	3.8	5.4	2.7	8.3	11.9	5.8	
2000.....	4.6	6.4	3.3	4.0	5.6	2.8	9.0	12.7	6.5	
1999.....	4.6	6.6	3.3	4.1	5.8	2.8	9.2	13.5	6.4	
Of small intestine (C17)										
2007.....	0.3	0.4	0.2	0.3	0.4	0.2	0.5	0.6	0.5	
2006.....	0.4	0.4	0.3	0.3	0.4	0.3	0.5	0.6	0.5	
2005.....	0.4	0.4	0.3	0.3	0.4	0.3	0.5	0.6	0.5	
2004.....	0.4	0.5	0.3	0.3	0.4	0.3	0.6	0.7	0.5	
2003.....	0.4	0.4	0.3	0.3	0.4	0.3	0.5	0.7	0.4	
2002.....	0.4	0.4	0.3	0.3	0.4	0.3	0.6	0.7	0.5	
2001.....	0.4	0.5	0.3	0.4	0.4	0.3	0.6	0.8	0.5	
2000.....	0.4	0.5	0.3	0.4	0.4	0.3	0.5	0.6	0.4	
1999.....	0.4	0.5	0.3	0.4	0.5	0.3	0.6	0.8	0.4	
Of colon, rectum and anus (C18-C21)										
2007.....	16.9	20.1	14.4	16.4	19.6	13.9	23.5	29.0	19.8	
2006.....	17.2	20.5	14.7	16.7	19.9	14.3	24.3	30.6	20.2	
2005.....	17.5	20.9	14.8	16.9	20.3	14.3	24.8	30.2	21.2	
2004.....	18.0	21.5	15.3	17.6	21.0	14.9	24.7	30.0	21.2	
2003.....	19.1	22.9	16.2	18.6	22.4	15.7	26.4	31.9	22.8	
2002.....	19.7	23.7	16.7	19.2	23.1	16.2	26.8	33.1	22.8	
2001.....	20.1	24.2	17.2	19.6	23.7	16.6	27.6	33.1	24.1	
2000.....	20.8	25.1	17.7	20.3	24.5	17.2	28.2	34.9	24.0	
1999.....	20.9	25.3	17.8	20.4	24.8	17.3	28.4	34.2	24.7	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Colon (C18)										
2007.....	13.9	16.3	12.0	13.4	15.8	11.5	19.9	24.2	17.2	
2006.....	14.2	16.7	12.3	13.7	16.1	11.9	20.5	25.8	17.3	
2005.....	14.4	17.1	12.4	13.9	16.6	11.9	21.3	25.5	18.5	
2004.....	14.9	17.6	12.9	14.5	17.1	12.5	21.3	25.5	18.5	
2003.....	16.0	19.0	13.8	15.5	18.5	13.2	22.7	27.1	19.9	
2002.....	16.5	19.7	14.2	16.0	19.1	13.7	23.2	28.3	19.9	
2001.....	16.9	20.1	14.7	16.4	19.6	14.1	24.0	28.4	21.3	
2000.....	17.6	21.0	15.1	17.1	20.4	14.7	24.6	29.9	21.3	
1999.....	17.7	21.2	15.3	17.3	20.8	14.7	24.8	29.2	22.0	
Rectosigmoid junction and rectum (C19-C20)										
2007.....	2.8	3.6	2.2	2.8	3.6	2.2	3.3	4.6	2.5	
2006.....	2.8	3.7	2.2	2.8	3.6	2.2	3.5	4.6	2.7	
2005.....	2.9	3.7	2.2	2.8	3.6	2.2	3.3	4.5	2.5	
2004.....	2.9	3.7	2.2	2.9	3.7	2.2	3.3	4.3	2.5	
2003.....	2.9	3.8	2.2	2.9	3.7	2.2	3.4	4.5	2.7	
2002.....	3.0	3.9	2.3	3.0	3.8	2.3	3.4	4.6	2.7	
2001.....	3.0	4.0	2.3	3.0	4.0	2.3	3.4	4.5	2.7	
2000.....	3.0	4.0	2.3	3.0	3.9	2.3	3.4	4.7	2.5	
1999.....	3.0	3.9	2.3	3.0	3.9	2.4	3.4	4.8	2.5	
Anus and anus canal (C21)										
2007.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	
2006.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	
2005.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	
2004.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2	
2003.....	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.2	0.2	
2002.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	
2001.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	*	0.2	
2000.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	
1999.....	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.2	0.2	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Of liver and intrahepatic bile ducts (C22)									
2007.....	5.4	7.9	3.2	5.0	7.2	3.1	7.1	11.5	3.8
2006.....	5.3	7.7	3.3	4.9	7.0	3.1	7.0	11.1	3.9
2005.....	5.2	7.7	3.2	4.8	7.1	3.0	7.0	11.3	3.8
2004.....	5.1	7.4	3.2	4.7	6.7	3.0	6.9	10.7	4.1
2003.....	5.0	7.4	3.1	4.6	6.7	2.8	6.6	10.2	4.0
2002.....	4.9	7.1	3.0	4.5	6.5	2.8	6.3	9.9	3.6
2001.....	4.7	6.9	3.0	4.3	6.3	2.7	6.2	9.1	4.1
2000.....	4.7	6.8	3.0	4.2	6.2	2.7	6.1	9.5	3.6
1999.....	4.5	6.5	3.0	4.2	5.9	2.7	5.9	8.7	3.8
Liver (C22.0,C22.2-C22.9)									
2007.....	4.2	6.5	2.2	3.8	5.8	2.0	6.0	10.2	2.8
2006.....	4.1	6.4	2.2	3.7	5.7	2.0	6.0	10.0	2.9
2005.....	4.1	6.4	2.2	3.7	5.7	2.0	5.9	9.9	2.8
2004.....	3.9	6.1	2.2	3.5	5.4	1.9	5.9	9.6	3.1
2003.....	3.9	6.0	2.1	3.5	5.4	1.9	5.6	8.9	3.1
2002.....	3.8	5.9	2.1	3.5	5.3	1.9	5.4	8.8	2.8
2001.....	3.7	5.7	2.1	3.3	5.2	1.9	5.3	8.0	3.3
2000.....	3.7	5.7	2.1	3.3	5.0	1.9	5.3	8.6	2.8
1999.....	3.6	5.4	2.1	3.2	4.9	1.9	5.1	7.9	3.1
Intrahepatic bile duct carcinoma (C22.1)									
2007.....	1.2	1.4	1.1	1.2	1.4	1.1	1.1	1.2	1.0
2006.....	1.2	1.3	1.1	1.2	1.4	1.1	1.0	1.1	0.9
2005.....	1.1	1.4	1.0	1.1	1.3	1.0	1.1	1.4	0.9
2004.....	1.2	1.3	1.1	1.2	1.3	1.0	1.0	1.1	0.9
2003.....	1.1	1.3	1.0	1.1	1.3	1.0	1.1	1.3	0.9
2002.....	1.0	1.2	0.9	1.0	1.2	0.9	0.9	1.1	0.8
2001.....	1.0	1.2	0.9	1.0	1.2	0.9	0.9	1.1	0.8
2000.....	1.0	1.1	0.9	1.0	1.2	0.9	0.8	0.9	0.8
1999.....	0.9	1.0	0.9	0.9	1.1	0.9	0.8	0.8	0.7

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Of gallbladder and extrahepatic bile ducts (C23-C24)										
2007.....	1.0	1.0	1.1	1.0	1.0	1.1	1.1	0.9	1.2	
2006.....	1.1	1.0	1.2	1.1	1.0	1.2	1.2	1.0	1.4	
2005.....	1.1	1.0	1.2	1.1	1.0	1.2	1.3	1.2	1.3	
2004.....	1.1	1.0	1.2	1.1	1.0	1.2	1.2	0.9	1.4	
2003.....	1.2	1.1	1.2	1.1	1.1	1.2	1.1	0.9	1.2	
2002.....	1.2	1.1	1.3	1.1	1.1	1.2	1.4	1.1	1.5	
2001.....	1.3	1.2	1.4	1.3	1.1	1.4	1.3	1.2	1.5	
2000.....	1.3	1.2	1.4	1.3	1.2	1.4	1.2	1.0	1.4	
1999.....	1.3	1.2	1.4	1.3	1.2	1.4	1.3	1.0	1.4	
Of pancreas (C25)										
2007.....	10.8	12.5	9.4	10.6	12.4	9.2	14.0	15.5	12.7	
2006.....	10.8	12.3	9.5	10.6	12.3	9.3	13.6	14.8	12.5	
2005.....	10.8	12.3	9.5	10.6	12.2	9.3	13.6	15.3	12.3	
2004.....	10.6	12.3	9.3	10.4	12.2	9.1	13.6	15.4	12.1	
2003.....	10.5	12.0	9.3	10.3	11.9	9.0	13.7	15.3	12.4	
2002.....	10.5	12.1	9.2	10.3	12.0	9.0	13.6	15.3	12.3	
2001.....	10.5	12.1	9.3	10.4	12.0	9.1	13.6	15.0	12.5	
2000.....	10.6	12.2	9.3	10.4	12.0	9.0	14.1	15.8	12.7	
1999.....	10.6	12.3	9.3	10.4	12.0	9.1	14.1	16.4	12.3	
Of other and ill-defined digestive organs (C26)										
2007.....	0.4	0.5	0.3	0.4	0.5	0.3	0.4	0.6	0.3	
2006.....	0.4	0.5	0.3	0.4	0.4	0.3	0.5	0.7	0.4	
2005.....	0.4	0.4	0.3	0.3	0.4	0.3	0.4	0.6	0.3	
2004.....	0.4	0.4	0.3	0.4	0.4	0.3	0.4	0.5	0.4	
2003.....	0.5	0.5	0.4	0.5	0.5	0.4	0.6	0.8	0.5	
2002.....	0.5	0.5	0.4	0.5	0.5	0.4	0.6	0.8	0.5	
2001.....	0.5	0.6	0.4	0.5	0.6	0.4	0.7	0.8	0.6	
2000.....	0.5	0.6	0.4	0.5	0.6	0.4	0.7	0.7	0.6	
1999.....	0.6	0.7	0.5	0.6	0.7	0.5	0.7	0.9	0.5	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/		Black 2/			
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)									
2007.....	52.1	67.5	40.6	52.6	67.0	41.8	58.1	87.0	39.0
2006.....	53.0	69.6	40.8	53.5	69.2	41.9	59.2	88.2	39.9
2005.....	54.2	71.6	41.2	54.6	71.1	42.2	61.1	91.4	41.0
2004.....	54.8	72.7	41.6	55.0	71.9	42.6	62.4	95.0	40.9
2003.....	55.7	74.5	42.0	56.0	73.6	43.0	63.7	98.1	41.1
2002.....	56.5	76.0	42.3	56.8	75.1	43.3	64.7	100.3	41.2
2001.....	57.0	78.1	41.8	57.1	76.9	42.8	65.4	104.7	39.9
2000.....	57.8	79.7	42.1	57.8	78.4	43.0	67.2	107.1	41.1
1999.....	57.2	79.8	41.1	57.0	78.4	41.8	67.7	108.5	41.1
Of nasal cavity, middle ear and accessory sinuses (C30-C31)									
2007.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.1
2006.....	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.1	0.1
2005.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.3	*
2004.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.3	*
2003.....	0.1	0.2	0.1	0.2	0.2	0.1	0.2	0.3	*
2002.....	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.2	*
2001.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.3	*
2000.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.1
1999.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	*
Of larynx (C32)									
2007.....	1.1	2.1	0.4	1.1	1.9	0.4	2.1	4.3	0.6
2006.....	1.2	2.2	0.5	1.2	2.0	0.5	2.1	4.1	0.7
2005.....	1.2	2.2	0.5	1.2	2.1	0.5	2.3	4.6	0.8
2004.....	1.2	2.2	0.4	1.1	2.1	0.4	2.2	4.6	0.7
2003.....	1.3	2.4	0.5	1.2	2.2	0.5	2.5	5.1	0.7
2002.....	1.3	2.3	0.5	1.2	2.1	0.5	2.5	4.8	0.8
2001.....	1.3	2.5	0.5	1.2	2.3	0.5	2.5	5.0	0.9
2000.....	1.4	2.5	0.5	1.3	2.3	0.5	2.8	5.5	0.9
1999.....	1.4	2.5	0.5	1.3	2.3	0.5	2.6	5.1	0.8

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Of trachea, bronchus and lung (C33-C34)									
2007.....	50.6	65.1	40.0	51.2	64.8	41.2	55.6	82.2	38.1
2006.....	51.5	67.0	40.0	52.1	66.8	41.1	56.8	83.7	39.0
2005.....	52.6	69.0	40.5	53.1	68.7	41.5	58.4	86.4	40.0
2004.....	53.2	70.1	40.9	53.6	69.4	41.9	59.8	90.0	39.9
2003.....	54.1	71.7	41.3	54.5	71.1	42.3	60.8	92.4	40.2
2002.....	54.9	73.2	41.6	55.3	72.5	42.6	61.9	95.0	40.1
2001.....	55.3	75.2	41.0	55.6	74.2	42.1	62.5	99.1	38.7
2000.....	56.1	76.7	41.3	56.2	75.7	42.3	64.0	101.1	39.8
1999.....	55.5	76.9	40.2	55.4	75.6	41.0	64.8	103.0	39.9
Of thymus, heart, mediastinum and pleura (C37-C38)									
2007.....	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.2	0.1
2006.....	0.1	0.1	0.1	0.1	0.2	0.1	0.2	0.2	0.2
2005.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.2
2004.....	0.2	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.2
2003.....	0.2	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.1
2002.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.1
2001.....	0.2	0.2	0.1	0.2	0.2	0.2	0.2	0.2	0.2
2000.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.2
1999.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.2
Of other and ill-defined sites in respiratory system and intrathoracic organs (C39)									
2007.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2006.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2005.....	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2002.....	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2001.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2000.....	0.0	0.0	*	0.0	0.0	*	*	*	*
1999.....	0.0	0.0	*	0.0	0.0	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/		Black 2/			
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of bone and articular cartilage (C40-C41)									
2007.....	0.4	0.5	0.3	0.5	0.6	0.4	0.4	0.5	0.3
2006.....	0.4	0.5	0.4	0.4	0.5	0.4	0.5	0.6	0.4
2005.....	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.5	0.4
2004.....	0.4	0.5	0.4	0.4	0.5	0.3	0.5	0.6	0.4
2003.....	0.4	0.5	0.3	0.4	0.5	0.4	0.4	0.5	0.3
2002.....	0.4	0.5	0.3	0.4	0.5	0.3	0.4	0.5	0.4
2001.....	0.5	0.6	0.4	0.5	0.6	0.3	0.5	0.6	0.4
2000.....	0.4	0.6	0.3	0.4	0.5	0.3	0.4	0.6	0.3
1999.....	0.4	0.5	0.4	0.4	0.5	0.4	0.4	0.5	0.3
Melanoma and other malignant neoplasms of skin (C43-C44)									
2007.....	3.5	5.4	2.1	4.0	6.0	2.4	1.0	1.3	0.7
2006.....	3.5	5.5	2.1	4.0	6.1	2.4	0.9	1.4	0.5
2005.....	3.5	5.4	2.2	4.0	5.9	2.5	0.9	1.3	0.7
2004.....	3.5	5.3	2.1	3.9	5.9	2.4	0.9	1.3	0.6
2003.....	3.5	5.2	2.2	3.9	5.8	2.4	0.9	1.1	0.7
2002.....	3.4	5.1	2.1	3.8	5.7	2.4	0.9	1.5	0.6
2001.....	3.5	5.3	2.2	3.9	5.9	2.5	1.0	1.4	0.7
2000.....	3.5	5.2	2.2	3.8	5.8	2.5	1.1	1.5	0.8
1999.....	3.5	5.2	2.2	3.9	5.8	2.5	0.9	1.6	0.6
Malignant melanoma of skin (C43)									
2007.....	2.7	4.0	1.7	3.0	4.5	1.9	0.5	0.5	0.4
2006.....	2.7	4.0	1.7	3.1	4.6	2.0	0.4	0.6	0.3
2005.....	2.7	4.0	1.8	3.1	4.5	2.1	0.4	0.5	0.4
2004.....	2.7	3.9	1.7	3.0	4.5	1.9	0.4	0.5	0.4
2003.....	2.7	3.9	1.8	3.0	4.4	2.0	0.4	0.4	0.4
2002.....	2.6	3.8	1.7	2.9	4.2	2.0	0.4	0.5	0.3
2001.....	2.7	3.9	1.8	3.0	4.3	2.0	0.4	0.3	0.4
2000.....	2.7	3.8	1.8	3.0	4.3	2.0	0.6	0.6	0.5
1999.....	2.6	3.8	1.7	3.0	4.2	2.0	0.4	0.5	0.3

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Other malignant neoplasms of skin (C44)										
2007.....	0.9	1.4	0.4	0.9	1.5	0.5	0.5	0.8	0.3	
2006.....	0.8	1.5	0.4	0.9	1.5	0.4	0.4	0.8	0.1	
2005.....	0.8	1.4	0.4	0.8	1.5	0.4	0.5	0.8	0.3	
2004.....	0.8	1.3	0.4	0.8	1.4	0.4	0.4	0.8	0.2	
2003.....	0.8	1.4	0.4	0.9	1.5	0.4	0.5	0.7	0.2	
2002.....	0.8	1.4	0.4	0.9	1.5	0.5	0.6	1.0	0.3	
2001.....	0.9	1.4	0.5	0.9	1.5	0.5	0.6	1.0	0.2	
2000.....	0.8	1.4	0.4	0.9	1.5	0.4	0.6	0.9	0.3	
1999.....	0.8	1.4	0.4	0.9	1.5	0.4	0.5	1.0	0.2	
Malignant neoplasms of mesothelial and soft tissue (C45-C49)										
2007.....	2.4	3.1	1.8	2.4	3.3	1.8	1.9	2.2	1.7	
2006.....	2.4	3.1	1.8	2.5	3.3	1.9	1.9	2.2	1.7	
2005.....	2.4	3.2	1.9	2.5	3.3	1.9	2.0	2.2	1.8	
2004.....	2.4	3.2	1.8	2.5	3.4	1.9	2.0	2.3	1.8	
2003.....	2.4	3.2	1.8	2.5	3.3	1.8	2.0	2.5	1.7	
2002.....	2.4	3.2	1.8	2.4	3.4	1.8	2.0	2.4	1.7	
2001.....	2.4	3.2	1.8	2.5	3.4	1.8	1.9	2.2	1.7	
2000.....	2.5	3.3	1.8	2.5	3.5	1.9	2.1	2.6	1.9	
1999.....	2.4	3.3	1.8	2.5	3.4	1.8	2.2	2.6	1.9	
Mesothelioma (C45)										
2007.....	0.8	1.5	0.3	0.8	1.6	0.3	0.3	0.7	*	
2006.....	0.8	1.5	0.3	0.9	1.7	0.3	0.4	0.9	0.2	
2005.....	0.9	1.6	0.3	0.9	1.7	0.3	0.4	0.8	0.1	
2004.....	0.8	1.6	0.3	0.9	1.7	0.3	0.3	0.7	*	
2003.....	0.8	1.6	0.3	0.9	1.8	0.3	0.4	0.9	*	
2002.....	0.9	1.7	0.2	0.9	1.8	0.3	0.4	0.8	*	
2001.....	0.8	1.6	0.3	0.9	1.8	0.3	0.2	0.5	*	
2000.....	0.9	1.7	0.3	0.9	1.8	0.3	0.4	0.8	0.1	
1999.....	0.9	1.6	0.3	0.9	1.8	0.3	0.4	0.8	0.2	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Kaposi's sarcoma (C46)									
2007.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2006.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2005.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2004.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2003.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2000.....	0.0	0.0	*	0.0	0.0	*	*	*	*
1999.....	0.0	0.0	*	0.0	0.0	*	*	*	*
Of other mesothelial and soft tissue (C47-C49)									
2007.....	1.6	1.6	1.5	1.6	1.6	1.5	1.6	1.5	1.6
2006.....	1.5	1.6	1.6	1.6	1.6	1.6	1.5	1.3	1.5
2005.....	1.5	1.5	1.5	1.5	1.6	1.5	1.6	1.4	1.7
2004.....	1.5	1.6	1.5	1.6	1.6	1.5	1.6	1.5	1.7
2003.....	1.5	1.5	1.5	1.5	1.5	1.5	1.6	1.6	1.6
2002.....	1.5	1.5	1.5	1.5	1.5	1.5	1.6	1.6	1.6
2001.....	1.5	1.5	1.5	1.5	1.6	1.5	1.6	1.7	1.6
2000.....	1.6	1.6	1.5	1.6	1.7	1.6	1.6	1.6	1.7
1999.....	1.6	1.6	1.5	1.5	1.6	1.5	1.7	1.8	1.7
Malignant neoplasm of breast (C50)									
2007.....	12.9	0.3	22.9	12.5	0.3	22.3	18.6	0.4	31.4
2006.....	13.2	0.3	23.5	12.8	0.3	22.9	18.7	0.5	31.6
2005.....	13.5	0.3	24.1	13.1	0.3	23.4	19.3	0.5	32.8
2004.....	13.8	0.3	24.4	13.4	0.3	23.9	19.0	0.5	32.2
2003.....	14.2	0.3	25.3	13.8	0.3	24.7	20.1	0.5	34.0
2002.....	14.5	0.3	25.6	14.1	0.3	25.0	20.2	0.6	34.0
2001.....	14.8	0.3	26.0	14.4	0.3	25.5	20.4	0.6	34.4
2000.....	15.3	0.4	26.8	14.9	0.4	26.3	20.5	0.6	34.5
1999.....	15.2	0.3	26.6	14.8	0.3	26.0	20.8	0.7	35.1

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Malignant neoplasms of female genital organs (C51-C58)										
2007.....	8.8	...	15.8	8.6	...	15.6	11.4	...	19.3	
2006.....	9.0	...	16.1	8.8	...	16.0	11.5	...	19.3	
2005.....	8.9	...	16.0	8.8	...	15.9	11.5	...	19.4	
2004.....	9.0	...	16.2	8.9	...	16.0	11.8	...	19.9	
2003.....	9.2	...	16.4	9.0	...	16.2	11.9	...	20.0	
2002.....	9.4	...	16.7	9.2	...	16.5	12.2	...	20.5	
2001.....	9.5	...	16.8	9.4	...	16.6	12.1	...	20.4	
2000.....	9.5	...	16.8	9.3	...	16.6	12.5	...	20.8	
1999.....	9.4	...	16.7	9.3	...	16.5	12.3	...	20.6	
Of vulva and vagina (C51-C52)										
2007.....	0.4	...	0.7	0.4	...	0.7	0.4	...	0.6	
2006.....	0.4	...	0.7	0.4	...	0.7	0.4	...	0.6	
2005.....	0.4	...	0.7	0.4	...	0.7	0.3	...	0.6	
2004.....	0.4	...	0.7	0.4	...	0.7	0.4	...	0.7	
2003.....	0.4	...	0.7	0.4	...	0.7	0.4	...	0.6	
2002.....	0.4	...	0.7	0.4	...	0.7	0.4	...	0.5	
2001.....	0.4	...	0.7	0.4	...	0.7	0.5	...	0.8	
2000.....	0.4	...	0.7	0.4	...	0.7	0.4	...	0.7	
1999.....	0.4	...	0.7	0.4	...	0.7	0.4	...	0.7	
Of cervix uteri (C53)										
2007.....	1.3	...	2.4	1.2	...	2.2	2.5	...	4.3	
2006.....	1.3	...	2.4	1.1	...	2.2	2.5	...	4.3	
2005.....	1.3	...	2.4	1.1	...	2.2	2.5	...	4.4	
2004.....	1.3	...	2.4	1.1	...	2.2	2.6	...	4.5	
2003.....	1.3	...	2.5	1.2	...	2.2	2.7	...	4.7	
2002.....	1.3	...	2.6	1.2	...	2.3	2.9	...	4.9	
2001.....	1.4	...	2.7	1.3	...	2.5	2.8	...	4.8	
2000.....	1.5	...	2.8	1.3	...	2.5	3.2	...	5.4	
1999.....	1.5	...	2.8	1.4	...	2.6	3.1	...	5.4	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Of corpus uteri and uterus, part unspecified (C54-C55)										
2007.....	2.3	...	4.2	2.2	...	3.9	4.5	...	7.5	
2006.....	2.4	...	4.2	2.2	...	3.9	4.4	...	7.3	
2005.....	2.3	...	4.1	2.2	...	3.8	4.2	...	7.1	
2004.....	2.3	...	4.1	2.2	...	3.9	4.1	...	6.9	
2003.....	2.3	...	4.1	2.2	...	3.9	4.2	...	7.0	
2002.....	2.4	...	4.2	2.2	...	3.9	4.4	...	7.3	
2001.....	2.4	...	4.2	2.3	...	4.0	4.2	...	7.0	
2000.....	2.4	...	4.1	2.2	...	3.9	4.4	...	7.2	
1999.....	2.4	...	4.1	2.2	...	3.9	4.2	...	6.8	
Of ovary (C56)										
2007.....	4.6	...	8.2	4.8	...	8.6	3.9	...	6.5	
2006.....	4.8	...	8.5	5.0	...	8.9	4.1	...	6.8	
2005.....	4.8	...	8.6	5.0	...	9.0	4.3	...	7.2	
2004.....	4.9	...	8.8	5.1	...	9.1	4.5	...	7.6	
2003.....	5.0	...	8.8	5.1	...	9.2	4.4	...	7.4	
2002.....	5.1	...	9.0	5.3	...	9.4	4.4	...	7.4	
2001.....	5.1	...	9.0	5.2	...	9.3	4.5	...	7.5	
2000.....	5.1	...	8.9	5.3	...	9.3	4.3	...	7.2	
1999.....	5.0	...	8.8	5.1	...	9.1	4.4	...	7.3	
Of other and unspecified female genital organs (C57)										
2007.....	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3	
2006.....	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3	
2005.....	0.1	...	0.1	0.1	...	0.1	0.1	...	0.1	
2004.....	0.1	...	0.1	0.1	...	0.1	0.1	...	0.1	
2003.....	0.1	...	0.2	0.1	...	0.2	0.1	...	0.3	
2002.....	0.1	...	0.3	0.1	...	0.3	0.2	...	0.3	
2001.....	0.1	...	0.2	0.1	...	0.2	0.1	...	0.2	
2000.....	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3	
1999.....	0.1	...	0.3	0.1	...	0.3	0.2	...	0.3	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Of placenta (C58)									
2007.....	*	...	*	*	...	*	*	...	*
2006.....	*	...	*	*	...	*	*	...	*
2005.....	*	...	*	*	...	*	*	...	*
2004.....	0.0	...	0.0	*	...	*	*	...	*
2003.....	*	...	*	*	...	*	*	...	*
2002.....	*	...	*	*	...	*	*	...	*
2001.....	*	...	*	*	...	*	*	...	*
2000.....	*	...	*	*	...	*	*	...	*
1999.....	*	...	*	*	...	*	*	...	*
Malignant neoplasms of male genital organs (C60-C63)									
2007.....	9.4	23.9	...	8.7	22.1	...	18.7	52.0	...
2006.....	9.4	23.9	...	8.7	22.2	...	18.4	50.9	...
2005.....	9.7	25.0	...	9.0	23.1	...	19.3	53.8	...
2004.....	10.0	25.9	...	9.2	23.9	...	19.9	55.9	...
2003.....	10.3	26.9	...	9.6	24.9	...	20.6	58.0	...
2002.....	10.8	28.4	...	10.0	26.2	...	22.1	62.4	...
2001.....	11.1	29.6	...	10.2	27.1	...	23.2	66.5	...
2000.....	11.5	30.8	...	10.5	28.3	...	24.1	68.6	...
1999.....	11.9	31.8	...	10.9	29.3	...	24.4	69.5	...
Of prostate (C61)									
2007.....	9.2	23.5	...	8.5	21.6	...	18.6	51.5	...
2006.....	9.2	23.5	...	8.5	21.7	...	18.2	50.5	...
2005.....	9.5	24.5	...	8.8	22.6	...	19.1	53.3	...
2004.....	9.8	25.4	...	9.0	23.4	...	19.7	55.5	...
2003.....	10.1	26.5	...	9.3	24.4	...	20.4	57.4	...
2002.....	10.6	27.9	...	9.8	25.7	...	22.0	62.0	...
2001.....	10.9	29.1	...	10.0	26.6	...	23.0	66.1	...
2000.....	11.3	30.4	...	10.3	27.8	...	23.9	68.1	...
1999.....	11.6	31.3	...	10.7	28.7	...	24.2	69.0	...

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Of testis (C62)									
2007.....	0.1	0.2	...	0.1	0.2	...	0.1	0.1	...
2006.....	0.1	0.2	...	0.1	0.3	...	0.1	0.1	...
2005.....	0.1	0.3	...	0.1	0.3	...	0.1	0.2	...
2004.....	0.1	0.3	...	0.1	0.3	...	0.1	0.2	...
2003.....	0.1	0.2	...	0.1	0.3	...	0.1	0.2	...
2002.....	0.1	0.3	...	0.2	0.3	...	0.1	0.2	...
2001.....	0.1	0.2	...	0.1	0.3	...	0.1	0.2	...
2000.....	0.1	0.3	...	0.1	0.3	...	0.1	0.2	...
1999.....	0.1	0.3	...	0.2	0.3	...	0.1	0.1	...
Of penis and other and unspecified male genital organs (C60,C63)									
2007.....	0.1	0.2	...	0.1	0.2	...	0.1	0.3	...
2006.....	0.1	0.2	...	0.1	0.2	...	0.1	0.3	...
2005.....	0.1	0.2	...	0.1	0.2	...	0.1	0.3	...
2004.....	0.1	0.2	...	0.1	0.2	...	0.1	0.2	...
2003.....	0.1	0.2	...	0.1	0.2	...	0.1	0.4	...
2002.....	0.1	0.2	...	0.1	0.2	...	0.1	0.2	...
2001.....	0.1	0.2	...	0.1	0.2	...	0.1	0.3	...
2000.....	0.1	0.2	...	0.1	0.2	...	0.1	0.3	...
1999.....	0.1	0.2	...	0.1	0.2	...	0.1	0.3	...
Malignant neoplasms of urinary tract (C64-C68)									
2007.....	8.7	13.7	5.0	8.9	14.2	5.1	8.0	11.5	5.8
2006.....	8.6	13.6	5.0	8.9	14.1	5.1	7.8	11.5	5.5
2005.....	8.7	13.7	5.1	8.9	14.2	5.2	7.9	11.9	5.3
2004.....	8.7	13.7	5.1	9.0	14.3	5.2	7.7	10.9	5.6
2003.....	8.7	13.7	5.1	8.9	14.2	5.2	8.0	11.6	5.7
2002.....	8.9	13.9	5.3	9.1	14.4	5.4	8.1	11.4	6.1
2001.....	8.8	14.1	5.2	9.0	14.5	5.2	8.2	11.7	5.9
2000.....	8.8	14.1	5.3	9.0	14.5	5.3	8.0	12.1	5.6
1999.....	8.7	13.8	5.2	8.8	14.2	5.1	8.3	12.0	6.0

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Of kidney and renal pelvis (C64-C65)										
2007.....	4.0	5.8	2.6	4.1	5.9	2.7	4.1	5.9	2.7	
2006.....	4.0	5.7	2.6	4.1	5.8	2.7	3.9	5.8	2.5	
2005.....	4.1	5.9	2.7	4.2	6.0	2.8	4.0	6.0	2.6	
2004.....	4.1	5.9	2.7	4.2	6.0	2.8	4.0	5.8	2.7	
2003.....	4.2	6.0	2.7	4.3	6.2	2.8	4.1	6.0	2.7	
2002.....	4.2	6.1	2.8	4.3	6.3	2.8	4.0	5.8	2.8	
2001.....	4.3	6.2	2.8	4.3	6.3	2.8	4.2	6.4	2.8	
2000.....	4.2	6.2	2.8	4.3	6.3	2.8	4.1	6.3	2.8	
1999.....	4.1	5.9	2.7	4.1	5.9	2.7	4.1	6.1	2.8	
Of bladder (C67)										
2007.....	4.4	7.6	2.2	4.6	8.0	2.2	3.8	5.4	2.8	
2006.....	4.3	7.5	2.2	4.5	7.9	2.2	3.8	5.5	2.8	
2005.....	4.3	7.5	2.2	4.5	7.9	2.2	3.7	5.7	2.6	
2004.....	4.4	7.6	2.2	4.5	8.0	2.3	3.6	4.9	2.7	
2003.....	4.3	7.3	2.2	4.4	7.7	2.2	3.7	5.4	2.7	
2002.....	4.4	7.5	2.4	4.5	7.8	2.4	3.9	5.5	3.0	
2001.....	4.3	7.5	2.3	4.5	7.9	2.2	3.7	5.1	2.9	
2000.....	4.3	7.6	2.3	4.5	7.9	2.3	3.7	5.6	2.7	
1999.....	4.4	7.6	2.3	4.5	7.9	2.3	4.0	5.8	3.0	
Of other and unspecified sites within the urinary tract (C66,C68)										
2007.....	0.2	0.3	0.2	0.3	0.4	0.2	0.2	0.2	0.2	
2006.....	0.3	0.4	0.2	0.3	0.4	0.2	0.2	*	0.3	
2005.....	0.2	0.3	0.2	0.2	0.3	0.2	0.2	*	0.2	
2004.....	0.2	0.3	0.2	0.2	0.3	0.1	0.2	*	0.2	
2003.....	0.2	0.3	0.2	0.2	0.3	0.2	0.2	*	0.2	
2002.....	0.2	0.3	0.2	0.2	0.3	0.2	0.2	*	0.2	
2001.....	0.2	0.3	0.2	0.2	0.3	0.2	0.2	0.2	0.2	
2000.....	0.2	0.3	0.2	0.2	0.3	0.2	0.2	*	0.2	
1999.....	0.2	0.3	0.2	0.2	0.3	0.2	0.2	*	0.2	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Malignant neoplasm of eye and adnexa (C69)									
2007.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2006.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2005.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2004.....	0.1	0.1	0.0	0.1	0.1	0.1	*	*	*
2003.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2001.....	0.1	0.1	0.0	0.1	0.1	0.1	*	*	*
2000.....	0.1	0.1	0.0	0.1	0.1	0.1	*	*	*
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)									
2007.....	4.2	5.1	3.5	4.6	5.5	3.8	2.4	2.8	2.1
2006.....	4.2	5.1	3.4	4.5	5.5	3.6	2.3	2.9	1.9
2005.....	4.3	5.3	3.5	4.7	5.7	3.8	2.5	3.2	2.0
2004.....	4.3	5.2	3.5	4.7	5.6	3.8	2.5	3.0	2.0
2003.....	4.4	5.3	3.6	4.7	5.7	3.9	2.7	3.4	2.1
2002.....	4.4	5.5	3.6	4.8	5.8	3.9	2.6	3.2	2.2
2001.....	4.5	5.5	3.6	4.8	5.9	3.9	2.6	3.1	2.2
2000.....	4.5	5.6	3.7	4.9	6.0	3.9	2.7	3.3	2.3
1999.....	4.6	5.6	3.8	5.0	6.0	4.1	2.7	3.3	2.3
Of brain (C71)									
2007.....	4.1	5.0	3.4	4.5	5.4	3.7	2.3	2.6	2.0
2006.....	4.1	5.0	3.3	4.4	5.4	3.6	2.3	2.9	1.9
2005.....	4.2	5.2	3.5	4.6	5.6	3.8	2.4	3.1	1.9
2004.....	4.2	5.1	3.4	4.6	5.5	3.8	2.4	2.9	2.0
2003.....	4.3	5.3	3.5	4.7	5.6	3.8	2.6	3.3	2.0
2002.....	4.3	5.3	3.5	4.7	5.7	3.8	2.5	3.0	2.1
2001.....	4.4	5.4	3.5	4.7	5.8	3.8	2.5	3.1	2.2
2000.....	4.5	5.5	3.6	4.8	5.9	3.9	2.6	3.2	2.2
1999.....	4.5	5.5	3.7	4.9	5.9	4.0	2.6	3.1	2.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Of other parts of central nervous system (C70,C72)										
2007.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*	
2006.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2005.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	*	*	
2004.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2003.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*	
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
Malignant neoplasms of thyroid and other endocrine glands (C73-C75)										
2007.....	0.7	0.7	0.7	0.7	0.7	0.7	0.8	0.6	0.9	
2006.....	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.5	0.8	
2005.....	0.7	0.7	0.7	0.7	0.7	0.7	0.8	0.6	0.9	
2004.....	0.7	0.7	0.7	0.7	0.7	0.7	0.6	0.6	0.7	
2003.....	0.7	0.7	0.7	0.7	0.7	0.7	0.6	0.5	0.6	
2002.....	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.6	0.7	
2001.....	0.8	0.8	0.7	0.7	0.8	0.7	0.7	0.6	0.8	
2000.....	0.8	0.8	0.7	0.7	0.8	0.7	0.8	0.6	0.9	
1999.....	0.7	0.8	0.7	0.7	0.8	0.7	0.7	0.7	0.7	
Other malignant neoplasms of other and unspecified sites (C76-C80,C97)										
2007.....	12.5	15.3	10.4	12.4	15.1	10.3	15.1	19.5	12.1	
2006.....	12.8	15.5	10.8	12.7	15.4	10.7	15.6	20.0	12.7	
2005.....	13.1	15.9	11.1	13.0	15.7	11.1	16.0	20.8	12.8	
2004.....	13.3	16.1	11.2	13.2	15.8	11.1	16.8	22.1	13.2	
2003.....	13.7	16.6	11.6	13.6	16.4	11.6	17.2	22.0	14.0	
2002.....	14.1	17.2	11.9	13.9	16.9	11.7	18.2	23.1	15.0	
2001.....	14.2	17.2	12.0	13.9	16.8	11.9	18.4	24.3	14.5	
2000.....	14.7	17.8	12.5	14.5	17.4	12.4	18.7	24.8	14.7	
1999.....	16.0	19.5	13.6	15.7	19.1	13.4	20.6	26.5	16.8	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/		Black 2/			
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)									
2007.....	17.6	22.8	13.7	17.8	23.2	13.8	17.6	21.9	14.8
2006.....	17.9	23.0	14.1	18.2	23.4	14.3	18.0	23.6	14.3
2005.....	18.2	23.5	14.3	18.5	24.0	14.5	17.6	22.5	14.3
2004.....	18.4	23.6	14.6	18.6	24.0	14.7	18.9	23.6	15.8
2003.....	19.1	24.5	15.1	19.4	25.1	15.2	18.8	22.9	16.1
2002.....	19.5	25.1	15.5	19.9	25.6	15.7	19.4	24.4	16.1
2001.....	20.0	25.5	16.0	20.2	25.9	16.2	20.0	25.3	16.5
2000.....	20.4	26.0	16.5	20.7	26.4	16.7	19.9	25.0	16.6
1999.....	20.6	26.1	16.7	20.8	26.5	16.8	20.3	25.6	16.8
Hodgkin's disease (C81)									
2007.....	0.4	0.5	0.3	0.4	0.5	0.3	0.4	0.5	0.3
2006.....	0.4	0.6	0.3	0.4	0.6	0.4	0.4	0.6	0.3
2005.....	0.4	0.5	0.4	0.4	0.5	0.4	0.4	0.5	0.3
2004.....	0.4	0.5	0.3	0.4	0.6	0.3	0.4	0.5	0.4
2003.....	0.5	0.5	0.4	0.5	0.6	0.4	0.4	0.5	0.3
2002.....	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.6	0.3
2001.....	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.5	0.4
2000.....	0.5	0.6	0.4	0.5	0.6	0.4	0.3	0.4	0.3
1999.....	0.5	0.6	0.4	0.5	0.7	0.4	0.4	0.5	0.3
Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)									
2007.....	6.6	8.3	5.2	6.8	8.6	5.5	4.6	5.8	3.6
2006.....	6.7	8.4	5.4	7.0	8.7	5.6	4.9	6.3	3.9
2005.....	6.9	8.8	5.5	7.2	9.1	5.8	4.7	6.0	3.8
2004.....	7.1	8.9	5.7	7.4	9.2	6.0	5.0	6.2	4.1
2003.....	7.3	9.3	5.9	7.7	9.7	6.2	4.9	5.8	4.3
2002.....	7.6	9.6	6.2	7.9	10.0	6.5	5.4	6.8	4.4
2001.....	7.9	10.0	6.4	8.2	10.3	6.7	5.6	7.3	4.5
2000.....	8.3	10.3	6.8	8.6	10.7	7.1	5.2	6.7	4.2
1999.....	8.4	10.4	6.9	8.7	10.7	7.2	5.6	7.5	4.4

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Non-Hodgkin's lymphoma (C82-C85)										
2007.....	6.5	8.3	5.2	6.8	8.6	5.5	4.5	5.7	3.6	
2006.....	6.7	8.4	5.4	7.0	8.7	5.6	4.9	6.2	3.9	
2005.....	6.9	8.7	5.5	7.2	9.1	5.8	4.7	6.0	3.8	
2004.....	7.0	8.8	5.7	7.3	9.2	6.0	5.0	6.2	4.1	
2003.....	7.3	9.3	5.9	7.7	9.7	6.1	4.9	5.8	4.2	
2002.....	7.6	9.6	6.2	7.9	10.0	6.5	5.4	6.8	4.4	
2001.....	7.9	9.9	6.4	8.2	10.2	6.7	5.6	7.2	4.5	
2000.....	8.2	10.2	6.7	8.6	10.7	7.1	5.2	6.6	4.2	
1999.....	8.3	10.3	6.9	8.7	10.7	7.2	5.6	7.4	4.4	
Of other and unspecified lymphoid, hematopoietic and related tissue (C96)										
2007.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2006.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2005.....	0.0	0.0	0.0	0.0	0.0	*	*	*	*	
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2003.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
Malignant immunoproliferative diseases (C88)										
2007.....	0.1	0.2	0.1	0.1	0.2	0.1	*	*	*	
2006.....	0.1	0.2	0.1	0.1	0.2	0.1	*	*	*	
2005.....	0.1	0.2	0.1	0.1	0.2	0.1	*	*	*	
2004.....	0.1	0.1	0.1	0.1	0.2	0.1	*	*	*	
2003.....	0.1	0.2	0.1	0.1	0.2	0.1	*	*	*	
2002.....	0.1	0.2	0.1	0.1	0.2	0.1	*	*	*	
2001.....	0.1	0.2	0.1	0.1	0.2	0.1	*	*	*	
2000.....	0.1	0.2	0.1	0.1	0.2	0.1	*	*	*	
1999.....	0.2	0.2	0.1	0.2	0.2	0.1	0.1	*	*	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/		Black 2/			
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Multiple myeloma and malignant plasma cell neoplasms (C90)									
2007.....	3.5	4.4	2.9	3.2	4.2	2.6	6.6	7.8	5.9
2006.....	3.5	4.4	2.9	3.3	4.1	2.6	6.5	8.1	5.4
2005.....	3.6	4.5	2.9	3.3	4.2	2.7	6.5	8.0	5.5
2004.....	3.6	4.4	3.0	3.3	4.2	2.7	6.8	8.2	6.0
2003.....	3.7	4.6	3.1	3.5	4.3	2.8	7.1	8.3	6.3
2002.....	3.8	4.8	3.2	3.6	4.5	2.9	6.9	8.2	6.0
2001.....	3.8	4.7	3.2	3.6	4.4	2.9	7.3	8.7	6.3
2000.....	3.9	4.7	3.3	3.6	4.4	3.0	7.6	8.8	6.8
1999.....	3.9	4.7	3.3	3.5	4.4	2.9	7.5	8.8	6.6
Leukemia (C91-C95)									
2007.....	7.0	9.4	5.3	7.2	9.7	5.4	6.1	7.9	5.0
2006.....	7.1	9.5	5.4	7.4	9.8	5.6	6.1	8.6	4.6
2005.....	7.1	9.6	5.4	7.4	10.0	5.6	5.9	7.8	4.7
2004.....	7.2	9.6	5.5	7.4	9.9	5.6	6.7	8.7	5.3
2003.....	7.4	9.9	5.6	7.6	10.2	5.7	6.4	8.3	5.2
2002.....	7.5	10.0	5.7	7.7	10.3	5.8	6.6	8.6	5.3
2001.....	7.6	10.1	5.9	7.8	10.4	6.0	6.7	8.7	5.4
2000.....	7.7	10.3	5.9	7.9	10.5	6.1	6.7	9.0	5.2
1999.....	7.6	10.2	6.0	7.8	10.4	6.1	6.7	8.7	5.4
Lymphoid leukemia (C91)									
2007.....	2.0	2.8	1.4	2.1	3.0	1.5	1.8	2.6	1.3
2006.....	2.1	2.9	1.5	2.1	3.0	1.5	1.9	3.0	1.2
2005.....	2.1	2.9	1.5	2.2	3.0	1.5	1.8	2.6	1.2
2004.....	2.1	2.9	1.5	2.1	3.0	1.5	1.9	2.9	1.3
2003.....	2.1	3.1	1.5	2.2	3.2	1.5	1.8	2.6	1.3
2002.....	2.2	3.1	1.6	2.3	3.2	1.6	1.9	2.7	1.4
2001.....	2.2	3.1	1.6	2.3	3.1	1.6	2.1	2.9	1.5
2000.....	2.2	3.1	1.6	2.3	3.2	1.6	1.9	2.8	1.3
1999.....	2.3	3.2	1.6	2.4	3.3	1.7	2.0	2.9	1.4

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Myeloid leukemia (C92)										
2007.....	3.2	4.3	2.5	3.4	4.5	2.6	2.7	3.0	2.4	
2006.....	3.3	4.3	2.6	3.5	4.5	2.7	2.8	3.6	2.3	
2005.....	3.3	4.3	2.6	3.4	4.5	2.6	2.6	3.1	2.2	
2004.....	3.4	4.3	2.7	3.4	4.5	2.7	2.9	3.4	2.5	
2003.....	3.4	4.4	2.7	3.5	4.6	2.7	2.9	3.7	2.4	
2002.....	3.4	4.5	2.7	3.5	4.6	2.8	2.8	3.7	2.3	
2001.....	3.5	4.5	2.8	3.6	4.7	2.9	2.9	3.6	2.4	
2000.....	3.5	4.6	2.8	3.6	4.7	2.8	3.0	3.7	2.6	
1999.....	3.4	4.3	2.7	3.4	4.4	2.8	2.8	3.6	2.3	
Monocytic leukemia (C93)										
2007.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2006.....	0.0	0.1	0.0	0.0	0.1	0.0	*	*	*	
2005.....	0.0	0.1	0.0	0.0	0.1	0.0	*	*	*	
2004.....	0.0	0.1	0.0	0.0	0.1	0.0	*	*	*	
2003.....	0.0	0.1	0.0	0.1	0.1	0.0	*	*	*	
2002.....	0.0	0.1	0.0	0.0	0.1	0.0	*	*	*	
2001.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*	
2000.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*	
1999.....	0.0	0.1	0.0	0.1	0.1	0.0	*	*	*	
Other and unspecified leukemia (C94-C95)										
2007.....	1.7	2.2	1.3	1.7	2.2	1.3	1.6	2.2	1.3	
2006.....	1.7	2.2	1.3	1.8	2.3	1.3	1.5	2.0	1.1	
2005.....	1.7	2.3	1.3	1.8	2.4	1.4	1.6	2.1	1.2	
2004.....	1.7	2.4	1.3	1.8	2.4	1.3	1.8	2.3	1.5	
2003.....	1.8	2.3	1.4	1.8	2.4	1.4	1.7	2.0	1.4	
2002.....	1.8	2.4	1.4	1.8	2.4	1.4	1.8	2.2	1.5	
2001.....	1.8	2.4	1.4	1.9	2.5	1.4	1.7	2.2	1.4	
2000.....	1.9	2.5	1.5	1.9	2.5	1.6	1.7	2.4	1.3	
1999.....	1.9	2.5	1.5	2.0	2.6	1.5	1.8	2.1	1.6	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Benign neoplasms (D10-D36)										
2007.....	0.5	0.4	0.4	0.4	0.4	0.4	0.8	0.7	0.8	
2006.....	0.5	0.4	0.5	0.4	0.4	0.5	0.7	0.7	0.7	
2005.....	0.5	0.4	0.5	0.4	0.4	0.5	0.8	0.6	0.9	
2004.....	0.5	0.5	0.6	0.5	0.4	0.5	0.9	0.9	0.9	
2003.....	0.5	0.5	0.5	0.5	0.5	0.5	0.8	0.8	0.7	
2002.....	0.5	0.5	0.6	0.5	0.5	0.5	0.8	0.9	0.9	
2001.....	0.6	0.6	0.6	0.5	0.5	0.5	1.0	1.1	0.9	
2000.....	0.6	0.5	0.6	0.5	0.5	0.6	0.9	0.8	1.0	
1999.....	0.6	0.6	0.6	0.5	0.5	0.5	1.0	1.0	0.9	
Of female genital organs (D25-D28)										
2007.....	0.0	...	0.0	0.0	...	0.0	*	...	*	
2006.....	0.0	...	0.0	0.0	...	0.0	0.1	...	0.1	
2005.....	0.0	...	0.0	*	...	*	0.1	...	0.1	
2004.....	0.0	...	0.0	0.0	...	0.0	*	...	*	
2003.....	0.0	...	0.0	0.0	...	0.0	*	...	*	
2002.....	0.0	...	0.0	0.0	...	0.0	0.1	...	0.1	
2001.....	0.0	...	0.0	0.0	...	0.0	*	...	*	
2000.....	0.0	...	0.0	0.0	...	0.0	0.1	...	0.2	
1999.....	0.0	...	0.0	*	...	*	*	...	*	
Of eye, brain and other parts of central nervous system (D31-D33)										
2007.....	0.3	0.3	0.3	0.3	0.2	0.3	0.5	0.4	0.5	
2006.....	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.4	0.4	
2005.....	0.3	0.2	0.3	0.2	0.2	0.3	0.5	0.3	0.5	
2004.....	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.5	0.5	
2003.....	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.4	0.4	
2002.....	0.3	0.3	0.4	0.3	0.3	0.4	0.5	0.4	0.5	
2001.....	0.3	0.3	0.3	0.3	0.3	0.3	0.6	0.6	0.6	
2000.....	0.3	0.3	0.4	0.3	0.3	0.4	0.5	0.4	0.5	
1999.....	0.3	0.3	0.3	0.3	0.3	0.3	0.6	0.5	0.6	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Of other and unspecified sites (D10-D24,D29-D30,D34-D36)										
2007.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.3	0.2	
2006.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3	
2005.....	0.2	0.2	0.2	0.1	0.2	0.2	0.3	0.3	0.3	
2004.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.2	
2003.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.2	
2002.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.3	
2001.....	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.5	0.2	
2000.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.3	
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.3	
Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)										
2007.....	3.8	4.9	3.0	3.9	5.0	3.1	3.0	4.0	2.3	
2006.....	3.8	4.9	3.0	3.9	5.0	3.1	3.2	4.2	2.7	
2005.....	3.8	4.9	3.0	3.9	5.0	3.1	3.1	3.7	2.7	
2004.....	3.8	4.9	3.0	3.9	5.1	3.1	3.2	4.0	2.7	
2003.....	3.9	5.0	3.2	4.0	5.2	3.2	3.3	4.0	2.8	
2002.....	3.8	5.0	3.1	4.0	5.2	3.2	3.3	4.1	2.9	
2001.....	4.1	5.3	3.3	4.2	5.4	3.4	3.4	4.3	2.9	
2000.....	4.1	5.3	3.3	4.2	5.4	3.4	3.4	4.4	2.9	
1999.....	4.1	5.3	3.3	4.2	5.5	3.4	3.6	4.1	3.2	
Neoplasms of uncertain or unknown behavior of unspecified sites (D48)										
2007.....	0.3	0.3	0.3	0.3	0.3	0.2	0.4	0.4	0.3	
2006.....	0.3	0.3	0.3	0.3	0.2	0.3	0.4	0.4	0.4	
2005.....	0.3	0.3	0.2	0.3	0.3	0.3	0.3	0.3	0.2	
2004.....	0.3	0.2	0.2	0.3	0.2	0.2	0.3	0.3	0.3	
2003.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	
2002.....	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.2	
2001.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	
2000.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2	
1999.....	0.2	0.2	0.2	0.2	0.1	0.2	0.3	0.3	0.3	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)									
2007.....	3.1	3.2	3.0	2.7	2.8	2.6	6.5	6.7	6.4
2006.....	2.8	2.9	2.8	2.4	2.5	2.3	6.5	6.2	6.7
2005.....	3.1	3.2	3.1	2.7	2.7	2.6	7.1	7.2	7.0
2004.....	3.1	3.2	3.1	2.7	2.8	2.6	7.0	7.2	6.8
2003.....	3.3	3.3	3.2	2.8	2.9	2.6	7.5	7.7	7.3
2002.....	3.4	3.4	3.3	2.8	2.9	2.8	7.5	7.7	7.4
2001.....	3.4	3.4	3.3	2.9	3.0	2.8	7.4	7.2	7.5
2000.....	3.4	3.4	3.3	2.9	3.0	2.8	7.4	7.2	7.5
1999.....	3.3	3.4	3.2	2.8	3.0	2.7	7.3	7.3	7.2
Anemias (D50-D64)									
2007.....	1.5	1.5	1.5	1.3	1.3	1.3	3.0	3.1	2.9
2006.....	1.3	1.3	1.3	1.1	1.1	1.1	2.9	2.8	2.8
2005.....	1.5	1.5	1.5	1.3	1.2	1.3	3.3	3.4	3.2
2004.....	1.5	1.5	1.5	1.3	1.3	1.3	3.3	3.5	3.1
2003.....	1.6	1.6	1.5	1.3	1.3	1.3	3.5	3.9	3.2
2002.....	1.6	1.6	1.6	1.3	1.3	1.4	3.7	3.9	3.5
2001.....	1.6	1.6	1.6	1.4	1.4	1.4	3.5	3.6	3.4
2000.....	1.6	1.6	1.6	1.4	1.4	1.4	3.4	3.4	3.5
1999.....	1.6	1.7	1.6	1.4	1.4	1.4	3.6	3.7	3.5
Nutritional anemias (D50-D53)									
2007.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2006.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
2005.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2004.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2003.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	*	0.2
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	*	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Hemolytic anemias (D55-D59)									
2007.....	0.3	0.3	0.2	0.1	0.1	0.1	1.3	1.4	1.2
2006.....	0.3	0.3	0.3	0.1	0.1	0.1	1.3	1.3	1.3
2005.....	0.3	0.3	0.3	0.1	0.1	0.1	1.4	1.5	1.4
2004.....	0.3	0.3	0.3	0.1	0.1	0.1	1.5	1.6	1.4
2003.....	0.3	0.3	0.3	0.1	0.1	0.1	1.4	1.5	1.3
2002.....	0.3	0.3	0.3	0.1	0.1	0.1	1.5	1.5	1.5
2001.....	0.3	0.3	0.3	0.1	0.1	0.1	1.4	1.5	1.3
2000.....	0.3	0.3	0.3	0.1	0.1	0.1	1.4	1.3	1.5
1999.....	0.3	0.3	0.3	0.1	0.1	0.1	1.5	1.7	1.4
Aplastic anemias (D60-D61)									
2007.....	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.4	0.4
2006.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
2005.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.2
2004.....	0.3	0.4	0.3	0.3	0.3	0.3	0.3	0.4	0.3
2003.....	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.4	0.4
2002.....	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.5	0.4
2001.....	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.3
2000.....	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.6	0.4
1999.....	0.4	0.5	0.4	0.4	0.4	0.4	0.5	0.5	0.4
Acute posthemorrhagic and other anemias (D62,D64)									
2007.....	0.8	0.8	0.9	0.8	0.8	0.8	1.2	1.2	1.2
2006.....	0.6	0.6	0.6	0.6	0.6	0.6	1.1	1.1	1.1
2005.....	0.8	0.8	0.8	0.8	0.7	0.8	1.5	1.5	1.5
2004.....	0.8	0.8	0.8	0.8	0.7	0.8	1.4	1.5	1.3
2003.....	0.8	0.8	0.8	0.8	0.8	0.8	1.5	1.8	1.3
2002.....	0.8	0.8	0.9	0.8	0.7	0.8	1.7	1.8	1.5
2001.....	0.9	0.8	0.9	0.8	0.7	0.8	1.5	1.5	1.6
2000.....	0.8	0.8	0.8	0.8	0.7	0.8	1.4	1.4	1.4
1999.....	0.8	0.8	0.9	0.8	0.7	0.8	1.5	1.3	1.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)									
2007.....	0.8	0.9	0.8	0.8	0.9	0.8	1.3	1.4	1.3
2006.....	0.8	0.9	0.8	0.8	0.8	0.7	1.4	1.3	1.4
2005.....	0.9	1.0	0.8	0.8	0.9	0.8	1.4	1.5	1.3
2004.....	0.9	0.9	0.8	0.8	0.9	0.8	1.4	1.5	1.3
2003.....	0.9	1.0	0.9	0.9	0.9	0.8	1.5	1.5	1.4
2002.....	0.9	1.0	0.9	0.9	1.0	0.8	1.4	1.4	1.4
2001.....	0.9	1.0	0.9	0.9	0.9	0.8	1.5	1.4	1.5
2000.....	1.0	1.0	0.9	0.9	1.0	0.9	1.5	1.5	1.5
1999.....	1.0	1.0	0.9	0.9	0.9	0.8	1.6	1.8	1.5
Other diseases of blood and blood-forming organs (D70-D76)									
2007.....	0.3	0.3	0.2	0.2	0.3	0.2	0.3	0.4	0.3
2006.....	0.3	0.3	0.2	0.3	0.3	0.2	0.3	0.3	0.2
2005.....	0.3	0.3	0.2	0.3	0.3	0.2	0.3	0.3	0.3
2004.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.2
2003.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
2002.....	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.4	0.3
2001.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
2000.....	0.3	0.4	0.3	0.3	0.3	0.3	0.3	0.4	0.3
1999.....	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.4	0.3
Certain disorders involving the immune mechanism (D80-D89)									
2007.....	0.5	0.5	0.5	0.3	0.3	0.3	1.9	1.8	2.0
2006.....	0.5	0.4	0.5	0.3	0.3	0.3	2.1	1.8	2.2
2005.....	0.5	0.5	0.5	0.3	0.3	0.3	2.1	2.0	2.2
2004.....	0.5	0.5	0.5	0.3	0.3	0.3	2.0	1.9	2.1
2003.....	0.5	0.5	0.5	0.3	0.3	0.3	2.2	2.0	2.4
2002.....	0.5	0.5	0.5	0.3	0.3	0.3	2.2	2.0	2.3
2001.....	0.5	0.5	0.5	0.3	0.3	0.3	2.1	1.9	2.4
2000.....	0.5	0.5	0.5	0.3	0.3	0.3	2.1	1.9	2.2
1999.....	0.4	0.4	0.4	0.3	0.3	0.2	1.7	1.5	1.9

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
IV. Endocrine, nutritional and metabolic diseases (E00-E88)									
2007.....	31.2	35.7	27.5	29.0	33.8	25.0	54.5	58.4	51.1
2006.....	31.8	36.2	28.3	29.6	34.2	25.8	56.5	61.4	52.6
2005.....	33.7	37.8	30.3	31.4	35.8	27.7	59.2	63.1	56.0
2004.....	33.1	37.2	29.8	30.7	35.1	27.2	59.8	63.6	56.8
2003.....	34.2	38.0	31.1	31.7	36.0	28.3	61.3	63.1	59.3
2002.....	34.4	37.7	31.7	31.9	35.7	28.8	62.3	62.3	61.2
2001.....	34.3	37.1	31.8	31.8	35.0	29.0	61.7	63.0	60.1
2000.....	34.1	36.9	31.8	31.6	34.9	28.9	62.1	61.4	61.8
1999.....	34.0	36.6	31.8	31.3	34.4	28.8	62.2	63.0	61.3
Endocrine diseases (E00-E34)									
2007.....	23.4	27.0	20.5	21.3	25.2	18.2	44.1	47.1	41.6
2006.....	24.2	28.0	21.2	22.0	26.0	18.9	46.5	51.0	42.9
2005.....	25.6	29.1	22.7	23.4	27.2	20.4	48.3	52.0	45.3
2004.....	25.4	28.9	22.7	23.2	26.8	20.2	49.4	52.6	46.7
2003.....	26.3	29.6	23.7	24.0	27.6	21.1	50.5	51.9	48.9
2002.....	26.4	29.3	24.1	24.1	27.4	21.5	50.8	50.4	50.0
2001.....	26.3	28.9	24.2	24.0	27.0	21.7	50.6	51.2	49.5
2000.....	26.0	28.4	24.1	23.7	26.6	21.5	51.0	49.9	51.0
1999.....	26.0	28.5	24.1	23.5	26.5	21.3	51.2	50.8	51.0
Disorders of thyroid gland (E00-E07)									
2007.....	0.6	0.3	0.7	0.5	0.3	0.7	0.7	0.5	0.8
2006.....	0.6	0.3	0.7	0.6	0.3	0.7	0.7	0.5	0.7
2005.....	0.6	0.4	0.8	0.6	0.3	0.8	0.7	0.4	0.9
2004.....	0.6	0.4	0.7	0.6	0.4	0.7	0.8	0.5	0.9
2003.....	0.7	0.4	0.8	0.7	0.3	0.8	0.7	0.5	0.9
2002.....	0.7	0.4	0.8	0.7	0.4	0.8	0.7	0.4	0.9
2001.....	0.7	0.4	0.8	0.7	0.4	0.8	0.8	0.6	0.9
2000.....	0.6	0.4	0.8	0.6	0.3	0.8	0.8	0.6	1.0
1999.....	0.7	0.4	0.8	0.6	0.4	0.8	0.9	0.6	1.0

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Diseases of thymus (E32)									
2007.....	*	*	*	*	*	*	*	*	*
2006.....	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Other endocrine diseases (E15-E21,E28-E31,E34)									
2007.....	0.2	0.2	0.2	0.1	0.1	0.1	0.4	0.4	0.5
2006.....	0.2	0.1	0.1	0.1	0.1	0.1	0.4	0.5	0.4
2005.....	0.1	0.1	0.1	0.1	0.1	0.1	0.4	0.5	0.4
2004.....	0.1	0.2	0.1	0.1	0.1	0.1	0.4	0.5	0.3
2003.....	0.2	0.1	0.1	0.1	0.1	0.1	0.4	0.4	0.3
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.3	0.3	0.4
2001.....	0.2	0.2	0.1	0.1	0.1	0.1	0.3	0.4	0.3
2000.....	0.1	0.1	0.2	0.1	0.1	0.1	0.4	0.3	0.4
1999.....	0.2	0.2	0.1	0.1	0.1	0.1	0.3	0.5	0.3
Nutritional diseases (E40-E68)									
2007.....	2.4	2.5	2.3	2.3	2.3	2.2	3.9	3.8	4.0
2006.....	2.3	2.2	2.3	2.2	2.2	2.2	3.8	3.7	3.9
2005.....	2.5	2.5	2.6	2.4	2.4	2.4	4.3	3.7	4.7
2004.....	2.4	2.3	2.4	2.2	2.2	2.2	4.2	3.8	4.5
2003.....	2.5	2.4	2.5	2.4	2.3	2.4	4.0	3.8	4.2
2002.....	2.6	2.5	2.6	2.4	2.4	2.4	4.5	4.0	4.7
2001.....	2.4	2.2	2.5	2.3	2.1	2.3	4.1	4.0	4.3
2000.....	2.6	2.5	2.6	2.5	2.4	2.5	4.3	4.2	4.4
1999.....	2.5	2.5	2.5	2.4	2.3	2.4	4.4	4.5	4.3

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Malnutrition (E40-E46)									
2007.....	0.8	0.8	0.8	0.8	0.7	0.8	1.4	1.6	1.3
2006.....	0.7	0.7	0.8	0.7	0.6	0.7	1.2	1.5	1.0
2005.....	1.0	0.9	1.0	0.9	0.8	1.0	1.6	1.5	1.6
2004.....	0.9	0.9	1.0	0.9	0.8	0.9	1.6	1.8	1.5
2003.....	1.1	1.0	1.1	1.0	0.9	1.1	1.8	2.1	1.7
2002.....	1.2	1.1	1.2	1.1	1.1	1.2	2.0	2.0	2.0
2001.....	1.2	1.1	1.3	1.2	1.0	1.2	1.9	2.2	1.7
2000.....	1.4	1.3	1.5	1.4	1.2	1.4	2.2	2.5	2.1
1999.....	1.5	1.4	1.5	1.4	1.3	1.4	2.5	2.9	2.2
Nutritional marasmus (E41)									
2007.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2006.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2005.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.0	0.1	0.0	0.0	0.1	*	*	*
2002.....	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*	*
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
Other and unspecified malnutrition (E40,E42-E46)									
2007.....	0.8	0.8	0.8	0.7	0.7	0.8	1.4	1.6	1.2
2006.....	0.7	0.7	0.7	0.7	0.6	0.7	1.2	1.4	1.0
2005.....	0.9	0.9	1.0	0.9	0.8	0.9	1.6	1.5	1.6
2004.....	0.9	0.9	0.9	0.8	0.8	0.9	1.6	1.8	1.4
2003.....	1.0	0.9	1.0	1.0	0.9	1.0	1.8	2.0	1.6
2002.....	1.1	1.1	1.1	1.1	1.0	1.1	1.9	1.9	1.9
2001.....	1.1	1.0	1.2	1.1	1.0	1.2	1.8	2.1	1.6
2000.....	1.3	1.2	1.4	1.3	1.2	1.3	2.1	2.3	2.0
1999.....	1.4	1.3	1.4	1.3	1.2	1.3	2.3	2.8	2.0

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Other nutritional deficiencies (E50-E64)									
2007.....	0.1	0.1	0.1	0.1	0.1	0.0	*	*	*
2006.....	0.1	0.0	0.0	0.0	0.0	0.0	*	*	*
2005.....	0.1	0.0	0.1	0.1	0.0	0.1	*	*	*
2004.....	0.1	0.0	0.0	0.1	0.0	0.1	*	*	*
2003.....	0.1	0.1	0.0	0.1	0.1	0.1	*	*	*
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	*	0.2
Obesity and other hyperalimentation (E65-E68)									
2007.....	1.5	1.6	1.5	1.4	1.6	1.3	2.5	2.2	2.7
2006.....	1.5	1.5	1.5	1.4	1.5	1.4	2.5	2.2	2.8
2005.....	1.5	1.5	1.5	1.4	1.5	1.4	2.6	2.1	3.0
2004.....	1.4	1.4	1.4	1.3	1.3	1.3	2.6	2.0	3.0
2003.....	1.3	1.3	1.4	1.3	1.3	1.3	2.1	1.7	2.5
2002.....	1.3	1.2	1.3	1.2	1.2	1.1	2.3	1.8	2.7
2001.....	1.1	1.0	1.2	1.0	1.0	1.0	2.1	1.6	2.5
2000.....	1.1	1.1	1.1	1.0	1.0	0.9	1.9	1.6	2.2
1999.....	0.9	0.9	1.0	0.9	0.9	0.8	1.7	1.3	2.0
Metabolic disorders (E70-E88)									
2007.....	5.4	6.2	4.7	5.4	6.2	4.6	6.4	7.5	5.5
2006.....	5.3	6.0	4.7	5.3	6.0	4.7	6.2	6.8	5.8
2005.....	5.6	6.2	5.0	5.6	6.3	5.0	6.6	7.4	6.0
2004.....	5.3	6.0	4.7	5.3	6.0	4.7	6.2	7.1	5.5
2003.....	5.4	6.0	4.8	5.4	6.0	4.8	6.7	7.4	6.1
2002.....	5.4	5.9	5.0	5.4	5.9	4.9	7.1	8.0	6.5
2001.....	5.5	6.0	5.0	5.5	6.0	5.0	7.0	7.9	6.3
2000.....	5.5	5.9	5.0	5.5	5.9	5.0	6.8	7.3	6.3
1999.....	5.5	5.7	5.1	5.4	5.6	5.1	6.6	7.7	6.0

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Cystic fibrosis (E84)									
2007.....	0.2	0.2	0.1	0.2	0.2	0.2	*	*	*
2006.....	0.1	0.1	0.1	0.2	0.2	0.2	*	*	*
2005.....	0.1	0.1	0.1	0.2	0.2	0.2	*	*	*
2004.....	0.2	0.2	0.2	0.2	0.2	0.2	*	*	*
2003.....	0.2	0.2	0.1	0.2	0.2	0.2	0.0	*	*
2002.....	0.2	0.2	0.2	0.2	0.2	0.2	0.1	*	*
2001.....	0.2	0.2	0.2	0.2	0.2	0.2	*	*	*
2000.....	0.2	0.2	0.2	0.2	0.2	0.2	*	*	*
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	*	*	*
Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)									
2007.....	1.8	1.7	1.8	1.7	1.6	1.8	2.8	3.0	2.5
2006.....	1.9	1.7	2.0	1.9	1.6	2.0	3.0	3.1	2.9
2005.....	2.2	2.0	2.2	2.1	2.0	2.2	3.3	3.5	3.1
2004.....	2.2	2.0	2.2	2.1	1.9	2.2	3.2	3.5	2.9
2003.....	2.4	2.2	2.4	2.3	2.1	2.4	3.7	3.9	3.6
2002.....	2.5	2.3	2.6	2.4	2.1	2.6	4.0	4.2	3.8
2001.....	2.7	2.5	2.8	2.6	2.3	2.8	4.4	4.5	4.2
2000.....	2.8	2.7	2.8	2.7	2.6	2.7	4.2	4.5	4.0
1999.....	3.0	2.8	3.1	2.9	2.6	3.1	4.3	5.1	3.7
Other metabolic disorders (E70-E83,E85,E88)									
2007.....	3.4	4.3	2.7	3.4	4.4	2.7	3.6	4.4	3.0
2006.....	3.2	4.1	2.6	3.3	4.2	2.6	3.2	3.7	2.9
2005.....	3.3	4.1	2.6	3.3	4.2	2.6	3.3	3.9	2.8
2004.....	3.0	3.8	2.3	3.1	3.9	2.4	3.0	3.5	2.6
2003.....	2.9	3.7	2.3	2.9	3.8	2.3	2.9	3.4	2.5
2002.....	2.8	3.5	2.2	2.8	3.5	2.2	3.1	3.7	2.6
2001.....	2.6	3.4	2.0	2.7	3.4	2.1	2.6	3.3	2.1
2000.....	2.5	3.1	2.0	2.5	3.2	2.0	2.5	2.7	2.3
1999.....	2.3	2.8	1.9	2.3	2.8	1.9	2.4	2.5	2.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
V. Mental and behavioral disorders (F01-F99)									
2007.....	27.4	25.6	27.5	27.7	25.7	28.0	29.8	31.1	28.0
2006.....	28.6	27.4	28.2	28.8	27.3	28.7	31.1	33.6	28.4
2005.....	22.8	22.0	22.3	23.1	22.0	22.7	24.6	26.4	22.5
2004.....	20.3	19.9	19.5	20.5	19.7	19.9	21.7	25.1	18.6
2003.....	20.2	19.6	19.6	20.4	19.5	20.0	21.3	24.0	18.5
2002.....	19.2	18.8	18.4	19.3	18.6	18.8	20.1	23.5	17.0
2001.....	18.0	17.8	17.1	18.1	17.6	17.4	18.9	22.7	15.6
2000.....	16.7	16.8	15.7	16.8	16.5	16.0	17.9	22.0	14.4
1999.....	15.5	16.0	14.3	15.6	15.8	14.5	15.9	20.4	12.3
Organic dementia (F01,F03)									
2007.....	23.6	20.4	25.1	23.9	20.5	25.5	25.2	23.8	25.5
2006.....	23.6	20.4	25.1	23.8	20.4	25.5	25.0	24.2	24.9
2005.....	17.8	15.0	19.2	18.1	15.1	19.5	18.6	17.1	19.0
2004.....	15.3	12.9	16.4	15.6	13.0	16.8	15.5	15.1	15.4
2003.....	15.3	12.8	16.4	15.6	12.9	16.8	15.0	14.3	15.0
2002.....	14.3	12.2	15.3	14.6	12.3	15.7	13.5	13.2	13.4
2001.....	13.2	11.4	14.0	13.5	11.5	14.3	12.0	11.7	12.0
2000.....	12.0	10.3	12.6	12.2	10.3	12.9	11.1	11.3	10.8
1999.....	10.6	9.3	11.1	10.9	9.5	11.5	8.7	8.9	8.4
Other and unspecified organic mental disorders (F04-F09)									
2007.....	0.5	0.4	0.5	0.5	0.4	0.5	0.4	0.5	0.4
2006.....	0.5	0.5	0.5	0.5	0.5	0.5	0.4	0.5	0.4
2005.....	0.5	0.5	0.6	0.6	0.5	0.6	0.6	0.5	0.5
2004.....	0.6	0.6	0.6	0.6	0.6	0.6	0.5	0.5	0.5
2003.....	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.7	0.5
2002.....	0.7	0.6	0.7	0.7	0.6	0.7	0.6	0.5	0.6
2001.....	0.7	0.7	0.8	0.8	0.7	0.8	0.8	0.8	0.7
2000.....	0.8	0.7	0.9	0.8	0.7	0.9	0.8	0.8	0.7
1999.....	0.9	0.8	0.9	0.9	0.8	1.0	0.9	1.1	0.8

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Mental and behavioral disorders due to psychoactive substance use (F10-F19)										
2007.....	2.6	4.1	1.3	2.6	4.0	1.3	3.6	6.0	1.6	
2006.....	3.8	5.8	1.9	3.7	5.7	1.9	5.0	8.2	2.5	
2005.....	3.7	5.7	1.8	3.6	5.6	1.8	4.8	7.9	2.3	
2004.....	3.6	5.6	1.7	3.5	5.4	1.7	5.0	8.6	2.1	
2003.....	3.4	5.4	1.6	3.3	5.1	1.6	4.9	8.3	2.2	
2002.....	3.3	5.2	1.5	3.1	4.9	1.5	5.2	8.9	2.3	
2001.....	3.2	5.0	1.5	3.0	4.6	1.4	5.4	9.4	2.3	
2000.....	3.1	5.0	1.4	2.9	4.6	1.3	5.4	9.3	2.3	
1999.....	3.1	5.1	1.4	2.9	4.6	1.2	5.7	9.7	2.6	
Mental and behavioral disorders due to use of alcohol (F10)										
2007.....	1.9	3.1	0.8	1.9	3.1	0.8	2.1	3.9	0.7	
2006.....	2.4	3.9	1.0	2.4	3.9	1.1	2.7	4.8	1.1	
2005.....	2.4	3.9	1.0	2.4	3.9	1.0	2.7	4.9	1.1	
2004.....	2.4	3.9	1.0	2.4	3.9	1.0	3.0	5.5	1.0	
2003.....	2.4	3.8	1.0	2.3	3.7	1.0	3.0	5.5	1.1	
2002.....	2.3	3.9	1.0	2.3	3.7	0.9	3.3	6.1	1.2	
2001.....	2.3	3.8	1.0	2.2	3.5	0.9	3.7	6.7	1.3	
2000.....	2.2	3.7	0.9	2.1	3.5	0.8	3.6	6.6	1.3	
1999.....	2.2	3.7	0.9	2.1	3.4	0.8	3.8	6.8	1.5	
Mental and behavioral disorders due to other psychoactive substance use (F11-F19)										
2007.....	0.8	1.0	0.5	0.7	1.0	0.5	1.5	2.1	0.9	
2006.....	1.4	1.9	0.9	1.3	1.8	0.8	2.3	3.4	1.4	
2005.....	1.3	1.8	0.8	1.2	1.7	0.8	2.0	3.0	1.3	
2004.....	1.2	1.6	0.7	1.1	1.5	0.7	2.0	3.1	1.1	
2003.....	1.1	1.5	0.6	1.0	1.4	0.6	1.9	2.8	1.2	
2002.....	0.9	1.3	0.6	0.9	1.2	0.5	1.9	2.8	1.1	
2001.....	0.9	1.2	0.5	0.8	1.1	0.5	1.8	2.6	1.1	
2000.....	0.9	1.2	0.6	0.8	1.1	0.5	1.8	2.7	1.1	
1999.....	0.9	1.3	0.5	0.8	1.1	0.4	1.9	2.9	1.1	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Schizophrenia, schizotypal and delusional disorders (F20-F29)									
2007.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2
2006.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.2
2005.....	0.1	0.2	0.2	0.1	0.1	0.2	0.2	0.3	0.2
2004.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.4	0.2
2003.....	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.2	0.2
2002.....	0.2	0.2	0.2	0.2	0.1	0.2	0.3	0.3	0.2
2001.....	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.2	0.2
2000.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Mood (affective) disorders (F30-F39)									
2007.....	0.2	0.2	0.2	0.2	0.2	0.2	0.1	*	*
2006.....	0.2	0.2	0.2	0.2	0.2	0.3	0.1	*	0.1
2005.....	0.2	0.2	0.3	0.3	0.2	0.3	0.1	*	*
2004.....	0.3	0.2	0.3	0.3	0.2	0.3	0.1	*	0.1
2003.....	0.3	0.3	0.3	0.3	0.3	0.4	0.2	*	0.2
2002.....	0.3	0.2	0.3	0.3	0.2	0.4	0.2	*	0.1
2001.....	0.3	0.3	0.3	0.3	0.3	0.3	0.1	0.2	0.1
2000.....	0.3	0.3	0.3	0.3	0.3	0.3	0.1	*	0.1
1999.....	0.3	0.3	0.3	0.3	0.3	0.3	0.1	*	*
Neurotic, stress-related and somatoform disorders (F40-F48)									
2007.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2006.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2005.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Mental retardation (F70-F79)										
2007.....	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.3	*	
2006.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.2	
2005.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.1	
2004.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.1	
2003.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.1	
2002.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	*	
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.1	
2000.....	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.2	0.1	
1999.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.2	0.2	
Other and unspecified mental and behavioral disorders (F50-F69,F80-F99)										
2007.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1	
2006.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1	
2005.....	0.1	0.1	0.2	0.1	0.1	0.1	0.2	0.2	0.2	
2004.....	0.2	0.1	0.2	0.2	0.1	0.2	0.1	*	0.1	
2003.....	0.2	0.1	0.2	0.2	0.1	0.2	0.2	*	0.2	
2002.....	0.2	0.1	0.2	0.2	0.1	0.2	0.1	*	0.1	
2001.....	0.1	0.1	0.2	0.1	0.1	0.2	0.1	*	*	
2000.....	0.1	0.1	0.2	0.1	0.1	0.2	0.1	*	*	
1999.....	0.1	0.1	0.2	0.1	0.1	0.2	0.1	*	*	
VI. Diseases of the nervous system (G00-G98)										
2007.....	39.3	39.9	38.4	40.6	41.1	39.6	34.6	34.6	33.9	
2006.....	38.9	39.6	37.9	40.2	41.0	39.1	33.1	32.3	32.8	
2005.....	39.0	39.3	38.2	40.2	40.4	39.3	34.4	34.5	33.7	
2004.....	37.1	37.5	36.3	38.2	38.7	37.3	32.7	32.1	32.5	
2003.....	37.2	37.8	36.1	38.3	38.9	37.3	31.1	32.2	30.0	
2002.....	35.6	36.5	34.4	36.7	37.6	35.6	29.3	30.5	27.9	
2001.....	34.2	35.8	32.7	35.2	36.8	33.7	28.6	30.2	27.3	
2000.....	33.0	34.8	31.5	34.0	35.8	32.4	27.0	28.8	25.7	
1999.....	30.9	33.2	29.1	31.8	34.2	29.9	25.0	26.6	23.7	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Meningitis (G00,G03)										
2007.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.3	
2006.....	0.2	0.2	0.2	0.2	0.2	0.2	0.4	0.4	0.3	
2005.....	0.2	0.3	0.2	0.2	0.2	0.2	0.4	0.5	0.4	
2004.....	0.3	0.3	0.2	0.2	0.2	0.2	0.5	0.5	0.4	
2003.....	0.3	0.3	0.2	0.2	0.3	0.2	0.6	0.5	0.6	
2002.....	0.2	0.3	0.2	0.2	0.3	0.2	0.5	0.7	0.4	
2001.....	0.3	0.3	0.2	0.2	0.3	0.2	0.5	0.8	0.4	
2000.....	0.3	0.3	0.2	0.2	0.3	0.2	0.6	0.8	0.6	
1999.....	0.3	0.3	0.3	0.3	0.3	0.2	0.7	0.9	0.6	
Other inflammatory diseases of central nervous system (G04-G09)										
2007.....	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.4	0.3	
2006.....	0.2	0.3	0.2	0.2	0.3	0.2	0.4	0.4	0.4	
2005.....	0.2	0.3	0.2	0.2	0.2	0.2	0.4	0.5	0.3	
2004.....	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.4	0.3	
2003.....	0.3	0.3	0.2	0.2	0.3	0.2	0.3	0.4	0.3	
2002.....	0.2	0.3	0.2	0.2	0.3	0.2	0.4	0.5	0.3	
2001.....	0.2	0.2	0.2	0.2	0.2	0.2	0.4	0.4	0.4	
2000.....	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.4	0.3	
1999.....	0.2	0.3	0.2	0.2	0.2	0.2	0.3	0.4	0.3	
Parkinson's disease (G20-G21)										
2007.....	6.4	9.6	4.3	6.8	10.2	4.6	3.2	4.9	2.2	
2006.....	6.3	9.6	4.3	6.7	10.2	4.6	2.7	4.3	1.9	
2005.....	6.4	9.8	4.4	6.8	10.4	4.6	3.0	4.9	1.9	
2004.....	6.1	9.3	4.1	6.4	9.8	4.4	2.7	4.3	1.9	
2003.....	6.2	9.3	4.2	6.5	9.8	4.5	2.7	4.4	1.8	
2002.....	5.9	9.0	4.1	6.3	9.5	4.3	2.5	4.0	1.7	
2001.....	5.9	9.1	4.0	6.2	9.7	4.3	2.5	3.9	1.8	
2000.....	5.7	8.8	3.9	6.0	9.3	4.2	2.5	4.2	1.6	
1999.....	5.4	8.4	3.7	5.7	8.9	3.9	2.4	3.7	1.7	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Alzheimer's disease (G30)									
2007.....	22.7	18.5	24.9	23.5	19.1	25.9	19.0	15.6	20.6
2006.....	22.6	18.4	24.7	23.4	19.1	25.7	18.3	14.7	19.8
2005.....	22.9	18.5	25.1	23.7	19.0	26.1	19.4	16.4	20.7
2004.....	21.8	17.7	23.8	22.5	18.3	24.7	18.5	15.2	19.9
2003.....	21.4	17.5	23.3	22.2	18.2	24.3	16.6	14.7	17.3
2002.....	20.2	16.6	22.0	21.1	17.3	22.9	15.2	13.2	15.9
2001.....	19.1	16.1	20.5	19.8	16.7	21.3	14.6	13.1	15.1
2000.....	18.1	15.2	19.3	18.8	15.8	20.1	13.0	11.4	13.7
1999.....	16.5	14.4	17.4	17.2	15.0	18.1	11.3	10.2	11.8
Multiple sclerosis (G35)									
2007.....	1.1	0.8	1.4	1.2	0.8	1.5	1.1	0.8	1.4
2006.....	1.1	0.8	1.3	1.1	0.9	1.4	1.1	0.8	1.4
2005.....	1.1	0.8	1.4	1.2	0.8	1.5	1.1	0.8	1.4
2004.....	1.1	0.8	1.4	1.1	0.8	1.4	1.0	0.7	1.3
2003.....	1.1	0.8	1.3	1.1	0.9	1.4	1.1	0.8	1.3
2002.....	1.1	0.8	1.3	1.1	0.8	1.4	1.1	0.6	1.5
2001.....	1.1	0.8	1.3	1.1	0.8	1.4	1.0	0.6	1.3
2000.....	1.0	0.8	1.2	1.1	0.8	1.3	1.0	0.7	1.3
1999.....	1.0	0.7	1.2	1.0	0.7	1.2	0.8	0.6	1.0
Epilepsy (G40-G41)									
2007.....	0.5	0.5	0.4	0.4	0.5	0.4	0.9	1.1	0.7
2006.....	0.5	0.5	0.4	0.4	0.5	0.4	0.9	1.0	0.7
2005.....	0.5	0.5	0.5	0.4	0.5	0.4	1.0	1.1	0.9
2004.....	0.5	0.5	0.4	0.4	0.5	0.3	0.9	1.1	0.7
2003.....	0.5	0.5	0.4	0.4	0.5	0.4	1.0	1.3	0.8
2002.....	0.5	0.6	0.4	0.4	0.5	0.4	1.0	1.3	0.8
2001.....	0.4	0.5	0.4	0.4	0.5	0.4	0.9	1.2	0.8
2000.....	0.5	0.5	0.4	0.4	0.5	0.4	1.0	1.3	0.9
1999.....	0.5	0.6	0.4	0.4	0.5	0.4	1.1	1.3	0.9

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/			Black 2/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Transient cerebral ischemic attacks and related syndromes (G45)									
2007.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2006.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2005.....	0.1	0.0	0.1	0.1	0.0	0.1	*	*	*
2004.....	0.1	0.0	0.1	0.1	0.0	0.1	*	*	*
2003.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
Infantile cerebral palsy (G80)									
2007.....	0.5	0.6	0.4	0.5	0.6	0.4	0.6	0.8	0.5
2006.....	0.5	0.5	0.4	0.5	0.5	0.4	0.7	0.8	0.6
2005.....	0.5	0.5	0.4	0.4	0.5	0.4	0.7	0.8	0.6
2004.....	0.5	0.5	0.4	0.5	0.5	0.4	0.7	0.8	0.6
2003.....	0.5	0.5	0.4	0.4	0.5	0.4	0.6	0.8	0.6
2002.....	0.4	0.5	0.4	0.4	0.5	0.3	0.6	0.7	0.5
2001.....	0.4	0.5	0.3	0.4	0.4	0.3	0.5	0.6	0.4
2000.....	0.4	0.4	0.3	0.4	0.4	0.3	0.6	0.6	0.5
1999.....	0.4	0.4	0.3	0.3	0.4	0.3	0.5	0.6	0.4
All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)									
2007.....	7.7	9.3	6.5	7.8	9.4	6.5	9.1	10.6	7.9
2006.....	7.5	9.1	6.2	7.5	9.2	6.2	8.6	10.0	7.6
2005.....	7.1	8.6	6.0	7.1	8.7	5.9	8.4	9.6	7.5
2004.....	6.8	8.1	5.7	6.8	8.2	5.7	8.0	9.1	7.3
2003.....	7.0	8.4	5.9	7.0	8.5	5.9	8.0	9.1	7.2
2002.....	6.9	8.3	5.8	6.9	8.3	5.8	7.9	9.4	6.8
2001.....	6.7	8.1	5.7	6.7	8.1	5.6	8.1	9.6	7.2
2000.....	6.8	8.3	5.7	6.8	8.3	5.6	7.9	9.5	6.8
1999.....	6.6	8.1	5.6	6.6	8.1	5.5	7.7	8.8	7.0

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
VII. Diseases of the eye and adnexa (H00-H57)									
2007.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2006.....	0.0	0.0	0.0	0.0	*	0.0	*	*	*
2005.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2004.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2003.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2002.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2001.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	*	0.0	0.0	*	0.0	*	*	*
VIII. Diseases of the ear and mastoid process (H60-H93)									
2007.....	0.0	0.0	0.0	0.0	*	0.0	*	*	*
2006.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2005.....	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
IX. Diseases of the circulatory system (I00-I99)									
2007.....	251.2	300.2	211.6	245.4	294.0	205.8	337.2	406.0	286.1
2006.....	262.5	313.3	221.6	256.3	306.7	215.5	351.0	422.8	298.2
2005.....	278.8	331.1	237.1	272.5	324.7	230.4	370.6	438.4	319.8
2004.....	288.0	341.7	245.3	281.2	335.1	238.0	384.6	454.0	333.6
2003.....	307.7	365.0	262.3	300.4	357.9	254.5	409.0	482.3	355.1
2002.....	319.0	378.8	272.1	312.0	372.2	264.3	418.9	490.6	366.4
2001.....	328.2	389.8	280.4	320.8	382.6	272.3	430.2	508.3	374.9
2000.....	341.3	409.0	290.0	334.0	402.3	281.7	441.6	520.7	386.1
1999.....	350.8	420.5	297.9	343.3	413.2	289.8	450.0	534.9	390.8

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)										
2007.....	1.0	0.8	1.2	1.0	0.8	1.2	0.8	0.7	0.9	
2006.....	1.1	0.8	1.2	1.1	0.8	1.3	0.8	0.7	0.9	
2005.....	1.1	0.8	1.3	1.1	0.8	1.3	0.8	0.7	0.8	
2004.....	1.1	0.8	1.3	1.1	0.8	1.3	0.8	0.6	1.0	
2003.....	1.2	1.0	1.4	1.3	1.0	1.4	1.1	0.9	1.2	
2002.....	1.2	0.9	1.5	1.3	0.9	1.5	0.9	0.8	1.0	
2001.....	1.2	0.9	1.5	1.2	0.9	1.5	1.1	0.7	1.3	
2000.....	1.3	1.0	1.5	1.3	1.0	1.6	1.0	0.7	1.2	
1999.....	1.3	0.9	1.6	1.4	0.9	1.6	1.0	0.8	1.2	
Acute rheumatic fever (I00-I02)										
2007.....	0.0	*	0.0	0.0	*	0.0	*	*	*	
2006.....	0.0	*	*	*	*	*	*	*	*	
2005.....	0.0	*	*	0.0	*	*	*	*	*	
2004.....	0.0	*	0.0	0.0	*	*	*	*	*	
2003.....	0.0	*	0.0	0.0	*	*	*	*	*	
2002.....	0.0	*	0.0	0.0	*	0.0	*	*	*	
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2000.....	0.0	*	0.0	0.0	*	*	*	*	*	
1999.....	0.0	*	*	0.0	*	*	*	*	*	
Chronic rheumatic heart diseases (I05-I09)										
2007.....	1.0	0.8	1.1	1.0	0.8	1.2	0.8	0.7	0.9	
2006.....	1.0	0.8	1.2	1.1	0.8	1.3	0.8	0.6	0.9	
2005.....	1.1	0.8	1.3	1.1	0.8	1.3	0.7	0.7	0.8	
2004.....	1.1	0.8	1.3	1.1	0.8	1.3	0.8	0.6	1.0	
2003.....	1.2	1.0	1.4	1.2	1.0	1.4	1.0	0.9	1.2	
2002.....	1.2	0.9	1.5	1.2	0.9	1.5	0.9	0.8	1.0	
2001.....	1.2	0.9	1.4	1.2	0.9	1.4	1.0	0.7	1.3	
2000.....	1.3	0.9	1.5	1.3	0.9	1.6	1.0	0.7	1.2	
1999.....	1.3	0.9	1.6	1.4	0.9	1.6	1.0	0.7	1.2	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)										
2007.....	0.3	0.2	0.3	0.3	0.2	0.3	0.3	0.2	0.3	
2006.....	0.3	0.2	0.4	0.3	0.2	0.4	0.2	0.2	0.3	
2005.....	0.3	0.2	0.4	0.3	0.2	0.4	0.3	0.2	0.3	
2004.....	0.3	0.2	0.4	0.3	0.2	0.4	0.2	*	0.3	
2003.....	0.3	0.2	0.4	0.3	0.2	0.4	0.3	0.2	0.4	
2002.....	0.4	0.2	0.4	0.4	0.3	0.4	0.3	0.2	0.3	
2001.....	0.4	0.3	0.5	0.4	0.3	0.5	0.3	0.2	0.4	
2000.....	0.4	0.3	0.5	0.4	0.3	0.5	0.4	0.2	0.5	
1999.....	0.4	0.3	0.5	0.5	0.3	0.5	0.3	0.2	0.3	
Hypertensive diseases (I10-I15)										
2007.....	17.8	18.6	16.5	15.3	15.7	14.3	42.5	49.1	36.9	
2006.....	17.8	18.6	16.5	15.2	15.6	14.3	43.6	51.1	37.7	
2005.....	18.5	18.9	17.4	15.8	15.8	15.0	45.7	52.2	40.3	
2004.....	18.1	18.7	17.0	15.4	15.7	14.5	45.7	51.0	40.9	
2003.....	18.1	18.3	17.2	15.3	15.3	14.6	46.0	51.1	41.7	
2002.....	17.1	17.4	16.3	14.4	14.4	13.8	44.8	49.6	40.6	
2001.....	16.5	16.6	15.8	13.9	13.7	13.4	43.0	47.8	38.9	
2000.....	16.2	16.3	15.5	13.5	13.4	12.9	43.3	47.2	39.9	
1999.....	15.8	15.9	15.2	13.1	12.9	12.6	43.1	47.5	39.4	
Essential (primary) hypertension (I10)										
2007.....	4.4	4.0	4.5	4.1	3.7	4.2	7.9	8.1	7.6	
2006.....	4.4	4.0	4.5	4.0	3.6	4.1	8.5	8.7	8.1	
2005.....	4.8	4.1	5.0	4.4	3.7	4.7	8.6	8.9	8.2	
2004.....	4.4	4.0	4.6	4.1	3.6	4.2	8.2	7.9	8.1	
2003.....	4.4	3.7	4.6	4.0	3.3	4.3	8.4	8.6	8.1	
2002.....	4.0	3.5	4.2	3.7	3.1	3.9	7.7	7.5	7.5	
2001.....	3.9	3.5	4.0	3.6	3.1	3.7	7.6	8.2	7.1	
2000.....	3.7	3.4	3.8	3.4	3.0	3.5	7.3	7.6	6.9	
1999.....	3.6	3.3	3.7	3.3	2.9	3.4	7.2	7.4	7.0	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Hypertensive heart disease (I11)									
2007.....	9.5	10.3	8.4	8.1	8.6	7.2	23.3	28.4	19.3
2006.....	9.4	10.1	8.3	8.0	8.4	7.2	23.1	28.5	18.8
2005.....	9.4	10.1	8.5	7.9	8.3	7.2	24.1	28.4	20.4
2004.....	9.5	10.1	8.5	8.0	8.4	7.2	24.2	28.4	20.6
2003.....	9.6	10.1	8.8	8.0	8.3	7.3	25.0	28.7	21.9
2002.....	9.1	9.5	8.3	7.6	7.8	7.0	24.4	28.0	21.2
2001.....	8.7	8.9	8.1	7.3	7.3	6.8	23.2	26.3	20.5
2000.....	8.6	8.7	8.1	7.1	7.1	6.7	23.7	26.3	21.2
1999.....	8.3	8.3	7.9	6.9	6.7	6.6	22.9	25.7	20.5
Hypertensive renal disease and secondary hypertension (I12,I15)									
2007.....	3.0	3.3	2.8	2.4	2.7	2.2	8.4	9.3	7.7
2006.....	3.1	3.5	2.9	2.5	2.8	2.3	9.2	10.5	8.3
2005.....	3.3	3.6	3.0	2.6	2.9	2.4	9.8	11.0	8.9
2004.....	3.2	3.5	3.0	2.5	2.8	2.3	10.0	11.0	9.2
2003.....	3.1	3.4	2.8	2.4	2.7	2.2	9.2	10.0	8.5
2002.....	3.0	3.3	2.8	2.4	2.7	2.2	9.3	10.2	8.6
2001.....	2.9	3.2	2.7	2.3	2.5	2.1	8.9	9.7	8.3
2000.....	2.8	3.1	2.6	2.2	2.5	2.0	9.0	9.5	8.7
1999.....	2.6	2.9	2.4	2.0	2.3	1.8	8.8	9.8	8.0
Hypertensive heart and renal disease (I13)									
2007.....	0.9	1.0	0.9	0.7	0.8	0.7	2.8	3.4	2.4
2006.....	0.9	1.0	0.8	0.7	0.8	0.7	2.9	3.5	2.5
2005.....	1.0	1.1	0.9	0.8	0.9	0.8	3.2	3.8	2.8
2004.....	1.0	1.1	0.9	0.8	0.8	0.7	3.2	3.6	2.9
2003.....	1.1	1.1	1.0	0.8	0.9	0.7	3.4	3.7	3.1
2002.....	1.0	1.0	0.9	0.7	0.8	0.7	3.5	3.9	3.2
2001.....	1.0	1.1	1.0	0.8	0.8	0.7	3.3	3.6	3.0
2000.....	1.0	1.1	0.9	0.8	0.9	0.7	3.3	3.8	3.1
1999.....	1.2	1.4	1.1	0.9	1.0	0.9	4.2	4.5	3.9

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Ischemic heart diseases (I20-I25)									
2007.....	126.0	165.4	95.7	125.5	165.6	94.2	150.6	191.6	121.5
2006.....	134.9	176.5	103.1	134.2	176.3	101.5	161.6	206.4	130.0
2005.....	144.4	187.4	111.7	143.8	187.7	110.0	171.3	213.9	140.9
2004.....	150.2	194.1	116.7	149.2	194.2	114.7	179.8	223.9	148.7
2003.....	162.9	209.9	127.2	161.7	209.6	125.0	195.0	243.3	160.9
2002.....	170.8	220.4	133.6	169.8	220.5	131.2	203.0	250.6	169.7
2001.....	177.8	228.5	139.9	176.5	228.4	137.4	211.6	262.0	176.7
2000.....	186.8	241.4	146.5	185.6	241.6	143.9	218.3	269.5	183.2
1999.....	194.6	251.2	152.9	193.4	251.2	150.5	224.9	278.7	188.0
Acute myocardial infarction (I21-I22)									
2007.....	41.4	54.0	31.5	41.3	54.3	30.9	49.5	60.9	41.2
2006.....	45.0	58.8	34.2	45.0	59.2	33.7	53.3	66.3	44.0
2005.....	49.1	63.6	37.8	49.1	64.1	37.2	57.6	69.7	48.8
2004.....	52.2	67.8	40.2	52.2	68.3	39.5	62.0	74.8	52.7
2003.....	57.9	75.0	44.8	57.8	75.3	44.0	68.6	84.3	57.6
2002.....	62.1	80.6	48.1	61.9	81.0	47.1	73.9	89.5	63.2
2001.....	65.4	84.9	50.9	65.1	85.4	49.9	76.8	93.1	66.0
2000.....	69.9	91.3	54.1	69.6	91.8	53.1	81.0	98.9	69.2
1999.....	73.2	96.1	56.3	73.1	96.8	55.4	83.1	102.1	70.5
Other acute ischemic heart diseases (I24)									
2007.....	1.3	1.6	1.0	1.2	1.5	1.0	1.8	2.4	1.4
2006.....	1.2	1.5	1.0	1.2	1.5	1.0	1.8	2.4	1.4
2005.....	1.2	1.4	0.9	1.1	1.3	0.9	2.0	2.6	1.5
2004.....	0.8	1.0	0.6	0.8	1.0	0.6	1.4	2.1	1.0
2003.....	1.1	1.3	0.9	1.0	1.2	0.8	1.6	2.4	1.1
2002.....	1.2	1.5	0.9	1.1	1.4	0.9	1.7	2.3	1.3
2001.....	1.2	1.5	0.9	1.1	1.4	0.9	1.8	2.3	1.3
2000.....	1.2	1.5	0.9	1.2	1.5	0.9	1.8	2.5	1.3
1999.....	1.2	1.5	0.9	1.1	1.4	0.9	1.8	2.4	1.3

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/		White 2/		Black 2/				
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other forms of chronic ischemic heart disease (I20,I25)									
2007.....	83.3	109.8	63.2	82.9	109.8	62.3	99.3	128.2	78.9
2006.....	88.7	116.1	67.9	88.0	115.7	66.9	106.5	137.7	84.6
2005.....	94.2	122.4	73.0	93.6	122.3	72.0	111.8	141.6	90.5
2004.....	97.1	125.3	75.8	96.3	125.0	74.6	116.4	146.9	95.0
2003.....	103.8	133.5	81.5	102.9	133.0	80.1	124.8	156.7	102.2
2002.....	107.6	138.3	84.5	106.8	138.0	83.2	127.5	158.9	105.2
2001.....	111.2	142.1	88.1	110.2	141.6	86.7	133.0	166.7	109.3
2000.....	115.7	148.6	91.5	114.8	148.3	90.0	135.4	168.1	112.6
1999.....	120.2	153.6	95.6	119.2	153.0	94.2	140.0	174.2	116.2
Atherosclerotic cardiovascular disease, so described (I25.0)									
2007.....	18.2	24.3	13.1	17.3	23.0	12.3	30.3	42.3	21.6
2006.....	19.3	25.6	14.0	18.1	24.0	13.1	32.7	46.0	23.3
2005.....	20.3	26.6	15.0	19.1	25.0	14.0	34.5	46.9	25.3
2004.....	21.1	27.5	15.7	19.9	25.9	14.7	36.1	48.9	27.0
2003.....	22.7	29.6	17.1	21.4	27.8	16.1	38.2	52.7	28.0
2002.....	23.5	30.4	17.8	22.1	28.5	16.7	40.3	54.8	30.0
2001.....	24.1	30.8	18.5	22.5	28.8	17.3	41.8	56.0	31.7
2000.....	25.2	32.3	19.4	23.6	30.2	18.2	43.5	58.7	32.9
1999.....	26.4	33.5	20.6	24.7	31.2	19.3	45.8	61.5	34.9
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)									
2007.....	65.1	85.5	50.1	65.6	86.7	49.9	69.1	85.8	57.4
2006.....	69.4	90.6	53.9	69.9	91.7	53.8	73.8	91.7	61.4
2005.....	73.9	95.8	58.0	74.5	97.3	57.9	77.3	94.7	65.2
2004.....	76.0	97.8	60.1	76.5	99.1	59.9	80.3	98.0	68.0
2003.....	81.1	104.0	64.4	81.5	105.2	64.0	86.6	104.0	74.2
2002.....	84.1	108.0	66.7	84.7	109.5	66.4	87.2	104.1	75.2
2001.....	87.1	111.3	69.6	87.6	112.8	69.4	91.1	110.6	77.6
2000.....	90.5	116.3	72.0	91.2	118.1	71.8	91.9	109.5	79.7
1999.....	93.8	120.1	75.0	94.5	121.8	74.9	94.1	112.7	81.3

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Other heart diseases (I26-I51)									
2007.....	53.4	60.2	48.0	52.6	59.0	47.3	69.7	81.9	60.4
2006.....	53.9	60.0	48.7	53.0	58.8	48.0	69.3	81.6	60.3
2005.....	55.2	61.5	49.9	54.1	60.2	49.0	71.9	83.0	63.4
2004.....	55.2	61.7	49.9	54.1	60.3	49.0	72.5	85.6	63.2
2003.....	57.6	64.5	52.0	56.4	63.1	50.9	75.7	87.7	66.7
2002.....	58.5	65.5	52.8	57.3	64.1	51.7	76.6	87.8	68.1
2001.....	59.1	66.1	53.4	57.7	64.4	52.3	77.8	91.8	68.2
2000.....	60.0	67.9	53.8	58.6	66.3	52.6	78.4	92.2	69.0
1999.....	60.9	69.2	54.5	59.3	67.2	53.1	81.2	97.5	70.1
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)									
2007.....	4.0	3.6	4.2	3.7	3.4	4.0	6.8	6.4	7.1
2006.....	4.0	3.6	4.3	3.8	3.4	4.0	6.7	6.2	7.1
2005.....	4.4	4.1	4.7	4.2	3.9	4.4	7.5	7.1	7.8
2004.....	4.4	4.0	4.7	4.1	3.8	4.4	7.6	7.3	7.8
2003.....	4.8	4.6	5.0	4.5	4.3	4.7	8.2	8.1	8.4
2002.....	4.8	4.4	5.0	4.5	4.2	4.7	8.1	7.2	8.8
2001.....	4.7	4.5	4.9	4.5	4.2	4.6	8.0	7.9	8.3
2000.....	4.7	4.5	4.9	4.5	4.3	4.6	8.0	7.6	8.3
1999.....	4.8	4.7	5.0	4.5	4.5	4.6	8.7	8.4	8.9
Pulmonary embolism (I26)									
2007.....	2.3	2.2	2.3	2.1	2.1	2.2	3.9	4.2	3.7
2006.....	2.2	2.2	2.2	2.1	2.1	2.1	3.9	4.1	3.8
2005.....	2.7	2.7	2.7	2.6	2.6	2.6	4.8	5.0	4.6
2004.....	2.7	2.7	2.7	2.6	2.5	2.6	4.9	5.2	4.7
2003.....	3.0	3.0	2.9	2.8	2.8	2.7	5.4	5.8	5.2
2002.....	3.0	2.9	3.1	2.8	2.7	2.9	5.2	5.0	5.4
2001.....	3.1	3.0	3.1	2.9	2.8	2.9	5.3	5.7	5.2
2000.....	3.0	3.0	3.0	2.9	2.9	2.8	5.4	5.7	5.4
1999.....	3.3	3.3	3.3	3.1	3.1	3.1	6.1	6.5	5.9

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)										
2007.....	1.7	1.4	1.9	1.6	1.3	1.8	2.9	2.2	3.4	
2006.....	1.8	1.4	2.0	1.7	1.4	1.9	2.8	2.1	3.3	
2005.....	1.7	1.4	1.9	1.6	1.4	1.8	2.7	2.1	3.2	
2004.....	1.7	1.4	1.9	1.6	1.3	1.8	2.7	2.1	3.1	
2003.....	1.8	1.6	2.0	1.7	1.5	1.9	2.8	2.2	3.2	
2002.....	1.8	1.5	1.9	1.7	1.5	1.8	2.9	2.2	3.3	
2001.....	1.7	1.5	1.8	1.6	1.4	1.7	2.7	2.3	3.0	
2000.....	1.7	1.5	1.8	1.6	1.4	1.7	2.5	2.0	2.9	
1999.....	1.5	1.4	1.6	1.5	1.4	1.5	2.6	1.9	3.0	
Other non-pulmonary forms of heart disease (I30-I51)										
2007.....	49.5	56.6	43.7	48.8	55.5	43.3	62.9	75.5	53.4	
2006.....	49.9	56.4	44.4	49.2	55.4	44.0	62.6	75.3	53.1	
2005.....	50.7	57.3	45.2	49.9	56.3	44.6	64.4	75.9	55.6	
2004.....	50.8	57.7	45.3	50.0	56.5	44.6	65.0	78.3	55.4	
2003.....	52.8	60.0	47.0	51.9	58.8	46.2	67.4	79.7	58.3	
2002.....	53.8	61.1	47.8	52.8	59.9	47.0	68.5	80.6	59.4	
2001.....	54.4	61.6	48.5	53.3	60.2	47.7	69.8	83.9	60.0	
2000.....	55.2	63.4	48.9	54.2	62.0	48.0	70.5	84.5	60.7	
1999.....	56.1	64.5	49.6	54.9	62.7	48.6	72.5	89.1	61.2	
Acute and subacute endocarditis (I33)										
2007.....	0.4	0.5	0.3	0.4	0.5	0.3	0.7	1.0	0.4	
2006.....	0.4	0.5	0.3	0.4	0.4	0.3	0.7	0.8	0.6	
2005.....	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.9	0.6	
2004.....	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.8	0.6	
2003.....	0.4	0.5	0.3	0.4	0.5	0.3	0.8	0.9	0.6	
2002.....	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.8	0.6	
2001.....	0.4	0.5	0.3	0.4	0.5	0.3	0.8	1.1	0.6	
2000.....	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.9	0.7	
1999.....	0.4	0.5	0.3	0.4	0.5	0.3	0.8	0.9	0.6	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Diseases of pericardium and acute myocarditis (I30-I31,I40)										
2007.....	0.3	0.3	0.2	0.3	0.3	0.2	0.4	0.4	0.4	
2006.....	0.3	0.3	0.2	0.3	0.3	0.2	0.3	0.3	0.3	
2005.....	0.3	0.3	0.3	0.3	0.3	0.2	0.5	0.5	0.5	
2004.....	0.3	0.3	0.2	0.2	0.3	0.2	0.5	0.6	0.4	
2003.....	0.3	0.3	0.3	0.3	0.3	0.2	0.5	0.5	0.5	
2002.....	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.5	0.4	
2001.....	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.5	0.4	
2000.....	0.3	0.3	0.2	0.3	0.3	0.2	0.4	0.5	0.4	
1999.....	0.3	0.4	0.3	0.3	0.3	0.3	0.5	0.5	0.5	
Other diseases of endocardium (I34-I38)										
2007.....	7.1	7.7	6.7	7.5	8.0	7.1	4.9	5.4	4.5	
2006.....	6.7	7.0	6.5	7.0	7.3	6.8	4.4	4.6	4.2	
2005.....	6.7	6.8	6.6	7.0	7.1	6.8	4.5	4.5	4.4	
2004.....	6.7	6.9	6.5	7.0	7.1	6.7	4.9	5.1	4.6	
2003.....	6.8	7.0	6.6	7.0	7.3	6.8	4.9	5.0	4.7	
2002.....	6.9	7.1	6.7	7.2	7.4	7.0	4.8	4.6	4.8	
2001.....	6.9	7.2	6.6	7.2	7.5	6.9	4.9	5.3	4.6	
2000.....	7.2	7.5	6.9	7.4	7.7	7.1	5.3	5.4	5.1	
1999.....	7.2	7.6	6.9	7.4	7.8	7.1	5.3	5.4	5.1	
Nonrheumatic mitral valve disorders (I34)										
2007.....	0.8	0.8	0.9	0.8	0.8	0.9	0.6	0.5	0.7	
2006.....	0.8	0.7	0.9	0.9	0.8	0.9	0.5	0.4	0.5	
2005.....	0.8	0.7	0.9	0.9	0.8	0.9	0.7	0.4	0.8	
2004.....	0.8	0.8	0.9	0.9	0.8	0.9	0.7	0.6	0.8	
2003.....	0.9	0.8	0.9	0.9	0.8	0.9	0.6	0.6	0.7	
2002.....	1.0	0.8	1.0	1.0	0.9	1.1	0.6	0.6	0.7	
2001.....	1.0	0.9	1.1	1.0	0.9	1.1	0.7	0.5	0.8	
2000.....	1.0	1.0	1.1	1.1	1.0	1.1	0.8	0.6	0.9	
1999.....	1.1	1.0	1.1	1.1	1.0	1.2	0.8	0.6	0.9	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Nonrheumatic aortic valve disorders (I35)										
2007.....	4.6	5.2	4.2	4.9	5.5	4.5	2.6	2.8	2.3	
2006.....	4.3	4.6	4.0	4.5	4.9	4.3	2.2	2.5	2.1	
2005.....	4.2	4.4	4.0	4.5	4.7	4.2	2.2	2.4	2.0	
2004.....	4.2	4.4	4.0	4.4	4.6	4.2	2.4	2.6	2.2	
2003.....	4.2	4.6	4.0	4.5	4.8	4.2	2.4	2.4	2.3	
2002.....	4.3	4.6	4.0	4.5	4.9	4.2	2.4	2.4	2.4	
2001.....	4.3	4.7	4.0	4.5	4.9	4.2	2.5	3.0	2.1	
2000.....	4.5	4.9	4.2	4.7	5.1	4.4	2.7	2.9	2.5	
1999.....	4.5	5.0	4.2	4.7	5.2	4.3	2.7	3.1	2.5	
All other diseases of endocardium (I36-I38)										
2007.....	1.7	1.7	1.6	1.7	1.7	1.7	1.7	2.0	1.5	
2006.....	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.7	1.5	
2005.....	1.7	1.6	1.6	1.7	1.6	1.7	1.6	1.7	1.6	
2004.....	1.7	1.7	1.6	1.7	1.7	1.6	1.8	1.9	1.7	
2003.....	1.7	1.6	1.7	1.7	1.6	1.7	1.9	2.0	1.8	
2002.....	1.6	1.6	1.7	1.6	1.6	1.7	1.7	1.6	1.7	
2001.....	1.6	1.7	1.6	1.6	1.7	1.6	1.7	1.8	1.6	
2000.....	1.6	1.6	1.6	1.6	1.6	1.6	1.8	2.0	1.7	
1999.....	1.6	1.6	1.6	1.6	1.6	1.6	1.8	1.8	1.8	
Cardiomyopathy (I42)										
2007.....	7.6	10.7	5.2	7.0	10.0	4.8	13.5	19.0	9.4	
2006.....	7.9	11.0	5.6	7.3	10.2	5.2	13.9	19.3	9.9	
2005.....	8.3	11.7	5.8	7.7	10.9	5.3	14.7	20.6	10.3	
2004.....	8.5	11.9	6.1	7.9	11.1	5.6	14.9	20.6	10.7	
2003.....	9.1	12.6	6.5	8.5	11.8	6.0	15.9	22.1	11.5	
2002.....	9.6	13.4	6.8	8.9	12.5	6.2	16.6	23.0	12.0	
2001.....	9.5	13.1	6.8	8.7	12.2	6.2	16.9	23.3	12.5	
2000.....	9.8	13.8	7.0	9.0	12.7	6.4	17.8	24.9	12.8	
1999.....	10.0	13.9	7.1	9.1	12.7	6.5	18.2	25.9	12.9	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Conduction disorders and cardiac dysrhythmias (I44-I49)										
2007.....	12.9	14.1	11.8	12.6	13.6	11.6	18.1	21.2	15.9	
2006.....	12.0	13.1	10.9	11.6	12.6	10.6	17.1	20.3	14.8	
2005.....	12.5	13.7	11.3	12.1	13.3	11.0	17.1	20.0	15.1	
2004.....	12.5	13.8	11.3	12.1	13.3	10.9	17.6	21.1	15.3	
2003.....	13.2	14.6	12.0	12.9	14.1	11.6	18.4	21.3	16.1	
2002.....	13.6	15.1	12.2	13.2	14.6	11.9	18.8	21.7	16.7	
2001.....	13.6	15.1	12.3	13.2	14.5	12.0	19.1	22.8	16.6	
2000.....	13.9	15.6	12.4	13.6	15.2	12.1	18.8	21.8	16.6	
1999.....	14.7	16.7	13.0	14.3	16.2	12.8	20.0	24.6	17.1	
Heart failure (I50)										
2007.....	17.3	18.6	16.2	17.3	18.7	16.3	20.1	22.3	18.4	
2006.....	18.9	20.0	17.9	19.0	20.2	18.1	20.9	23.0	19.3	
2005.....	18.9	19.9	18.0	18.9	20.0	18.1	21.7	23.0	20.4	
2004.....	18.9	19.9	18.0	19.0	20.0	18.1	20.9	23.4	19.3	
2003.....	19.4	20.6	18.4	19.5	20.7	18.4	21.8	23.6	20.5	
2002.....	19.5	20.4	18.6	19.5	20.5	18.6	21.8	23.3	20.5	
2001.....	20.2	21.2	19.2	20.2	21.2	19.3	22.1	24.4	20.6	
2000.....	20.3	21.5	19.2	20.3	21.6	19.3	22.0	24.2	20.7	
1999.....	20.3	21.4	19.3	20.3	21.4	19.3	22.5	25.3	20.9	
Congestive heart failure (I50.0)										
2007.....	15.9	17.1	14.9	16.0	17.2	15.1	17.7	19.8	16.2	
2006.....	17.5	18.4	16.7	17.7	18.6	16.9	18.6	20.1	17.5	
2005.....	17.6	18.5	16.8	17.7	18.6	17.0	19.2	20.2	18.2	
2004.....	17.6	18.5	16.8	17.7	18.6	17.0	18.8	20.8	17.4	
2003.....	18.1	19.1	17.3	18.2	19.3	17.4	19.5	20.9	18.5	
2002.....	18.0	18.8	17.3	18.1	19.0	17.4	19.6	20.7	18.5	
2001.....	18.7	19.5	18.0	18.8	19.6	18.1	19.9	21.7	18.8	
2000.....	18.8	19.8	17.9	18.9	20.0	18.0	19.7	21.3	18.8	
1999.....	18.8	19.8	17.9	18.8	19.7	18.0	20.3	22.7	18.9	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Other and unspecified heart failure (I50.1-I50.9)										
2007.....	1.4	1.6	1.3	1.3	1.5	1.2	2.4	2.5	2.2	
2006.....	1.4	1.6	1.2	1.3	1.5	1.2	2.2	3.0	1.8	
2005.....	1.3	1.5	1.2	1.2	1.4	1.1	2.5	2.8	2.2	
2004.....	1.3	1.5	1.1	1.2	1.4	1.1	2.2	2.6	1.9	
2003.....	1.3	1.5	1.1	1.2	1.4	1.1	2.3	2.7	2.0	
2002.....	1.4	1.6	1.3	1.3	1.5	1.2	2.2	2.6	2.0	
2001.....	1.5	1.7	1.3	1.4	1.6	1.2	2.2	2.7	1.8	
2000.....	1.5	1.7	1.4	1.4	1.6	1.3	2.3	2.8	1.9	
1999.....	1.5	1.7	1.3	1.4	1.6	1.3	2.2	2.5	2.0	
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)										
2007.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.3	
2006.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3	
2005.....	0.2	0.3	0.2	0.2	0.2	0.2	0.4	0.5	0.4	
2004.....	0.2	0.2	0.2	0.2	0.2	0.2	0.4	0.5	0.4	
2003.....	0.2	0.2	0.2	0.2	0.2	0.2	0.4	0.4	0.3	
2002.....	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.6	0.4	
2001.....	0.3	0.3	0.2	0.2	0.2	0.2	0.5	0.6	0.4	
2000.....	0.3	0.3	0.2	0.2	0.3	0.2	0.4	0.4	0.3	
1999.....	0.3	0.3	0.2	0.2	0.3	0.2	0.4	0.5	0.3	
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)										
2007.....	3.6	4.4	2.9	3.5	4.3	2.9	4.9	5.8	4.1	
2006.....	3.5	4.3	2.8	3.4	4.2	2.7	5.0	6.6	3.9	
2005.....	3.4	4.2	2.7	3.3	4.0	2.7	4.8	5.9	3.9	
2004.....	3.3	4.0	2.7	3.2	3.9	2.6	5.1	6.4	4.1	
2003.....	3.3	4.0	2.7	3.2	3.9	2.6	4.9	6.0	4.0	
2002.....	3.2	4.0	2.6	3.1	3.8	2.5	4.9	6.1	3.9	
2001.....	3.2	3.8	2.6	3.0	3.7	2.5	5.0	6.0	4.2	
2000.....	3.2	3.9	2.6	3.0	3.7	2.4	5.0	6.5	4.0	
1999.....	3.0	3.7	2.4	2.8	3.5	2.2	4.7	6.0	3.8	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Cerebrovascular diseases (I60-I69)									
2007.....	42.2	42.5	41.3	40.5	40.2	39.9	60.3	67.1	55.0
2006.....	43.6	43.9	42.6	41.7	41.7	41.1	61.6	67.1	57.0
2005.....	46.6	46.9	45.6	44.7	44.7	44.0	65.2	70.5	60.7
2004.....	50.0	50.4	48.9	48.0	48.1	47.2	69.9	74.9	65.5
2003.....	53.5	54.1	52.3	51.4	51.7	50.5	74.3	79.5	69.8
2002.....	56.2	56.5	55.2	54.2	54.2	53.4	76.3	81.7	71.8
2001.....	57.9	59.0	56.4	55.8	56.5	54.5	78.8	85.4	73.7
2000.....	60.9	62.4	59.1	58.8	59.8	57.3	81.9	89.6	76.2
1999.....	61.6	63.2	59.8	59.6	60.8	58.0	81.8	89.6	76.2
Subarachnoid hemorrhage (I60)									
2007.....	1.8	1.5	2.1	1.7	1.4	2.0	2.5	2.0	2.8
2006.....	1.9	1.5	2.2	1.8	1.4	2.1	2.6	2.0	3.1
2005.....	2.0	1.5	2.3	1.9	1.5	2.2	2.7	2.1	3.2
2004.....	2.0	1.6	2.4	1.9	1.5	2.3	2.7	2.0	3.2
2003.....	2.0	1.6	2.4	1.9	1.5	2.2	2.9	2.3	3.4
2002.....	2.1	1.6	2.6	2.0	1.5	2.4	2.9	2.1	3.5
2001.....	2.2	1.7	2.5	2.0	1.6	2.4	3.1	2.3	3.7
2000.....	2.4	1.8	2.8	2.2	1.7	2.6	3.4	2.3	4.2
1999.....	2.4	1.8	2.8	2.2	1.7	2.6	3.3	2.4	4.0
Intracerebral and other intracranial hemorrhage (I61-I62)									
2007.....	8.7	9.6	7.9	8.1	8.9	7.5	12.6	14.6	10.9
2006.....	8.7	9.8	7.9	8.2	9.0	7.4	12.7	14.9	11.0
2005.....	9.1	10.3	8.2	8.5	9.6	7.7	13.2	15.5	11.3
2004.....	9.2	10.3	8.4	8.6	9.6	7.8	13.5	15.8	11.8
2003.....	9.3	10.4	8.4	8.6	9.6	7.8	14.1	16.7	12.2
2002.....	9.4	10.4	8.5	8.8	9.7	8.0	13.2	15.3	11.6
2001.....	9.4	10.5	8.6	8.8	9.8	8.0	13.7	15.8	12.1
2000.....	9.9	11.1	9.0	9.2	10.3	8.5	14.3	17.0	12.3
1999.....	9.3	10.3	8.5	8.7	9.6	8.0	13.3	15.5	11.7

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Cerebral infarction (I63)										
2007.....	2.1	2.1	2.0	2.0	2.1	2.0	2.8	3.0	2.5	
2006.....	2.2	2.4	2.1	2.2	2.3	2.1	2.9	3.4	2.6	
2005.....	2.4	2.5	2.4	2.4	2.4	2.3	3.3	3.6	3.1	
2004.....	3.6	3.8	3.5	3.5	3.6	3.4	5.1	5.8	4.5	
2003.....	4.1	4.2	3.9	4.0	4.1	3.9	5.4	5.8	5.0	
2002.....	4.3	4.4	4.1	4.1	4.3	4.0	5.5	6.1	5.0	
2001.....	4.5	4.6	4.3	4.4	4.5	4.2	5.8	6.8	5.2	
2000.....	5.0	5.3	4.7	4.9	5.1	4.6	6.6	7.5	5.9	
1999.....	5.4	5.7	5.2	5.3	5.5	5.1	7.1	8.3	6.3	
Stroke, not specified as hemorrhage or infarction (I64)										
2007.....	22.0	21.5	21.9	21.2	20.4	21.3	31.8	35.7	28.8	
2006.....	22.9	22.4	22.8	22.1	21.4	22.2	32.4	34.9	30.2	
2005.....	24.1	23.7	23.8	23.3	22.7	23.2	33.5	36.1	31.2	
2004.....	25.4	24.9	25.3	24.6	23.9	24.5	35.8	37.0	34.3	
2003.....	28.0	27.7	27.6	27.1	26.6	26.9	38.7	40.6	36.8	
2002.....	30.0	29.6	29.8	29.1	28.4	29.0	41.2	44.1	38.8	
2001.....	31.4	31.4	30.9	30.5	30.2	30.2	42.7	46.4	39.8	
2000.....	32.8	33.0	32.2	31.9	31.8	31.4	44.2	48.5	41.1	
1999.....	33.6	34.0	32.8	32.7	32.8	32.0	44.6	48.4	41.8	
Other cerebrovascular diseases and their sequelae (I67,I69)										
2007.....	7.7	7.8	7.5	7.4	7.5	7.3	10.8	11.8	9.9	
2006.....	7.7	7.9	7.6	7.5	7.5	7.3	10.9	12.0	10.2	
2005.....	8.9	8.9	8.9	8.7	8.5	8.6	12.4	13.1	11.8	
2004.....	9.6	9.7	9.4	9.3	9.4	9.2	12.8	14.2	11.7	
2003.....	10.1	10.2	9.9	9.9	9.9	9.7	13.3	14.2	12.5	
2002.....	10.4	10.6	10.2	10.2	10.3	10.0	13.4	14.0	12.9	
2001.....	10.4	10.7	10.1	10.2	10.5	9.8	13.5	14.1	13.0	
2000.....	10.8	11.3	10.3	10.5	11.0	10.1	13.4	14.4	12.7	
1999.....	10.9	11.4	10.5	10.7	11.1	10.3	13.6	15.0	12.5	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Diseases of arteries, arterioles and capillaries (I70-I78)										
2007.....	9.4	11.4	7.9	9.4	11.4	7.8	11.0	13.2	9.4	
2006.....	9.9	12.2	8.2	9.9	12.2	8.2	11.7	13.6	10.2	
2005.....	11.5	13.9	9.7	11.5	14.0	9.7	13.0	15.0	11.3	
2004.....	11.9	14.3	10.1	11.9	14.4	10.0	13.2	15.1	11.7	
2003.....	12.8	15.7	10.7	12.9	15.8	10.6	14.0	16.6	12.1	
2002.....	13.4	16.4	11.2	13.5	16.6	11.2	14.4	17.0	12.6	
2001.....	14.0	17.1	11.7	14.0	17.2	11.7	15.1	17.4	13.4	
2000.....	14.7	18.4	12.0	14.7	18.6	11.9	15.7	18.4	13.9	
1999.....	15.0	18.5	12.5	15.1	18.7	12.5	15.5	18.2	13.7	
Atherosclerosis (I70)										
2007.....	2.5	2.6	2.4	2.6	2.7	2.4	2.4	2.6	2.3	
2006.....	2.7	2.8	2.6	2.8	2.8	2.6	2.7	3.3	2.4	
2005.....	3.8	3.9	3.7	3.9	4.0	3.7	3.4	3.4	3.3	
2004.....	3.9	4.1	3.7	4.0	4.2	3.8	3.5	3.5	3.3	
2003.....	4.4	4.5	4.3	4.5	4.5	4.4	4.2	4.7	3.8	
2002.....	4.7	4.9	4.6	4.8	5.0	4.7	4.4	4.9	4.0	
2001.....	5.0	5.1	4.8	5.1	5.2	5.0	4.3	4.6	4.1	
2000.....	5.2	5.5	5.0	5.3	5.6	5.1	4.8	5.4	4.4	
1999.....	5.5	5.7	5.3	5.6	5.8	5.4	5.0	5.8	4.5	
Aortic aneurysm and dissection (I71)										
2007.....	4.1	5.7	2.9	4.1	5.8	2.9	4.2	5.5	3.2	
2006.....	4.3	6.0	3.0	4.3	6.1	3.0	4.4	5.5	3.5	
2005.....	4.6	6.5	3.1	4.6	6.6	3.1	4.6	6.0	3.6	
2004.....	4.6	6.7	3.1	4.7	6.8	3.2	4.5	6.1	3.3	
2003.....	5.1	7.5	3.3	5.1	7.7	3.3	4.6	6.1	3.5	
2002.....	5.2	7.7	3.3	5.2	7.9	3.4	4.6	6.1	3.5	
2001.....	5.4	8.0	3.6	5.5	8.2	3.6	5.0	6.4	3.9	
2000.....	5.7	8.7	3.7	5.8	9.0	3.6	5.2	6.8	4.1	
1999.....	5.8	8.8	3.7	5.9	9.1	3.7	4.9	6.3	3.9	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Other diseases of arteries, arterioles and capillaries (I72-I78)										
2007.....	2.8	3.1	2.5	2.7	3.0	2.5	4.4	5.1	4.0	
2006.....	2.9	3.3	2.7	2.9	3.2	2.6	4.6	4.8	4.3	
2005.....	3.2	3.5	2.9	3.1	3.4	2.8	5.0	5.6	4.5	
2004.....	3.3	3.5	3.1	3.2	3.4	3.0	5.2	5.6	5.0	
2003.....	3.3	3.7	3.1	3.2	3.6	3.0	5.2	5.7	4.9	
2002.....	3.5	3.8	3.3	3.4	3.7	3.1	5.4	6.0	5.0	
2001.....	3.6	3.9	3.3	3.4	3.8	3.2	5.9	6.4	5.5	
2000.....	3.7	4.2	3.4	3.6	4.0	3.2	5.7	6.2	5.3	
1999.....	3.7	4.0	3.4	3.5	3.9	3.3	5.6	6.0	5.3	
Other disorders of circulatory system (I80-I99)										
2007.....	1.3	1.4	1.2	1.2	1.3	1.1	2.2	2.4	1.9	
2006.....	1.3	1.3	1.2	1.2	1.2	1.1	2.3	2.4	2.2	
2005.....	1.6	1.6	1.5	1.5	1.5	1.4	2.7	3.1	2.3	
2004.....	1.5	1.6	1.5	1.5	1.5	1.4	2.7	2.9	2.6	
2003.....	1.6	1.6	1.6	1.5	1.4	1.5	2.9	3.2	2.8	
2002.....	1.6	1.6	1.6	1.5	1.5	1.6	2.9	3.2	2.6	
2001.....	1.6	1.7	1.6	1.5	1.6	1.5	2.7	3.1	2.5	
2000.....	1.7	1.7	1.6	1.6	1.6	1.5	2.9	3.0	2.8	
1999.....	1.5	1.5	1.4	1.4	1.4	1.3	2.4	2.6	2.2	
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)										
2007.....	1.0	1.0	0.9	0.9	0.9	0.9	1.7	1.8	1.6	
2006.....	0.9	0.9	0.9	0.8	0.8	0.8	1.8	1.9	1.7	
2005.....	1.1	1.1	1.1	1.0	1.0	1.1	2.0	2.2	1.9	
2004.....	1.2	1.2	1.2	1.1	1.1	1.0	2.2	2.2	2.1	
2003.....	1.2	1.1	1.3	1.1	1.0	1.2	2.3	2.4	2.3	
2002.....	1.2	1.1	1.3	1.2	1.0	1.2	2.3	2.4	2.2	
2001.....	1.2	1.2	1.3	1.2	1.1	1.2	2.2	2.3	2.1	
2000.....	1.2	1.2	1.2	1.1	1.1	1.2	2.2	2.1	2.3	
1999.....	1.0	0.9	1.0	0.9	0.9	0.9	1.7	1.7	1.6	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Acute pharyngitis and tonsillitis (J02-J03)									
2007.....	0.0	*	*	0.0	*	*	*	*	*
2006.....	*	*	*	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*	*	*
2004.....	0.0	*	*	0.0	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	0.0	*	*	*	*	*
2000.....	0.0	*	*	0.0	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Acute laryngitis and tracheitis (J04)									
2007.....	0.0	0.0	*	0.0	*	*	*	*	*
2006.....	0.0	*	*	0.0	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*	*	*
2004.....	0.0	*	*	0.0	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	0.0	*	*	*	*	*
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)									
2007.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2006.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2005.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.1	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.1	*	*	*
1999.....	0.0	0.1	0.0	0.0	0.0	0.0	0.1	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Other diseases of the respiratory system (J09-J98,U04)									
2007.....	71.6	86.5	61.8	73.8	88.2	64.4	60.7	81.4	48.2
2006.....	72.8	88.0	63.1	75.0	89.6	65.7	61.7	82.3	49.4
2005.....	78.4	94.2	68.2	80.6	95.6	70.9	67.6	91.2	53.6
2004.....	76.1	92.9	65.5	78.1	94.4	67.9	65.3	89.6	51.1
2003.....	80.4	97.9	69.5	82.5	99.1	72.1	68.4	94.9	53.0
2002.....	81.3	100.4	69.7	83.3	101.7	72.1	70.5	97.0	55.3
2001.....	81.5	101.2	69.7	83.2	102.2	71.8	71.8	99.4	56.2
2000.....	83.7	105.7	70.8	85.3	106.7	72.9	73.5	102.8	56.8
1999.....	84.3	107.6	70.5	85.8	108.6	72.4	75.4	106.8	57.8
Influenza (J09-J11)									
2007.....	0.1	0.1	0.1	0.1	0.1	0.1	0.0	*	*
2006.....	0.3	0.3	0.2	0.3	0.3	0.3	0.1	0.2	0.1
2005.....	0.6	0.6	0.6	0.6	0.6	0.6	0.2	0.3	0.2
2004.....	0.4	0.4	0.3	0.4	0.4	0.4	0.2	0.2	0.1
2003.....	0.6	0.6	0.6	0.6	0.6	0.6	0.3	0.3	0.3
2002.....	0.2	0.3	0.2	0.2	0.3	0.2	0.1	*	*
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.6	0.7	0.6	0.6	0.7	0.6	0.3	0.4	0.3
1999.....	0.6	0.6	0.6	0.6	0.6	0.6	0.3	0.3	0.3
Pneumonia (J12-J18)									
2007.....	16.1	19.2	14.1	15.9	18.8	14.0	18.4	23.6	15.2
2006.....	17.5	20.9	15.3	17.4	20.6	15.2	19.5	24.2	16.6
2005.....	19.7	23.3	17.3	19.6	23.0	17.3	21.5	26.6	18.2
2004.....	19.4	23.3	16.9	19.2	22.9	16.8	22.1	28.9	18.0
2003.....	21.4	25.5	18.8	21.3	25.0	18.9	23.0	30.6	18.4
2002.....	22.4	26.7	19.6	22.3	26.4	19.7	23.9	30.7	19.9
2001.....	21.9	26.5	19.1	21.6	25.9	19.0	24.1	32.2	19.4
2000.....	23.1	28.2	20.1	22.9	27.7	20.0	25.3	33.5	20.5
1999.....	22.9	27.9	20.0	22.7	27.5	19.9	25.1	33.2	20.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Viral pneumonia, not elsewhere classified (J12)										
2007.....	0.0	0.1	0.0	0.0	0.1	0.0	0.0	*	*	
2006.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2005.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	
2004.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2003.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.1	
Bacterial pneumonia (J13-J15)										
2007.....	0.9	1.1	0.8	0.9	1.1	0.7	1.2	1.6	0.9	
2006.....	1.5	1.9	1.2	1.5	1.9	1.2	1.7	2.2	1.4	
2005.....	1.6	2.0	1.3	1.5	2.0	1.3	1.8	2.2	1.5	
2004.....	1.5	1.9	1.2	1.4	1.8	1.2	2.0	2.6	1.6	
2003.....	1.6	2.1	1.3	1.6	2.0	1.3	1.9	2.5	1.5	
2002.....	1.6	2.0	1.3	1.6	2.0	1.3	1.7	2.3	1.4	
2001.....	1.6	2.0	1.2	1.5	2.0	1.3	1.9	2.8	1.3	
2000.....	1.6	2.2	1.3	1.6	2.1	1.2	2.0	2.9	1.5	
1999.....	1.4	1.8	1.2	1.4	1.7	1.2	1.7	2.3	1.3	
Pneumonia due to other or unspecified organisms (J16,J18)										
2007.....	15.1	18.0	13.2	15.0	17.7	13.2	17.1	22.0	14.2	
2006.....	16.0	18.9	14.0	15.8	18.7	14.0	17.7	21.9	15.1	
2005.....	18.1	21.2	16.0	18.0	20.9	16.0	19.6	24.3	16.7	
2004.....	17.9	21.4	15.6	17.7	21.0	15.6	20.1	26.2	16.4	
2003.....	19.7	23.3	17.4	19.6	22.9	17.5	21.0	27.9	16.8	
2002.....	20.7	24.6	18.2	20.6	24.3	18.3	22.1	28.3	18.5	
2001.....	20.2	24.3	17.7	20.0	23.8	17.7	22.2	29.3	18.0	
2000.....	21.3	25.9	18.7	21.1	25.4	18.6	23.2	30.5	18.9	
1999.....	21.3	26.0	18.6	21.2	25.6	18.6	23.3	30.7	19.0	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Other acute lower respiratory infections (J20-J22,U04)										
2007.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2006.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	
2005.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1	
2004.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2003.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.2	
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	
Acute bronchitis and bronchiolitis (J20-J21)										
2007.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	*	*	
2006.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	
2005.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2004.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2003.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1	
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2	
Other and unspecified acute lower respiratory infections (J22,U04)										
2007.....	0.0	*	0.0	0.0	*	0.0	*	*	*	
2006.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2005.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2003.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2000.....	0.0	0.0	0.0	0.1	0.0	0.0	*	*	*	
1999.....	0.1	0.0	0.1	0.1	0.0	0.1	*	*	*	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Other diseases of upper respiratory tract (J30-J39)										
2007.....	0.1	0.2	0.1	0.1	0.1	0.1	0.2	0.2	0.2	
2006.....	0.1	0.2	0.1	0.1	0.2	0.1	0.3	0.3	0.2	
2005.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.2	
2004.....	0.1	0.2	0.1	0.1	0.1	0.1	0.2	0.3	0.2	
2003.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.2	
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.1	
2001.....	0.2	0.2	0.1	0.1	0.2	0.1	0.3	0.4	0.2	
2000.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.3	0.2	
1999.....	0.2	0.2	0.1	0.1	0.2	0.1	0.3	0.3	0.2	
Chronic lower respiratory diseases (J40-J47)										
2007.....	40.8	48.0	36.0	43.0	49.8	38.5	28.1	39.6	21.4	
2006.....	40.5	47.6	35.9	42.6	49.2	38.4	28.1	39.5	21.3	
2005.....	43.2	51.2	38.1	45.4	52.8	40.7	30.6	44.1	22.8	
2004.....	41.1	49.5	36.0	43.2	51.1	38.4	28.2	40.9	20.9	
2003.....	43.3	52.3	37.8	45.4	53.8	40.3	30.1	44.4	22.0	
2002.....	43.5	53.5	37.4	45.4	54.9	39.7	31.2	46.3	22.6	
2001.....	43.7	54.0	37.6	45.6	55.5	39.8	30.9	46.3	22.4	
2000.....	44.2	55.8	37.4	46.0	57.2	39.5	31.6	47.5	22.7	
1999.....	45.4	58.7	37.7	47.1	60.0	39.7	33.5	51.5	23.4	
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)										
2007.....	40.5	47.9	35.6	42.7	49.7	38.1	28.1	39.5	21.4	
2006.....	40.1	47.4	35.5	42.3	49.0	38.0	27.9	39.3	21.3	
2005.....	42.9	51.0	37.7	45.1	52.6	40.3	30.5	44.0	22.7	
2004.....	40.8	49.2	35.6	42.9	50.8	38.0	28.1	40.8	20.8	
2003.....	43.0	52.1	37.4	45.1	53.5	39.9	30.0	44.3	21.8	
2002.....	43.1	53.2	37.0	45.1	54.7	39.3	31.0	46.2	22.4	
2001.....	43.3	53.8	37.2	45.3	55.2	39.4	30.8	46.1	22.3	
2000.....	43.8	55.5	37.0	45.6	56.9	39.1	31.4	47.3	22.6	
1999.....	45.1	58.4	37.3	46.8	59.7	39.3	33.3	51.3	23.2	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Bronchitis, not specified as acute or chronic (J40)										
2007.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2006.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1	
2005.....	0.1	0.2	0.1	0.1	0.2	0.1	0.1	*	*	
2004.....	0.1	0.2	0.1	0.1	0.2	0.1	0.1	*	*	
2003.....	0.1	0.1	0.1	0.2	0.1	0.1	0.1	*	0.2	
2002.....	0.2	0.2	0.2	0.2	0.2	0.2	0.1	*	0.1	
2001.....	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2	*	
2000.....	0.2	0.2	0.2	0.2	0.3	0.2	0.2	0.2	0.2	
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2	
Chronic bronchitis (J41-J42)										
2007.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2006.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2005.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2004.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	
2003.....	0.1	0.2	0.1	0.1	0.2	0.1	0.1	*	0.1	
2002.....	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.2	*	
2001.....	0.1	0.2	0.1	0.2	0.2	0.1	0.1	*	*	
2000.....	0.2	0.2	0.2	0.2	0.2	0.2	0.1	*	*	
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.1	*	*	
Emphysema (J43)										
2007.....	4.1	5.1	3.4	4.3	5.3	3.7	2.6	4.2	1.6	
2006.....	4.1	5.0	3.5	4.4	5.2	3.8	2.6	4.0	1.7	
2005.....	4.6	5.8	3.8	4.9	6.0	4.1	3.1	4.9	1.9	
2004.....	4.6	5.8	3.8	4.9	6.1	4.1	2.7	4.6	1.6	
2003.....	5.1	6.3	4.3	5.4	6.6	4.6	3.2	5.3	2.0	
2002.....	5.4	7.0	4.4	5.7	7.2	4.8	3.4	5.8	2.0	
2001.....	5.8	7.4	4.7	6.1	7.8	5.1	3.3	5.3	2.2	
2000.....	6.0	7.8	4.9	6.4	8.1	5.3	3.8	6.2	2.4	
1999.....	6.5	8.6	5.2	6.9	8.9	5.6	4.0	6.7	2.3	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Other chronic obstructive pulmonary disease (J44)									
2007.....	35.1	41.7	30.8	37.2	43.5	33.2	22.7	32.6	17.0
2006.....	34.6	41.2	30.5	36.7	42.9	32.8	22.4	32.5	16.5
2005.....	36.7	43.9	32.2	38.9	45.5	34.6	24.2	36.1	17.4
2004.....	34.7	42.0	30.1	36.7	43.7	32.4	22.1	32.9	15.9
2003.....	36.2	44.2	31.3	38.2	45.8	33.6	23.3	35.9	16.2
2002.....	35.9	44.7	30.7	37.8	46.1	32.8	24.0	36.8	16.8
2001.....	35.7	44.8	30.3	37.5	46.2	32.4	23.6	37.3	16.1
2000.....	35.8	45.9	29.9	37.5	47.3	31.9	23.5	37.3	15.8
1999.....	36.5	47.9	29.8	38.1	49.3	31.6	25.1	40.6	16.5
Asthma (J45-J46)									
2007.....	1.1	0.9	1.2	0.9	0.7	1.0	2.5	2.5	2.6
2006.....	1.2	0.9	1.3	0.9	0.7	1.1	2.7	2.5	2.8
2005.....	1.3	1.0	1.5	1.0	0.7	1.3	3.0	2.7	3.3
2004.....	1.3	1.0	1.4	1.0	0.8	1.2	3.0	3.0	3.1
2003.....	1.4	1.1	1.6	1.1	0.9	1.3	3.2	2.9	3.3
2002.....	1.5	1.2	1.7	1.2	0.9	1.4	3.4	3.3	3.4
2001.....	1.5	1.2	1.7	1.2	0.9	1.5	3.6	3.2	3.8
2000.....	1.6	1.3	1.8	1.3	1.0	1.5	3.9	3.5	4.1
1999.....	1.7	1.3	2.0	1.4	1.0	1.7	3.9	3.5	4.1
Bronchiectasis (J47)									
2007.....	0.3	0.2	0.4	0.3	0.2	0.4	*	*	*
2006.....	0.3	0.2	0.4	0.3	0.2	0.4	0.1	0.2	*
2005.....	0.3	0.2	0.4	0.3	0.2	0.4	0.1	*	*
2004.....	0.3	0.2	0.4	0.3	0.2	0.4	0.1	*	0.1
2003.....	0.3	0.2	0.4	0.3	0.2	0.4	0.1	*	*
2002.....	0.3	0.2	0.4	0.3	0.2	0.4	0.1	*	*
2001.....	0.3	0.2	0.4	0.4	0.2	0.4	0.1	*	*
2000.....	0.3	0.2	0.4	0.4	0.3	0.4	0.1	*	*
1999.....	0.3	0.3	0.4	0.4	0.3	0.4	0.2	*	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Lung diseases due to external agents (J60-J70)										
2007.....	5.5	8.0	4.0	5.6	8.2	4.1	5.5	8.0	4.0	
2006.....	5.7	8.3	4.0	5.8	8.4	4.1	5.7	8.5	4.1	
2005.....	5.9	8.4	4.4	6.0	8.5	4.4	6.6	9.5	4.9	
2004.....	6.0	8.6	4.3	6.0	8.7	4.4	6.3	9.3	4.6	
2003.....	6.3	9.0	4.7	6.3	9.1	4.6	6.9	9.7	5.4	
2002.....	6.5	9.4	4.8	6.5	9.5	4.8	6.8	9.9	5.2	
2001.....	6.6	9.5	4.9	6.6	9.6	4.8	7.3	10.1	5.8	
2000.....	6.5	9.7	4.7	6.5	9.8	4.7	7.0	10.2	5.2	
1999.....	6.1	9.2	4.4	6.0	9.1	4.3	7.1	10.7	5.1	
Pneumoconioses and chemical effects (J60-J66,J68)										
2007.....	0.3	0.7	0.0	0.3	0.8	0.0	0.2	0.4	*	
2006.....	0.3	0.7	0.0	0.3	0.8	0.0	0.2	0.5	*	
2005.....	0.3	0.8	0.0	0.4	0.9	0.0	0.2	0.5	*	
2004.....	0.3	0.9	0.0	0.4	0.9	0.0	0.2	0.6	*	
2003.....	0.4	1.0	0.0	0.4	1.0	0.0	0.2	0.5	*	
2002.....	0.4	1.0	0.0	0.4	1.0	0.0	0.2	0.6	*	
2001.....	0.4	1.0	0.0	0.4	1.1	0.0	0.2	0.7	*	
2000.....	0.4	1.1	0.0	0.4	1.1	0.0	0.2	0.7	*	
1999.....	0.4	1.0	0.0	0.4	1.1	0.0	0.3	0.8	*	
Pneumonitis due to solids and liquids (J69)										
2007.....	5.2	7.3	4.0	5.3	7.4	4.0	5.3	7.5	4.0	
2006.....	5.3	7.5	4.0	5.4	7.6	4.1	5.5	8.0	4.1	
2005.....	5.6	7.6	4.4	5.6	7.6	4.4	6.4	9.0	4.9	
2004.....	5.6	7.7	4.3	5.6	7.7	4.3	6.0	8.6	4.6	
2003.....	5.9	8.0	4.6	5.8	8.0	4.6	6.7	9.1	5.3	
2002.....	6.1	8.4	4.7	6.1	8.5	4.7	6.6	9.2	5.1	
2001.....	6.1	8.5	4.8	6.1	8.5	4.8	7.0	9.4	5.7	
2000.....	6.1	8.6	4.6	6.0	8.6	4.6	6.8	9.4	5.2	
1999.....	5.6	8.1	4.3	5.6	8.0	4.3	6.8	9.9	5.1	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Pneumonitis due to food and vomit (J69.0)										
2007.....	5.2	7.3	4.0	5.3	7.4	4.0	5.3	7.5	4.0	
2006.....	5.3	7.5	4.0	5.4	7.6	4.1	5.5	8.0	4.1	
2005.....	5.6	7.6	4.4	5.6	7.6	4.4	6.4	9.0	4.9	
2004.....	5.6	7.7	4.3	5.6	7.7	4.3	6.0	8.6	4.6	
2003.....	5.9	8.0	4.6	5.8	8.0	4.6	6.7	9.1	5.3	
2002.....	6.1	8.4	4.7	6.1	8.5	4.7	6.6	9.2	5.1	
2001.....	6.1	8.5	4.8	6.1	8.5	4.8	7.0	9.4	5.7	
2000.....	6.1	8.6	4.6	6.0	8.6	4.6	6.8	9.4	5.2	
1999.....	5.6	8.1	4.3	5.6	8.0	4.3	6.8	9.9	5.1	
Other pneumonitis due to solids and liquids (J69.1-J69.8)										
2007.....	*	*	*	*	*	*	*	*	*	
2006.....	*	*	*	*	*	*	*	*	*	
2005.....	*	*	*	*	*	*	*	*	*	
2004.....	*	*	*	*	*	*	*	*	*	
2003.....	*	*	*	*	*	*	*	*	*	
2002.....	*	*	*	*	*	*	*	*	*	
2001.....	*	*	*	*	*	*	*	*	*	
2000.....	*	*	*	*	*	*	*	*	*	
1999.....	*	*	*	*	*	*	*	*	*	
Other lung diseases due to external agents (J67,J70)										
2007.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2006.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2005.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2003.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2001.....	0.0	0.0	0.0	0.0	0.1	0.0	*	*	*	
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)										
2007.....	0.2	0.3	0.1	0.2	0.3	0.1	0.3	0.5	0.2	
2006.....	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.4	0.3	
2005.....	0.2	0.3	0.1	0.2	0.3	0.2	0.3	0.5	0.1	
2004.....	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.4	0.2	
2003.....	0.2	0.3	0.2	0.2	0.3	0.2	0.4	0.6	0.2	
2002.....	0.2	0.4	0.2	0.2	0.3	0.1	0.4	0.7	0.3	
2001.....	0.3	0.4	0.2	0.2	0.3	0.2	0.4	0.6	0.2	
2000.....	0.2	0.4	0.2	0.2	0.4	0.2	0.4	0.7	0.3	
1999.....	0.3	0.4	0.2	0.3	0.4	0.2	0.5	0.8	0.3	
Pleural effusion and plaque (J90,J92)										
2007.....	0.4	0.4	0.3	0.4	0.4	0.3	0.3	0.4	0.2	
2006.....	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.4	0.3	
2005.....	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.4	0.2	
2004.....	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.4	0.3	
2003.....	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.4	0.3	
2002.....	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.4	0.3	
2001.....	0.3	0.4	0.3	0.3	0.4	0.3	0.5	0.5	0.4	
2000.....	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.5	0.3	
1999.....	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.4	0.3	
All other diseases of respiratory system (J80-J84,J93-J98)										
2007.....	8.2	10.1	7.0	8.4	10.2	7.0	7.7	9.1	6.8	
2006.....	8.1	9.9	6.9	8.3	10.1	7.0	7.3	8.7	6.4	
2005.....	8.1	9.7	7.0	8.2	9.7	7.1	7.8	9.4	6.8	
2004.....	8.4	10.1	7.2	8.5	10.3	7.3	7.6	9.2	6.6	
2003.....	7.9	9.5	6.9	8.0	9.6	6.9	7.1	8.6	6.1	
2002.....	7.9	9.4	6.9	8.0	9.5	6.9	7.3	8.5	6.6	
2001.....	8.4	10.0	7.3	8.5	10.1	7.3	8.2	9.2	7.7	
2000.....	8.4	10.2	7.2	8.5	10.3	7.3	8.1	9.6	7.2	
1999.....	8.3	10.0	7.2	8.3	10.1	7.1	8.2	9.5	7.5	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
XI. Diseases of the digestive system (K00-K92)									
2007.....	27.9	32.7	23.3	28.2	33.0	23.6	28.7	35.6	23.4
2006.....	27.9	32.5	23.7	28.2	32.7	23.9	28.9	35.5	23.7
2005.....	28.4	33.3	24.0	28.5	33.2	24.1	30.7	38.0	25.0
2004.....	28.8	33.4	24.5	28.8	33.2	24.5	31.6	39.6	25.5
2003.....	29.8	34.8	25.1	29.8	34.7	25.2	32.7	41.1	26.2
2002.....	29.9	35.0	25.3	30.0	34.9	25.4	32.4	40.1	26.5
2001.....	30.3	35.5	25.7	30.1	34.9	25.6	34.6	44.0	27.5
2000.....	30.4	35.8	25.5	30.0	35.3	25.3	35.3	43.8	28.8
1999.....	30.3	36.0	25.3	30.0	35.5	25.0	35.3	44.1	28.7
Diseases of oral cavity, salivary glands and jaws (K00-K14)									
2007.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2006.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2005.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2004.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2003.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.1	0.0	0.0	0.1	0.0	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Diseases of other parts of digestive system (K20-K92)									
2007.....	27.8	32.7	23.3	28.1	32.9	23.5	28.7	35.5	23.3
2006.....	27.9	32.5	23.6	28.1	32.6	23.9	28.8	35.4	23.7
2005.....	28.4	33.2	23.9	28.4	33.1	24.0	30.6	37.9	25.0
2004.....	28.7	33.3	24.4	28.7	33.1	24.4	31.6	39.6	25.4
2003.....	29.8	34.8	25.1	29.7	34.6	25.2	32.6	41.1	26.2
2002.....	29.9	35.0	25.3	29.9	34.8	25.3	32.3	40.1	26.4
2001.....	30.2	35.4	25.6	30.1	34.9	25.6	34.5	44.0	27.4
2000.....	30.3	35.7	25.5	30.0	35.2	25.2	35.2	43.8	28.7
1999.....	30.3	35.9	25.3	30.0	35.4	25.0	35.3	44.0	28.6

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Diseases of esophagus (K20-K22)									
2007.....	0.6	0.7	0.5	0.6	0.7	0.5	0.5	0.6	0.4
2006.....	0.6	0.7	0.5	0.6	0.7	0.5	0.4	0.6	0.3
2005.....	0.6	0.8	0.5	0.6	0.8	0.5	0.5	0.6	0.5
2004.....	0.6	0.7	0.6	0.6	0.7	0.6	0.6	0.8	0.4
2003.....	0.7	0.8	0.6	0.7	0.9	0.6	0.6	0.7	0.5
2002.....	0.7	0.9	0.6	0.8	0.9	0.6	0.7	0.8	0.5
2001.....	0.7	0.9	0.7	0.8	0.9	0.7	0.6	0.9	0.5
2000.....	0.8	1.0	0.7	0.8	1.0	0.7	0.7	1.0	0.5
1999.....	0.8	1.0	0.7	0.8	1.0	0.7	0.7	1.0	0.5
Peptic ulcer (K25-K28)									
2007.....	1.0	1.1	0.8	1.0	1.1	0.8	1.0	1.4	0.7
2006.....	1.1	1.2	0.9	1.0	1.2	0.9	1.2	1.7	0.9
2005.....	1.1	1.4	0.9	1.1	1.3	0.9	1.3	1.8	0.9
2004.....	1.2	1.4	1.1	1.2	1.4	1.1	1.3	1.7	1.1
2003.....	1.3	1.6	1.1	1.3	1.5	1.2	1.4	2.1	1.0
2002.....	1.4	1.7	1.2	1.4	1.7	1.2	1.4	2.0	1.1
2001.....	1.6	1.8	1.4	1.6	1.8	1.4	1.6	2.3	1.1
2000.....	1.6	2.0	1.4	1.6	2.0	1.4	1.7	2.3	1.3
1999.....	1.7	2.0	1.4	1.7	2.0	1.4	1.9	2.4	1.4
Gastric ulcer (K25)									
2007.....	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.6	0.3
2006.....	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.5	0.3
2005.....	0.3	0.4	0.3	0.3	0.4	0.3	0.5	0.6	0.4
2004.....	0.4	0.4	0.3	0.4	0.4	0.4	0.5	0.6	0.4
2003.....	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.7	0.4
2002.....	0.4	0.5	0.4	0.4	0.4	0.4	0.5	0.6	0.4
2001.....	0.5	0.4	0.4	0.4	0.4	0.4	0.6	0.8	0.4
2000.....	0.5	0.5	0.4	0.5	0.5	0.4	0.7	1.0	0.4
1999.....	0.5	0.5	0.4	0.5	0.5	0.4	0.7	1.0	0.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Duodenal ulcer (K26)									
2007.....	0.3	0.4	0.3	0.4	0.4	0.3	0.3	0.4	0.2
2006.....	0.4	0.5	0.3	0.4	0.5	0.3	0.4	0.6	0.3
2005.....	0.4	0.5	0.3	0.4	0.5	0.3	0.4	0.7	0.2
2004.....	0.4	0.5	0.3	0.4	0.5	0.3	0.4	0.5	0.2
2003.....	0.4	0.6	0.4	0.5	0.6	0.3	0.5	0.7	0.3
2002.....	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.6	0.3
2001.....	0.5	0.7	0.4	0.6	0.7	0.4	0.4	0.6	0.3
2000.....	0.5	0.7	0.4	0.6	0.7	0.4	0.5	0.6	0.3
1999.....	0.5	0.7	0.4	0.5	0.7	0.4	0.5	0.6	0.4
Peptic ulcer, site unspecified (K27)									
2007.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.2
2006.....	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.5	0.3
2005.....	0.4	0.4	0.3	0.4	0.4	0.3	0.4	0.5	0.3
2004.....	0.4	0.5	0.4	0.4	0.4	0.4	0.4	0.5	0.4
2003.....	0.5	0.6	0.4	0.5	0.6	0.4	0.5	0.7	0.3
2002.....	0.5	0.6	0.4	0.5	0.6	0.4	0.5	0.7	0.3
2001.....	0.6	0.7	0.5	0.6	0.7	0.5	0.6	0.9	0.4
2000.....	0.6	0.8	0.5	0.6	0.8	0.5	0.6	0.7	0.5
1999.....	0.6	0.8	0.6	0.6	0.8	0.6	0.6	0.8	0.5
Gastrojejunal ulcer (K28)									
2007.....	0.0	*	*	0.0	*	*	*	*	*
2006.....	0.0	*	*	0.0	*	*	*	*	*
2005.....	0.0	*	*	0.0	*	*	*	*	*
2004.....	0.0	0.0	0.0	0.0	*	0.0	*	*	*
2003.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2001.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
1999.....	0.0	*	*	0.0	*	*	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Gastritis and duodenitis (K29)										
2007.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2006.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2005.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2004.....	0.1	0.2	0.1	0.1	0.1	0.1	0.2	0.2	0.1	
2003.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.1	
2002.....	0.1	0.1	0.1	0.1	0.2	0.1	0.1	*	0.1	
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2000.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	*	0.1	
1999.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.2	
Dyspepsia and other diseases of stomach and duodenum (K30-K31)										
2007.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2	
2006.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	*	0.2	
2005.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2	
2004.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.2	
2003.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2	0.3	
2002.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3	
2001.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	
2000.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2	
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	
Diseases of appendix (K35-K38)										
2007.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.2	
2006.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.2	
2005.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.1	
2004.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.1	
2003.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.2	
2002.....	0.2	0.2	0.1	0.1	0.2	0.1	0.2	0.4	0.1	
2001.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.4	0.2	
2000.....	0.1	0.2	0.1	0.1	0.2	0.1	0.3	0.4	0.2	
1999.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.4	0.1	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Hernia (K40-K46)										
2007.....	0.5	0.5	0.5	0.5	0.5	0.5	0.6	0.7	0.5	
2006.....	0.5	0.6	0.5	0.6	0.6	0.5	0.6	0.7	0.6	
2005.....	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.6	0.5	
2004.....	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.6	0.5	
2003.....	0.6	0.6	0.5	0.6	0.6	0.5	0.6	0.7	0.6	
2002.....	0.5	0.5	0.6	0.6	0.5	0.6	0.5	0.6	0.5	
2001.....	0.5	0.5	0.5	0.5	0.5	0.5	0.6	0.8	0.5	
2000.....	0.5	0.5	0.5	0.6	0.6	0.6	0.6	0.6	0.6	
1999.....	0.5	0.6	0.5	0.5	0.6	0.5	0.6	0.6	0.6	
Crohn's disease and ulcerative colitis (K50-K51)										
2007.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	
2006.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	
2005.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	*	0.2	
2004.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.1	
2003.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	
2002.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.1	
2001.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	
2000.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	*	0.2	
1999.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	
Vascular disorders and obstruction of intestine without hernia (K55-K56)										
2007.....	4.5	4.1	4.7	4.5	4.0	4.7	5.2	5.4	5.0	
2006.....	4.6	4.2	4.9	4.6	4.1	5.0	5.4	5.9	5.0	
2005.....	4.7	4.3	5.0	4.7	4.1	5.0	5.7	6.1	5.3	
2004.....	4.8	4.3	5.1	4.8	4.2	5.1	6.0	6.6	5.6	
2003.....	5.1	4.6	5.3	5.0	4.4	5.4	6.0	6.9	5.5	
2002.....	5.0	4.5	5.3	5.0	4.4	5.3	6.0	6.3	5.7	
2001.....	5.1	4.6	5.3	5.0	4.5	5.3	6.2	7.0	5.8	
2000.....	5.2	4.7	5.4	5.1	4.5	5.4	6.6	7.3	6.1	
1999.....	5.1	4.6	5.4	5.1	4.4	5.4	6.4	6.6	6.2	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Diverticular disease of intestine (K57)										
2007.....	1.0	0.8	1.1	1.0	0.8	1.1	0.7	0.6	0.8	
2006.....	1.0	0.8	1.1	1.1	0.8	1.2	0.9	0.8	1.0	
2005.....	1.0	0.8	1.1	1.0	0.8	1.2	0.9	0.8	1.0	
2004.....	1.1	0.9	1.3	1.2	0.9	1.3	1.0	0.9	1.0	
2003.....	1.2	1.0	1.3	1.2	1.0	1.3	1.0	0.8	1.1	
2002.....	1.2	0.9	1.3	1.2	1.0	1.3	1.1	1.0	1.1	
2001.....	1.2	1.0	1.3	1.2	1.0	1.4	1.1	0.8	1.2	
2000.....	1.2	1.0	1.3	1.2	1.0	1.3	1.0	0.8	1.2	
1999.....	1.3	1.0	1.4	1.3	1.1	1.4	1.1	0.8	1.2	
Other diseases of intestines and peritoneum (K52,K58-K63)										
2007.....	2.0	1.9	2.0	2.0	1.8	2.0	2.1	2.2	2.0	
2006.....	1.9	1.7	1.9	1.8	1.7	1.9	2.0	2.1	1.9	
2005.....	1.8	1.7	1.9	1.8	1.7	1.9	2.1	2.4	1.8	
2004.....	1.7	1.5	1.7	1.7	1.5	1.8	1.7	1.9	1.5	
2003.....	1.7	1.6	1.7	1.7	1.6	1.7	1.9	2.0	1.8	
2002.....	1.7	1.6	1.7	1.7	1.6	1.7	1.7	1.9	1.5	
2001.....	1.7	1.6	1.7	1.7	1.5	1.8	1.9	2.1	1.8	
2000.....	1.6	1.5	1.7	1.6	1.5	1.7	1.8	1.9	1.7	
1999.....	1.6	1.6	1.7	1.6	1.5	1.7	1.7	2.0	1.5	
Diseases of peritoneum (K65-K66)										
2007.....	0.6	0.6	0.5	0.5	0.6	0.5	0.9	1.0	0.8	
2006.....	0.6	0.6	0.6	0.6	0.5	0.6	0.7	0.7	0.6	
2005.....	0.6	0.6	0.6	0.6	0.6	0.6	0.8	0.8	0.9	
2004.....	0.6	0.6	0.6	0.6	0.6	0.5	0.8	0.8	0.8	
2003.....	0.6	0.7	0.6	0.6	0.6	0.6	0.9	0.9	1.0	
2002.....	0.7	0.6	0.7	0.6	0.6	0.7	1.0	0.9	1.0	
2001.....	0.7	0.7	0.6	0.6	0.7	0.6	1.1	1.2	1.0	
2000.....	0.7	0.7	0.7	0.7	0.7	0.6	1.1	1.3	1.0	
1999.....	0.7	0.8	0.6	0.7	0.7	0.6	1.1	1.2	1.0	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Diseases of liver (K70-K76)										
2007.....	12.2	16.4	8.4	12.5	16.8	8.5	11.0	15.6	7.4	
2006.....	11.8	15.8	8.2	12.1	16.2	8.4	10.5	14.6	7.3	
2005.....	12.0	16.2	8.2	12.2	16.4	8.3	11.4	16.1	7.8	
2004.....	12.0	16.2	8.2	12.1	16.4	8.2	11.8	16.6	7.9	
2003.....	12.4	16.8	8.4	12.5	17.0	8.5	12.2	17.6	8.1	
2002.....	12.3	16.7	8.4	12.4	16.9	8.4	12.3	16.9	8.6	
2001.....	12.3	16.7	8.3	12.3	16.7	8.3	13.0	18.9	8.5	
2000.....	12.2	16.7	8.2	12.1	16.7	8.0	13.2	18.4	9.1	
1999.....	12.3	17.0	8.1	12.1	16.9	7.9	13.9	19.6	9.5	
Alcoholic liver disease (K70)										
2007.....	4.5	6.8	2.3	4.7	7.1	2.4	3.4	5.5	1.8	
2006.....	4.1	6.3	2.2	4.3	6.5	2.3	3.0	4.7	1.7	
2005.....	4.2	6.4	2.2	4.3	6.6	2.3	3.5	5.4	1.9	
2004.....	4.2	6.3	2.2	4.3	6.5	2.2	3.5	5.6	1.9	
2003.....	4.2	6.4	2.1	4.2	6.6	2.1	3.8	6.0	2.0	
2002.....	4.2	6.3	2.2	4.2	6.5	2.2	3.8	5.8	2.3	
2001.....	4.3	6.6	2.2	4.3	6.6	2.2	4.4	6.9	2.4	
2000.....	4.3	6.8	2.2	4.3	6.9	2.0	4.6	7.0	2.8	
1999.....	4.3	6.9	2.1	4.3	6.8	2.0	5.1	8.1	2.8	
Chronic hepatitis, not elsewhere classified (K73)										
2007.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2006.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2005.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2003.....	0.0	0.1	0.0	0.0	0.1	0.0	*	*	*	
2002.....	0.2	0.1	0.2	0.2	0.1	0.3	0.2	*	0.2	
2001.....	0.2	0.1	0.2	0.2	0.1	0.2	0.2	*	0.2	
2000.....	0.2	0.1	0.2	0.1	0.1	0.2	0.2	*	0.2	
1999.....	0.2	0.1	0.2	0.1	0.1	0.2	0.2	*	0.3	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Fibrosis and cirrhosis of liver (K74)									
2007.....	4.6	5.8	3.5	4.7	6.0	3.6	4.0	5.5	2.7
2006.....	4.6	5.8	3.5	4.8	6.0	3.7	3.9	5.5	2.7
2005.....	4.7	6.0	3.6	4.8	6.1	3.7	4.2	6.0	2.8
2004.....	4.8	6.1	3.6	4.9	6.2	3.7	4.3	6.1	2.9
2003.....	5.1	6.5	3.8	5.3	6.6	4.0	4.5	6.3	3.1
2002.....	5.0	6.4	3.8	5.2	6.6	3.9	4.5	6.1	3.2
2001.....	5.1	6.5	3.8	5.2	6.5	3.9	4.7	6.8	3.0
2000.....	5.1	6.5	3.8	5.1	6.6	3.8	4.7	6.4	3.3
1999.....	5.0	6.6	3.7	5.1	6.6	3.8	4.8	6.7	3.2
Other diseases of liver (K71-K72,K75-K76)									
2007.....	3.1	3.7	2.5	3.0	3.7	2.4	3.6	4.6	2.9
2006.....	3.0	3.7	2.5	3.0	3.7	2.4	3.5	4.4	2.9
2005.....	3.0	3.8	2.4	2.9	3.7	2.2	3.7	4.6	3.0
2004.....	3.0	3.7	2.4	2.9	3.6	2.3	3.9	4.9	3.1
2003.....	3.0	3.8	2.4	3.0	3.7	2.3	3.9	5.1	3.0
2002.....	2.9	3.8	2.1	2.8	3.7	2.1	3.8	5.0	2.9
2001.....	2.7	3.5	2.1	2.7	3.4	2.0	3.8	5.0	2.9
2000.....	2.6	3.3	2.0	2.5	3.2	1.9	3.8	4.9	2.9
1999.....	2.7	3.4	2.1	2.5	3.3	1.9	3.9	4.7	3.2
Cholelithiasis and other disorders of gallbladder (K80-K82)									
2007.....	1.0	1.1	0.9	1.0	1.1	0.9	1.1	1.1	1.0
2006.....	1.0	1.1	0.9	1.0	1.1	0.9	1.2	1.1	1.2
2005.....	1.0	1.1	0.9	1.0	1.1	0.9	1.2	1.3	1.2
2004.....	1.0	1.2	0.9	1.0	1.2	0.9	1.2	1.2	1.2
2003.....	1.0	1.2	0.9	1.0	1.1	0.9	1.1	1.3	1.0
2002.....	1.0	1.2	0.9	1.0	1.2	0.9	1.1	1.1	1.1
2001.....	1.0	1.2	0.9	1.0	1.2	0.9	1.3	1.4	1.2
2000.....	1.0	1.2	0.9	1.0	1.2	0.9	1.2	1.1	1.2
1999.....	1.0	1.2	0.9	1.0	1.2	0.9	1.1	1.1	1.0

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Disorders of biliary tract and pancreas (K83-K86)										
2007.....	1.5	1.9	1.2	1.5	1.8	1.2	2.0	2.4	1.7	
2006.....	1.6	1.9	1.3	1.6	1.9	1.3	2.0	2.5	1.5	
2005.....	1.6	2.0	1.3	1.5	1.9	1.3	2.1	2.7	1.7	
2004.....	1.6	2.0	1.3	1.6	1.9	1.3	2.4	3.0	1.9	
2003.....	1.7	2.0	1.4	1.6	1.9	1.3	2.4	3.2	1.8	
2002.....	1.7	2.1	1.3	1.6	2.0	1.3	2.2	3.1	1.7	
2001.....	1.8	2.1	1.5	1.7	2.0	1.4	2.5	3.1	2.0	
2000.....	1.7	2.1	1.4	1.6	1.9	1.3	2.5	3.3	1.9	
1999.....	1.6	2.0	1.3	1.5	1.9	1.3	2.4	3.2	1.8	
Pancreatitis (K85,K86.0-K86.1)										
2007.....	1.0	1.3	0.8	1.0	1.3	0.8	1.4	1.8	1.1	
2006.....	1.1	1.4	0.9	1.1	1.4	0.9	1.4	1.9	1.0	
2005.....	1.1	1.4	0.8	1.1	1.4	0.8	1.5	2.0	1.1	
2004.....	1.2	1.4	0.9	1.1	1.4	0.8	1.8	2.5	1.4	
2003.....	1.2	1.5	0.9	1.1	1.4	0.9	1.8	2.6	1.3	
2002.....	1.2	1.5	0.9	1.2	1.5	0.9	1.6	2.3	1.1	
2001.....	1.3	1.5	1.1	1.2	1.5	1.0	1.9	2.4	1.5	
2000.....	1.2	1.5	1.0	1.2	1.4	0.9	1.9	2.6	1.3	
1999.....	1.2	1.5	1.0	1.1	1.4	0.9	1.9	2.6	1.3	
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)										
2007.....	0.5	0.5	0.4	0.5	0.5	0.4	0.6	0.6	0.6	
2006.....	0.5	0.5	0.4	0.5	0.5	0.4	0.5	0.6	0.5	
2005.....	0.5	0.6	0.4	0.5	0.5	0.4	0.6	0.7	0.5	
2004.....	0.5	0.5	0.4	0.5	0.5	0.4	0.5	0.6	0.5	
2003.....	0.5	0.6	0.4	0.5	0.6	0.4	0.6	0.6	0.6	
2002.....	0.5	0.6	0.4	0.5	0.6	0.4	0.6	0.7	0.6	
2001.....	0.5	0.5	0.4	0.4	0.5	0.4	0.6	0.7	0.5	
2000.....	0.5	0.5	0.4	0.4	0.5	0.4	0.6	0.7	0.5	
1999.....	0.4	0.5	0.4	0.4	0.5	0.4	0.5	0.6	0.5	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
All other diseases of digestive system (K90,K92)										
2007.....	2.3	2.7	2.0	2.3	2.7	2.0	2.9	3.6	2.5	
2006.....	2.5	3.0	2.1	2.4	2.9	2.1	3.3	4.0	2.8	
2005.....	2.6	3.0	2.3	2.6	2.9	2.3	3.3	4.0	2.9	
2004.....	2.6	3.1	2.3	2.6	3.0	2.3	3.4	4.1	2.8	
2003.....	2.8	3.2	2.4	2.8	3.2	2.4	3.6	4.2	3.1	
2002.....	2.9	3.3	2.5	2.8	3.2	2.5	3.6	4.4	3.0	
2001.....	2.9	3.4	2.5	2.9	3.3	2.5	3.8	4.7	3.1	
2000.....	2.9	3.4	2.6	2.9	3.3	2.5	4.0	4.8	3.4	
1999.....	2.9	3.3	2.5	2.8	3.3	2.5	3.6	4.4	3.0	
XIII. Diseases of the skin and subcutaneous tissue (L00-L98)										
2007.....	1.2	1.2	1.2	1.1	1.1	1.0	2.6	2.4	2.7	
2006.....	1.2	1.2	1.3	1.1	1.1	1.1	2.8	2.6	2.9	
2005.....	1.4	1.3	1.4	1.2	1.2	1.2	3.3	3.0	3.5	
2004.....	1.4	1.3	1.5	1.2	1.1	1.2	3.6	3.1	3.8	
2003.....	1.4	1.3	1.5	1.2	1.1	1.3	3.6	3.5	3.7	
2002.....	1.4	1.3	1.4	1.1	1.1	1.1	3.9	3.3	4.2	
2001.....	1.3	1.2	1.4	1.1	1.0	1.2	3.6	3.4	3.7	
2000.....	1.4	1.2	1.4	1.1	1.0	1.2	3.9	3.7	4.0	
1999.....	1.3	1.3	1.4	1.1	1.0	1.1	4.2	4.0	4.4	
Infections of skin and subcutaneous tissue (L00-L08)										
2007.....	0.6	0.6	0.5	0.6	0.6	0.5	0.8	0.6	0.8	
2006.....	0.6	0.6	0.5	0.6	0.6	0.5	0.8	0.7	0.8	
2005.....	0.5	0.6	0.5	0.5	0.6	0.5	0.7	0.7	0.7	
2004.....	0.5	0.6	0.5	0.5	0.6	0.5	0.7	0.8	0.7	
2003.....	0.5	0.6	0.5	0.5	0.5	0.5	0.7	0.8	0.6	
2002.....	0.5	0.5	0.4	0.4	0.5	0.4	0.6	0.6	0.6	
2001.....	0.4	0.4	0.4	0.4	0.4	0.4	0.6	0.5	0.6	
2000.....	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.5	0.5	
1999.....	0.4	0.4	0.3	0.4	0.4	0.3	0.5	0.6	0.5	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/		Black 2/			
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)									
2007.....	0.6	0.6	0.7	0.5	0.5	0.5	1.8	1.7	1.9
2006.....	0.7	0.6	0.7	0.5	0.5	0.6	2.0	1.8	2.1
2005.....	0.8	0.7	0.9	0.6	0.6	0.7	2.6	2.3	2.8
2004.....	0.8	0.7	0.9	0.6	0.5	0.7	2.8	2.4	3.0
2003.....	0.8	0.7	1.0	0.7	0.6	0.7	3.0	2.8	3.1
2002.....	0.9	0.8	1.0	0.7	0.6	0.8	3.3	2.7	3.6
2001.....	0.9	0.7	1.0	0.7	0.6	0.8	3.1	2.9	3.1
2000.....	0.9	0.8	1.1	0.7	0.6	0.8	3.4	3.2	3.5
1999.....	1.0	0.9	1.1	0.8	0.7	0.8	3.8	3.4	3.9
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)									
2007.....	4.4	3.3	5.1	4.2	3.2	5.0	5.7	4.9	6.5
2006.....	4.5	3.3	5.3	4.3	3.2	5.1	5.8	4.5	6.8
2005.....	4.7	3.5	5.6	4.6	3.4	5.4	6.0	4.4	7.3
2004.....	4.8	3.5	5.7	4.6	3.3	5.4	6.5	5.2	7.5
2003.....	4.9	3.6	5.8	4.7	3.4	5.6	6.5	5.4	7.4
2002.....	5.0	3.6	5.9	4.8	3.5	5.7	6.4	5.1	7.4
2001.....	5.0	3.6	5.9	4.8	3.5	5.7	6.4	5.1	7.5
2000.....	5.0	3.7	5.9	4.8	3.5	5.7	6.6	5.7	7.4
1999.....	4.9	3.5	5.9	4.7	3.4	5.6	6.3	5.0	7.4
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)									
2007.....	0.8	0.4	1.0	0.8	0.4	1.0	0.6	0.4	0.7
2006.....	0.8	0.5	1.1	0.9	0.5	1.1	0.6	0.3	0.8
2005.....	0.8	0.5	1.1	0.9	0.5	1.1	0.6	0.3	0.8
2004.....	0.9	0.5	1.1	0.9	0.5	1.2	0.7	0.4	0.9
2003.....	0.9	0.5	1.2	1.0	0.5	1.2	0.7	0.5	0.8
2002.....	1.0	0.6	1.2	1.0	0.6	1.3	0.7	0.4	0.8
2001.....	0.9	0.5	1.2	1.0	0.5	1.3	0.7	0.4	0.8
2000.....	1.0	0.6	1.2	1.0	0.6	1.3	0.6	0.4	0.8
1999.....	1.0	0.6	1.3	1.0	0.6	1.3	0.6	0.4	0.7

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Systemic lupus erythematosus (M32)										
2007.....	0.4	0.1	0.6	0.3	0.1	0.5	1.1	0.3	1.8	
2006.....	0.4	0.1	0.7	0.3	0.1	0.5	1.2	0.4	1.9	
2005.....	0.4	0.1	0.8	0.3	0.1	0.5	1.3	0.4	2.1	
2004.....	0.4	0.2	0.7	0.3	0.1	0.5	1.4	0.5	2.1	
2003.....	0.5	0.1	0.7	0.3	0.1	0.5	1.2	0.3	2.0	
2002.....	0.4	0.1	0.7	0.3	0.1	0.5	1.4	0.2	2.3	
2001.....	0.5	0.2	0.8	0.3	0.1	0.6	1.3	0.3	2.1	
2000.....	0.5	0.2	0.8	0.3	0.1	0.6	1.3	0.4	2.2	
1999.....	0.5	0.2	0.8	0.4	0.1	0.6	1.3	0.3	2.2	
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)										
2007.....	1.4	1.0	1.7	1.3	1.0	1.6	1.9	1.6	2.2	
2006.....	1.4	1.0	1.8	1.4	1.0	1.7	1.9	1.4	2.3	
2005.....	1.6	1.1	1.9	1.5	1.1	1.8	2.1	1.4	2.5	
2004.....	1.6	1.1	1.9	1.5	1.1	1.8	2.1	1.5	2.6	
2003.....	1.6	1.2	1.9	1.5	1.1	1.8	2.3	1.9	2.7	
2002.....	1.7	1.2	2.0	1.6	1.1	1.9	2.2	1.7	2.6	
2001.....	1.7	1.3	2.0	1.6	1.2	1.9	2.3	1.8	2.6	
2000.....	1.7	1.3	2.0	1.7	1.2	2.0	2.4	2.1	2.6	
1999.....	1.7	1.3	2.0	1.6	1.2	1.9	2.4	1.9	2.9	
Dorsopathies (M40-M54)										
2007.....	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.3	
2006.....	0.4	0.3	0.4	0.4	0.4	0.4	0.4	0.4	0.3	
2005.....	0.3	0.4	0.3	0.3	0.4	0.4	0.4	0.5	0.3	
2004.....	0.4	0.3	0.4	0.4	0.3	0.4	0.3	0.5	0.2	
2003.....	0.3	0.4	0.3	0.3	0.3	0.3	0.4	0.5	0.3	
2002.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.2	
2001.....	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.4	0.3	
2000.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.2	
1999.....	0.3	0.3	0.2	0.3	0.3	0.3	0.3	0.4	0.2	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Soft tissue disorders (M60-M79)										
2007.....	0.4	0.5	0.4	0.4	0.5	0.4	0.6	0.9	0.4	
2006.....	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.8	0.6	
2005.....	0.4	0.5	0.3	0.4	0.5	0.3	0.6	0.8	0.5	
2004.....	0.4	0.5	0.4	0.4	0.5	0.3	0.8	1.0	0.6	
2003.....	0.4	0.5	0.4	0.4	0.5	0.3	0.7	0.8	0.7	
2002.....	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.9	0.5	
2001.....	0.4	0.5	0.3	0.3	0.5	0.3	0.7	0.9	0.6	
2000.....	0.4	0.5	0.3	0.3	0.4	0.2	0.8	0.9	0.7	
1999.....	0.4	0.5	0.3	0.3	0.4	0.2	0.7	1.0	0.5	
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)										
2007.....	1.0	0.8	1.1	1.0	0.7	1.1	1.1	1.2	1.0	
2006.....	1.0	0.8	1.1	1.0	0.7	1.1	1.0	1.2	0.9	
2005.....	1.0	0.8	1.2	1.1	0.8	1.2	1.1	1.0	1.1	
2004.....	1.1	0.8	1.2	1.1	0.8	1.3	1.1	1.3	1.0	
2003.....	1.1	0.8	1.3	1.2	0.8	1.3	1.1	1.5	0.9	
2002.....	1.2	0.8	1.3	1.2	0.8	1.3	1.1	1.3	1.0	
2001.....	1.1	0.8	1.3	1.2	0.8	1.3	1.1	1.2	1.0	
2000.....	1.1	0.8	1.2	1.1	0.8	1.3	1.1	1.4	0.9	
1999.....	1.1	0.7	1.2	1.1	0.7	1.3	0.9	1.0	0.9	
Osteoporosis (M80-M81)										
2007.....	0.5	0.2	0.6	0.5	0.2	0.7	0.1	*	*	
2006.....	0.5	0.2	0.6	0.5	0.2	0.7	0.1	*	*	
2005.....	0.5	0.2	0.7	0.6	0.2	0.8	0.1	*	0.1	
2004.....	0.6	0.2	0.7	0.6	0.2	0.8	0.1	*	0.1	
2003.....	0.6	0.2	0.8	0.6	0.2	0.9	*	*	*	
2002.....	0.6	0.2	0.8	0.7	0.2	0.9	0.1	*	0.2	
2001.....	0.6	0.2	0.8	0.6	0.2	0.8	0.1	*	0.1	
2000.....	0.6	0.2	0.8	0.7	0.2	0.9	0.1	*	*	
1999.....	0.6	0.2	0.8	0.7	0.2	0.9	0.1	*	*	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)										
2007.....	0.5	0.6	0.4	0.5	0.6	0.4	1.0	1.2	0.9	
2006.....	0.5	0.6	0.5	0.5	0.6	0.4	0.9	1.2	0.8	
2005.....	0.5	0.6	0.4	0.5	0.6	0.4	1.0	1.0	0.9	
2004.....	0.5	0.6	0.5	0.5	0.6	0.5	1.0	1.2	0.9	
2003.....	0.6	0.6	0.5	0.5	0.6	0.5	1.1	1.4	0.8	
2002.....	0.5	0.6	0.4	0.5	0.6	0.4	1.0	1.3	0.8	
2001.....	0.5	0.6	0.5	0.5	0.6	0.4	1.0	1.2	0.9	
2000.....	0.5	0.6	0.4	0.4	0.5	0.4	1.0	1.3	0.8	
1999.....	0.4	0.5	0.4	0.4	0.5	0.4	0.8	0.9	0.7	
XIV. Diseases of the genitourinary system (N00-N98)										
2007.....	20.2	23.1	18.3	18.7	21.6	16.8	36.9	41.2	34.2	
2006.....	20.3	23.2	18.3	18.7	21.7	16.8	37.8	42.7	34.5	
2005.....	20.4	23.3	18.6	18.8	21.8	17.0	37.9	42.4	35.0	
2004.....	20.1	23.0	18.3	18.5	21.4	16.7	37.9	43.0	34.8	
2003.....	20.4	23.4	18.5	18.8	21.8	16.9	38.8	44.4	35.2	
2002.....	20.2	23.2	18.3	18.6	21.6	16.7	38.2	44.4	34.6	
2001.....	20.2	23.2	18.5	18.5	21.6	16.7	38.6	43.1	36.0	
2000.....	19.8	23.0	18.0	18.1	21.4	16.3	37.7	42.6	34.8	
1999.....	19.5	22.7	17.7	17.8	20.9	16.1	37.9	44.6	34.2	
Diseases of urinary system (N00-N39)										
2007.....	19.9	22.5	18.1	18.4	21.1	16.6	36.4	40.5	33.8	
2006.....	19.9	22.6	18.1	18.3	21.0	16.5	37.3	42.0	34.1	
2005.....	20.0	22.7	18.3	18.5	21.2	16.7	37.4	41.7	34.6	
2004.....	19.8	22.4	18.1	18.2	20.8	16.5	37.4	42.1	34.4	
2003.....	20.1	22.9	18.3	18.4	21.3	16.7	38.4	43.7	34.9	
2002.....	19.9	22.6	18.1	18.3	21.0	16.6	37.7	43.5	34.2	
2001.....	19.9	22.6	18.2	18.2	21.0	16.5	38.1	42.4	35.6	
2000.....	19.5	22.4	17.7	17.8	20.8	16.1	37.2	41.8	34.4	
1999.....	19.2	22.0	17.5	17.5	20.2	15.9	37.3	43.7	33.7	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Glomerular and renal tubulo-interstitial diseases (N00-N15)										
2007.....	1.3	1.6	1.1	1.2	1.5	1.1	2.3	2.8	2.0	
2006.....	1.0	1.1	0.9	0.9	1.0	0.8	1.6	2.1	1.4	
2005.....	0.7	0.8	0.7	0.7	0.8	0.6	1.1	1.3	1.0	
2004.....	0.6	0.7	0.6	0.6	0.7	0.6	0.9	1.0	0.8	
2003.....	0.7	0.7	0.6	0.6	0.7	0.6	0.9	1.1	0.8	
2002.....	0.7	0.8	0.6	0.6	0.7	0.6	0.8	1.0	0.7	
2001.....	0.7	0.8	0.6	0.6	0.7	0.6	0.8	1.0	0.7	
2000.....	0.7	0.8	0.7	0.6	0.7	0.6	1.1	1.2	1.0	
1999.....	0.7	0.9	0.7	0.7	0.9	0.7	1.1	1.5	0.9	
Glomerular diseases (N00-N07)										
2007.....	1.0	1.2	0.8	0.9	1.1	0.7	2.0	2.3	1.8	
2006.....	0.6	0.8	0.5	0.5	0.7	0.4	1.3	1.7	1.1	
2005.....	0.3	0.4	0.2	0.3	0.4	0.2	0.6	0.8	0.5	
2004.....	0.2	0.3	0.2	0.2	0.2	0.2	0.4	0.4	0.4	
2003.....	0.2	0.3	0.2	0.2	0.2	0.2	0.3	0.4	0.2	
2002.....	0.2	0.3	0.2	0.2	0.3	0.2	0.4	0.4	0.4	
2001.....	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.4	0.3	
2000.....	0.2	0.3	0.2	0.2	0.3	0.2	0.4	0.5	0.4	
1999.....	0.3	0.3	0.2	0.2	0.3	0.2	0.5	0.6	0.4	
Acute and rapidly progressive nephritic syndrome (N00-N01)										
2007.....	0.0	*	*	0.0	*	*	*	*	*	
2006.....	0.0	*	*	0.0	*	*	*	*	*	
2005.....	*	*	*	*	*	*	*	*	*	
2004.....	0.0	*	*	0.0	*	*	*	*	*	
2003.....	*	*	*	*	*	*	*	*	*	
2002.....	0.0	*	*	0.0	*	*	*	*	*	
2001.....	0.0	*	*	0.0	*	*	*	*	*	
2000.....	0.0	*	*	0.0	*	*	*	*	*	
1999.....	0.0	*	*	0.0	*	*	*	*	*	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Nephrotic syndrome (N04)										
2007.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	*	*	
2006.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2005.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2003.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*	
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2000.....	0.0	0.1	0.0	0.0	0.1	0.0	0.1	*	*	
1999.....	0.1	0.1	0.0	0.0	0.0	0.0	0.1	*	*	
Other and unspecified glomerular diseases (N02-N03,N05-N07)										
2007.....	0.9	1.1	0.8	0.8	1.0	0.7	1.9	2.1	1.7	
2006.....	0.6	0.7	0.5	0.5	0.6	0.4	1.2	1.6	1.1	
2005.....	0.3	0.3	0.2	0.2	0.3	0.2	0.6	0.7	0.5	
2004.....	0.2	0.2	0.2	0.2	0.2	0.1	0.3	0.3	0.3	
2003.....	0.2	0.2	0.1	0.1	0.2	0.1	0.3	0.3	0.2	
2002.....	0.2	0.2	0.1	0.2	0.2	0.1	0.3	0.3	0.2	
2001.....	0.2	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.2	
2000.....	0.2	0.2	0.2	0.2	0.2	0.1	0.3	0.4	0.3	
1999.....	0.2	0.2	0.1	0.2	0.2	0.1	0.4	0.4	0.3	
Renal tubulo-interstitial diseases (N10-N15)										
2007.....	0.4	0.4	0.3	0.4	0.4	0.3	0.3	0.5	0.3	
2006.....	0.4	0.4	0.3	0.4	0.4	0.4	0.3	0.4	0.3	
2005.....	0.4	0.4	0.4	0.4	0.4	0.4	0.5	0.6	0.5	
2004.....	0.4	0.5	0.4	0.4	0.4	0.4	0.5	0.6	0.4	
2003.....	0.4	0.5	0.4	0.4	0.5	0.4	0.6	0.6	0.6	
2002.....	0.4	0.5	0.4	0.4	0.5	0.4	0.4	0.5	0.4	
2001.....	0.4	0.5	0.4	0.4	0.4	0.4	0.5	0.6	0.5	
2000.....	0.5	0.5	0.5	0.4	0.4	0.4	0.6	0.7	0.6	
1999.....	0.5	0.6	0.5	0.5	0.6	0.5	0.6	0.9	0.5	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Renal failure (N17-N19)										
2007.....	13.6	16.5	11.6	12.3	15.3	10.3	27.4	31.4	24.9	
2006.....	13.9	17.0	11.8	12.4	15.6	10.4	28.9	33.3	26.0	
2005.....	14.0	17.1	12.0	12.6	15.8	10.5	29.0	33.3	26.3	
2004.....	13.9	17.0	12.0	12.5	15.6	10.6	29.1	33.8	26.3	
2003.....	14.2	17.5	12.1	12.7	16.1	10.7	30.0	35.6	26.5	
2002.....	13.9	17.3	11.9	12.5	15.8	10.5	29.3	34.8	25.9	
2001.....	13.7	17.1	11.8	12.3	15.8	10.2	29.2	33.4	26.7	
2000.....	13.2	16.6	11.3	11.8	15.2	9.8	28.2	32.6	25.5	
1999.....	12.8	16.1	10.8	11.3	14.6	9.4	27.9	33.0	24.9	
Urolithiasis (N20-N23)										
2007.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2006.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2005.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2004.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2003.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*	
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*	
Other and unspecified disorders of kidney (N25-N27)										
2007.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2006.....	0.0	0.0	0.0	0.0	0.0	*	*	*	*	
2005.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2003.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2001.....	0.0	*	0.0	0.0	*	0.0	*	*	*	
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Urinary tract infection, site not specified (N39.0)										
2007.....	4.5	3.8	4.9	4.4	3.7	4.8	5.9	5.4	6.2	
2006.....	4.5	3.8	4.9	4.5	3.8	4.9	5.9	5.7	6.1	
2005.....	4.8	4.1	5.2	4.7	3.9	5.2	6.5	6.0	6.6	
2004.....	4.6	4.0	5.0	4.5	3.8	4.9	6.5	6.4	6.5	
2003.....	4.7	3.9	5.1	4.6	3.9	4.9	6.7	6.1	7.0	
2002.....	4.7	4.0	5.2	4.6	3.8	5.0	6.8	6.7	6.9	
2001.....	5.0	4.1	5.4	4.8	4.0	5.2	7.3	7.0	7.4	
2000.....	5.0	4.4	5.3	4.9	4.2	5.2	7.2	7.0	7.1	
1999.....	5.2	4.4	5.5	5.0	4.2	5.4	7.5	8.0	7.2	
Other diseases of urinary system (N28-N36,N39.1-N39.9)										
2007.....	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.8	0.6	
2006.....	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.9	0.5	
2005.....	0.4	0.5	0.3	0.4	0.5	0.3	0.6	0.8	0.5	
2004.....	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.8	0.7	
2003.....	0.4	0.5	0.3	0.4	0.4	0.3	0.6	0.8	0.5	
2002.....	0.4	0.5	0.3	0.4	0.4	0.3	0.7	0.8	0.6	
2001.....	0.4	0.4	0.3	0.4	0.4	0.3	0.7	0.8	0.6	
2000.....	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.9	0.6	
1999.....	0.4	0.5	0.3	0.3	0.5	0.3	0.7	0.9	0.5	
Diseases of male genital organs (N40-N50)										
2007.....	0.2	0.5	...	0.2	0.5	...	0.2	0.7	...	
2006.....	0.2	0.6	...	0.2	0.6	...	0.3	0.7	...	
2005.....	0.2	0.6	...	0.2	0.6	...	0.2	0.7	...	
2004.....	0.2	0.6	...	0.2	0.6	...	0.3	0.8	...	
2003.....	0.2	0.6	...	0.2	0.6	...	0.2	0.6	...	
2002.....	0.2	0.6	...	0.2	0.5	...	0.3	0.8	...	
2001.....	0.2	0.6	...	0.2	0.6	...	0.3	0.8	...	
2000.....	0.2	0.6	...	0.2	0.6	...	0.3	0.8	...	
1999.....	0.2	0.6	...	0.2	0.6	...	0.3	0.9	...	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Hyperplasia of prostate (N40)									
2007.....	0.1	0.4	...	0.1	0.4	...	0.2	0.5	...
2006.....	0.2	0.5	...	0.2	0.5	...	0.2	0.4	...
2005.....	0.2	0.5	...	0.2	0.5	...	0.1	0.4	...
2004.....	0.2	0.5	...	0.2	0.5	...	0.1	0.4	...
2003.....	0.2	0.5	...	0.2	0.5	...	0.1	0.4	...
2002.....	0.1	0.4	...	0.1	0.4	...	0.2	0.5	...
2001.....	0.1	0.5	...	0.2	0.5	...	0.1	0.4	...
2000.....	0.2	0.5	...	0.2	0.5	...	0.2	0.5	...
1999.....	0.2	0.5	...	0.2	0.5	...	0.2	0.5	...
Other diseases of male genital organs (N41-N50)									
2007.....	0.1	0.1	...	0.1	0.1	...	0.1	0.3	...
2006.....	0.1	0.2	...	0.1	0.2	...	0.1	0.3	...
2005.....	0.1	0.2	...	0.1	0.2	...	0.1	0.3	...
2004.....	0.1	0.2	...	0.1	0.1	...	0.1	0.4	...
2003.....	0.0	0.1	...	0.0	0.1	...	0.1	0.2	...
2002.....	0.1	0.1	...	0.0	0.1	...	0.1	0.3	...
2001.....	0.1	0.1	...	0.0	0.1	...	0.1	0.3	...
2000.....	0.0	0.1	...	0.0	0.1	...	0.1	0.3	...
1999.....	0.1	0.2	...	0.1	0.1	...	0.1	0.4	...
Disorders of breast (N60-N64)									
2007.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2006.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2005.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2004.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2003.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2002.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2001.....	0.0	*	0.0	0.0	*	0.0	*	*	*
2000.....	0.0	*	0.0	0.0	*	0.0	*	*	*
1999.....	0.0	*	0.0	0.0	*	0.0	*	*	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Disorders of female genital organs (N70-N98)										
2007.....	0.1	...	0.2	0.1	...	0.2	0.2	...	0.4	
2006.....	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3	
2005.....	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3	
2004.....	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3	
2003.....	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3	
2002.....	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3	
2001.....	0.1	...	0.2	0.1	...	0.2	0.2	...	0.4	
2000.....	0.1	...	0.2	0.1	...	0.2	0.2	...	0.3	
1999.....	0.1	...	0.2	0.1	...	0.2	0.3	...	0.4	
Inflammatory diseases of female pelvic organs (N70-N76)										
2007.....	0.0	...	0.1	0.0	...	0.1	0.0	...	0.1	
2006.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1	
2005.....	0.0	...	0.1	0.0	...	0.1	*	...	*	
2004.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.1	
2003.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1	
2002.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.1	
2001.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1	
2000.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1	
1999.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2	
Noninflammatory disorders of female genital tract (N80-N98)										
2007.....	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2	
2006.....	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2	
2005.....	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2	
2004.....	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2	
2003.....	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2	
2002.....	0.1	...	0.1	0.0	...	0.1	0.1	...	0.2	
2001.....	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2	
2000.....	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2	
1999.....	0.1	...	0.1	0.1	...	0.1	0.1	...	0.3	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
XV. Pregnancy, childbirth and the puerperium (O00-O99)										
2007.....	0.3	...	0.5	0.2	...	0.4	0.6	...	1.2	
2006.....	0.2	...	0.5	0.2	...	0.4	0.6	...	1.2	
2005.....	0.3	...	0.5	0.2	...	0.4	0.7	...	1.3	
2004.....	0.2	...	0.5	0.2	...	0.4	0.6	...	1.2	
2003.....	0.2	...	0.4	0.1	...	0.3	0.5	...	0.9	
2002.....	0.1	...	0.3	0.1	...	0.2	0.4	...	0.8	
2001.....	0.1	...	0.3	0.1	...	0.2	0.4	...	0.8	
2000.....	0.1	...	0.3	0.1	...	0.2	0.4	...	0.7	
1999.....	0.2	...	0.3	0.1	...	0.2	0.4	...	0.8	
Pregnancy with abortive outcome (O00-O07)										
2007.....	0.0	...	0.0	*	...	*	*	...	*	
2006.....	0.0	...	0.0	*	...	*	*	...	*	
2005.....	0.0	...	0.0	*	...	*	*	...	*	
2004.....	0.0	...	0.0	*	...	*	*	...	*	
2003.....	0.0	...	0.0	*	...	*	*	...	*	
2002.....	0.0	...	0.0	*	...	*	*	...	*	
2001.....	0.0	...	0.0	0.0	...	0.0	*	...	*	
2000.....	0.0	...	0.0	*	...	*	*	...	*	
1999.....	0.0	...	0.0	*	...	*	*	...	*	
Ectopic pregnancy (O00)										
2007.....	*	...	*	*	...	*	*	...	*	
2006.....	*	...	*	*	...	*	*	...	*	
2005.....	*	...	*	*	...	*	*	...	*	
2004.....	*	...	*	*	...	*	*	...	*	
2003.....	0.0	...	0.0	*	...	*	*	...	*	
2002.....	*	...	*	*	...	*	*	...	*	
2001.....	0.0	...	0.0	*	...	*	*	...	*	
2000.....	0.0	...	0.0	*	...	*	*	...	*	
1999.....	*	...	*	*	...	*	*	...	*	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Spontaneous abortion (O03)										
2007.....	*	...	*	*	...	*	*	...	*	
2006.....	*	...	*	*	...	*	*	...	*	
2005.....	*	...	*	*	...	*	*	...	*	
2004.....	*	...	*	*	...	*	*	...	*	
2003.....	*	...	*	*	...	*	*	...	*	
2002.....	*	...	*	*	...	*	*	...	*	
2001.....	*	...	*	*	...	*	*	...	*	
2000.....	*	...	*	*	...	*	*	...	*	
1999.....	*	...	*	*	...	*	*	...	*	
Medical abortion (O04)										
2007.....	*	...	*	*	...	*	*	...	*	
2006.....	*	...	*	*	...	*	*	...	*	
2005.....	*	...	*	*	...	*	*	...	*	
2004.....	*	...	*	*	...	*	*	...	*	
2003.....	*	...	*	*	...	*	*	...	*	
2002.....	*	...	*	*	...	*	*	...	*	
2001.....	*	...	*	*	...	*	*	...	*	
2000.....	*	...	*	*	...	*	*	...	*	
1999.....	*	...	*	*	...	*	*	...	*	
Other abortion (O05)										
2007.....	*	...	*	*	...	*	*	...	*	
2006.....	*	...	*	*	...	*	*	...	*	
2005.....	*	...	*	*	...	*	*	...	*	
2004.....	*	...	*	*	...	*	*	...	*	
2003.....	*	...	*	*	...	*	*	...	*	
2002.....	*	...	*	*	...	*	*	...	*	
2001.....	*	...	*	*	...	*	*	...	*	
2000.....	*	...	*	*	...	*	*	...	*	
1999.....	*	...	*	*	...	*	*	...	*	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/		White 2/		Black 2/				
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other and unspecified pregnancy with abortive outcome (O01-O02,006-O07)									
2007.....	*	...	*	*	...	*	*	...	*
2006.....	*	...	*	*	...	*	*	...	*
2005.....	*	...	*	*	...	*	*	...	*
2004.....	*	...	*	*	...	*	*	...	*
2003.....	*	...	*	*	...	*	*	...	*
2002.....	*	...	*	*	...	*	*	...	*
2001.....	*	...	*	*	...	*	*	...	*
2000.....	*	...	*	*	...	*	*	...	*
1999.....	*	...	*	*	...	*	*	...	*
Other direct obstetric deaths (O10-O92)									
2007.....	0.1	...	0.3	0.1	...	0.2	0.3	...	0.5
2006.....	0.1	...	0.3	0.1	...	0.2	0.4	...	0.8
2005.....	0.1	...	0.3	0.1	...	0.2	0.4	...	0.7
2004.....	0.1	...	0.2	0.1	...	0.2	0.3	...	0.7
2003.....	0.1	...	0.2	0.1	...	0.2	0.3	...	0.7
2002.....	0.1	...	0.2	0.1	...	0.1	0.3	...	0.6
2001.....	0.1	...	0.2	0.1	...	0.2	0.3	...	0.6
2000.....	0.1	...	0.2	0.1	...	0.2	0.2	...	0.5
1999.....	0.1	...	0.2	0.1	...	0.1	0.3	...	0.6
Eclampsia and pre-eclampsia (O11,O13-O16)									
2007.....	0.0	...	0.0	0.0	...	0.0	*	...	*
2006.....	0.0	...	0.0	0.0	...	0.0	0.1	...	0.1
2005.....	0.0	...	0.0	0.0	...	0.0	0.0	...	0.1
2004.....	0.0	...	0.0	0.0	...	0.0	0.1	...	0.1
2003.....	0.0	...	0.0	0.0	...	0.0	0.1	...	0.1
2002.....	0.0	...	0.0	0.0	...	0.0	0.1	...	0.1
2001.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.1
2000.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1
1999.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,072)									
2007.....	0.0	...	0.0	0.0	...	0.0	*	...	*
2006.....	0.0	...	0.0	0.0	...	0.0	*	...	*
2005.....	0.0	...	0.0	0.0	...	0.0	*	...	*
2004.....	0.0	...	0.0	0.0	...	0.0	*	...	*
2003.....	0.0	...	0.0	0.0	...	0.0	*	...	*
2002.....	0.0	...	0.0	*	...	*	*	...	*
2001.....	0.0	...	0.0	0.0	...	0.0	*	...	*
2000.....	0.0	...	0.0	0.0	...	0.0	*	...	*
1999.....	0.0	...	0.0	0.0	...	0.0	*	...	*
Complications predominately related to the puerperium (O85-O92)									
2007.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.1
2006.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
2005.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
2004.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.2
2003.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
2002.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
2001.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.2
2000.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.1
1999.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2
Obstetric embolism (O88)									
2007.....	0.0	...	0.0	*	...	*	*	...	*
2006.....	0.0	...	0.0	0.0	...	0.0	*	...	*
2005.....	0.0	...	0.0	0.0	...	0.0	*	...	*
2004.....	0.0	...	0.0	0.0	...	0.0	*	...	*
2003.....	0.0	...	0.0	0.0	...	0.0	*	...	*
2002.....	0.0	...	0.0	0.0	...	0.0	*	...	*
2001.....	0.0	...	0.0	0.0	...	0.0	0.1	...	0.1
2000.....	0.0	...	0.0	0.0	...	0.0	*	...	*
1999.....	0.0	...	0.0	0.0	...	0.0	*	...	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Other complications predominately related to the puerperium (O85-O87,O89-O92)										
2007.....	0.0	...	0.0	0.0	...	0.0	0.1	...	0.1	
2006.....	0.0	...	0.0	0.0	...	0.0	0.1	...	0.2	
2005.....	0.0	...	0.0	0.0	...	0.0	0.1	...	0.1	
2004.....	0.0	...	0.0	0.0	...	0.0	0.1	...	0.1	
2003.....	0.0	...	0.0	0.0	...	0.0	*	...	*	
2002.....	0.0	...	0.0	*	...	*	0.1	...	0.1	
2001.....	0.0	...	0.0	*	...	*	*	...	*	
2000.....	0.0	...	0.0	*	...	*	*	...	*	
1999.....	0.0	...	0.0	*	...	*	*	...	*	
All other direct obstetric causes (O10,O12,O21-O43,O47-O66,O68-O71,O73-O75)										
2007.....	0.1	...	0.1	0.0	...	0.1	0.1	...	0.3	
2006.....	0.1	...	0.1	0.0	...	0.1	0.2	...	0.4	
2005.....	0.1	...	0.1	0.0	...	0.1	0.2	...	0.4	
2004.....	0.1	...	0.1	0.0	...	0.1	0.1	...	0.3	
2003.....	0.0	...	0.1	0.0	...	0.1	0.2	...	0.3	
2002.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.2	
2001.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2	
2000.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.2	
1999.....	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2	
Obstetric death of unspecified cause (O95)										
2007.....	0.0	...	0.0	*	...	*	*	...	*	
2006.....	0.0	...	0.0	*	...	*	*	...	*	
2005.....	0.0	...	0.0	0.0	...	0.0	*	...	*	
2004.....	0.0	...	0.0	*	...	*	*	...	*	
2003.....	0.0	...	0.0	*	...	*	*	...	*	
2002.....	*	...	*	*	...	*	*	...	*	
2001.....	*	...	*	*	...	*	*	...	*	
2000.....	*	...	*	*	...	*	*	...	*	
1999.....	*	...	*	*	...	*	*	...	*	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Other deaths related to pregnancy, childbirth and the puerperium (O96-O97)										
2007.....	0.1	...	0.2	0.1	...	0.1	0.2	...	0.3	
2006.....	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2	
2005.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.2	
2004.....	0.1	...	0.1	0.1	...	0.1	0.1	...	0.2	
2003.....	0.0	...	0.0	0.0	...	0.0	*	...	*	
2002.....	0.0	...	0.0	*	...	*	*	...	*	
2001.....	*	...	*	*	...	*	*	...	*	
2000.....	*	...	*	*	...	*	*	...	*	
1999.....	*	...	*	*	...	*	*	...	*	
Indirect obstetric deaths (O98-O99)										
2007.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.2	
2006.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.2	
2005.....	0.1	...	0.1	0.0	...	0.1	0.1	...	0.3	
2004.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.2	
2003.....	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1	
2002.....	0.0	...	0.0	0.0	...	0.0	*	...	*	
2001.....	0.0	...	0.0	0.0	...	0.0	0.0	...	0.1	
2000.....	0.0	...	0.0	0.0	...	0.0	*	...	*	
1999.....	0.0	...	0.0	0.0	...	0.0	*	...	*	
XVI. Certain conditions originating in the perinatal period (P00-P96)										
2007.....	4.7	5.2	4.2	3.7	4.1	3.3	9.8	10.8	8.7	
2006.....	4.8	5.3	4.3	3.8	4.2	3.3	10.4	11.2	9.6	
2005.....	4.9	5.4	4.4	3.8	4.2	3.4	10.5	11.5	9.5	
2004.....	4.8	5.4	4.2	3.8	4.3	3.3	10.3	11.3	9.2	
2003.....	4.9	5.5	4.3	4.0	4.4	3.5	10.2	11.4	9.0	
2002.....	4.9	5.4	4.3	3.9	4.3	3.4	10.1	10.9	9.2	
2001.....	4.7	5.3	4.2	3.7	4.1	3.3	10.3	11.3	9.1	
2000.....	5.1	5.6	4.5	4.0	4.4	3.5	11.3	12.6	9.9	
1999.....	5.2	5.7	4.6	4.0	4.4	3.6	11.8	13.1	10.4	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Intrauterine hypoxia and birth asphyxia (P20-P21)										
2007.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2	
2006.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.1	
2005.....	0.2	0.2	0.2	0.2	0.2	0.1	0.3	0.3	0.2	
2004.....	0.2	0.2	0.2	0.2	0.2	0.1	0.3	0.3	0.3	
2003.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3	
2002.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3	
2001.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3	
2000.....	0.2	0.3	0.2	0.2	0.2	0.2	0.4	0.4	0.3	
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.4	0.4	0.3	
Respiratory distress of newborn (P22)										
2007.....	0.3	0.3	0.2	0.2	0.2	0.2	0.5	0.7	0.4	
2006.....	0.3	0.3	0.2	0.2	0.3	0.2	0.6	0.6	0.6	
2005.....	0.3	0.3	0.2	0.2	0.3	0.2	0.6	0.7	0.5	
2004.....	0.3	0.3	0.3	0.2	0.3	0.2	0.6	0.7	0.5	
2003.....	0.3	0.3	0.2	0.2	0.3	0.2	0.5	0.6	0.5	
2002.....	0.3	0.4	0.3	0.3	0.3	0.2	0.7	0.7	0.7	
2001.....	0.3	0.4	0.3	0.3	0.3	0.3	0.7	0.9	0.6	
2000.....	0.4	0.5	0.3	0.3	0.4	0.2	0.8	0.9	0.6	
1999.....	0.4	0.5	0.3	0.3	0.4	0.3	0.9	1.0	0.7	
Other respiratory conditions originating in the perinatal period (P23-P28)										
2007.....	0.4	0.4	0.3	0.3	0.3	0.3	0.7	0.8	0.7	
2006.....	0.4	0.5	0.4	0.3	0.4	0.3	0.8	0.9	0.7	
2005.....	0.4	0.5	0.3	0.3	0.4	0.3	0.8	0.9	0.8	
2004.....	0.4	0.5	0.4	0.3	0.4	0.3	0.9	0.9	0.8	
2003.....	0.4	0.5	0.3	0.4	0.4	0.3	0.8	1.0	0.7	
2002.....	0.4	0.5	0.4	0.4	0.4	0.3	0.9	1.0	0.8	
2001.....	0.5	0.6	0.5	0.4	0.5	0.4	1.1	1.2	1.0	
2000.....	0.6	0.6	0.5	0.5	0.5	0.4	1.2	1.4	1.0	
1999.....	0.6	0.7	0.5	0.5	0.6	0.4	1.2	1.4	1.1	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/		Black 2/			
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Infections specific to the perinatal period (P35-P39)									
2007.....	0.3	0.4	0.3	0.3	0.3	0.3	0.7	0.7	0.6
2006.....	0.3	0.4	0.3	0.3	0.3	0.3	0.7	0.8	0.6
2005.....	0.3	0.4	0.3	0.3	0.3	0.2	0.7	0.8	0.6
2004.....	0.4	0.4	0.3	0.3	0.3	0.2	0.7	0.8	0.6
2003.....	0.3	0.4	0.3	0.3	0.3	0.2	0.7	0.8	0.6
2002.....	0.3	0.4	0.3	0.3	0.3	0.2	0.6	0.6	0.6
2001.....	0.3	0.3	0.3	0.2	0.3	0.2	0.6	0.6	0.6
2000.....	0.3	0.4	0.3	0.3	0.3	0.2	0.7	0.8	0.6
1999.....	0.3	0.4	0.3	0.2	0.3	0.2	0.7	0.8	0.6
Other and unspecified conditions originating in the perinatal period (P05,P08,P29,P50-P96)									
2007.....	1.0	1.1	0.9	0.8	0.9	0.7	1.7	1.9	1.5
2006.....	1.0	1.1	0.9	0.8	0.9	0.7	1.9	2.1	1.7
2005.....	1.0	1.1	0.8	0.8	0.9	0.7	1.8	2.1	1.5
2004.....	0.9	1.0	0.8	0.7	0.9	0.6	1.7	1.9	1.5
2003.....	0.9	1.0	0.8	0.8	0.9	0.7	1.7	1.9	1.4
2002.....	0.8	0.9	0.7	0.7	0.8	0.6	1.4	1.6	1.3
2001.....	0.8	1.0	0.7	0.7	0.8	0.6	1.5	1.8	1.3
2000.....	1.0	1.1	0.8	0.8	0.9	0.7	1.9	2.3	1.6
1999.....	1.0	1.1	0.8	0.8	0.9	0.7	2.0	2.2	1.9
XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)									
2007.....	3.4	3.6	3.2	3.3	3.5	3.1	3.9	4.0	3.8
2006.....	3.5	3.7	3.2	3.4	3.7	3.2	4.0	4.4	3.8
2005.....	3.5	3.7	3.3	3.5	3.6	3.3	3.9	4.1	3.7
2004.....	3.6	3.7	3.4	3.5	3.7	3.4	4.0	4.4	3.7
2003.....	3.6	3.8	3.4	3.6	3.7	3.4	4.0	4.4	3.6
2002.....	3.7	3.8	3.5	3.7	3.8	3.5	4.1	4.4	3.9
2001.....	3.6	3.8	3.4	3.6	3.8	3.4	4.1	4.3	3.9
2000.....	3.8	4.1	3.6	3.7	4.0	3.5	4.4	4.8	4.1
1999.....	3.8	3.9	3.6	3.7	3.9	3.6	4.3	4.7	4.1

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Anencephaly and similar malformations (Q00)										
2007.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
2006.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
2005.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
2004.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2003.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
Spina bifida (Q05)										
2007.....	0.0	0.0	0.0	0.0	0.1	0.0	*	*	*	*
2006.....	0.1	0.1	0.1	0.1	0.1	0.1	0.0	*	*	*
2005.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2004.....	0.0	0.0	0.0	0.1	0.0	0.1	*	*	*	*
2003.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*	*
2001.....	0.0	0.0	0.1	0.0	0.0	0.1	*	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
1999.....	0.0	0.0	0.0	0.1	0.0	0.1	*	*	*	*
All other congenital malformations of nervous system (Q01-Q04,Q06-Q07)										
2007.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4
2006.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.4	0.4
2005.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.3	0.3
2004.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.4	0.4
2003.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.3
2002.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
2001.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.4
2000.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.4	0.3
1999.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.3	0.4

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Congenital malformations of heart (Q20-Q24)										
2007.....	0.9	1.1	0.8	0.9	1.0	0.8	1.1	1.2	1.1	
2006.....	0.9	1.0	0.8	0.9	1.0	0.8	1.2	1.3	1.1	
2005.....	1.0	1.0	0.9	1.0	1.1	0.9	1.1	1.1	1.1	
2004.....	1.0	1.1	0.9	1.0	1.1	0.9	1.2	1.3	1.0	
2003.....	1.1	1.2	1.0	1.1	1.2	1.0	1.2	1.3	1.1	
2002.....	1.1	1.2	1.0	1.1	1.2	1.0	1.2	1.3	1.2	
2001.....	1.1	1.2	1.0	1.1	1.2	1.0	1.4	1.4	1.3	
2000.....	1.2	1.3	1.2	1.2	1.3	1.1	1.4	1.6	1.3	
1999.....	1.3	1.4	1.2	1.3	1.4	1.2	1.5	1.7	1.3	
Other congenital malformations of circulatory system (Q25-Q28)										
2007.....	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.3	0.3	
2006.....	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.4	0.2	
2005.....	0.2	0.3	0.2	0.2	0.2	0.2	0.3	0.3	0.3	
2004.....	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.5	0.3	
2003.....	0.3	0.3	0.3	0.3	0.3	0.3	0.4	0.4	0.3	
2002.....	0.3	0.4	0.3	0.3	0.3	0.3	0.5	0.6	0.4	
2001.....	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.4	0.3	
2000.....	0.3	0.4	0.3	0.3	0.4	0.3	0.4	0.5	0.4	
1999.....	0.3	0.4	0.3	0.3	0.3	0.3	0.4	0.5	0.4	
Congenital malformations of respiratory system (Q30-Q34)										
2007.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.2	
2006.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.1	
2005.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.2	
2004.....	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.4	0.3	
2003.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.2	
2002.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3	
2001.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.2	
2000.....	0.3	0.3	0.2	0.2	0.3	0.2	0.4	0.4	0.3	
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.2	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Other congenital malformations and deformations (Q10-Q18,Q35-Q89)										
2007.....	1.0	1.1	0.9	1.0	1.1	0.9	1.2	1.3	1.0	
2006.....	1.0	1.1	0.9	1.0	1.1	0.9	1.1	1.3	1.0	
2005.....	0.9	1.1	0.8	0.9	1.0	0.8	1.1	1.3	0.9	
2004.....	0.9	1.0	0.8	0.9	1.0	0.8	1.0	1.2	0.8	
2003.....	0.9	1.0	0.8	0.9	1.0	0.8	1.1	1.3	0.9	
2002.....	0.9	1.0	0.8	0.9	1.0	0.8	1.0	1.2	0.9	
2001.....	0.9	1.0	0.8	0.9	1.0	0.8	1.0	1.2	0.9	
2000.....	0.9	1.1	0.8	0.9	1.0	0.8	1.1	1.2	0.9	
1999.....	0.9	1.1	0.9	0.9	1.0	0.8	1.1	1.2	0.9	
Down's syndrome (Q90)										
2007.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	
2006.....	0.3	0.3	0.2	0.3	0.3	0.2	0.2	0.2	0.2	
2005.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.3	
2004.....	0.3	0.3	0.2	0.3	0.3	0.3	0.2	0.3	0.2	
2003.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.2	
2002.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.2	
2001.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	
2000.....	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.2	0.2	
1999.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	
All other chromosomal abnormalities, not elsewhere classified (Q91-Q99)										
2007.....	0.4	0.3	0.4	0.4	0.3	0.4	0.4	0.3	0.5	
2006.....	0.4	0.3	0.4	0.4	0.3	0.4	0.5	0.4	0.6	
2005.....	0.3	0.3	0.4	0.3	0.3	0.4	0.4	0.3	0.5	
2004.....	0.4	0.3	0.4	0.3	0.3	0.4	0.4	0.4	0.5	
2003.....	0.4	0.3	0.4	0.4	0.3	0.4	0.4	0.3	0.5	
2002.....	0.4	0.3	0.4	0.4	0.3	0.4	0.4	0.3	0.5	
2001.....	0.3	0.3	0.4	0.3	0.3	0.4	0.4	0.3	0.4	
2000.....	0.4	0.3	0.4	0.3	0.3	0.4	0.4	0.4	0.5	
1999.....	0.3	0.3	0.4	0.3	0.3	0.4	0.4	0.3	0.5	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)										
2007.....	10.4	10.5	9.8	10.0	10.2	9.5	13.9	15.2	12.4	
2006.....	10.1	10.5	9.3	9.8	10.1	9.1	13.3	15.4	11.6	
2005.....	10.4	11.1	9.4	10.0	10.6	9.1	14.6	17.1	12.5	
2004.....	10.0	10.7	9.0	9.6	10.2	8.7	14.0	16.0	12.1	
2003.....	10.6	11.4	9.5	10.2	11.0	9.2	14.8	17.4	12.5	
2002.....	10.3	11.4	9.0	9.9	10.8	8.7	14.5	18.0	11.8	
2001.....	11.3	12.7	9.8	11.0	12.2	9.5	15.1	17.7	12.8	
2000.....	11.4	13.1	9.7	11.0	12.6	9.3	16.5	19.7	13.6	
1999.....	9.7	11.0	8.4	9.1	10.1	7.9	15.4	19.1	12.4	
Senility (R54)										
2007.....	1.5	1.2	1.7	1.6	1.3	1.7	1.3	1.0	1.5	
2006.....	1.5	1.3	1.6	1.6	1.3	1.7	1.3	1.0	1.4	
2005.....	1.5	1.2	1.6	1.5	1.2	1.6	1.4	1.2	1.5	
2004.....	1.6	1.3	1.7	1.6	1.3	1.7	1.3	1.1	1.4	
2003.....	1.6	1.2	1.7	1.6	1.3	1.8	1.1	0.9	1.2	
2002.....	1.5	1.2	1.6	1.5	1.3	1.6	1.2	1.1	1.2	
2001.....	1.5	1.2	1.6	1.5	1.3	1.6	1.3	1.3	1.3	
2000.....	1.4	1.1	1.5	1.4	1.2	1.5	1.4	1.0	1.5	
1999.....	1.2	1.0	1.2	1.2	1.0	1.3	1.0	1.1	1.0	
Ill-defined and unknown causes of mortality (R95-R99)										
2007.....	4.4	5.3	3.6	4.2	5.0	3.4	6.3	7.8	5.0	
2006.....	4.6	5.5	3.7	4.5	5.3	3.6	6.1	7.6	4.8	
2005.....	5.6	6.7	4.4	5.3	6.4	4.2	8.5	10.6	6.8	
2004.....	5.2	6.3	4.1	4.9	6.0	3.9	7.7	9.4	6.2	
2003.....	5.9	7.2	4.7	5.6	6.8	4.4	8.7	11.2	6.8	
2002.....	5.9	7.2	4.6	5.6	6.8	4.4	8.9	11.7	6.7	
2001.....	7.1	8.7	5.6	6.8	8.4	5.3	9.5	11.6	7.7	
2000.....	7.4	9.1	5.8	7.0	8.7	5.4	10.9	13.9	8.4	
1999.....	6.0	7.3	4.8	5.5	6.6	4.4	10.1	13.3	7.6	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Sudden infant death syndrome (R95)									
2007.....	0.8	0.9	0.7	0.7	0.8	0.6	1.4	1.6	1.3
2006.....	0.8	0.9	0.7	0.7	0.8	0.5	1.5	1.6	1.4
2005.....	0.8	0.9	0.6	0.6	0.8	0.5	1.4	1.5	1.3
2004.....	0.8	0.9	0.6	0.6	0.7	0.5	1.5	1.8	1.2
2003.....	0.7	0.9	0.6	0.6	0.7	0.5	1.4	1.5	1.4
2002.....	0.8	0.9	0.7	0.7	0.8	0.5	1.4	1.6	1.3
2001.....	0.8	0.9	0.6	0.6	0.7	0.5	1.6	1.7	1.4
2000.....	0.9	1.1	0.8	0.7	0.9	0.6	1.9	2.1	1.6
1999.....	1.0	1.1	0.8	0.8	0.9	0.7	1.9	2.1	1.7
Other ill-defined and unknown causes of mortality (R96-R99)									
2007.....	3.6	4.4	2.9	3.5	4.2	2.9	4.8	6.2	3.7
2006.....	3.8	4.7	3.0	3.8	4.6	3.0	4.6	6.1	3.4
2005.....	4.9	5.8	3.9	4.7	5.6	3.7	7.1	9.1	5.5
2004.....	4.4	5.4	3.5	4.3	5.2	3.3	6.2	7.6	5.0
2003.....	5.2	6.3	4.0	5.0	6.1	4.0	7.3	9.6	5.5
2002.....	5.1	6.3	4.0	4.9	6.0	3.8	7.5	10.0	5.4
2001.....	6.3	7.8	4.9	6.2	7.6	4.8	7.9	9.8	6.3
2000.....	6.5	8.1	5.0	6.3	7.8	4.8	9.0	11.8	6.8
1999.....	5.1	6.2	4.0	4.7	5.7	3.7	8.2	11.1	6.0
All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)									
2007.....	4.4	4.0	4.5	4.2	3.9	4.4	6.3	6.4	6.1
2006.....	3.9	3.7	3.9	3.8	3.5	3.8	5.9	6.7	5.3
2005.....	3.3	3.1	3.3	3.2	3.0	3.3	4.7	5.3	4.2
2004.....	3.2	3.1	3.2	3.1	2.9	3.2	5.0	5.4	4.5
2003.....	3.2	3.0	3.1	3.0	2.9	3.0	4.9	5.3	4.5
2002.....	2.9	3.0	2.8	2.8	2.8	2.7	4.5	5.3	3.9
2001.....	2.8	2.8	2.7	2.6	2.6	2.6	4.3	4.8	3.8
2000.....	2.7	2.8	2.5	2.6	2.7	2.4	4.2	4.8	3.7
1999.....	2.6	2.7	2.4	2.4	2.5	2.3	4.2	4.8	3.8

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/		White 2/		Black 2/				
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
XX. External causes of mortality (*U01-*U03,V01-Y89)									
2007.....	60.1	86.6	35.1	60.3	85.7	36.2	66.4	105.8	31.8
2006.....	59.6	86.2	34.6	59.5	84.6	35.5	68.5	109.7	32.5
2005.....	58.7	85.0	34.0	58.3	83.3	34.7	68.7	109.3	33.3
2004.....	57.3	82.7	33.7	57.2	81.5	34.3	65.5	103.8	32.5
2003.....	57.0	82.9	33.0	56.5	81.2	33.4	66.0	104.9	32.4
2002.....	56.7	82.7	32.6	55.9	80.7	32.9	67.2	107.7	32.6
2001 3/.....	56.1	82.6	31.8	55.2	80.3	31.8	68.2	108.4	33.7
2000.....	53.9	79.5	30.5	52.5	76.6	30.4	67.9	109.1	32.9
1999.....	54.3	80.1	31.0	52.7	76.9	30.7	69.8	111.7	34.6
Accidents (unintentional injuries) (V01-X59,Y85-Y86)									
2007.....	40.0	55.2	25.8	41.5	56.8	26.9	36.6	54.8	21.6
2006.....	39.8	55.2	25.5	41.0	56.4	26.4	38.3	57.5	22.2
2005.....	39.1	54.2	25.0	40.1	55.2	25.8	38.7	58.1	22.8
2004.....	37.7	52.1	24.5	38.8	53.2	25.3	36.3	54.4	21.7
2003.....	37.3	51.8	24.1	38.2	52.8	24.7	36.1	53.9	21.6
2002.....	36.9	51.5	23.5	37.5	52.0	24.0	36.9	56.2	21.3
2001.....	35.7	50.2	22.5	36.0	50.3	22.8	37.6	56.9	22.0
2000.....	34.9	49.3	22.0	35.1	49.2	22.2	37.7	58.1	21.3
1999.....	35.3	49.8	22.3	35.2	49.3	22.4	40.1	61.3	23.1
Transport accidents (V01-V99,Y85)									
2007.....	15.3	22.4	8.6	15.8	22.9	8.9	15.0	24.1	7.3
2006.....	16.0	23.0	9.2	16.4	23.5	9.5	15.5	24.2	8.1
2005.....	16.2	23.4	9.3	16.7	23.9	9.7	15.4	24.1	8.0
2004.....	16.2	23.1	9.7	16.7	23.6	10.0	15.7	24.4	8.4
2003.....	16.4	23.4	9.8	16.8	23.9	10.0	15.8	24.5	8.6
2002.....	16.7	23.9	9.9	17.1	24.2	10.2	16.0	25.0	8.5
2001.....	16.6	23.7	9.8	16.8	24.0	10.0	16.5	25.5	8.9
2000.....	16.6	23.7	10.0	16.8	23.8	10.2	16.7	26.1	8.7
1999.....	16.6	23.6	10.2	16.8	23.6	10.4	17.1	26.5	9.5

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Railway accidents (V05,V15,V80.6,V81.2-V81.9)									
2007.....	0.1	0.2	0.0	0.1	0.2	0.0	0.2	0.4	*
2006.....	0.1	0.3	0.0	0.1	0.2	0.0	0.2	0.4	*
2005.....	0.2	0.2	0.1	0.1	0.2	0.0	0.2	0.4	*
2004.....	0.1	0.3	0.1	0.1	0.3	0.0	0.2	0.4	*
2003.....	0.1	0.2	0.0	0.1	0.3	0.0	0.2	0.4	*
2002.....	0.2	0.3	0.0	0.2	0.3	0.0	0.3	0.5	*
2001.....	0.2	0.3	0.1	0.2	0.3	0.0	0.2	0.4	*
2000.....	0.2	0.3	0.1	0.2	0.3	0.0	0.3	0.5	*
1999.....	0.2	0.3	0.0	0.2	0.3	0.1	0.2	0.4	*
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)									
2007.....	14.4	20.9	8.2	14.8	21.3	8.5	14.1	22.5	7.0
2006.....	15.0	21.4	8.8	15.4	21.8	9.1	14.6	22.6	7.8
2005.....	15.2	21.7	8.9	15.6	22.2	9.2	14.5	22.5	7.6
2004.....	15.2	21.4	9.3	15.6	21.8	9.5	14.8	22.7	8.1
2003.....	15.3	21.6	9.3	15.7	21.9	9.5	14.9	22.7	8.3
2002.....	15.7	22.1	9.6	16.0	22.4	9.8	15.0	23.2	8.2
2001.....	15.3	21.8	9.3	15.6	22.0	9.5	15.4	23.5	8.5
2000.....	15.4	21.7	9.5	15.6	21.8	9.8	15.7	24.4	8.4
1999.....	15.2	21.3	9.6	15.3	21.3	9.7	15.9	24.3	9.0
Pedestrian involved in collision with motor vehicle (V02-V04)									
2007.....	1.0	1.5	0.6	0.9	1.3	0.5	1.5	2.4	0.8
2006.....	1.1	1.6	0.6	1.0	1.5	0.6	1.5	2.3	0.9
2005.....	1.1	1.6	0.7	1.1	1.5	0.6	1.5	2.4	0.7
2004.....	1.1	1.6	0.7	1.1	1.6	0.6	1.5	2.4	0.8
2003.....	1.2	1.8	0.7	1.2	1.7	0.7	1.6	2.6	0.8
2002.....	1.3	1.8	0.8	1.2	1.7	0.7	1.7	2.8	0.9
2001.....	1.3	1.8	0.7	1.2	1.7	0.6	1.8	2.7	1.0
2000.....	1.2	1.8	0.7	1.1	1.6	0.7	1.8	3.0	0.9
1999.....	1.2	1.8	0.7	1.1	1.6	0.7	1.9	3.2	0.9

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/		Black 2/			
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Motorcyclist involved in collision with railway train (V25,V35)									
2007.....	*	*	*	*	*	*	*	*	*
2006.....	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Other motor vehicle accident involving collision with railway train (V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)									
2007.....	0.0	0.1	0.0	0.1	0.1	0.0	0.1	*	*
2006.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	*	*
2005.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2004.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2003.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	*	*
2002.....	0.1	0.1	0.0	0.1	0.1	0.1	0.1	*	*
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.0	0.1	0.2	0.0	0.1	*	*
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47, V50-V51,V56-V57,V60-V61,V66-V67,V70-V71,V76-V77)									
2007.....	1.3	2.0	0.6	1.3	2.0	0.7	1.4	2.3	0.5
2006.....	1.4	2.1	0.8	1.5	2.1	0.8	1.5	2.4	0.8
2005.....	1.4	2.2	0.7	1.5	2.2	0.8	1.5	2.4	0.6
2004.....	1.6	2.3	0.8	1.6	2.4	0.8	1.7	2.6	0.9
2003.....	1.7	2.5	0.9	1.8	2.6	0.9	1.6	2.7	0.7
2002.....	1.7	2.6	0.9	1.8	2.6	1.0	1.9	3.1	0.8
2001.....	1.5	2.2	0.8	1.5	2.2	0.8	1.7	2.8	0.8
2000.....	1.4	2.0	0.7	1.4	2.0	0.8	1.6	2.6	0.7
1999.....	1.3	2.0	0.7	1.4	2.0	0.7	1.7	2.7	0.8

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)										
2007.....	2.7	3.5	2.1	2.8	3.5	2.2	2.6	3.7	1.7	
2006.....	3.1	3.8	2.4	3.2	3.9	2.5	2.9	4.0	2.1	
2005.....	3.4	4.3	2.6	3.5	4.4	2.7	3.1	4.2	2.2	
2004.....	3.6	4.5	2.8	3.7	4.6	2.9	3.2	4.4	2.3	
2003.....	3.7	4.6	2.8	3.8	4.7	3.0	3.4	4.6	2.5	
2002.....	3.9	5.0	2.9	4.0	5.1	3.0	3.5	4.8	2.4	
2001.....	3.7	4.7	2.8	3.8	4.8	2.9	3.6	4.9	2.4	
2000.....	3.7	4.6	2.9	3.8	4.7	3.0	3.5	4.9	2.3	
1999.....	3.7	4.6	2.9	3.7	4.6	3.0	3.6	5.1	2.4	
Occupant of heavy transport vehicle or bus involved in collision with other motor vehicle (V62-V64,V69,V72-V74,V79)										
2007.....	0.1	0.1	*	0.1	0.1	*	0.1	0.2	*	
2006.....	0.1	0.1	*	0.1	0.1	*	0.1	0.2	*	
2005.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*	
2004.....	0.1	0.1	0.0	0.1	0.1	*	0.1	0.2	*	
2003.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*	
2002.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*	
2001.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*	
2000.....	0.1	0.1	0.0	0.1	0.1	*	0.1	0.2	*	
1999.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	*	*	
Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)										
2007.....	1.2	1.7	0.7	1.2	1.8	0.7	1.0	1.7	0.4	
2006.....	1.3	1.8	0.8	1.3	1.9	0.8	1.1	1.6	0.6	
2005.....	1.3	1.8	0.8	1.4	2.0	0.8	1.1	1.6	0.7	
2004.....	1.4	1.9	0.8	1.4	2.0	0.8	1.2	1.7	0.7	
2003.....	1.5	2.0	0.9	1.5	2.1	1.0	1.2	1.8	0.7	
2002.....	1.5	2.1	0.9	1.6	2.2	1.0	1.2	1.9	0.6	
2001.....	1.4	1.9	0.8	1.4	2.0	0.8	1.1	1.7	0.5	
2000.....	1.3	1.9	0.7	1.4	2.0	0.8	1.1	1.7	0.6	
1999.....	1.3	1.9	0.8	1.3	1.9	0.8	1.2	1.7	0.7	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Other and unspecified land transport accidents (V01,V06,V09.1,V09.3,V09.9,V10-V11,V16-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.7-V80.9,V87.9,V88.9,V89.1,V89.3,V89.9)										
2007.....	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.3	0.1	
2006.....	0.2	0.4	0.1	0.2	0.4	0.1	0.2	0.3	0.1	
2005.....	0.3	0.4	0.1	0.3	0.4	0.2	0.3	0.5	0.1	
2004.....	0.3	0.3	0.1	0.3	0.3	0.2	0.3	0.5	0.1	
2003.....	0.3	0.5	0.2	0.3	0.5	0.2	0.3	0.6	0.1	
2002.....	0.2	0.4	0.1	0.2	0.3	0.1	0.2	0.4	*	
2001.....	0.3	0.5	0.2	0.3	0.5	0.1	0.4	0.6	0.2	
2000.....	0.4	0.5	0.2	0.4	0.5	0.2	0.3	0.5	0.2	
1999.....	0.6	0.9	0.3	0.6	0.9	0.3	0.5	0.8	0.2	
Water transport accidents (V90-V94)										
2007.....	0.2	0.3	0.0	0.2	0.3	0.0	0.2	0.3	*	
2006.....	0.2	0.3	0.0	0.2	0.3	0.1	0.1	0.3	*	
2005.....	0.2	0.3	0.0	0.2	0.3	0.0	0.1	0.2	*	
2004.....	0.2	0.3	0.1	0.2	0.4	0.1	0.1	0.3	*	
2003.....	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.4	*	
2002.....	0.2	0.4	0.0	0.2	0.4	0.0	0.2	0.4	*	
2001.....	0.2	0.4	0.1	0.2	0.4	0.1	0.2	0.4	*	
2000.....	0.2	0.4	0.0	0.2	0.4	0.0	0.2	0.4	*	
1999.....	0.2	0.4	0.1	0.2	0.4	0.1	0.3	0.5	*	
Air and space transport accidents (V95-V97)										
2007.....	0.2	0.3	0.1	0.2	0.3	0.1	*	*	*	
2006.....	0.2	0.4	0.1	0.3	0.4	0.1	*	*	*	
2005.....	0.2	0.3	0.1	0.2	0.4	0.1	*	*	*	
2004.....	0.2	0.4	0.1	0.3	0.5	0.1	*	*	*	
2003.....	0.3	0.4	0.1	0.3	0.5	0.1	*	*	*	
2002.....	0.2	0.4	0.0	0.3	0.5	0.1	*	*	*	
2001.....	0.3	0.5	0.2	0.4	0.6	0.2	0.1	*	*	
2000.....	0.3	0.5	0.1	0.3	0.5	0.1	*	*	*	
1999.....	0.2	0.4	0.1	0.3	0.5	0.1	*	*	*	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/		Black 2/			
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other and unspecified transport accidents and their sequelae (V98-V99,Y85)									
2007.....	0.2	0.4	0.1	0.2	0.4	0.1	0.3	0.5	0.1
2006.....	0.2	0.4	0.1	0.2	0.4	0.1	0.3	0.6	0.1
2005.....	0.2	0.4	0.1	0.2	0.4	0.1	0.3	0.4	0.1
2004.....	0.2	0.4	0.1	0.2	0.3	0.1	0.2	0.4	*
2003.....	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.4	*
2002.....	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.5	*
2001.....	0.2	0.3	0.1	0.2	0.3	0.1	0.3	0.5	*
2000.....	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.4	*
1999.....	0.2	0.3	0.1	0.2	0.3	0.1	0.3	0.4	0.1
Falls (W00-W19)									
2007.....	7.0	9.0	5.5	7.5	9.5	5.9	3.4	4.9	2.4
2006.....	6.6	8.6	5.1	7.0	8.9	5.5	3.7	5.4	2.5
2005.....	6.4	8.3	4.9	6.7	8.6	5.2	3.7	5.2	2.6
2004.....	6.3	8.2	4.8	6.6	8.6	5.1	3.6	5.5	2.3
2003.....	5.9	7.6	4.5	6.1	7.9	4.8	3.4	4.7	2.4
2002.....	5.6	7.4	4.3	5.9	7.6	4.6	3.1	4.9	2.0
2001.....	5.3	7.2	3.9	5.6	7.5	4.1	3.3	5.1	2.0
2000.....	4.8	6.6	3.6	5.0	6.8	3.7	2.9	4.6	1.8
1999.....	4.8	6.6	3.5	5.0	6.8	3.7	3.4	5.3	2.1
Fall on same level (W00-W09,W18)									
2007.....	2.6	3.1	2.2	2.7	3.3	2.3	1.0	1.3	0.9
2006.....	2.4	2.9	2.0	2.5	3.0	2.1	1.1	1.6	0.7
2005.....	2.3	2.9	1.9	2.4	3.0	2.0	1.2	1.7	0.8
2004.....	2.1	2.6	1.7	2.2	2.7	1.8	1.1	1.5	0.8
2003.....	1.9	2.4	1.6	2.0	2.5	1.7	0.7	1.0	0.5
2002.....	1.8	2.3	1.5	1.9	2.4	1.6	0.8	1.3	0.5
2001.....	1.4	1.8	1.1	1.5	1.9	1.2	0.7	1.0	0.5
2000.....	1.2	1.5	1.0	1.2	1.5	1.0	0.5	0.7	0.4
1999.....	0.8	1.1	0.7	0.8	1.1	0.7	0.6	0.8	0.4

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Fall from one level to another (W10-W17)										
2007.....	1.2	1.9	0.5	1.2	2.0	0.6	0.6	1.2	0.2	
2006.....	1.2	1.9	0.5	1.2	2.0	0.6	0.8	1.4	0.3	
2005.....	1.1	1.8	0.5	1.2	2.0	0.6	0.7	1.1	0.3	
2004.....	1.1	1.8	0.5	1.2	1.9	0.6	0.7	1.3	0.3	
2003.....	1.1	1.8	0.5	1.2	1.9	0.6	0.8	1.4	0.4	
2002.....	1.1	1.9	0.5	1.2	2.0	0.6	0.7	1.2	0.4	
2001.....	1.2	2.0	0.5	1.2	2.0	0.6	0.9	1.5	0.3	
2000.....	1.0	1.8	0.4	1.1	1.9	0.5	0.7	1.3	0.3	
1999.....	1.1	1.9	0.5	1.2	1.9	0.6	0.8	1.3	0.3	
Unspecified fall (W19)										
2007.....	3.3	4.0	2.8	3.5	4.1	3.0	1.8	2.4	1.3	
2006.....	3.1	3.7	2.6	3.2	3.9	2.8	1.8	2.4	1.4	
2005.....	2.9	3.5	2.5	3.1	3.6	2.7	1.8	2.4	1.4	
2004.....	3.1	3.8	2.5	3.2	4.0	2.7	1.8	2.7	1.2	
2003.....	2.8	3.4	2.4	2.9	3.5	2.5	1.8	2.4	1.5	
2002.....	2.6	3.2	2.2	2.8	3.3	2.4	1.6	2.4	1.1	
2001.....	2.7	3.5	2.2	2.8	3.6	2.3	1.7	2.6	1.2	
2000.....	2.6	3.3	2.1	2.7	3.4	2.2	1.7	2.5	1.2	
1999.....	2.9	3.7	2.3	3.0	3.8	2.4	2.1	3.1	1.4	
Accidental discharge of firearms (W32-W34)										
2007.....	0.2	0.4	0.0	0.2	0.3	0.1	0.3	0.5	*	
2006.....	0.2	0.4	0.0	0.2	0.3	0.0	0.3	0.6	*	
2005.....	0.3	0.5	0.1	0.3	0.4	0.1	0.4	0.7	*	
2004.....	0.2	0.4	0.0	0.2	0.4	0.1	0.3	0.6	*	
2003.....	0.2	0.5	0.1	0.2	0.5	0.1	0.3	0.6	*	
2002.....	0.3	0.4	0.1	0.2	0.4	0.1	0.4	0.7	*	
2001.....	0.3	0.5	0.1	0.3	0.5	0.1	0.3	0.5	*	
2000.....	0.3	0.5	0.1	0.3	0.5	0.1	0.3	0.6	0.1	
1999.....	0.3	0.5	0.1	0.3	0.5	0.1	0.4	0.7	0.1	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Accidental explosion (W39-W40)									
2007.....	0.0	0.1	0.0	0.0	0.1	*	0.1	*	*
2006.....	0.0	0.1	0.0	0.1	0.1	0.0	*	*	*
2005.....	0.0	0.1	0.0	0.0	0.1	*	0.1	*	*
2004.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2003.....	0.1	0.1	0.0	0.1	0.1	*	*	*	*
2002.....	0.0	0.1	*	0.1	0.1	*	*	*	*
2001.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2000.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
1999.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
Accidental drowning and submersion (W65-W74)									
2007.....	1.1	1.8	0.5	1.1	1.7	0.5	1.2	2.1	0.4
2006.....	1.2	1.9	0.5	1.2	1.8	0.6	1.3	2.3	0.4
2005.....	1.2	1.9	0.5	1.2	1.8	0.5	1.4	2.5	0.4
2004.....	1.1	1.8	0.5	1.1	1.7	0.5	1.3	2.2	0.5
2003.....	1.1	1.8	0.5	1.1	1.7	0.5	1.2	2.2	0.4
2002.....	1.2	2.0	0.5	1.1	1.8	0.4	1.6	2.7	0.5
2001.....	1.1	1.8	0.5	1.1	1.7	0.5	1.5	2.6	0.5
2000.....	1.2	2.0	0.5	1.2	1.8	0.5	1.5	2.6	0.6
1999.....	1.3	2.0	0.5	1.2	1.9	0.5	1.7	3.0	0.5
Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)									
2007.....	1.4	1.7	1.1	1.4	1.7	1.1	1.7	2.3	1.4
2006.....	1.4	1.8	1.1	1.4	1.8	1.1	1.7	2.1	1.5
2005.....	1.4	1.8	1.2	1.4	1.7	1.2	2.0	2.4	1.6
2004.....	1.5	1.9	1.2	1.5	1.9	1.2	1.9	2.5	1.6
2003.....	1.5	1.9	1.2	1.4	1.9	1.2	1.8	2.3	1.4
2002.....	1.4	1.8	1.2	1.4	1.8	1.1	2.1	2.7	1.6
2001.....	1.5	1.9	1.2	1.4	1.8	1.2	2.2	3.1	1.6
2000.....	1.6	2.0	1.3	1.5	1.9	1.3	2.1	2.9	1.7
1999.....	1.4	1.8	1.2	1.4	1.7	1.2	2.2	2.9	1.7

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Accidental exposure to smoke, fire and flames (X00-X09)										
2007.....	1.1	1.3	0.8	0.9	1.2	0.7	2.2	3.0	1.6	
2006.....	1.0	1.3	0.8	0.9	1.2	0.7	2.1	2.7	1.6	
2005.....	1.1	1.4	0.8	0.9	1.2	0.7	2.2	3.1	1.6	
2004.....	1.1	1.4	0.8	0.9	1.2	0.7	2.4	3.3	1.8	
2003.....	1.2	1.4	0.9	1.1	1.3	0.8	2.4	3.2	1.9	
2002.....	1.1	1.4	0.8	0.9	1.3	0.7	2.4	3.3	1.7	
2001.....	1.2	1.5	0.9	1.0	1.3	0.7	2.7	3.8	1.9	
2000.....	1.2	1.6	0.9	1.0	1.3	0.7	3.0	4.2	2.2	
1999.....	1.2	1.6	0.9	1.0	1.4	0.7	3.0	4.0	2.3	
Accidental contact with heat and hot substances (X10-X19)										
2007.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2006.....	0.0	0.1	0.0	0.0	0.0	0.0	0.1	*	*	
2005.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*	
2003.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*	
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*	
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*	
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*	
Victim of lightning (X33)										
2007.....	0.0	0.0	*	0.0	0.0	*	*	*	*	
2006.....	0.0	0.0	*	0.0	0.0	*	*	*	*	
2005.....	0.0	0.0	*	0.0	0.0	*	*	*	*	
2004.....	0.0	0.0	*	0.0	0.0	*	*	*	*	
2003.....	0.0	0.0	*	0.0	0.0	*	*	*	*	
2002.....	0.0	0.0	*	0.0	0.1	*	*	*	*	
2001.....	0.0	0.0	*	0.0	0.0	*	*	*	*	
2000.....	0.0	0.0	*	0.0	0.0	*	*	*	*	
1999.....	0.0	0.1	*	0.0	0.1	*	*	*	*	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)									
2007.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2006.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2005.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2004.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2003.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2002.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2001.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2000.....	0.0	0.0	*	0.0	0.0	*	*	*	*
1999.....	0.0	0.0	*	0.0	0.0	*	*	*	*
Cataclysmic storm and flood (X37-X38)									
2007.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2006.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2005.....	0.3	0.3	0.2	0.2	0.2	0.2	1.3	2.0	0.9
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	*	*	*	*
1999.....	0.1	0.1	0.1	0.1	0.0	0.1	0.1	*	*
Accidental poisoning and exposure to noxious substances (X40-X49)									
2007.....	9.8	13.0	6.6	10.6	14.0	7.2	8.6	12.2	5.5
2006.....	9.1	12.4	5.9	9.7	13.0	6.3	9.4	14.1	5.5
2005.....	7.9	10.7	5.1	8.4	11.3	5.5	8.2	11.9	4.9
2004.....	7.1	9.5	4.7	7.6	10.1	5.0	6.9	9.9	4.5
2003.....	6.7	9.1	4.3	7.1	9.6	4.5	6.8	9.9	4.1
2002.....	6.1	8.4	3.7	6.3	8.7	3.9	6.8	10.1	3.9
2001.....	4.9	7.0	2.9	5.0	7.1	2.9	6.3	9.5	3.7
2000.....	4.5	6.6	2.5	4.5	6.6	2.5	6.0	9.3	3.3
1999.....	4.4	6.4	2.3	4.3	6.3	2.3	6.2	9.8	3.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)									
2007.....	9.1	12.0	6.2	9.9	12.9	6.8	8.0	11.3	5.2
2006.....	8.8	11.8	5.7	9.3	12.4	6.1	9.1	13.5	5.2
2005.....	7.6	10.1	5.0	8.0	10.7	5.3	7.7	11.2	4.7
2004.....	6.8	9.0	4.5	7.2	9.5	4.8	6.6	9.3	4.3
2003.....	6.3	8.5	4.1	6.7	9.0	4.3	6.4	9.2	4.0
2002.....	5.7	7.8	3.5	5.9	8.1	3.7	6.3	9.3	3.7
2001.....	4.6	6.5	2.7	4.7	6.5	2.7	5.9	8.7	3.3
2000.....	4.1	6.0	2.3	4.2	6.0	2.3	5.6	8.6	3.1
1999.....	4.0	5.9	2.1	3.9	5.8	2.1	5.8	9.0	2.9
Accidental poisoning by and exposure to other and unspecified solid or liquids substances (X45-X46,X48-X49)									
2007.....	0.5	0.8	0.2	0.5	0.8	0.3	0.3	0.5	0.2
2006.....	0.2	0.2	0.1	0.2	0.2	0.1	0.1	0.3	*
2005.....	0.2	0.3	0.1	0.2	0.3	0.1	0.1	0.2	*
2004.....	0.2	0.3	0.1	0.2	0.3	0.1	0.1	0.2	*
2003.....	0.2	0.3	0.1	0.2	0.3	0.1	0.1	0.3	*
2002.....	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.4	0.1
2001.....	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.1
2000.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.3	*
1999.....	0.2	0.2	0.1	0.2	0.2	0.1	0.2	0.3	*
Accidental poisoning by and exposure to other gases and vapors (X47)									
2007.....	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.3	0.2
2006.....	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.4	0.2
2005.....	0.2	0.3	0.1	0.2	0.3	0.1	0.3	0.5	0.2
2004.....	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.3	0.1
2003.....	0.2	0.3	0.1	0.2	0.3	0.1	0.3	0.4	0.1
2002.....	0.2	0.4	0.1	0.3	0.3	0.1	0.2	0.4	0.1
2001.....	0.2	0.3	0.1	0.2	0.3	0.1	0.3	0.5	0.2
2000.....	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.4	0.1
1999.....	0.2	0.3	0.1	0.2	0.3	0.1	0.3	0.4	0.1

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All races 1/			White 2/		Black 2/			
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)									
2007.....	3.2	4.4	2.2	3.3	4.4	2.2	3.1	4.4	2.0
2006.....	3.4	4.7	2.3	3.5	4.7	2.4	3.2	4.9	2.0
2005.....	3.6	4.9	2.4	3.7	5.0	2.4	3.4	5.0	2.1
2004.....	3.4	4.6	2.4	3.5	4.7	2.4	3.3	4.9	2.0
2003.....	3.6	4.9	2.5	3.7	4.9	2.6	3.6	5.3	2.3
2002.....	3.7	5.0	2.6	3.8	5.0	2.7	3.7	5.4	2.4
2001.....	4.0	5.3	2.9	4.1	5.4	2.9	3.9	5.5	2.7
2000.....	4.0	5.3	2.8	4.0	5.3	2.8	4.0	6.5	2.3
1999.....	4.4	5.9	3.1	4.4	5.8	3.1	4.9	7.7	3.0
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)									
2007.....	11.3	18.4	4.7	12.5	20.2	5.2	5.0	8.8	1.7
2006.....	10.9	18.0	4.5	12.1	19.6	5.1	5.1	9.4	1.4
2005.....	10.9	18.0	4.4	12.0	19.6	4.9	5.2	9.2	1.9
2004.....	10.9	18.0	4.5	12.0	19.6	5.0	5.3	9.6	1.8
2003.....	10.8	18.0	4.2	11.8	19.6	4.6	5.2	9.2	1.9
2002.....	10.9	18.4	4.2	12.0	20.0	4.7	5.3	9.8	1.6
2001 3/.....	10.7	18.2	4.0	11.7	19.6	4.5	5.5	9.8	1.8
2000.....	10.4	17.7	4.0	11.3	19.1	4.3	5.5	10.0	1.8
1999.....	10.5	17.8	4.0	11.3	19.0	4.4	5.6	10.3	1.6
Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)									
2007.....	1.5	1.5	1.6	1.8	1.6	1.9	0.4	0.4	0.5
2006.....	1.5	1.4	1.6	1.7	1.6	1.8	0.4	0.5	0.4
2005.....	1.4	1.3	1.5	1.6	1.5	1.7	0.5	0.5	0.5
2004.....	1.4	1.3	1.5	1.6	1.5	1.7	0.4	0.4	0.5
2003.....	1.3	1.3	1.4	1.5	1.5	1.5	0.5	0.4	0.6
2002.....	1.3	1.3	1.4	1.5	1.4	1.6	0.4	0.5	0.4
2001.....	1.3	1.2	1.3	1.4	1.4	1.4	0.5	0.4	0.5
2000.....	1.2	1.1	1.2	1.3	1.3	1.4	0.5	0.5	0.4
1999.....	1.1	1.1	1.2	1.3	1.2	1.3	0.4	0.5	0.4

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)									
2007.....	0.1	0.1	0.1	0.1	0.1	0.0	0.1	*	*
2006.....	0.1	0.1	0.1	0.1	0.2	0.0	*	*	*
2005.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2004.....	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2003.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2002.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2001.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2000.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
1999.....	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)									
2007.....	0.4	0.7	0.2	0.5	0.8	0.2	0.1	0.2	*
2006.....	0.4	0.7	0.2	0.5	0.8	0.2	0.1	0.2	*
2005.....	0.4	0.7	0.2	0.5	0.8	0.2	0.1	0.2	*
2004.....	0.5	0.7	0.2	0.5	0.9	0.2	0.1	0.2	*
2003.....	0.5	0.8	0.2	0.5	0.9	0.2	0.1	0.2	*
2002.....	0.5	0.8	0.2	0.6	0.9	0.2	0.1	0.2	*
2001.....	0.5	0.8	0.2	0.6	0.9	0.3	0.1	0.2	*
2000.....	0.5	0.8	0.2	0.6	0.9	0.3	0.1	0.2	*
1999.....	0.5	0.9	0.3	0.6	1.0	0.3	0.1	0.2	*
Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)									
2007.....	2.7	4.4	1.0	2.9	4.8	1.0	1.3	2.2	0.4
2006.....	2.5	4.2	0.9	2.7	4.5	0.9	1.2	2.3	0.3
2005.....	2.4	4.0	0.9	2.6	4.3	0.9	1.2	2.1	0.5
2004.....	2.5	4.1	0.9	2.6	4.4	0.9	1.4	2.4	0.4
2003.....	2.3	3.9	0.8	2.4	4.0	0.8	1.3	2.3	0.4
2002.....	2.2	3.7	0.8	2.4	4.0	0.8	1.1	2.1	0.3
2001.....	2.2	3.7	0.7	2.3	3.9	0.7	1.1	2.0	0.3
2000.....	2.0	3.4	0.7	2.1	3.6	0.7	1.2	2.1	0.3
1999.....	1.9	3.3	0.6	2.0	3.4	0.7	1.0	2.0	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Intentional self-harm (suicide) by discharge of firearms (X72-X74)										
2007.....	5.6	10.3	1.4	6.3	11.4	1.6	2.5	4.9	0.5	
2006.....	5.5	10.2	1.4	6.2	11.2	1.6	2.6	5.3	0.5	
2005.....	5.7	10.5	1.4	6.3	11.5	1.6	2.7	5.3	0.5	
2004.....	5.6	10.3	1.5	6.3	11.4	1.7	2.7	5.4	0.5	
2003.....	5.8	10.7	1.4	6.4	11.8	1.6	2.7	5.3	0.6	
2002.....	5.9	11.1	1.4	6.5	12.1	1.6	3.0	5.9	0.5	
2001.....	5.9	11.0	1.4	6.5	12.0	1.6	3.1	6.0	0.6	
2000.....	5.9	11.0	1.5	6.5	11.9	1.6	3.1	6.0	0.7	
1999.....	6.0	11.1	1.5	6.5	12.0	1.7	3.3	6.5	0.6	
Intentional self-harm (suicide) by jumping from a high place (X80)										
2007.....	0.2	0.4	0.1	0.3	0.4	0.1	0.1	0.3	*	
2006.....	0.2	0.4	0.1	0.2	0.4	0.1	0.2	0.3	*	
2005.....	0.2	0.3	0.1	0.2	0.4	0.1	0.2	0.3	0.1	
2004.....	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.3	*	
2003.....	0.2	0.3	0.1	0.2	0.4	0.1	0.2	0.3	*	
2002.....	0.2	0.4	0.1	0.3	0.4	0.1	0.2	0.3	*	
2001.....	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.3	*	
2000.....	0.2	0.3	0.1	0.2	0.3	0.1	0.2	0.3	*	
1999.....	0.3	0.4	0.1	0.3	0.4	0.1	0.2	0.3	*	
Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,Y87.0)										
2007.....	0.7	1.0	0.3	0.7	1.0	0.3	0.4	0.7	0.3	
2006.....	0.7	1.1	0.3	0.7	1.1	0.3	0.4	0.7	0.2	
2005.....	0.7	1.0	0.3	0.7	1.0	0.3	0.5	0.8	0.3	
2004.....	0.6	1.0	0.3	0.7	1.0	0.3	0.5	0.8	0.2	
2003.....	0.6	0.9	0.3	0.6	0.9	0.3	0.5	0.7	0.2	
2002.....	0.6	1.0	0.3	0.7	1.0	0.3	0.5	0.8	0.2	
2001 3/.....	0.6	0.9	0.3	0.6	1.0	0.3	0.5	0.7	0.3	
2000.....	0.6	0.9	0.3	0.6	0.9	0.3	0.5	0.8	0.2	
1999.....	0.6	0.9	0.3	0.6	0.9	0.3	0.5	0.8	0.3	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)										
2007.....	6.1	9.6	2.5	3.7	5.4	2.0	21.1	37.1	6.1	
2006.....	6.2	9.7	2.5	3.7	5.4	1.9	21.6	37.8	6.4	
2005.....	6.1	9.6	2.5	3.7	5.3	1.9	21.1	37.3	6.1	
2004.....	5.9	9.2	2.5	3.6	5.3	1.9	20.1	35.1	6.3	
2003.....	6.0	9.4	2.6	3.7	5.3	2.0	21.0	36.7	6.4	
2002.....	6.1	9.4	2.8	3.7	5.3	2.0	21.0	36.4	6.9	
2001 3/.....	7.1	10.8	3.3	4.9	7.1	2.6	21.2	36.2	7.4	
2000.....	5.9	9.0	2.8	3.6	5.2	2.1	20.5	35.4	7.1	
1999.....	6.0	9.1	2.9	3.8	5.4	2.1	20.1	34.2	7.3	
Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (*U01.6-*U01.7,X85-X90)										
2007.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2006.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*	
2005.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*	
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*	
2003.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*	
2002.....	0.0	0.0	0.0	0.0	0.0	*	*	*	*	
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
2000.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	
Assault (homicide) by hanging, strangulation and suffocation (X91)										
2007.....	0.2	0.1	0.3	0.2	0.1	0.2	0.5	0.4	0.5	
2006.....	0.2	0.2	0.2	0.2	0.1	0.2	0.5	0.4	0.6	
2005.....	0.2	0.1	0.3	0.2	0.1	0.2	0.5	0.4	0.6	
2004.....	0.2	0.1	0.3	0.2	0.1	0.2	0.5	0.4	0.7	
2003.....	0.2	0.1	0.3	0.2	0.1	0.2	0.6	0.4	0.7	
2002.....	0.2	0.2	0.3	0.2	0.1	0.2	0.6	0.4	0.7	
2001.....	0.2	0.2	0.3	0.2	0.1	0.2	0.6	0.4	0.7	
2000.....	0.2	0.2	0.3	0.2	0.1	0.2	0.6	0.3	0.8	
1999.....	0.2	0.2	0.3	0.2	0.2	0.2	0.6	0.4	0.8	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)										
2007.....	4.2	7.1	1.2	2.2	3.5	1.0	16.3	30.1	3.2	
2006.....	4.3	7.2	1.3	2.2	3.4	0.9	16.7	30.8	3.5	
2005.....	4.2	7.0	1.2	2.2	3.5	0.9	16.0	29.8	3.0	
2004.....	4.0	6.6	1.1	2.2	3.4	0.8	15.0	28.0	2.9	
2003.....	4.1	6.8	1.2	2.2	3.4	0.9	15.7	29.0	3.2	
2002.....	4.1	6.8	1.3	2.2	3.4	1.0	15.6	28.7	3.5	
2001.....	3.9	6.6	1.3	2.2	3.5	1.0	14.8	27.4	3.0	
2000.....	3.8	6.3	1.2	2.1	3.2	0.9	14.6	26.9	3.1	
1999.....	3.8	6.3	1.3	2.1	3.3	1.0	14.3	26.1	3.5	
Assault (homicide) by sharp object (X99)										
2007.....	0.6	1.0	0.3	0.4	0.7	0.2	2.0	3.2	1.0	
2006.....	0.7	1.0	0.4	0.5	0.7	0.3	2.0	3.1	1.0	
2005.....	0.7	1.0	0.4	0.5	0.7	0.3	2.2	3.4	1.1	
2004.....	0.7	1.0	0.4	0.5	0.7	0.3	2.2	3.3	1.2	
2003.....	0.7	1.0	0.4	0.5	0.7	0.3	2.2	3.5	1.1	
2002.....	0.7	1.0	0.4	0.5	0.7	0.3	2.2	3.3	1.2	
2001.....	0.7	1.0	0.4	0.5	0.6	0.3	2.1	3.1	1.2	
2000.....	0.6	0.9	0.4	0.5	0.7	0.3	2.0	2.9	1.1	
1999.....	0.7	1.0	0.4	0.4	0.6	0.3	2.0	3.3	1.0	
Assault (homicide) by blunt object (Y00)										
2007.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	*	
2006.....	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.1	*	
2005.....	0.0	0.1	0.0	0.0	0.1	0.0	0.1	0.1	*	
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	*	
2003.....	0.0	0.1	0.0	0.0	0.1	0.0	0.1	0.2	*	
2002.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.3	*	
2001.....	0.1	0.1	0.1	0.1	0.1	0.0	0.2	0.3	*	
2000.....	0.1	0.1	0.1	0.1	0.1	0.0	0.2	0.3	0.1	
1999.....	0.1	0.1	0.0	0.1	0.1	0.0	0.2	0.3	0.1	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Assault (homicide) by bodily force (Y04)									
2007.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2006.....	0.0	0.0	*	0.0	0.0	*	0.1	0.1	*
2005.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2004.....	0.0	0.0	*	0.0	0.0	*	0.1	*	*
2003.....	0.0	0.1	*	0.0	0.1	*	0.1	0.2	*
2002.....	0.0	0.1	*	0.0	0.0	*	0.1	0.2	*
2001.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2000.....	0.1	0.1	*	0.0	0.1	*	0.1	0.2	*
1999.....	0.1	0.1	0.0	0.0	0.1	*	0.2	0.3	*
Sexual assault (homicide) by bodily force (Y05)									
2007.....	*	*	*	*	*	*	*	*	*
2006.....	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*
Neglect, abandonment and other maltreatment syndromes (Y06-Y07)									
2007.....	0.1	0.1	0.0	0.0	0.0	0.0	0.1	0.1	0.1
2006.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	*
2005.....	0.1	0.1	0.1	0.0	0.0	0.1	0.1	0.2	0.1
2004.....	0.1	0.1	0.1	0.0	0.1	0.0	0.1	0.1	0.1
2003.....	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.2	0.1
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2
2001.....	0.1	0.1	0.1	0.1	0.1	0.0	0.2	0.2	0.2
2000.....	0.1	0.1	0.1	0.1	0.1	0.0	0.2	0.2	0.2
1999.....	0.1	0.1	0.1	0.0	0.1	0.0	0.2	0.2	0.2

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female

Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

2007.....	0.9	1.2	0.6	0.7	0.9	0.4	2.0	3.0	1.2
2006.....	0.9	1.2	0.5	0.7	0.9	0.5	2.0	3.0	1.1
2005.....	0.9	1.2	0.5	0.7	0.9	0.4	2.1	3.2	1.2
2004.....	0.9	1.1	0.6	0.7	0.9	0.4	2.0	2.9	1.2
2003.....	0.9	1.2	0.6	0.7	0.9	0.5	2.1	3.2	1.2
2002.....	0.8	1.1	0.5	0.6	0.9	0.4	2.2	3.4	1.2
2001 3/.....	2.0	2.8	1.1	1.8	2.6	1.0	3.2	4.6	2.1
2000.....	1.0	1.4	0.7	0.7	0.9	0.5	2.9	4.4	1.7
1999.....	1.0	1.3	0.7	0.8	1.0	0.5	2.5	3.6	1.5

Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)

2007.....	1.8	2.2	1.3	1.8	2.2	1.4	2.0	3.0	1.1
2006.....	1.7	2.2	1.2	1.7	2.1	1.3	1.9	2.9	1.0
2005.....	1.6	1.9	1.2	1.6	1.9	1.3	1.8	2.5	1.2
2004.....	1.7	2.2	1.2	1.7	2.2	1.3	1.9	2.7	1.1
2003.....	1.7	2.3	1.2	1.8	2.3	1.3	1.9	2.9	1.0
2002.....	1.7	2.2	1.2	1.7	2.2	1.2	2.1	3.1	1.2
2001.....	1.5	2.0	0.9	1.5	2.0	1.0	1.8	2.9	0.9
2000.....	1.3	1.9	0.8	1.3	1.8	0.8	1.9	3.0	1.0
1999.....	1.4	2.0	0.9	1.4	1.8	0.9	2.0	3.2	0.9

Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)

2007.....	1.1	1.3	1.0	1.3	1.4	1.0	1.0	1.5	0.7
2006.....	1.1	1.3	0.9	1.2	1.4	1.0	1.0	1.4	0.7
2005.....	1.0	1.2	0.9	1.1	1.2	1.0	1.0	1.2	0.8
2004.....	1.1	1.4	0.9	1.2	1.4	1.0	1.1	1.5	0.7
2003.....	1.2	1.5	0.9	1.3	1.6	1.0	1.2	1.8	0.7
2002.....	1.1	1.3	0.8	1.1	1.4	0.9	1.3	1.7	0.9
2001.....	1.0	1.3	0.7	1.0	1.3	0.7	1.1	1.8	0.5
2000.....	0.8	1.2	0.6	0.9	1.2	0.6	1.1	1.6	0.6
1999.....	0.9	1.2	0.6	0.9	1.1	0.6	1.2	1.9	0.6

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/	
		Male	Female		Male	Female		Male	Female
Poisoning by and exposure to other and unspecified solid or liquid substance, undetermined intent (Y15-Y16,Y18-Y19)									
2007.....	0.0	0.1	0.0	0.0	0.1	0.0	0.1	0.1	*
2006.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2005.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2003.....	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2002.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2001.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2000.....	0.0	0.0	*	0.0	0.0	*	*	*	*
1999.....	0.0	0.0	*	0.0	0.0	*	*	*	*
Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)									
2007.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2006.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2005.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2003.....	0.0	0.1	0.0	0.0	0.1	0.0	*	*	*
2002.....	0.0	0.0	*	0.0	0.0	*	*	*	*
2001.....	0.0	0.1	*	0.0	0.1	*	*	*	*
2000.....	0.0	0.0	0.0	0.0	0.0	*	*	*	*
1999.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
Discharge of firearms, undetermined intent (Y22-Y24)									
2007.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2006.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2005.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*
2004.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*
2003.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2002.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2001.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2000.....	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.3	*
1999.....	0.1	0.2	0.0	0.1	0.2	0.0	0.2	0.3	*

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Falling, jumping or pushed from a high place, undetermined intent (Y30)										
2007.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2006.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2005.....	0.0	0.0	*	0.0	0.0	0.0	*	*	*	*
2004.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2003.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2002.....	0.0	0.0	0.0	0.0	0.1	0.0	*	*	*	*
2001.....	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*	*
2000.....	0.0	0.0	*	0.0	0.0	0.0	*	*	*	*
1999.....	0.0	0.0	*	0.0	0.0	0.0	*	*	*	*
All other and unspecified events of undetermined intent and their sequelae (Y20-Y21,Y25-Y29,Y31-Y34,Y87.2,Y89.9)										
2007.....	0.4	0.6	0.3	0.4	0.5	0.2	0.7	1.1	0.4	
2006.....	0.4	0.6	0.3	0.4	0.5	0.2	0.7	1.1	0.3	
2005.....	0.4	0.6	0.3	0.3	0.5	0.2	0.7	1.1	0.3	
2004.....	0.4	0.6	0.3	0.4	0.5	0.2	0.7	1.0	0.4	
2003.....	0.4	0.5	0.2	0.3	0.5	0.2	0.5	0.8	0.3	
2002.....	0.4	0.5	0.2	0.4	0.5	0.2	0.6	1.1	0.3	
2001.....	0.3	0.5	0.2	0.3	0.4	0.2	0.5	0.8	0.3	
2000.....	0.3	0.5	0.2	0.3	0.4	0.2	0.6	1.0	0.3	
1999.....	0.3	0.5	0.2	0.3	0.4	0.2	0.5	0.9	0.2	
Legal intervention (Y35,Y89.0)										
2007.....	0.1	0.3	*	0.1	0.2	*	0.3	0.6	*	
2006.....	0.2	0.3	*	0.1	0.2	*	0.3	0.6	*	
2005.....	0.1	0.3	*	0.1	0.2	*	0.3	0.6	*	
2004.....	0.1	0.3	*	0.1	0.2	*	0.2	0.5	*	
2003.....	0.1	0.3	*	0.1	0.2	*	0.3	0.6	*	
2002.....	0.1	0.3	*	0.1	0.2	*	0.3	0.6	*	
2001.....	0.1	0.3	*	0.1	0.2	*	0.2	0.5	*	
2000.....	0.1	0.2	*	0.1	0.2	*	0.3	0.6	*	
1999.....	0.1	0.3	*	0.1	0.2	*	0.3	0.7	*	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Legal execution (Y35.5)										
2007.....	0.0	0.0	*	0.0	0.0	*	*	*	*	
2006.....	0.0	0.0	*	0.0	0.0	*	*	*	*	
2005.....	0.0	0.0	*	0.0	0.0	*	*	*	*	
2004.....	0.0	0.0	*	0.0	0.0	*	*	*	*	
2003.....	0.0	0.1	*	0.0	0.0	*	*	*	*	
2002.....	0.0	0.1	*	0.0	0.0	*	*	*	*	
2001.....	0.0	0.1	*	0.0	0.0	*	*	*	*	
2000.....	0.0	0.0	*	0.0	0.0	*	0.1	0.2	*	
1999.....	0.0	0.1	*	0.0	0.1	*	0.1	0.2	*	
Other legal intervention and their sequelae (Y35.0-Y35.4,Y35.6-Y35.7,Y89.0)										
2007.....	0.1	0.2	*	0.1	0.2	*	0.3	0.5	*	
2006.....	0.1	0.3	*	0.1	0.2	*	0.2	0.5	*	
2005.....	0.1	0.2	*	0.1	0.2	*	0.2	0.5	*	
2004.....	0.1	0.2	*	0.1	0.2	*	0.2	0.4	*	
2003.....	0.1	0.2	*	0.1	0.2	*	0.3	0.5	*	
2002.....	0.1	0.2	*	0.1	0.2	*	0.2	0.5	*	
2001.....	0.1	0.2	*	0.1	0.2	*	0.2	0.4	*	
2000.....	0.1	0.2	*	0.1	0.2	*	0.2	0.4	*	
1999.....	0.1	0.2	*	0.1	0.2	*	0.2	0.5	*	
Operations of war and their sequelae (Y36,Y89.1)										
2007.....	0.0	0.0	*	0.0	0.0	*	*	*	*	
2006.....	0.0	0.0	*	0.0	0.0	*	*	*	*	
2005.....	0.0	0.0	*	0.0	0.0	*	*	*	*	
2004.....	0.0	0.0	*	0.0	0.0	*	*	*	*	
2003.....	*	*	*	*	*	*	*	*	*	
2002.....	0.0	0.0	*	*	*	*	*	*	*	
2001.....	*	*	*	*	*	*	*	*	*	
2000.....	*	*	*	*	*	*	*	*	*	
1999.....	0.0	0.0	*	0.0	0.0	*	*	*	*	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Complications of medical and surgical care (Y40-Y84,Y88)										
2007.....	0.8	0.9	0.8	0.8	0.9	0.7	1.4	1.6	1.4	
2006.....	0.8	0.8	0.8	0.8	0.8	0.7	1.4	1.4	1.4	
2005.....	0.9	0.9	0.8	0.8	0.9	0.8	1.5	1.5	1.5	
2004.....	1.0	1.1	0.9	0.9	1.1	0.8	1.5	1.5	1.6	
2003.....	1.0	1.1	0.9	0.9	1.0	0.8	1.5	1.5	1.6	
2002.....	1.0	1.0	1.0	0.9	1.0	0.9	1.6	1.6	1.6	
2001.....	1.1	1.1	1.0	1.0	1.1	0.9	2.0	2.3	1.7	
2000.....	1.1	1.2	1.0	1.0	1.2	0.9	1.9	1.9	1.9	
1999.....	1.0	1.1	0.9	1.0	1.1	0.9	1.8	2.0	1.7	
Drugs, medicaments and biological substances causing adverse effects in therapeutic use, and their sequelae (Y40-Y59,Y88.0)										
2007.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	*	
2006.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.2	
2005.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2	
2004.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2	
2003.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	*	
2002.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1	
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2	
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.2	
1999.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	
Misadventures to patients during medical and surgical care and their sequelae (Y60-Y69,Y88.1)										
2007.....	0.0	0.0	0.1	0.0	0.0	0.0	0.1	*	0.2	
2006.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	0.2	
2005.....	0.0	0.0	0.0	0.0	0.0	0.0	0.1	*	0.1	
2004.....	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*	*	
2003.....	0.1	0.0	0.1	0.0	0.0	0.1	0.1	*	0.1	
2002.....	0.0	0.0	0.1	0.0	0.0	0.0	0.1	*	0.1	
2001.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	
2000.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.2	
1999.....	0.1	0.1	0.1	0.0	0.1	0.1	0.1	*	0.2	

Worktable 13. Age-adjusted death rates for 358 selected causes, by race and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	Both sexes	All races 1/		Both sexes	White 2/		Both sexes	Black 2/		
		Male	Female		Male	Female		Male	Female	
Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)										
2007.....	0.7	0.7	0.6	0.6	0.7	0.6	1.2	1.3	1.1	
2006.....	0.7	0.7	0.6	0.6	0.7	0.6	1.1	1.1	1.0	
2005.....	0.7	0.8	0.7	0.7	0.7	0.6	1.2	1.3	1.2	
2004.....	0.8	0.9	0.7	0.8	0.9	0.6	1.3	1.3	1.3	
2003.....	0.8	0.9	0.8	0.8	0.9	0.7	1.3	1.2	1.3	
2002.....	0.8	0.9	0.8	0.8	0.9	0.8	1.4	1.4	1.3	
2001.....	0.9	1.0	0.9	0.8	0.9	0.8	1.6	2.0	1.4	
2000.....	0.9	1.1	0.9	0.9	1.0	0.8	1.6	1.7	1.6	
1999.....	0.9	1.0	0.8	0.8	0.9	0.8	1.5	1.7	1.4	

* Figure does not meet standards of reliability or precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

1/ Includes races other than white and black.

2/ Multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming in 2007, by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming in 2006, by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming in 2005, by California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and by California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality."

3/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

Source: CDC/NCHS, National Vital Statistics System, Mortality.