


Figure 6.1. Prevalence of obesity among adults aged 20 years and over: United States, 1997–June 2010




NOTES: Obesity is defined as a body mass index (BMI) of 30 kg/m² or more. The measure is based on self-reported height and weight. Estimates of obesity are restricted to adults aged 20 years and over for consistency with the Healthy People 2010 (3) program. The analyses excluded people with unknown height or weight (about 6% of respondents each year). Beginning with the 2003 data, the National Health Interview Survey (NHIS) transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. See "About This Early Release" for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997–June 2010, Sample Adult Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For January–June 2010, 28.2% (95% confidence interval = 27.28%–29.15%) of U.S. adults aged 20 years and over were obese, which was not significantly different from the 2009 estimate of 28.0%.
- The prevalence of obesity among U.S. adults aged 20 years and over has increased over time, from 19.4% in 1997 to 28.2% for January–June 2010.

Figure 6.2. Prevalence of obesity among adults aged 20 years and over, by age group and sex: United States, January–June 2010




NOTES: Obesity is defined as a body mass index (BMI) of 30 kg/m² or more. The measure is based on self-reported height and weight. Estimates of obesity are restricted to adults aged 20 years and over for consistency with the Healthy People 2010 (3) program. The analyses excluded 585 people (4.2%) with unknown height or weight.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–June 2010, Sample Adult Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- For both sexes combined, the prevalence of obesity was higher among adults aged 40–59 years (31.9%) than among adults aged 20–39 (24.6%) and those aged 60 and over (28.0%).
- For adults aged 20–39 years, 40–59 years, and 60 years and over, there was no significant difference between women and men in the prevalence of obesity.

Figure 6.3. Age-adjusted prevalence of obesity among adults aged 20 years and over, by sex and race/ethnicity: United States, January–June 2010



NOTES: Obesity is defined as a body mass index (BMI) of 30 kg/m² or more. The measure is based on self-reported height and weight. Estimates of obesity are restricted to adults aged 20 years and over for consistency with the Healthy People 2010 (3) program. The analyses excluded 585 people (4.2%) with unknown height or weight. Estimates are age-adjusted using the projected 2000 U.S. population as the standard population and using five age groups: 20–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–June 2010, Sample Adult Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

- Non-Hispanic black women were most likely to be obese, followed by Hispanic women and non-Hispanic white women.
- Non-Hispanic white men were less likely than Hispanic men and non-Hispanic black men to be obese.

Data tables for Figures 6.1–6.3:

Data table for Figure 6.1. Prevalence of obesity among adults aged 20 years and over: United States, 1997–June 2010

Year	Crude ¹ percent (95% confidence interval)	Age-adjusted ² percent (95% confidence interval)
1997	19.4 (18.9-19.9)	19.5 (18.9-20.0)
1998	20.6 (20.1-21.1)	20.6 (20.0-21.1)
1999	21.5 (20.9-22.1)	21.5 (20.9-22.1)
2000	21.8 (21.2-22.4)	21.8 (21.2-22.3)
2001	23.0 (22.4-23.6)	22.9 (22.3-23.5)
2002	23.9 (23.3-24.6)	23.8 (23.2-24.5)
2003	23.7 (23.1-24.3)	23.5 (22.9-24.2)
2004	24.5 (23.9-25.1)	24.3 (23.8-25.0)
2005	25.4 (24.77-26.09)	25.3 (24.66-25.96)
2006	26.4 (25.62-27.09)	26.2 (25.44-26.90)
2007	26.7 (25.82-27.50)	26.6 (25.78-27.49)
2008	27.6 (26.80-28.50)	27.5 (26.69-28.36)
2009	28.0 (27.20-28.76)	27.9 (27.13-28.71)
January-June 2010	28.2 (27.28-29.15)	28.0 (27.05-28.95)

¹Crude estimates are presented in the figure.

²Estimates for this *Healthy People 2010* Leading Health Indicator are age-adjusted using the projected 2000 U.S. population as the standard population and using seven age groups: 20–29 years, 30–39 years, 40–49 years, 50–59 years, 60–69 years, 70–79 years, and 80 years and over.

NOTES: Beginning with the 2003 data, the National Health Interview Survey transitioned to weights derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. See "About This Early Release" for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997–June 2010, Sample Adult Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

Data table for Figure 6.2. Prevalence of obesity among adults aged 20 years and over, by age group and sex: United States, January–June 2010

Age and sex	Percent	95% confidence interval
20-39 years, total	24.6	23.06-26.13
20-39 years, male	25.2	23.11-27.36
20-39 years, female	23.9	21.95-25.95
40-59 years, total	31.9	30.40-33.44
40-59 years, male	32.0	29.82-34.14
40-59 years, female	31.9	29.82-33.90
60 years and over, total	28.0	26.53-29.50
60 years and over, male	28.2	25.70-30.80
60 years and over, female	27.8	25.80-29.85
20 years and over (crude ¹), total	28.2	27.28-29.15
20 years and over (crude ¹), male	28.5	27.18-29.90
20 years and over (crude ¹), female	27.9	26.68-29.12
20 years and over (age-adjusted ²), total	28.0	27.05-28.95
20 years and over (age-adjusted ²), male	28.2	26.83-29.53
20 years and over (age-adjusted ²), female	27.8	26.52-29.03

¹Crude estimates are presented in the figure.

²Estimates for this *Healthy People 2010* Leading Health Indicator are age-adjusted using the projected 2000 U.S. population as the standard population and using seven age groups: 20–29 years, 30–39 years, 40–49 years, 50–59 years, 60–69 years, 70–79 years, and 80 years and over.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–June 2010, Sample Adult Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.

Data table for Figure 6.3. Age-adjusted prevalence of obesity among adults aged 20 years and over, by sex and race/ethnicity: United States, January–June 2010

Sex and race/ethnicity	Percent¹	95% confidence interval
Male, Hispanic or Latino	33.0	29.20-36.89
Male, not Hispanic or Latino, single race, white	27.3	25.74-28.94
Male, not Hispanic or Latino, single race, black	35.2	31.05-39.33
Female, Hispanic or Latina	32.5	29.32-35.66
Female, not Hispanic or Latina, single race, white	24.7	23.04-26.35
Female, not Hispanic or Latina, single race, black	42.1	38.97-45.31

¹Estimates for this *Healthy People 2010* Leading Health Indicator are age-adjusted using the projected 2000 U.S. population as the standard population and using five age groups: 20–24 years, 25–34 years, 35–44 years, 45–64 years, and 65 years and over.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–June 2010, Sample Adult Core component. Data are based on household interviews of a sample of the civilian noninstitutionalized population.