


## Receipt of pneumococcal vaccination

**Figure 5.1. Percentage of adults aged 65 and over who had ever received a pneumococcal vaccination: United States, 1997–September 2015**




NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. The analyses exclude those with unknown pneumococcal vaccination status (about 5% of respondents each year). See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997–September 2015, Sample Adult Core component.

- For January–September 2015, the percentage of adults aged 65 and over who had ever received a pneumococcal vaccination was 62.6% (95% confidence interval = 60.92%–64.28%), which was higher than, but not significantly different from the 2014 estimate of 61.3%.
- The percentage of adults aged 65 and over who had ever received a pneumococcal vaccination increased rapidly from 42.4% in 1997 to 53.1% in 2000, and then increased more slowly to 62.6% in January–September 2015.

**Figure 5.2. Percentage of adults aged 65 and over who had ever received a pneumococcal vaccination, by age group and sex: United States, January–September 2015**




NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. The analyses exclude the 4.0% of adults aged 65 and over with unknown pneumococcal vaccination status. See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–September 2015, Sample Adult Core component.

- For both sexes combined, the percentages of adults by age group who had ever received a pneumococcal vaccination were 59.0% for persons aged 65–74 and 67.8% for persons aged 75 and over.
- For both sexes combined, adults aged 75 and over were more likely than those aged 65–74 to have ever received a pneumococcal vaccination. This pattern was found in both men and women.

**Figure 5.3. Percentage of adults aged 65 and over who had ever received a pneumococcal vaccination, by race/ethnicity: United States, January–September 2015**



NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. The analyses exclude the 4.0% of adults aged 65 and over with unknown pneumococcal vaccination status. See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–September 2015, Sample Adult Core component.

- The percentages of adults aged 65 and over by race/ethnicity who had ever received a pneumococcal vaccination were 41.1% for Hispanic persons, 67.1% for non-Hispanic white persons, and 49.4% for non-Hispanic black persons.
- Of the three race/ethnicity groups, non-Hispanic white adults were more likely to have ever received a pneumococcal vaccination compared with Hispanic adults and non-Hispanic black adults.

## Data tables for Figures 5.1–5.3:

### Data table for Figure 5.1. Percentage of adults aged 65 and over who had ever received a pneumococcal vaccination: United States, 1997–September 2015

Year	Crude <sup>1</sup> percent (95% confidence interval)	Age-adjusted <sup>2</sup> percent (95% confidence interval)
1997	42.4 (40.9-43.9)	42.6 (41.1-44.1)
1998	46.0 (44.5-47.5)	46.3 (44.8-47.8)
1999	49.7 (48.1-51.3)	49.9 (48.3-51.5)
2000	53.1 (51.5-54.7)	53.4 (51.8-55.0)
2001	54.0 (52.4-55.6)	54.2 (52.6-55.7)
2002	56.0 (54.5-57.5)	56.2 (54.8-57.7)
2003	55.6 (54.1-57.1)	55.7 (54.2-57.1)
2004	56.8 (55.3-58.3)	57.0 (55.5-58.5)
2005	56.2 (54.82-57.64)	56.3 (54.95-57.72)
2006	57.1 (55.25-58.88)	57.2 (55.43-59.04)
2007	57.7 (55.68-59.69)	57.8 (55.78-59.80)
2008	59.6 (57.68-61.48)	59.8 (57.98-61.64)
2009	60.6 (58.74-62.51)	61.0 (59.15-62.79)
2010	59.4 (57.61-61.17)	59.8 (58.02-61.51)
2011	62.3 (60.55-63.95)	62.7 (61.04-64.31)
2012	59.9 (58.29-61.49)	60.4 (58.86-62.01)
2013	59.7 (58.02-61.36)	60.4 (58.77-62.05)
2014	61.3 (59.88-62.80)	62.3 (60.82-63.70)
January–September 2015	62.6 (60.92-64.28)	63.2 (61.50-64.86)

<sup>1</sup>Crude estimates are presented.

<sup>2</sup>Estimates are age-adjusted using the projected 2000 U.S. population as the standard population and two age groups: 65–74 and 75 and over.

NOTES: Data are based on household interviews of a sample of the civilian noninstitutionalized population. Beginning with 2012 data, the National Health Interview Survey transitioned to weights derived from the 2010 census. For 2003–2011 data, weights were derived from the 2000 census. In this Early Release, estimates for 2000–2002 were recalculated using weights derived from the 2000 census. For 1997–1999 data, weights were derived from the 1990 census. See [Technical Notes](#) for more details.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, 1997–September 2015, Sample Adult Core component.

**Data table for Figure 5.2. Percentage of adults aged 65 and over who had ever received a pneumococcal vaccination, by age group and sex: United States, January–September 2015**

Age (years) and sex	Percent	95% confidence interval
65–74, total	59.0	56.70-61.31
65–74, male	57.9	54.25-61.65
65–74, female	59.9	56.94-62.90
75 and over, total	67.8	65.24-70.27
75 and over, male	66.2	62.07-70.31
75 and over, female	68.9	66.03-71.71
65 and over (crude <sup>1</sup> ), total	62.6	60.92-64.28
65 and over (crude <sup>1</sup> ), male	61.1	58.35-63.89
65 and over (crude <sup>1</sup> ), female	63.8	61.71-65.88
65 and over (age-adjusted <sup>2</sup> ), total	63.2	61.50-64.86
65 and over (age-adjusted <sup>2</sup> ), male	61.9	59.14-64.63
65 and over (age-adjusted <sup>2</sup> ), female	64.2	62.15-66.24

<sup>1</sup>Crude estimates are presented.

<sup>2</sup>Estimates are age-adjusted using the projected 2000 U.S. population as the standard population and two age groups: 65–74 and 75 and over.

NOTE: Data are based on household interviews of a sample of the civilian noninstitutionalized population.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–September 2015, Sample Adult Core component.

**Data table for Figure 5.3. Percentage of adults aged 65 and over who had ever received a pneumococcal vaccination, by race/ethnicity: United States, January–September 2015**

Race/ethnicity	Crude <sup>1</sup> percent (95% confidence interval)	Age-adjusted <sup>2</sup> percent (95% confidence interval)
Hispanic or Latino	41.1 (36.05-46.15)	41.7 (36.75-46.70)
Not Hispanic or Latino, single race, white	67.1 (65.19-68.97)	67.6 (65.65-69.47)
Not Hispanic or Latino, single race, black	49.4 (45.00-53.80)	49.7 (45.24-54.17)

<sup>1</sup>Crude estimates are presented.

<sup>2</sup>Estimates are age-adjusted using the projected 2000 U.S. population as the standard population and two age groups: 65–74 and 75 and over.

NOTE: Data are based on household interviews of a sample of the civilian noninstitutionalized population.

DATA SOURCE: CDC/NCHS, National Health Interview Survey, January–September 2015, Sample Adult Core component.