

**Table 13. National Tuberculosis Genotyping Surveillance Coverage¹:
United States, 2004–2016**

Year	Reported TB cases	Reported culture positive cases	Cases with genotype result	Genotype surveillance coverage
	No.	No.	No.	(%)
2004	14,499	11,327	5,954	(52.6)
2005	14,065	10,957	7,500	(68.4)
2006	13,728	10,745	7,529	(70.1)
2007	13,281	10,425	8,428	(80.8)
2008	12,890	10,021	8,178	(81.6)
2009	11,517	8,883	7,718	(86.9)
2010	11,157	8,457	7,748	(91.6)
2011	10,509	8,087	7,620	(94.2)
2012	9,940	7,628	7,236	(94.9)
2013	9,561	7,368	7,063	(95.9)
2014	9,398	7,232	6,991	(96.7)
2015	9,547	7,408	7,194	(97.1)
2016	9,272	7,162	6,903	(96.4)

¹Genotype surveillance coverage is defined as the percentage of all culture positive tuberculosis (TB) cases for which there was a genotyped isolate.

Note: This table reflects genotyping surveillance coverage for the 50 states and the District of Columbia; for genotyping surveillance coverage of the U.S.-affiliated areas, please see Table 14.

See Surveillance Slide #33.