



Promoting Preventive Services for Adults 50-64: Community and Clinical Partnerships

State-by-State Percentages for Key Indicators

State-by-State Percentages for Key Indicators

	Healthy People 2010* (%)	AL	AK	AZ	AR	CA	CO	CT	DE	D.C.	FL	GA	HI	ID	IL	IN	IA	KS	KY	LA	ME	MD	MA	MI	MN	MS	MO	MT	NE	NV	NH	NJ	NM	NY	NC	ND	OH	OK	OR	PA	RI	SC	SD	TN	TX	UT	VT	VA	WA	WV	WI	WY	Summary Data (%)
SCREENING AND COUNSELING (%)																																																					
Mammogram within past two years*	≥70	75.9	75.6	81.0	75.0	83.5	75.4	88.1	87.1	82.9	80.6	84.1	83.3	71.4	78.4	75.8	80.3	79.2	78.0	80.1	85.3	80.1	89.6	84.0	80.2	71.0	78.3	74.0	77.1	70.3	86.2	80.0	73.6	84.7	82.2	81.6	79.8	71.3	82.2	81.0	86.8	79.4	80.2	77.5	74.4	72.4	82.5	80.4	79.9	78.8	72.4	80.3	
Pap test within past three years†*	≥90	84.4	86.1	81.8	79.7	87.2	86.7	88.6	—	89.6	83.3	87.1	86.4	80.5	86.4	81.1	84.3	87.0	83.2	79.2	90.7	86.8	92.5	89.1	88.0	77.7	81.1	84.2	84.3	72.2	90.5	83.0	84.2	87.5	87.3	86.2	83.9	75.6	87.6	85.1	93.3	85.4	86.9	84.0	82.8	78.9	89.1	91.4	87.5	82.8	85.6	81.2	85.5
Colorectal cancer screening**	≥50	49.8	44.4	50.2	44.1	51.9	51.8	61.0	64.3	60.3	55.9	55.5	49.2	44.1	48.6	48.0	54.7	52.3	53.5	48.6	65.5	61.0	66.8	58.1	57.9	47.3	50.9	45.7	49.0	42.5	62.7	53.9	45.1	56.7	57.7	44.8	50.8	43.7	54.6	53.8	59.8	56.9	50.7	54.1	46.3	54.6	59.0	59.4	55.0	48.5	53.6	44.1	53.3
Cholesterol screening within past five years§	≥80	90.3	89.3	87.5	85.6	90.0	90.2	93.5	95.3	92.1	89.1	92.0	88.7	83.8	88.8	90.7	89.8	89.4	87.0	88.5	91.6	90.4	94.2	92.0	90.1	86.3	87.0	85.2	87.6	86.8	91.7	92.1	86.4	91.3	91.7	88.3	89.9	85.8	89.0	88.4	92.9	91.2	87.8	90.3	86.6	87.3	90.2	92.6	89.9	90.1	88.5	89.4	89.7
IMMUNIZATIONS (%)																																																					
Influenza vaccination within past year*	≥60	41.8	43.0	39.4	44.9	39.5	48.6	45.7	46.7	44.2	32.4	38.6	47.4	39.5	38.6	39.9	48.2	43.5	43.6	43.6	47.3	46.3	45.8	41.8	50.4	38.4	45.1	40.9	51.3	29.6	49.4	39.9	42.3	43.9	47.3	45.4	42.0	51.3	43.2	43.2	49.9	42.9	53.6	42.7	42.1	48.6	46.9	48.0	44.2	46.5	44.4	44.8	42.3
Pneumococcal immunization ever among persons at risk¶*	≥60	32.9	39.3	35.4	31.4	29.0	37.9	27.8	29.0	31.5	30.4	33.0	30.5	31.4	26.4	34.2	36.2	31.8	33.4	36.9	35.0	35.2	35.7	35.7	32.1	33.4	34.4	37.5	35.5	34.2	37.8	27.6	33.3	30.0	34.5	37.7	33.0	39.7	40.6	35.5	32.6	38.9	32.9	33.6	33.2	36.4	37.3	33.9	38.1	41.7	33.2	35.8	32.9
UP TO DATE WITH SELECT CLINICAL PREVENTIVE SERVICES (%)																																																					
Women: Influenza vaccination and breast, cervical and colorectal cancer screenings*	—	24.2	21.9	22.8	21.6	25.8	29.2	31.2	—	28.5	21.1	26.0	27.0	22.5	19.8	21.6	29.7	27.0	25.1	23.5	34.9	30.0	32.0	28.3	32.8	19.5	24.5	21.4	29.7	14.4	33.2	24.7	20.5	25.7	31.8	25.2	24.8	24.2	27.3	25.6	32.8	27.8	32.5	25.6	24.7	28.1	29.9	34.7	27.4	24.3	28.9	22.0	25.9
Men: Influenza vaccination and colorectal cancer screening*	—	22.3	25.2	27.5	27.4	22.7	29.6	32.9	30.5	30.5	23.3	22.6	27.9	22.8	23.2	22.9	29.4	25.9	29.2	25.4	35.2	32.6	35.2	27.3	32.7	23.0	24.1	23.8	28.5	16.9	34.3	26.3	23.4	31.6	30.4	22.0	25.0	26.6	27.3	26.1	30.7	28.9	30.2	25.5	24.5	31.5	32.9	29.3	28.9	26.9	26.3	23.2	26.5
RISK FACTORS (%)																																																					
No leisure-time physical activity within past month*	≤20	33.9	27.3	21.1	33.8	23.6	19.0	23.4	23.9	23.0	27.1	28.5	20.1	23.5	29.7	32.0	26.4	27.6	34.5	31.7	23.0	24.8	23.3	25.0	20.0	35.4	31.1	22.9	25.0	29.8	22.8	26.8	24.2	29.6	27.9	26.5	28.3	35.5	20.7	28.3	25.5	28.4	26.8	32.0	32.4	21.9	19.8	23.5	18.6	33.7	23.5	26.5	27.1
Smoking - current*	≤12	23.7	19.0	17.1	21.3	13.5	14.8	13.9	19.4	18.3	19.2	20.5	15.3	15.7	21.3	24.4	18.9	17.1	23.7	18.7	16.1	16.1	15.3	18.4	18.7	22.9	21.4	19.0	17.1	23.1	16.6	16.3	18.4	18.4	20.5	17.1	19.5	25.7	14.7	20.3	16.2	18.5	17.8	24.3	19.4	9.9	14.2	15.1	14.8	22.7	18.4	18.3	18.4
Binge drinking within past 30 days*	≤13.4¶	8.7	12.8	10.6	8.8	11.4	9.9	11.6	13.7	8.5	10.2	8.7	13.3	9.7	12.9	11.1	14.3	8.9	7.0	9.5	10.8	9.6	12.2	12.9	14.4	8.0	10.3	13.3	13.3	13.5	10.9	10.3	9.3	9.9	8.7	13.9	10.9	8.2	9.8	10.1	12.5	8.8	12.0	7.9	11.0	6.2	10.6	7.6	11.1	5.0	15.8	11.5	10.6
Obesity – current*	≤15	35.4	30.9	26.3	36.4	31.1	23.1	24.7	33.6	28.4	30.3	32.1	24.9	30.6	32.2	36.5	31.7	35.0	36.0	35.0	31.2	32.4	24.8	34.5	30.1	36.9	34.0	26.1	32.6	31.6	27.4	28.7	27.7	29.1	34.5	33.3	33.8	36.1	28.9	33.2	27.3	35.7	33.4	35.3	32.5	31.6	25.3	32.7	30.8	38.3	28.9	30.5	31.7
High blood pressure ever§	—	49.3	35.4	38.0	44.8	36.4	33.4	33.4	42.3	44.2	38.1	47.1	39.8	37.9	41.4	43.1	38.6	36.9	46.3	47.3	38.6	41.6	36.8	40.9	32.9	51.1	44.6	33.5	40.5	41.0	37.5	38.0	36.4	41.0	44.0	35.7	39.5	46.0	37.8	39.5	41.0	43.7	35.8	48.1	43.1	33.7	32.2	42.3	36.3	44.4	39.1	34.7	40.4
Moderate depressive symptoms within past two weeks*	—	13.7	5.7	—	11.6	10.9	—	6.3	8.3	6.3	8.1	9.9	7.6	—	—	10.7	5.1	6.5	—	9.0	5.4	7.0	—	10.3	6.1	13.6	10.6	6.8	6.3	9.9	7.2	—	10.8	—	—	6.0	—	12.2	6.9	—	9.0	9.9	—	11.4	10.7	8.7	6.0	5.1	6.1	15.2	6.1	8.7	9.3

* Based on 2008 data from 50 states and the District of Columbia

† Among women with intact cervix

‡ Had home blood stool test within past year or colonoscopy or sigmoidoscopy within past 10 years

§ Based on 2007 data from 50 states and the District of Columbia

¶ Smoke currently or have diabetes, asthma or cardiovascular disease

‡ Reflects change from original target after Midpoint Review

Based on 2006 data from 38 states and the District of Columbia

The cornerstone of this Report is 14 key indicators for monitoring the use of clinical preventive services in adults aged 50 to 64. The indicators are organized into four categories: screenings, immunizations, up to date with select clinical preventive services, and risk factors. The first three categories focus on services that need to be promoted. The final category includes risk factors that should be prevented or decreased. Ten indicators have Healthy People 2010 (HP2010) targets for the adult population, which is a set of health objectives for the nation (www.healthypeople.gov).

These state-by-state data serve as a baseline and the Behavioral Risk Factor Surveillance System offers a reliable source to track progress. Please note confidence intervals should be used if comparing states because differences may not be significant if the confidence intervals for two states overlap. See appendices in the full Report for detailed information on HP2010, methods, and state-by-state data with confidence intervals.

These data can help practitioners identify gaps and barriers and monitor trends over time. The data can also help build the case for efforts to enhance clinical preventive services through promoting public programs and policies and health system change.



Suggested Citation: Centers for Disease Control and Prevention, AARP, and the American Medical Association. *Promoting Preventive Services for Adults 50-64: Community and Clinical Partnerships*. Atlanta, GA: National Association of Chronic Disease Directors; 2009. Available at www.cdc.gov/aging