

School Health Profiles (Profiles)

2010 State Results

National Center for Chronic Disease Prevention and Health Promotion
Division of Adolescent and School Health



This slide set contains information about and data from the School Health Profiles or Profiles. It includes state data from 2010.

What is the School Health Profiles (Profiles)?

- Profiles is a system of surveys assessing school health policies and practices in states, territories, large urban school districts, and tribal governments
- Profiles surveys are conducted biennially (during even-numbered years) by state, territorial, local, and tribal government education and health agencies with technical assistance from CDC
- Profiles questionnaires are administered to middle and high school principals and lead health education teachers

National Center for Chronic Disease Prevention and Health Promotion
Division of Adolescent and School Health



What is the School Health Profiles (Profiles)?

- Profiles is a system of surveys assessing school health policies and practices in states, territories, large urban school districts, and tribal governments.
- Profiles surveys are conducted biennially (during even-numbered years) by state, territorial, local, and tribal government education and health agencies with technical assistance from CDC.
- Profiles questionnaires are administered to middle and high school principals and lead health education teachers.

Topics Monitored by Profiles

- School health education requirements and content
- Physical education requirements
- School health policies related to HIV infection/AIDS, tobacco-use prevention, and nutrition
- Asthma management activities
- Family and community involvement in school health programs

National Center for Chronic Disease Prevention and Health Promotion
Division of Adolescent and School Health



The topics monitored by Profiles are:

- School health education requirements and content
- Physical education requirements
- School health policies related to HIV infection/AIDS, tobacco-use prevention, and nutrition
- Asthma management activities
- Family and community involvement in school health programs

Profiles Participation, 1998 – 2010


	1998	2000	2002	2004	2006	2008	2010
# of states	38	38	43	41	44	50	50
# of cities	10	13	14	13	16	21	20
# of territories	--	--	--	--	--	5	6
# of tribal governments	--	--	--	--	--	2	2
Total # of sites	48	51	57	54	60	78	78
Total % of sites with weighted data	77%	75%	77%	70%	82%	94%	99%

National Center for Chronic Disease Prevention and Health Promotion
Division of Adolescent and School Health



This slide contains state and local Profiles participation data from 1998 through 2010.

- In 1998, 38 states and 10 cities (including Washington, D.C.) participated in Profiles, for a total number of 48 states and cities. The percent of sites with weighted data in 1998 was 77%.
- In 2000, 38 states and 13 cities (including Washington, D.C.) participated in Profiles, for a total number of 51 states and cities. The percent of sites with weighted data in 2000 was 75%.
- In 2002, 43 states and 14 cities (including Washington, D.C.) participated in Profiles, for a total number of 57 states and cities. The percent of sites with weighted data in 2002 was 77%.
- In 2004, 41 states and 13 cities (including Washington, D.C.) participated in Profiles, for a total number of 54 states and cities. The percent of sites with weighted data in 2004 was 70%.
- In 2006, 44 states and 16 cities (including Washington, D.C.) participated in Profiles, for a total number of 60 states and cities. The percent of sites with weighted data in 2006 was 82%.
- In 2008, 50 states and 21 cities (including Washington, D.C.) participated in Profiles, for a total number of 71 states and cities. The percent of sites with weighted data in 2008 was 94%.
- In 2010, 50 states and 20 cities (including Washington, D.C.) participated in Profiles, for a total number of 70 states and cities. The percent of sites with weighted data in 2010 was 99%.



This slide shows the states within the U.S. that participated in the 2010 School Health Profiles. The states that participated and had weighted results include: Washington, Oregon, California, Idaho, Nevada, Montana, Wyoming, Utah, Arizona, New Mexico, Alaska, Hawaii, Colorado, North Dakota, South Dakota, Nebraska, Kansas, Oklahoma, Texas, Minnesota, Iowa, Missouri, Arkansas, Wisconsin, Louisiana, Mississippi, Michigan, Indiana, Kentucky, Tennessee, Alabama, Georgia, Ohio, West Virginia, Virginia, North Carolina, South Carolina, Florida, Maryland, Delaware, New Jersey, Pennsylvania, New York, Vermont, Connecticut, Rhode Island, Massachusetts, New Hampshire, and Maine. Illinois is the only state with unweighted results.

Uses of Profiles Data

Education and health officials use Profiles data to

- Describe school health policies and practices and compare them across jurisdictions
- Identify professional development needs
- Plan and monitor programs
- Support health-related policies and legislation
- Seek funding
- Garner support for future surveys

National Center for Chronic Disease Prevention and Health Promotion
Division of Adolescent and School Health



Education and health officials use Profiles data to

- Describe school health policies and practices and compare them across jurisdictions
- Identify professional development needs
- Plan and monitor programs
- Support health-related policies and legislation
- Seek funding
- Garner support for future surveys

Examples of Uses of Profiles Data

→ Michigan

- Used Profiles data to write a state Senate bill requiring health and physical education in each of grades K-8

→ North Carolina

- Used Profiles data along with other data sources to develop objectives and standards for the state's health improvement plan

National Center for Chronic Disease Prevention and Health Promotion
Division of Adolescent and School Health



Examples of uses of Profiles data are:

Michigan

Used Profiles data to write a state Senate bill requiring health and physical education in each of grades K-8.

North Carolina

Used Profiles data along with other data sources to develop objectives and standards for the state's health improvement plan.

Examples of Uses of Profiles Data

→ Delaware

- Used Profiles and other data sources to identify professional development needs related to sexual health curricula and to target professional development efforts to that area.

National Center for Chronic Disease Prevention and Health Promotion
Division of Adolescent and School Health



Another example of uses of Profiles data is:

Delaware

Used Profiles and other data sources to identify professional development needs related to sexual health curricula and to target professional development efforts to that area.

Profiles Information

<http://www.cdc.gov/schoolhealthprofiles>

Background on the School Health Profiles

Questionnaires and item rationales

Comprehensive results

Fact sheets

Publications and journal articles

Participation history

National Center for Chronic Disease Prevention and Health Promotion
Division of Adolescent and School Health



For more information regarding Profiles please go to the following website:
<http://www.cdc.gov/schoolhealthprofiles>

This website will provide you with information regarding:


- Background on the School Health Profiles
- Questionnaires and item rationales
- Comprehensive results
- Fact sheets
- Publications and journal articles
- Participation history

Physical Activity And Physical Education

National Center for Chronic Disease Prevention and Health Promotion
Division of Adolescent and School Health




Physical Activity and Physical Education




This slide shows the percentage of secondary schools in which students could not be exempted from taking required physical education for certain reasons*. The values range from 15% to 89%. Alaska, Arkansas, Colorado, Florida, Georgia, Michigan, Mississippi, Nevada, Oklahoma, South Carolina, Texas, Washington, range from 15% to 44%. Alabama, Arizona, California, Hawaii, Idaho, Iowa, Kentucky, Louisiana, New Mexico, Oregon, Tennessee, Utah, range from 45% to 64%. Indiana, Kansas, Maryland, Massachusetts, Missouri, Nebraska, New Hampshire, North Carolina, North Dakota, Ohio, Vermont, Wyoming, range from 65% to 77%. Connecticut, Delaware, Maine, Minnesota, Montana, New Jersey, New York, Pennsylvania, Rhode Island, South Dakota, Virginia, West Virginia, Wisconsin, range from 78% to 89%. Illinois did not have weighted results.


*Enrollment in other courses, participation in school sports, participation in other school activities, participation in community sports activities, high physical fitness competency test score, participation in vocational training, and participation in community service activities.




This slide shows the percentage of secondary schools that taught 12 key physical activity topics in a required course. The values range from 39% to 75%. Alaska, Arizona, Connecticut, Iowa, Kansas, Maine, Massachusetts, Minnesota, Oklahoma, Oregon, Vermont, range from 39% to 50%. California, Michigan, Nevada, New Hampshire, North Dakota, Ohio, Pennsylvania, Rhode Island, South Dakota, Tennessee, Washington, Wisconsin, Wyoming, range from 51% to 56%. Georgia, Hawaii, Idaho, Indiana, Louisiana, Maryland, Montana, Nebraska, New Jersey, North Carolina, Utah, Wyoming, range from 57% to 62%. Alabama, Arkansas, Delaware, Florida, Kentucky, Mississippi, Missouri, New York, South Carolina, Texas, Virginia, West Virginia, range from 63% to 75%. Colorado, Illinois and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that taught about developing an individualized physical activity plan in a required course. The values range from 50% to 87%. Alaska, Arizona, Connecticut, Kansas, Massachusetts, Michigan, Nevada, New Hampshire, Oklahoma, South Dakota, Tennessee, range from 50% to 71%. Alabama, California, Iowa, Louisiana, Minnesota, Montana, Nebraska, North Dakota, Ohio, Pennsylvania, Vermont, Wisconsin, range from 72% to 76%. Delaware, Florida, Georgia, Hawaii, Idaho, Maine, Maryland, New Jersey, North Carolina, Oregon, Rhode Island, Texas, range from 77% to 79%. Arkansas, Indiana, Kentucky, Mississippi, Missouri, New York, South Carolina, Utah, Virginia, Washington, West Virginia, Wyoming, range from 80% to 87%. Colorado, Illinois and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools in which those who teach physical education were provided with key materials for teaching physical education. The values range from 39% to 94%. Alabama, Alaska, Arizona, Idaho, Indiana, Nebraska, North Dakota, Ohio, Oklahoma, Oregon, South Dakota, Washington, range from 39% to 62%. Arkansas, California, Colorado, Iowa, Kansas, Louisiana, Michigan, Minnesota, Mississippi, Montana, New Mexico, Rhode Island, range from 63% to 71%. Hawaii, Kentucky, Maine, Massachusetts, Missouri, Nevada, New Hampshire, North Carolina, Tennessee, Texas, Vermont, Wyoming, range from 72% to 77%. Connecticut, Delaware, Florida, Georgia, Maryland, New Jersey, New York, Pennsylvania, South Carolina, Utah, Virginia, West Virginia, Wisconsin, range from 78% to 94%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools in which at least one physical education teacher or specialist received professional development on physical education during the 2 years before the survey. The values range from 38% to 98%. Alaska, Arizona, Idaho, Indiana, Kentucky, Nebraska, New Mexico, North Dakota, Ohio, Oklahoma, Oregon, South Dakota, range from 38% to 80%. Alabama, California, Hawaii, Iowa, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Montana, Washington, Wyoming, range from 81% to 87%. Colorado, Georgia, Kansas, Massachusetts, Nevada, North Carolina, Pennsylvania, Rhode Island, Tennessee, Utah, West Virginia, range from 88% to 92%. Arkansas, Connecticut, Delaware, Florida, Maine, Maryland, New Hampshire, New Jersey, New York, South Carolina, Texas, Vermont, Virginia, Wisconsin, range from 93% to 98%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools that offered opportunities for all students to participate in intramural activities or physical activity clubs. The values range from 41% to 87%. Alabama, Arkansas, Indiana, Iowa, Kansas, Louisiana, Nebraska, North Dakota, Ohio, Oklahoma, South Dakota, Texas, range from 41% to 56%. Georgia, Idaho, Kentucky, Michigan, Mississippi, Montana, New Mexico, North Carolina, South Carolina, Tennessee, West Virginia, Wyoming, range from 57% to 62%. Arizona, Colorado, Connecticut, Delaware, Minnesota, Missouri, Oregon, Rhode Island, Utah, Virginia, Washington, Wisconsin, range from 63% to 75%. Alaska, California, Florida, Hawaii, Maine, Maryland, Massachusetts, Nevada, New Hampshire, New Jersey, New York, Pennsylvania, Vermont, range from 76% to 87%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools in which children or adolescents use the school's indoor physical activity or athletic facilities for community-sponsored physical activity classes or lessons. The values range from 55% to 97%. Alabama, Arizona, Arkansas, Florida, Georgia, Louisiana, Mississippi, Nevada, North Carolina, Ohio, South Carolina, Tennessee, range from 55% to 71%. Delaware, Hawaii, Idaho, Indiana, Kentucky, Michigan, Missouri, New Mexico, North Dakota, Oklahoma, Rhode Island, Texas, range from 72% to 80%. California, Connecticut, Iowa, Kansas, Maine, Massachusetts, Montana, New Hampshire, Oregon, Pennsylvania, South Dakota, Wyoming, range from 81% to 86%. Alaska, Colorado, Maryland, Minnesota, Nebraska, New Jersey, New York, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, range from 87% to 97%. Illinois did not have weighted results.



This slide shows the percentage of secondary schools that provided parents and families with health information to increase parent and family knowledge of physical activity. The values range from 32% to 61%. Alaska, Georgia, Hawaii, Indiana, Iowa, Massachusetts, Minnesota, Oregon, South Dakota, Utah, range from 32% to 40%. Alabama, Arizona, California, Idaho, Louisiana, Michigan, Montana, Nevada, New Hampshire, Ohio, Pennsylvania, Rhode Island, Washington, range from 41% to 44%. Connecticut, Florida, Kansas, Maine, Missouri, Nebraska, New Jersey, North Dakota, Oklahoma, Texas, Vermont, West Virginia, range from 45% to 47%. Arkansas, Delaware, Kentucky, Maryland, Mississippi, New York, North Carolina, South Carolina, Tennessee, Virginia, Wisconsin, Wyoming, range from 48% to 61%. Colorado, Illinois and New Mexico did not have weighted results.




Tobacco Use




This slide shows the percentage of secondary schools that prohibited all tobacco use at all times in all locations*. The values range from 24% to 79%. Alaska, Idaho, Kansas, Kentucky, Missouri, Nebraska, North Dakota, Ohio, Oklahoma, South Dakota, Tennessee, Wyoming, range from 24% to 49%. Connecticut, Florida, Indiana, Michigan, Minnesota, Montana, Nevada, New Jersey, New Mexico, New York, Virginia, Wisconsin, range from 50% to 56%. Arizona, Arkansas, Colorado, Iowa, Louisiana, Massachusetts, New Hampshire, Oregon, Pennsylvania, Rhode Island, Utah, Vermont, range from 57% to 62%. Alabama, California, Delaware, Georgia, Hawaii, Maine, Maryland, Mississippi, North Carolina, South Carolina, Texas, Washington, West Virginia, range from 63% to 79%. Illinois did not have weighted results.


*Prohibited the use of all tobacco, including cigarettes, smokeless tobacco, cigars, and pipes, by students, faculty and school staff, and visitors, in school buildings, outside on school grounds, on school buses or other vehicles used to transport students, and at off-campus, school-sponsored events, during school hours and non-school hours.




This slide shows the percentage of secondary schools that implement a tobacco-free environment policy addressing 7 key elements. The values range from 5% to 34%. Alabama, Arkansas, Connecticut, Florida, Georgia, Kentucky, Maryland, Michigan, Missouri, Nevada, Texas, range from 5% to 10%. Colorado, Hawaii, Kansas, Massachusetts, Mississippi, Nebraska, New York, North Carolina, Oklahoma, South Carolina, South Dakota, Tennessee, Virginia, range from 11% to 13%. Arizona, Delaware, Indiana, Iowa, Louisiana, New Mexico, North Dakota, Ohio, Oregon, Rhode Island, Utah, range from 14% to 16%. Alaska, California, Idaho, Maine, Minnesota, Montana, New Hampshire, New Jersey, Pennsylvania, Vermont, Washington, West Virginia, Wisconsin, Wyoming, range from 17% to 34%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools that sometimes, almost always, or always required students who were caught smoking to participate in an assistance, education, or cessation program. The values range from 17% to 86%. Alabama, Arkansas, Georgia, Iowa, Kansas, Louisiana, Mississippi, Missouri, Nebraska, Oklahoma, South Dakota, Texas, range from 17% to 31%. Arizona, Colorado, Connecticut, Delaware, Hawaii, Massachusetts, Michigan, Minnesota, New York, North Dakota, Tennessee, Wisconsin, range from 32% to 41%. Alaska, Florida, Indiana, Kentucky, Nevada, New Jersey, New Mexico, North Carolina, Ohio, Pennsylvania, South Carolina, Virginia, range from 42% to 49%. California, Idaho, Maine, Maryland, Montana, New Hampshire, Oregon, Rhode Island, Utah, Vermont, Washington, West Virginia, Wyoming, range from 50% to 86%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools that provided tobacco cessation services for students, faculty, and staff at school or through arrangements with providers not on school property. The values range from 14% to 43%. Arizona, Arkansas, Connecticut, Georgia, Hawaii, Kansas, Louisiana, Michigan, Nebraska, Nevada, Oklahoma, Texas, range from 14% to 19%. California, Colorado, Delaware, Kentucky, Maryland, Minnesota, Mississippi, Missouri, Montana, New York, South Dakota, Virginia, range from 20% to 25%. Alaska, Idaho, Indiana, Iowa, Massachusetts, New Hampshire, New Jersey, New Mexico, Ohio, Oregon, Tennessee, Wisconsin, range from 26% to 31%. Alabama, Florida, Maine, North Carolina, North Dakota, Pennsylvania, Rhode Island, South Carolina, Utah, Vermont, Washington, West Virginia, Wyoming, range from 32% to 43%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools that posted signs marking a tobacco-free school zone. The values range from 41% to 95%. Connecticut, Delaware, Idaho, Kentucky, Michigan, Missouri, Nebraska, Nevada, North Dakota, South Carolina, Utah, Vermont, range from 41% to 72%. Florida, Indiana, Kansas, Massachusetts, Minnesota, New Jersey, New Mexico, New York, Ohio, Pennsylvania, Rhode Island, Virginia, range from 73% to 79%. Alabama, Alaska, Georgia, Hawaii, Maine, Maryland, Oklahoma, Oregon, South Dakota, Texas, Wisconsin, Wyoming, range from 80% to 86%. Arizona, Arkansas, California, Colorado, Iowa, Louisiana, Mississippi, Montana, New Hampshire, North Carolina, Tennessee, Washington, West Virginia, range from 87% to 95%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools that taught 15 key tobacco-use prevention topics in a required course. The values range from 34% to 78%. Alaska, Arizona, California, Connecticut, Maine, Massachusetts, Minnesota, Oklahoma, Oregon, South Dakota, Vermont, range from 34% to 45%. Florida, Hawaii, Iowa, Kansas, Michigan, Nebraska, New Hampshire, Pennsylvania, Rhode Island, Utah, Washington, Wisconsin, range from 46% to 52%. Alabama, Georgia, Idaho, Indiana, Maryland, Montana, North Carolina, North Dakota, Ohio, South Carolina, Tennessee, Wyoming, range from 53% to 56%. Arkansas, Delaware, Kentucky, Louisiana, Mississippi, Missouri, Nevada, New Jersey, New York, Texas, Virginia, West Virginia, range from 57% to 78%. Colorado, Illinois and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools in which the lead health education teacher received professional development during the 2 years before the survey on tobacco-use prevention. The values range from 17% to 51%. Alaska, Connecticut, Indiana, Iowa, Massachusetts, Nebraska, Ohio, Oregon, Pennsylvania, Rhode Island, South Dakota, range from 17% to 27%. Arizona, California, Georgia, Kansas, Kentucky, Maine, Michigan, Minnesota, Missouri, Vermont, Virginia, Washington, range from 28% to 32%. Alabama, Delaware, Florida, Idaho, Montana, New Hampshire, New Jersey, North Dakota, South Carolina, Texas, Wisconsin, Wyoming, range from 33% to 40%. Arkansas, Hawaii, Louisiana, Maryland, Mississippi, Nevada, New York, North Carolina, Oklahoma, Tennessee, Utah, West Virginia, range from 41% to 51%. Colorado, Illinois and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that provided parents and families of students with health information to increase parent and family knowledge of tobacco-use prevention. The values range from 24% to 53%. Connecticut, Georgia, Indiana, Iowa, Massachusetts, Minnesota, New Hampshire, Oregon, Pennsylvania, Rhode Island, Utah, range from 24% to 31%. Arizona, Kansas, Maine, Michigan, Missouri, Nebraska, Nevada, Ohio, South Carolina, South Dakota, Virginia, Wisconsin, range from 32% to 33%. Alaska, California, Florida, Hawaii, Idaho, Louisiana, Montana, New Jersey, North Carolina, Texas, Washington, Wyoming, range from 34% to 40%. Alabama, Arkansas, Delaware, Kentucky, Maryland, Mississippi, New York, North Dakota, Oklahoma, Tennessee, Vermont, West Virginia, range from 41% to 53%. Colorado, Illinois and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that used the School Health Index or a similar self-assessment tool to assess their policies, activities, and programs in tobacco-use prevention. The values range from 25% to 75%. Alaska, Arizona, Iowa, Minnesota, Nebraska, Nevada, North Carolina, Ohio, Oregon, Pennsylvania, Rhode Island, South Dakota, range from 25% to 35%. California, Connecticut, Idaho, Louisiana, Massachusetts, Michigan, New Hampshire, New Jersey, South Carolina, Texas, Virginia, Wyoming, range from 36% to 42%. Alabama, Colorado, Florida, Indiana, Kansas, Kentucky, Missouri, New Mexico, New York, North Dakota, Utah, Washington, range from 43% to 50%. Arkansas, Delaware, Georgia, Hawaii, Maine, Maryland, Mississippi, Montana, Oklahoma, Tennessee, Vermont, West Virginia, Wisconsin, range from 51% to 75%. Illinois did not have weighted results.




Nutrition



This slide shows the percentage of secondary schools in which students could not purchase chocolate candy from vending machines or at the school store, canteen, or snack bar. The values range from 25% to 100%. Georgia, Idaho, Indiana, Kansas, Louisiana, Michigan, Minnesota, Montana, Nebraska, Oklahoma, South Carolina, Utah, range from 25% to 66%. Colorado, Florida, Maryland, Missouri, North Carolina, North Dakota, Ohio, Tennessee, Virginia, Washington, Wisconsin, Wyoming, range from 67% to 79%. Alaska, Arizona, Arkansas, Delaware, Iowa, Kentucky, New Mexico, Oregon, Pennsylvania, South Dakota, Texas, Vermont, range from 80% to 89%. Alabama, California, Connecticut, Hawaii, Maine, Massachusetts, Mississippi, Nevada, New Hampshire, New Jersey, New York, Rhode Island, West Virginia, range from 90% to 100%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools in which students could not purchase other kinds of candy from vending machines or at the school store, canteen, or snack bar. The values range from 26% to 98%. Georgia, Idaho, Indiana, Kansas, Louisiana, Michigan, Minnesota, Montana, Nebraska, Oklahoma, South Carolina, Utah, range from 26% to 62%. Colorado, Florida, Kentucky, Maryland, Missouri, Nevada, North Carolina, Ohio, Virginia, Washington, Wisconsin, Wyoming, range from 63% to 74%. Alaska, Arizona, Arkansas, Delaware, Iowa, New Mexico, North Dakota, Oregon, Pennsylvania, South Dakota, Tennessee, Texas, range from 75% to 84%. Alabama, California, Connecticut, Hawaii, Maine, Massachusetts, Mississippi, New Hampshire, New Jersey, New York, Rhode Island, Vermont, West Virginia, range from 85% to 98%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools in which students could not purchase salty snacks that are not low in fat* from vending machines or at the school store, canteen, or snack bar. The values range from 32% to 92%. Georgia, Idaho, Indiana, Louisiana, Maryland, Michigan, Minnesota, North Carolina, Ohio, South Carolina, Utah, Washington, range from 32% to 60%. Colorado, Florida, Kansas, Missouri, Montana, Nebraska, New York, Oklahoma, Pennsylvania, Virginia, Wisconsin, Wyoming, range from 61% to 72%. Alaska, Arizona, Arkansas, Iowa, Massachusetts, New Hampshire, New Jersey, New Mexico, South Dakota, Tennessee, Texas, Vermont, range from 73% to 79%. Alabama, California, Connecticut, Delaware, Hawaii, Kentucky, Maine, Mississippi, Nevada, North Dakota, Oregon, Rhode Island, West Virginia, range from 80% to 92%. Illinois did not have weighted results.


*Such as regular potato chips.



This slide shows the percentage of secondary schools in which students could not purchase cookies, crackers, cakes, pastries, or other baked goods that are not low in fat from vending machines or at the school store, canteen, or snack bar. The values range from 30% to 94%. Florida, Georgia, Idaho, Indiana, Kansas, Maryland, Michigan, Minnesota, Montana, Ohio, South Carolina, Utah, range from 30% to 57%. Colorado, Louisiana, Missouri, Nebraska, New York, North Carolina, Oklahoma, Pennsylvania, Vermont, Virginia, Washington, Wisconsin, range from 58% to 67%. Arizona, Arkansas, Iowa, Massachusetts, New Hampshire, New Jersey, North Dakota, Oregon, South Dakota, Tennessee, Texas, Wyoming, range from 68% to 79%. Alabama, Alaska, California, Connecticut, Delaware, Hawaii, Kentucky, Maine, Mississippi, Nevada, New Mexico, Rhode Island, West Virginia, range from 80% to 94%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools in which students could not purchase soda pop or fruit drinks that are not 100% juice from vending machines or at the school store, canteen, or snack bar. The values range from 46% to 97%. Georgia, Idaho, Indiana, Kansas, Louisiana, Minnesota, Missouri, Montana, Nebraska, Oklahoma, South Carolina, Utah, range from 46% to 59%. Arkansas, Florida, Iowa, Kentucky, Maryland, Michigan, North Carolina, North Dakota, South Dakota, Virginia, Washington, Wyoming, range from 60% to 69%. Alabama, Alaska, Arizona, Colorado, New Hampshire, New York, Ohio, Oregon, Pennsylvania, Tennessee, Texas, Wisconsin, range from 70% to 81%. California, Connecticut, Delaware, Hawaii, Maine, Massachusetts, Mississippi, Nevada, New Jersey, New Mexico, Rhode Island, Vermont, West Virginia, range from 82% to 97%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools in which students could not purchase sports drinks* from vending machines or at the school store, canteen, or snack bar. The values range from 27% to 92%. Indiana, Iowa, Kansas, Minnesota, Missouri, Montana, Nebraska, Nevada, South Dakota, Utah, Washington, Wyoming, range from 27% to 37%. California, Florida, Georgia, Idaho, Louisiana, Michigan, North Carolina, North Dakota, Ohio, Oklahoma, South Carolina, Wisconsin, range from 38% to 48%. Colorado, Kentucky, Maine, Maryland, Mississippi, New Hampshire, New Jersey, New York, Pennsylvania, Texas, Vermont, Virginia, range from 49% to 59%. Alabama, Alaska, Arizona, Arkansas, Connecticut, Delaware, Hawaii, Massachusetts, New Mexico, Oregon, Rhode Island, Tennessee, West Virginia, range from 60% to 92%. Illinois did not have weighted results.


*Such as Gatorade.



This slide shows the percentage of secondary schools in which students could not purchase 2% or whole milk from vending machines or at the school store, canteen, or snack bar. The values range from 47% to 94%. Florida, Georgia, Indiana, Maryland, Michigan, Nevada, New Hampshire, Ohio, Pennsylvania, Utah, Vermont, Wisconsin, range from 47% to 59%. California, Idaho, Iowa, Minnesota, Missouri, New Jersey, New York, South Carolina, South Dakota, Texas, Virginia, Washington, range from 60% to 64%. Delaware, Kansas, Kentucky, Massachusetts, Mississippi, Montana, Nebraska, New Mexico, North Carolina, Oklahoma, Tennessee, Wyoming, range from 65% to 72%. Alabama, Alaska, Arizona, Arkansas, Colorado, Connecticut, Hawaii, Louisiana, Maine, North Dakota, Oregon, Rhode Island, West Virginia, range from 73% to 94%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools in which students could not purchase foods or beverages containing caffeine from vending machines or at the school store, canteen, or snack bar. The values range from 51% to 99%. Georgia, Idaho, Indiana, Kansas, Louisiana, Minnesota, Missouri, Montana, Nebraska, Oklahoma, South Carolina, Utah, range from 51% to 62%. Arkansas, Florida, Iowa, Kentucky, Michigan, North Carolina, North Dakota, Ohio, South Dakota, Washington, Wisconsin, Wyoming, range from 63% to 71%. Alabama, Alaska, Arizona, Colorado, Maryland, Mississippi, New Hampshire, Oregon, Pennsylvania, Tennessee, Texas, Virginia, range from 72% to 80%. California, Connecticut, Delaware, Hawaii, Maine, Massachusetts, Nevada, New Jersey, New Mexico, New York, Rhode Island, Vermont, West Virginia, range from 81% to 99%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools that did not sell less nutritious foods and beverages* anywhere outside the school food service program. The values range from 8% to 82%. Idaho, Kansas, Louisiana, Michigan, Minnesota, Missouri, Montana, Nebraska, Nevada, Oklahoma, Utah, Washington, range from 8% to 28%. Florida, Georgia, Indiana, Iowa, New Hampshire, New York, Ohio, South Carolina, South Dakota, Virginia, Wisconsin, Wyoming, range from 29% to 37%. California, Colorado, Kentucky, Maryland, Massachusetts, New Jersey, North Carolina, North Dakota, Oregon, Pennsylvania, Texas, Vermont, range from 38% to 50%. Alabama, Alaska, Arizona, Arkansas, Connecticut, Delaware, Hawaii, Maine, Mississippi, New Mexico, Rhode Island, Tennessee, West Virginia, range from 51% to 82%. Illinois did not have weighted results.

*Baked goods not low in fat, salty snacks not low in fat, chocolate candy, other kinds of candy, and soda pop or fruit drinks that are not 100% juice, and sports drinks




This slide shows the percentage of secondary schools that always offered fruits or non-fried vegetables in vending machines or school stores, and during celebrations when foods and beverages are offered. The values range from 1% to 23%. Alabama, Alaska, Arkansas, Hawaii, Kansas, Kentucky, Montana, Nebraska, North Dakota, Oklahoma, South Dakota, West Virginia, range from 1% to 6%. Colorado, Georgia, Idaho, Indiana, Iowa, Louisiana, Mississippi, Missouri, North Carolina, Ohio, Texas, Wyoming, range from 7% to 9%. Arizona, Connecticut, Delaware, Maine, Maryland, Minnesota, New Mexico, Oregon, Tennessee, Virginia, Washington, Wisconsin, range from 10% to 14%. California, Florida, Massachusetts, Michigan, Nevada, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, South Carolina, Utah, Vermont, range from 15% to 23%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools that prohibited all forms of advertising and promotion of candy, fast food restaurants, or soft drinks in all locations*. The values range from 32% to 73%. Georgia, Indiana, Iowa, Kansas, Kentucky, Missouri, Montana, Nebraska, North Dakota, Ohio, Oklahoma, South Dakota, range from 32% to 38%. Alabama, Arkansas, Colorado, Florida, Idaho, South Carolina, Tennessee, Utah, Virginia, West Virginia, Wisconsin, Wyoming, range from 39% to 48%. Delaware, Maryland, Michigan, Minnesota, Mississippi, New Mexico, North Carolina, Oregon, Pennsylvania, Texas, Vermont, Washington, range from 49% to 58%. Alaska, Arizona, California, Connecticut, Hawaii, Louisiana, Maine, Massachusetts, Nevada, New Hampshire, New Jersey, New York, Rhode Island, range from 59% to 73%. Illinois did not have weighted results.


*In school buildings; on school grounds, including on the outside of the school building, on playing fields, or other areas of the campus; on school buses or other vehicles used to transport students; and in school publications.




This slide shows the percentage of secondary schools that used at least 3 different strategies to promote healthy eating. The values range from 9% to 50%. Alaska, Arkansas, Hawaii, Kentucky, Louisiana, Nevada, North Carolina, South Dakota, Utah, Washington, West Virginia, Wyoming, range from 9% to 16%. Alabama, Indiana, Kansas, Maryland, Mississippi, Missouri, Montana, Nebraska, New Mexico, Oregon, South Carolina, Texas, range from 17% to 20%. Arizona, Colorado, Connecticut, Georgia, Idaho, Iowa, Michigan, North Dakota, Ohio, Oklahoma, Tennessee, Virginia, range from 21% to 27%. California, Delaware, Florida, Maine, Massachusetts, Minnesota, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, Vermont, Wisconsin, range from 28% to 50%. Illinois did not have weighted results.



This slide shows the percentage of secondary schools that taught 14 key nutrition and dietary behavior topics in a required course. The values range from 41% to 88%. Alaska, Arizona, California, Connecticut, Maine, Massachusetts, Minnesota, Oklahoma, Oregon, Rhode Island, Vermont, range from 41% to 57%. Delaware, Florida, Hawaii, Kansas, Maryland, Michigan, New Hampshire, North Carolina, Ohio, South Dakota, Tennessee, Washington, range from 58% to 65%. Alabama, Georgia, Iowa, Louisiana, Nebraska, Nevada, Pennsylvania, South Carolina, Texas, Utah, Wisconsin, Wyoming, range from 66% to 70%. Arkansas, Idaho, Indiana, Kentucky, Mississippi, Missouri, Montana, New Jersey, New York, North Dakota, Virginia, West Virginia, range from 71% to 88%. Colorado, Illinois and New Mexico did not have weighted results.



This slide shows the percentage of secondary schools in which the lead health education teacher received professional development during the 2 years before the survey on nutrition and dietary behavior. The values range from 21% to 59%. Alabama, Alaska, California, Connecticut, Georgia, Indiana, Iowa, Montana, Nebraska, Oregon, South Dakota, range from 21% to 33%. Arizona, Delaware, Florida, Idaho, Kentucky, Minnesota, Pennsylvania, Rhode Island, South Carolina, Virginia, Washington, West Virginia, range from 34% to 40%. Arkansas, Kansas, Louisiana, Massachusetts, Michigan, New Jersey, North Carolina, North Dakota, Ohio, Texas, Utah, range from 41% to 43%. Hawaii, Maine, Maryland, Mississippi, Missouri, Nevada, New Hampshire, New York, Oklahoma, Tennessee, Vermont, Wisconsin, Wyoming, range from 44% to 59%. Colorado, Illinois and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that used the School Health Index or a similar self-assessment tool to assess their policies, activities, and programs in nutrition and physical activity. The values range from 20% to 78%. Alabama, Alaska, Arizona, Idaho, Iowa, Minnesota, Nevada, North Carolina, Oregon, South Dakota, Washington, Wyoming, range from 20% to 31%. California, Connecticut, Florida, Indiana, Kansas, Louisiana, Michigan, Nebraska, New Jersey, Ohio, Rhode Island, Utah, range from 32% to 39%. Colorado, Kentucky, Maryland, Massachusetts, New Hampshire, New Mexico, North Dakota, Oklahoma, Pennsylvania, South Carolina, Texas, Virginia, range from 40% to 44%. Arkansas, Delaware, Georgia, Hawaii, Maine, Mississippi, Missouri, Montana, New York, Tennessee, Vermont, West Virginia, Wisconsin, range from 45% to 78%. Illinois did not have weighted results.

School Health Coordination


National Center for Chronic Disease Prevention and Health Promotion
Division of Adolescent and School Health




School Health Coordination




This slide shows the percentage of secondary schools that have a designated individual responsible for coordinating school health and safety programs and activities. The values range from 71% to 93%. Alaska, Arizona, California, Colorado, Maine, Michigan, North Dakota, Ohio, Oregon, South Dakota, Vermont, Wyoming, range from 71% to 79%. Alabama, Connecticut, Idaho, Iowa, Kansas, Maryland, Massachusetts, Montana, New Hampshire, North Carolina, Rhode Island, Washington, range from 80% to 84%. Florida, Georgia, Hawaii, Indiana, Kentucky, Louisiana, Minnesota, Nebraska, New Mexico, South Carolina, Utah, Wisconsin, range from 85% to 87%. Arkansas, Delaware, Mississippi, Missouri, Nevada, New Jersey, New York, Oklahoma, Pennsylvania, Tennessee, Texas, Virginia, West Virginia, range from 88% to 93%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools that had a school health council, committee, or team that offers guidance on the development of policies or coordinates activities on health topics . The values range from 33% to 80%. Alabama, Alaska, Arizona, California, Florida, Idaho, Kentucky, Louisiana, Nevada, Oregon, Utah, Washington, range from 33% to 49%. Colorado, Connecticut, Georgia, Maryland, Montana, Nebraska, North Carolina, North Dakota, Ohio, Rhode Island, South Dakota, Wyoming, range from 50% to 57%. Indiana, Iowa, Kansas, Massachusetts, Michigan, Minnesota, Missouri, New Jersey, New Mexico, South Carolina, Virginia, Wisconsin, range from 58% to 64%. Arkansas, Delaware, Hawaii, Maine, Mississippi, New Hampshire, New York, Oklahoma, Pennsylvania, Tennessee, Texas, Vermont, West Virginia, range from 65% to 80%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools that had one or more than one group (e.g., a school health council, committee, or team) that offers guidance on the development of policies or coordinates activities on health topics with representation from at least 10 of 14 groups. The values range from 6% to 44%. Alabama, Alaska, Arizona, California, Colorado, Florida, Louisiana, Maryland, Nevada, North Dakota, Oregon, Washington, range from 6% to 13%. Connecticut, Georgia, Idaho, Kansas, Kentucky, Montana, Nebraska, New Jersey, North Carolina, Ohio, South Dakota, Utah, range from 14% to 17%. Delaware, Hawaii, Maine, Massachusetts, Michigan, Minnesota, New Hampshire, South Carolina, Vermont, Virginia, Wisconsin, Wyoming, range from 18% to 21%. Arkansas, Indiana, Iowa, Mississippi, Missouri, New Mexico, New York, Oklahoma, Pennsylvania, Rhode Island, Tennessee, Texas, West Virginia, range from 22% to 44%. Illinois did not have weighted results.



This slide shows the percentage of secondary schools in which students' families and community members helped develop or implement policies and programs on tobacco-use prevention, physical activity, and nutrition and healthy eating. The values range from 4% to 31%. Arizona, Connecticut, Georgia, Louisiana, Massachusetts, Nebraska, Nevada, New Hampshire, New Jersey, Ohio, Oregon, Washington, range from 4% to 10%. California, Colorado, Florida, Hawaii, Idaho, Maryland, North Carolina, North Dakota, Vermont, Wisconsin, Wyoming, range from 11% to 12%. Alabama, Delaware, Indiana, Kansas, Kentucky, Maine, Michigan, Montana, Pennsylvania, South Carolina, Utah, Virginia, West Virginia, range from 13% to 16%. Alaska, Arkansas, Iowa, Minnesota, Mississippi, Missouri, New Mexico, New York, Oklahoma, Rhode Island, South Dakota, Tennessee, Texas, range from 17% to 31%. Illinois did not have weighted results.



This slide shows the percentage of secondary schools that ever used the School Health Index or other self-assessment tool to assess their policies, activities, and programs in physical activity, nutrition, and tobacco-use prevention. The values range from 18% to 72%. Alabama, Alaska, Arizona, Iowa, Minnesota, Nebraska, Nevada, North Carolina, Ohio, Oregon, Washington, Wyoming, range from 18% to 26%. California, Connecticut, Florida, Idaho, Indiana, Louisiana, Michigan, New Jersey, Pennsylvania, Rhode Island, South Dakota, Utah, range from 27% to 33%. Colorado, Kansas, Kentucky, Maryland, Massachusetts, Missouri, New Hampshire, New Mexico, North Dakota, South Carolina, Texas, Virginia, range from 34% to 38%. Arkansas, Delaware, Georgia, Hawaii, Maine, Mississippi, Montana, New York, Oklahoma, Tennessee, Vermont, West Virginia, Wisconsin, range from 39% to 72%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools with a School Improvement Plan that includes any health-related goals and objectives. The values range from 29% to 94%. Alaska, Kansas, Maine, Minnesota, New Hampshire, New Jersey, New York, Pennsylvania, South Dakota, West Virginia, Wisconsin, Wyoming, range from 29% to 58%. Arizona, California, Connecticut, Idaho, Michigan, Montana, Nebraska, Nevada, North Dakota, Ohio, Oklahoma, Vermont, range from 59% to 71%. Colorado, Georgia, Indiana, Kentucky, Mississippi, New Mexico, Oregon, Rhode Island, South Carolina, Utah, Virginia, Washington, range from 72% to 79%. Alabama, Arkansas, Delaware, Florida, Hawaii, Iowa, Louisiana, Maryland, Massachusetts, Missouri, North Carolina, Tennessee, Texas, range from 80% to 94%. Illinois did not have weighted results.

HIV, STD, and Pregnancy Prevention


National Center for Chronic Disease Prevention and Health Promotion
Division of Adolescent and School Health




HIV, STD, and Pregnancy Prevention




This slide shows the percentage of secondary schools that tried to increase student knowledge on HIV prevention in a required course. The values range from 53% to 97%. Alabama, Alaska, Arizona, Florida, Louisiana, Massachusetts, Oklahoma, South Dakota, Tennessee, Texas, Vermont, range from 53% to 86%. California, Connecticut, Georgia, Iowa, Kansas, Kentucky, Maryland, Michigan, Mississippi, North Carolina, Ohio, Wyoming, range from 87% to 89%. Delaware, Maine, Minnesota, Missouri, Montana, Nebraska, New Hampshire, North Dakota, Oregon, Pennsylvania, South Carolina, Virginia, range from 90% to 93%. Arkansas, Hawaii, Idaho, Indiana, Nevada, New Jersey, New York, Rhode Island, Utah, Washington, West Virginia, Wisconsin, range from 94% to 97%. Colorado, Illinois and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that tried to increase student knowledge on human sexuality in a required course. The values range from 49% to 95%. Alabama, Alaska, Arizona, Florida, Georgia, Louisiana, North Carolina, Oklahoma, South Dakota, Tennessee, Texas, range from 49% to 79%. Arkansas, California, Kentucky, Massachusetts, Michigan, Mississippi, Missouri, Montana, North Dakota, Ohio, Virginia, Wyoming, range from 80% to 86%. Connecticut, Idaho, Iowa, Maine, Maryland, Nebraska, Pennsylvania, Rhode Island, South Carolina, Vermont, Washington, West Virginia, range from 87% to 90%. Delaware, Hawaii, Indiana, Kansas, Minnesota, Nevada, New Hampshire, New Jersey, New York, Oregon, Utah, Wisconsin, range from 91% to 95%. Colorado, Illinois and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that tried to increase student knowledge on pregnancy prevention in a required course. The values range from 47% to 94%. Alaska, Arizona, California, Connecticut, Florida, Louisiana, Michigan, North Dakota, Oklahoma, South Dakota, Vermont, range from 47% to 77%. Iowa, Kansas, Massachusetts, Montana, North Carolina, Pennsylvania, Rhode Island, Tennessee, Texas, Utah, Washington, Wyoming, range from 78% to 82%. Alabama, Idaho, Kentucky, Maine, Maryland, Minnesota, Mississippi, Missouri, Nebraska, New Hampshire, Ohio, Virginia, range from 83% to 87%. Arkansas, Delaware, Georgia, Hawaii, Indiana, Nevada, New Jersey, New York, Oregon, South Carolina, West Virginia, Wisconsin, range from 88% to 94%. Colorado, Illinois and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that tried to increase student knowledge on STD prevention in a required course. The values range from 50% to 97%. Alaska, Arizona, Florida, Louisiana, Massachusetts, Michigan, North Dakota, Oklahoma, South Dakota, Tennessee, Vermont, range from 50% to 85%. California, Connecticut, Iowa, Kansas, Kentucky, Mississippi, Montana, New Hampshire, North Carolina, Ohio, Texas, Wyoming, range from 86% to 88%. Alabama, Georgia, Hawaii, Idaho, Maine, Maryland, Minnesota, Pennsylvania, Rhode Island, South Carolina, Virginia, West Virginia, range from 89% to 91%. Arkansas, Delaware, Indiana, Missouri, Nebraska, Nevada, New Jersey, New York, Oregon, Utah, Washington, Wisconsin, range from 92% to 97%. Colorado, Illinois and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that taught 11 key HIV, STD, and pregnancy prevention topics in a required course during grades 6, 7, or 8. The values range from 13% to 66%. Alaska, Arizona, Connecticut, Kansas, Louisiana, Minnesota, Mississippi, Nebraska, South Dakota, Vermont, Wyoming, range from 13% to 39%. Idaho, Iowa, Maine, Massachusetts, Michigan, Montana, North Dakota, Oklahoma, Oregon, Pennsylvania, Tennessee, Utah, range from 40% to 42%. Alabama, California, Delaware, Florida, Kentucky, Missouri, New Hampshire, North Carolina, Ohio, Texas, Washington, Wisconsin, range from 43% to 53%. Arkansas, Georgia, Hawaii, Indiana, Maryland, Nevada, New Jersey, New York, Rhode Island, South Carolina, Virginia, West Virginia, range from 54% to 66%. Colorado, Illinois and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that taught 8 key HIV, STD, and pregnancy prevention topics in a required course during grades 9, 10, 11, or 12. The values range from 45% to 96%. Alaska, Arizona, Iowa, Louisiana, Montana, Nebraska, North Carolina, North Dakota, Oklahoma, South Dakota, Wyoming, range from 45% to 70%. Alabama, Connecticut, Florida, Kansas, Maine, Massachusetts, Michigan, Minnesota, Texas, Utah, Virginia, Wisconsin, range from 71% to 79%. California, Hawaii, Idaho, New Hampshire, Ohio, Oregon, Pennsylvania, Rhode Island, South Carolina, Tennessee, Vermont, Washington, range from 80% to 84%. Arkansas, Delaware, Georgia, Indiana, Kentucky, Maryland, Mississippi, Missouri, Nevada, New Jersey, New York, West Virginia, range from 85% to 96%. Colorado, Illinois and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that taught 4 key topics related to condom use in a required course during grades 9, 10, 11, or 12. The values range from 9% to 88%. Alabama, Alaska, Arizona, Georgia, North Carolina, North Dakota, Oklahoma, South Dakota, Texas, Utah, Wyoming, range from 9% to 31%. Florida, Idaho, Indiana, Kansas, Louisiana, Michigan, Mississippi, Missouri, Montana, Nebraska, Ohio, Tennessee, range from 32% to 44%. Arkansas, Iowa, Kentucky, Minnesota, Nevada, Pennsylvania, Rhode Island, South Carolina, Virginia, Washington, West Virginia, Wisconsin, range from 45% to 66%. California, Connecticut, Delaware, Hawaii, Maine, Maryland, Massachusetts, New Hampshire, New Jersey, New York, Oregon, Vermont, range from 67% to 88%. Colorado, Illinois and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that taught how to access valid and reliable health information, products, or services related to HIV, other STDs, and pregnancy in a required course. The values range from 42% to 92%. Alaska, Arizona, Florida, Louisiana, Michigan, Montana, North Carolina, Oklahoma, South Dakota, Tennessee, Wyoming, range from 42% to 73%. California, Connecticut, Iowa, Kansas, Kentucky, Maine, Massachusetts, Mississippi, New Hampshire, North Dakota, Ohio, Texas, range from 74% to 78%. Alabama, Delaware, Indiana, Maryland, Missouri, Nebraska, Pennsylvania, South Carolina, Utah, Vermont, Virginia, Wisconsin, range from 79% to 82%. Arkansas, Georgia, Hawaii, Idaho, Minnesota, Nevada, New Jersey, New York, Oregon, Rhode Island, Washington, West Virginia, range from 83% to 92%. Colorado, Illinois and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools in which the lead health education teacher received professional development during the 2 years before the survey on HIV prevention. The values range from 15% to 67%. Alaska, Arizona, Arkansas, Connecticut, Louisiana, Nebraska, Ohio, Oregon, Rhode Island, South Dakota, Wyoming, range from 15% to 27%. Indiana, Iowa, Kansas, Kentucky, Massachusetts, Mississippi, Missouri, North Dakota, Vermont, Virginia, West Virginia, Wisconsin, range from 28% to 37%. Alabama, California, Idaho, Maine, Minnesota, Montana, New Hampshire, New Jersey, North Carolina, Pennsylvania, Tennessee, Texas, range from 38% to 45%. Delaware, Florida, Georgia, Hawaii, Maryland, Michigan, Nevada, New York, Oklahoma, South Carolina, Utah, Washington, range from 46% to 67%. Colorado, Illinois and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools in which the lead health education teacher received professional development during the 2 years before the survey on 4 key HIV prevention topics. The values range from 7% to 48%. Alaska, Arizona, Connecticut, Indiana, Kansas, Kentucky, Louisiana, Nebraska, Oregon, South Dakota, Wyoming, range from 7% to 15%. Alabama, Arkansas, Iowa, Minnesota, Mississippi, Missouri, North Dakota, Ohio, Rhode Island, Virginia, West Virginia, Wisconsin, range from 16% to 21%. Florida, Georgia, Maine, Massachusetts, Montana, New Hampshire, New Jersey, North Carolina, Pennsylvania, Tennessee, Texas, Vermont, range from 22% to 28%. California, Delaware, Hawaii, Idaho, Maryland, Michigan, Nevada, New York, Oklahoma, South Carolina, Utah, Washington, range from 29% to 48%. Colorado, Illinois and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools in which the lead health education teacher received professional development during the 2 years before the survey on at least 6 of 11 key HIV prevention topics. The values range from 5% to 49%. Alaska, Arizona, Connecticut, Indiana, Kansas, Minnesota, Nebraska, Rhode Island, South Dakota, Wyoming, range from 5% to 13%. Alabama, Iowa, Kentucky, Louisiana, Massachusetts, Missouri, New Hampshire, North Dakota, Ohio, Oregon, Virginia, West Virginia, Wisconsin, range from 14% to 18%. Arkansas, Georgia, Idaho, Maine, Mississippi, Montana, North Carolina, Pennsylvania, Tennessee, Texas, Utah, Vermont, range from 19% to 24%. California, Delaware, Florida, Hawaii, Maryland, Michigan, Nevada, New Jersey, New York, Oklahoma, South Carolina, Washington, range from 25% to 49%. Colorado, Illinois and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools with a gay/straight alliance or similar club. The values range from 9% to 50%. Arkansas, Georgia, Kansas, Kentucky, Louisiana, Mississippi, Nebraska, North Dakota, Oklahoma, South Carolina, South Dakota, Tennessee, range from 9% to 18%. Alabama, Alaska, Arizona, Idaho, Iowa, Missouri, Montana, New Mexico, North Carolina, Ohio, Texas, West Virginia, range from 19% to 25%. Delaware, Hawaii, Indiana, Michigan, Minnesota, Nevada, Oregon, Pennsylvania, Utah, Virginia, Wisconsin, Wyoming, range from 26% to 31%. California, Colorado, Connecticut, Florida, Maine, Maryland, Massachusetts, New Hampshire, New Jersey, New York, Rhode Island, Vermont, Washington, range from 32% to 50%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools that provide curricula or supplementary materials and engage in 5 practices related to LGBTQ youth. The values range from 1% to 21%. Alabama, Arizona, Georgia, Idaho, Indiana, Kansas, Louisiana, Mississippi, Montana, South Carolina, South Dakota, Utah, range from 1% to 2%. Alaska, Colorado, Kentucky, Missouri, New Mexico, North Carolina, North Dakota, Ohio, Oklahoma, Texas, Virginia, Wyoming, range from 3% to 5%. Arkansas, Delaware, Florida, Maryland, Michigan, Minnesota, Nebraska, Nevada, Oregon, Tennessee, Washington, West Virginia, range from 6% to 8%. California, Connecticut, Hawaii, Iowa, Maine, Massachusetts, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, Vermont, Wisconsin, range from 9% to 21%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools that deliver HIV, STD, and pregnancy prevention programs (including after school or supplemental programs) that meet the needs of ethnic/racial minority youth at high risk. The values range from 0% to 15%. Connecticut, Indiana, Kansas, Maine, Minnesota, New Hampshire, Pennsylvania, South Dakota, Utah, Vermont, Virginia, Wyoming, range from 0% to 2%. Alaska, Arizona, Colorado, Idaho, Missouri, Montana, Nebraska, New Jersey, North Dakota, Ohio, Oregon, range from 3% to 4%. Delaware, Georgia, Hawaii, Iowa, Kentucky, Louisiana, Massachusetts, Michigan, Rhode Island, Washington, West Virginia, Wisconsin, range from 5% to 6%. Alabama, Arkansas, California, Florida, Maryland, Mississippi, Nevada, New Mexico, New York, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, range from 7% to 15%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools with a policy on students or staff who have HIV infection or AIDS that addresses attendance of students with HIV infection, procedures to protect HIV-infected students and staff from discrimination, and maintaining confidentiality of HIV-infected students and staff. The values range from 25% to 85%. Alaska, Arkansas, California, Colorado, Delaware, Kentucky, Louisiana, Michigan, Mississippi, North Carolina, North Dakota, Texas, range from 25% to 50%. Alabama, Hawaii, Iowa, Kansas, Maryland, Missouri, Nebraska, New Mexico, Ohio, West Virginia, Wisconsin, Wyoming, range from 51% to 55%. Arizona, Florida, Georgia, Idaho, Indiana, Massachusetts, Minnesota, Montana, New Jersey, New York, South Dakota, Utah, range from 56% to 63%. Connecticut, Maine, Nevada, New Hampshire, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, Tennessee, Vermont, Virginia, Washington, range from 64% to 85%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools that provide parents and families with health information to increase parent and family knowledge of HIV prevention, STD prevention, or teen pregnancy prevention. The values range from 11% to 93%. Alaska, Arizona, Connecticut, Indiana, Kansas, Louisiana, Nebraska, New Hampshire, South Dakota, Vermont, Wyoming, range from 11% to 19%. Iowa, Maine, Massachusetts, Minnesota, Mississippi, Missouri, Montana, North Dakota, Ohio, Oregon, Pennsylvania, Rhode Island, range from 20% to 27%. Florida, Georgia, Idaho, Michigan, New Jersey, North Carolina, South Carolina, Tennessee, Texas, Virginia, West Virginia, Wisconsin, range from 28% to 33%. Alabama, Arkansas, California, Delaware, Hawaii, Kentucky, Maryland, Nevada, New York, Oklahoma, Utah, Washington, range from 34% to 93%. Colorado, Illinois and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools in which students' families and community members helped develop or implement HIV prevention, STD prevention, or teen pregnancy prevention policies and programs. The values range from 5% to 37%. Connecticut, Delaware, Hawaii, Kansas, Louisiana, Maine, Nebraska, New Hampshire, North Dakota, Ohio, Vermont, Wyoming, range from 5% to 8%. Alaska, Arizona, Colorado, Idaho, Massachusetts, Nevada, New Jersey, Oregon, Rhode Island, South Dakota, Utah, West Virginia, range from 9% to 10%. California, Florida, Georgia, Iowa, Kentucky, Mississippi, Missouri, Montana, North Carolina, Pennsylvania, Washington, Wisconsin, range from 11% to 14%. Alabama, Arkansas, Indiana, Maryland, Michigan, Minnesota, New Mexico, New York, Oklahoma, South Carolina, Tennessee, Texas, Virginia, range from 15% to 37%. Illinois did not have weighted results.




Health Education




This slide shows the percentage of secondary schools that required students to take 2 or more health education courses. The values range from 12% to 92%. Alabama, Arizona, California, Florida, Louisiana, Mississippi, Nevada, Oklahoma, South Dakota, Tennessee, Texas, range from 12% to 38%. Alaska, Georgia, Hawaii, Iowa, Kansas, Kentucky, Maryland, Michigan, North Carolina, Ohio, South Carolina, Utah, range from 39% to 51%. Arkansas, Delaware, Idaho, Indiana, Maine, Massachusetts, Missouri, Nebraska, New York, Vermont, Washington, Wyoming, range from 52% to 62%. Connecticut, Minnesota, Montana, New Hampshire, New Jersey, North Dakota, Oregon, Pennsylvania, Rhode Island, Virginia, West Virginia, Wisconsin, range from 63% to 92%. Colorado, Illinois and New Mexico did not have weighted results.




This slide shows the among secondary schools that required a health education course, percentage that required students who fail the course to repeat it. The values range from 21% to 85%. California, Connecticut, Maine, Massachusetts, New Jersey, North Carolina, Oklahoma, South Carolina, Utah, Virginia, West Virginia, range from 21% to 50%. Arizona, Delaware, Georgia, Iowa, Maryland, New Hampshire, Oregon, Pennsylvania, Rhode Island, Tennessee, Vermont, Washington, range from 51% to 60%. Florida, Indiana, Kansas, Kentucky, Louisiana, Minnesota, Missouri, Nebraska, Nevada, North Dakota, Ohio, Wyoming, range from 61% to 67%. Alabama, Alaska, Arkansas, Hawaii, Idaho, Michigan, Mississippi, Montana, New York, South Dakota, Texas, Wisconsin, range from 68% to 85%. Colorado, Illinois and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools with a health education curriculum that addresses all 8 national standards for health education. The values range from 37% to 83%. Alaska, Arizona, California, Florida, Kansas, Minnesota, North Dakota, Oklahoma, South Dakota, Vermont, Washington, range from 37% to 57%. Alabama, Delaware, Hawaii, Iowa, Louisiana, Massachusetts, Michigan, Nebraska, New Hampshire, Ohio, Oregon, Rhode Island, range from 58% to 66%. Connecticut, Georgia, Idaho, Maine, Montana, Nevada, Pennsylvania, South Carolina, Tennessee, Texas, Utah, Wisconsin, range from 67% to 70%. Arkansas, Indiana, Kentucky, Maryland, Mississippi, Missouri, New Jersey, New York, North Carolina, Virginia, West Virginia, Wyoming, range from 71% to 83%. Colorado, Illinois and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools in which the lead health education teacher had professional preparation in health education or in health and physical education combined. The values range from 17% to 89%. Alaska, Arizona, California, Florida, Hawaii, Iowa, Mississippi, Nebraska, North Dakota, Vermont, Washington, range from 17% to 49%. Alabama, Connecticut, Kansas, Michigan, Missouri, New Hampshire, Oklahoma, Oregon, South Carolina, South Dakota, Texas, Wyoming, range from 50% to 61%. Idaho, Kentucky, Maine, Maryland, Massachusetts, Montana, Nevada, New York, North Carolina, Tennessee, Utah, Wisconsin, range from 62% to 74%. Arkansas, Delaware, Georgia, Indiana, Louisiana, Minnesota, New Jersey, Ohio, Pennsylvania, Rhode Island, Virginia, West Virginia, range from 75% to 89%. Colorado, Illinois and New Mexico did not have weighted results.



This slide shows the percentage of secondary schools in which all staff who teach health education were licensed, certified, or endorsed by the state in health education. The values range from 26% to 99%. Alabama, Alaska, Arizona, California, Colorado, Florida, Hawaii, New Hampshire, Oregon, Vermont, Washington, range from 26% to 78%. Connecticut, Kansas, Maine, Maryland, Massachusetts, Nebraska, New Mexico, New York, North Carolina, Ohio, South Carolina, Tennessee, range from 79% to 85%. Delaware, Idaho, Iowa, Kentucky, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Oklahoma, Texas, Wisconsin, range from 86% to 92%. Arkansas, Georgia, Indiana, Montana, Nevada, New Jersey, North Dakota, Pennsylvania, Rhode Island, South Dakota, Utah, Virginia, West Virginia, Wyoming, range from 93% to 99%. Illinois did not have weighted results.



This slide shows the percentage of secondary schools in which those who teach health education were provided with key materials for teaching health education. The values range from 33% to 70%. Alabama, Alaska, Arizona, California, Kansas, Minnesota, Nebraska, New Hampshire, North Dakota, Oklahoma, South Dakota, range from 33% to 43%. Connecticut, Florida, Indiana, Iowa, Maine, Massachusetts, North Carolina, Ohio, Oregon, Utah, Vermont, Washington, range from 44% to 51%. Arkansas, Delaware, Georgia, Hawaii, Louisiana, Michigan, Montana, Nevada, South Carolina, Tennessee, West Virginia, Wisconsin, range from 52% to 56%. Idaho, Kentucky, Maryland, Mississippi, Missouri, New Jersey, New York, Pennsylvania, Rhode Island, Texas, Virginia, Wyoming, range from 57% to 70%. Colorado, Illinois and New Mexico did not have weighted results.




This slide shows the percentage of secondary schools that provided parents and families health information to increase parent and family knowledge of tobacco-use prevention, physical activity, and nutrition and healthy eating. The values range from 16% to 46%. Alaska, Connecticut, Georgia, Indiana, Iowa, Massachusetts, Minnesota, New Hampshire, Oregon, Pennsylvania, Rhode Island, range from 16% to 23%. Arizona, Idaho, Kansas, Maine, Michigan, Nevada, Ohio, South Dakota, Utah, Vermont, Washington, Wisconsin, range from 24% to 27%. Alabama, California, Florida, Hawaii, Louisiana, Missouri, Montana, Nebraska, North Carolina, South Carolina, Texas, Virginia, range from 28% to 30%. Arkansas, Delaware, Kentucky, Maryland, Mississippi, New Jersey, New York, North Dakota, Oklahoma, Tennessee, West Virginia, Wyoming, range from 31% to 46%. Colorado, Illinois and New Mexico did not have weighted results.

Asthma


National Center for Chronic Disease Prevention and Health Promotion
Division of Adolescent and School Health




Asthma




This slide shows the percentage of secondary schools with a full-time registered nurse who provides health services to students at school. The values range from 5% to 99%. Alaska, California, Hawaii, Idaho, Michigan, Montana, Nevada, North Dakota, Oregon, Utah, Washington, Wisconsin, range from 5% to 23%. Arizona, Colorado, Kansas, Louisiana, Minnesota, Nebraska, North Carolina, Ohio, Oklahoma, South Dakota, West Virginia, Wyoming, range from 24% to 43%. Alabama, Arkansas, Florida, Georgia, Indiana, Iowa, Kentucky, Maine, Maryland, Mississippi, New Mexico, Tennessee, range from 44% to 73%. Connecticut, Delaware, Massachusetts, Missouri, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, South Carolina, Texas, Vermont, Virginia, range from 74% to 99%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools that implemented a policy permitting students to carry and self-administer asthma medications by communicating the policy to students, parents, and families, and by designating an individual responsible for implementing the policy. The values range from 22% to 70%. Alaska, Arizona, California, Hawaii, Idaho, Maryland, Michigan, Montana, North Dakota, Oregon, South Dakota, Utah, range from 22% to 45%. Alabama, Colorado, Connecticut, Florida, Kansas, Kentucky, Louisiana, Mississippi, Missouri, New York, Ohio, Wisconsin, range from 46% to 52%. Arkansas, Delaware, Georgia, Iowa, Massachusetts, Minnesota, Nebraska, New Mexico, Oklahoma, Pennsylvania, Washington, Wyoming, range from 53% to 58%. Indiana, Maine, Nevada, New Hampshire, New Jersey, North Carolina, Rhode Island, South Carolina, Tennessee, Texas, Vermont, Virginia, West Virginia, range from 59% to 70%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools that had an asthma action plan on file for all students with known asthma. The values range from 31% to 88%. Alaska, Arizona, Hawaii, Idaho, Michigan, Mississippi, Montana, North Dakota, South Dakota, Utah, Vermont, Wyoming, range from 31% to 49%. California, Iowa, Kansas, Minnesota, New Mexico, New York, Ohio, Oklahoma, Oregon, Pennsylvania, Texas, Wisconsin, range from 50% to 58%. Arkansas, Connecticut, Florida, Georgia, Indiana, Kentucky, Maine, Massachusetts, Missouri, Nebraska, New Hampshire, Virginia, range from 59% to 67%. Alabama, Colorado, Delaware, Louisiana, Maryland, Nevada, New Jersey, North Carolina, Rhode Island, South Carolina, Tennessee, Washington, West Virginia, range from 68% to 88%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools that identified students with poorly controlled asthma by keeping track of them in at least 3 different ways. The values range from 41% to 76%. Alabama, Alaska, Arkansas, Kentucky, Louisiana, Mississippi, Montana, North Dakota, Oklahoma, South Dakota, Tennessee, West Virginia, range from 41% to 57%. Arizona, Florida, Idaho, Iowa, Kansas, Michigan, Minnesota, Nebraska, Ohio, Texas, Utah, Wisconsin, range from 58% to 63%. Colorado, Delaware, Hawaii, Indiana, Massachusetts, Missouri, Nevada, New Hampshire, North Carolina, Oregon, South Carolina, Wyoming, range from 64% to 70%. California, Connecticut, Georgia, Maine, Maryland, New Jersey, New Mexico, New York, Pennsylvania, Rhode Island, Vermont, Virginia, Washington, range from 71% to 76%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools that provided intensive case management for students with poorly controlled asthma at school. The values range from 10% to 51%. Alaska, Arizona, Florida, Kansas, Louisiana, Michigan, Montana, North Dakota, Ohio, Oklahoma, South Dakota, Utah, range from 10% to 18%. Alabama, Arkansas, California, Colorado, Georgia, Hawaii, Idaho, Indiana, Iowa, Nebraska, Oregon, Wyoming, range from 19% to 25%. Kentucky, Minnesota, Mississippi, Missouri, Nevada, North Carolina, Pennsylvania, Tennessee, Texas, Virginia, Washington, Wisconsin, range from 26% to 33%. Connecticut, Delaware, Maine, Maryland, Massachusetts, New Hampshire, New Jersey, New Mexico, New York, Rhode Island, South Carolina, Vermont, West Virginia, range from 34% to 51%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools that required all school staff members to receive annual training on recognizing and responding to severe asthma symptoms. The values range from 12% to 66%. Alaska, California, Hawaii, Idaho, Indiana, Iowa, Kansas, Michigan, Montana, North Dakota, Ohio, South Dakota, range from 12% to 25%. Arizona, Delaware, Maine, Minnesota, Mississippi, New Hampshire, New York, Oregon, Pennsylvania, Utah, Vermont, Wisconsin, range from 26% to 30%. Arkansas, Colorado, Connecticut, Florida, Georgia, Maryland, Massachusetts, Nevada, New Mexico, Oklahoma, Rhode Island, Wyoming, range from 31% to 40%. Alabama, Kentucky, Louisiana, Missouri, Nebraska, New Jersey, North Carolina, South Carolina, Tennessee, Texas, Virginia, Washington, West Virginia, range from 41% to 66%. Illinois did not have weighted results.



This slide shows the percentage of secondary schools that provided parents and families of students with health information to increase their knowledge of asthma . The values range from 7% to 37%. Alaska, Idaho, Indiana, Iowa, Kansas, Minnesota, New Hampshire, Oregon, South Dakota, Utah, Wyoming, range from 7% to 15%. Connecticut, Georgia, Hawaii, Maine, Massachusetts, Michigan, Montana, Nevada, North Dakota, Pennsylvania, Rhode Island, Wisconsin, range from 16% to 19%. Arizona, California, Florida, Kentucky, Louisiana, Maryland, North Carolina, Ohio, South Carolina, Texas, Vermont, Washington, range from 20% to 24%. Alabama, Arkansas, Delaware, Mississippi, Missouri, Nebraska, New Jersey, New York, Oklahoma, Tennessee, Virginia, West Virginia, range from 25% to 37%. Colorado, Illinois and New Mexico did not have weighted results.



This slide shows the percentage of secondary schools in which students' families and community members helped develop or implement asthma management policies and programs. The values range from 3% to 25%. Alaska, Colorado, Connecticut, Kansas, Louisiana, Maine, Nevada, New Hampshire, Ohio, Oregon, Wyoming, range from 3% to 6%. Arizona, California, Hawaii, Idaho, Indiana, Iowa, Kentucky, Nebraska, North Dakota, Rhode Island, South Dakota, Washington, Wisconsin, range from 7% to 9%. Florida, Georgia, Maryland, Massachusetts, Michigan, Minnesota, Montana, New Jersey, Pennsylvania, Utah, Vermont, Virginia, range from 10% to 12%. Alabama, Arkansas, Delaware, Mississippi, Missouri, New Mexico, New York, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, West Virginia, range from 13% to 25%. Illinois did not have weighted results.




This slide shows the percentage of secondary schools that have ever used the School Health Index or other self-assessment tool to assess their asthma policies, activities, and programs. The values range from 12% to 59%. Alaska, Arizona, California, Idaho, Iowa, Kansas, Nevada, Ohio, Oregon, South Dakota, Washington, range from 12% to 17%. Alabama, Connecticut, Indiana, Kentucky, Louisiana, Minnesota, Nebraska, North Carolina, North Dakota, Pennsylvania, Utah, Wyoming, range from 18% to 22%. Colorado, Delaware, Florida, Massachusetts, Michigan, New Hampshire, Oklahoma, Rhode Island, South Carolina, Texas, Virginia, West Virginia, Wisconsin, range from 23% to 29%. Arkansas, Georgia, Hawaii, Maine, Maryland, Mississippi, Missouri, Montana, New Jersey, New Mexico, New York, Tennessee, Vermont, range from 30% to 59%. Illinois did not have weighted results.

Violence Prevention


National Center for Chronic Disease Prevention and Health Promotion
Division of Adolescent and School Health




Violence Prevention



This slide shows the percentage of secondary schools that have or participate in a safe passages to school program. The values range from 13% to 64%. Alaska, Connecticut, Delaware, Indiana, Iowa, Kansas, Maine, New Hampshire, North Dakota, South Dakota, Washington, Wyoming, range from 13% to 18%. Alabama, Georgia, Hawaii, Massachusetts, Missouri, Nebraska, North Carolina, Oregon, Pennsylvania, Rhode Island, Vermont, Wisconsin, range from 19% to 20%. Arkansas, California, Florida, Idaho, Kentucky, Louisiana, Michigan, Minnesota, Montana, Ohio, Oklahoma, South Carolina, range from 21% to 26%. Arizona, Colorado, Maryland, Mississippi, Nevada, New Jersey, New Mexico, New York, Tennessee, Texas, Utah, Virginia, West Virginia, range from 27% to 64%. Illinois did not have weighted results.



This slide shows the percentage of secondary schools that have or participate in a program to prevent bullying. The values range from 69% to 96%. Alabama, Alaska, California, Hawaii, Louisiana, Michigan, Mississippi, Nevada, North Dakota, Ohio, Oregon, Washington, range from 69% to 78%. Arizona, Arkansas, Georgia, Idaho, Indiana, Iowa, Maine, Missouri, Montana, New Mexico, South Carolina, South Dakota, range from 79% to 81%. Kentucky, Massachusetts, Minnesota, Nebraska, New Hampshire, North Carolina, Oklahoma, Rhode Island, Texas, Virginia, Wisconsin, Wyoming, range from 82% to 85%. Colorado, Connecticut, Delaware, Florida, Kansas, Maryland, New Jersey, New York, Pennsylvania, Tennessee, Utah, Vermont, West Virginia, range from 86% to 96%. Illinois did not have weighted results.



This slide shows the percentage of secondary schools that have or participate in a program to prevent dating violence. The values range from 20% to 78%. Alaska, California, Delaware, Georgia, Kansas, Nevada, North Dakota, Oregon, South Dakota, Washington, Wyoming, range from 20% to 26%. Alabama, Arizona, Arkansas, Connecticut, Florida, Iowa, Kentucky, Louisiana, Michigan, Mississippi, North Carolina, Pennsylvania, South Carolina, range from 27% to 31%. Colorado, Indiana, Maine, Minnesota, Missouri, Montana, New Hampshire, Ohio, Oklahoma, Tennessee, Vermont, Virginia, range from 32% to 37%. Hawaii, Idaho, Maryland, Massachusetts, Nebraska, New Jersey, New Mexico, New York, Rhode Island, Texas, Utah, West Virginia, Wisconsin, range from 38% to 78%. Illinois did not have weighted results.