

## **Appendixes**



**Appendix I. Published Reports of Less Common Causes of Elevated Blood Lead Levels (EBLLs) in Children.**

<b>Exposure Source</b>	<b>Description/Exposure Pathway</b>	<b>Study Type*</b>	<b>Study Description</b>	<b>Ref. #</b>
Occupational Take Home Exposures				
<i>Battery reclamation</i>	Lead carried home by battery workers. (Only a minority of battery workers showered or changed clothes before going home.)	E	Twelve (75%) of 16 children of lead-exposed workers had EBLLs and a higher average BLL than neighborhood controls (22.4 vs. 9.8 µg/dL, $p=.049$ ).	28
<i>Ceramics</i>	Ceramic-coated capacitors made with fritted glass containing lead.	E	Case-control study of 51 children under 6 years (20 exposed, 31 controls) showed higher average BLLs in exposed children (13.4 vs. 7.1 µg/dL, $p<.001$ ).	31
<i>Furniture refinishing</i>	Lead carried home by workers who restored furniture that had undergone chemical stripping and was thought to be lead-free.	CR	Report of six workers and three of their children aged 4-18 months.	19
<i>Construction</i>	Lead dust on skin and clothes taken home.	E	Case-control study of 50 children under 6 years (31 exposed, 19 controls) showed 25.8% of workers' children had EBLLs compared to 5.3% of control children (OR=6.1).	53
<i>Radiator repair</i>	Lead carried home by workers who did soldering to repair radiator.	E	The mean BLL for 18 children (under 7 years) of lead-exposed workers was 10 µg/dL.	42, 34

Exposure Source	Description/Exposure Pathway	Study Type*	Study Description	Ref. #
Imported Cosmetics				
<i>Kohl (Middle East, India, Pakistan, some parts of Africa)</i>	A gray or black eye cosmetic applied to the conjunctival margins of the eyes. Can contain up to 83% lead. It is believed to strengthen and protect the eyes against disease. Also known as Al Kohl.	E	A study of 538 girls aged 6 to 12 years demonstrated that the application of kohl was associated with higher BLLs ( $p=0.0461$ ).	3, 37
<i>Pakistani eye cosmetics</i>	Eye cosmetics are often applied to the eyes of children.	E	Retrospective chart review of 175 children aged 8 months to 6 years showed an average BLL of 4.3 $\mu\text{g}/\text{dL}$ for Pakistani/Indian children not using eye cosmetics and 12.9 $\mu\text{g}/\text{dL}$ for those using eye cosmetics ( $p=0.03$ ).	50, 52
<i>Surma (India)</i>	A black fine powder applied to the eyes for medicinal and cosmetic reasons.	E	A case-control study of 62 children demonstrated higher BLLs in children using surma ( $p<.001$ ).	2, 15
Contaminated Foods				
<i>Apple cider</i>	Cider was made in a maple syrup evaporator that had lead solder joining the interior seams.	CR	Report of a 7-year-old child.	8
<i>Flour (Middle East)</i>	Lead fillings used in stone mills contaminated flour.	E	Investigation of 43 symptomatic patients aged zero to 80 years and their families and of 563 children aged 10 to 18 years demonstrated that 33 (23%) of 146 community stone mills had lead contamination and that 171 (30.4%) of 563 children had BLLs exceeding 30 $\mu\text{g}/\text{dL}$ .	25

Exposure Source	Description/Exposure Pathway	Study Type*	Study Description	Ref. #
<i>Lozeena</i>	An orange powder used to color rice and meat that contains 7.8%-8.9% lead.	CR	Report of brothers aged 2 and 3 years and their parents. In addition, 9 of 18 extended family members had EBLLs.	15
<i>Infant formula</i>	Infant formula was made with contaminated tap water from copper pipes with lead solder.	CR	Report (with environmental sampling data) of a 13-month-old child.	46, 47
<i>Tamarind candy (Mexico)</i>	Tamarind candy jam products from Mexico. During the manufacturing process, the candied jam is packaged in stoneware or terra cotta ceramic jars that can leach lead.	CR	Report of two children under 6 years old, six older children, and one adult.	14
<b>Beverage Containers</b>				
<i>Bulk-water storage tank</i>	Lead leached from soldered seams and brass fittings in bulk-water storage tanks.	CR	Report of three children aged 6, 12, and 14 months.	11
<i>Ceramic glaze</i>	Lead in ceramic glaze can leach into stored beverages, especially juices since they are acidic. The risk is highest for improperly fired containers.	CR	Multiple reports.	7, 12, 49
<i>Cocktail glass</i>	Lead leached from cocktail glass.	CR	Report of a family with one adult and children aged 4, 5, and 14 years.	22
<i>Iranian urn (samovar)</i>	Lead spot solder from the original manufacturing process leached into water used to make baby formula.	CR	Reports of a 10-week-old child with seizures and of a 4-month-old child.	33, 48

Exposure Source	Description/Exposure Pathway	Study Type*	Study Description	Ref. #
<i>Lead-soldered kettle</i>	Lead leached into infant formulas.	CR	Reports of a 3-month-old child and of a 1-day-old child.	40
Folk Remedies				
<i>Azarcon</i>	Also known as <b>alarcon, coral, luiga, maria luisa, or rueda</b> . Bright orange powder used to treat empacho (an illness believed to be caused by something stuck in the gastrointestinal tract, resulting in diarrhea and vomiting). Azarcon is 95% lead.	E	Report of 15-month-old and 3-year-old siblings who expired with seizures and a subsequent survey of 545 systematically selected households for azarcon and greta usage.	16, 15, 18, 4, 51
<i>Ayurvedic medicine (Tibet)</i>	Unnamed traditional medicine.	CR	Single case.	15
<i>Ba-Baw-San (China)</i>	Herbal medicine used to treat colic pain or to pacify young children.	E	Study of 319 children aged 1 to 7 years demonstrated that consumption was associated with increased BLLs ( $p=.038$ ).	21
<i>Bint Al Zahab (Iran)</i>	Rock ground into a powder and mixed with honey and butter given to newborn babies for colic and early passage of meconium after birth.	CR	Report of six children aged 2 days to 3 months.	44
<i>Bint Dahab</i> <i>(Saudi Arabia: means "gold")</i>	A yellow lead oxide used by local jewelers and as a home remedy.	CR	Report of 10 children aged 7 days to 13 months, including three who took bint dahab.	35

"daughter of

Exposure Source	Description/Exposure Pathway	Study Type*	Study Description	Ref. #
<i>Bokhoor (Kuwait)</i>	A traditional practice of burning wood and lead sulphide to produce pleasant fumes to calm infants.	CR	Report of four children aged 16 days to 4.5 months.	27
<i>Ghasard</i>	Brown powder used as a tonic to aid in digestion.	CR	Report of a 9-month-old child who died.	17
<i>Greta (Mexico)</i>	Yellow powder used to treat empacho (see <b>azarcon</b> ); can be obtained through pottery suppliers, as it is also used as a glaze for low-fired ceramics. Greta is 97% lead.	E	See azarcon.	4, 16, 18, 51
<i>Jin Bu Huan (China)</i>	An herbal medicine used to relieve pain.	CR	Report of three children aged 13 and 23 months and 2.5 years.	10
<i>Pay-loo-ah (Vietnam)</i>	A red powder given to children to cure fever or rash.	CR	Report of a 6-month-old child.	13, 15
<i>Po Ying Tan (China)</i>	An herbal medicine used to treat minor ailments in children.	CR	Report of a 4-month-old child.	20
<i>Santrinj (Saudi Arabia)</i>	An amorphous red powder containing 98% lead oxide used principally as a primer for paint for metallic surfaces, but also as a home remedy for “gum boils” and “teething.”	CR	Report of 10 children aged 7 days to 13 months, including 7 who took santrinj.	35
<i>Surma (India)</i>	Black powder used as a cosmetic and as teething powder.	E	A case-control study of 62 children demonstrated higher BLLs in children using surma (p<.001).	2, 15
<i>Tibetan herbal vitamin</i>	Used to strengthen the brain.	CR	Report of a 5-year-old child.	38

Exposure Source	Description/Exposure Pathway	Study Type*	Study Description	Ref. #
<i>Traditional Saudi medicine</i>	Orange powder prescribed by a traditional medicine practitioner for teething; also has an antidiarrheal effect.	CR	Report of three children aged 11, 22, and 44 months.	1
Miscellaneous				
<i>Automobile key-chain emblem</i>	Ingestion of lead-containing automobile key-chain emblem.	CR	Report of a 23-month-old child.	5
<i>Clothing accessory</i>	Ingestion of a "simulated watch."	CR	Report of 3-year-old child who required endoscopy.	26
<i>Curtain weights</i>	Ingestion of lead-containing curtain weights.	CR	Report of deaths of a 23-month-old child and a 2-year-old child.	30, 6
<i>Fishing sinkers</i>	Ingestion of a lead-containing fishing sinker.	CR	Report of an 8-year-old.	39
<i>Gasoline sniffing</i>	Lead in gasoline absorbed through gasoline sniffing.	CR	Report of six of seven siblings aged 10 to 17 years.	9, 24
<i>Lead bullet</i>	Lead absorbed from a retained bullet.	CR	Report of one adult and review of 18 other cases including seven children under 2 years old.	23, 32
<i>Lead pellets</i>	Ingestion of lead pellets from pellet gun.	CR	Report of a 6-year-old child.	45
<i>Lead shot (boat keel)</i>	Lead shot used in a toy boat keel that was eaten by a child.	CR	Report of a 4-year-old child.	28
<i>Lead shot (boat keel) and toy newsprint fireplace log</i>	Lead inhaled during burning of a log made from old newsprint.	CR	Report of a 6-month-old child.	43

Exposure Source	Description/Exposure Pathway	Study Type*	Study Description	Ref. #
<i>Pool cue chalk</i>	Lead contained in pool cue chalk.	CR	Report of two children aged 28 and 27 months.	36
<i>Vinyl miniblinds</i>	Lead dust from vinyl miniblinds.	E	A study of 92 children aged 6 to 72 months attributed 9% of lead poisoning cases to vinyl miniblind exposure.	41

\*CR = case report, E = epidemiological study

**References**

1. Abu Melha A, Ahmed NA, el Hassan AY. Traditional remedies and lead intoxication. *Trop Geogr Med* 1987;39:100-3.
2. Ali AR, Smales OR, Aslam M. Surma and lead poisoning. *BMJ* 1978;2:915-6.
3. Al-Saleh I, Nester M, DeVol E, et al. Determinants of blood lead levels in Saudi Arabian schoolgirls. *Int J Occup Environ Health* 1999;5:107-14.
4. Baer RD, De Alba JG, Cueto LM, et al. Lead-based remedies for empacho: patterns and consequences. *Soc Sci Med* 1989;29:1373-9.
5. Biehusen FC, Pulaski EJ. Lead poisoning after ingestion of a foreign body retained in the stomach. *N Engl J Med* 1956;254:1179-81.
6. Blank E, Howieson J. Lead poisoning from a curtain weight. *JAMA*. 1983;249:2176-7.
7. Browder AA. Lead poisoning from glazes. *Ann Intern Med* 1972;76:665.
8. Carney JK, Garbarino KM. Childhood lead poisoning from apple cider. *Pediatrics* 1997;100:1048-9.
9. CDC. Epidemiologic notes and reports: Gasoline sniffing and lead toxicity among siblings—Virginia. *MMWR* 1985;34:449-50,455.
10. CDC. Epidemiologic notes and reports - Jin Bu Huan toxicity in children—Colorado, 1993. *MMWR* 1993;42:633-6.
11. CDC. Epidemiologic notes and reports: Lead-contaminated drinking water in bulk-water storage tanks—Arizona and California, 1993. *MMWR* 1994; 43:751, 757-8.
12. CDC. Epidemiologic notes and reports: Lead poisoning following ingestion of homemade beverage stored in a ceramic jug. *MMWR* 1989;38:379-80.
13. CDC. Folk Remedy-associated lead poisoning in Hmong children—Minnesota. *MMWR* 1983;32:555-6.
14. CDC. Lead poisoning associated with imported candy and powdered food coloring—California and Michigan. *MMWR* 1998;47:1041-3.

15. CDC. Lead poisoning associated with use of traditional ethnic remedies—California, 1991-1992. *MMWR* 1993;42:521-4.
16. CDC. Lead poisoning from Mexican folk remedies—California. *MMWR* 1983;32:554-5.
17. CDC. Lead poisoning-associated death from Asian Indian folk remedies—Florida. *MMWR* 1984;33:638,643-5.
18. CDC. Use of lead tetroxide as a folk remedy for gastrointestinal illness. *MMWR* 1981;30:546-7.
19. CDC. Occupational and take-home lead poisoning associated with restoring chemically stripped furniture—California, 1998. *MMWR* 2001;50:246-8.
20. Chan H, Billmeier GJ Jr, Evans WE, et al. Lead poisoning from ingestion of Chinese herbal medicine. *Clin Toxicol* 1977;10:273-81.
21. Cheng TJ, Wong RH, Lin YP, et al. Chinese herbal medicine, sibship, and blood lead in children. *Occup Environ Med* 1998;55:573-6.
22. Dickinson L, Reichert EL, Ho RC, et al. Lead poisoning in a family due to cocktail glasses. *Am J Med* 1972;52:391-4.
23. Dillman RO, Crumb CK, Lidsky MJ. Lead poisoning from a gunshot wound. Report of a case and review of the literature. *Am J Med* 1979;66:509-14.
24. Eastwell HD. Elevated lead levels in petrol “sniffers.” *Med J Aust* 1985;143(9 suppl):S63-S64.
25. Eisenberg A, Avni A, Grauer F, et al. Identification of community flour mills as the source of lead poisoning in West Bank Arabs. *Arch Intern Med* 1985;145:1848-51.
26. Esernio-Jenssen D, Donatelli-Guagenti A, Mofenson HC. Severe lead poisoning from an imported clothing accessory: “watch” out for lead. *J Toxicol Clin Toxicol* 1996;34:329-33.
27. Fernando NP, Healy MA, Aslam M, et al. Lead poisoning and traditional practices: the consequences for world health. A study in Kuwait. *Public Health* 1981;95:250-60.

## Appendixes

28. Gittleman JL, Engalgau MM, Shaw J, et al. Lead poisoning among battery reclamation workers in Alabama. *J Occup Med* 1994;36:526-32.
29. Greensher J, Mofenson HC, Balakrishnan C, et al. Leading poisoning from ingestion of lead shot. *Pediatrics* 1974;54:641-3.
30. Hugelmeyer CD, Moorhead JC, Horenblas L, et al. Fatal lead encephalopathy following foreign body ingestion: case report. *J Emerg Med* 1988;6:397-400.
31. Kaye WE, Novotny TE, Tucker M. New ceramics-related industry implicated in elevated blood lead levels in children. *Arch Environ Health* 1987;42:161-4.
32. Kikano GE, Stange KC. Lead poisoning in a child after a gunshot injury. *J Fam Pract* 1992;34:498-500, 502, 504.
33. Lockitch G, Berry B, Roland E, et al. Seizures in a 10-week-old infant: lead poisoning from an unexpected source. *CMAJ* 1991;145:1465-8.
34. Lussenhop DH, Parker DL, Barklind A, et al. Lead exposure and radiator repair work. *Am J Public Health* 1989;79:1558-60.
35. McNiel JR, Reinhard MC. Lead poisoning from home remedies. *Clin Pediatr (Phila)* 1967;6:150-6.
36. Miller MB, Curry SC, Kunkel DB, et al. Pool cue chalk: a source of environmental lead. *Pediatrics* 1996;97:916-7.
37. Mojdehi GM, Gurtner J. Childhood lead poisoning through kohl. *Am J Public Health* 1996;86:587-8.
38. Moore C, Adler R. Herbal vitamins: lead toxicity and developmental delay. *Pediatrics* 2000;106:600-2.
39. Mowad E, Haddad I, Gemmel DJ. Management of lead poisoning from ingested fishing sinkers. *Arch Pediatr Adolesc Med* 1998;152:485-8.
40. Ng R, Martin DJ. Lead poisoning from lead-soldered electric kettles. *Can Med Assoc J* 1977;116:508-9.

41. Norman EH, Hertz-Picciotto I, Salmen DA, et al. Childhood lead poisoning and vinyl miniblind exposure. *Arch Pediatr Adolesc Med* 1997;151:1033-7.
42. Nunez CM, Klitzman S, Goodman A. Lead exposure among automobile radiator repair workers and their children in New York City. *Am J Ind Med* 1993;23:763-77.
43. Perkins KC, Oski FA. Elevated blood lead in a 6-month-old breast-fed infant: the role of newsprint logs. *Pediatrics* 1976;57:426-7.
44. Rahman H, Al Khayat A, Menon N. Lead poisoning in infancy—unusual causes in the U.A.E. *Ann Trop Paediatr* 1986;6:213-7.
45. Roberts JR, Landers KM, Fargason CA Jr. An unusual source of lead poisoning. *Clin Pediatr (Phila)* 1998;37:377-9.
46. Shannon M, Graef J. Hazard of lead in infant formula (letter). *N Engl J Med* 1992;326:137.
47. Shannon M, Graef JW. Lead intoxication from lead-contaminated water used to reconstitute infant formula. *Clin Pediatr (Phila)* 1989;28:380-2.
48. Shannon M. Lead poisoning from an unexpected source in a 4-month-old infant. *Environ Health Perspect* 1998;106:313-6.
49. Sitarz AL. Letter: Severe lead poisoning in a 6-month-old infant. *J Pediatr* 1975;86:810-1.
50. Sprinkle RV. Leaded eye cosmetics: a cultural cause of elevated lead levels in children. *J Fam Pract* 1995;40:358-62.
51. Trotter RT 2d. Greta and Azarcon: a survey of episodic lead poisoning from a folk remedy. *Hum Organ* 1985;44:64-72.
52. Warley MA, Blackledge P, O’Gorman P. Lead poisoning from eye cosmetic. *BMJ* 1968;1:117.
53. Whelan EA, Piacitelli GM, Gerwel B, et al. Elevated blood lead levels in children of construction workers. *Am J Public Health* 1997;87:1352-5.

## Appendix II. Sources of Information on Lead Abatement

- The Office of Lead Hazard Control of the U.S. Department of Housing and Urban Development (HUD) publishes a manual that explains how renovations and remodeling projects can be safely conducted. Single copies of this publication (*Lead Paint Safety—A Field Guide for Painting, Home Maintenance and Renovation Work*) can be ordered from the National Lead Information Center at 1-800-424-5323 or downloaded from the HUD Office of Lead Hazard Control Web site at [www.hud.gov/lea/leahome.html](http://www.hud.gov/lea/leahome.html).
- The Center for National Lead-Safe Housing provides information about safe repair. The Web site is [www.leadshousing.org/html/tech\\_assistance.htm](http://www.leadshousing.org/html/tech_assistance.htm).
- The Alliance to End Childhood Lead Poisoning provides information about safe repair. The Web site is [www.aeclp.org/painting/index.html](http://www.aeclp.org/painting/index.html).
- HUD offers a web-based, one-hour training course on how to visually assess the condition of paint films. The Web site is [www.hud.gov/lea/lbptraining.html](http://www.hud.gov/lea/lbptraining.html).

