

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
TENTH REVISION, INTERNATIONAL
CLASSIFICATION OF DISEASES,
1992), HISPANIC ORIGIN,
RACE FOR NON-HISPANIC
POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
All causes											
All origins 3/	854.0	736.7	32.4	15.8	20.3	67.1	93.6	91.2	111.0	159.2	239.1
Male	853.0	806.5	35.9	17.6	24.2	93.3	138.0	128.8	147.9	205.4	305.9
Female	855.0	663.4	28.7	14.0	16.1	39.3	47.1	52.8	73.6	113.2	173.4
Hispanic	303.8	596.3	29.6	14.1	17.5	61.6	83.3	76.7	93.2	129.4	192.2
Male	331.3	637.1	31.5	15.0	21.3	90.5	123.1	110.3	131.1	172.5	259.1
Female	274.6	553.6	27.5	13.2	13.6	28.7	34.6	37.5	49.8	82.4	122.2
Mexican	257.3	585.7	30.2	14.3	17.9	64.2	84.1	76.6	88.4	120.3	172.8
Male	283.3	590.1	32.2	15.7	21.0	95.8	124.8	112.9	125.0	158.9	229.5
Female	228.6	580.5	28.2	12.7	14.3	28.3	32.6	34.8	44.4	75.0	108.7
Puerto Rican	437.4	*	28.5	18.6	17.0	51.2	84.4	91.6	150.4	243.3	365.9
Male	482.9	*	*	*	*	62.0	124.8	129.1	212.9	336.9	551.2
Female	391.0	*	*	*	*	38.0	*	53.5	91.1	158.8	212.5
Cuban	803.6	*	*	*	*	*	*	*	*	105.5	174.7
Male	842.0	*	*	*	*	*	*	*	*	*	*
Female	765.7	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	298.2	536.8	29.2	11.8	15.9	61.0	81.9	73.4	89.5	117.1	178.0
Male	328.5	*	28.7	12.5	21.9	91.5	119.8	97.9	127.5	161.8	246.8
Female	269.3	*	29.9	11.1	11.3	27.3	40.3	40.5	48.4	74.5	116.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
All causes											
Non-Hispanic 5/	929.6	760.4	32.9	16.1	20.7	67.7	95.3	93.8	113.5	162.4	242.8
Male	928.1	836.8	36.8	18.1	24.7	93.4	140.9	132.2	150.3	208.8	309.0
Female	931.0	680.1	28.8	14.0	16.4	41.0	49.3	55.6	77.1	116.8	178.2
Non-Hispanic white	993.2	596.5	28.5	14.7	19.4	64.0	84.1	80.6	98.7	144.5	216.4
Male	978.5	658.7	32.4	16.6	23.3	86.1	122.9	112.7	132.2	188.0	278.5
Female	1,007.3	530.9	24.4	12.6	15.2	41.0	44.5	48.0	64.7	100.8	154.5
Non-Hispanic black	805.5	1,477.8	51.7	22.6	27.5	89.9	160.5	175.2	213.6	290.5	443.1
Male	859.5	1,623.2	56.1	25.4	33.0	134.0	249.2	256.9	283.2	371.1	558.6
Female	756.7	1,327.7	47.1	19.7	21.7	44.9	76.5	101.3	151.1	219.2	341.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Septicemia (A40-A41)

Non-Hispanic 5/	12.1	6.9	0.7	0.2	0.1	0.2	0.3	0.5	0.9	1.5	2.5
Male	10.7	6.7	0.7	0.2	*	0.2	0.4	0.7	1.0	1.5	2.6
Female	13.5	7.0	0.6	*	*	*	*	0.4	0.9	1.5	2.4
Non-Hispanic white	11.9	4.8	0.5	0.2	*	0.2	0.2	0.4	0.7	1.1	2.0
Male	10.4	4.7	0.5	*	*	*	0.3	0.4	0.7	1.2	2.1
Female	13.4	4.8	0.6	*	*	*	*	*	0.6	1.0	1.9
Non-Hispanic black	16.7	16.6	1.1	*	*	*	0.8	1.5	2.6	3.8	5.9
Male	15.3	16.4	*	*	*	*	*	2.2	2.6	3.6	6.0
Female	17.9	16.9	*	*	*	*	*	*	2.6	4.1	5.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/ YEAR 2/ YEARS	UNDER 1 YEAR 2/ YEARS	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
------------------------------------	-----------------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Non-Hispanic 5/	5.0	*	*	*	0.1	0.2	0.8	3.4	8.7	12.5	12.9
Male	7.7	*	*	*	*	*	0.8	3.8	11.7	18.6	19.5
Female	2.5	*	*	*	*	*	0.8	3.0	5.7	6.4	6.4
Non-Hispanic white	2.3	*	*	*	*	*	0.2	1.1	3.5	6.3	6.1
Male	4.0	*	*	*	*	*	*	1.5	5.6	10.6	10.2
Female	0.7	*	*	*	*	*	*	0.7	1.5	2.1	1.9
Non-Hispanic black	22.0	*	*	*	*	0.7	3.9	15.8	38.4	51.0	58.2
Male	31.7	*	*	*	*	*	3.8	17.1	49.4	73.0	85.3
Female	13.1	*	*	*	*	*	4.0	14.6	28.5	31.5	34.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms (C00-C97)

All origins 3/	196.5	2.4	2.7	2.4	2.6	3.7	5.1	7.0	12.5	24.5	48.8
Male	207.2	2.6	3.0	2.6	2.9	4.3	5.9	7.1	11.3	21.2	44.4
Female	186.2	2.3	2.5	2.2	2.2	3.0	4.2	7.0	13.7	27.8	53.2
Hispanic	59.9	*	2.5	2.5	3.4	3.8	4.5	5.8	9.0	18.2	33.9
Male	61.3	*	2.9	3.0	4.1	4.3	4.8	5.7	8.2	13.8	28.0
Female	58.5	*	2.2	2.0	2.6	3.4	4.1	5.8	9.9	23.0	40.0
Mexican	48.6	*	2.6	2.4	3.4	4.0	4.2	5.2	8.6	18.5	30.1
Male	49.4	*	2.9	2.9	4.3	4.7	4.5	5.7	8.2	13.0	24.9
Female	47.7	*	2.4	1.8	2.3	3.2	3.8	4.7	9.1	25.0	36.0
Puerto Rican	81.3	*	*	*	*	*	*	10.1	11.8	19.4	43.7
Male	86.2	*	*	*	*	*	*	*	*	*	*
Female	76.2	*	*	*	*	*	*	*	*	*	*
Cuban	173.8	*	*	*	*	*	*	*	*	*	*
Male	196.8	*	*	*	*	*	*	*	*	*	*
Female	151.1	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	64.9	*	*	*	3.7	4.0	4.4	5.7	9.1	16.4	39.5
Male	64.6	*	*	*	*	*	*	*	*	11.5	30.5
Female	65.2	*	*	*	*	*	*	*	11.3	21.1	47.6

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms (C00-C97)

Non-Hispanic 5/	215.6	2.5	2.7	2.3	2.4	3.6	5.2	7.2	13.1	25.3	50.3
Male	228.7	2.6	2.9	2.5	2.6	4.3	6.2	7.3	11.8	22.3	46.2
Female	203.1	2.4	2.6	2.2	2.2	3.0	4.2	7.2	14.3	28.3	54.4
Non-Hispanic white	232.2	2.6	2.8	2.4	2.4	3.5	5.1	7.3	12.7	24.0	46.9
Male	244.4	2.9	3.1	2.5	2.6	4.1	6.1	7.4	11.8	21.5	42.9
Female	220.6	2.4	2.4	2.4	2.2	2.9	4.1	7.1	13.5	26.5	50.9
Non-Hispanic black	174.9	*	2.8	2.2	2.4	4.5	6.1	8.3	16.3	35.4	76.1
Male	194.9	*	2.6	2.5	2.7	5.5	7.4	7.6	13.3	28.3	71.8
Female	156.9	*	3.0	1.8	2.0	3.6	4.8	8.9	18.9	41.8	79.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/ YEAR 2/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

Non-Hispanic 5/	22.4	*	*	*	*	*	0.1	0.4	1.0	2.2	4.0
Male	22.8	*	*	*	*	*	*	0.5	1.1	2.5	4.6
Female	22.1	*	*	*	*	*	*	0.4	0.8	1.8	3.5
Non-Hispanic white	24.0	*	*	*	*	*	*	0.4	0.9	1.9	3.6
Male	24.2	*	*	*	*	*	*	0.4	1.1	2.3	4.1
Female	23.8	*	*	*	*	*	*	0.4	0.6	1.5	3.2
Non-Hispanic black	19.4	*	*	*	*	*	*	*	1.6	3.8	6.7
Male	19.8	*	*	*	*	*	*	*	1.8	4.2	8.2
Female	19.1	*	*	*	*	*	*	*	1.4	3.5	5.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Non-Hispanic 5/	61.5	*	*	*	*	*	*	0.2	0.8	3.2	10.0
Male	73.2	*	*	*	*	*	*	0.3	0.8	3.6	11.3
Female	50.4	*	*	*	*	*	*	*	0.9	2.8	8.7
Non-Hispanic white	67.4	*	*	*	*	*	*	0.2	0.9	3.0	9.4
Male	78.9	*	*	*	*	*	*	*	0.8	3.2	10.0
Female	56.4	*	*	*	*	*	*	*	0.9	2.7	8.7
Non-Hispanic black	45.4	*	*	*	*	*	*	*	0.9	5.0	16.1
Male	60.3	*	*	*	*	*	*	*	*	6.0	21.9
Female	31.9	*	*	*	*	*	*	*	*	4.1	10.9

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS	
Malignant neoplasm of breast (C50)												
Non-Hispanic 5/	16.5	*	*	*	*	*	*	0.4	2.0	4.5	8.6	
Male	0.3	*	*	*	*	*	*	*	*	*	*	
Female	31.8	*	*	*	*	*	*	0.8	4.0	8.8	16.9	
Non-Hispanic white	17.4	*	*	*	*	*	*	0.4	1.7	4.0	7.7	
Male	0.4	*	*	*	*	*	*	*	*	*	*	
Female	33.8	*	*	*	*	*	*	0.7	3.4	7.8	15.3	
Non-Hispanic black	15.3	*	*	*	*	*	*	0.8	3.8	8.0	15.2	
Male	0.4	*	*	*	*	*	*	*	*	*	*	
Female	28.9	*	*	*	*	*	*	*	7.2	14.8	28.4	

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasm of cervix uteri (C53)

Non-Hispanic 5/	1.5	*	*	*	*	*	*	0.4	0.8	1.2	1.7
Female	3.0	*	*	*	*	*	*	0.8	1.5	2.4	3.4
Non-Hispanic white	1.4	*	*	*	*	*	*	0.4	0.7	1.1	1.4
Female	2.7	*	*	*	*	*	*	0.8	1.4	2.1	2.9
Non-Hispanic black	2.5	*	*	*	*	*	*	*	1.5	2.1	3.2
Female	4.7	*	*	*	*	*	*	*	2.8	4.0	6.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Non-Hispanic 5/	21.8	1.0	0.9	0.8	0.9	1.6	2.1	2.2	2.6	3.5	4.8
Male	23.7	*	1.0	1.0	1.0	1.9	2.6	2.3	3.2	4.0	5.7
Female	20.1	*	0.9	0.6	0.8	1.2	1.5	2.1	2.0	2.9	3.9
Non-Hispanic white	24.2	1.3	1.0	0.8	1.0	1.5	1.9	2.1	2.5	3.3	4.6
Male	26.2	*	1.1	1.0	1.1	1.8	2.5	2.2	3.0	3.9	5.7
Female	22.3	*	0.9	0.7	0.9	1.1	1.4	1.9	2.0	2.8	3.5
Non-Hispanic black	14.3	*	*	0.6	0.7	2.0	2.7	2.9	3.2	4.5	6.4
Male	15.2	*	*	*	*	2.4	3.3	2.7	4.6	4.8	6.7
Female	13.5	*	*	*	*	1.5	2.1	3.0	1.9	4.3	6.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Leukemia (C91-C95)											
Non-Hispanic 5/	8.2	1.0	0.9	0.7	0.7	1.1	1.1	1.2	1.2	1.6	1.9
Male	9.3	*	0.9	0.8	0.9	1.4	1.4	1.3	1.5	1.6	2.1
Female	7.2	*	0.9	0.5	0.6	0.9	0.7	1.1	0.9	1.5	1.7
Non-Hispanic white	9.2	1.3	0.9	0.7	0.8	1.0	1.1	1.1	1.1	1.5	1.8
Male	10.4	*	1.0	0.8	0.9	1.2	1.4	1.2	1.3	1.6	2.1
Female	8.1	*	0.9	0.6	0.7	0.8	0.7	1.0	0.8	1.5	1.6
Non-Hispanic black	4.9	*	*	*	0.6	1.5	1.3	1.5	1.9	1.7	2.5
Male	5.5	*	*	*	*	2.1	1.8	*	2.6	1.5	2.5
Female	4.4	*	*	*	*	*	*	1.6	*	1.9	2.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97
)

Non-Hispanic 5/	24.4	0.7	0.8	0.6	0.6	1.2	1.5	1.5	2.0	3.6	6.1
Male	25.2	*	0.9	0.6	0.7	1.3	2.0	2.0	2.4	4.0	6.7
Female	23.7	*	0.8	0.6	0.6	1.0	1.0	1.1	1.7	3.2	5.5
Non-Hispanic white	26.5	*	0.8	0.7	0.6	1.1	1.6	1.7	2.0	3.4	5.6
Male	27.2	*	1.0	0.7	0.7	1.3	2.0	2.2	2.5	3.8	6.0
Female	25.8	*	0.6	0.7	0.6	0.9	1.1	1.1	1.6	2.9	5.2
Non-Hispanic black	18.9	*	0.9	*	0.7	1.6	1.4	1.3	2.0	5.4	10.1
Male	20.0	*	*	*	*	1.8	2.3	1.6	2.1	5.6	12.0
Female	17.8	*	*	*	*	1.4	*	*	1.9	5.2	8.5

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Anemias (D50-D64)

Non-Hispanic 5/	1.8	*	0.3	0.1	0.1	0.2	0.3	0.3	0.5	0.4	0.5
Male	1.3	*	*	*	*	*	0.4	0.3	0.5	0.3	0.4
Female	2.2	*	*	*	*	*	0.3	0.3	0.5	0.5	0.6
Non-Hispanic white	1.7	*	*	*	*	*	*	*	*	*	0.1
Male	1.2	*	*	*	*	*	*	*	*	*	*
Female	2.1	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black	2.7	*	*	*	*	*	1.7	1.8	2.9	2.3	2.7
Male	2.4	*	*	*	*	*	1.7	1.6	2.8	2.0	2.3
Female	3.1	*	*	*	*	*	1.7	2.0	3.0	2.6	3.1

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Diabetes mellitus (E10-E14)

Non-Hispanic 5/	25.9	*	*	*	*	0.3	0.7	1.2	2.1	3.2	5.6
Male	24.2	*	*	*	*	0.3	0.8	1.3	2.6	3.7	6.9
Female	27.6	*	*	*	*	*	0.6	1.0	1.7	2.7	4.4
Non-Hispanic white	25.4	*	*	*	*	0.2	0.5	0.9	1.8	2.8	5.0
Male	24.3	*	*	*	*	*	0.7	1.0	2.3	3.2	6.3
Female	26.5	*	*	*	*	*	0.4	0.8	1.3	2.4	3.7
Non-Hispanic black	34.0	*	*	*	*	*	1.5	2.9	4.3	6.2	10.4
Male	28.3	*	*	*	*	*	1.6	3.3	4.7	7.4	12.0
Female	39.1	*	*	*	*	*	1.5	2.6	3.9	5.1	8.9

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Major cardiovascular diseases (I00-I78)

All origins 3/	332.9	16.7	1.5	0.7	1.1	2.4	4.3	6.7	12.2	24.2	49.6
Male	315.8	17.5	1.6	0.7	1.2	2.8	5.0	8.2	15.3	31.3	68.0
Female	349.5	15.9	1.4	0.6	1.0	2.0	3.5	5.2	9.0	17.1	31.5
Hispanic	96.5	13.7	1.3	0.6	0.8	1.9	3.2	4.8	8.2	14.5	29.7
Male	96.0	13.7	1.4	*	1.2	2.4	3.7	6.0	10.4	18.2	40.0
Female	96.9	13.8	*	*	*	*	2.6	3.4	5.7	10.5	18.9
Mexican	76.6	13.4	1.1	*	*	2.1	2.9	4.8	8.5	12.9	28.9
Male	77.4	13.4	*	*	*	2.8	3.3	5.9	10.6	16.2	39.5
Female	75.7	13.5	*	*	*	*	2.4	3.5	6.0	9.0	17.0
Puerto Rican	138.2	*	*	*	*	*	*	*	11.4	28.5	47.9
Male	137.0	*	*	*	*	*	*	*	*	*	*
Female	139.4	*	*	*	*	*	*	*	*	*	*
Cuban	351.7	*	*	*	*	*	*	*	*	*	*
Male	346.2	*	*	*	*	*	*	*	*	*	*
Female	357.2	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	93.0	*	*	*	*	*	4.7	4.4	5.6	13.9	23.3
Male	93.0	*	*	*	*	*	*	6.2	7.6	19.0	30.5
Female	93.0	*	*	*	*	*	*	*	*	9.0	16.9

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Major cardiovascular diseases (I00-I78)

Non-Hispanic 5/	365.6	17.3	1.6	0.7	1.1	2.5	4.5	7.0	12.8	25.4	51.5
Male	347.7	18.3	1.7	0.8	1.2	2.9	5.4	8.6	16.2	33.0	70.8
Female	382.6	16.3	1.5	0.6	1.1	2.1	3.7	5.5	9.5	18.0	32.7
Non-Hispanic white	395.5	14.6	1.4	0.6	0.9	2.1	3.5	5.6	10.4	21.4	43.9
Male	373.6	15.5	1.6	0.7	1.1	2.4	4.3	7.0	13.7	28.9	62.9
Female	416.5	13.6	1.1	0.4	0.8	1.7	2.8	4.2	7.0	13.8	25.0
Non-Hispanic black	294.2	28.8	2.5	1.0	1.9	4.8	9.5	15.7	28.8	52.8	106.7
Male	284.3	30.0	2.3	*	1.9	5.3	10.8	18.9	33.9	63.5	133.3
Female	303.1	27.5	2.9	*	2.0	4.4	8.4	12.9	24.2	43.3	83.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Diseases of heart (I00-I09,I11,I13,I20-I51)											
Non-Hispanic 5/	277.3	13.3	1.2	0.5	0.8	2.1	3.5	5.6	10.2	20.3	41.3
Male	275.2	13.7	1.4	0.6	0.9	2.4	4.3	7.0	13.4	27.7	59.7
Female	279.3	12.8	1.1	0.4	0.7	1.7	2.7	4.2	7.0	12.9	23.3
Non-Hispanic white	301.3	11.4	1.1	0.4	0.7	1.7	2.7	4.3	8.3	17.6	36.5
Male	297.5	12.0	1.2	0.5	0.8	1.9	3.3	5.5	11.4	24.9	54.8
Female	304.9	10.8	0.9	0.3	0.5	1.4	2.0	3.1	5.2	10.2	18.2
Non-Hispanic black	218.6	20.5	2.0	0.9	1.4	4.2	7.8	13.4	22.3	39.4	78.8
Male	217.6	20.1	2.0	*	1.5	4.9	9.2	16.4	27.6	50.0	102.5
Female	219.5	20.8	2.0	*	1.4	3.4	6.5	10.6	17.5	30.0	58.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS	
Hypertensive heart disease (I11)												
Non-Hispanic 5/	9.2	*	*	*	*	*	0.2	0.4	0.8	1.5	3.0	
Male	7.8	*	*	*	*	*	0.3	0.4	1.0	1.9	4.0	
Female	10.6	*	*	*	*	*	*	0.3	0.5	1.0	2.1	
Non-Hispanic white	8.3	*	*	*	*	*	*	0.2	0.4	0.9	2.0	
Male	6.6	*	*	*	*	*	*	*	0.6	1.3	2.9	
Female	9.9	*	*	*	*	*	*	*	*	0.5	1.2	
Non-Hispanic black	16.6	*	*	*	*	*	*	1.4	2.9	5.1	9.9	
Male	16.1	*	*	*	*	*	*	1.6	3.7	6.0	12.6	
Female	17.1	*	*	*	*	*	*	*	2.2	4.3	7.6	

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Ischemic heart diseases (I20-I25)											
Non-Hispanic 5/	200.5	0.8	*	*	*	0.2	0.5	1.3	3.7	10.6	24.7
Male	207.6	*	*	*	*	*	0.7	2.1	5.5	15.8	38.9
Female	193.8	*	*	*	*	*	0.3	0.6	1.9	5.4	10.9
Non-Hispanic white	220.2	*	*	*	*	0.2	0.5	1.1	3.5	10.2	23.4
Male	227.6	*	*	*	*	*	0.6	1.8	5.3	15.5	37.8
Female	213.1	*	*	*	*	*	*	0.5	1.6	4.9	9.0
Non-Hispanic black	144.4	*	*	*	*	*	0.8	2.6	6.2	15.0	37.7
Male	145.7	*	*	*	*	*	*	3.9	8.2	20.8	52.5
Female	143.2	*	*	*	*	*	*	*	4.4	9.9	24.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Acute myocardial infarction (I21-I22)											
Non-Hispanic 5/	75.3	*	*	*	*	*	0.2	0.6	1.4	4.3	10.5
Male	80.3	*	*	*	*	*	*	0.9	2.0	6.3	16.4
Female	70.6	*	*	*	*	*	*	0.3	0.8	2.3	4.7
Non-Hispanic white	82.7	*	*	*	*	*	0.2	0.5	1.4	4.4	10.1
Male	88.5	*	*	*	*	*	*	0.7	2.0	6.5	16.2
Female	77.2	*	*	*	*	*	*	*	0.7	2.2	3.9
Non-Hispanic black	54.0	*	*	*	*	*	*	1.2	2.3	5.1	15.3
Male	53.7	*	*	*	*	*	*	1.7	2.7	6.7	20.9
Female	54.2	*	*	*	*	*	*	*	1.9	3.7	10.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Other forms of chronic ischemic heart disease (I20,I25)

Non-Hispanic 5/	123.9	*	*	*	*	*	0.3	0.7	2.2	6.1	14.0
Male	126.0	*	*	*	*	*	0.4	1.2	3.4	9.3	22.0
Female	122.0	*	*	*	*	*	*	*	1.1	3.0	6.1
Non-Hispanic white	136.1	*	*	*	*	*	0.2	0.6	2.0	5.7	13.1
Male	137.7	*	*	*	*	*	0.4	1.0	3.2	8.8	21.2
Female	134.6	*	*	*	*	*	*	*	0.9	2.6	4.9
Non-Hispanic black	89.1	*	*	*	*	*	*	1.3	3.7	9.7	21.9
Male	90.4	*	*	*	*	*	*	2.2	5.1	13.7	30.8
Female	88.0	*	*	*	*	*	*	*	2.5	6.0	14.1

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Atherosclerotic cardiovascular disease, so described (I25.0)

Non-Hispanic 5/	26.9	*	*	*	*	*	*	0.2	0.6	1.8	4.6
Male	28.2	*	*	*	*	*	*	0.3	1.0	2.9	7.4
Female	25.6	*	*	*	*	*	*	*	0.3	0.7	1.9
Non-Hispanic white	27.6	*	*	*	*	*	*	*	0.6	1.6	4.2
Male	28.4	*	*	*	*	*	*	*	0.9	2.7	6.9
Female	26.9	*	*	*	*	*	*	*	*	0.6	1.5
Non-Hispanic black	29.3	*	*	*	*	*	*	*	0.9	3.0	7.6
Male	33.2	*	*	*	*	*	*	*	*	4.8	11.2
Female	25.8	*	*	*	*	*	*	*	*	1.5	4.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Non-Hispanic 5/	97.0	*	*	*	*	*	0.2	0.5	1.6	4.3	9.4
Male	97.8	*	*	*	*	*	0.3	0.9	2.4	6.4	14.7
Female	96.3	*	*	*	*	*	*	*	0.8	2.3	4.2
Non-Hispanic white	108.4	*	*	*	*	*	*	0.5	1.5	4.0	8.9
Male	109.2	*	*	*	*	*	*	0.8	2.3	6.1	14.3
Female	107.7	*	*	*	*	*	*	*	0.7	2.0	3.5
Non-Hispanic black	59.8	*	*	*	*	*	*	0.9	2.8	6.6	14.3
Male	57.3	*	*	*	*	*	*	*	3.9	8.9	19.6
Female	62.1	*	*	*	*	*	*	*	1.9	4.6	9.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Other heart diseases (I26-I51)											
Non-Hispanic 5/	65.1	12.4	1.1	0.5	0.8	1.8	2.8	3.8	5.5	7.9	12.8
Male	58.0	12.7	1.2	0.6	0.9	2.1	3.3	4.4	6.6	9.6	16.1
Female	71.8	12.0	1.0	0.4	0.7	1.5	2.3	3.2	4.4	6.2	9.6
Non-Hispanic white	70.3	10.5	0.9	0.4	0.6	1.4	2.1	2.9	4.4	6.2	10.6
Male	61.6	10.9	1.1	0.5	0.7	1.6	2.5	3.4	5.4	7.8	13.6
Female	78.7	10.1	0.8	*	0.5	1.2	1.6	2.4	3.4	4.7	7.6
Non-Hispanic black	54.6	19.6	2.0	0.8	1.4	3.8	6.3	9.1	12.4	18.5	29.1
Male	53.3	19.1	2.0	*	1.5	4.5	7.1	10.5	14.9	22.4	35.3
Female	55.8	20.1	1.9	*	1.3	3.1	5.5	7.8	10.2	15.0	23.6

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)											
Non-Hispanic 5/	42.3	11.0	0.9	0.4	0.6	1.6	2.5	3.4	5.0	7.2	11.4
Male	40.0	10.9	1.0	0.5	0.7	1.9	3.0	3.9	6.1	8.8	14.3
Female	44.5	11.0	0.8	0.2	0.5	1.3	2.1	2.9	4.0	5.6	8.6
Non-Hispanic white	44.9	9.6	0.8	0.3	0.5	1.2	1.9	2.6	4.1	5.7	9.5
Male	41.7	9.6	0.8	0.4	0.6	1.4	2.3	3.1	5.0	7.1	12.0
Female	47.9	9.6	0.7	*	0.3	1.0	1.5	2.1	3.1	4.3	6.9
Non-Hispanic black	39.5	16.1	1.5	0.7	1.1	3.4	5.7	8.3	11.3	16.5	25.7
Male	40.5	15.0	1.7	*	1.2	4.0	6.4	9.5	13.7	20.0	31.2
Female	38.6	17.3	*	*	*	2.7	5.1	7.1	9.0	13.4	20.9

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

Non-Hispanic 5/	7.0	*	*	*	*	*	*	*	0.3	0.5	1.1
Male	5.4	*	*	*	*	*	*	*	0.3	0.5	1.4
Female	8.5	*	*	*	*	*	*	*	0.2	0.4	0.8
Non-Hispanic white	6.5	*	*	*	*	*	*	*	*	0.2	0.6
Male	4.9	*	*	*	*	*	*	*	*	0.3	0.8
Female	8.1	*	*	*	*	*	*	*	*	*	0.4
Non-Hispanic black	11.0	*	*	*	*	*	*	*	1.1	2.1	4.2
Male	9.5	*	*	*	*	*	*	*	*	2.2	5.4
Female	12.4	*	*	*	*	*	*	*	*	2.1	3.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Cerebrovascular diseases (I60-I69)											
Non-Hispanic 5/	65.4	3.6	0.3	0.1	0.3	0.4	0.7	1.1	1.8	3.9	7.6
Male	51.5	4.2	*	*	*	0.4	0.7	1.1	1.8	3.8	7.7
Female	78.7	2.8	0.4	*	0.3	0.3	0.7	1.0	1.9	4.1	7.6
Non-Hispanic white	70.0	2.6	0.3	*	0.2	0.4	0.6	1.0	1.4	2.9	5.6
Male	53.9	3.3	*	*	*	0.5	0.6	1.1	1.4	2.9	5.7
Female	85.5	1.9	*	*	*	*	0.6	0.9	1.4	3.0	5.5
Non-Hispanic black	54.3	8.0	*	*	*	*	1.1	1.6	4.5	9.9	20.2
Male	47.7	9.2	*	*	*	*	*	*	4.2	9.5	21.0
Female	60.3	*	*	*	*	*	*	1.7	4.7	10.2	19.5

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Influenza and pneumonia (J10-J18)

Non-Hispanic 5/	25.4	7.5	0.7	0.2	0.2	0.4	0.7	0.8	1.1	1.8	3.0
Male	22.7	8.5	0.7	0.2	*	0.4	0.7	0.9	1.2	1.9	3.8
Female	27.9	6.5	0.6	*	*	0.3	0.6	0.6	1.0	1.8	2.2
Non-Hispanic white	27.9	5.0	0.5	0.2	0.2	0.3	0.6	0.6	1.0	1.7	2.5
Male	24.5	5.8	0.5	*	*	0.4	0.7	0.7	1.0	1.7	3.2
Female	31.3	4.2	0.6	*	*	*	0.5	0.6	1.0	1.7	1.9
Non-Hispanic black	16.9	17.5	1.2	*	*	*	1.1	1.4	2.1	2.9	6.3
Male	17.3	18.8	1.7	*	*	*	*	2.1	2.6	3.2	8.6
Female	16.5	16.2	*	*	*	*	*	*	1.7	2.7	4.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Pneumonia (J12-J18)											
Non-Hispanic 5/	24.7	7.3	0.6	0.2	0.1	0.3	0.7	0.7	1.1	1.8	2.9
Male	22.2	8.2	0.7	*	*	0.4	0.7	0.8	1.2	1.9	3.7
Female	27.1	6.4	0.5	*	*	0.3	0.6	0.6	1.0	1.8	2.2
Non-Hispanic white	27.1	4.8	0.5	0.2	0.2	0.3	0.6	0.6	1.0	1.7	2.4
Male	23.8	5.4	0.4	*	*	0.3	0.7	0.6	1.0	1.7	3.1
Female	30.3	4.2	0.5	*	*	*	0.5	0.6	0.9	1.6	1.8
Non-Hispanic black	16.6	17.2	1.2	*	*	*	1.1	1.4	2.1	2.9	6.2
Male	17.1	18.8	1.7	*	*	*	*	2.1	2.6	3.2	8.3
Female	16.3	15.5	*	*	*	*	*	*	1.7	2.7	4.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/ YEAR 2/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Chronic lower respiratory diseases (J40-J47)

Non-Hispanic 5/	48.3	1.0	0.3	0.2	0.5	0.5	0.6	0.7	0.9	1.4	2.9
Male	48.7	*	0.4	0.3	0.5	0.6	0.7	0.8	1.0	1.4	2.8
Female	48.0	*	*	*	0.4	0.3	0.5	0.6	0.8	1.5	3.0
Non-Hispanic white	55.7	*	*	*	0.2	0.3	0.4	0.5	0.6	1.1	2.4
Male	55.1	*	*	*	*	0.3	0.5	0.5	0.6	0.9	2.4
Female	56.2	*	*	*	*	*	0.4	0.5	0.7	1.3	2.5
Non-Hispanic black	21.5	*	0.9	0.7	1.4	1.5	1.4	1.9	2.5	3.6	6.3
Male	25.1	*	*	*	1.8	2.1	1.6	2.4	3.3	3.8	6.1
Female	18.2	*	*	*	*	*	*	1.5	1.7	3.4	6.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Asthma (J45-J46)											
Non-Hispanic 5/	1.7	*	0.2	0.2	0.4	0.4	0.5	0.6	0.8	0.9	1.3
Male	1.2	*	*	0.2	0.5	0.5	0.6	0.7	0.8	0.8	1.1
Female	2.1	*	*	*	0.4	0.3	0.5	0.5	0.7	1.0	1.5
Non-Hispanic white	1.4	*	*	*	0.2	0.2	0.4	0.4	0.5	0.5	0.9
Male	1.0	*	*	*	*	*	0.4	0.4	0.5	0.4	0.7
Female	1.9	*	*	*	*	*	0.3	*	0.5	0.7	1.0
Non-Hispanic black	3.3	*	*	0.7	1.4	1.3	1.2	1.8	2.1	2.7	3.7
Male	2.8	*	*	*	1.7	1.8	*	2.3	2.7	2.9	3.2
Female	3.7	*	*	*	*	*	*	*	1.5	2.7	4.1

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Non-Hispanic 5/	9.8	7.5	0.6	0.2	0.2	0.3	0.4	0.5	0.8	1.0	1.8
Male	9.8	8.4	0.6	0.2	*	0.3	0.5	0.5	0.8	1.1	1.9
Female	9.9	6.6	0.5	*	*	0.3	0.3	0.6	0.7	0.9	1.6
Non-Hispanic white	10.8	5.4	0.5	0.2	0.2	0.3	0.3	0.5	0.7	0.8	1.6
Male	10.7	6.6	0.5	*	*	*	0.4	0.4	0.7	0.9	1.7
Female	11.0	4.1	0.5	*	*	*	*	0.5	0.6	0.7	1.5
Non-Hispanic black	6.7	17.7	0.8	*	*	*	0.8	0.9	1.4	2.4	3.1
Male	6.8	17.1	*	*	*	*	*	*	*	3.0	3.5
Female	6.6	18.3	*	*	*	*	*	*	*	1.9	2.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Non-Hispanic 5/	9.4	*	*	*	*	*	0.1	0.5	1.4	4.6	9.7
Male	12.4	*	*	*	*	*	*	0.6	1.7	5.9	13.3
Female	6.7	*	*	*	*	*	*	0.3	1.2	3.3	6.1
Non-Hispanic white	10.0	*	*	*	*	*	*	0.3	1.3	4.4	9.4
Male	13.1	*	*	*	*	*	*	0.5	1.6	5.7	13.1
Female	7.0	*	*	*	*	*	*	*	1.0	3.1	5.8
Non-Hispanic black	7.7	*	*	*	*	*	*	0.8	1.4	5.0	11.2
Male	10.3	*	*	*	*	*	*	*	*	6.8	14.8
Female	5.4	*	*	*	*	*	*	*	1.5	3.5	8.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS	
Alcoholic liver disease (K70)												
Non-Hispanic 5/	4.2	*	*	*	*	*	*	0.3	1.0	3.1	6.2	
Male	6.2	*	*	*	*	*	*	0.4	1.2	4.0	8.6	
Female	2.3	*	*	*	*	*	*	*	0.7	2.3	3.9	
Non-Hispanic white	4.3	*	*	*	*	*	*	0.2	0.9	3.0	6.0	
Male	6.5	*	*	*	*	*	*	0.4	1.2	3.8	8.3	
Female	2.2	*	*	*	*	*	*	*	0.6	2.2	3.7	
Non-Hispanic black	3.9	*	*	*	*	*	*	*	1.0	3.2	7.1	
Male	5.5	*	*	*	*	*	*	*	*	4.4	9.9	
Female	2.5	*	*	*	*	*	*	*	*	2.2	4.6	

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Non-Hispanic 5/	14.4	4.4	*	*	*	0.1	0.3	0.5	0.8	1.3	2.1
Male	14.1	5.4	*	*	*	*	0.3	0.5	0.9	1.5	2.5
Female	14.7	3.4	*	*	*	*	0.4	0.4	0.7	1.2	1.7
Non-Hispanic white	14.2	2.7	*	*	*	*	0.3	0.4	0.5	0.8	1.4
Male	14.1	3.5	*	*	*	*	*	0.4	0.6	0.9	1.7
Female	14.2	1.8	*	*	*	*	*	*	0.4	0.7	1.2
Non-Hispanic black	19.5	12.0	*	*	*	*	*	1.1	2.7	4.5	6.5
Male	18.3	13.6	*	*	*	*	*	*	3.0	5.3	8.2
Female	20.7	10.2	*	*	*	*	*	*	2.5	3.9	5.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Renal failure (N17-N19)											
Non-Hispanic 5/	14.1	4.2	*	*	*	0.1	0.3	0.4	0.7	1.2	2.0
Male	13.8	5.2	*	*	*	*	*	0.4	0.8	1.4	2.4
Female	14.4	3.1	*	*	*	*	0.3	0.4	0.7	1.1	1.7
Non-Hispanic white	13.9	2.5	*	*	*	*	0.2	0.3	0.4	0.8	1.3
Male	13.7	3.3	*	*	*	*	*	0.3	0.5	0.8	1.6
Female	14.0	*	*	*	*	*	*	*	0.4	0.7	1.1
Non-Hispanic black	19.2	11.4	*	*	*	*	*	1.0	2.6	4.2	6.3
Male	17.9	13.6	*	*	*	*	*	*	2.8	5.1	8.0
Female	20.3	9.2	*	*	*	*	*	*	2.4	3.5	4.9

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Pregnancy, childbirth and the puerperium (000-099)

All origins 3/	0.1	*	*	0.4	0.4	0.5	0.4	0.2
Female	0.3	*	*	0.8	0.9	0.9	0.8	0.3
Hispanic	0.2	*	*	0.6	0.6	*	*	*
Female	0.5	*	*	1.4	1.3	*	*	*
Mexican	0.2	*	*	*	*	*	*	*
Female	0.5	*	*	*	*	*	*	*
Puerto Rican	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
Cuban	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
Other Hispanic 4/	0.3	*	*	*	*	*	*	*
Female	0.6	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Pregnancy, childbirth and the puerperium (000-099)

Non-Hispanic 5/	0.1	*	*	0.4	0.4	0.5	0.3	0.2
Female	0.3	*	*	0.7	0.8	0.9	0.7	0.3
Non-Hispanic white	0.1	*	*	0.2	0.3	0.3	0.2	*
Female	0.2	*	*	0.4	0.5	0.6	0.5	*
Non-Hispanic black	0.4	*	*	1.1	1.2	1.3	0.9	*
Female	0.8	*	*	2.2	2.3	2.5	1.7	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Pregnancy with abortive outcome (000-007)

All origins 3/	0.0	*	*	*	*	*	*	*
Female	0.0	*	*	*	*	*	*	*
Hispanic	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
Mexican	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
Puerto Rican	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
Cuban	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
Other Hispanic 4/	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy with abortive outcome (000-007)

Non-Hispanic 5/	0.0	*	*	*	*	*	*	*
Female	0.0	*	*	*	*	*	*	*
Non-Hispanic white	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
Non-Hispanic black	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Other complications of pregnancy, childbirth and the puerperium (010-099)

All origins 3/	0.1	*	*	0.4	0.4	0.4	0.3	0.1
Female	0.3	*	*	0.8	0.8	0.9	0.7	0.3
Hispanic	0.2	*	*	0.6	*	*	*	*
Female	0.5	*	*	1.3	*	*	*	*
Mexican	0.2	*	*	*	*	*	*	*
Female	0.5	*	*	*	*	*	*	*
Puerto Rican	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
Cuban	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
Other Hispanic 4/	0.3	*	*	*	*	*	*	*
Female	0.6	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Other complications of pregnancy, childbirth and the puerperium (O10-099)

Non-Hispanic 5/	0.1	*	*	0.3	0.4	0.4	0.3	0.1
Female	0.2	*	*	0.7	0.7	0.8	0.6	0.3
Non-Hispanic white	0.1	*	*	0.2	0.2	0.3	0.2	*
Female	0.1	*	*	0.4	0.5	0.6	0.4	*
Non-Hispanic black	0.3	*	*	1.0	1.0	1.1	0.7	*
Female	0.7	*	*	1.9	1.9	2.2	1.4	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Non-Hispanic 5/	3.6	148.8	3.1	1.0	1.0	1.2	1.2	1.2	1.4	1.3	1.4
Male	3.9	154.4	3.1	1.0	1.0	1.4	1.4	1.3	1.7	1.4	1.5
Female	3.4	142.8	3.0	0.9	1.1	1.0	1.0	1.0	1.1	1.1	1.2
Non-Hispanic white	3.4	141.2	2.6	0.9	1.0	1.0	1.2	1.2	1.5	1.3	1.3
Male	3.6	147.9	2.6	0.8	0.9	1.1	1.5	1.3	1.8	1.4	1.5
Female	3.2	134.2	2.6	0.9	1.0	0.9	1.0	1.0	1.2	1.1	1.2
Non-Hispanic black	5.0	184.1	5.2	1.3	1.4	1.9	1.3	1.4	1.4	1.6	1.7
Male	5.5	186.0	5.6	1.7	1.3	2.6	*	1.8	1.7	1.7	2.0
Female	4.5	182.2	4.8	*	1.5	*	*	*	*	1.5	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Non-Hispanic 5/	12.0	102.4	1.6	0.4	0.4	1.6	2.9	3.4	4.3	5.5	7.5
Male	11.9	118.0	1.7	0.5	0.5	1.9	3.9	4.5	5.6	7.2	10.0
Female	12.1	86.1	1.5	0.3	0.4	1.2	1.9	2.2	2.9	3.8	5.1
Non-Hispanic white	12.2	79.7	1.4	0.3	0.4	1.5	2.8	3.1	4.0	5.4	7.2
Male	11.8	91.8	1.7	0.4	0.4	1.9	3.8	4.2	5.4	7.2	9.8
Female	12.6	66.9	1.2	*	0.4	1.0	1.7	2.0	2.6	3.5	4.7
Non-Hispanic black	13.9	205.6	2.3	0.6	0.8	2.2	4.3	5.5	6.4	7.3	11.3
Male	15.7	236.8	2.2	*	*	2.1	5.7	7.3	8.2	8.5	13.3
Female	12.4	173.4	2.4	*	*	2.2	3.0	3.8	4.8	6.3	9.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)											
Non-Hispanic 5/	69.3	29.6	4.0	1.9	2.6	3.8	5.8	6.8	9.3	14.3	22.5
Male	56.9	33.6	4.4	1.9	2.7	4.3	6.9	7.9	10.7	16.8	26.5
Female	81.0	25.4	3.5	1.8	2.4	3.3	4.6	5.6	7.9	11.8	18.6
Non-Hispanic white	74.3	24.0	3.4	1.7	2.4	3.7	5.4	5.8	8.1	12.3	19.9
Male	59.6	26.7	3.8	1.9	2.7	4.3	6.6	7.3	9.7	14.9	23.9
Female	88.3	21.2	2.9	1.6	2.1	3.0	4.1	4.3	6.6	9.7	15.9
Non-Hispanic black	60.1	51.3	6.1	2.4	3.5	4.9	8.6	13.4	17.8	27.8	42.7
Male	56.1	61.4	6.4	2.1	3.2	5.1	9.4	13.6	18.9	30.9	48.8
Female	63.7	40.9	5.8	2.7	3.7	4.8	7.8	13.2	16.8	25.0	37.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All origins 3/	34.8	23.1	11.9	6.8	7.7	33.4	38.8	30.4	28.6	32.9	35.4
Male	46.2	25.7	14.0	7.8	9.9	45.8	58.9	47.0	43.1	48.9	53.1
Female	23.8	20.5	9.7	5.7	5.5	20.3	17.8	13.5	13.9	17.0	18.0
Hispanic	25.0	15.3	10.4	5.7	5.6	27.9	35.5	27.9	26.6	30.3	33.1
Male	36.9	17.0	11.6	5.7	6.5	40.8	54.8	44.0	41.7	46.3	53.9
Female	12.4	13.5	9.2	5.6	4.7	13.1	11.9	9.0	9.4	12.9	11.3
Mexican	25.1	16.1	11.3	6.2	6.1	30.1	37.2	28.3	27.5	30.7	31.8
Male	36.7	16.4	12.5	6.5	7.1	44.0	57.6	45.0	43.0	45.8	50.0
Female	12.2	15.8	10.0	5.9	5.1	14.3	11.2	9.2	8.9	12.9	11.2
Puerto Rican	25.1	*	*	*	*	16.0	30.9	27.7	31.9	40.1	45.6
Male	37.7	*	*	*	*	*	44.5	*	51.3	61.8	*
Female	12.3	*	*	*	*	*	*	*	*	*	*
Cuban	26.2	*	*	*	*	*	*	*	*	*	*
Male	34.3	*	*	*	*	*	*	*	*	*	*
Female	18.1	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	24.6	*	8.0	3.9	4.4	26.5	32.3	26.1	24.4	28.2	32.5
Male	37.6	*	*	*	*	41.7	49.9	39.7	38.5	45.1	56.9
Female	12.2	*	*	*	*	9.8	12.9	*	9.1	12.1	10.6

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Non-Hispanic 5/	36.0	25.0	12.2	7.0	8.1	34.3	39.3	30.8	28.8	33.1	35.4
Male	47.3	27.7	14.5	8.3	10.5	46.6	59.6	47.4	43.1	49.0	52.5
Female	25.2	22.2	9.7	5.6	5.6	21.5	18.9	14.3	14.6	17.5	18.6
Non-Hispanic white	37.3	20.1	11.1	6.3	8.0	37.3	41.4	31.1	29.1	33.4	35.0
Male	48.1	22.9	13.4	7.5	10.2	50.0	62.6	47.2	43.4	49.3	51.5
Female	26.9	17.1	8.7	5.0	5.6	24.0	19.6	14.7	14.6	17.5	18.5
Non-Hispanic black	34.4	46.5	16.8	10.0	9.1	24.3	35.5	34.2	32.8	36.7	44.6
Male	50.3	49.5	19.4	12.2	12.5	35.3	54.5	55.9	51.1	56.9	69.9
Female	20.1	43.3	14.2	7.8	5.5	13.0	17.5	14.6	16.5	18.8	22.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Transport accidents (V01-V99,Y85)											
All origins 3/	16.6	4.6	4.5	4.1	5.3	27.1	29.3	20.4	16.8	17.1	16.8
Male	23.3	4.8	5.0	4.4	6.4	35.5	43.3	30.8	24.7	24.8	24.8
Female	10.2	4.3	4.0	3.8	4.1	18.1	14.6	9.8	8.7	9.5	9.0
Hispanic	14.8	5.4	5.2	4.0	4.3	21.9	27.0	19.4	15.0	15.6	16.9
Male	21.6	6.1	5.6	3.8	5.2	31.3	40.4	30.2	22.1	22.9	25.6
Female	7.6	*	4.8	4.1	3.3	11.3	10.7	6.8	6.8	7.8	7.7
Mexican	15.9	6.5	5.8	4.6	4.8	23.9	29.1	20.3	17.3	17.1	17.4
Male	23.1	*	6.4	4.5	5.8	33.9	44.0	32.0	26.0	24.2	25.3
Female	8.0	*	5.3	4.7	3.7	12.5	10.3	7.0	6.8	8.7	8.5
Puerto Rican	10.2	*	*	*	*	12.7	18.7	15.7	11.1	11.2	*
Male	14.4	*	*	*	*	*	*	*	*	*	*
Female	6.0	*	*	*	*	*	*	*	*	*	*
Cuban	14.7	*	*	*	*	*	*	*	*	*	*
Male	21.3	*	*	*	*	*	*	*	*	*	*
Female	8.3	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	13.2	*	*	*	*	19.7	23.3	17.3	11.4	14.2	17.4
Male	19.9	*	*	*	*	30.6	34.8	25.8	15.7	21.8	28.3
Female	6.9	*	*	*	*	*	10.8	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Transport accidents (V01-V99,Y85)											
Non-Hispanic 5/	16.8	4.3	4.3	4.1	5.4	27.9	29.7	20.5	17.0	17.3	16.7
Male	23.4	4.5	4.8	4.5	6.6	36.2	43.9	30.8	25.1	24.9	24.5
Female	10.5	4.2	3.8	3.7	4.2	19.3	15.3	10.4	9.0	9.8	9.1
Non-Hispanic white	17.1	3.5	3.8	4.0	5.6	30.9	30.9	20.2	17.0	17.4	16.6
Male	23.7	3.8	4.3	4.5	6.8	39.6	45.6	29.7	24.8	24.8	24.1
Female	10.9	3.2	3.4	3.5	4.4	21.8	15.9	10.5	9.0	10.0	9.1
Non-Hispanic black	16.4	6.9	5.9	4.6	5.0	17.8	27.7	24.4	20.0	18.4	19.7
Male	24.8	7.2	6.4	5.0	6.7	24.6	41.7	39.5	31.7	29.5	31.6
Female	8.8	*	5.4	4.2	3.2	10.8	14.4	10.7	9.6	8.6	9.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL UNDER 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44
 AGES 1/ YEAR 2/ YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86
 ,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

All origins 3/	15.4	4.4	4.2	3.8	4.8	26.0	28.0	19.2	15.5	15.6	15.1
Male	21.3	4.6	4.7	4.1	5.8	33.9	41.2	28.6	22.6	22.1	21.9
Female	9.7	4.2	3.8	3.5	3.8	17.5	14.2	9.5	8.2	9.1	8.4
Hispanic	13.9	5.2	4.9	3.7	4.0	21.2	26.0	18.5	14.0	14.4	15.2
Male	20.1	6.1	5.2	3.5	4.8	30.2	38.7	28.6	20.7	20.6	22.9
Female	7.2	*	4.6	4.0	3.1	10.9	10.4	6.7	6.4	7.6	7.1
Mexican	15.0	6.2	5.4	4.3	4.6	23.1	28.2	19.4	16.2	16.1	15.8
Male	21.7	*	5.7	4.1	5.4	32.8	42.5	30.3	24.4	22.4	23.0
Female	7.6	*	5.1	4.5	3.5	12.1	10.0	6.8	6.4	8.6	7.7
Puerto Rican	9.6	*	*	*	*	12.1	17.6	15.3	11.1	*	*
Male	13.1	*	*	*	*	*	*	*	*	*	*
Female	6.0	*	*	*	*	*	*	*	*	*	*
Cuban	13.5	*	*	*	*	*	*	*	*	*	*
Male	18.8	*	*	*	*	*	*	*	*	*	*
Female	8.1	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	12.3	*	*	*	*	18.9	22.2	16.6	10.4	12.9	15.1
Male	18.3	*	*	*	*	29.7	32.8	24.5	14.3	19.8	24.2
Female	6.6	*	*	*	*	*	10.5	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)											
Non-Hispanic 5/	15.6	4.2	4.1	3.8	5.0	26.8	28.3	19.2	15.6	15.7	15.0
Male	21.4	4.2	4.5	4.2	6.0	34.5	41.6	28.5	22.8	22.3	21.6
Female	10.0	4.2	3.6	3.4	3.9	18.7	14.9	10.0	8.5	9.3	8.5
Non-Hispanic white	15.8	3.3	3.7	3.7	5.1	29.7	29.4	18.7	15.5	15.8	14.8
Male	21.5	3.4	4.1	4.2	6.1	37.9	43.0	27.2	22.4	22.1	21.1
Female	10.3	3.2	3.2	3.2	4.1	21.1	15.4	10.1	8.5	9.5	8.5
Non-Hispanic black	15.5	6.9	5.6	4.3	4.7	16.9	26.9	23.7	18.6	16.9	18.3
Male	23.1	7.2	6.0	4.5	6.4	23.2	40.3	38.0	29.3	26.9	29.0
Female	8.5	*	5.2	4.1	3.1	10.5	14.2	10.6	9.0	8.2	8.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Nontransport accidents (W00-X59,Y86)											
Non-Hispanic 5/	19.2	20.7	7.9	2.9	2.7	6.4	9.7	10.3	11.8	15.8	18.7
Male	23.9	23.2	9.7	3.7	3.8	10.4	15.7	16.7	18.0	24.1	28.0
Female	14.7	18.0	6.0	1.9	1.4	2.2	3.6	3.9	5.6	7.7	9.6
Non-Hispanic white	20.1	16.6	7.3	2.3	2.3	6.4	10.5	10.9	12.1	16.0	18.4
Male	24.5	19.1	9.1	3.0	3.4	10.4	17.0	17.5	18.6	24.5	27.4
Female	16.0	14.0	5.4	1.5	1.2	2.3	3.8	4.2	5.6	7.5	9.3
Non-Hispanic black	18.0	39.5	10.9	5.4	4.1	6.5	7.8	9.8	12.8	18.3	24.9
Male	25.5	42.3	12.9	7.2	5.8	10.7	12.8	16.4	19.4	27.5	38.2
Female	11.3	36.6	8.8	3.6	2.4	2.2	3.1	3.9	6.9	10.2	13.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
Accidental exposure to smoke, fire and flames (X00-X09)											
Non-Hispanic 5/	1.3	1.0	2.2	1.0	0.5	0.4	0.7	0.7	0.7	0.9	1.0
Male	1.6	*	2.6	1.2	0.4	0.4	0.9	0.9	0.9	1.3	1.2
Female	1.0	*	1.7	0.8	0.5	0.3	0.5	0.5	0.5	0.5	0.7
Non-Hispanic white	1.1	*	1.5	0.7	0.4	0.4	0.6	0.6	0.6	0.8	0.9
Male	1.4	*	1.8	0.8	0.3	0.4	0.9	0.7	0.8	1.1	1.1
Female	0.9	*	1.1	0.6	0.4	*	0.4	0.4	0.4	0.4	0.6
Non-Hispanic black	2.6	*	5.1	2.2	0.8	*	1.0	1.2	1.3	1.6	2.0
Male	3.2	*	5.8	2.7	*	*	*	*	1.6	2.3	2.6
Female	2.0	*	4.3	1.7	*	*	*	*	*	*	1.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Non-Hispanic 5/	4.6	*	0.2	*	0.1	1.8	4.6	5.3	6.8	9.4	11.2
Male	6.6	*	*	*	*	2.8	7.2	8.3	9.8	13.5	16.2
Female	2.7	*	*	*	*	0.8	1.9	2.2	3.8	5.5	6.3
Non-Hispanic white	4.6	*	*	*	*	2.2	5.4	5.9	7.2	9.7	11.0
Male	6.6	*	*	*	*	3.3	8.6	9.2	10.4	14.0	15.8
Female	2.7	*	*	*	*	1.0	2.0	2.5	3.9	5.4	6.2
Non-Hispanic black	5.7	*	*	*	*	*	2.2	4.3	7.0	10.9	15.6
Male	8.4	*	*	*	*	*	2.8	7.1	9.8	15.0	23.2
Female	3.2	*	*	*	*	*	1.5	1.8	4.4	7.2	8.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/ YEAR 2/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
---------------------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Intentional self-harm (suicide) (X60-X84,Y87.0)

Non-Hispanic 5/	11.1	*	1.6	8.4	13.5	13.1	12.9	14.6	16.1
Male	18.3	*	2.4	13.8	23.3	21.9	20.9	23.2	24.9
Female	4.3	*	0.6	2.8	3.5	4.4	5.0	6.1	7.5
Non-Hispanic white	12.5	*	1.7	9.0	14.3	14.0	14.4	16.4	18.0
Male	20.4	*	2.6	14.8	24.8	23.2	22.9	25.7	27.6
Female	4.9	*	0.7	3.0	3.7	4.7	5.7	7.1	8.5
Non-Hispanic black	5.5	*	1.2	5.7	11.4	10.2	7.9	7.2	8.2
Male	9.6	*	2.1	9.9	20.1	18.2	14.1	12.9	13.7
Female	1.7	*	*	1.5	3.1	3.0	2.3	2.2	3.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (X60-X71,X75-X84,Y87.0)

Non-Hispanic 5/	4.8	*	1.0	3.7	5.5	6.2	6.3	7.3	8.2
Male	6.9	*	1.5	5.6	9.0	9.6	9.7	11.1	11.6
Female	2.7	*	0.4	1.7	2.1	2.8	2.9	3.6	4.9
Non-Hispanic white	5.2	*	0.9	4.0	5.8	6.6	6.9	8.1	9.1
Male	7.5	*	1.4	6.2	9.5	10.3	10.4	12.1	12.6
Female	3.0	*	0.4	1.7	2.1	2.9	3.4	4.0	5.5
Non-Hispanic black	2.4	*	1.1	1.8	4.3	4.0	3.7	3.7	4.8
Male	3.9	*	1.9	2.6	7.1	6.4	6.6	6.3	7.4
Female	1.1	*	*	*	1.7	1.9	*	1.4	2.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Assault (homicide) (X85-Y09,Y87.1)

Non-Hispanic 5/	5.6	9.6	2.4	0.7	1.0	8.4	15.1	11.5	8.6	7.4	6.4
Male	8.6	11.2	2.6	0.7	1.3	13.4	25.5	18.9	13.2	10.4	9.2
Female	2.7	7.9	2.2	0.7	0.8	3.0	4.7	4.2	4.0	4.5	3.6
Non-Hispanic white	2.8	6.2	1.6	0.6	0.7	2.7	4.4	4.0	3.9	4.1	3.8
Male	3.6	8.3	1.8	0.6	0.8	3.5	6.0	5.2	5.2	5.1	5.3
Female	1.9	4.1	1.3	0.5	0.6	1.9	2.8	2.8	2.6	3.1	2.3
Non-Hispanic black	22.1	24.1	6.1	1.2	2.6	34.1	66.4	50.4	34.3	26.6	22.3
Male	38.3	24.9	6.1	*	3.3	59.0	122.1	93.5	59.9	43.8	34.8
Female	7.4	23.3	6.0	1.5	1.8	8.6	13.7	11.4	11.3	11.3	11.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Assault (homicide) by discharge of firearms (X93-X95)

Non-Hispanic 5/	3.6	*	0.2	0.2	0.6	6.7	12.2	9.0	6.0	4.6	3.6
Male	6.0	*	*	0.2	0.9	11.5	21.6	15.7	10.0	7.3	5.5
Female	1.2	*	*	*	0.3	1.7	2.7	2.3	2.1	2.0	1.8
Non-Hispanic white	1.4	*	*	0.2	0.3	1.7	2.7	2.5	2.4	2.4	2.1
Male	2.0	*	*	*	0.5	2.5	4.0	3.5	3.4	3.4	3.0
Female	0.9	*	*	*	*	0.9	1.5	1.4	1.4	1.5	1.3
Non-Hispanic black	16.0	*	*	*	1.8	29.5	57.3	42.5	26.6	17.8	12.9
Male	30.1	*	*	*	2.8	53.3	109.1	82.3	49.4	32.6	22.0
Female	3.3	*	*	*	*	5.2	8.4	6.5	6.2	4.6	4.9

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Assault (homicide) by other and unspecified means and their sequelae (X85-X92,X96-Y09,Y87.1)

Non-Hispanic 5/	2.0	9.2	2.2	0.4	0.4	1.7	3.0	2.6	2.6	2.8	2.8
Male	2.6	10.6	2.4	0.4	0.3	1.9	3.9	3.2	3.2	3.1	3.8
Female	1.5	7.8	2.0	0.5	0.5	1.4	2.0	1.9	1.9	2.5	1.8
Non-Hispanic white	1.3	6.0	1.4	0.3	0.4	1.0	1.7	1.6	1.5	1.7	1.7
Male	1.6	8.1	1.7	0.4	0.3	1.0	2.0	1.7	1.8	1.8	2.3
Female	1.0	3.9	1.2	*	0.4	1.0	1.4	1.4	1.2	1.7	1.0
Non-Hispanic black	6.1	22.9	5.7	0.9	0.8	4.6	9.0	7.8	7.7	8.8	9.4
Male	8.2	22.5	5.7	*	*	5.7	13.0	11.2	10.5	11.2	12.8
Female	4.1	23.3	5.8	*	*	3.4	5.3	4.8	5.1	6.7	6.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Non-Hispanic 5/	1.5	1.9	0.3	*	0.2	0.6	1.5	1.7	1.9	2.8	3.3
Male	2.0	1.6	0.4	*	0.4	1.0	2.2	2.5	2.7	3.9	4.5
Female	0.9	2.2	*	*	*	0.3	0.7	0.9	1.2	1.7	2.1
Non-Hispanic white	1.4	1.5	0.2	*	0.2	0.7	1.5	1.8	2.0	2.8	3.2
Male	2.0	*	*	*	0.4	1.1	2.2	2.6	2.7	3.9	4.3
Female	0.9	*	*	*	*	0.4	0.8	1.1	1.3	1.8	2.0
Non-Hispanic black	1.9	*	*	*	*	*	1.4	1.7	2.2	3.4	4.9
Male	2.9	*	*	*	*	*	2.6	2.8	3.5	5.0	7.3
Female	1.0	*	*	*	*	*	*	*	*	2.0	2.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-9 YEARS	10-14 YEARS	15-19 YEARS	20-24 YEARS	25-29 YEARS	30-34 YEARS	35-39 YEARS	40-44 YEARS
----------------	--------------------	--------------	--------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Non-Hispanic 5/	1.4	1.9	0.3	*	0.2	0.5	1.2	1.5	1.8	2.7	3.2
Male	1.9	1.6	0.4	*	0.3	0.7	1.8	2.2	2.5	3.8	4.4
Female	0.9	2.2	*	*	*	0.3	0.6	0.9	1.1	1.7	2.0
Non-Hispanic white	1.4	1.5	0.2	*	0.2	0.6	1.3	1.7	1.9	2.7	3.1
Male	1.8	*	*	*	0.3	0.9	1.9	2.3	2.5	3.8	4.1
Female	0.9	*	*	*	*	0.3	0.7	1.0	1.3	1.7	2.0
Non-Hispanic black	1.7	*	*	*	*	*	0.9	1.4	2.0	3.3	4.8
Male	2.6	*	*	*	*	*	1.6	2.2	3.1	4.7	7.2
Female	0.9	*	*	*	*	*	*	*	*	2.0	2.6

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 YEARS
 YEARS YEARS YEARS YEARS YEARS YEARS YEARS YEARS AND OVER

All causes

All origins 3/	352.5	509.0	792.6	1,241.0	1,906.3	2,929.4	4,553.8	7,335.0	15,524.4
Male	456.3	642.2	989.8	1,535.9	2,362.0	3,676.1	5,693.2	9,095.4	17,501.4
Female	252.0	381.3	608.1	973.7	1,515.7	2,341.2	3,760.2	6,296.6	14,719.2
Hispanic	281.1	394.3	612.1	913.8	1,425.9	2,278.4	3,543.6	5,777.6	11,822.9
Male	378.6	519.5	801.5	1,181.3	1,804.0	2,918.9	4,480.2	7,220.4	13,086.2
Female	183.8	274.9	440.6	683.1	1,119.7	1,793.2	2,880.3	4,899.7	11,202.8
Mexican	267.9	364.6	601.8	1,022.7	1,454.8	2,368.7	3,807.5	*	*
Male	351.2	444.0	765.4	1,239.1	1,765.6	2,996.8	*	*	*
Female	180.2	279.0	445.5	814.2	1,192.9	1,875.9	*	*	*
Puerto Rican	463.1	616.4	945.1	1,058.3	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Cuban	225.1	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	251.0	366.9	464.6	710.3	1,330.3	2,182.6	*	*	*
Male	343.9	516.5	650.2	*	*	*	*	*	*
Female	171.2	243.3	324.9	469.8	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
All causes									
Non-Hispanic 5/	357.3	515.8	802.9	1,259.9	1,931.9	2,956.6	4,586.6	7,375.0	15,619.8
Male	460.9	648.3	998.9	1,554.1	2,388.4	3,704.3	5,730.8	9,144.1	17,639.8
Female	257.3	388.8	619.1	992.4	1,539.1	2,366.3	3,790.4	6,332.6	14,802.9
Non-Hispanic white	318.7	470.1	749.9	1,199.4	1,865.4	2,889.5	4,525.0	7,369.2	15,799.7
Male	415.7	588.9	928.4	1,477.3	2,305.7	3,615.5	5,653.4	9,137.4	17,853.2
Female	222.8	353.6	579.3	941.7	1,478.3	2,305.2	3,731.9	6,318.4	14,971.7
Non-Hispanic black	679.7	956.6	1,368.9	1,959.9	2,809.4	4,065.0	5,837.2	8,234.9	14,837.5
Male	869.4	1,248.2	1,802.9	2,498.8	3,559.1	5,258.8	7,462.1	10,491.3	16,944.8
Female	515.5	708.3	1,013.7	1,534.7	2,256.3	3,272.3	4,857.7	7,099.0	14,054.9

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Septicemia (A40-A41)

Non-Hispanic 5/	4.0	6.1	9.5	15.4	24.6	38.8	64.4	105.6	217.5
Male	4.5	7.0	10.2	16.9	26.9	43.6	74.3	122.0	224.8
Female	3.5	5.2	9.0	14.0	22.5	35.0	57.6	95.9	214.5
Non-Hispanic white	3.0	4.8	7.9	12.7	20.9	34.8	58.2	97.7	204.3
Male	3.4	5.3	8.4	14.1	22.9	39.7	66.7	114.0	211.2
Female	2.5	4.2	7.3	11.4	19.1	30.8	52.1	88.1	201.6
Non-Hispanic black	11.3	17.3	24.7	40.5	60.8	86.7	148.0	218.1	395.9
Male	12.8	21.9	27.9	45.9	69.8	96.7	185.0	251.5	432.1
Female	10.0	13.4	22.0	36.2	54.1	80.1	125.7	201.3	382.5

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Viral hepatitis (B15-B19)

Non-Hispanic 5/	5.0	4.6	3.4	3.3	3.9	4.5	5.1	4.9	3.8
Male	7.7	7.0	4.8	3.9	4.2	4.6	6.2	5.9	5.4
Female	2.4	2.3	2.0	2.7	3.6	4.4	4.3	4.3	3.2
Non-Hispanic white	4.7	3.9	2.7	2.7	3.3	4.1	4.9	4.6	3.7
Male	7.3	6.1	3.9	3.3	3.6	4.2	6.0	5.6	5.4
Female	2.1	1.8	1.5	2.3	3.2	4.0	4.0	4.0	3.1
Non-Hispanic black	7.4	8.8	6.9	4.5	5.3	6.3	5.8	7.2	*
Male	11.2	13.8	10.3	5.3	6.9	7.5	*	*	*
Female	4.2	4.5	4.0	3.9	4.1	5.4	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Non-Hispanic 5/	12.0	8.4	5.5	3.7	2.4	1.6	0.8	*	*
Male	19.2	13.8	9.3	6.5	4.1	2.7	1.6	*	*
Female	5.1	3.2	1.9	1.2	0.9	0.7	*	*	*
Non-Hispanic white	5.5	3.7	2.7	1.5	1.0	0.6	0.4	*	*
Male	9.6	6.5	4.9	2.8	2.0	0.9	0.8	*	*
Female	1.4	0.8	0.7	*	*	*	*	*	*
Non-Hispanic black	59.5	46.9	29.6	22.2	14.7	11.5	6.0	*	*
Male	93.8	77.5	51.3	40.2	25.3	21.9	11.6	*	*
Female	29.8	20.8	11.8	8.1	6.9	4.5	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Non-Hispanic 5/	1.3	1.7	2.3	3.6	4.7	7.0	9.9	12.8	21.2
Male	1.6	2.2	2.6	4.0	5.9	8.8	12.1	15.9	27.8
Female	1.1	1.2	2.1	3.2	3.8	5.6	8.4	10.9	18.6
Non-Hispanic white	1.2	1.5	2.2	3.2	4.6	6.8	9.8	12.6	21.0
Male	1.5	1.9	2.5	3.6	5.6	8.6	11.6	15.8	27.0
Female	1.0	1.0	1.9	2.8	3.7	5.3	8.5	10.8	18.6
Non-Hispanic black	2.7	3.5	4.0	6.0	6.2	10.1	11.6	16.1	24.2
Male	3.3	4.8	4.6	7.4	8.8	12.6	16.9	19.7	31.7
Female	2.1	2.3	3.5	4.9	4.3	8.4	8.4	14.3	21.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Malignant neoplasms (C00-C97)									
All origins 3/	93.6	166.2	289.0	463.4	690.6	951.6	1,222.5	1,505.3	1,819.4
Male	93.5	173.9	321.3	535.7	834.7	1,190.4	1,586.0	2,050.4	2,710.7
Female	93.7	158.8	258.9	398.0	567.2	763.5	969.3	1,183.8	1,456.4
Hispanic	64.3	105.4	180.7	272.5	422.5	633.8	851.8	1,138.5	1,383.5
Male	60.9	104.2	199.7	323.3	521.3	820.6	1,140.5	1,637.2	2,049.7
Female	67.6	106.5	163.5	228.8	342.4	492.3	647.3	835.0	1,056.5
Mexican	62.6	97.5	168.8	291.6	405.9	638.3	875.3	*	*
Male	58.7	85.7	184.5	323.5	481.7	822.6	*	*	*
Female	66.7	110.3	153.7	260.9	342.0	493.7	*	*	*
Puerto Rican	83.8	145.2	238.6	271.2	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Cuban	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	61.7	102.3	156.6	240.2	445.0	668.5	*	*	*
Male	50.3	108.6	173.4	*	*	*	*	*	*
Female	71.5	97.1	143.9	181.9	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms (C00-C97)

Non-Hispanic 5/	96.1	170.7	296.5	476.3	706.7	967.3	1,236.6	1,515.9	1,831.1
Male	96.4	179.1	329.4	549.5	852.4	1,207.8	1,602.6	2,061.5	2,732.4
Female	95.8	162.7	265.6	409.7	581.3	777.5	981.8	1,194.4	1,466.6
Non-Hispanic white	90.3	163.6	288.2	470.0	701.3	959.9	1,227.7	1,512.6	1,826.6
Male	90.1	168.5	314.5	535.3	837.1	1,182.9	1,574.8	2,036.1	2,714.1
Female	90.5	158.8	263.2	409.5	582.0	780.3	983.7	1,201.6	1,468.7
Non-Hispanic black	148.6	257.2	415.1	616.2	874.9	1,193.3	1,497.2	1,718.5	2,015.1
Male	155.3	299.9	521.3	783.3	1,154.5	1,675.4	2,183.1	2,709.5	3,290.1
Female	142.8	220.9	328.1	484.4	668.7	873.2	1,083.7	1,219.7	1,541.6

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Non-Hispanic 5/	2.1	3.9	5.8	8.3	10.0	11.7	12.8	14.1	19.5
Male	3.3	6.2	9.6	13.0	15.3	17.1	19.0	20.1	26.5
Female	1.0	1.7	2.2	4.0	5.5	7.5	8.4	10.5	16.6
Non-Hispanic white	1.8	3.3	5.1	7.4	9.5	11.3	12.7	14.1	19.8
Male	2.8	5.0	8.3	11.4	14.1	16.1	18.3	19.7	27.4
Female	0.9	1.6	2.0	3.7	5.5	7.4	8.7	10.7	16.8
Non-Hispanic black	4.1	8.5	12.3	15.6	15.0	16.5	14.5	16.1	14.9
Male	6.3	16.3	22.0	28.1	27.7	29.7	28.9	29.2	*
Female	2.2	*	4.3	5.7	5.7	7.7	5.8	9.5	14.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Malignant neoplasm of esophagus (C15)									
Non-Hispanic 5/	2.5	5.2	9.1	13.7	18.5	22.5	24.7	27.4	26.1
Male	4.5	9.1	15.9	23.8	31.5	38.6	44.2	49.3	47.8
Female	0.7	1.5	2.8	4.5	7.3	9.8	11.1	14.5	17.3
Non-Hispanic white	2.4	4.8	8.4	12.9	17.9	22.2	24.2	27.4	26.7
Male	4.4	8.4	14.7	22.8	30.8	38.2	43.4	50.1	49.4
Female	0.4	1.2	2.4	3.7	6.6	9.3	10.7	13.8	17.5
Non-Hispanic black	4.2	10.0	17.4	23.7	27.5	29.9	36.4	30.5	22.6
Male	6.5	17.8	31.3	39.1	45.6	50.2	66.0	46.3	38.7
Female	2.2	3.4	6.0	11.6	14.2	16.3	18.6	22.5	16.6

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Malignant neoplasm of stomach (C16)									
Non-Hispanic 5/	2.2	3.4	5.4	9.4	13.1	18.7	26.4	35.6	52.0
Male	2.8	4.6	7.6	13.6	18.4	26.8	38.6	49.3	74.5
Female	1.5	2.2	3.4	5.6	8.6	12.3	17.8	27.6	42.9
Non-Hispanic white	1.6	2.8	4.6	8.1	11.0	16.0	22.6	31.2	46.3
Male	2.1	4.0	6.5	11.9	15.7	23.2	33.5	43.8	67.3
Female	1.1	1.7	2.8	4.5	6.9	10.2	15.0	23.8	37.9
Non-Hispanic black	4.8	7.2	11.2	17.6	27.0	38.7	57.3	75.6	102.7
Male	6.7	9.9	16.6	26.4	38.6	57.7	83.3	104.7	138.6
Female	3.1	4.8	6.8	10.7	18.5	26.1	41.5	60.9	89.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

Non-Hispanic 5/	8.4	14.9	25.6	43.0	63.3	94.1	128.6	176.5	273.1
Male	9.3	17.3	31.1	52.0	78.7	117.9	161.4	214.0	318.6
Female	7.5	12.6	20.4	34.8	50.1	75.4	105.7	154.4	254.7
Non-Hispanic white	7.5	13.9	23.9	41.4	61.5	92.3	126.6	173.7	271.9
Male	8.5	16.3	29.0	49.9	76.3	115.6	159.5	209.2	315.8
Female	6.6	11.7	19.1	33.5	48.4	73.5	103.5	152.6	254.1
Non-Hispanic black	15.6	25.4	40.7	63.9	88.5	127.9	169.6	228.8	312.9
Male	17.0	29.9	52.3	78.0	111.1	162.6	211.9	302.1	381.6
Female	14.4	21.6	31.2	52.7	71.8	104.8	144.0	191.8	287.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Non-Hispanic 5/	3.5	5.3	7.4	10.8	15.4	20.5	26.2	30.0	31.7
Male	5.8	8.3	11.2	15.9	22.7	29.5	38.0	44.6	43.4
Female	1.4	2.4	3.9	6.2	9.1	13.4	18.0	21.4	27.0
Non-Hispanic white	2.8	4.2	6.1	9.4	13.8	18.8	24.7	29.2	30.8
Male	4.4	6.4	8.9	13.6	20.3	27.1	36.5	43.5	41.6
Female	1.2	2.0	3.4	5.5	8.0	12.1	16.3	20.8	26.5
Non-Hispanic black	7.1	10.9	13.0	16.2	20.7	27.3	29.2	29.0	31.8
Male	12.6	18.1	22.7	25.9	33.6	42.0	37.1	46.3	48.1
Female	2.3	4.8	5.1	8.6	11.2	17.5	24.4	20.3	25.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of pancreas (C25)

Non-Hispanic 5/	4.2	8.9	15.5	25.2	37.7	51.6	70.9	85.9	96.6
Male	5.5	11.4	18.9	30.6	44.9	57.2	81.4	95.5	103.5
Female	3.1	6.5	12.2	20.3	31.4	47.3	63.7	80.3	93.8
Non-Hispanic white	4.1	8.6	14.9	24.6	36.6	50.5	69.6	84.7	94.7
Male	5.3	11.0	18.2	30.1	44.5	56.0	80.9	94.5	103.0
Female	3.0	6.3	11.7	19.6	29.8	46.0	61.6	78.9	91.4
Non-Hispanic black	5.9	13.5	23.0	33.9	51.1	70.6	94.9	105.2	121.2
Male	7.4	18.0	29.5	41.0	56.8	80.0	102.6	117.6	110.4
Female	4.5	9.7	17.7	28.3	47.0	64.4	90.3	98.9	125.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Malignant neoplasm of larynx (C32)									
Non-Hispanic 5/	0.8	1.6	3.0	4.7	6.1	7.4	8.2	8.0	6.5
Male	1.3	2.6	5.0	8.1	10.0	13.1	15.1	16.7	17.3
Female	0.4	0.6	1.0	1.5	2.7	2.9	3.4	2.8	2.1
Non-Hispanic white	0.7	1.3	2.5	4.0	5.7	6.8	7.6	7.8	6.5
Male	1.1	2.1	4.1	6.9	9.2	11.9	14.1	16.3	17.4
Female	0.3	0.6	0.9	1.4	2.6	2.7	3.1	2.8	2.1
Non-Hispanic black	2.1	4.1	7.7	11.0	10.7	15.5	15.9	11.8	7.0
Male	3.3	7.7	14.7	21.0	20.0	30.4	29.9	27.5	*
Female	*	*	*	*	3.9	5.7	7.6	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Non-Hispanic 5/	22.3	45.9	96.8	167.6	255.5	335.4	376.0	379.5	304.5
Male	27.2	55.6	120.7	210.9	334.0	451.9	525.2	564.0	526.2
Female	17.5	36.6	74.4	128.2	188.0	243.4	272.1	270.8	214.8
Non-Hispanic white	20.8	44.4	96.3	169.9	258.9	339.0	378.9	385.7	306.8
Male	25.0	51.8	116.5	209.8	331.3	448.7	520.4	565.8	527.3
Female	16.7	37.2	77.0	133.0	195.2	250.6	279.5	278.7	218.0
Non-Hispanic black	36.4	70.5	124.8	192.2	284.2	363.5	399.7	347.2	291.0
Male	48.8	102.3	187.8	276.6	438.7	568.2	664.3	623.1	567.1
Female	25.7	43.4	73.3	125.6	170.3	227.5	240.2	208.2	188.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN,

RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of breast (C50)

All origins 3/	13.6	20.8	28.1	35.0	43.4	55.2	70.9	90.9	147.4
Male	*	0.3	0.7	0.8	1.1	1.7	2.1	2.4	4.2
Female	26.6	40.4	53.8	66.0	79.6	97.3	118.8	143.1	205.7
Hispanic	10.4	14.5	21.2	20.7	30.5	32.3	41.3	53.5	71.7
Male	*	*	*	*	*	*	*	*	*
Female	20.7	28.0	39.9	38.2	54.7	55.2	69.5	84.2	105.8
Mexican	10.3	14.5	17.8	23.9	29.7	25.4	34.9	*	*
Male	*	*	*	*	*	*	*	*	*
Female	21.2	30.2	34.8	46.3	53.5	43.9	*	*	*
Puerto Rican	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Cuban	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	10.2	12.7	19.9	20.4	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	19.0	22.6	34.2	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Malignant neoplasm of breast (C50)									
Non-Hispanic 5/	13.9	21.3	28.6	35.9	44.1	56.4	72.0	92.1	149.7
Male	*	0.3	0.7	0.8	1.1	1.7	2.2	2.4	4.2
Female	27.1	41.3	54.8	67.8	81.1	99.6	120.6	145.0	208.6
Non-Hispanic white	12.5	20.1	27.2	35.3	44.0	56.8	72.3	92.9	149.6
Male	*	0.3	0.6	0.8	1.1	1.7	2.2	2.2	4.0
Female	24.7	39.4	52.6	67.2	81.7	101.1	121.5	146.9	208.3
Non-Hispanic black	24.3	33.7	44.9	48.2	53.5	66.5	85.5	101.4	178.7
Male	*	*	*	*	*	*	*	*	*
Female	45.0	61.9	80.5	84.9	92.4	108.9	135.6	150.4	241.6

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasm of cervix uteri (C53)

Non-Hispanic 5/	2.2	2.6	3.0	2.9	3.5	3.7	4.1	5.5	7.1
Female	4.3	5.1	5.8	5.5	6.5	6.7	7.0	8.7	9.9
Non-Hispanic white	1.9	2.3	2.6	2.4	2.8	3.1	3.6	4.8	6.0
Female	3.8	4.5	5.1	4.6	5.2	5.5	6.1	7.6	8.4
Non-Hispanic black	3.8	5.0	6.2	6.6	9.6	10.0	10.9	13.8	19.7
Female	7.0	9.3	11.4	11.7	16.7	16.6	17.4	20.7	27.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Non-Hispanic 5/	1.0	1.8	3.6	6.1	8.6	11.2	14.2	18.8	24.9
Female	1.9	3.5	6.9	11.6	15.9	20.0	24.0	29.9	34.9
Non-Hispanic white	1.0	1.7	3.4	5.4	7.4	9.9	13.1	17.7	24.5
Female	1.9	3.4	6.7	10.5	13.8	17.9	22.3	28.2	34.3
Non-Hispanic black	1.0	2.4	5.3	12.3	20.0	26.5	28.3	35.6	33.7
Female	2.0	4.5	9.7	22.0	34.8	44.0	45.3	53.6	46.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Malignant neoplasm of ovary (C56)									
Non-Hispanic 5/	3.1	5.6	8.5	12.6	17.8	22.1	31.2	36.8	41.2
Female	6.1	11.0	16.5	24.0	33.1	39.5	52.9	58.4	57.8
Non-Hispanic white	3.3	5.8	8.9	13.1	18.3	22.7	32.3	37.9	42.2
Female	6.6	11.4	17.5	25.2	34.4	41.0	54.9	60.4	59.2
Non-Hispanic black	2.2	4.8	6.6	11.2	16.9	19.9	26.6	27.9	34.7
Female	4.1	9.0	11.9	20.0	29.3	33.1	42.7	41.9	47.5

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Malignant neoplasm of prostate (C61)									
Non-Hispanic 5/	0.6	1.5	4.8	12.4	26.5	49.5	85.8	137.4	210.4
Male	1.1	3.1	9.9	26.0	57.3	112.1	209.2	370.6	730.7
Non-Hispanic white	0.4	1.3	4.0	10.6	22.8	43.3	78.1	126.3	199.3
Male	0.8	2.6	8.1	22.0	48.7	97.0	189.3	338.7	693.6
Non-Hispanic black	1.5	3.6	13.3	30.0	66.1	122.3	190.2	295.1	359.6
Male	3.3	7.8	29.5	68.0	155.6	306.5	505.8	881.4	1,328.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Non-Hispanic 5/	2.2	4.6	7.7	10.9	15.2	19.5	24.6	29.5	33.9
Male	3.1	6.8	11.2	15.8	21.7	27.5	34.5	44.1	55.4
Female	1.4	2.4	4.4	6.4	9.6	13.3	17.7	20.9	25.2
Non-Hispanic white	2.3	4.7	7.9	11.2	15.5	19.9	24.9	29.9	34.6
Male	3.2	6.9	11.4	16.2	21.7	27.8	34.9	44.5	56.9
Female	1.4	2.5	4.6	6.4	10.1	13.5	17.9	21.3	25.7
Non-Hispanic black	2.5	4.6	7.5	10.7	15.2	18.7	23.9	29.0	29.6
Male	3.3	7.1	11.5	15.7	25.6	28.4	32.8	48.1	49.3
Female	1.9	2.5	4.3	6.7	7.5	12.3	18.6	19.4	22.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Malignant neoplasm of bladder (C67)									
Non-Hispanic 5/	0.8	1.7	3.7	7.1	12.3	20.1	29.7	46.2	70.3
Male	1.2	2.6	5.7	11.1	19.9	33.0	51.5	83.5	141.3
Female	0.5	0.8	1.7	3.4	5.7	9.9	14.4	24.2	41.6
Non-Hispanic white	0.9	1.8	3.9	7.5	12.9	20.8	30.4	47.1	72.8
Male	1.2	2.8	6.0	11.9	21.1	34.5	53.6	85.9	148.0
Female	0.5	0.8	1.8	3.4	5.8	9.8	14.0	24.1	42.5
Non-Hispanic black	0.9	1.3	3.0	5.5	9.7	18.0	28.8	44.5	49.6
Male	*	*	4.7	7.4	12.8	24.3	41.4	73.0	88.1
Female	*	*	*	4.0	7.5	13.8	21.2	30.2	35.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Non-Hispanic 5/	4.2	6.5	8.6	11.5	15.1	18.8	19.5	21.3	16.9
Male	5.1	8.1	10.6	14.1	18.2	23.3	23.3	26.5	22.3
Female	3.4	4.9	6.7	9.1	12.4	15.2	16.8	18.3	14.7
Non-Hispanic white	4.7	7.2	9.2	12.5	16.4	20.3	20.6	22.1	17.5
Male	5.7	8.7	11.4	15.3	19.8	24.9	24.5	27.7	23.5
Female	3.7	5.6	7.2	9.9	13.4	16.6	17.8	18.8	15.1
Non-Hispanic black	2.2	3.7	5.5	6.2	7.6	9.0	10.5	14.7	10.5
Male	2.2	5.7	6.6	7.9	7.7	12.6	12.0	*	*
Female	2.3	2.0	4.6	4.9	7.5	6.6	9.6	14.3	10.5

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Non-Hispanic 5/	8.4	13.6	23.8	38.4	63.2	92.3	132.7	178.1	206.5
Male	10.2	16.4	28.7	46.0	79.4	119.1	172.1	233.3	285.0
Female	6.6	10.9	19.2	31.4	49.2	71.2	105.3	145.6	174.8
Non-Hispanic white	8.0	13.3	23.6	38.6	64.5	94.1	135.7	183.7	210.7
Male	9.8	15.9	28.4	46.2	81.5	120.4	176.5	240.8	293.8
Female	6.2	10.8	19.0	31.5	49.5	73.0	107.0	149.8	177.2
Non-Hispanic black	12.6	18.0	29.7	44.8	63.5	91.5	115.8	132.8	177.1
Male	15.0	22.4	37.6	55.1	74.9	127.1	149.8	170.8	226.6
Female	10.5	14.3	23.3	36.7	55.0	67.8	95.2	113.6	158.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Non-Hodgkin's lymphoma (C82-C85)									
Non-Hispanic 5/	3.7	5.8	10.8	16.1	25.4	36.3	54.3	72.7	80.9
Male	4.8	7.6	13.4	19.1	31.0	45.3	67.2	89.3	106.7
Female	2.6	4.2	8.3	13.4	20.7	29.1	45.3	62.9	70.4
Non-Hispanic white	3.7	6.0	11.4	16.9	26.8	38.1	57.1	76.9	85.1
Male	4.7	7.6	13.9	19.9	32.8	46.9	70.8	94.1	113.0
Female	2.6	4.4	9.0	14.1	21.6	31.1	47.6	66.6	73.8
Non-Hispanic black	4.5	5.6	9.1	12.8	17.0	22.9	26.3	28.5	36.6
Male	6.2	8.0	13.9	15.9	19.7	32.5	31.3	32.6	43.4
Female	3.1	3.6	5.1	10.4	14.9	16.6	23.2	26.4	34.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Leukemia (C91-C95)									
Non-Hispanic 5/	3.1	4.5	7.8	12.9	22.2	33.8	48.1	66.0	84.7
Male	3.5	5.1	9.6	16.3	30.0	47.3	66.4	94.5	121.5
Female	2.6	4.0	6.0	9.8	15.5	23.1	35.4	49.2	69.8
Non-Hispanic white	2.9	4.5	7.8	13.1	23.2	35.1	49.9	68.5	87.3
Male	3.4	5.0	9.7	16.5	31.3	48.6	68.9	97.9	125.5
Female	2.5	4.0	6.0	9.9	16.0	24.2	36.6	51.1	72.0
Non-Hispanic black	4.7	5.3	8.1	14.3	18.1	29.0	35.3	44.3	66.1
Male	5.1	6.3	10.2	18.9	22.9	44.4	50.6	65.2	98.6
Female	4.3	4.5	6.4	10.6	14.5	18.8	26.1	33.7	54.1

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Non-Hispanic 5/	1.2	2.7	4.6	8.5	14.5	20.5	28.4	36.8	38.0
Male	1.4	3.0	5.0	9.5	17.0	24.5	36.0	46.5	52.5
Female	1.0	2.3	4.2	7.6	12.3	17.4	23.1	31.1	32.1
Non-Hispanic white	1.0	2.3	3.9	7.7	13.3	19.2	26.5	35.6	35.2
Male	1.2	2.6	4.3	8.8	15.8	22.9	34.1	45.7	50.9
Female	0.8	2.0	3.5	6.7	11.1	16.2	21.2	29.7	28.9
Non-Hispanic black	3.0	6.5	11.9	16.9	27.9	38.0	53.8	58.3	72.8
Male	3.1	7.4	12.7	18.7	31.7	48.5	67.4	70.4	81.0
Female	2.9	5.7	11.2	15.4	25.2	31.1	45.6	52.3	69.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Non-Hispanic 5/	10.9	18.5	30.8	49.3	72.6	101.4	136.7	177.9	241.0
Male	12.5	21.2	35.8	58.6	87.5	123.8	168.2	222.5	303.0
Female	9.3	16.0	26.1	40.9	59.7	83.6	114.9	151.7	215.9
Non-Hispanic white	10.2	17.5	30.1	48.3	72.4	100.8	136.2	179.9	244.9
Male	11.5	20.0	34.7	56.1	87.6	123.1	166.3	226.2	309.5
Female	8.8	14.9	25.7	41.0	59.1	82.9	115.1	152.4	218.9
Non-Hispanic black	17.1	29.5	42.1	65.9	86.8	118.5	156.7	174.4	211.1
Male	21.4	33.9	52.1	90.4	103.1	150.3	211.9	212.0	253.6
Female	13.5	25.7	33.8	46.6	74.7	97.3	123.4	155.5	195.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85 YEARS
	YEARS	AND OVER							

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Non-Hispanic 5/	1.5	2.3	3.9	6.8	10.8	17.6	29.2	51.2	85.2
Male	1.7	2.7	4.2	7.8	13.8	21.6	37.2	66.6	109.4
Female	1.3	1.9	3.6	5.8	8.2	14.6	23.7	42.1	75.4
Non-Hispanic white	1.4	2.3	4.0	6.8	11.0	17.6	29.8	52.8	87.6
Male	1.6	2.7	4.4	7.5	13.9	21.3	38.0	69.4	113.1
Female	1.2	1.8	3.6	6.1	8.5	14.6	24.2	43.0	77.3
Non-Hispanic black	2.4	3.2	4.2	7.3	10.9	20.3	27.0	38.2	64.2
Male	2.2	3.7	4.2	10.4	16.0	23.9	34.7	39.5	76.3
Female	2.5	2.8	4.2	4.9	7.1	17.9	22.4	37.6	59.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Anemias (D50-D64)

Non-Hispanic 5/	0.6	0.7	1.0	1.2	2.2	3.4	6.6	13.9	43.2
Male	0.6	0.7	1.1	1.3	2.2	3.5	6.6	14.4	41.0
Female	0.6	0.7	0.9	1.2	2.3	3.4	6.6	13.6	44.1
Non-Hispanic white	0.2	0.4	0.7	0.9	2.0	3.2	6.3	13.8	42.9
Male	*	0.4	0.8	0.9	2.1	3.3	6.3	14.1	39.9
Female	*	0.4	0.6	0.9	1.8	3.1	6.3	13.5	44.1
Non-Hispanic black	3.0	3.2	3.0	4.0	4.9	6.3	10.5	18.1	51.8
Male	3.3	3.5	*	*	*	*	11.1	18.9	56.4
Female	2.8	3.1	3.0	4.0	5.9	6.4	10.2	17.7	50.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diabetes mellitus (E10-E14)

Non-Hispanic 5/	9.5	17.0	28.3	46.4	71.1	103.6	151.7	209.0	313.4
Male	11.4	19.5	31.8	51.9	77.6	114.4	168.6	237.2	339.7
Female	7.7	14.6	25.1	41.4	65.5	95.1	139.9	192.4	302.8
Non-Hispanic white	7.8	14.0	23.7	39.7	61.2	91.7	137.5	193.0	295.5
Male	9.5	16.7	27.3	45.7	69.6	105.1	157.9	227.9	330.4
Female	6.1	11.3	20.4	34.2	53.7	80.9	123.2	172.2	281.4
Non-Hispanic black	22.4	41.7	67.7	103.7	160.7	226.7	320.5	409.2	530.1
Male	26.1	44.4	73.5	110.1	155.4	227.2	314.1	384.5	456.8
Female	19.3	39.4	63.0	98.6	164.6	226.4	324.4	421.7	557.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Nutritional deficiencies (E40-E64)

Non-Hispanic 5/	0.2	0.3	0.5	1.0	2.0	3.1	6.8	15.8	50.3
Male	0.3	*	0.7	1.1	2.0	3.4	6.7	14.3	43.9
Female	0.2	0.4	0.4	0.9	2.0	2.8	6.9	16.6	52.8
Non-Hispanic white	0.2	0.2	0.5	0.9	1.9	2.7	6.4	15.3	49.0
Male	*	*	0.5	0.9	1.8	2.8	6.1	13.8	43.0
Female	*	0.3	0.4	0.9	1.9	2.6	6.5	16.1	51.5
Non-Hispanic black	*	*	*	2.2	3.3	6.7	12.1	23.0	67.4
Male	*	*	*	*	*	9.9	14.9	23.2	64.6
Female	*	*	*	*	*	4.5	10.5	22.9	68.5

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Malnutrition (E40-E46)

Non-Hispanic 5/	0.2	0.3	0.5	0.9	1.9	2.8	6.4	14.6	46.5
Male	0.2	*	0.6	1.0	1.8	3.2	6.3	13.0	40.9
Female	0.2	0.3	0.3	0.9	1.9	2.5	6.6	15.5	48.7
Non-Hispanic white	0.2	0.2	0.4	0.8	1.7	2.4	6.1	14.1	45.3
Male	*	*	0.4	0.8	1.7	2.6	5.8	12.3	39.8
Female	*	*	*	0.9	1.8	2.2	6.3	15.1	47.4
Non-Hispanic black	*	*	*	2.0	3.2	6.7	11.2	21.3	64.2
Male	*	*	*	*	*	9.9	14.0	23.2	62.2
Female	*	*	*	*	*	4.5	9.6	20.3	65.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Parkinson's disease (G20-G21)

Non-Hispanic 5/	*	0.2	0.6	1.8	6.6	17.1	45.5	88.5	133.3
Male	*	*	0.9	2.3	9.1	24.7	68.5	136.7	225.2
Female	*	*	0.4	1.3	4.4	11.0	29.4	60.2	96.1
Non-Hispanic white	*	0.2	0.7	1.9	7.1	18.2	48.4	93.7	141.2
Male	*	*	0.9	2.5	9.8	26.2	72.0	143.4	238.6
Female	*	*	0.5	1.4	4.8	11.8	31.8	64.1	101.9
Non-Hispanic black	*	*	*	*	3.2	8.9	21.0	37.4	52.5
Male	*	*	*	*	*	14.0	37.6	65.2	88.1
Female	*	*	*	*	*	5.4	11.0	23.3	39.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Alzheimer's disease (G30)

Non-Hispanic 5/	*	0.3	1.1	3.3	9.2	29.5	86.5	224.5	676.7
Male	*	0.3	1.0	3.0	9.8	29.3	81.0	209.5	531.0
Female	*	0.3	1.2	3.6	8.7	29.6	90.3	233.3	735.6
Non-Hispanic white	*	0.3	1.2	3.5	9.7	30.5	89.5	232.7	705.1
Male	*	*	1.0	3.2	10.3	30.2	83.8	217.3	551.7
Female	*	*	1.4	3.9	9.2	30.8	93.4	241.9	767.0
Non-Hispanic black	*	*	*	2.2	7.6	26.5	72.3	173.9	450.9
Male	*	*	*	*	8.0	28.4	65.5	157.9	378.1
Female	*	*	*	*	7.3	25.2	76.4	181.9	478.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Major cardiovascular diseases (I00-I78)

All origins 3/	88.5	146.7	248.4	407.5	647.9	1,066.0	1,824.4	3,253.7	8,030.3
Male	126.8	209.0	350.6	556.9	857.2	1,403.8	2,305.7	3,956.7	8,448.1
Female	51.4	87.0	152.8	272.0	468.5	799.9	1,489.2	2,839.0	7,860.1
Hispanic	56.1	99.6	178.2	310.5	501.0	861.4	1,451.4	2,583.3	6,168.2
Male	80.2	144.5	257.4	426.9	658.5	1,145.9	1,849.2	3,116.9	6,304.3
Female	32.0	56.8	106.6	210.2	373.5	645.9	1,169.6	2,258.6	6,101.4
Mexican	55.5	91.8	173.7	343.4	503.0	883.7	1,551.3	*	*
Male	75.9	126.1	240.8	440.7	633.4	1,155.4	*	*	*
Female	33.9	54.8	109.7	249.7	393.2	670.6	*	*	*
Puerto Rican	83.3	139.0	299.6	380.8	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Cuban	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	44.6	92.7	133.5	233.6	461.3	781.2	*	*	*
Male	66.2	140.1	215.2	*	*	*	*	*	*
Female	26.0	53.6	72.0	135.4	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,

1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Major cardiovascular diseases (I00-I78)

Non-Hispanic 5/	91.1	149.9	252.6	412.4	655.5	1,073.4	1,835.5	3,270.0	8,076.4
Male	130.7	213.2	355.8	563.0	866.4	1,411.9	2,318.5	3,979.0	8,513.1
Female	53.0	89.2	155.8	275.5	473.9	806.2	1,499.3	2,852.2	7,899.8
Non-Hispanic white	78.1	132.4	228.1	378.9	617.5	1,032.1	1,791.9	3,258.1	8,179.0
Male	117.3	193.1	326.7	527.5	829.2	1,372.2	2,284.2	3,984.9	8,650.5
Female	39.4	73.0	133.8	241.2	431.4	758.3	1,445.9	2,826.1	7,988.9
Non-Hispanic black	195.9	310.0	497.1	748.1	1,078.7	1,639.8	2,519.2	3,750.0	7,576.5
Male	252.1	415.3	680.4	967.2	1,346.6	2,057.4	3,018.8	4,434.6	7,755.5
Female	147.3	220.3	347.1	575.2	881.0	1,362.6	2,218.0	3,405.4	7,510.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN,

RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

All origins 3/	71.2	120.4	205.5	330.6	515.8	826.7	1,372.3	2,392.1	5,926.1
Male	107.1	178.1	298.9	463.9	702.0	1,119.7	1,787.5	3,012.3	6,430.0
Female	36.4	65.0	118.1	209.8	356.3	596.0	1,083.0	2,026.2	5,720.9
Hispanic	39.9	76.8	141.1	237.3	388.4	654.8	1,096.6	1,940.5	4,701.8
Male	60.8	115.6	208.7	338.5	525.4	892.6	1,411.5	2,374.9	4,897.5
Female	19.1	39.8	80.0	150.1	277.5	474.6	873.7	1,676.2	4,605.8
Mexican	38.9	70.9	134.5	260.7	384.3	650.8	1,158.8	*	*
Male	56.9	102.4	193.1	344.5	495.4	877.7	*	*	*
Female	19.9	37.0	78.5	180.0	290.7	472.8	*	*	*
Puerto Rican	65.0	104.8	249.7	296.6	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Cuban	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	30.6	69.9	102.4	172.0	347.1	571.8	*	*	*
Male	49.8	111.4	167.9	*	*	*	*	*	*
Female	*	35.6	53.1	89.7	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Diseases of heart (I00-I09,I11,I13,I20-I51)									
Non-Hispanic 5/	73.9	123.4	209.4	335.7	522.3	833.0	1,380.1	2,402.3	5,953.6
Male	111.1	182.4	304.2	470.1	710.1	1,127.0	1,798.4	3,028.8	6,474.6
Female	37.9	66.9	120.5	213.4	360.8	600.9	1,089.1	2,033.1	5,742.9
Non-Hispanic white	65.8	112.1	192.8	313.1	497.3	805.7	1,353.2	2,402.0	6,047.9
Male	102.8	169.3	284.7	446.8	686.5	1,102.9	1,780.9	3,046.4	6,606.6
Female	29.1	56.0	105.0	189.2	330.9	566.6	1,052.6	2,019.1	5,822.7
Non-Hispanic black	145.2	236.1	386.8	579.6	826.1	1,242.2	1,852.8	2,693.6	5,424.6
Male	195.0	325.5	540.6	763.9	1,047.1	1,582.4	2,260.1	3,247.6	5,677.2
Female	102.0	159.9	260.9	434.1	663.2	1,016.4	1,607.3	2,414.7	5,330.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Non-Hispanic 5/	0.6	0.8	1.4	2.2	3.4	5.5	8.3	11.8	17.1
Male	0.5	0.6	1.1	1.8	2.3	3.5	5.9	8.1	12.5
Female	0.7	0.9	1.6	2.5	4.5	7.1	10.0	13.9	19.0
Non-Hispanic white	0.5	0.7	1.3	2.2	3.6	5.6	8.9	12.5	18.0
Male	0.5	0.6	1.1	1.9	2.3	3.5	6.3	8.7	13.0
Female	0.6	0.8	1.4	2.4	4.7	7.3	10.7	14.8	20.0
Non-Hispanic black	1.0	1.1	2.0	2.1	3.5	3.7	3.8	*	8.6
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	4.5	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Hypertensive heart disease (I11)									
Non-Hispanic 5/	4.8	6.7	9.5	13.3	16.5	22.7	36.1	66.0	194.3
Male	6.4	9.6	12.3	17.6	20.6	26.7	36.6	61.5	139.6
Female	3.3	3.8	6.8	9.3	13.0	19.5	35.8	68.6	216.5
Non-Hispanic white	3.1	4.3	6.2	9.1	12.2	17.0	29.7	58.0	182.5
Male	4.5	6.5	8.5	12.4	15.6	20.7	30.2	52.9	129.5
Female	1.7	2.1	4.0	6.0	9.2	14.0	29.4	61.0	203.9
Non-Hispanic black	18.1	25.9	38.4	48.9	55.6	82.1	111.8	165.8	344.1
Male	22.0	36.5	48.8	65.1	71.2	95.7	117.1	179.4	275.9
Female	14.8	16.9	29.9	36.1	44.0	73.1	108.6	159.0	369.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Hypertensive heart and renal disease (I13)

Non-Hispanic 5/	0.3	0.4	0.8	1.0	1.6	3.0	5.0	9.8	24.2
Male	0.4	0.5	0.9	1.0	1.7	3.1	4.8	10.7	27.9
Female	0.2	0.4	0.7	0.9	1.5	2.8	5.2	9.3	22.8
Non-Hispanic white	*	0.2	0.4	0.4	0.8	2.1	3.9	8.0	21.8
Male	*	*	0.4	*	1.0	2.2	4.2	9.0	24.3
Female	*	*	*	*	0.7	2.0	3.7	7.4	20.8
Non-Hispanic black	2.0	2.5	4.1	5.4	8.4	11.9	17.0	31.3	50.9
Male	2.8	3.0	5.1	5.7	9.3	13.3	12.5	33.5	64.6
Female	*	2.1	3.3	5.2	7.7	10.9	19.7	30.2	45.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Ischemic heart diseases (I20-I25)									
All origins 3/	48.0	86.7	153.9	250.0	390.8	628.0	1,032.5	1,761.7	4,182.5
Male	76.9	134.2	232.8	362.9	550.5	879.6	1,394.4	2,305.0	4,701.7
Female	20.0	41.3	80.0	147.7	253.9	429.9	780.5	1,441.2	3,971.0
Hispanic	27.4	55.4	109.9	186.8	312.5	521.9	873.5	1,544.0	3,697.2
Male	44.8	86.7	167.7	276.6	435.1	726.3	1,161.2	1,936.1	3,865.6
Female	10.0	25.5	57.6	109.4	213.2	367.1	669.8	1,305.3	3,614.6
Mexican	26.4	49.9	104.6	208.1	307.5	519.3	922.8	*	*
Male	42.2	74.1	154.9	285.9	406.6	723.3	*	*	*
Female	9.6	23.8	56.6	133.1	224.0	359.3	*	*	*
Puerto Rican	47.7	78.8	192.3	232.0	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Cuban	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	19.7	51.3	80.1	133.8	285.0	454.0	*	*	*
Male	33.9	88.8	136.3	*	*	*	*	*	*
Female	*	20.3	37.7	64.4	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Ischemic heart diseases (I20-I25)									
Non-Hispanic 5/	49.7	88.9	156.4	253.1	394.4	631.3	1,036.0	1,764.3	4,187.0
Male	79.6	137.3	236.5	366.8	555.3	883.8	1,399.7	2,312.5	4,721.1
Female	20.8	42.4	81.4	149.8	255.9	431.9	782.9	1,441.2	3,971.0
Non-Hispanic white	46.9	84.3	148.3	240.8	380.8	617.2	1,021.9	1,770.0	4,261.3
Male	77.2	132.8	226.8	355.2	545.0	873.4	1,394.6	2,335.6	4,833.4
Female	17.0	36.7	73.2	134.8	236.5	410.9	759.9	1,433.9	4,030.7
Non-Hispanic black	79.8	142.9	254.5	398.2	573.6	874.6	1,307.5	1,894.4	3,701.7
Male	113.9	201.0	371.1	536.0	735.3	1,148.8	1,639.7	2,329.3	3,911.2
Female	50.2	93.4	159.1	289.5	454.4	692.6	1,107.3	1,675.5	3,623.9

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
TENTH REVISION, INTERNATIONAL
CLASSIFICATION OF DISEASES,
1992), HISPANIC ORIGIN,
RACE FOR NON-HISPANIC
POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Acute myocardial infarction (I21-I22)									
Non-Hispanic 5/	22.0	39.1	68.7	113.4	169.5	257.1	407.8	652.8	1,340.8
Male	35.1	59.5	102.2	162.2	235.0	353.2	538.5	840.4	1,576.2
Female	9.3	19.4	37.4	69.0	113.2	181.3	316.8	542.3	1,245.6
Non-Hispanic white	21.6	37.9	67.3	111.3	166.6	253.9	403.8	654.9	1,355.3
Male	35.3	59.3	101.2	161.8	234.8	352.6	539.5	847.9	1,600.7
Female	8.0	17.0	34.9	64.5	106.7	174.5	308.4	540.2	1,256.3
Non-Hispanic black	30.2	56.7	96.2	154.4	222.8	333.5	495.3	705.5	1,281.8
Male	41.9	74.6	133.6	200.3	275.0	426.1	596.8	865.1	1,426.6
Female	20.1	41.4	65.7	118.3	184.2	272.0	434.1	625.2	1,228.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other acute ischemic heart diseases (I24)

Non-Hispanic 5/	0.5	1.1	1.7	2.6	3.2	4.0	6.1	9.6	22.2
Male	0.8	1.7	2.6	3.7	4.6	5.3	7.8	11.9	21.0
Female	*	0.6	0.8	1.6	2.1	3.0	4.9	8.2	22.7
Non-Hispanic white	0.4	1.0	1.4	2.4	3.0	3.8	5.7	9.5	22.5
Male	0.8	1.6	2.3	3.4	4.2	5.0	7.4	11.8	21.3
Female	*	0.5	0.6	1.4	2.0	2.9	4.6	8.2	23.0
Non-Hispanic black	1.0	2.1	3.9	4.9	5.2	6.1	10.1	11.8	21.9
Male	*	3.5	5.9	6.8	8.3	9.2	13.5	*	*
Female	*	*	*	3.4	*	*	8.1	9.9	22.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other forms of chronic ischemic heart disease (I20,I25)

Non-Hispanic 5/	27.2	48.7	86.0	137.1	221.6	370.1	622.1	1,101.9	2,824.0
Male	43.7	76.1	131.7	200.9	315.7	525.3	853.4	1,460.2	3,123.9
Female	11.3	22.4	43.2	79.1	140.7	247.6	461.2	890.7	2,702.7
Non-Hispanic white	24.9	45.3	79.5	127.1	211.2	359.4	612.4	1,105.6	2,883.6
Male	41.1	72.0	123.2	189.9	306.1	515.8	847.8	1,475.9	3,211.3
Female	8.9	19.2	37.8	68.9	127.8	233.5	447.0	885.5	2,751.4
Non-Hispanic black	48.5	84.1	154.4	238.9	345.6	535.0	802.0	1,177.1	2,398.0
Male	70.5	122.9	231.6	328.9	452.0	713.5	1,029.4	1,448.7	2,463.5
Female	29.5	51.1	91.2	167.9	267.2	416.5	665.0	1,040.4	2,373.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Atherosclerotic cardiovascular disease, so described (I25.0)

Non-Hispanic 5/	9.6	17.7	28.6	40.3	55.3	81.3	125.0	207.9	535.9
Male	15.8	28.4	45.7	60.2	80.4	114.3	165.7	252.5	551.0
Female	3.7	7.4	12.5	22.3	33.8	55.2	96.7	181.6	529.8
Non-Hispanic white	8.5	15.7	24.5	34.3	48.6	72.3	115.5	198.2	530.2
Male	14.2	25.6	39.5	52.4	71.8	102.1	152.6	240.8	544.7
Female	2.8	5.9	10.2	17.4	28.2	48.3	89.5	172.9	524.4
Non-Hispanic black	19.7	36.9	66.3	94.4	120.8	178.9	240.8	342.8	652.2
Male	30.5	57.2	107.3	137.8	170.0	259.7	339.1	429.1	711.6
Female	10.4	19.5	32.7	60.2	84.5	125.2	181.5	299.4	630.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Non-Hispanic 5/	17.6	31.0	57.4	96.8	166.3	288.8	497.1	894.0	2,288.1
Male	27.9	47.7	86.0	140.8	235.3	411.0	687.7	1,207.7	2,572.9
Female	7.6	15.0	30.7	56.9	106.9	192.4	364.5	709.1	2,172.9
Non-Hispanic white	16.4	29.6	55.0	92.8	162.6	287.1	496.9	907.4	2,353.3
Male	26.9	46.3	83.7	137.5	234.2	413.6	695.2	1,235.1	2,666.6
Female	6.1	13.2	27.6	51.5	99.6	185.3	357.5	712.6	2,227.0
Non-Hispanic black	28.8	47.3	88.1	144.5	224.8	356.1	561.3	834.3	1,745.8
Male	40.0	65.6	124.3	191.1	282.0	453.8	690.3	1,019.6	1,751.9
Female	19.2	31.6	58.5	107.7	182.6	291.3	483.5	741.0	1,743.5

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Other heart diseases (I26-I51)									
Non-Hispanic 5/	18.4	26.7	41.3	66.1	106.4	170.6	294.7	550.4	1,530.9
Male	24.1	34.3	53.3	83.0	130.2	209.8	351.4	636.0	1,573.6
Female	12.8	19.3	30.1	50.8	86.0	139.6	255.3	500.0	1,513.7
Non-Hispanic white	15.2	22.6	36.7	60.6	99.8	163.9	288.8	553.5	1,564.2
Male	20.5	29.2	47.9	76.8	122.7	203.1	345.6	640.2	1,606.4
Female	9.9	16.2	26.1	45.6	79.8	132.3	248.8	502.0	1,547.2
Non-Hispanic black	44.3	63.6	87.8	125.0	185.1	270.0	412.7	597.8	1,319.4
Male	55.6	84.4	114.4	156.1	229.2	321.9	487.5	704.6	1,416.1
Female	34.5	45.9	66.0	100.5	152.6	235.5	367.6	544.0	1,283.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Heart failure (I50)									
Non-Hispanic 5/	1.5	3.2	6.3	13.1	24.7	47.5	95.2	202.8	694.8
Male	2.1	3.8	7.5	15.5	28.9	56.8	110.3	233.4	683.3
Female	0.9	2.6	5.2	10.9	21.1	40.1	84.7	184.8	699.5
Non-Hispanic white	1.1	2.7	5.5	12.0	23.3	45.8	94.5	204.9	714.3
Male	1.7	3.2	6.9	14.2	27.8	55.5	110.1	235.8	701.6
Female	0.6	2.1	4.2	9.9	19.3	38.1	83.6	186.6	719.4
Non-Hispanic black	4.3	7.9	14.4	24.8	42.0	73.4	121.4	211.8	564.8
Male	5.7	9.3	15.2	30.0	46.4	84.1	137.3	246.3	589.4
Female	3.1	6.6	13.7	20.6	38.7	66.2	111.8	194.4	555.6

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Non-Hispanic 5/	16.2	22.7	34.0	52.0	80.1	120.9	196.2	343.5	831.4
Male	21.1	29.7	44.5	66.2	99.4	150.3	236.4	396.9	883.9
Female	11.4	16.1	24.2	39.0	63.5	97.7	168.2	312.1	810.2
Non-Hispanic white	13.5	19.3	30.4	47.6	75.0	115.9	191.0	344.5	845.3
Male	18.2	25.2	39.8	61.4	93.3	144.9	231.0	398.7	898.3
Female	8.9	13.6	21.4	34.9	58.9	92.6	162.9	312.3	824.0
Non-Hispanic black	38.1	53.9	70.9	98.3	140.3	193.5	287.0	382.5	748.9
Male	47.4	72.9	96.8	124.0	177.8	235.1	344.9	454.0	821.9
Female	30.0	37.6	49.6	78.0	112.7	165.8	252.1	346.5	721.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

Non-Hispanic 5/	1.8	3.0	4.5	7.6	11.5	18.9	33.5	62.9	162.9
Male	2.3	3.8	5.4	8.7	11.5	20.4	33.9	62.4	142.0
Female	1.3	2.2	3.5	6.6	11.5	17.8	33.3	63.2	171.3
Non-Hispanic white	1.0	1.7	2.9	5.0	8.4	14.9	28.3	57.0	152.7
Male	1.3	2.3	3.7	5.9	8.7	16.3	29.1	56.5	131.8
Female	0.6	1.1	2.2	4.2	8.0	13.8	27.8	57.2	161.1
Non-Hispanic black	7.3	13.1	18.3	29.5	41.0	59.9	93.8	132.8	285.9
Male	9.0	15.9	22.9	35.3	40.8	65.3	93.9	139.9	268.9
Female	5.9	10.6	14.5	25.0	41.1	56.3	93.8	129.2	292.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Cerebrovascular diseases (I60-I69)									
Non-Hispanic 5/	12.9	19.4	30.9	53.5	91.3	169.6	335.1	658.3	1,604.5
Male	13.9	21.3	34.9	62.0	102.6	192.7	369.1	698.8	1,499.9
Female	12.0	17.6	27.1	45.7	81.7	151.4	311.4	634.5	1,646.8
Non-Hispanic white	9.2	15.1	25.0	45.6	81.4	158.8	323.1	651.1	1,616.8
Male	10.0	16.4	27.5	52.8	90.8	179.5	355.4	689.2	1,505.9
Female	8.3	13.8	22.5	38.8	73.2	142.2	300.4	628.4	1,661.6
Non-Hispanic black	38.0	52.4	78.6	115.6	178.0	284.5	479.0	775.9	1,545.8
Male	41.5	61.7	99.4	138.8	216.4	343.4	548.2	872.8	1,479.5
Female	34.9	44.5	61.7	97.3	149.6	245.5	437.3	727.1	1,570.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Atherosclerosis (I70)									
Non-Hispanic 5/	0.4	0.6	1.3	2.7	5.2	10.6	23.0	52.8	190.3
Male	0.5	0.8	1.8	3.7	6.7	13.1	27.5	60.2	178.7
Female	0.2	0.4	0.9	1.7	4.0	8.6	19.8	48.5	195.1
Non-Hispanic white	0.3	0.5	1.3	2.5	5.3	10.5	23.3	53.4	196.8
Male	0.4	0.7	1.8	3.6	6.6	13.3	28.3	60.8	183.6
Female	*	0.3	0.8	1.5	4.1	8.3	19.8	49.1	202.1
Non-Hispanic black	*	1.5	2.1	4.8	5.8	13.1	23.6	54.6	140.2
Male	*	*	*	6.2	8.8	14.0	25.0	67.8	150.3
Female	*	*	*	3.7	*	12.5	22.7	48.0	136.5

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of circulatory system (I71-I78)

Non-Hispanic 5/	2.2	3.5	6.5	13.0	25.0	41.3	63.8	93.7	165.0
Male	2.9	5.0	9.5	18.5	35.6	58.7	89.6	128.8	217.9
Female	1.5	2.1	3.7	8.1	16.0	27.5	45.8	73.0	143.6
Non-Hispanic white	1.9	3.0	6.1	12.7	25.2	42.1	64.0	94.6	164.8
Male	2.7	4.4	9.0	18.4	36.6	60.3	90.5	132.0	222.7
Female	1.1	1.7	3.3	7.5	15.2	27.5	45.3	72.3	141.5
Non-Hispanic black	4.6	6.9	11.3	18.6	27.8	40.1	69.9	93.1	180.0
Male	5.6	10.2	15.2	23.0	33.6	52.3	91.5	106.4	179.7
Female	3.8	4.2	8.0	15.1	23.6	32.0	56.9	86.4	180.1

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Aortic aneurysm and dissection (I71)

Non-Hispanic 5/	1.4	2.2	4.5	8.7	16.8	28.4	41.6	57.3	80.3
Male	2.0	3.5	6.9	13.5	25.7	42.9	62.6	86.5	129.6
Female	0.8	1.0	2.2	4.3	9.2	16.9	27.0	40.2	60.4
Non-Hispanic white	1.2	2.0	4.2	8.8	17.3	29.7	42.6	59.1	82.0
Male	1.9	3.2	6.7	13.8	27.0	45.0	64.3	90.3	135.3
Female	0.5	0.8	1.9	4.1	8.8	17.4	27.4	40.5	60.5
Non-Hispanic black	2.9	3.7	7.2	9.5	14.6	20.5	34.1	41.1	63.0
Male	3.6	6.1	10.3	13.2	18.1	28.4	47.7	46.3	70.5
Female	2.4	*	4.7	6.5	12.0	15.2	25.8	38.5	60.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Non-Hispanic 5/	0.8	1.3	2.1	4.4	8.2	12.9	22.2	36.3	84.7
Male	0.9	1.5	2.6	4.9	9.9	15.8	27.0	42.3	88.2
Female	0.7	1.1	1.6	3.8	6.8	10.6	18.8	32.8	83.3
Non-Hispanic white	0.7	1.0	1.9	3.9	7.8	12.4	21.3	35.5	82.8
Male	0.8	1.2	2.4	4.5	9.5	15.3	26.2	41.7	87.4
Female	0.6	0.9	1.3	3.4	6.4	10.1	17.9	31.8	81.0
Non-Hispanic black	1.7	3.2	4.0	9.1	13.2	19.6	35.9	52.0	117.0
Male	2.0	4.1	4.9	9.8	15.5	23.9	43.8	60.1	109.2
Female	*	2.4	3.3	8.6	11.6	16.8	31.1	48.0	119.9

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other disorders of circulatory system (I80-I99)

Non-Hispanic 5/	1.3	1.5	1.8	2.7	3.4	5.4	7.7	12.0	25.6
Male	1.5	1.7	2.0	2.7	3.9	5.7	8.0	12.3	22.1
Female	1.1	1.3	1.6	2.7	3.0	5.1	7.5	11.8	27.0
Non-Hispanic white	1.2	1.4	1.6	2.4	2.9	5.1	7.3	11.9	25.5
Male	1.4	1.6	1.7	2.4	3.4	5.6	7.8	12.1	21.8
Female	1.0	1.1	1.4	2.4	2.5	4.7	7.0	11.8	27.0
Non-Hispanic black	2.5	3.2	4.0	5.3	8.7	9.3	14.5	14.4	29.9
Male	3.2	3.8	4.9	5.7	9.3	8.9	13.0	*	30.5
Female	2.0	2.7	3.3	4.9	8.3	9.5	15.4	14.7	29.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN,

RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Influenza and pneumonia (J10-J18)

Non-Hispanic 5/	3.9	5.6	8.8	15.6	26.6	52.2	106.7	241.0	745.7
Male	4.7	7.0	10.8	18.7	32.9	65.2	134.0	307.9	894.9
Female	3.3	4.4	7.0	12.7	21.2	41.9	87.7	201.6	685.4
Non-Hispanic white	3.3	4.9	7.9	14.2	24.4	49.7	105.5	240.7	759.8
Male	3.7	5.7	9.2	16.9	29.5	61.4	131.6	305.1	908.1
Female	2.9	4.0	6.7	11.7	20.0	40.2	87.2	202.5	700.0
Non-Hispanic black	8.6	12.4	18.1	27.6	46.2	78.5	124.8	247.4	604.5
Male	11.7	17.6	26.2	35.5	65.0	108.7	173.4	345.9	763.2
Female	6.0	8.1	11.4	21.3	32.2	58.5	95.5	197.9	545.6

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Pneumonia (J12-J18)									
Non-Hispanic 5/	3.8	5.5	8.5	15.1	25.8	50.7	104.1	234.8	725.2
Male	4.5	6.8	10.5	18.2	31.9	63.5	131.3	301.6	873.8
Female	3.1	4.3	6.8	12.2	20.5	40.6	85.2	195.4	665.2
Non-Hispanic white	3.2	4.7	7.6	13.7	23.6	48.2	102.7	234.2	738.1
Male	3.7	5.6	8.9	16.4	28.5	59.6	128.7	298.4	886.1
Female	2.8	3.9	6.3	11.1	19.3	39.0	84.5	196.0	678.4
Non-Hispanic black	8.4	12.3	17.9	27.2	45.6	77.0	123.9	245.1	595.0
Male	11.4	17.4	25.9	35.3	64.5	107.6	171.5	345.0	750.3
Female	5.8	8.1	11.4	20.8	31.7	56.7	95.2	194.9	537.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Chronic lower respiratory diseases (J40-J47)

Non-Hispanic 5/	6.1	12.3	31.0	65.3	129.3	226.0	335.4	480.7	653.2
Male	6.5	12.6	33.4	70.3	144.8	265.6	412.8	637.1	978.0
Female	5.7	12.1	28.8	60.8	115.9	194.7	281.5	388.6	521.9
Non-Hispanic white	5.5	12.0	31.7	68.8	136.5	238.6	351.8	503.7	680.8
Male	5.9	12.1	33.6	72.5	148.9	274.6	425.1	658.0	1,008.6
Female	5.1	11.9	29.9	65.3	125.7	209.7	300.2	412.0	548.7
Non-Hispanic black	11.3	18.5	33.8	58.1	98.7	152.8	213.1	273.6	376.2
Male	11.8	19.6	40.3	74.2	140.2	228.6	328.0	451.4	696.3
Female	10.8	17.4	28.4	45.5	68.0	102.5	143.7	184.1	257.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Emphysema (J43)									
Non-Hispanic 5/	0.8	1.9	5.4	11.0	21.8	36.9	48.7	61.5	64.3
Male	1.1	2.2	6.0	11.7	25.6	44.0	61.9	85.1	105.9
Female	0.6	1.6	4.8	10.3	18.5	31.3	39.5	47.7	47.5
Non-Hispanic white	0.8	1.9	5.6	11.9	23.2	39.5	51.6	64.9	66.9
Male	0.9	2.1	6.2	12.3	26.3	46.3	64.8	88.6	107.3
Female	0.6	1.6	5.1	11.5	20.5	34.1	42.3	50.8	50.6
Non-Hispanic black	1.6	2.6	4.8	7.8	15.3	19.2	24.5	31.0	37.5
Male	2.5	3.6	6.1	10.8	25.9	29.4	34.7	53.2	89.2
Female	*	*	3.7	5.4	7.5	12.5	18.3	19.9	18.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Asthma (J45-J46)									
Non-Hispanic 5/	1.6	1.7	2.0	2.3	3.3	4.5	5.5	8.7	16.6
Male	1.3	1.2	1.6	1.6	2.6	3.2	3.8	6.7	12.7
Female	1.9	2.1	2.4	3.0	3.9	5.5	6.7	9.8	18.2
Non-Hispanic white	1.0	1.2	1.4	1.8	2.5	3.9	5.0	8.4	15.8
Male	0.9	0.9	1.0	1.1	1.8	2.9	3.5	6.3	11.1
Female	1.2	1.4	1.7	2.3	3.2	4.8	6.1	9.7	17.7
Non-Hispanic black	5.6	5.3	6.7	7.2	9.1	9.8	10.9	9.8	22.3
Male	4.1	3.2	6.3	6.0	8.8	6.8	*	*	24.7
Female	6.9	7.1	7.1	8.2	9.2	11.8	14.5	9.9	21.4

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other chronic lower respiratory diseases (J44,J47)

Non-Hispanic 5/	3.5	8.7	23.5	51.5	103.5	183.6	279.1	407.0	560.5
Male	4.1	9.2	25.7	56.4	115.7	217.5	344.3	540.5	846.6
Female	3.0	8.2	21.5	47.0	93.0	156.8	233.7	328.4	444.8
Non-Hispanic white	3.6	8.9	24.5	54.6	110.1	194.1	292.9	426.8	585.5
Male	4.0	9.0	26.3	58.6	119.8	224.5	353.8	558.0	876.5
Female	3.2	8.7	22.9	51.0	101.5	169.7	250.1	348.8	468.2
Non-Hispanic black	4.0	10.3	22.2	42.9	73.5	122.6	175.9	229.9	312.6
Male	5.2	12.7	27.9	56.8	105.0	191.0	286.6	386.2	578.9
Female	2.9	8.3	17.5	31.9	50.3	77.1	109.2	151.2	213.7

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Pneumonitis due to solids and liquids (J69)

Non-Hispanic 5/	0.8	1.2	1.9	3.2	7.0	14.5	30.8	66.4	189.0
Male	1.0	1.5	2.3	4.0	9.6	20.6	42.6	96.0	279.7
Female	0.6	1.0	1.5	2.5	4.7	9.6	22.6	48.9	152.3
Non-Hispanic white	0.7	1.1	1.8	3.0	6.6	13.6	30.5	66.3	192.5
Male	0.9	1.3	2.3	3.6	8.8	19.0	41.7	96.9	288.1
Female	0.5	0.9	1.4	2.3	4.6	9.3	22.6	48.2	153.9
Non-Hispanic black	1.7	2.3	3.0	5.9	11.5	25.5	39.1	72.4	160.6
Male	2.4	3.2	3.4	7.4	17.9	42.0	62.1	97.8	210.2
Female	*	*	*	4.7	6.9	14.5	25.3	59.6	142.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Non-Hispanic 5/	2.7	4.7	7.6	12.7	23.0	37.0	60.7	89.1	141.6
Male	2.9	5.1	8.7	14.6	27.0	46.0	77.5	112.3	181.3
Female	2.4	4.3	6.6	11.1	19.6	30.0	49.0	75.5	125.5
Non-Hispanic white	2.2	4.4	7.4	12.5	22.9	37.9	62.1	91.0	145.7
Male	2.4	4.7	8.4	14.2	27.0	46.6	79.0	114.9	185.9
Female	2.1	4.0	6.4	10.8	19.3	30.8	50.3	76.9	129.5
Non-Hispanic black	6.1	7.9	11.8	15.9	28.2	33.5	50.2	71.8	99.8
Male	7.4	8.7	13.7	18.1	32.2	41.3	68.9	83.3	135.0
Female	5.1	7.2	10.3	14.3	25.2	28.4	38.9	66.1	86.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Peptic ulcer (K25-K28)

Non-Hispanic 5/	0.7	1.0	1.5	2.2	3.2	5.1	8.8	15.5	35.7
Male	0.9	1.6	2.1	2.8	3.9	6.9	11.3	18.1	41.0
Female	0.5	0.5	1.0	1.7	2.5	3.6	7.0	14.0	33.5
Non-Hispanic white	0.6	0.9	1.4	2.2	3.1	5.1	8.7	15.9	36.3
Male	0.8	1.3	2.0	2.8	3.8	6.7	11.0	18.1	41.0
Female	0.4	0.5	0.9	1.7	2.4	3.7	7.0	14.5	34.4
Non-Hispanic black	1.4	2.1	2.6	2.7	4.2	5.7	9.4	11.2	26.7
Male	*	3.4	3.6	*	*	10.3	12.5	*	31.7
Female	*	*	*	*	*	*	7.6	8.6	24.9

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Non-Hispanic 5/	15.9	17.5	20.4	24.7	26.9	29.0	30.3	28.3	22.1
Male	23.7	26.4	29.1	35.3	36.0	38.5	38.4	33.4	33.0
Female	8.4	9.0	12.3	15.1	19.1	21.5	24.7	25.3	17.7
Non-Hispanic white	15.7	16.7	19.9	24.7	27.2	29.7	31.6	29.1	22.8
Male	23.8	25.1	28.5	35.4	36.4	39.2	39.7	34.7	34.1
Female	7.7	8.4	11.8	14.8	19.0	22.1	26.0	25.9	18.2
Non-Hispanic black	19.8	26.3	26.0	26.3	27.8	25.0	18.3	17.8	14.0
Male	27.2	41.2	38.9	37.8	38.4	37.9	27.9	19.7	*
Female	13.3	13.6	15.4	17.3	20.1	16.3	12.5	16.9	11.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Alcoholic liver disease (K70)									
Non-Hispanic 5/	9.4	10.0	10.8	12.0	10.4	8.5	6.6	4.2	2.2
Male	14.0	15.5	16.1	18.4	16.4	14.3	12.4	7.3	5.0
Female	4.8	4.8	5.8	6.1	5.2	3.9	2.6	2.4	1.0
Non-Hispanic white	9.2	9.7	10.4	11.9	10.4	8.6	6.7	4.1	2.2
Male	14.1	15.0	15.8	18.4	16.5	14.3	12.6	7.3	5.1
Female	4.4	4.4	5.3	5.8	5.1	4.0	2.5	2.3	1.0
Non-Hispanic black	11.3	13.6	14.0	12.8	12.0	8.6	6.3	*	*
Male	15.5	21.0	20.5	18.5	19.5	17.1	11.6	*	*
Female	7.7	7.3	8.7	8.4	6.5	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other chronic liver disease and cirrhosis (K73-K74)

Non-Hispanic 5/	6.6	7.5	9.6	12.7	16.5	20.5	23.7	24.1	19.9
Male	9.7	11.0	13.0	16.9	19.6	24.1	26.0	26.1	28.0
Female	3.5	4.3	6.5	9.0	14.0	17.6	22.1	22.9	16.7
Non-Hispanic white	6.5	7.0	9.5	12.8	16.7	21.1	25.0	25.0	20.6
Male	9.7	10.1	12.7	16.9	20.0	24.8	27.1	27.4	29.0
Female	3.3	4.0	6.5	9.0	13.8	18.1	23.5	23.6	17.2
Non-Hispanic black	8.5	12.7	12.0	13.5	15.8	16.4	12.0	13.5	12.4
Male	11.8	20.2	18.5	19.3	18.9	20.8	16.4	*	*
Female	5.6	6.2	6.7	8.9	13.6	13.4	9.3	14.3	11.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

Non-Hispanic 5/	0.2	0.3	0.6	1.0	1.8	2.9	5.5	10.1	24.6
Male	*	0.3	0.7	1.1	2.1	3.5	7.0	12.6	27.7
Female	*	0.4	0.5	0.9	1.5	2.5	4.5	8.6	23.4
Non-Hispanic white	0.1	0.3	0.5	0.9	1.7	2.6	5.5	9.9	25.3
Male	*	*	0.7	1.0	2.1	3.2	7.1	12.8	28.9
Female	*	0.3	0.4	0.8	1.3	2.2	4.3	8.2	23.9
Non-Hispanic black	*	*	*	*	2.8	4.9	6.5	11.2	18.1
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	4.8	6.4	11.7	19.6

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Non-Hispanic 5/	3.5	5.5	9.4	16.6	29.2	47.2	79.6	133.4	279.0
Male	4.5	6.4	10.8	18.3	32.7	54.3	100.3	174.1	384.0
Female	2.6	4.7	8.2	15.1	26.2	41.6	65.2	109.4	236.6
Non-Hispanic white	2.2	3.6	6.6	12.5	23.0	40.2	72.7	125.6	269.4
Male	2.8	4.2	7.7	14.3	27.3	48.4	93.5	167.6	378.9
Female	1.7	3.1	5.6	10.9	19.3	33.6	58.2	100.6	225.2
Non-Hispanic black	13.1	20.1	33.0	51.3	84.6	121.2	167.2	245.1	410.2
Male	18.0	24.3	38.3	54.9	87.9	128.5	201.4	292.7	496.7
Female	8.9	16.4	28.7	48.5	82.2	116.4	146.6	221.2	378.1

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Renal failure (N17-N19)

Non-Hispanic 5/	3.4	5.3	9.2	16.3	28.6	46.2	78.3	130.6	274.9
Male	4.3	6.1	10.5	17.9	31.9	53.2	98.4	171.3	377.7
Female	2.5	4.6	8.0	14.8	25.8	40.7	64.3	106.7	233.4
Non-Hispanic white	2.1	3.6	6.4	12.2	22.6	39.2	71.5	122.8	265.7
Male	2.6	4.1	7.4	14.0	26.6	47.3	91.7	164.6	373.4
Female	1.6	3.0	5.4	10.6	19.0	32.7	57.3	98.0	222.2
Non-Hispanic black	12.7	19.2	32.6	50.6	83.2	120.1	164.9	241.7	404.2
Male	17.6	22.9	37.9	53.8	85.8	127.8	198.9	290.1	487.3
Female	8.5	16.0	28.3	48.1	81.2	115.0	144.3	217.3	373.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Non-Hispanic 5/	1.5	1.9	2.2	2.5	2.6	2.5	3.2	4.9	7.5
Male	1.6	2.2	2.0	2.5	2.8	2.6	3.1	5.2	10.4
Female	1.3	1.6	2.4	2.5	2.5	2.4	3.3	4.8	6.4
Non-Hispanic white	1.5	2.0	2.2	2.6	2.7	2.5	3.3	5.1	7.8
Male	1.6	2.3	2.0	2.7	2.6	2.5	3.2	5.5	10.8
Female	1.4	1.8	2.4	2.5	2.7	2.5	3.4	4.9	6.6
Non-Hispanic black	1.7	1.5	2.4	2.4	2.8	*	*	*	*
Male	2.1	*	*	*	*	*	*	*	*
Female	*	*	2.9	3.4	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Non-Hispanic 5/	8.6	7.9	8.5	10.5	14.5	19.3	30.6	57.4	211.6
Male	11.5	10.9	11.3	13.5	19.0	22.6	36.6	63.0	185.4
Female	5.7	5.0	5.8	7.7	10.5	16.7	26.5	54.1	222.3
Non-Hispanic white	8.0	7.1	7.7	9.7	13.2	18.7	30.0	57.0	215.8
Male	10.9	9.7	10.0	12.5	17.1	21.6	35.9	62.6	189.6
Female	5.2	4.5	5.6	7.1	9.7	16.3	25.9	53.6	226.4
Non-Hispanic black	13.9	16.0	15.8	19.1	28.3	30.3	44.7	74.1	189.2
Male	18.3	23.5	24.7	25.5	41.0	38.6	55.4	85.0	163.2
Female	10.1	9.7	8.5	14.1	18.9	24.7	38.3	68.7	198.9

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN,

RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All origins 3/	31.0	40.4	53.1	74.1	106.0	168.2	288.9	537.8	1,445.7
Male	38.5	47.6	60.9	83.1	117.6	183.1	303.4	537.6	1,266.2
Female	23.7	33.4	45.8	65.9	96.1	156.5	278.8	537.8	1,518.8
Hispanic	24.8	31.0	45.2	50.5	84.3	127.4	217.6	404.9	913.0
Male	33.3	38.7	59.2	55.8	105.9	138.1	221.5	378.4	870.7
Female	16.3	23.7	32.5	45.9	66.7	119.3	214.9	421.1	933.8
Mexican	25.9	26.4	45.1	59.1	88.1	133.4	242.0	*	*
Male	34.6	30.4	55.1	58.1	110.7	143.8	*	*	*
Female	16.8	22.1	35.5	60.2	69.1	125.3	*	*	*
Puerto Rican	32.2	54.1	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Cuban	*	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*
Other Hispanic 4/	22.3	34.9	32.7	41.2	66.8	*	*	*	*
Male	29.3	48.5	*	*	*	*	*	*	*
Female	16.3	23.7	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN,

RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Non-Hispanic 5/	31.3	40.9	53.4	75.6	107.1	169.8	291.3	541.5	1,462.0
Male	38.7	48.0	60.8	84.6	117.8	184.8	306.1	542.6	1,279.8
Female	24.2	34.1	46.5	67.3	97.8	158.0	281.0	540.9	1,535.7
Non-Hispanic white	27.3	37.3	49.3	71.2	103.2	165.9	287.8	540.7	1,481.3
Male	34.7	44.1	56.1	78.5	112.4	178.8	300.6	538.2	1,288.3
Female	19.9	30.7	42.7	64.4	95.2	155.5	278.8	542.2	1,559.1
Non-Hispanic black	65.4	77.2	97.4	124.6	159.8	237.4	375.9	624.5	1,403.3
Male	75.8	90.8	116.7	153.5	191.1	284.0	432.1	698.6	1,340.9
Female	56.4	65.7	81.6	101.8	136.6	206.4	342.1	587.2	1,426.5

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Non-Hispanic 5/	34.3	30.5	29.8	32.3	35.5	48.6	76.0	125.4	276.1
Male	51.4	45.4	42.8	45.0	47.9	64.0	100.1	166.5	363.1
Female	17.7	16.2	17.5	20.6	24.9	36.5	59.2	101.1	240.9
Non-Hispanic white	33.0	28.7	28.6	31.7	34.5	47.8	76.2	128.1	285.6
Male	49.1	42.2	40.2	43.3	45.4	62.5	98.8	168.3	373.8
Female	17.1	15.5	17.6	20.9	24.9	35.9	60.3	104.3	250.0
Non-Hispanic black	47.9	47.6	42.7	39.6	46.8	60.4	75.9	100.3	183.2
Male	76.2	77.4	72.6	65.3	75.2	87.1	120.4	156.2	277.1
Female	23.5	22.3	18.3	19.3	26.0	42.7	49.1	72.2	148.3

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Transport accidents (V01-V99,Y85)									
Non-Hispanic 5/	16.1	15.2	15.6	16.3	16.7	18.9	25.1	30.0	31.7
Male	23.6	21.9	22.2	22.3	22.5	25.5	34.3	44.3	60.4
Female	8.9	8.9	9.5	10.9	11.8	13.8	18.8	21.7	20.1
Non-Hispanic white	15.9	14.8	15.4	16.4	16.5	18.7	25.3	30.9	32.4
Male	23.2	21.1	21.4	21.9	21.6	24.9	34.0	44.8	62.3
Female	8.7	8.5	9.6	11.3	12.0	13.7	19.2	22.6	20.4
Non-Hispanic black	19.8	19.6	18.4	16.4	18.6	21.5	19.4	20.7	21.9
Male	30.0	30.3	31.5	27.2	31.7	33.8	34.7	39.5	45.8
Female	11.0	10.5	7.8	7.9	8.8	13.4	10.2	11.2	13.1

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.

RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE TENTH REVISION, INTERNATIONAL CLASSIFICATION OF DISEASES, 1992), HISPANIC ORIGIN, RACE FOR NON-HISPANIC POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)									
Non-Hispanic 5/	14.5	13.6	14.1	14.7	15.1	17.7	23.9	28.6	30.6
Male	21.1	19.2	19.6	19.5	19.7	23.2	32.0	41.7	58.3
Female	8.2	8.2	9.0	10.3	11.1	13.3	18.2	20.9	19.4
Non-Hispanic white	14.3	13.0	13.9	14.7	14.9	17.4	24.1	29.3	31.3
Male	20.7	18.3	18.9	18.9	18.8	22.7	31.7	42.0	60.0
Female	8.0	7.8	9.1	10.7	11.4	13.2	18.7	21.8	19.8
Non-Hispanic black	18.0	18.6	17.0	15.5	16.9	20.3	18.1	19.8	20.7
Male	27.0	28.5	28.6	25.5	28.2	31.8	32.3	38.6	44.6
Female	10.3	10.2	7.5	7.5	8.5	12.7	9.6	10.4	11.8

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Nontransport accidents (W00-X59,Y86)									
Non-Hispanic 5/	18.1	15.2	14.2	15.9	18.8	29.7	50.8	95.3	244.4
Male	27.8	23.5	20.6	22.7	25.4	38.5	65.9	122.2	302.7
Female	8.8	7.3	8.1	9.7	13.1	22.7	40.4	79.5	220.8
Non-Hispanic white	17.2	13.9	13.2	15.3	18.0	29.1	50.9	97.3	253.2
Male	25.9	21.1	18.8	21.4	23.8	37.6	64.9	123.5	311.5
Female	8.5	6.9	7.9	9.6	12.9	22.2	41.0	81.7	229.7
Non-Hispanic black	28.1	28.0	24.3	23.2	28.3	38.9	56.5	79.6	161.2
Male	46.2	47.1	41.1	38.1	43.4	53.3	85.7	116.7	231.3
Female	12.5	11.8	10.5	11.4	17.1	29.3	38.9	60.9	135.2

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Falls (W00-W19)									
Non-Hispanic 5/	2.0	2.6	3.3	4.6	6.2	12.1	22.2	44.5	113.5
Male	3.3	4.0	4.9	6.8	8.4	15.7	29.5	57.9	146.5
Female	0.8	1.2	1.8	2.5	4.3	9.3	17.2	36.7	100.1
Non-Hispanic white	2.1	2.5	3.2	4.5	6.1	12.5	23.2	46.7	120.2
Male	3.4	3.7	4.6	6.8	8.2	16.2	30.4	60.3	154.1
Female	0.8	1.2	1.8	2.5	4.3	9.5	18.2	38.6	106.5
Non-Hispanic black	2.0	3.6	4.6	5.1	7.1	9.7	11.8	20.7	39.1
Male	3.3	6.3	8.0	8.5	10.7	12.3	18.8	26.6	61.1
Female	*	*	*	*	4.5	7.9	7.6	17.7	31.0

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental exposure to smoke, fire and flames (X00-X09)

Non-Hispanic 5/	1.2	1.3	1.4	1.7	2.0	2.4	3.4	4.3	6.3
Male	1.6	1.9	2.0	2.3	2.5	3.2	4.4	5.9	9.3
Female	0.7	0.8	0.9	1.2	1.7	1.7	2.8	3.4	5.0
Non-Hispanic white	1.0	1.2	1.1	1.4	1.7	2.0	2.8	3.5	5.1
Male	1.4	1.7	1.6	1.8	2.0	2.6	3.6	4.9	8.1
Female	0.6	0.7	0.7	1.0	1.5	1.4	2.3	2.6	3.9
Non-Hispanic black	2.2	2.5	3.9	4.6	5.1	7.2	11.1	16.1	20.7
Male	3.2	4.0	5.6	7.2	7.7	10.6	15.9	21.5	28.2
Female	*	*	*	*	*	5.0	8.1	13.4	17.9

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Non-Hispanic 5/	9.9	5.9	3.2	2.2	1.5	1.5	1.6	2.6	4.4
Male	14.4	8.5	3.8	2.7	1.7	1.7	1.7	2.9	6.2
Female	5.5	3.3	2.6	1.8	1.4	1.3	1.6	2.4	3.6
Non-Hispanic white	9.2	5.0	2.9	2.1	1.5	1.4	1.7	2.7	4.1
Male	13.2	7.0	3.0	2.3	1.4	1.6	1.7	3.0	5.8
Female	5.3	3.0	2.7	1.9	1.5	1.3	1.7	2.5	3.4
Non-Hispanic black	16.8	14.3	6.9	3.8	2.6	*	*	*	8.3
Male	27.0	23.2	11.7	6.8	*	*	*	*	*
Female	8.1	6.7	3.0	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Non-Hispanic 5/	4.0	4.2	5.1	6.4	7.9	12.4	22.3	42.0	118.5
Male	6.6	6.8	7.9	9.1	10.9	15.8	28.5	52.4	136.9
Female	1.4	1.7	2.4	3.9	5.4	9.8	18.0	35.8	111.1
Non-Hispanic white	3.8	4.1	4.9	6.2	7.5	12.0	21.9	42.5	122.0
Male	6.1	6.5	7.4	8.6	10.2	15.3	27.5	52.0	139.7
Female	1.5	1.7	2.4	4.0	5.2	9.4	18.0	36.9	114.9
Non-Hispanic black	5.8	5.8	8.1	8.8	12.6	18.1	30.8	39.4	90.9
Male	10.4	10.3	14.2	14.2	18.7	23.6	47.7	64.4	124.5
Female	1.8	*	3.0	4.5	8.1	14.5	20.6	26.8	78.5

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
Intentional self-harm (suicide) (X60-X84,Y87.0)									
Non-Hispanic 5/	15.4	14.4	13.1	11.8	11.5	14.0	16.8	19.4	19.9
Male	24.2	22.0	20.6	19.1	20.0	26.5	35.5	44.9	58.4
Female	7.0	7.1	6.1	5.2	4.1	4.1	3.8	4.4	4.3
Non-Hispanic white	17.3	15.9	14.4	13.1	12.6	15.0	18.0	20.6	21.3
Male	26.8	24.0	22.4	21.1	21.6	28.0	37.9	47.9	62.7
Female	8.0	7.9	6.8	5.8	4.6	4.5	4.0	4.4	4.6
Non-Hispanic black	6.6	6.4	4.6	3.8	4.4	6.3	5.4	6.6	*
Male	11.6	10.4	8.3	6.6	9.6	13.7	12.0	*	*
Female	2.2	3.0	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Assault (homicide) (X85-Y09,Y87.1)

Non-Hispanic 5/	5.0	3.8	3.2	2.5	2.4	2.3	2.3	2.6	2.3
Male	7.4	5.6	4.9	3.7	3.2	3.2	2.7	3.8	3.1
Female	2.6	2.0	1.6	1.4	1.6	1.5	2.0	1.9	2.0
Non-Hispanic white	3.2	2.6	2.4	1.8	1.7	1.8	2.0	2.0	2.1
Male	4.3	3.5	3.5	2.6	2.1	2.4	2.3	2.6	2.6
Female	2.0	1.7	1.4	1.2	1.4	1.3	1.8	1.6	1.9
Non-Hispanic black	17.1	12.4	9.2	7.5	8.1	7.0	5.4	9.8	*
Male	29.2	22.5	16.3	13.4	14.9	12.6	*	19.7	*
Female	6.5	3.8	3.5	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
 RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
 OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
 BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
 HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
 TENTH REVISION, INTERNATIONAL
 CLASSIFICATION OF DISEASES,
 1992), HISPANIC ORIGIN,
 RACE FOR NON-HISPANIC
 POPULATION, AND SEX

	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Assault (homicide) by other and unspecified means and their sequelae (X85-X92,X96-Y09,Y87.1)

Non-Hispanic 5/	2.3	1.8	1.5	1.3	1.4	1.3	1.5	2.1	1.6
Male	3.2	2.5	2.4	1.9	1.9	2.0	1.9	3.1	2.4
Female	1.4	1.0	0.7	0.8	0.9	0.8	1.2	1.5	1.3
Non-Hispanic white	1.5	1.2	1.1	0.9	0.9	1.0	1.3	1.6	1.5
Male	1.9	1.6	1.6	1.3	1.1	1.5	1.5	2.2	1.9
Female	1.1	0.9	0.5	0.5	0.7	0.7	1.1	1.2	1.3
Non-Hispanic black	7.4	5.5	5.0	4.8	5.3	4.2	4.2	6.9	*
Male	12.2	9.5	9.0	7.9	9.6	6.8	*	*	*
Female	3.3	2.0	*	*	*	*	*	*	*

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP.)

RATES ARE BASED ON POPULATIONS ENUMERATED AS OF APRIL 1, 2000 AND MAY DIFFER FROM RATES PREVIOUSLY PUBLISHED.
RACE AND HISPANIC ORIGIN ARE REPORTED SEPARATELY ON THE DEATH CERTIFICATE. PERSONS OF HISPANIC ORIGIN MAY BE
OF ANY RACE. DATA FOR HISPANIC PERSONS ARE NOT TABULATED SEPARATELY BY RACE; DATA FOR NON-HISPANIC PERSONS ARE TABULATED
BY RACE. DATA FOR HISPANIC ORIGIN SHOULD BE INTERPRETED WITH CAUTION BECAUSE OF INCONSISTENCIES BETWEEN REPORTING
HISPANIC ORIGIN ON DEATH CERTIFICATES AND ON CENSUSES AND SURVEYS)

CAUSE OF DEATH (BASED ON THE
TENTH REVISION, INTERNATIONAL
CLASSIFICATION OF DISEASES,
1992), HISPANIC ORIGIN,

RACE FOR NON-HISPANIC POPULATION, AND SEX	45-49 YEARS	50-54 YEARS	55-59 YEARS	60-64 YEARS	65-69 YEARS	70-74 YEARS	75-79 YEARS	80-84 YEARS	85 YEARS AND OVER
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Complications of medical and surgical care (Y40-Y84,Y88)

Non-Hispanic 5/	0.7	1.0	1.2	2.1	2.7	4.4	5.8	8.1	13.7
Male	0.6	1.0	1.2	2.1	2.8	5.3	6.9	10.4	16.1
Female	0.7	0.9	1.2	2.1	2.6	3.7	5.0	6.8	12.7
Non-Hispanic white	0.6	0.9	1.1	1.8	2.5	4.1	5.5	8.0	13.2
Male	0.6	1.0	1.2	1.8	2.5	5.0	6.8	10.6	15.9
Female	0.6	0.9	1.0	1.8	2.5	3.5	4.5	6.5	12.0
Non-Hispanic black	1.2	1.7	2.7	4.8	4.8	8.0	10.5	10.6	21.3
Male	*	*	*	4.7	6.4	9.6	10.1	*	*
Female	*	*	3.2	4.9	*	7.0	10.7	10.8	22.2

* FIGURE DOES NOT MEET STANDARDS OF RELIABILITY OR PRECISION.

0.0 QUANTITY MORE THAN ZERO BUT LESS THAN 0.05.

... CATEGORY NOT APPLICABLE.

1/ FIGURES FOR AGE NOT STATED INCLUDED IN "ALL AGES" BUT NOT DISTRIBUTED AMONG AGE GROUPS.

2/ DEATH RATES FOR "UNDER 1 YEAR" (BASED ON POPULATION ESTIMATES) DIFFER FROM INFANT MORTALITY RATES (BASED ON LIVE BIRTHS).

3/ ALL ORIGINS INCLUDES ORIGIN NOT STATED; SPECIFIED ORIGINS EXCLUDE ORIGIN NOT STATED.

4/ INCLUDES CENTRAL AND SOUTH AMERICAN AND OTHER AND UNKNOWN HISPANIC.

5/ INCLUDES RACES OTHER THAN WHITE AND BLACK.

SOURCE: CDC/NCHS, NATIONAL VITAL STATISTICS SYSTEM, MORTALITY.