

Indiana, 2009

Primary and secondary syphilis

- CASES: Indiana reported 158 cases in 2009.
- RATES: Indiana ranked 28 among 50 states; Washington, DC; and 3 territories with 2.5 cases per 100,000 population compared to the U.S. rate of 4.6 cases per 100,000 population.

Rates of primary and secondary syphilis, by sex

- In Indiana, the rate among males was 4.6 per 100,000 population compared to the U.S. male rate of 7.8 per 100,000.
- The rate among females was 0.4 per 100,000 compared to the U.S. female rate of 1.4 per 100,000.


Rates of primary and secondary syphilis, by race/ethnicity

- In Indiana, the race/ethnicity adjusted rates per 100,000 population were 1.4 among whites, 12.5 among blacks, 1.8 among Hispanics, 3.2 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.
- The rate among blacks was 8.9 times that of whites.
- The rate among Hispanics was 1.3 times that of whites.
- The rate among Asian/Pacific Islanders was 2.3 times that of whites.


Congenital syphilis

- CASES: Indiana reported 1 case in 2009.
- RATES: Indiana ranked 34 among among 34 areas (33 states and 1 territory) reporting congenital syphilis cases with a rate of 1.1 cases per 100,000 live births compared to the U.S. rate of 10.0 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Indiana, 2000-2009



Total P&S Syphilis Rates by County, Indiana, 2009



Rate (per 100,000 population)

