

# Kentucky, 2007

## Primary and secondary syphilis

- **CASES:** Kentucky reported 56 cases in 2007.
- **RATES:** Kentucky ranked 33 among 50 states, DC, and 3 territories with 1.3 cases per 100,000 population compared to the total rate of 3.8 cases per 100,000 population.

## Rates of primary and secondary syphilis, by sex

- In Kentucky, the rate among males was 2.3 per 100,000 population compared to the total male rate of 6.6 per 100,000.
- The rate among females was 0.4 per 100,000 compared to the total female rate of 1.1 per 100,000.

## Rates of primary and secondary syphilis, by race/ethnicity


In Kentucky, the race/ethnicity adjusted rates per 100,000 population were 0.9 among whites, 6.5 among blacks, 1.2 among Hispanics, 0.0 among Asian/Pacific Islanders, and 11.0 among American Indians/Alaska Natives.

- The rate among blacks was 7.2 times that of whites.
- The rate among Hispanics was 1.3 times that of whites.
- The rate among American Indians/Alaska Natives was 12.2 times that of whites.

## Congenital syphilis

- **CASES:** Kentucky reported 0 cases in 2007.
- **RATES:** Kentucky had a rate of 0.0 cases per 100,000 live births compared to the total rate of 10.5 cases per 100,000.

## Syphilis: P&S Rates and Congenital Cases, Kentucky, 1998-2007



## Total P&S Syphilis Rates by County, Kentucky, 2007

