

Topic

- Cases and Deaths
- Case Rates
- Death Rates
- Syndromes
- Serotypes
- Age and Serotypes
- Surveillance Report

Year

2024

Dataset version: Feb 2026
Final run: Mar 23, 2026

Note: Click [here](#) to access and download Surveillance Reports.

Group A *Streptococcus* (GAS) | Group B *Streptococcus* (GBS) | ***Haemophilus influenzae* (HFlu)** | *Neisseria meningitidis* (NMen) | *Streptococcus pneumoniae* (SPN) | Bact Facts

Active Bacterial Core Surveillance (ABCs) Report
Emerging Infections Program Network
***Haemophilus influenzae*, 2024**

ABCs Areas: California (3 county San Francisco Bay area); Colorado (5 county Denver area); Connecticut; Georgia; Maryland; Minnesota; New Mexico; New York (15 county Rochester and Albany areas); Oregon; Tennessee (20 urban counties)

ABCs Population: The surveillance areas represent 46,473,142 persons. Source: Census Bureau's Vintage 2024 population estimates.

ABCs Case Definition: For routine ABCs surveillance, a case of invasive bacterial disease is defined as isolation of *Haemophilus influenzae* from a normally sterile site or detection of ABCs pathogen-specific nucleic acid in a specimen obtained from a normally sterile body site, using a validated molecular test in a resident of one of the surveillance areas.

ABCs Methodology: ABCs personnel routinely contacted all microbiology laboratories serving acute care hospitals in their area to identify cases. Standardized case report forms that include information on demographic characteristics, clinical syndrome, and outcome of illness were completed for each identified case. Serotyping was done on *Haemophilus influenzae* isolates at CDC and state laboratories. Regular laboratory audits assessed completeness of active surveillance and detected additional cases.

All rates of invasive *Haemophilus influenzae* disease were calculated using population estimates from the Vintage 2024 file. For national estimates, race- and age-specific rates of disease were applied from the aggregate surveillance areas to the race- and age-specific distribution of the U.S. population. Cases with missing data, excluding ethnicity, were multiply imputed using sequential regression imputation methods.† Cases with unknown serotypes were distributed based on age and race distribution for known cases.

ABCs Profiles

Race	No.	Rate*
Black	272	3.2
White	765	2.4
Other	79	1.4
Total	1,116	2.4

National Estimates of Invasive Disease

Total Cases: 8,170 (2.4/100,000 population)

Deaths: 880 (0.26/100,000 population)

Serotype	B		Non-B		Non-Type†	
	No.	Rate*	No.	Rate*	No.	Rate*
<1	2	0.45	6	1.23	34	6.95
1	0	0.04	5	1.00	10	1.97
2-4	0	0.00	5	0.33	14	0.92
5-17	0	0.00	15	0.20	20	0.27
18-34	1	0.01	14	0.13	65	0.61
35-49	1	0.01	30	0.32	115	1.24
50-64	4	0.05	67	0.78	142	1.65
65-74	4	0.09	61	1.28	166	3.49
75-84	1	0.04	35	1.37	164	6.39
≥85	0	0.00	16	1.91	117	13.97
Total	14	0.03	254	0.55	846	1.82

*Per 100,000 population for ABCs areas †Non-typeable isolates

Syndromes

Syndrome	Cases		Deaths	
	No.	%*	No.	Rate†
Bacteremia Without Focus	192	17.2	18	9.6
Meningitis	98	8.8	2	2.0
Pneumonia With Bacteremia	627	56.2	77	12.3

*Percent of cases
†Deaths per 100 cases with known outcome

† Surveillance Note

Missing race (n=146) data were multiply imputed using sequential regression imputation methods.

Citation

Centers for Disease Control and Prevention. 2024. Active Bacterial Core Surveillance Report, Emerging Infections Program Network, *Haemophilus influenzae*, 2024. www.cdc.gov/abcs/downloads/HFLU_Surveillance_Report_2024.pdf