

Adolescent Behaviors and Experiences Survey

	Grade 9		Grade 10		Grade 11		Grade 12	
Mental Health	Percentage	Confidence Interval	Percentage	Confidence Interval	Percentage	Confidence Interval	Percentage	Confidence Interval
Who felt sad or hopeless (almost every day for two or more weeks in a row so that they stopped doing some usual activities, ever during the 12 months before the survey)	43.3	38.6 - 48.1	44.6	40.8 - 48.4	44.7	40.6 - 48.8	44.4	40.7 - 48.2
Seriously considered attempting suicide (during the 12 months before the survey)	22.1	18.3 - 26.4	19.6	17.0 - 22.5	18.5	15.9 - 21.3	19.2	16.3 - 22.6
Made a plan about how they would attempt suicide (during the 12 months before the survey)	15.6	12.7 - 18.9	15.9	13.7 - 18.3	15.0	12.2 - 18.3	14.8	11.4 - 18.9
Actually attempted suicide (one or more times during the 12 months before the survey)	10.3	7.8 - 13.6	9.9	8.1 - 12.2	8.4	6.4 - 10.9	7.2	4.9 - 10.4
Had a suicide attempt that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)	2.0	1.2 - 3.3	1.7	1.0 - 2.7	2.0	1.2 - 3.3	1.8	1.1 - 3.0
Reported that their mental health was most of the time or always not good (including stress, anxiety, and depression, during the 30 days before the survey)	30.2	26.7 - 33.9	31.3	27.8 - 35.0	30.2	26.7 - 34.0	32.8	29.0 - 36.8