

Adolescent Behaviors and Experiences Survey														
Sexual Behaviors	American Indian or Alaska Native*		Asian*		Black or African American*		Hispanic or Latino*		Native Hawaiian or Other Pacific Islander*		White*		Multiple Race*	
	Percentage	Confidence Interval	Percentage	Confidence Interval	Percentage	Confidence Interval	Percentage	Confidence Interval	Percentage	Confidence Interval	Percentage	Confidence Interval	Percentage	Confidence Interval
Ever had sexual intercourse	43.4	25.2 - 63.6	13.7	9.6 - 19.2	28.9	24.1 - 34.2	26.7	21.9 - 32.1	N/A	N/A	29.9	26.9 - 33.0	36.5	27.2 - 46.8
Had sexual intercourse for the first time before age 13 years	8.2	3.0 - 20.4	0.6	0.2 - 2.3	4.3	3.0 - 6.0	2.6	1.9 - 3.6	N/A	N/A	1.2	0.8 - 2.0	5.1	2.8 - 9.1
Had sexual intercourse with four or more persons during their life	11.9	6.0 - 22.3	0.8	0.2 - 3.3	7.8	5.5 - 10.8	3.8	2.7 - 5.3	N/A	N/A	6.2	4.6 - 8.2	10.6	7.1 - 15.6
Were currently sexually active (had sexual intercourse with at least one person, during the 3 months before the survey)	27.5	17.6 - 40.2	10.1	7.2 - 14.1	18.3	14.9 - 22.2	14.7	11.9 - 18.0	N/A	N/A	20.9	18.4 - 23.6	26.2	18.1 - 36.4
Drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)	N/A	N/A	1.6	0.2 - 10.1	17.8	13.0 - 23.9	16.5	11.6 - 22.9	N/A	N/A	17.1	13.4 - 21.5	18.4	9.7 - 32.3
Used a condom during last sexual intercourse (among students who were currently sexually active)	N/A	N/A	49.1	32.2 - 66.2	54.3	45.1 - 63.2	53.3	45.7 - 60.8	N/A	N/A	55.0	50.9 - 59.0	50.4	37.0 - 63.6
Used birth control pills before last sexual intercourse with opposite-sex partner (to prevent pregnancy, not counting emergency contraception such as Plan B or the "morning after" pill, among students who were currently sexually active)	N/A	N/A	N/A	N/A	11.5	6.6 - 19.2	13.7	9.2 - 19.9	N/A	N/A	26.5	21.9 - 31.7	25.6	17.4 - 36.0
Were ever tested for human immunodeficiency virus (HIV) (not counting tests done if they donated blood)	7.4	3.2 - 16.0	5.2	2.9 - 9.0	7.2	5.8 - 8.9	4.6	3.3 - 6.2	N/A	N/A	6.3	5.1 - 7.9	5.6	3.2 - 9.5
Were tested for a sexually transmitted disease (STD) other than HIV, such as chlamydia or gonorrhea (during the 12 months before the survey)	8.5	2.9 - 22.4	4.2	1.6 - 10.1	7.1	4.8 - 10.3	2.9	2.1 - 4.0	N/A	N/A	4.8	3.8 - 6.2	5.6	2.7 - 11.1
Used an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon) before last sexual intercourse with an opposite-sex partner (to prevent pregnancy, among students who were currently sexually active)	N/A	N/A	N/A	N/A	3.4	1.1 - 10.1	6.7	3.7 - 11.7	N/A	N/A	6.4	4.5 - 9.1	4.7	2.4 - 8.8
Used birth control pills; an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon); or a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse with an opposite-sex partner (to prevent pregnancy, among students who were currently sexually active)	N/A	N/A	N/A	N/A	19.9	13.9 - 27.7	26.2	20.3 - 33.2	N/A	N/A	38.6	32.9 - 44.6	39.1	27.3 - 52.5
Used both a condom during last sexual intercourse and birth control pills; an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon); or a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse with an opposite-sex partner (to prevent pregnancy, among students who were currently sexually active)	N/A	N/A	3.3	0.4 - 21.2	5.8	2.5 - 12.6	7.9	5.0 - 12.1	N/A	N/A	13.9	11.2 - 17.2	15.9	8.8 - 27.0
Did not use any method to prevent pregnancy during last sexual intercourse with an opposite-sex partner (among students who were currently sexually active)	N/A	N/A	N/A	N/A	23.7	18.4 - 30.0	14.1	9.0 - 21.4	N/A	N/A	11.1	8.3 - 14.5	18.0	11.0 - 28.0

*Non-Hispanic

N/A: Less than 30 respondents for the subgroup